BIOLOGICAL SCIENCES: Biochemistry

Inhibitory Proteins Block Substrate Access to the Active Site of *Bacillus subtilis*

Intramembrane Metalloprotease SpoIVFB

Sandra Olenic[a], Lim Heo[a], Michael Feig[a], and Lee Kroos[a,1]

[a]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing,

MI 48824

[1]To whom correspondence should be addressed. Email: kroos@msu.edu

Abstract

Intramembrane proteases of diverse signaling pathways use membrane-embedded active sites to

cleave membrane-associated substrates. Interactions of intramembrane metalloproteases with

modulators are poorly understood. Inhibition of *Bacillus subtilis* intramembrane metalloprotease

SpoIVFB requires BofA and SpoIVFA, which transiently prevent cleavage of Pro-σ^K during

endosporulation. Three conserved BofA residues (N48, N61, T64) in or near predicted

transmembrane segment (TMS) 2 were found to be required for SpoIVFB inhibition. Disulfide

cross-linking indicated that BofA TMS2 occupies the SpoIVFB active site region. BofA and

SpoIVFA neither prevented SpoIVFB from interacting with Pro-σ^K in co-purification assays nor

interfered with cross-linking between the C-terminal regions of Pro-σ^K and SpoIVFB. However,

BofA and SpoIVFA did interfere with cross-linking between the N-terminal Proregion of Pro-σ^K

and the SpoIVFB active site region and interdomain linker. A BofA variant lacking predicted

TMS1, in combination with SpoIVFA, was less effective at interfering with some of the cross-

links and slightly less effective at inhibiting cleavage of Pro-σ^K by SpoIVFB. A structural model

was built of SpoIVFB in complex with BofA and parts of SpoIVFA and Pro-σ^K, using partial

homology and constraints from cross-linking and co-evolutionary analyses. The model predicts

that N48 in BofA TMS2 interacts with T64 (and possibly N61) of BofA to stabilize a membrane-

embedded C-terminal region. SpoIVFA is predicted to bridge the BofA C-terminal region and

SpoIVFB. Thus, the two inhibitory proteins block access of the Pro-σ^K N-terminal region to the

SpoIVFB active site region. Our findings may inform efforts to develop selective inhibitors of

intramembrane metalloproteases.

Significance

Intramembrane metalloproteases (IMMPs) function in numerous signaling processes that impact

health. For example, IMMPs function in pathways regulating human cholesterol levels and

bacterial pathogenesis. Knowledge of IMMP interactions with modulators could inform design

of therapeutics, but structures of complexes have not been reported. We found that a

transmembrane segment of BofA occupies the active site region to inhibit SpoIVFB, an IMMP

crucial for endosporulation of *Bacillus subtilis*. Because endosporulation enhances persistence

of related pathogenic bacteria, our structural model of SpoIVFB in complex with BofA and parts

of its other inhibitory protein SpoIVFA and its substrate Pro-σ^K, may lead to strategies to control

endosporulation. More broadly, insights from the SpoIVFB inhibition mechanism could guide

efforts to develop inhibitors of other IMMPs.

Introduction

Intramembrane proteases (IPs) are membrane proteins containing a membrane-embedded active site. IPs cleave membrane-associated substrates within a transmembrane segment (TMS) or near the membrane surface in a process referred to as regulated intramembrane proteolysis (RIP) (1). Released substrate fragments impact diverse signaling pathways in a wide variety of organisms (2). There are four known IP families: intramembrane metalloproteases (IMMPs) like SpoIVFB, aspartyl proteases like presenilins, serine proteases (rhomboids), and the glutamyl protease Rce1 (2, 3). Crystal structures have been solved for one or more IP in each family (3-8), revealing that TMSs arrange to form a channel that delivers water to the active site for hydrolysis of a substrate peptide bond. Structures have also been solved for rhomboid·peptide inhibitor complexes (9, 10) and for γ-secretase·substrate complexes (11, 12), which may guide the design of IP modulators as therapeutics or for other purposes.

IMMPs activate transcription factors via RIP in all three domains of life (1, 2). In humans, S2P is involved in regulation of cholesterol homeostasis and responses to endoplasmic reticulum stress and viral infection (13, 14). S2P homologs in plants and in the pathogenic fungus *Cryptococcus neoformans* play important roles in chloroplast development and virulence, respectively (15, 16). In bacteria, IMMPs enhance pathogenicity, control stress responses and polar morphogenesis, produce mating signals, and clear signal peptides from the membrane (17-19). In *Bacillus* and *Clostridium* species, SpoIVFB cleaves inactive Pro-σ^K to σ^K, which directs RNA polymerase to transcribe genes necessary for endospore formation (18, 20). Endospores are dormant and are able to survive harsh environmental conditions (21), enhancing the persistence of pathogenic species (22, 23). Knowledge of SpoIVFB interactions with its

inhibitory proteins, BofA and SpoIVFA (24-26), could lead to new strategies to manipulate

endosporulation and other processes involving IMMPs in bacteria and eukaryotes.

During endospore formation, *B. subtilis* forms a polar septum that divides the mother cell

(MC) and forespore (FS) compartments (27) (*SI Appendix*, Fig. S1). The MC engulfs the FS,

surrounding it with a second membrane and pinching it off within the MC. SpoIVFB, BofA, and

SpoIVFA form a complex in the outer FS membrane during engulfment (28-30). BofA was

implicated as the direct inhibitor of SpoIVFB (31), while SpoIVFA appeared to localize and

stabilize the heterotrimer (30, 32). Signaling from the FS relieves inhibition of SpoIVFB (*SI*

Appendix, Fig. S1). Two proteases, SpoIVB and CtpB, are exported into the space between the

two membranes surrounding the FS (33, 34). SpoIVB cleaves the C-terminal end of SpoIVFA

(35-38) and CtpB can cleave the C-terminal ends of both SpoIVFA and BofA (36, 37, 39). Once

inhibition is removed, SpoIVFB cleaves the N-terminal 21-residue Proregion from Pro-σ^K,

releasing σ^K into the MC (40-42). σ^K directs RNA polymerase to transcribe genes whose

products form the spore coat and lyse the MC, releasing a mature spore (43, 44).

Cleavage of Pro-σ^K by SpoIVFB can be reproduced by expressing these proteins in *Escherichia*

coli (31) (Fig. 1A). Cleavage could be inhibited by additionally producing GFPΔ27BofA (a

functional fusion protein lacking predicted TMS1 of BofA) and SpoIVFA (31, 39). However,

we found that the extent of cleavage inhibition was incomplete and variable. Here, we describe

an improved cleavage inhibition assay. Using the assay and bioinformatics, we identified three

conserved residues of BofA important for inhibition of SpoIVFB in *E. coli* and in sporulating *B.*

subtilis. One of the residues, N48, is near the middle of BofA TMS2, close to the sole Cys

residue of BofA, C46. We exploited C46 for disulfide cross-linking experiments that showed

TMS2 of BofA can occupy the SpoIVFB active site region. These experiments relied on single-

Cys variants of SpoIVFB created previously and used to cross-link residues predicted to be near

the active site with residues near the cleavage site in single-Cys variants of Pro-σ^K (45). We

created additional single-Cys variants of SpoIVFB, Pro-σ^K, and BofA, as well as functional Cys-

less SpoIVFA, and performed disulfide cross-linking experiments. We found that BofA and

SpoIVFA interfere with cross-linking between the N-terminal Proregion of Pro-σ^K and the

SpoIVFB active site region, indicating that the inhibitory proteins block substrate access to the

IMMP active site. The results were used in combination with prior cross-linking studies (45,

46), partial homology, and evolutionary co-variation of amino acid residues, to constrain a

structural model of SpoIVFB in complex with BofA and parts of SpoIVFA and Pro-σ^K. The

model predicts that conserved residues in TMS2 and the C-terminal region of BofA interact, and

that SpoIVFA bridges the BofA C-terminal region and SpoIVFB to form a membrane-embedded

inhibitory complex.

Results

BofA and SpoIVFA Can Prevent Cleavage of Pro-σ^K by SpoIVFB in *E. coli*. To study RIP

of Pro-σ^K, *E. coli* was engineered to synthesize variants of SpoIVFB and Pro-σ^K, and the two

inhibitory proteins, BofA and SpoIVFA, in various combinations. The SpoIVFB variant cytTM-

SpoIVFB-FLAG$_2$-His$_6$ (cytTM-SpoIVFB) (Fig. 1A) contains cytTM (47), which improves

accumulation (42). The substrate variant Pro-σ^K(1-127)-His$_6$ [Pro-σ^K(1-127)] lacks the C-

terminal half of Pro-σ^K, but can be cleaved by SpoIVFB and the cleavage product can easily be

separated from Pro-σ^K(1-127) by SDS-PAGE (48) [Note: Pro-σ^K(1-126) was renamed Pro-σ^K(1-

127) as explained in (46)]. When Pro-σ^K(1-127) was produced with cytTM-SpoIVFB, cleavage

was abundant (Fig. 1B, lane 1), as shown previously (42). To quantify cleavage, we measured

the abundance of cleavage product and divided that by the amount of Pro-σ^K(1-127) plus

cleavage product, which resulted in a cleavage ratio of 0.78 $\pm$ 0.02. The additional production of

GFPΔ27BofA and SpoIVFA from a second plasmid reduced the cleavage ratio to 0.3 $\pm$ 0.08, but

cleavage product was visible (lane 2). A similar result was seen when full-length BofA (for

which no antibody has been described) and SpoIVFA were produced from a second plasmid

(lane 3). Since the second plasmid had the same replication origin as the plasmid used to

produce Pro-σ^K(1-127) and cytTM-SpoIVFB (i.e., the two plasmids were "incompatible"), the

copy number of the two plasmids may be unequal in some cells, despite using different selection

markers to try to maintain both plasmids. Unequal copy numbers could result in Pro-σ^K(1-127)

cleavage in cells producing too little of the inhibitory proteins. Therefore, we designed single

plasmids to synthesize all four proteins, which we called "pET Quartet plasmids." When pET

Quartets produced either GFPΔ27BofA (lane 4) or BofA (lane 5), inhibition improved, resulting

in very little Pro-σ^K(1-127) cleavage. A longer exposure of the immunoblot revealed a faint

cleavage product with pET Quartet GFPΔ27BofA, but not with pET Quartet BofA, indicating

that full-length BofA inhibits cleavage slightly better than GFPΔ27BofA.

We tested whether both GFPΔ27BofA and SpoIVFA are required for cleavage inhibition

by engineering "pET Triplet plasmids" to synthesize Pro-σ^K(1-127), cytTM-SpoIVFB, and either

GFPΔ27BofA or SpoIVFA. With pET Triplet GFPΔ27BofA, cleavage of Pro-σ^K(1-127) was

observed (ratio = 0.7 $\pm$ 0.1), but accumulation of cytTM-SpoIVFB, Pro-σ^K(1-127), and cleavage

product was greatly diminished (Fig. 1C, lane 2), suggesting that GFPΔ27BofA inhibits

synthesis and/or enhances degradation of the other proteins when SpoIVFA is absent. With pET

Triplet SpoIVFA (lane 3), the cleavage ratio (0.6 $\pm$ 0.05) and accumulation of cytTM-SpoIVB,

Pro-σ^K(1-127), and cleavage product were similar to the control plasmid that synthesized only

cytTM-SpoIVFB and Pro-σ^K(1-127) (lane 1). We conclude that both GFPΔ27BofA and

SpoIVFA are necessary to strongly inhibit Pro-σ^K(1-127) cleavage by cytTM-SpoIVB in *E. coli*

(lane 4). Both inhibitory proteins are required to prevent cleavage of Pro-σ^K by SpoIVFB during

B. subtilis sporulation (24, 30, 49).

139   Efficient relief of SpoIVFB inhibition by BofA and SpoIVFA during *B. subtilis*

sporulation requires SpoIVB and CtpB (*SI Appendix*, Fig. S1), but SpoIVB alone can partially

relieve inhibition (34, 36, 37, 39). Likewise, SpoIVB partially relieved inhibition when

produced in *E. coli* from "pET Quintet plasmids" (i.e., pET Quartet plus SpoIVB) (*SI Appendix*,

Results and Fig. S2). In the absence of SpoIVB, an F66A substitution in SpoIVFB-YFP can

partially overcome inhibition by BofA and SpoIVFA during *B. subtilis* sporulation (49). The

F66A substitution in cytTM-SpoIVFB likewise partially overcame inhibition by GFPΔ27BofA

and SpoIVFA when produced from pET Quartet in *E. coli* (*SI Appendix*, Results and Fig. S3).

Hence, the heterologous system mimics the endogenous pathway of *B. subtilis* in these respects.

We next exploited the improved cleavage inhibition assay based on pET Quartet plasmids in *E.*

coli to identify residues of BofA important for SpoIVFB inhibition.

Three Conserved Residues of BofA Are Important for Inhibition of SpoIVFB in *E. coli*. To

identify residues of BofA that may play a role in SpoIVFB inhibition, an alignment of 70 BofA

orthologs was made (*SI Appendix*, Fig. S4A). The majority of conserved residues reside in

predicted TMS2 and the C-terminal region, which could explain why GFPΔ27BofA (lacking

predicted TMS1) is a functional inhibitor (30, 31) (Fig. 1B). Most endospore formers encode

BofA, but about half lack a recognizable gene for SpoIVFA (20, 50). A smaller alignment (*SI*

Appendix, Fig. S4B) was made to see if additional residues are conserved in BofA orthologs of

species encoding SpoIVFA. In total, 17 residues of interest were conserved in BofA: 13 within

TMS2 and the C-terminal region of the larger alignment, and 4 from the C-terminal region of the

smaller alignment.

We substituted conserved residues in GFPΔ27BofA with alanine to investigate their

roles. Since deletion of three residues from the C-terminal end of BofA caused a loss of function

or stability in *B. subtilis* (26, 51), we also made Ala substitutions for F85 and I87 (as well as I86,

which is conserved) in GFPΔ27BofA. Each GFPΔ27BofA variant was co-produced with Pro-

σ^K(1-127), cytTM-SpoIVFB, and SpoIVFA from pET Quartet plasmids in *E. coli*, and the Pro-

σ^K(1-127) cleavage ratio was quantified (Fig. 2). Three GFPΔ27BofA variants, N48A, N61A,

and T64A (lanes 4, 7 & 8), led to the greatest cleavage ratios, indicating that these substitutions

strongly impaired inhibition of cytTM-SpoIVFB. However, less of the N61A and T64A variants

accumulated than most of the other GFPΔ27BofA variants, and for all three variants, less

SpoIVFA and cytTM-SpoIVFB accumulated, indicating that the variants reduce the synthesis

and/or stability of proteins that normally form the SpoIVFB inhibition complex (24, 30, 31, 39,

49). We reasoned that accumulation of the proteins may differ in *E. coli* and sporulating *B.*

subtilis, so we examined the effects of GFPΔ27BofA N48A, N61A, and T64A variants during

sporulation (see below).

Importantly, deletion of three residues from the C-terminal end of GFPΔ27BofA strongly

impaired inhibition of cytTM-SpoIVFB in *E. coli* (*SI Appendix*, Results and Fig. S5A),

mimicking results obtained for the endogenous pathway of *B. subtilis* (26, 51). We also used

pET Quartet plasmids in *E. coli* to show that nine residues preceding predicted TMS2 of

GFPΔ27BofA contribute to its inhibitory function (*SI Appendix*, Results and Fig. S5B).

The Conserved Residues of BofA Are Important for SpoIVFB Inhibition during *B. subtilis*

Sporulation. A *B. subtilis spoIVB165 bofA::erm* double mutant provides a genetic background

to test the effects of Ala substitutions on BofA function during sporulation. The *spoIVB165* null

mutation alone would block Pro-σ^K cleavage, but cleavage is restored in the *spoIVB165*

bofA::erm double mutant, and ectopic production of GFPΔ27BofA under control of an MC-

specific promoter in the double mutant leads to inhibition of SpoIVFB (30, 31).

　　　　To test the effects of the three GFPΔ27BofA variants (N48A, N61A, and T64A) during

sporulation, each was produced ectopically under control of the *bofA* promoter in the *spoIVB165*

bofA::erm double mutant background (Fig. 3A). As a control, in wild-type *B. subtilis* strain

PY79, BofA and SpoIVFA inhibited Pro-σ^K cleavage until between 4 and 5 h poststarvation

(PS), and the levels of SpoIVFA and SpoIVFB decreased between 4 and 5 h PS (lanes 1 & 2).

For the *spoIVB165* single mutant, very little Pro-σ^K cleavage was observed, as expected, and the

SpoIVFA and SpoIVFB levels did not decrease at 5 h (lanes 3 & 4). For the *spoIVB165*

bofA::erm double mutant, inhibition of SpoIVFB was expected to be lost, and in agreement the

Pro-σ^K cleavage ratio at 4 h (lane 5) was greater than for the wild-type strain at 4 h (lane 1) and

similar to the ratios at 5 h (lanes 2 and 6). Very little SpoIVFA and SpoIVFB was detected

(lanes 5 & 6), consistent with the need for BofA to stabilize these proteins (30). As expected,

ectopic production of GFPΔ27BofA in the double mutant inhibited SpoIVFB so that little Pro-σ^K

cleavage was observed, and the SpoIVFA and SpoIVFB levels did not decrease at 5 h (lanes 7 &

8), similar to the *spoIVB165* single mutant (lanes 3 & 4). GFPΔ27BofA allowed a greater Pro-

σ^K cleavage ratio at 5 h (lane 8) than BofA (lane 4), suggesting that full-length BofA is a slightly

better inhibitor of SpoIVFB during *B. subtilis* sporulation, as observed in *E. coli* (Fig. 1B, longer

exposure).

Strikingly, ectopic production of the GFPΔ27BofA N48A or N61A variant in the double

mutant did not inhibit SpoIVFB, as Pro-σ^K was cleaved at 4 h (Fig. 3A, lanes 9 & 11). The level

of the N48A variant (lanes 9 & 10) was comparable with GFPΔ27BofA (lanes 7 & 8), but less of

the N61A variant was observed, especially at 5 h (lanes 11 & 12). In comparison with the strain

that produced GFPΔ27BofA, the SpoIVFA level was normal in the strain that produced the

N48A variant, but less SpoIVFB was observed, and less of both SpoIVFA and SpoIVFB was

observed in the strain producing the N61A variant. Production of the T64A variant in the double

mutant (lanes 13 & 14) showed a pattern similar to the wild-type strain (lanes 1 & 2); Pro-σ^K

cleavage increased between 4 and 5 h, and the levels of SpoIVFA and SpoIVFB decreased. The

levels of all three proteins were slightly lower in the strain producing the T64A variant than in

the wild-type strain. The level of the T64A variant was similar to the N48A and N61A variants

at 4 h, yet the Pro-σ^K cleavage ratio was lower (lanes 9, 11 & 13), suggesting that the T64A

variant inhibits SpoIVFB, although not as well as GFPΔ27BofA (lane 7). We conclude that the

three conserved residues of BofA are important for SpoIVFB inhibition during sporulation.

Since GFPΔ27BofA co-localizes with SpoIVFA and SpoIVFB to the outer FS membrane

during sporulation (28-30), we examined the ability of the GFPΔ27BofA variants to localize to

the FS (Fig. 3B). As a control, GFPΔ27BofA produced in the *spoIVB165 bofA::erm* double

mutant localized discretely to the FS at 3 h PS. Strains producing the N48A and T64A variants

showed GFP signal localized to the FS, but signal was also observed in the cytoplasm and

possibly at the membrane of the MC, suggesting partial mislocalization. The strain producing

the N61A variant appeared to have GFP signal dispersed evenly throughout the MC cytoplasm,

suggesting a loss of FS localization. The fusion proteins were intact, suggesting the cytoplasmic

signal was not attributable to breakdown (*SI Appendix*, Fig. S6). Inability of the N61A variant to

localize to the FS may explain the loss of SpoIVFB inhibition and the abundant cleavage of Pro-

σ^K at 4 h (Fig. 3A, lane 11). Importantly, the similar ability of the N48A and T64A variants to

localize to the FS (Fig. 3B) does not account for their differential effects on the level of

SpoIVFB and its ability to cleave Pro-σ^K (Fig. 3A). The strain producing GFPΔ27BofA N48A

exhibited less SpoIVFB yet more Pro-σ^K cleavage at 4 h (lane 9) than the strain producing

GFPΔ27BofA T64A (lane 13), so the N48A substitution more severely impairs the ability of

GFPΔ27BofA to inhibit SpoIVFB.

BofA TMS2 Can Occupy the Active Site Region of SpoIVFB. N48 within TMS2 is conserved

among BofA orthologs (*SI Appendix*, Fig. S4) and potentially in proximity to the active site of

SpoIVFB. The SpoIVFB catalytic core formed by a zinc ion near E44 of the HELGH

metalloprotease motif within TMS2 is predicted to be surrounded by five other TMSs in the

membrane (Fig. 1A and *SI Appendix*, Fig. S7A). The model of the SpoIVFB membrane domain

is based on the crystal structure of an archaeal homolog (6). The model is supported by cross-

linking studies of catalytically-inactive SpoIVFB E44Q in complex with Pro-σ^K(1-127), whose

Proregion is predicted to occupy the SpoIVFB active site region (45, 46, 52). If BofA TMS2

instead occupied the SpoIVFB active site region, it would likely inhibit Pro-σ^K cleavage.

To determine whether BofA TMS2 can occupy the SpoIVFB active site region, we

performed disulfide cross-linking experiments that took advantage of the sole Cys residue of

BofA, C46, located near N48 in TMS2. We created MBPΔ27BofA because MBP (unlike GFP)

contains no Cys residues. As a negative control, we replaced C46 with Ser to obtain Cys-less

MBPΔ27BofA C46S. We also created Cys-less SpoIVFA C77L C82L, which cannot form

disulfide cross-links. Cys-less SpoIVFA and either MBPΔ27BofA or Cys-less MBPΔ27BofA

were functional when produced from a pET Quartet plasmid, since Pro-σ^K(1-127) cleavage by

cytTM-SpoIVFB was inhibited in *E. coli* (*SI Appendix*, Fig. S8). For the disulfide cross-linking

experiments, we used Cys-less Pro-σ^K(1-127) C110S and single-Cys variants of cytTM-

SpoIVFB created previously (45).

To test whether BofA TMS2 C46 could be cross-linked to a Cys located at the SpoIVFB

active site, MBPΔ27BofA C46 and single-Cys E44C cytTM-SpoIVFB were co-produced with

Cys-less variants of SpoIVFA and Pro-σ^K(1-127) from a pET Quartet plasmid in *E. coli*. Cells

were treated with the oxidant Cu^{2+}(phenanthroline)$_3$ to promote disulfide bond formation. For

MBPΔ27BofA C46 (Fig. 4A, lane 2), but not the Cys-less C46S negative control (lane 5),

treatment with oxidant caused formation of a species of the expected size for a cross-linked

complex with single-Cys E44C cytTM-SpoIVFB, which was detected by immunoblotting with

anti-FLAG antibodies. Treatment with the reducing agent DTT greatly diminished the

abundance of the apparent complex, consistent with cross-link reversal (lane 3). A species of the

expected size for a cross-linked dimer of single-Cys E44C cytTM-SpoIVFB was also observed.

Formation of the apparent dimer varies (see below), as reported previously (45). As expected,

anti-MBP antibodies detected the presumptive cross-linked complex of MBPΔ27BofA C46 with

single-Cys E44C cytTM-SpoIVFB, albeit weakly, and the negative control with E44Q rather

than E44C failed to form the complex (*SI Appendix*, Fig. S9, lanes 2 & 5). Since the signal for

the complex was stronger with anti-FLAG antibodies (Fig. 4A), we used those antibodies in the

cross-linking experiments reported below.

In addition to E44 of SpoIVFB, V70 in a predicted membrane-reentrant loop and P135 in

a predicted short loop interrupting TMS4 (Fig. 1A and *SI Appendix*, Fig. S7A) were shown to be

in proximity to the Proregion of Pro-σ^K(1-127) (45). Therefore, we tested whether

MBPΔ27BofA C46 could be cross-linked to single-Cys V70C or P135C cytTM-SpoIVFB E44Q

variants. We included the inactivating E44Q substitution since the V70C and P135C variants

(unlike the E44C variant) could cleave Cys-less Pro-σ^K(1-127) (45), even though Cys-less

SpoIVFA and either MBPΔ27BofA or Cys-less MBPΔ27BofA were expected to almost

completely inhibit cleavage (*SI Appendix*, Fig. S8). MBPΔ27BofA C46 formed a complex with

the P135C variant, but not with the V70C variant (Fig. 4B, lanes 2 & 8). As expected, Cys-less

MBPΔ27BofA C46S failed to form a complex with either variant (lanes 5 & 11). Full-length

BofA C46 also formed a complex of the expected (smaller) size with the E44C and P135C

variants, but not with the V70C variant (*SI Appendix*, Fig. S10, lanes 2, 14, & 8). As expected,

Cys-less BofA C46S failed to form a complex with any of the variants (lanes 5, 11, & 17). Our

cross-linking results show that BofA TMS2 can occupy the active site region of SpoIVFB,

placing BofA C46 in proximity to SpoIVFB E44 and P135, but not V70.

Based on our cross-linking results, we modeled BofA TMS2 in the SpoIVFB active site

region, and tested predictions of the model using additional disulfide cross-linking experiments

(*SI Appendix*, Results and Fig. S11). The results confirmed predictions of the initial model and

suggested a preferred orientation of BofA TMS2 in the SpoIVFB active site region, which led to

a refined model (*SI Appendix*, Fig. S7B).

BofA and SpoIVFA Do Not Prevent Pro-σ^K(1-127) from Interacting with SpoIVFB. Since

our cross-linking results show that BofA TMS2 can occupy the SpoIVFB active site region, we

tested whether BofA and SpoIVFA prevent Pro-σ^K(1-127) from interacting with SpoIVFB in *E.*

coli. Using a catalytically-inactive E44C cytTM-SpoIVFB variant containing the FLAG$_2$ epitope

tag but lacking His$_6$, co-produced Pro-σ^K(1-127), SpoIVFA, and GFPΔ27BofA co-purified with

the cytTM-SpoIVFB variant in pull-down assays with anti-FLAG antibody beads (Fig. 5). All

four proteins were seen in the bound sample (lane 4), indicating that GFPΔ27BofA and

SpoIVFA did not completely prevent Pro-σ^K(1-127) from interacting with the cytTM-SpoIVFB

variant. However, only the cytTM-SpoIVFB variant was detected when the bound sample was

diluted tenfold (lane 3) to match the input sample concentration (lane 1), and portions of the

other three proteins were observed in the unbound sample (lane 2). The results indicate

inefficient co-purification and/or recovery from the beads. A negative control with the cytTM-

SpoIVFB variant lacking the FLAG$_2$ epitope tag showed no Pro-σ^K(1-127) in the bound sample,

but small amounts of GFPΔ27BofA and SpoIVFA were detected (lane 8), indicative of weak,

nonspecific binding to the beads.

We also performed pull-down assays with cobalt resin, which binds to the His$_6$ tag on

Pro-σ^K(1-127), and we performed both types of pull-down assays (i.e., anti-FLAG antibody

beads and cobalt resin) when full-length BofA rather than GFPΔ27BofA was co-produced with

the other three proteins (*SI Appendix*, Results and Fig. S12). Neither BofA nor GFPΔ27BofA

when co-produced with SpoIVFA prevented Pro-σ^K(1-127) from interacting with the cytTM-

SpoIVFB variant. However, GFPΔ27BofA and SpoIVFA reduced co-purification of full-length

Pro-σ^K-His$_6$ with the cytTM-SpoIVFB variant in both types of pull-down assays (*SI Appendix*,

Results and Fig. S13), as compared with Pro-σ^K(1-127) (Fig. 5 and *SI Appendix*, Fig. S12A),

suggesting that the C-terminal half of Pro-σ^K affects complex formation (*SI Appendix*,

Discussion). As expected, GFPΔ27BofA and SpoIVFA strongly inhibited cytTM-SpoIVFB

cleavage of full-length Pro-σ^K-His$_6$ (*SI Appendix*, Fig. S14), similar to the inhibition of Pro-σ^K(1-

127) cleavage we observed (Fig. 1B, lane 4). To further investigate the strong inhibition of

cleavage despite the presence of substrate in the pulled-down protein complexes, we next

examined the effects of full-length BofA or MBPΔ27BofA co-produced with SpoIVFA on the

interaction of Pro-σ^K(1-127) with cytTM-SpoIVFB using disulfide cross-linking approaches.

Full-length BofA and SpoIVFA Interfere with Cross-Linking between the Proregion of

Pro-σ^K(1-127) and the SpoIVFB Active Site Region, and MBPΔ27BofA in Combination

with SpoIVFA Is Less Effective at Interfering with Some of the Cross-links. SpoIVFB

cleaves Pro-σ^K (43) and Pro-σ^K(1-127) (31) between residues S21 and Y22. In disulfide cross-

linking experiments, Cys substitutions for several residues near the cleavage site of otherwise

Cys-less Pro-σ^K(1-127) formed a cross-linked complex with single-Cys (E44C, V70C, P135C)

cytTM-SpoIVFB variants (45). The complex was most abundant with the E44C and V70C

variants, so we compared these interactions in the presence or absence of Cys-less inhibitory

proteins.

To examine proximity between single-Cys E44C cytTM-SpoIVFB and the Proregion, we

first co-produced single-Cys (F18C, V20C, S21C, K24C) Pro-σ^K(1-127) variants in the absence

of inhibitory proteins and measured time-dependent cross-linking. We quantified the cross-

linking by measuring the abundance of the complex and dividing that by the total amount of the

cytTM-SpoIVFB variant monomer, dimer, and complex. The ratio over time was plotted with a

best-fit trend line. Single-Cys E44C cytTM-SpoIVFB formed abundant complex with single-

Cys V20C and K24C Pro-σ^K(1-127) variants, and the amount of complex increased over time,

but with the F18C and S21C variants much less complex formed, only slightly more than with

the Cys-less Pro-σ^K(1-127) negative control (Fig. 6A and *SI Appendix*, Fig. S15A). Figure 6B

shows a representative immunoblot (60-min oxidant treatment).

Co-production of Cys-less inhibitory proteins greatly reduced complex formation

between single-Cys E44C cytTM-SpoIVFB and the V20C and K24C Pro-σ^K(1-127) variants.

When Cys-less MBPΔ27BofA and SpoIVFA were co-produced, much less complex formed and

its abundance did not increase over time (Fig. 6C and *SI Appendix*, Fig. S15B). The decrease in

abundance of the complex is most evident comparing representative immunoblots (60-min

oxidant treatment) with (Fig. 6D) or without (Fig. 6B) co-produced inhibitory proteins. When

Cys-less full-length BofA and SpoIVFA were co-produced, similar decreases in abundance of

the complex were observed (*SI Appendix*, Fig. S16). Figure 6E shows a comparison of the cross-

linking time courses with and without inhibitory proteins for single-Cys E44C cytTM-SpoIVFB

with single-Cys V20C Pro-σ^K(1-127). The effects of Cys-less MBPΔ27BofA and Cys-less full-

length BofA were indistinguishable. Similar results were observed with single-Cys K24C Pro-

σ^K(1-127) (compare Fig. 6C and *SI Appendix*, Fig. S16B). We conclude that MBPΔ27BofA

(lacking TMS1) is as effective as full-length BofA at interfering with cross-linking between the

Proregion of Pro-σ^K and single-Cys E44C cytTM-SpoIVFB. These results likely explain why

BofA TMS1 is dispensable for most of the inhibitory function of BofA (30, 31) (Figs. 1B and

3A). SpoIVFB E44 normally activates a water molecule to hydrolyze the bond between S21 and

Y22 of Pro-σ^K. Interfering with this interaction is expected to inhibit cleavage.

We next examined proximity between V70 of SpoIVFB and the Proregion of Pro-σ^K(1-

127). V70 is located in a predicted long and potentially flexible membrane-reentrant loop in the

SpoIVFB active site region (Fig. 1A and *SI Appendix*, Fig. S7A), perhaps explaining why single-

Cys V70C cytTM-SpoIVFB E44Q formed abundant cross-linked complex with both the F18C

and K24C Pro-σ^K(1-127) variants in the absence of inhibitory proteins (45) (Fig. 6 F and G; *SI*

Appendix, Fig. S17 A and B). With co-production of Cys-less variants of MBPΔ27BofA and

SpoIVFA, a lower relative amount of complex was measured, which did not increase over time

(Fig. 6 F and G; *SI Appendix*, Fig. S17 C and D), but the impact on complex formation was not

as great as when Cys-less variants of full-length BofA and SpoIVFA were co-produced (Fig. 6 F

and G; *SI Appendix*, Fig. S17 E and F). Full-length BofA and SpoIVFA also caused four novel

species to be observed, including with the Cys-less Pro-σ^K(1-127) negative control (*SI Appendix*,

Fig. S17G), perhaps due to cross-linking of the V70C cytTM-SpoIVFB variant to *E. coli* proteins

(i.e., BofA may cause the SpoIVFB membrane-reentrant loop to be exposed). In any case, full-

length BofA hindered cross-linking more than MBPΔ27BofA (lacking TMS1) (Fig. 6 F and G),

suggesting that both TMSs of BofA interfere with the normal interaction between the membrane-

reentrant loop of SpoIVFB and the Proregion of Pro-σ^K.

Since full-length BofA also inhibited cleavage of Pro-σ^K(1-127) in *E. coli* (Fig. 1B) and

Pro-σ^K in *B. subtilis* (Fig. 3A) slightly more than GFPΔ27BofA (lacking TMS1), we compared

time-dependent cross-linking of BofA and MBPΔ27BofA to single-Cys cytTM-SpoIVFB

variants. The results suggest that full-length BofA C46 forms slightly more complex with

single-Cys E44C cytTM-SpoIVFB and single-Cys P135C cytTM-SpoIVFB E44Q over time,

whereas the abundance of complex did not increase over time in the case of MBPΔ27BofA C46

(*SI Appendix*, Fig. S18). Hence, BofA TMS1 may slightly enhance TMS2 occupancy in the

active site region of SpoIVFB.

BofA and SpoIVFA Interfere with Cross-Linking between Pro-σ^K(1-127) and the SpoIVFB

Interdomain Linker, but Not with Cross-Linking between Pro-σ^K(1-127) and the SpoIVFB

CBS Domain. To further examine the effects of the inhibitory proteins on the interaction of Pro-

σ^K(1-127) with SpoIVFB, we identified additional residues that formed disulfide cross-linked

complexes. A model of catalytically-inactive SpoIVFB E44Q in complex with Pro-σ^K(1-127)

(46) predicts interactions between the C-terminal part of SpoIVFB and parts of Pro-σ^K(1-127).

The C-terminal part of SpoIVFB includes a 26-residue interdomain linker and a CBS domain

(Fig. 1A and *SI Appendix*, Fig. S7A). For both the linker and the CBS domain, we used the

model to predict residues of SpoIVFB in proximity to residues of Pro-σ^K(1-127). We tested the

predictions by co-producing single-Cys variants of the two proteins and performing disulfide

cross-linking experiments. We discovered that a single-Cys Y214C linker variant of cytTM-

SpoIVFB E44Q formed abundant complex with single-Cys L41C Pro-σ^K(1-127), and a single-

Cys A231C CBS domain variant of cytTM-SpoIVFB E44Q formed complex with single-Cys

A97C Pro-σ^K(1-127) (*SI Appendix*, Fig. S19 A and B). We showed that the Cys substitutions do

not impair cytTM-SpoIVFB activity or Pro-σ^K(1-127) susceptibility to cleavage (*SI Appendix*,

Fig. S19C). Finally, we measured time-dependent cross-linking in the presence or absence of

Cys-less inhibitory proteins.

Interestingly, formation of cross-linked complex between the Y214C linker variant of

cytTM-SpoIVFB and single-Cys L41C Pro-σ^K(1-127) was hindered more by Cys-less full-length

BofA than by Cys-less MBPΔ27BofA, when co-produced with Cys-less SpoIVFA (Fig. 6H and

SI Appendix, Fig. S20A). The pattern is similar to that observed for the V70C membrane-

reentrant loop variant of cytTM-SpoIVFB and single-Cys F18C or K24C Pro-σ^K(1-127) (Fig. 6 F

and G). The similar pattern suggests that in both cases BofA TMS2 partially interferes with the

interaction and BofA TMS1 augments the interference.

Strikingly, co-production of Cys-less inhibitory proteins had little or no effect on

formation of cross-linked complex between the A231C CBS domain variant of cytTM-SpoIVFB

and single-Cys A97C Pro-σ^K(1-127) (Fig. 6I and *SI Appendix*, Fig. S20B). These results suggest

that neither full-length BofA nor MBPΔ27BofA, when co-produced with SpoIVFA, prevent Pro-

σ^K(1-127) from interacting with the CBS domain of SpoIVFB in *E. coli*, consistent with our

finding that inhibitory proteins did not completely prevent co-purification of SpoIVFB and Pro-

σ^K(1-127) (Fig. 5 and *SI Appendix*, Fig. S12).

A Model of SpoIVFB in Complex with BofA and Parts of SpoIVFA and Pro-σ^K.

Computational models were generated using a similar protocol as described previously (46), but

including additional constraints reflecting experimental cross-linking data reported herein, as

well as newly predicted intra- and inter-chain contacts based on co-evolutionary couplings. Two

final models were generated: 1) Full-length SpoIVFB modeled as a tetramer, with part of one

Pro-σ^K molecule (residues 1-114), referred to as 'fb.sigk'; 2) Full-length SpoIVFB, again

modeled as a tetramer, with one molecule each of full-length BofA and parts of Pro-σ^K (residues

38-114) and SpoIVFA (residues 65-111), referred to as 'fb.sigk.bofa.fa'. The omitted residues of

Pro-σ^K and SpoIVFA could not be placed with sufficient confidence. The first model is

described in *SI Appendix*, Methods and Fig. S21.

Relevant features of the second model are illustrated in Figure 7 (only one molecule of

SpoIVFB is shown). The first side view shows SpoIVFB with BofA TMS2 and the C-terminal

part of Pro-σ^K(1-127) (Fig. 7A). The membrane domain of SpoIVFB (green) interacts with

BofA TMS2 (cyan) while the interdomain linker and CBS domain of SpoIVFB interact with the

modeled portion of Pro-σ^K(1-127) (red). The enlarged view of the SpoIVFB active site region

shows BofA TMS2 surrounded by SpoIVFB (TMSs labeled 1-6). The top view of the membrane

domain emphasizes proximity between BofA TMS2, SpoIVFB TMSs 2-4, and the zinc ion

(magenta) involved in catalysis. The second side view shows SpoIVFB with full-length BofA

and the C-terminal part of Pro-σ^K(1-127) (Fig. 7B). Our model predicts that BofA contains 2

TMSs and a membrane-embedded C-terminal region (labeled C near the C-terminus in the

enlarged view of the SpoIVFB active site region) that forms two short α-helices connected by a

turn. The enlarged and top views show that BofA interacts extensively with SpoIVFB and

occupies its active site region, which would sterically hinder access of the Proregion of Pro-σ^K.

Figure 7C shows the addition of the modeled portion of SpoIVFA (purple), which is predicted by

co-evolutionary couplings to contact the BofA C-terminal region and SpoIVFB TMS4. Hence,

the model predicts that SpoIVFA stabilizes the interaction of BofA with SpoIVFB. Since

GFPΔ27BofA has the C-terminal region predicted to interact with SpoIVFA, the interaction of

GFPΔ27BofA with SpoIVFB is likewise predicted to be stabilized by SpoIVFA, which may

explain the dependence of SpoIVFB inhibition on both GFPΔ27BofA and SpoIVFA (Fig. 1C),

and why GFPΔ27BofA inhibits substrate cleavage (Fig. 1B and 3A) and MBPΔ27BofA occupies

the SpoIVFB active site region (*SI Appendix*, Fig. S18) nearly as well as full-length BofA.

Discussion

Our results provide evidence that BofA occupies the active site region of SpoIVFB. BofA

residue C46 near the middle of TMS2 formed disulfide cross-links with SpoIVFB variants

containing a single-Cys at (E44C) or near (P135C) its active site. BofA variants containing a

single-Cys near the ends of TMS2 formed cross-links with single-Cys SpoIVFB variants that

suggest BofA TMS2 adopts a preferred orientation in the SpoIVFB active site region. N48

within BofA TMS2 and both N61 and T64 near its C-terminal end are highly conserved among

BofA orthologs. We showed that these residues are important for SpoIVFB inhibition during *B.*

subtilis sporulation and upon heterologous expression in *E. coli*. Structural modeling based on

partial homology and constraints from cross-linking and co-evolutionary analyses predicts that

BofA N48 and T64 interact, and that nearby N61 may also interact to stabilize the BofA

structure. SpoIVFA is also important for SpoIVFB inhibition and the model predicts that

SpoIVFA bridges the BofA C-terminal region and SpoIVFB to form a membrane-embedded

inhibitory complex (Fig. 7C). We found that BofA and SpoIVFA interfere with cross-linking

between the Pro-σ^K N-terminal region and the SpoIVFB active site region, supporting a steric

hindrance mechanism of SpoIVFB inhibition. BofA TMS1 enhanced interference with some

cross-links between Pro-σ^K and SpoIVFB, and slightly enhanced inhibition of SpoIVFB,

suggesting that BofA TMS1 enhances steric hindrance of substrate access to the enzyme active

site. Our findings have implications for the design of IMMP inhibitors.

Conserved Residues May Stabilize BofA in a Membrane-Embedded Inhibitory Complex

with SpoIVFA that Sterically Hinders Substrate Access to the SpoIVFB Active Site. BofA,

SpoIVFA, and SpoIVFB form a heterotrimeric membrane complex that improves accumulation

of each protein likely by inhibiting proteolytic degradation (30). Ala substitutions for N48, N61,

and T64 in GFPΔ27BofA reduced its level as well as the levels of SpoIVFA and SpoIVFB, both

upon heterologous expression in *E. coli* (Fig. 2) and during *B. subtilis* sporulation (Fig. 3A),

suggesting that assembly of the inhibitory complex was impaired. In agreement, the Ala

substitutions caused partial or complete mislocalization of GFPΔ27BofA during sporulation (Fig.

3B), which normally relies on SpoIVFA (30). Impaired assembly of the inhibitory complex can

explain the observed cleavage of Pro-σ^K(1-127) in *E. coli* (Fig. 2) and premature cleavage of

Pro-σ^K in sporulating *B. subtilis* (Fig. 3A).

Our structural model predicts proximity within BofA between the N48 side chain and a

loop that includes N61 and T64. In particular, the T64 side chain is oriented toward the N48 side

chain in the model, predicting an interaction. The N61 side chain points away from the N48 and

T64 side chains in the model, but the orientation of the N61 side chain is uncertain given the

predicted loop structure, so an interaction with the side chains of N48 and/or T64 remains

possible. Alternatively, the three conserved residues may contact other residues within BofA,

based on co-evolutionary couplings (*SI Appendix*, Dataset S4), to stabilize the BofA structure

and promote interactions with SpoIVFA and SpoIVFB. For example, disulfide cross-linking

supports proximity between BofA L62 and V63 (in the predicted loop that includes N61 and

T64) and SpoIVFB M30, which is located in a predicted short loop between TMS1 and TMS2

(*SI Appendix*, Fig. S11). Our model satisfies restraints based on this observed cross-linking, as

well as many other experimental cross-linking results (45, 46) (*SI Appendix*, Dataset S3). The

model also predicts numerous interactions between BofA, SpoIVFA, and SpoIVFB that remain

to be tested, which will be facilitated by approaches described herein. Of note, the model

predicts that BofA C-terminal residues 85 to 87 interact with SpoIVFA and SpoIVFB, and that

BofA residues 28 to 36 preceding predicted TMS2 interact with SpoIVFA (Fig. 7C), which may

explain the effects of deleting or changing those BofA residues, including loss of SpoIVFB

inhibition resulting in Pro-σ^K(1-127) cleavage and reduced levels of SpoIVFA and SpoIVFB in

E. coli (*SI Appendix*, Results and Fig. S5). The predicted interactions can be tested further by

disulfide cross-linking.

BofA and SpoIVFA interfere with cross-linking between the Pro-σ^K N-terminal region

and the SpoIVFB active site region, supporting a steric hindrance mechanism of SpoIVFB

inhibition. Many latent zymogens likewise use a prosegment or prodomain to sterically hinder

substrate access to their active site (53). Many metalloprotease zymogens in addition use a

ligand switch mechanism of inhibition and activation. Inhibition involves an amino acid side

chain of a residue in the prosegment acting as a metal ligand to prevent catalysis. Activation

requires replacement of the side chain metal ligand with a water molecule necessary for substrate

peptide bond hydrolysis. BofA residue C46 near the middle of TMS2 is well-positioned in our

model to act as a zinc ligand, but C46 is not conserved among BofA orthologs (*SI Appendix*, Fig.

S4). Therefore, we favor a simple steric hindrance mechanism of SpoIVFB inhibition. During

sporulation, SpoIVFB would be activated when SpoIVB cleaves the C-terminal end of SpoIVFA

(35-38) and CtpB cleaves the C-terminal ends of both SpoIVFA and BofA (36, 37, 39) (*SI*

Appendix, Fig. S1). These cleavages result in a conformational change in the inhibition complex

(36), which we propose allows the Proregion of Pro-σ^K to be positioned in the active site for

cleavage. After cleavage, the conformational change may also aid in release of the product, σ^K.

BofA TMS1 Appears to Enhance Inhibition of SpoIVFB Slightly by Partially Interfering

with Its Interaction with Pro-σ^K. Full-length BofA inhibited cleavage of Pro-σ^K(1-127) in *E.*

coli (Fig. 1B) and Pro-σ^K in *B. subtilis* (Fig. 3A) slightly more than GFPΔ27BofA. The most

likely explanation for the difference, based on our results, is that TMS1 of BofA sterically

hinders some interactions between the Pro-σ^K N-terminal region and SpoIVFB more than BofA

variants lacking TMS1 (such as GFPΔ27BofA and MBPΔ27BofA). The interactions hindered

more by full-length BofA included cross-links between the Pro-σ^K Proregion (F18C and K24C)

and the SpoIVFB membrane-reentrant loop (V70C) (Fig. 6F and G), which may bind to the

Proregion and present it to the SpoIVFB active site for cleavage (46) based on a study of the

homologous membrane-reentrant loop of *E. coli* RseP (54). Our model predicts that both TMSs

of BofA, as well as its C-terminal region, occupy the active site region of SpoIVFB (Fig. 7B),

which would sterically hinder interactions between the SpoIVFB membrane-reentrant loop and

the Pro-σ^K Proregion.

Full-length BofA also hindered cross-links between Pro-σ^K L41C and the SpoIVFB

interdomain linker (Y214C) more than MBPΔ27BofA (Fig. 6H). The interdomain linker has

been proposed to allow coordination between ATP binding by the CBS domain and Pro-σ^K

cleavage by the membrane domain (46). Thus, BofA TMS1 may interfere with coordination

between the domains of SpoIVFB by inhibiting movement of the linker, in addition to hindering

linker interaction with the Pro-σ^K N-terminal region.

Design of IMMP Inhibitors. Our findings reveal concepts that may inform efforts to design

selective inhibitors of IMMPs. BofA TMS2 appears to block the SpoIVFB active site (Fig. 7A),

similar to the prosegment of many latent zymogens (53). Like BofA, some prosegments can act

in *trans*, an observation that has led to the demonstration of prodomains as selective inhibitors of

A Disintegrin And Metalloproteinase (ADAM) family enzymes (55, 56). Selectivity relies on

the extensive interaction surface between the prodomain and the enzyme, including features

specific to the pair, which can boost efficacy and diminish off-target effects in therapeutic

strategies (57). A peptide encompassing residues 48 to 64 of BofA may inhibit SpoIVFB, but

our model also predicts extensive interactions between the BofA C-terminal region (residues 65

to 87) and SpoIVFB (Fig. 7B), so a longer BofA peptide, perhaps flexibly linked to SpoIVFA

residues 96 to 109, which are also predicted to interact with SpoIVFB (Fig. 7C), may exhibit

improved inhibition and selectivity. Since cleavage of Pro-σ^K(1-127) by cytTM-SpoIVFB has

been reconstituted *in vitro* (42), it will be possible to test peptides for inhibition. Structure

determination of SpoIVFB in complex with an inhibitory peptide, GFPΔ27BofA, or full-length

BofA is an important goal to facilitate design of IMMP inhibitors. In particular, it may inform

efforts to design inhibitors of SpoIVFB orthologs in *Bacillus* and *Clostridium* species, many of

which cause disease or provide benefits, and persist by forming endospores (58). Such efforts

could lead to new strategies to control endosporulation.

Our findings also suggest that BofA TMS1 interferes with interactions between the Pro-

σ^K N-terminal region and SpoIVFB. Therefore, a cyclic peptide inhibitor may prove to be more

effective than a linear one. The desirable characteristics of cyclic peptides as therapeutics, and

new methods of producing and screening cyclic peptide libraries, make this an attractive

possibility (59, 60).

The inhibitor design concepts discussed above could apply equally well to IMMPs that

normally use one or more PDZ domains to exclude substrate from their active site region. *E. coli*

RseP appears to use two PDZ domains to block access of substrate RseA (61-63). Cleavage of

the RseA periplasmic domain by DegS (referred to as site-1 cleavage) (64-66) is proposed to

allow entry of the RseA TMS into the RseP active site region for site-2 cleavage (67). Recently,

batimastat, a hydroxamic acid derivative that mimics the peptide structure of natural substrates,

was shown to selectively inhibit RseP *in vivo* (68). Two other peptidomimetic hydroxamic acids,

ilomastat and miramastat, exhibited very little RseP inhibition. A lack of selectivity has been a

problem in efforts toward using peptidomimetic hydroxamic acids as matrix metalloprotease

inhibitors for treatment of cancer, arthritis, and other diseases (57, 69, 70). The hydroxamic acid

group of these compounds strongly coordinates the catalytic zinc ion, and the peptidomimetic

portions have not provided enough specificity to prevent off-target effects. Even so, the side

effects may not preclude using hydroxamic acid-based peptidomimetics to treat bacterial

infections topically, and systemically if administered briefly. Perhaps weaker zinc ligands based

on amino acid side chains would reduce side effects. A cyclic peptidomimetic with a zinc ligand

could presumably block access of the RseA TMS to the RseP active site, or substrate access to

the active site of other IMMPs. Other small-molecule matrix metalloprotease inhibitors that

have already been developed (69) should be screened for activity against bacteria and fungi with

IMMPs known to be involved in pathogenesis (16-19).

Methods

Plasmids, Primers, and Strains. Plasmids used in this study are described in *SI Appendix*,

Table S1. Primers used in plasmid construction are listed in *SI Appendix*, Table S2. Plasmids

were cloned in *E. coli* strain DH5α (71). Relevant parts of plasmids were verified by DNA

sequencing with primers listed in *SI Appendix*, Table S3. *B. subtilis* strains used in this study are

described in *SI Appendix*, Table S4.

Pro-σ$^{\mathrm{K}}$(1-127) Cleavage in *E. coli*. Strain BL21(DE3) (Novagen) was used to produce proteins

in *E. coli*. Two plasmids with different antibiotic resistance genes were cotransformed (31) or a

single plasmid was transformed, with selection on Luria-Bertani (LB) agar supplemented with

kanamycin sulfate (50 μg/mL) and/or ampicillin (100 μg/mL). Transformants (4-5 colonies)

were grown in LB medium with 50 μg/mL kanamycin sulfate and/or 200 μg/mL ampicillin at

37°C with shaking (200 rpm). Typically, overnight culture (200 μL) was transferred to 10 mL of

LB medium with antibiotics, cultures were grown at 37°C with shaking (250 rpm) to an optical

density of 60-80 Klett units, and isopropyl β-D-thiogalactopyranoside (IPTG) (0.5 mM) was

added to induce protein production for 2 h. For transformants with either pET Quintet or full-

length Pro-σ^K-His$_6$, overnight growth was avoided. Transformants were transferred directly to

10 mL of LB medium with antibiotic, and cultures were grown and induced as described above.

Equivalent amounts of cells (based on optical density in Klett units) were collected (12,000 $\times$ g

for 1 min) and extracts were prepared (31), then subjected to immunoblot analysis (*SI Appendix*,

Methods).

BofA Sequence Analysis. Orthologs of *B. subtilis bofA*, which are present in the genomes of

most endospore-forming bacteria (20), were collected from the NCBI and Uniprot databases.

The protein sequences of BofA orthologs were aligned using the T-Coffee multiple sequence

alignment package (72). Residues identical in at least 70% of the sequences were considered

conserved.

B. subtilis **Sporulation and GFPΔBofA Localization.** GFPΔ27BofA or its variants were

expressed under control of the *bofA* promoter (P*$_{bofA}$*) after ectopic chromosomal integration.

Plasmids bearing P*$_{bofA}$*-*gfpΔ27bofA* or its variant, bordered by regions of homology to *B. subtilis*

amyE, were transformed into strain ZR264. Transformants with a gene replacement at *amyE*

were selected on LB agar with spectinomycin sulfate (100 μg/mL) and identified by loss of

amylase activity (73). Sporulation was induced by growing cells in the absence of antibiotics

followed by the resuspension of cells in SM medium (73). At indicated times PS, samples (50

μL) were centrifuged (12,000 $\times$ g for 1 min), supernatants were removed, and cell pellets were

stored at -80°C. Whole-cell extracts were prepared as described for *E. coli* (31), except samples

were incubated at 50°C for 3 min instead of boiling for 3 min (46), and proteins were subjected

to immunoblot analysis (*SI Appendix*, Methods).

To image GFPΔ27BofA localization, samples collected at 3 h PS were examined by

fluorescence microscopy using an Olympus FluoView FV-1000 filter-based confocal

microscope. GFPΔ27BofA (ex/em ~488/507 nm) was excited using a 458 nM argon laser and

fluorescence was captured using a BA465-495 nm band pass filter. The lipophilic dye FM 4-64

(1 μg/mL) (AAT Bioquest) was used to stain membranes. FM 4-64 (ex/em ~515/640 nm) was

excited using a 515 nm argon laser and fluorescence was captured using a BA560IF band pass

filter (74).

Disulfide Cross-Linking. A method described previously (75) was used with slight

modifications (45). Briefly, as described above for Pro-σ^K(1-127) cleavage, *E. coli* BL21(DE3)

was transformed with a plasmid, grown in LB (10 mL), induced with IPTG, and equivalent

amounts of cells were collected. Cells were subjected to disulfide cross-linking, then lysed and

proteins were precipitated by addition of trichloroacetic acid (5%), and proteins were subjected

to immunoblot analysis (*SI Appendix*, Methods).

Co-Immunoprecipitation (FLAG₂ Pull-Down Assays). *E. coli* BL21(DE3) was transformed

with a plasmid, grown in LB (1 L), and induced with IPTG as described above. The culture was

split, cells were harvested, and cell pellets were stored at -80°C. Cell lysates were prepared as

described (52), except that each cell pellet was resuspended in 20 mL of lysis buffer containing

50 mM Tris-HCl pH 7.1 rather than PBS. Cell lysates were centrifuged (15,000 × g for 15 min

at 4 °C) to sediment cell debris and protein inclusion bodies. The supernatant was treated with

1% *n*-dodecyl-β-D-maltoside (DDM) (Anatrace) for 1 h at 4 °C to solubilize membrane proteins,

then centrifuged at 150,000 × g for 1 h at 4 °C. The supernatant was designated the input sample

and 1 mL was mixed with 50 uL anti-DYKDDDDK magnetic agarose (Pierce) that had been

equilibrated with buffer (50 mM Tris-HCl pH 7.1, 0.1% DDM, 5 mM 2-mercaptoethanol, 10%

glycerol) and the mixture was rotated for 1 h at 25°C. The magnetic agarose was removed with a

DynaMag-2 magnet (Invitrogen) and the supernatant was saved (unbound sample). The

magnetic agarose was washed three times by gently vortexing with 500 μL wash buffer (50 mM

Tris-HCl pH 7.1, 150 mM NaCl, 10% glycerol, 0.1% DDM), then washed once with 500 μL

water. The magnetic agarose was mixed with 50 μL of 2× sample buffer (50 mM Tris·HCl pH

6.8, 4% SDS, 20% glycerol, 200 mM DTT 0.03% bromophenol blue) and boiled for 3 min

(bound sample). A portion of the bound sample was diluted tenfold (1/10 bound sample) with

1× sample buffer to match the concentration of the input sample. Samples were subjected to

immunoblot analysis (*SI Appendix*, Methods).

Colbalt Affinity Purification (His$_6$ Pull-Down Assays). Input sample (15 mL) prepared as

described above was mixed with imidazole (5 mM) and 0.5 mL of Talon superflow metal affinity

resin (Clontech) that had been equilibrated with buffer (as above for magnetic agarose). The

mixture was rotated for 1 h at 4°C. The cobalt resin was sedimented by centrifugation at $708 \times g$

for 2 min at 4 °C and the supernatant was saved (unbound sample). The resin was washed three

times with 5 mL wash buffer (as above plus 5 mM imidazole), each time rotating the mixture for

10 min at 4 °C and sedimenting resin as above. The resin was mixed with 0.5 mL 2× sample

buffer and boiled for 3 min (bound sample). A portion of the bound sample was diluted

fifteenfold (1/15 bound sample) with 1× sample buffer to match the concentration of the input

sample. Samples were subjected to immunoblot analysis (*SI Appendix*, Methods).

Modeling of Complexes Containing SpoIVFB, BofA, and Parts of SpoIVFA and Pro-σ^K.

The modeling proceeded through stages where initial monomeric models were assembled step-

by-step into multimeric complexes guided primarily by the restraints from cross-linking

experiments and predicted contacts from co-evolutionary coupling analysis. More specifically, a

SpoIVFB monomer was first assembled from the membrane and CBS domains. Two monomers

were combined into a dimer and the dimer was assembled into a plausible tetramer. Part of one

molecule of Pro-σ^K (residues 1-114) was subsequently added, which resulted in 'fb.sigk'. The

'fb.sigk.bofa.fa' model was developed by starting from 'fb.sigk', truncating the Pro-σ^K N-

terminus (residues 1-37), and subsequently adding first BofA and finally SpoIVFA (residues 65-

111). See *SI Appendix*, Methods for additional description of the modeling.

ACKNOWLEDGMENTS. We thank Daniel Parrell for assistance with fluorescence

microscopy, Jon Kaguni for helpful comments on the manuscript, and David Rudner, Simon

Cutting, Ruanbao Zhou, and Yang Zhang for plasmids. This research was supported by National

Institutes of Health Grants R01 GM43585 (to L.K.) and R35 GM126948 (to M.F.), and by

Michigan State University AgBioResearch.

References

1. Brown MS, Ye J, Rawson RB, Goldstein JL (2000) Regulated intramembrane proteolysis: a

control mechanism conserved from bacteria to humans. *Cell* 100:391-398.

2. Urban S (2013) Mechanisms and cellular functions of intramembrane proteases. *Biochim.*

Biophys. Acta - Biomembr. 1828:2797-2800.

3.   Manolaridis I, *et al.* (2013) Mechanism of farnesylated CAAX protein processing by the

intramembrane protease Rce1. *Nature* 504:301-305.

4.   Hu J, Xue Y, Lee S, Ha Y (2011) The crystal structure of GXGD membrane protease FlaK.

Nature 475:528-531.

5.   Li X, *et al.* (2013) Structure of a presenilin family intramembrane aspartate protease. *Nature*

493:56-61.

6.   Feng L, *et al.* (2007) Structure of a site-2 protease family intramembrane metalloprotease.

Science 318:1608-1612.

7.   Wang Y, Zhang Y, Ha Y (2006) Crystal structure of a rhomboid family intramembrane

protease. *Nature* 444:179-183.

8.   Wu Z, *et al.* (2006) Structural analysis of a rhomboid family intramembrane protease reveals

a gating mechanism for substrate entry. *Nat. Struct. Mol. Biol.* 13:1084-1091.

9.   Cho S, Dickey SW, Urban S (2016) Crystal structures and inhibition kinetics reveal a two-

stage catalytic mechanism with drug design implications for rhomboid proteolysis. *Mol. Cell*

61:329-340.

10. Zoll S, *et al.* (2014) Substrate binding and specificity of rhomboid intramembrane protease

revealed by substrate-peptide complex structures. *Embo J.* 33:2408-2421.

11. Yang G, *et al.* (2019) Structural basis of Notch recognition by human γ-secretase. *Nature*

565:192-197.

12. Zhou R, *et al.* (2019) Recognition of the amyloid precursor protein by human γ-secretase.

Science 363:eaaw0930.

13. Rawson RB (2013) The site-2 protease. *Biochim. Biophys. Acta - Biomembr.* 1828:2801-

2807.

14. Ye J (2013) Roles of regulated intramembrane proteolysis in virus infection and antiviral

immunity. *Biochim. Biophys. Acta - Biomembr.* 1828:2926-2932.

15. Adam Z (2013) Emerging roles for diverse intramembrane proteases in plant biology.

Biochim. Biophys. Acta - Biomembr. 1828:2933-2936.

16. Chang YC, Bien CM, Lee H, Espenshade PJ, Kwon-Chung KJ (2007) Sre1p, a regulator of

oxygen sensing and sterol homeostasis, is required for virulence in *Cryptococcus*

neoformans. *Mol. Microbiol.* 64:614-629.

17. Urban S (2009) Making the cut: central roles of intramembrane proteolysis in pathogenic

microorganisms. *Nat. Rev. Microbiol.* 7:411-423.

18. Kroos L, Akiyama Y (2013) Biochemical and structural insights into intramembrane

metalloprotease mechanisms. *Biochim. Biophys. Acta - Biomembr.* 1828:2873-2885.

19. Schneider JS, Glickman MS (2013) Function of site-2 proteases in bacteria and bacterial

pathogens. *Biochim. Biophys. Acta - Biomembr.* 1828:2808-2814.

20. Galperin MY*, et al.* (2012) Genomic determinants of sporulation in *Bacilli* and *Clostridia*:

towards the minimal set of sporulation-specific genes. *Environ. Microbiol.* 14:2870-2890.

21. McKenney PT, Driks A, Eichenberger P (2013) The *Bacillus subtilis* endospore: assembly

and functions of the multilayered coat. *Nat. Rev. Microbiol.* 11:33-44.

22. Al-Hinai MA, Jones SW, Papoutsakis ET (2015) The *Clostridium* sporulation programs:

diversity and preservation of endospore differentiation. *Microbiol. Mol. Biol. Rev.* 79:19-37.

23. Checinska A, Paszczynski A, Burbank M (2015) *Bacillus* and other spore-forming genera:

variations in responses and mechanisms for survival. *Annu. Rev. Food Sci. Technol.* 6:351-

369.

24. Cutting S*, et al.* (1990) A forespore checkpoint for mother-cell gene expression during

development in *Bacillus subtilis*. *Cell* 62:239-250.

25. Cutting S, Roels S, Losick R (1991) Sporulation operon *spoIVF* and the characterization of

mutations that uncouple mother-cell from forespore gene expression in *Bacillus subtilis*. *J.*

Mol. Biol. 221:1237-1256.

26. Ricca E, Cutting S, Losick R (1992) Characterization of *bofA*, a gene involved in

intercompartmental regulation of pro-σ^K processing during sporulation in *Bacillus subtilis*. *J.*

Bacteriol. 174:3177-3184.

27. Tan IS, Ramamurthi KS (2014) Spore formation in *Bacillus subtilis*. *Environ. Microbiol.*

Rep. 6:212-225.

28. Resnekov O, Alper S, Losick R (1996) Subcellular localization of proteins governing the

proteolytic activation of a developmental transcription factor in *Bacillus subtilis*. *Genes Cells*

1:529-542.

29. Rudner DZ, Pan Q, Losick RM (2002) Evidence that subcellular localization of a bacterial

membrane protein is achieved by diffusion and capture. *Proc Natl Acad Sci U S A* 99:8701-

8706.

30. Rudner DZ, Losick R (2002) A sporulation membrane protein tethers the pro-σ^K processing

enzyme to its inhibitor and dictates its subcellular localization. *Genes Dev.* 16:1007-1018.

31. Zhou R, Kroos L (2004) BofA protein inhibits intramembrane proteolysis of pro-σ^K in an

intercompartmental signaling pathway during *Bacillus subtilis* sporulation. *Proc. Natl. Acad.*

Sci. USA 101:6385-6390.

32. Kroos L, Yu YT, Mills D, Ferguson-Miller S (2002) Forespore signaling is necessary for

pro-σ^K processing during *Bacillus subtilis* sporulation despite the loss of SpoIVFA upon

translational arrest. *J. Bacteriol.* 184:5393-5401.

33. Cutting S, Driks A, Schmidt R, Kunkel B, Losick R (1991) Forespore-specific transcription

of a gene in the signal transduction pathway that governs pro-σ^K processing in *Bacillus*

subtilis. *Genes Dev.* 5:456-466.

34. Pan Q, Losick R, Rudner DZ (2003) A second PDZ-containing serine protease contributes to

activation of the sporulation transcription factor σ^K in *Bacillus subtilis*. *J. Bacteriol.*

185:6051-6056.

35. Dong TC, Cutting SM (2003) SpoIVB-mediated cleavage of SpoIVFA could provide the

intercellular signal to activate processing of Pro-σ^K in *Bacillus subtilis*. *Mol. Microbiol.*

49:1425-1434.

36. Campo N, Rudner DZ (2006) A branched pathway governing the activation of a

developmental transcription factor by regulated intramembrane proteolysis. *Mol. Cell* 23:25-

35.

37. Campo N, Rudner DZ (2007) SpoIVB and CtpB are both forespore signals in the activation

of the sporulation transcription factor σ^K in *Bacillus subtilis*. *J. Bacteriol.* 189:6021-6027.

38. Mastny M, *et al.* (2013) CtpB assembles a gated protease tunnel regulating cell-cell signaling

during spore formation in *Bacillus subtilis*. *Cell* 155:647-658.

39. Zhou R, Kroos L (2005) Serine proteases from two cell types target different components of

a complex that governs regulated intramembrane proteolysis of pro-σ^K during *Bacillus*

subtilis development. *Mol. Microbiol.* 58:835-846.

40. Rudner D, Fawcett P, Losick R (1999) A family of membrane-embedded metalloproteases

involved in regulated proteolysis of membrane-associated transcription factors. *Proc. Natl.*

Acad. Sci. USA 96:14765-14770.

41. Yu Y-TN, Kroos L (2000) Evidence that SpoIVFB is a novel type of membrane

metalloprotease governing intercompartmental communication during *Bacillus subtilis*

sporulation. *J. Bacteriol.* 182:3305-3309.

42. Zhou R, Cusumano C, Sui D, Garavito RM, Kroos L (2009) Intramembrane proteolytic

cleavage of a membrane-tethered transcription factor by a metalloprotease depends on ATP.

Proc. Natl. Acad. Sci. USA 106:16174-16179.

43. Kroos L, Kunkel B, Losick R (1989) Switch protein alters specificity of RNA polymerase

containing a compartment-specific sigma factor. *Science* 243:526-529.

44. Eichenberger P*, et al.* (2004) The program of gene transcription for a single differentiating

cell type during sporulation in *Bacillus subtilis*. *PLoS Biol.* 2:e328.

45. Zhang Y, Luethy PM, Zhou R, Kroos L (2013) Residues in conserved loops of

intramembrane metalloprotease SpoIVFB interact with residues near the cleavage site in Pro-

σ^K. *J. Bacteriol.* 195:4936-4946.

46. Halder S, Parrell D, Whitten D, Feig M, Kroos L (2017) Interaction of intramembrane

metalloprotease SpoIVFB with substrate Pro-σ^K. *Proc. Natl. Acad. Sci. USA* 114:E10677-

E10686.

47. Saribas AS, Gruenke L, Waskell L (2001) Overexpression and purification of the membrane-

bound cytochrome P450 2B4. *Protein Expr. Purif.* 21:303-309.

48. Prince H, Zhou R, Kroos L (2005) Substrate requirements for regulated intramembrane

proteolysis of *Bacillus subtilis* pro-σ^K. *J. Bacteriol.* 187:961-971.

49. Ramirez-Guadiana FH, *et al.* (2018) Evidence that regulation of intramembrane proteolysis is

mediated by substrate gating during sporulation in *Bacillus subtilis*. *PLoS Genet.*

14:e1007753.

50. de Hoon MJ, Eichenberger P, Vitkup D (2010) Hierarchical evolution of the bacterial

sporulation network. *Curr. Biol.* 20:R735-R745.

51. Varcamonti M, Marasco R, De Felice M, Sacco M (1997) Membrane topology analysis of

the *Bacillus subtilis* BofA protein involved in pro-σ^K processing. *Microbiol.* 143:1053-1058.

52. Zhang Y, *et al.* (2016) Complex formed between intramembrane metalloprotease SpoIVFB

and its substrate, Pro-σ^K. *J. Biol. Chem.* 291:10347-10362.

53. Arolas JL, Goulas T, Cuppari A, Gomis-Ruth FX (2018) Multiple architectures and

mechanisms of latency in metallopeptidase zymogens. *Chem. Rev.* 118:5581-5597.

54. Akiyama K, *et al.* (2015) Roles of the membrane-reentrant β-hairpin-like loop of RseP

protease in selective substrate cleavage. *eLife* 4:e08928

55. Moss ML, *et al.* (2007) The ADAM10 prodomain is a specific inhibitor of ADAM10

proteolytic activity and inhibits cellular shedding events. *J. Biol. Chem.* 282:35712-35721.

56. Wong E, *et al.* (2016) Harnessing the natural inhibitory domain to control TNFα Converting

Enzyme (TACE) activity *in vivo*. *Sci. Rep.* 6:35598.

57. Gomis-Ruth FX (2017) Third time lucky? Getting a grip on matrix metalloproteinases. *J.*

Biol. Chem. 292:17975-17976.

58. Driks A, Eichenberger P eds (2016) *The bacterial spore: from molecules to systems* (ASM

Press, Washington, DC), p 397.

59. Zorzi A, Deyle K, Heinis C (2017) Cyclic peptide therapeutics: past, present and future.

Curr. Opin. Chem. Biol. 38:24-29.

60. Sohrabi C, Foster A, Tavassoli A (2020) Methods for generating and screening libraries of

genetically encoded cyclic peptides in drug discovery. *Nat. Rev. Chem.* 4:90-101.

61. Grigorova IL*, et al.* (2004) Fine-tuning of the *Escherichia coli* σ^E envelope stress response

relies on multiple mechanisms to inhibit signal-independent proteolysis of the

transmembrane anti-sigma factor, RseA. *Genes Dev.* 18:2686-2697.

62. Inaba K*, et al.* (2008) A pair of circularly permutated PDZ domains control RseP, the S2P

family intramembrane protease of *Escherichia coli*. *J. Biol. Chem.* 283:35042-35052.

63. Kanehara K, Ito K, Akiyama Y (2003) YaeL proteolysis of RseA is controlled by the PDZ

domain of YaeL and a Gln-rich region of RseA. *EMBO J.* 22:6389-6398.

64. Ades SE, Connolly LE, Alba BM, Gross CA (1999) The *Escherichia coli* σ^E-dependent

extracytoplasmic stress response is controlled by the regulated proteolysis of an anti-sigma

factor. *Genes Dev.* 13:2449-2461.

65. Alba BM, Leeds JA, Onufryk C, Lu CZ, Gross CA (2002) DegS and YaeL participate

sequentially in the cleavage of RseA to activate the σ^E-dependent extracytoplasmic stress

response. *Genes Dev.* 16:2156-2168.

66. Kanehara K, Ito K, Akiyama Y (2002) YaeL (EcfE) activates the σ^E pathway of stress

response through a site-2 cleavage of anti-σ^E, RseA. *Genes Dev.* 16:2147-2155.

67. Hizukuri Y*, et al.* (2014) A structure-based model of substrate discrimination by a

noncanonical PDZ tandem in the intramembrane-cleaving protease RseP. *Structure* 22:326-

336.

68. Konovalova A*, et al.* (2018) Inhibitor of intramembrane protease RseP blocks the σ^E

response causing lethal accumulation of unfolded outer membrane proteins. *Proc. Natl.*

Acad. Sci. USA 115:E6614-E6621.

69. Jacobsen JA, Major Jourden JL, Miller MT, Cohen SM (2010) To bind zinc or not to bind

zinc: an examination of innovative approaches to improved metalloproteinase inhibition.

Biochim. Biophys. Acta 1803:72-94.

70. Fields GB (2015) New strategies for targeting matrix metalloproteinases. *Matrix Biol.* 44-

46:239-246.

71. Hanahan D (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.*

166:557-580.

72. Notredame C, Higgins DG, Heringa J (2000) T-Coffee: A novel method for fast and accurate

multiple sequence alignment. *J. Mol. Biol.* 302:205-217.

73. Harwood CR, Cutting SM (1990) *Molecular Biological Methods for Bacillus* (John Wiley &

Sons, Chichester, England) p 581.

74. Parrell D, Kroos L (2020) Channels modestly impact compartment-specific ATP levels

during *Bacillus subtilis* sporulation and a rise in the mother cell ATP level is not necessary

for Pro-σ^K cleavage. *Mol. Microbiol.* 114:563-581.

75. Koide K, Ito K, Akiyama Y (2008) Substrate recognition and binding by RseP, an

Escherichia coli intramembrane protease. *J. Biol. Chem.* 283:9562-9570.

 Figure Legends

Fig. 1. Inhibition of Pro-σ^K cleavage. (*A*) Diagram of SpoIVFB inhibition during *B. subtilis*

endosporulation and upon heterologous expression in *E. coli*. During endosporulation (*Left*),

SpoIVFB and its inhibitory proteins BofA and SpoIVFA are produced in the mother cell (MC)

and localize to the outer forespore (FS) membrane (OFM). Pro-σ^K is also produced in the MC

and associates with membranes. When these proteins are synthesized in *E. coli* (*Right*), they

localize to the inner membrane (IM). The expanded view of the membranes (*Center*) shows a

SpoIVFB variant with an extra N-terminal transmembrane segment (cytTM), and highlights

several residues (E44, V70, P135) at or near the active site in the membrane domain, which is

connected to the CBS domain by an interdomain linker. When produced in *E. coli*, cytTM-

SpoIVFB cleaves Pro-σ^K(1-127), removing its N-terminal Proregion. Co-production of

SpoIVFA and either full-length BofA or GFPΔ27BofA (lacking predicted TMS1) inhibits Pro-

σ^K(1-127) cleavage. (*B*) Cleavage assays comparing inhibition by SpoIVFA and either

GFPΔ27BofA or full-length BofA in *E. coli*. Pro-σ^K(1-127) and cytTM-SpoIVFB were

produced alone (lane 1, pYZ2) or in combination with GFPΔ27BofA and SpoIVFA (lane 2,

pYZ46) or full-length BofA and SpoIVFA (lane 3, pSO212). Alternatively, "pET Quartet"

plasmids were used to produce Pro-σ^K(1-127), cytTM-SpoIVFB, SpoIVFA, and either

GFPΔ27BofA (lane 4, pSO40) or full-length BofA (lane 5, pSO213). Samples collected after 2

h of IPTG induction were subjected to immunoblot analysis with SpoIVFA (*Top*), GFP (*Middle*),

or penta-His antibodies (*Bottom*, 2 and 30 s exposures). The single star (*) indicates cross-

reacting proteins below SpoIVFA and the double star (**) indicates breakdown species of

GFPΔ27BofA. A breakdown species below SpoIVFA (not indicated) is observed in some

experiments. The graph shows quantification of the cleavage ratio (cleavage product/[Pro-σ^K(1-

127) + cleavage product]) for three biological replicates. Error bars, 1 standard deviation. (*C*)

Cleavage assays comparing inhibition by either GFPΔ27BofA or SpoIVFA. pET Triplet

plasmids were used to produce Pro-σ^K(1-127), cytTM-SpoIVFB, and either GFPΔ27BofA (lane

2, pSO64) or SpoIVFA (lane 3, pSO65). Samples were subjected to immunoblot analysis and

quantification as in *B*.

Fig. 2. Effects of alanine substitutions in GFPΔ27BofA on inhibition of Pro-σ^K(1-127) cleavage

in *E. coli*. pET Quartet plasmids were used to produce Pro-σ^K(1-127), cytTM-SpoIVFB,

SpoIVFA, and GFPΔ27BofA (lane 1, pSO40) or Ala-substituted GFPΔ27BofA (lanes 2-20,

pSO44-pSO58 and pSO60-pSO63). Samples collected after 2 h of IPTG induction were

subjected to immunoblot analysis with SpoIVFA, GFP, and penta-His antibodies. Single (*) and

double (**) stars are explained in the Figure 1B legend, as is the graph.

Fig. 3. Effects of alanine substitutions in GFPΔ27BofA during *B. subtilis* sporulation. (*A*)

Effects of GFPΔ27BofA variants (N48A, N61A, T64A) on Pro-σ^K cleavage. Wild-type strain

PY79, a *spoIVB165* null mutant, a *spoIVB165 bofA::erm* double mutant, and the double mutant

with P*$_{bofA}$*-*gfpΔ27bofA* integrated ectopically at *amyE* to express GFPΔ27BofA with no

substitution (none) or the indicated Ala substitution, were starved to induce sporulation.

Samples collected at 4 and 5 h poststarvation (PS) were subjected to immunoblot analysis with

antibodies against SpoIVFA, GFP, SpoIVFB, and Pro-σ^K. The graph shows quantification of the

cleavage ratio [cleavage product/(Pro-σ^K + cleavage product)] for three biological replicates.

Error bars, 1 standard deviation. Student's two-tailed *t* tests were performed to compare certain

cleavage ratios (*p* values). (*B*) Localization of GFPΔ27BofA and the three variants. Samples

collected at 3 h PS were treated with FM 4-64 to stain membranes. Confocal microscopy images

of fluorescence from GFPΔ27BofA, membranes, and merged images are shown for

representative sporangia with discrete (no substitution in GFPΔ27BofA, designated "none"),

partial (N48A and T64A), or no forespore (FS) localization (N61A). Scale bar, 1 μm. The

percentage of sporangia (44-93 counted; non-sporulating cells were not counted) with each

localization pattern is shown.

Fig. 4. MBPΔ27BofA TMS2 is near the cytTM-SpoIVFB active site. (*A*) Disulfide cross-

linking of single-Cys E44C cytTM-SpoIVFB to MBPΔ27BofA C46. pET Quartet plasmids

were used to produce single-Cys E44C cytTM-SpoIVFB in combination with MBPΔ27BofA

C46 (pSO91) or Cys-less MBPΔ27BofA C46S as a negative control (pSO110), and Cys-less

variants of SpoIVFA and Pro-σ^K(1-127) in *E. coli*. Samples collected after 2 h of IPTG

induction were treated with Cu^{2+}(phenanthroline)$_3$ (Cu +) for 60 min or with 2-phenanthroline

(Cu –) as a control. Samples were treated with TCA to precipitate proteins and resuspended in

sample buffer with DTT (+) or without (–) and subjected to immunoblot analysis with FLAG

antibodies to visualize cytTM-SpoIVFB monomer, dimer, and complex with MBPΔ27BofA. (*B*)

Disulfide cross-linking of single-Cys V70C or P135C cytTM-SpoIVFB E44Q variants to

MBPΔ27BofA C46. pET Quartet plasmids were used to produce single-Cys V70C or P135C

cytTM-SpoIVFB E44Q variants in combination with MBPΔ27BofA (pSO92, pSO93) or Cys-

less MBPΔ27BofA C46S as a negative control (pSO111, pSO112), and Cys-less variants of

SpoIVFA and Pro-σ^K(1-127) in *E. coli*. Samples collected after 2 h of IPTG induction were

treated and subjected to immunoblot analysis as in *A*. A representative result from at least two

biological replicates is shown in *A* and *B*.

Fig. 5. GFPΔ27BofA and SpoIVFA do not prevent Pro-σ^K(1-127) from co-purifying with

SpoIVFB. pET Quartet plasmids were used to produce a catalytically-inactive E44C cytTM-

SpoIVFB variant containing FLAG$_2$ but lacking His$_6$ (pSO73), or a variant also lacking FLAG$_2$

as a negative control (pSO149), in combination with Pro-σ^K(1-127), GFPΔ27BofA, and

SpoIVFA in *E. coli*. Samples collected after 2 h of IPTG induction were subjected to co-

immunoprecipitation with anti-FLAG antibody beads. Input, unbound, 1/10 bound (diluted to

match input), and (undiluted) bound samples were subjected to immunoblot analysis with FLAG,

penta-His, SpoIVFA, and GFP antibodies. The single star (*) indicates cross-reacting proteins

below SpoIVFA and the double star (**) indicates a cross-reacting protein or breakdown species

of GFPΔ27BofA that fail to co-purify. A representative result from two biological replicates is

shown.

Fig. 6. Effects of inhibitory proteins on cross-linking between cytTM-SpoIVFB and Pro-σ^K(1-

127). (*A*) Quantification of disulfide cross-linking between single-Cys E44C cytTM-SpoIVFB

and single-Cys Pro-σ^K(1-127) variants in the absence of inhibitory proteins. pET Duet plasmids

were used to produce single-Cys E44C cytTM-SpoIVFB in combination with single-Cys F18C

(pSO167), V20C (pSO169), S21C (pSO170), or K24C (pSO128) Pro-σ^K(1-127), or with Cys-

less Pro-σ^K(1-127) (pSO79) as a negative control, in *E. coli*. Samples collected after 2 h of IPTG

induction were treated with Cu^{2+}(phenanthroline)$_3$ (oxidant; Cu + in *B*) for 15, 30, 45, or 60 min

or with 2-phenanthroline (Cu – in *B*) as a control. Samples were treated with TCA to precipitate

proteins and resuspended in sample buffer with DTT (+ in *B*) or without (– in *B*) and subjected to

immunoblot analysis with FLAG antibodies to visualize the cytTM-SpoIVFB monomer, dimer,

and complex with Pro-σ^K(1-127) (*SI Appendix*, Fig. S15A). Abundance of the complex was

divided by the total amount of cytTM-SpoIVFB monomer, dimer, and complex. The ratio over

time was plotted (n=2) with a best-fit trend line. (*B*) Representative immunoblots of 60-min

samples from the experiment described in *A*. (*C*) Quantification of cross-linking between single-

Cys E44C cytTM-SpoIVFB and single-Cys Pro-σ^K(1-127) variants in the presence of Cys-less

variants of MBPΔ27BofA and SpoIVFA. pET Quartet plasmids were used to produce single-

Cys E44C cytTM-SpoIVFB in combination with single-Cys F18C (pSO163), V20C (pSO165),

S21C (pSO166), or K24C (pSO131) Pro-σ^K(1-127), or with Cys-less Pro-σ^K(1-127) (pSO110) as

a negative control, and Cys-less variants of MBPΔ27BofA and SpoIVFA in *E. coli*. Samples

collected after 2 h of IPTG induction were treated and subjected to immunoblot analysis as in *A*

(*SI Appendix*, Fig. S15B). (*D*) Representative immunoblots of 60-min samples from the

experiment described in *C*. (*E*) Quantification of cross-linking between single-Cys E44C

cytTM-SpoIVFB and single-Cys V20C Pro-σ^K(1-127) in the absence or presence of inhibitory

proteins. Data from *A* (No BofA), *C* (MBPΔ27BofA), and *SI Appendix*, Fig. S16A (full-length

BofA) are plotted along with Cys-less Pro-σ^K(1-127) as a negative control. (*F* and *G*)

Quantification of cross-linking between single-Cys V70C cytTM-SpoIVFB E44Q and single-

Cys F18C or K24C Pro-σ^K(1-127) variants in the absence or presence of inhibitory proteins.

Data from *SI Appendix*, Fig. S17 are plotted using symbols and lines as in *E*. (*H* and *I*)

Quantification of cross-linking between single-Cys Y214C or A231C cytTM-SpoIVFB E44Q

and single-Cys L41C or A97C Pro-σ^K(1-127) in the absence or presence of inhibitory proteins.

Data from *SI Appendix*, Fig. S20 are plotted using symbols and lines as in *E*.

Fig. 7. Model of SpoIVFB monomer with BofA and parts of SpoIVFA and Pro-σ^K. (*A*) Model

of SpoIVFB, BofA TMS2, and the C-terminal part of Pro-σ^K(1-127). At *Left*, a side view of the

complex, showing the six TMSs of the SpoIVFB membrane domain with the active site zinc ion

(magenta), the interdomain linker, and the CBS domain (green), BofA TMS2 (cyan), and Pro-

σ^K(38-114) (red). In the enlarged view of the active site region (*Center*), TMSs 1– 6 of

SpoIVFB and TMS2 of BofA are numbered. At *Right*, a top view is shown. (*B*) Model of

SpoIVFB with full-length BofA and Pro-σ^K(38-114). Predicted TMSs 1 and 2 of BofA are

numbered and its C-terminal region is labeled "C" near the C-terminus in the views shown in the

Center and at *Right*. (*C*) Model of SpoIVFB with full-length BofA, SpoIVFA(65-111) (purple,

residue 111 is numbered), and Pro-σ^K(38-114).

Fig. 1

Fig. 2

Fig. 3

A

spoIVB bofA double mutant

Substitution in GFPΔ27BofA

	Wild type		*spoIVB*		none		N48A		N61A		T64A	
Time PS (h)	4	5	4	5	4	5	4	5	4	5	4	5

αSpoIVFA ← SpoIVFA

αGFP ← GFPΔ27BofA

αSpoIVFB ← * SpoIVFB

αPro-σ^K ← Pro-σ^K ← σ^K

Cleavage ratio

p= 0.02   n= 3   *p*= 0.005   *p*= 0.001   *p*= 0.04

Lane: 1 2 3 4 5 6 7 8 9 10 11 12 13 14

B

Substitution in GFPΔ27BofA	GFP	Membranes	Merge	FS localization of GFPΔ27BofA (%)		
				Discrete	Partial	None
none				99	1	0
N48A				0	84	16
N61A				0	1	99
T64A				0	77	23

Fig. 4

Fig. 5

Fig. 6

Fig. 7

A

Side view   Active site region   Top view

Membrane domain

Linker

CBS domain

90°

B

Side view   Active site region   Top view

Membrane domain

Linker

CBS domain

90°

C

Side view   Active site region   Top view

Membrane domain

Linker

CBS domain

90°