Inhibitory proteins block substrate access by occupying the active site cleft of *Bacillus*

***subtilis* intramembrane protease SpoIVFB**

Sandra Olenic[a], Lim Heo[a], Michael Feig[a], and Lee Kroos[a,1]

[a]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing,

MI 48824

[1]To whom correspondence should be addressed. Email: kroos@msu.edu

Abstract

Intramembrane proteases function in numerous signaling pathways that impact health, but knowledge about regulation of intramembrane proteolysis is limited. We examined inhibition of intramembrane metalloprotease SpoIVFB by proteins BofA and SpoIVFA. We found that BofA residues in and near a predicted transmembrane segment are required for SpoIVFB inhibition, and cross-linking experiments indicated that this transmembrane segment occupies the SpoIVFB active site cleft. SpoIVFA is also required for SpoIVFB inhibition. The inhibitory proteins block access of the substrate N-terminal Proregion to the membrane-embedded SpoIVFB active site, based on additional cross-linking experiments; however, the inhibitory proteins did not prevent interaction between the substrate C-terminal region and the SpoIVFB soluble domain. A structural model was built of SpoIVFB in complex with BofA and parts of SpoIVFA and substrate, using partial homology and constraints from cross-linking and co-evolutionary analyses. The model predicts that conserved BofA residues interact to stabilize a transmembrane segment and a membrane-embedded C-terminal region. SpoIVFA is predicted to bridge the BofA C-terminal region and SpoIVFB, forming a membrane-embedded inhibition complex. Our results reveal a novel mechanism of intramembrane protease inhibition with clear implications for relief from inhibition *in vivo* and design of inhibitors as potential therapeutics.

Introduction

Intramembrane proteases (IPs) are membrane proteins containing a membrane-embedded active site. IPs cleave membrane-associated substrates within a transmembrane segment (TMS) or near the membrane surface in a process referred to as regulated intramembrane proteolysis (RIP) (1). Released substrate fragments impact diverse signaling pathways in a wide variety of organisms (2). IPs also profoundly impact protein degradation (3). There are four known IP families: metallo IPs like SpoIVFB, aspartyl IPs like γ-secretase, serine IPs (rhomboids), and the glutamyl IP Rce1 (2, 4). Crystal structures have been solved for one or more IP in each family (4-9), revealing that TMSs arrange to form a channel that delivers water to the active site for hydrolysis of a substrate peptide bond. Structures have also been solved for rhomboid·peptide (inhibitor and substrate) complexes (10-12) and for γ-secretase·substrate complexes (13, 14), which may guide the design of IP modulators as therapeutics.

Metallo IPs activate transcription factors via RIP in all three domains of life (1, 2). In mammals, S2P is involved in regulation of cholesterol homeostasis and responses to endoplasmic reticulum stress and viral infection (15, 16). S2P homologs in plants and in the pathogenic fungus *Cryptococcus neoformans* play important roles in chloroplast development and virulence, respectively (17, 18). In bacteria, metallo IPs enhance pathogenicity, control stress responses and polar morphogenesis, produce mating signals, and clear signal peptides from the membrane (19-21). In bacilli and most clostridia that form spores, SpoIVFB cleaves inactive Pro-σ^K to σ^K, which directs RNA polymerase to transcribe genes necessary for endospore formation (22-25). Endospores are dormant and are able to survive harsh environmental conditions (26), enhancing the persistence of pathogenic species (27-29). Understanding

regulation of SpoIVFB could lead to new strategies to manipulate endosporulation and other

processes involving IPs in bacteria and eukaryotes.

During endospore formation, *B. subtilis* forms a polar septum that divides the mother cell

(MC) and forespore (FS) compartments (30) (Figure 1-figure supplement 1). The MC engulfs

the FS, surrounding it with a second membrane and pinching it off within the MC. Regulation of

SpoIVFB involves inhibition by BofA and SpoIVFA (31-33). The three proteins form a

complex in the outer FS membrane during engulfment (34-36). BofA was implicated as the

direct inhibitor of SpoIVFB (37), while SpoIVFA appeared to localize and stabilize the

inhibition complex (36, 38). Signaling from the FS relieves inhibition of SpoIVFB (Figure 1-

figure supplement 1). Two proteases, SpoIVB and CtpB, are exported into the space between the

two membranes surrounding the FS (39, 40). SpoIVB cleaves the C-terminal end of SpoIVFA

(41-44) and CtpB can cleave the C-terminal ends of both SpoIVFA and BofA (42, 43, 45). Once

inhibition is removed, SpoIVFB cleaves the N-terminal 21-residue Proregion from Pro-σ^K,

releasing σ^K into the MC (46-48). σ^K directs RNA polymerase to transcribe genes whose

products form the spore coat and lyse the MC, releasing a mature spore (49, 50).

Regulation of SpoIVFB by direct interaction with the inhibitory proteins BofA and

SpoIVFA differs from known regulation of other IPs. Rather, in eukaryotic cells some IP

substrates initially localize to a different organelle than their cognate IP, and regulation involves

membrane trafficking to deliver the substrate to the enzyme (3, 51). In bacteria as well as

eukaryotes, many IP substrates require an initial extramembrane cleavage by another protease

prior to the intramembrane cleavage, and the initial cleavage is typically the regulated step (19-

21, 52, 53). In contrast, Pro-σ^K is cleaved only once, by SpoIVFB. Cleavage of SpoIVFA and

BofA by SpoIVB and CtpB is regulated (Figure 1-figure supplement 1), but how this relieves

inhibition of SpoIVFB has been unknown. Understanding the mechanism of inhibition could

have broad implications since the catalytic core of metallo IPs includes three conserved TMSs

(54) and general principles may emerge that inform potential inhibition strategies for other IP

types.

Cleavage of Pro-σ^K by SpoIVFB can be reproduced by expressing these proteins in

Escherichia coli (37) (Figure 1A). Cleavage could be inhibited by additionally producing

GFPΔ27BofA (a functional fusion protein lacking predicted TMS1 of BofA) and SpoIVFA (37,

45). However, we found that the extent of cleavage inhibition was incomplete and variable.

Here, we describe an improved cleavage inhibition assay. Using the assay and bioinformatics,

we identified three conserved residues of BofA important for inhibition of SpoIVFB in *E. coli*

and in sporulating *B. subtilis.* One of the residues is near the middle of predicted TMS2 of

BofA, close to the sole Cys residue of BofA, C46. We exploited C46 for disulfide cross-linking

experiments that showed BofA TMS2 can occupy the SpoIVFB active site cleft. Additional

experiments in *E. coli* indicate that BofA and SpoIVFA block access of the N-terminal Proregion

of Pro-σ^K to the membrane-embedded SpoIVFB active site, yet the C-terminal soluble regions of

Pro-σ^K and SpoIVFB can interact. These results reveal a novel mechanism of IP inhibition and

were used in combination with prior cross-linking studies (55, 56), partial homology, and

evolutionary co-variation of amino acid residues, to constrain a structural model of SpoIVFB in

complex with BofA and parts of SpoIVFA and Pro-σ^K. The model predicts that conserved

residues in TMS2 and the C-terminal region of BofA interact, and that SpoIVFA bridges the

BofA C-terminal region and SpoIVFB to form an inhibition complex. The predicted inhibition

complex has implications for activation of SpoIVFB and its orthologs in other endospore

formers, as well as for translational efforts to design therapeutic IP inhibitors.

Results

Both BofA and SpoIVFA Are Required to Inhibit SpoIVFB in *E. coli*. To study RIP of Pro-

σ^K, *E. coli* was engineered to synthesize variants of SpoIVFB and Pro-σ^K, and the inhibitory

proteins BofA and SpoIVFA, in various combinations. The SpoIVFB variant contains the extra

TMS cytTM (57) (Figure 1A), which improves accumulation (48). The substrate variant Pro-

σ^K(1-127) lacks the C-terminal half of Pro-σ^K, but is accurately cleaved by SpoIVFB and the

cleavage product is easily separated from Pro-σ^K(1-127) by SDS-PAGE (58). When Pro-σ^K(1-

127) and cytTM-SpoIVFB were produced from the same plasmid, about 80% of the substrate

was cleaved (Figure 1B). The additional production of GFPΔ27BofA and SpoIVFA from a

second plasmid reduced cleavage to about 30%. A similar result was seen when full-length

BofA (for which no antibody has been described) and SpoIVFA were produced from a second

plasmid.

Since the two plasmids had the same replication origin, their copy number may be

unequal in some cells, despite using different selection markers. Unequal copy numbers could

result in Pro-σ^K(1-127) cleavage in cells producing too little of the inhibitory proteins.

Therefore, we designed single "pET Quartet" plasmids to synthesize all four proteins. When

pET Quartets produced either GFPΔ27BofA or BofA, very little Pro-σ^K(1-127) was cleaved

(Figure 1B). A longer exposure of the immunoblot revealed a faint cleavage product with pET

Quartet GFPΔ27BofA, but not with pET Quartet BofA, indicating that full-length BofA inhibits

cleavage slightly better than GFPΔ27BofA.

We tested whether both inhibitory proteins are required for cleavage inhibition by

engineering "pET Triplet" plasmids to synthesize Pro-σ^K(1-127), cytTM-SpoIVFB, and either

GFPΔ27BofA, full-length BofA, or SpoIVFA. With pET Triplets containing GFPΔ27BofA

(Figure 1C) or BofA (Figure 1-figure supplement 2A), cleavage inhibition was lost and

accumulation of cytTM-SpoIVFB, Pro-σ^K(1-127), and cleavage product was greatly diminished,

suggesting that BofA inhibits synthesis and/or enhances degradation of the other proteins when

SpoIVFA is absent. With pET Triplet SpoIVFA, cleavage inhibition was also nearly lost (Figure

1C). We conclude that both inhibitory proteins are necessary to prevent Pro-σ^K(1-127) cleavage

by cytTM-SpoIVB in *E. coli*. Both inhibitory proteins are also required to prevent cleavage of

full-length Pro-σ^K in *E. coli* (Figure 1-figure supplement 2B and Supplementary File 1). The

heterologous system mimics the endogenous pathway of *B. subtilis*, since both inhibitory

proteins are necessary to prevent cleavage of Pro-σ^K by SpoIVFB during sporulation (31, 36,

59).

We also showed that the *E. coli* system mimics the endogenous pathway with respect to

partial relief from inhibition by SpoIVB (40, 42, 43, 45) (Figure 1-figure supplement 3 and

Supplementary File 1) or an F66A substitution in SpoIVFB (59) (Figure 1-figure supplement 4

and Supplementary File 1). We reasoned that the strong cleavage inhibition observed with pET

Quartet plasmids in *E. coli* (Figure 1B) could be used to test the effects of Ala substitutions in

the inhibitory proteins. Since most endospore formers encode BofA, but about half lack a

recognizable gene for SpoIVFA (22, 23), we focused first on identifying residues of BofA

important for inhibition of SpoIVFB.

Three Conserved Residues of BofA Are Important for Inhibition of SpoIVFB in *E. coli*. To

identify residues of BofA that may play a role in SpoIVFB inhibition, an alignment of 70 BofA

orthologs was analyzed (Figure 2-figure supplement 1A). Thirteen highly conserved residues

reside in predicted TMS2 and the C-terminal region. The lack of highly conserved residues in

TMS1 may explain why GFPΔ27BofA (lacking predicted TMS1) is a functional inhibitor (36,

37) (Figure 1B). An alignment of just the 31 BofA orthologs from species with a recognizable

gene for SpoIVFA revealed an additional four conserved residues that we deemed of interest for

Ala substitutions (Figure 2-figure supplement 1B). We also substituted Ala for F85 and I87 (as

well as for I86, which is conserved), since deletion of three residues from the C-terminal end of

BofA caused a loss of function or stability in *B. subtilis* (33, 60).

The Ala substitutions were made in GFPΔ27BofA and co-produced with Pro-σ^K(1-127),

cytTM-SpoIVFB, and SpoIVFA from pET Quartet plasmids in *E. coli*. Three GFPΔ27BofA

variants, N48A, N61A, and T64A led to 40-60% cleavage of Pro-σ^K(1-127), indicating that these

substitutions strongly impaired inhibition of cytTM-SpoIVFB (Figure 2). Less of the N61A and

T64A variants was observed than most of the other GFPΔ27BofA variants, and for all three

variants, less SpoIVFA and cytTM-SpoIVFB was observed, perhaps indicative of reduced

complex formation leading to protein instability, as inferred from studies in *B. subtilis* (36, 61).

We also examined the effects of the GFPΔ27BofA N48A, N61A, and T64A variants during *B.*

subtilis sporulation (see the next section).

The other GFPΔ27BofA variants had less effect on Pro-σ^K(1-127) cleavage, indicating

less effect on cytTM-SpoIVFB inhibition in *E. coli* (Figure 2). Although the F85A, I86A, and

I87A variants had little effect, both a triple-Ala variant and a variant lacking residues 85-87

strongly impaired inhibition of cytTM-SpoIVFB (Figure 2-figure supplement 2A and

Supplementary File 1); the latter mimicking the results in *B. subtilis* (33, 60). We also used pET

Quartet plasmids in *E. coli* to show for the first time that the nine residues preceding predicted

TMS2 of GFPΔ27BofA contribute to its inhibitory function, perhaps by moving the GFP tag

away from the membrane (Figure 2-figure supplement 2B and Supplementary File 1).

The Conserved Residues of BofA Are Important for SpoIVFB Inhibition during *B. subtilis*

Sporulation. To test the effects of the three GFPΔ27BofA variants (N48A, N61A, and T64A)

during sporulation, each was produced ectopically under control of the MC-specific *bofA*

promoter in a *spoIVB165 bofA::erm* double mutant. The *spoIVB165* null mutation alone would

block Pro-σ^K cleavage, but the *bofA::erm* null mutation allows unregulated cleavage in the

double mutant, providing a genetic background in which ectopic production of GFPΔ27BofA

under control of a MC-specific promoter leads to inhibition of SpoIVFB and loss of Pro-σ^K

cleavage (36, 37). As anticipated, Pro-σ^K was cleaved primarily between 4 and 5 h

poststarvation (PS) in wild type, very little cleavage was observed in the *spoIVB165* single

mutant, and cleavage occurred prematurely at 4 h in the *spoIVB165 bofA::erm* double mutant

(Figure 3A). Very little SpoIVFA and SpoIVFB was detected in the double mutant, consistent

with the need for BofA to stabilize these proteins (36). Also as expected, ectopic production of

GFPΔ27BofA in the double mutant inhibited SpoIVFB so that little Pro-σ^K cleavage was

observed (lanes 7 & 8). GFPΔ27BofA allowed slightly more Pro-σ^K cleavage at 5 h (lane 8)

than BofA in the *spoIVB165* single mutant (lane 4), suggesting that full-length BofA is a slightly

better inhibitor of SpoIVFB during *B. subtilis* sporulation, as observed in *E. coli* (Figure 1B,

longer exposure).

Strikingly, ectopic production of the GFPΔ27BofA N48A or N61A variant in the double

mutant did not inhibit SpoIVFB, as Pro-σ^K was cleaved prematurely at 4 h (Figure 3A, lanes 9 &

11). The level of the N48A variant was comparable with GFPΔ27BofA, but less of the N61A

variant was observed, especially at 5 h (lane 12). In comparison with the strain that produced

GFPΔ27BofA, the SpoIVFA level was normal in the strain that produced the N48A variant, but

less SpoIVFB was observed, and less of both SpoIVFA and SpoIVFB was observed in the strain

producing the N61A variant. The reduced levels of SpoIVFB and SpoIVFA may reflect protein

instability due to less complex formation with the GFPΔ27BofA variants, as inferred from

reduced SpoIVFB and SpoIVFA levels in the absence of BofA during *B. subtilis* sporulation

(36).

Production of the T64A variant in the double mutant (Figure 3A, lanes 13 & 14) showed

a pattern similar to wild type; Pro-σ^K cleavage increased between 4 and 5 h, and the levels of

SpoIVFA and SpoIVFB decreased. The levels of all three proteins were slightly lower in the

strain producing the T64A variant than in wild type. The level of the T64A variant was similar

to the N48A and N61A variants at 4 h, yet a lower percentage of the Pro-σ^K was cleaved in the

T64A variant (lanes 9, 11 & 13), suggesting that the T64A variant inhibits SpoIVFB, although

not as well as GFPΔ27BofA (lane 7). We conclude that the three conserved residues of BofA

are important for SpoIVFB inhibition during sporulation.

Since GFPΔ27BofA co-localizes with SpoIVFA and SpoIVFB to the outer FS membrane

during sporulation (34-36), we examined the ability of the GFPΔ27BofA variants to localize to

the FS. As a control, GFPΔ27BofA produced in the *spoIVB165 bofA::erm* double mutant

localized discretely to the FS at 3 h PS (Figure 3B). The N48A and T64A variants localized

partially to the FS, but GFP fluorescence was also observed in the MC cytoplasm, suggesting

partial mislocalization. The N61A variant appeared to be dispersed evenly throughout the MC

cytoplasm. The fusion proteins were intact, demonstrating that the cytoplasmic GFP

fluorescence was not attributable to breakdown (Figure 3-figure supplement 1). Inability of the

N61A variant to localize to the FS may explain the loss of SpoIVFB inhibition and the abundant

cleavage of Pro-σ^K at 4 h (Figure 3A, lane 11). Importantly, the similar ability of the N48A and

T64A variants to localize to the FS (Figure 3B) does not account for their differential effects on

the level of SpoIVFB and its ability to cleave Pro-σ^K (Figure 3A). The strain producing

GFPΔ27BofA N48A exhibited less SpoIVFB yet more Pro-σ^K cleavage at 4 h (lane 9) than the

strain producing GFPΔ27BofA T64A (lane 13), so the N48A substitution more severely impairs

the ability of GFPΔ27BofA to inhibit SpoIVFB.

BofA TMS2 Occupies the SpoIVFB Active Site Cleft. A common mechanism of

metalloprotease inhibition involves a protein residue side chain coordinating the catalytic metal

ion in place of a water molecule necessary for substrate peptide bond hydrolysis (62, 63). Early

work suggested that BofA H57 might inhibit SpoIVFB by coordinating zinc (37). Although His

is highly conserved at the corresponding position of BofA orthologs (Figure 2-figure supplement

1) and GFPΔ27BofA H57A allowed about 20% cleavage of Pro-σ^K(1-127) in *E. coli* (Figure 2),

H57 is near the predicted C-terminal end of BofA TMS2 (Figure 2-figure supplement 1) and

therefore presumably near the membrane surface, whereas a zinc ion at the SpoIVFB active site

is predicted to be near the middle of the lipid bilayer (56). The sole Cys of *B. subtilis* BofA,

C46, is a potential zinc ligand located near the center of TMS2, but Cys is not conserved at the

corresponding position in BofA orthologs (Figure 2-figure supplement 1). However, nearby N48

is highly conserved among BofA orthologs and is also a potential zinc ligand. For example, the

latent form of *Staphylococcus aureus* LytM has a prosegment with an Asn residue as a zinc

ligand (64). We hypothesized that BofA N48 is a zinc ligand and that BofA TMS2 occupies the

SpoIVFB active site cleft in the inhibition complex.

To begin testing our hypothesis, we devised a strategy based on a model of the SpoIVFB

membrane domain derived from the crystal structure of an archaeal homolog (7). The model is

supported by cross-linking studies of catalytically-inactive SpoIVFB E44Q in complex with Pro-

σ^K(1-127), whose Proregion appears to occupy the SpoIVFB active site cleft (55, 56, 65). A

cleft between TMS1 and TMS6 of SpoIVFB is proposed to gate substrate access to the active site

(59), which mutational evidence supports is formed by a zinc ion near E44 of the HELGH

metalloprotease motif within SpoIVFB TMS2 (46, 47) (Figure 4-figure supplement 1A). If

BofA N48 is a zinc ligand and BofA TMS2 occupies the SpoIVFB active site cleft in the

inhibition complex, we reasoned that BofA C46 may be near SpoIVFB E44 (Figure 4-figure

supplement 1B) and we used a disulfide cross-linking approach to test this possibility.

We tested whether BofA C46 could be cross-linked to SpoIVFB E44C in the inhibition

complex formed with pET Quartet plasmids in *E. coli* as in Figure 1B, but with appropriate

variants of the four proteins. Single-Cys E44C cytTM-SpoIVFB (which is catalytically-inactive)

and Cys-less Pro-σ^K(1-127) (which can be cleaved by active SpoIVFB) were described

previously (55). We created functional variants of Cys-less SpoIVFA and single-Cys

MBPΔ27BofA (GFP has Cys residues but MBP lacks them, leaving BofA C46 as the sole Cys

residue) (Figure 4-figure supplement 2). As a negative control, we replaced C46 with Ser to

obtain functional Cys-less MBPΔ27BofA. *E. coli* induced to co-produce combinations of four

proteins were treated with the oxidant Cu^{2+}(phenanthroline)$_3$ to promote disulfide bond

formation. For MBPΔ27BofA C46 (Figure 4A, lane 2), but not the Cys-less C46S negative

control (lane 5), treatment with oxidant caused formation of a species of the expected size for a

cross-linked complex with single-Cys E44C cytTM-SpoIVFB, which was detected by

immunoblotting with anti-FLAG antibodies. Treatment with the reducing agent DTT greatly

diminished the abundance of the apparent complex, consistent with cross-link reversal (lane 3).

A species of the expected size for a cross-linked dimer of single-Cys E44C cytTM-SpoIVFB was

also observed. Formation of the apparent dimer varies, as reported previously (55). As

expected, anti-MBP antibodies detected the presumptive cross-linked complex of MBPΔ27BofA

C46 with single-Cys E44C cytTM-SpoIVFB, albeit weakly, and the negative control with E44Q

rather than E44C failed to form the complex (Figure 4-figure supplement 3, lanes 2 & 5). Since

the signal for the complex was stronger with anti-FLAG antibodies (Figure 4A), we used those

antibodies in the cross-linking experiments reported below.

In addition to E44 of SpoIVFB, V70 in a predicted membrane-reentrant loop and P135 in

a predicted short loop interrupting TMS4 (Figure 1A and Figure 4-figure supplement 1A) were

shown to be in proximity to the Proregion of Pro-σ^K(1-127) (55). Therefore, we tested whether

MBPΔ27BofA C46 could be cross-linked to single-Cys V70C or P135C cytTM-SpoIVFB E44Q

variants. We included the inactivating E44Q substitution since the V70C and P135C variants

(unlike the E44C variant) could cleave Cys-less Pro-σ^K(1-127) (55), even though the inhibitory

proteins were expected to almost completely inhibit cleavage (Figure 4-figure supplement 2).

MBPΔ27BofA C46 formed a complex with the P135C variant, but not with the V70C variant

(Figure 4B, lanes 2 & 8). As expected, Cys-less MBPΔ27BofA C46S failed to form a complex

with either variant (lanes 5 & 11). Full-length BofA C46 (lacking MBP) also formed a complex

of the expected (smaller) size with the E44C and P135C variants, but not with the V70C variant

(Figure 4-figure supplement 4, lanes 2, 14, & 8). As expected, Cys-less BofA C46S failed to

form a complex with any of the variants (lanes 5, 11, & 17). Our cross-linking results show that

BofA TMS2 occupies the SpoIVFB active site cleft in the inhibition complex, placing BofA C46

in proximity to SpoIVFB E44 and P135, but not V70.

Based on our initial cross-linking results, we modeled BofA TMS2 in the SpoIVFB

active site cleft, and tested predictions of the model using additional disulfide cross-linking

experiments (Figure 4-figure supplement 5 and Supplementary File 1). The results confirmed

predictions of the initial model and suggested a preferred orientation of BofA TMS2 in the

SpoIVFB active site cleft, which led to the refined model shown in Figure 4-figure supplement

1B.

BofA and SpoIVFA Do Not Prevent Pro-σ^K(1-127) from Interacting with SpoIVFB. Since

our cross-linking results show that BofA TMS2 occupies the SpoIVFB active site cleft, we tested

whether BofA and SpoIVFA prevent Pro-σ^K(1-127) from interacting with SpoIVFB in *E. coli*. A

catalytically-inactive E44C cytTM-SpoIVFB variant with a FLAG$_2$ epitope tag was co-produced

with Pro-σ^K(1-127), SpoIVFA, and GFPΔ27BofA. Cell lysates were prepared and detergent-

solubilized proteins were co-immunoprecipitated in pull-down assays with anti-FLAG antibody

beads. All four proteins were seen in the bound sample (Figure 5A, lane 4), indicating that

GFPΔ27BofA and SpoIVFA did not completely prevent Pro-σ^K(1-127) from interacting with the

cytTM-SpoIVFB variant. However, only the cytTM-SpoIVFB variant was detected when the

bound sample was diluted tenfold (lane 3) to match the input sample concentration (lane 1), and

portions of the other three proteins were observed in the unbound sample (lane 2), indicating that

co-purification was inefficient. A negative control with the cytTM-SpoIVFB variant lacking the

FLAG$_2$ epitope tag showed no Pro-σ^K(1-127) in the bound sample, but small amounts of

GFPΔ27BofA and SpoIVFA were detected (lane 8), indicative of weak, nonspecific binding to

the beads.

We also performed pull-down assays with cobalt resin, which binds to the His_6 tag on

Pro-σ^K(1-127), and we performed both types of pull-down assays (i.e., anti-FLAG antibody

beads and cobalt resin) when full-length BofA rather than GFPΔ27BofA was co-produced with

the other three proteins (Figure 5-figure supplement 1 and Supplementary File 1). Neither BofA

nor GFPΔ27BofA when co-produced with SpoIVFA prevented Pro-σ^K(1-127) from interacting

with the cytTM-SpoIVFB variant. However, GFPΔ27BofA and SpoIVFA reduced co-

purification of full-length Pro-σ^K-His_6 with the cytTM-SpoIVFB variant in both types of pull-

down assays (Figure 5-figure supplement 2 and Supplementary File 1), as compared with Pro-

σ^K(1-127) (Figure 5A and Figure 5-figure supplement 1), suggesting that the C-terminal half of

Pro-σ^K affects complex formation (Supplementary File 2).

To explain the presence of substrate in the pulled-down protein complexes, but the

absence of substrate cleavage (Figure 1B), we hypothesized that inhibitory proteins block

substrate access to the SpoIVFB active site cleft, but do not prevent substrate interaction with the

soluble C-terminal CBS domain of SpoIVFB (Figure 1A). The model of catalytically-inactive

SpoIVFB E44Q in complex with Pro-σ^K(1-127) predicts extensive interactions between the

SpoIVFB CBS domain and the Pro-σ^K(1-127) C-terminal region (56). We used the model to

guide testing of single-Cys variants of the two proteins for disulfide cross-link formation. We

discovered that A231C in the cytTM-SpoIVFB CBS domain can be cross-linked to A97C in the

Pro-σ^K(1-127) C-terminal region when *E. coli* co-producing the proteins are treated with oxidant

(Figure 5-figure supplement 3A). We showed that the Cys substitutions do not impair cytTM-

SpoIVFB activity or Pro-σ^K(1-127) susceptibility to cleavage (Figure 5-figure supplement 3B).

Finally, we measured time-dependent cross-linking in the presence or absence of Cys-less

inhibitory proteins. Co-production of inhibitory proteins had little or no effect on formation of

cross-linked complex (Figure 5B and Figure 5-figure supplement 3C). These results suggest that

neither full-length BofA nor MBPΔ27BofA, when co-produced with SpoIVFA, prevent Pro-

σ^K(1-127) from interacting with the CBS domain of SpoIVFB in *E. coli*, consistent with the

results of our pull-down assays (Figure 5A and Figure 5-figure supplement 1).

Inhibitory Proteins Block Access of the Substrate N-terminal Proregion to the SpoIVFB

Active Site. SpoIVFB cleaves Pro-σ^K (49) and Pro-σ^K(1-127) (37) between residues S21 and

Y22. In disulfide cross-linking experiments, Cys substitutions for several residues near the

cleavage site in otherwise Cys-less Pro-σ^K(1-127) formed a cross-linked complex with single-

Cys (E44C, V70C, or P135C) cytTM-SpoIVFB variants (55). The complex was most abundant

with the E44C and V70C variants, so we compared these interactions in the presence or absence

of Cys-less inhibitory proteins.

SpoIVFB E44 is presumed to activate a water molecule for substrate peptide bond

hydrolysis at the enzyme active site (46, 47). To test access of the substrate Proregion to the

enzyme active site, we first measured time-dependent crosslinking between single-Cys E44C

cytTM-SpoIVFB and single-Cys (F18C, V20C, S21C, K24C) Pro-σ^K(1-127) variants co-

produced in the absence of inhibitory proteins in *E. coli*. The V20C and K24C Pro-σ^K(1-127)

variants formed abundant complex that increased over time, but the F18C and S21C variants

formed much less complex, only slightly more than the Cys-less Pro-σ^K(1-127) negative control

(Figure 6A and Figure 6-figure supplement 1A). Figure 6B shows a representative immunoblot

(60-min oxidant treatment). When Cys-less MBPΔ27BofA and SpoIVFA were co-produced, the

V20C and K24C Pro-σ^K(1-127) variants formed much less complex and its abundance did not

increase over time (Figure 6C and Figure 6-figure supplement 1B). Figure 6D shows a

representative immunoblot (60-min oxidant treatment) for comparison with Figure 6B. When

Cys-less full-length BofA and SpoIVFA were co-produced, similar decreases in abundance of

the complex were observed (Figure 6-figure supplement 2). Figure 6 E and F summarize the

cross-linking time courses with and without inhibitory proteins for the V20C and K24C Pro-

σ^K(1-127) variants. The effects of MBPΔ27BofA (lacking TMS1) and full-length BofA were

indistinguishable. These results likely explain why BofA TMS1 is dispensable for most of the

inhibitory function of BofA (36, 37) (Figure 1B and 3A). We conclude that inhibitory proteins

block access of the substrate N-terminal Proregion to the SpoIVFB active site.

SpoIVFB V70 is located in a predicted membrane-reentrant loop (Figure 1A), which may

bind to the Proregion and present it to the active site for cleavage (56) based on a study *E. coli*

RseP (66). Single-Cys V70C cytTM-SpoIVFB E44Q formed abundant cross-linked complex

with F18C and K24C Pro-σ^K(1-127) variants in the absence of inhibitory proteins (55) (Figure 6

G and H; Figure 6-figure supplement 3 A and B). When Cys-less MBPΔ27BofA and SpoIVFA

were co-produced, less complex formed (Figure 6 G and H; Figure 6-figure supplement 3 C and

D). Cys-less full-length BofA and SpoIVFA further decreased complex formation (Figure 6 G

and H; Figure 6-figure supplement 3 E and F) and caused four novel species to be observed,

including with the Cys-less Pro-σ^K(1-127) negative control (Figure 6-figure supplement 3G),

perhaps due to cross-linking of the V70C cytTM-SpoIVFB variant to *E. coli* proteins (i.e., BofA

may cause the SpoIVFB membrane-reentrant loop to be exposed). Since full-length BofA

hindered cross-linking more than MBPΔ27BofA (lacking TMS1) (Figure 6 G and H), both

TMSs of BofA appear to interfere with the normal interaction between the SpoIVFB membrane-

reentrant loop and the substrate Proregion.

Since full-length BofA also inhibited cleavage of Pro-σ^K(1-127) in *E. coli* (Figure 1B)

and Pro-σ^K in *B. subtilis* (Figure 3A) slightly more than GFPΔ27BofA (lacking TMS1), we

compared time-dependent cross-linking of C46 of BofA and MBPΔ27BofA to single-Cys E44C

and P135C cytTM-SpoIVFB variants. The results suggest that full-length BofA forms slightly

more complex over time, whereas complex abundance did not increase over time with

MBPΔ27BofA (Figure 6-figure supplement 4). Hence, BofA TMS1 may slightly enhance TMS2

occupancy in the SpoIVFB active site cleft.

To examine the extent to which inhibitory proteins hinder the interaction of the substrate

with SpoIVFB, we identified a residue in the linker between the membrane and CBS domains of

SpoIVFB (Figure 1A) that forms disulfide cross-linked complexes with Pro-σ^K(1-127). The

model of catalytically-inactive SpoIVFB E44Q in complex with Pro-σ^K(1-127) (56) was used to

guide testing for disulfide cross-link formation and we found that A214C in the cytTM-SpoIVFB

linker can be cross-linked to A41C in Pro-σ^K(1-127) (Figure 6-figure supplement 5A). We

showed that the Cys substitutions do not impair cytTM-SpoIVFB activity or Pro-σ^K(1-127)

susceptibility to cleavage (Figure 5-figure supplement 3B). Finally, we measured time-

dependent cross-linking in the presence or absence of Cys-less inhibitory proteins. Interestingly,

formation of cross-linked complex was hindered more by full-length BofA than by

MBPΔ27BofA, when co-produced with SpoIVFA (Figure 6I and Figure 6-figure supplement

5B), similar to cross-linking between V70C in the cytTM-SpoIVFB membrane-reentrant loop

and F18C or K24C near the cleavage site in Pro-σ^K(1-127) (Figure 6 G and H). The similar

pattern suggests that in both cases BofA TMS2 partially interferes with the interaction and BofA

TMS1 augments the interference.

A Model of SpoIVFB in Complex with BofA and Parts of SpoIVFA and Pro-σ^K.

Computational models were generated using a similar protocol as described previously (56), but

including additional constraints reflecting experimental cross-linking data reported herein, as

well as newly predicted intra- and inter-chain contacts based on co-evolutionary couplings. Two

final models were generated: 1) Full-length SpoIVFB modeled as a tetramer, with part of one

Pro-σ^K molecule (residues 1-114), referred to as 'fb.sigk'; 2) Full-length SpoIVFB, again

modeled as a tetramer, with one molecule each of full-length BofA and parts of Pro-σ^K (residues

38-114) and SpoIVFA (residues 65-111), referred to as 'fb.sigk.bofa.fa'. The omitted residues of

Pro-σ^K and SpoIVFA could not be placed with sufficient confidence. The first model is

described in Figure 7-figure supplement 1 and Supplementary File 3.

Relevant features of the second model are illustrated in Figure 7 (only one molecule of

SpoIVFB is shown). The first side view shows SpoIVFB with BofA TMS2 and the C-terminal

part of Pro-σ^K(1-127) (Figure 7A). The membrane domain of SpoIVFB (green) interacts with

BofA TMS2 (cyan) while the interdomain linker and CBS domain of SpoIVFB interact with the

modeled portion of Pro-σ^K(1-127) (red). The enlarged view of the SpoIVFB active site cleft

shows BofA TMS2 surrounded by SpoIVFB (TMSs labeled 1-6). The top view of the membrane

domain emphasizes proximity between BofA TMS2, SpoIVFB TMSs 2-4, and the zinc ion

(magenta) involved in catalysis. The second side view shows SpoIVFB with full-length BofA

and the C-terminal part of Pro-σ^K(1-127) (Figure 7B). Our model predicts that BofA contains 2

TMSs and a membrane-embedded C-terminal region (labeled C near the C-terminus in the

enlarged view of the SpoIVFB active site cleft) that forms two short α-helices connected by a

turn. The enlarged and top views show that BofA interacts extensively with SpoIVFB and

occupies its active site cleft, which would sterically hinder access of the Proregion of Pro-σ^K.

Figure 7C shows the addition of the modeled portion of SpoIVFA (purple), which is predicted by

co-evolutionary couplings to contact the BofA C-terminal region and SpoIVFB TMS4. Hence,

the model predicts that SpoIVFA stabilizes the interaction of BofA with SpoIVFB. Since

GFPΔ27BofA has the C-terminal region predicted to interact with SpoIVFA, the interaction of

GFPΔ27BofA with SpoIVFB is likewise predicted to be stabilized by SpoIVFA, which may

explain the dependence of SpoIVFB inhibition on both GFPΔ27BofA and SpoIVFA (Figure 1C),

and why GFPΔ27BofA inhibits substrate cleavage (Figure 1B and 3A) and MBPΔ27BofA

occupies the SpoIVFB active site cleft (Figure 6-figure supplement 4) nearly as well as full-

length BofA.

Discussion

Our results provide evidence that BofA TMS2 occupies the SpoIVFB active site cleft. Both

inhibitory proteins block access of the substrate N-terminal Proregion to the SpoIVFB active site,

but do not prevent interaction between the C-terminal region of Pro-σ^K(1-127) and the SpoIVFB

CBS domain. The mechanism of SpoIVFB inhibition is novel in comparison with previously

known mechanisms of IP regulation. Structural modeling predicts that three conserved BofA

residues (N48, N61, and T64) interact to stabilize TMS2 and a membrane-embedded C-terminal

region. SpoIVFA is predicted to contact the BofA C-terminal region and SpoIVFB TMS4,

bridging the two proteins to stabilize the inhibition complex. The model has clear implications

for relief of SpoIVFB and its orthologs from inhibition during sporulation, as well as for IP

inhibitor design.

A Novel Mechanism of Intramembrane Protease Regulation. IPs were known to be regulated

by three mechanisms – substrate localization, substrate extramembrane domain cleavage, and

substrate interaction with an adapter protein. Substrate localization to a different organelle than

its cognate IP is common in eukaryotic cells (3, 51). An early example of this regulatory

mechanism emerged from studies of S2P. SREBP substrates are retained in the endoplasmic

reticulum by SCAP and Insig proteins, then transported to the Golgi when more cholesterol is

needed (67) (Figure 8).

RIP of SREBPs also involves extramembrane loop cleavage, referred to as site-1

cleavage, prior to site-2 cleavage by S2P (1) (Figure 8). Substrates of bacterial metallo IPs (with

the exception of SpoIVFB) typically also require initial regulated site-1 cleavage by another

protease (19). A well-studied example in *E. coli* involves envelope stress regulating site-1

cleavage of the RseA extramembrane domain by DegS, allowing site-2 cleavage by RseP (20,

68) (Figure 8).   Substrates of aspartyl IPs likewise require an initial cleavage by another

protease, referred to as ectodomain shedding (52, 53). Shedding and site-1 cleavage of RseA

allow substrate entry to the IP active site cleft. Without the initial cleavage, the substrate

extramembrane domain prevents entry, due to size exclusion by the IP. In RseP, tandem PDZ

domains act as a size-exclusion filter (69) (Figure 8). The nicastrin subunit plays a similar role

in the metazoan aspartyl IP γ-secretase (70).

Only two cases of IP regulation by substrate interaction with an adapter protein have been

reported. In the bacterium *Mycobacterium tuberculosis*, the adapter protein Ppr1 bridges the

metallo IP Rip1 to one of its substrates, but not to others (21). The γ-secretase subunits nicastrin

and PEN-2 also appear to interact with substrate before passage to the active site (71).

Our results support a novel mechanism of IP regulation – inhibitory proteins block

substrate access to the SpoIVFB active site (Figure 8). Many soluble proteases are regulated by

inhibitory proteins, prodomains, or prosegments that sterically hinder substrate access to their

active site (62), but analogous regulation of an IP has not been reported previously. In addition

to steric hindrance, inhibition of soluble metalloproteases often involves a protein residue side

chain coordinating the catalytic metal ion (62, 63). We hypothesized that BofA N48 coordinates

the predicted zinc ion at the SpoIVFB active site since Asn is highly conserved at the

corresponding position in BofA orthologs (Figure 2-figure supplement 1) and BofA N48 is

crucial for SpoIVFB inhibition (Figure 2 and 3). Consistent with our hypothesis, BofA TMS2

with N48 near its center occupies the SpoIVFB active site cleft, based on disulfide cross-linking

experiments (Figure 4 and Figure 4-figure supplement 3-5). However, the N48 side chain points

away from the zinc ion in our structural model, which instead predicts that the N48 side chain

interacts with those of other conserved BofA residues shown to be important for SpoIVFB

inhibition (Supplementary File 2). Therefore, we favor a simple steric hindrance mechanism of

inhibition not involving a BofA residue acting as a zinc ligand.

In addition to BofA TMS2 sterically hindering substrate access to the SpoIVFB active

site, what are the roles of other parts of BofA in the novel inhibition mechanism? It was known

that deletion of three residues from the C-terminal end of BofA caused a loss of function or

stability in *B. subtilis* (33, 60). We found that deleting the three residues or changing them to

Ala in GFPΔ27BofA caused loss of SpoIVFB inhibition in our *E. coli* system (Figure 2-figure

supplement 2A). Our structural model predicts that the three BofA residues interact with

SpoIVFA and SpoIVFB (Figure 7C), which can be tested by cross-linking experiments.

BofA TMS1 appears to sterically hinder some interactions between the Pro-σ^K N-

terminal region and SpoIVFB, based on comparison between full-length BofA and

MBPΔ27BofA (lacking TMS1) in cross-linking time courses. The interactions hindered more by

full-length BofA included cross-links between the Pro-σ^K Proregion and the SpoIVFB

membrane-reentrant loop (Figure 6G and H), which may bind to the Proregion and present it to

the SpoIVFB active site for cleavage (56) based on a study of the homologous membrane-

reentrant loop of *E. coli* RseP (66). Full-length BofA also hindered cross-links between Pro-σ^K

L41C and the SpoIVFB interdomain linker (Y214C) more than MBPΔ27BofA (Figure 6I),

suggesting that BofA TMS1 weakens the interaction between the interdomain linker and Pro-σ^K,

which is crucial for cleavage (56, 59). Our model predicts that both TMSs of BofA, as well as its

C-terminal region, occupy the SpoIVFB active site cleft (Figure 7B), plausibly explaining the

additional steric hindrance that BofA TMS1 appears to provide. Yet, TMS1 plays a minor role

in SpoIVFB inhibition compared with BofA TMS2 and its C-terminal region, since full-length

BofA inhibited cleavage of Pro-σ^K(1-127) in *E. coli* (Figure 1B) and Pro-σ^K in *B. subtilis* (Figure

3A) only slightly more than GFPΔ27BofA. Nevertheless, the minor role of BofA TMS1 may be

important as premature σ^K production can reduce sporulation efficiency (31).

Implications for Activation of SpoIVFB and Its Orthologs in Other Endospore Formers.

Inhibition of SpoIVFB is relieved during *B. subtilis* sporulation when SpoIVB cleaves the C-

terminal end of SpoIVFA (41-44) and CtpB cleaves the C-terminal ends of both SpoIVFA and

BofA (42, 43, 45) (Figure 1-figure supplement 1). The levels of inhibitory proteins decrease

coincident with increasing Pro-σ^K cleavage (38, 41, 42, 45), suggesting that degradation of

SpoIVFA and BofA permits SpoIVFB activity. Alternatively, a conformational change in the

inhibition complex may suffice to trigger SpoIVFB activity, based on *in vitro* experiments

indicating that SpoIVB cleavage of SpoIVFA alters susceptibility of the complex to general

proteolysis but does not disrupt the complex (42).

Our results imply that activation of SpoIVFB requires removal or substantial movement

of inhibitory proteins to allow substrate access to the active site. Foremost, steric hindrance by

BofA TMS2 must be relieved. Additionally, the critical roles of the BofA C-terminal region and

SpoIVFA must be explained. Our structural model predicts that SpoIVFA bridges the BofA C-

terminal region and SpoIVFB TMS4 to stabilize the inhibition complex (Figure 7C). We

propose that cleavage of SpoIVFA by SpoIVB initiates destabilization of the inhibition complex.

The initial cleavage, perhaps followed by further proteolysis of SpoIVFA, may destabilize the

membrane-embedded BofA C-terminal region and expose it to cleavage by CtpB, which hastens

SpoIVFB activation but is not essential (40, 42, 43, 45). Finally, degradation of BofA or a

conformational change would relieve steric hindrance by BofA TMS2 and allow substrate access

to the SpoIVFB active site.

The mechanisms of inhibition and activation of *B. subtilis* SpoIVFB likely apply to its

orthologs in spore-forming bacilli since these bacteria encode BofA, SpoIVFA, SpoIVB, and

CtpB (23, 25). This group includes well-known human or plant pathogens such as *Bacillus*

anthracis, *Bacillus cereus*, and *Bacillus thuringiensis*. In contrast, clostridia that form

endospores do not have a recognizable gene for SpoIVFA (22, 23). In some clostridia, a non-

orthologous gene has been proposed to code for a protein that performs the same function as

SpoIVFA (23), so similar mechanisms of SpoIVFB regulation may apply. The *E. coli* system

described herein will facilitate further testing of heterologous protein function. Many spore-

forming clostridia also lack a recognizable gene for BofA (25). Our results imply that SpoIVFB

activity would be unregulated in these bacteria (i.e., coordination between forespore and mother

cell gene expression would be lost). A precedence for less precise control of σ^K-dependent gene

expression is known from studies of the human pathogen *Clostridioides difficile* (72, 73), in

which the gene for σ^K does not encode a Proregion and a gene for SpoIVFB is absent (74).

Based on phylogenomic analysis, the precise control of σ^K-dependent gene expression observed

in *B. subtilis* emerged early in evolution and has been maintained in bacilli which propagate

primarily in aerobic environments, but has been modified or lost in many clostridia which are

found mainly in anaerobic habitats (25).

Design of IP Inhibitors. Efforts toward development of therapeutic peptidyl inhibitors of serine

IPs (rhomboids) are well-advanced (11, 12, 75). Translational work on aspartyl IPs has focused

on γ-secretase owing to its processing of the amyloid precursor protein (APP) associated with

Alzheimer's disease (76). Wide-spectrum inhibitors of γ-secretase exhibit toxicity in clinical

trials, mainly due to inhibition of signaling *via* Notch receptors, which are also substrates of γ-

secretase (77). The structures of γ-secretase complexes with APP and Notch reveal differences

in binding that may allow substrate-specific inhibitors to be developed as therapeutics (13, 14).

Stabilization of complexes in which γ-secretase progressively cleaves APP is another promising

approach toward development of drugs to treat Alzheimer's disease (78).

In comparison with rhomboids and γ-secretase, translational work on metallo IP

inhibitors is in its infancy. Recently, batimastat, a peptidic hydroxamate known to inhibit

eukaryotic matrix metalloproteases (MMPs), was shown to selectively inhibit *E. coli* RseP *in*

vivo (79). A lack of selectivity has been a problem in efforts toward using peptidic

hydroxamates as MMP inhibitors for treatment of cancer, arthritis, and other diseases (80-82).

The hydroxamate strongly coordinates the catalytic metal ion and the peptidic portions have not

provided enough specificity to prevent off-target effects. Even so, the side effects may not

preclude using peptidic hydroxamates to treat bacterial infections topically, and systemically if

administered briefly. Other small-molecule MMP inhibitors that have already been developed

(80) should be screened for activity against bacteria and fungi with metallo IPs known to be

involved in pathogenesis (18-21).

Our findings reveal concepts that may inform efforts to design selective inhibitors of IPs.

BofA TMS2 appears to block the SpoIVFB active site (Figure 7A), similar to the prosegment of

many latent proteases (62). Like BofA, some prosegments can act in *trans*, an observation that

has led to the demonstration of prodomains as selective inhibitors of A Disintegrin And

Metalloproteinase (ADAM) family enzymes (83, 84). Selectivity relies on the extensive

interaction surface between the prodomain and the enzyme, including features specific to the

pair, which can boost efficacy and diminish off-target effects in therapeutic strategies (82). A

peptide encompassing residues 48 to 64 of BofA may inhibit SpoIVFB, but our model also

predicts extensive interactions between the BofA C-terminal region (residues 65 to 87) and

SpoIVFB (Figure 7B), so a longer BofA peptide, perhaps flexibly linked to SpoIVFA residues 96

to 109, which are also predicted to interact with SpoIVFB (Figure 7C), may exhibit improved

inhibition and selectivity. Since cleavage of Pro-σ^K(1-127) by cytTM-SpoIVFB has been

reconstituted *in vitro* (48), it will be possible to test peptides for inhibition. Structure

determination of SpoIVFB in complex with an inhibitory peptide, GFPΔ27BofA, or full-length

BofA is an important goal to facilitate design of metallo IP inhibitors. In particular, it may

inform efforts to design inhibitors of SpoIVFB orthologs in pathogenic bacilli and clostridia,

which persist by forming endospores (27-29). Such efforts could lead to new strategies to

control endosporulation.

Our findings also suggest that BofA TMS1 interferes with interactions between the Pro-

σ^K N-terminal region and SpoIVFB. Therefore, a cyclic peptide inhibitor may prove to be more

effective than a linear one. The desirable characteristics of cyclic peptides as therapeutics, and

new methods of producing and screening cyclic peptide libraries, make this an attractive

possibility (85, 86). Although we favor a simple steric hindrance mechanism of inhibition not

involving a BofA residue acting as a zinc ligand (Supplementary File 2), it is possible that

inclusion of such a residue in a BofA TMS2 peptide mimic would improve SpoIVFB inhibition.

A similar strategy could be used to design inhibitors of other metallo IPs.

Materials and Methods

Plasmids, Primers, and Strains. Plasmids used in this study are described in Supplementary

File 4, as are primers used in plasmid construction. Plasmids were cloned in *E. coli* strain DH5α

(87). Relevant parts of plasmids were verified by DNA sequencing with primers listed in

Supplementary File 4 and *B. subtilis* strains used in this study are also described therein.

Pro-σ^K(1-127) Cleavage in *E. coli*. Strain BL21(DE3) (Novagen) was used to produce proteins

in *E. coli*. Two plasmids with different antibiotic resistance genes were cotransformed (37) or a

single plasmid was transformed, with selection on Luria-Bertani (LB) agar supplemented with

kanamycin sulfate (50 μg/mL) and/or ampicillin (100 μg/mL). Transformants (4-5 colonies)

were grown in LB medium with 50 μg/mL kanamycin sulfate and/or 200 μg/mL ampicillin at

37°C with shaking (200 rpm). Typically, overnight culture (200 μL) was transferred to 10 mL of

LB medium with antibiotics, cultures were grown at 37°C with shaking (250 rpm) to an optical

density of 60-80 Klett units, and isopropyl β-D-thiogalactopyranoside (IPTG) (0.5 mM) was

added to induce protein production for 2 h. For transformants with either pET Quintet or full-

length Pro-σ^K-His$_6$, overnight growth was avoided. Transformants were transferred directly to

10 mL of LB medium with antibiotic, and cultures were grown and induced as described above.

Equivalent amounts of cells (based on optical density in Klett units) were collected ($12,000 \times g$

for 1 min) and extracts were prepared (37), then subjected to immunoblot analysis

(Supplementary File 3).

BofA Sequence Analysis. Orthologs of *B. subtilis bofA*, which are present in the genomes of

most endospore-forming bacteria (23), were collected from the NCBI and Uniprot databases.

The protein sequences of BofA orthologs were aligned using the T-Coffee multiple sequence

alignment package (88). Residues identical in at least 70% of the sequences were considered

conserved.

B. subtilis **Sporulation and GFPΔBofA Localization.** GFPΔ27BofA or its variants were

expressed under control of the *bofA* promoter (P$_{bofA}$) after ectopic chromosomal integration.

Plasmids bearing P$_{bofA}$-*gfpΔ27bofA* or its variant, bordered by regions of homology to *B. subtilis*

amyE, were transformed into strain ZR264. Transformants with a gene replacement at *amyE*

were selected on LB agar with spectinomycin sulfate (100 μg/mL) and identified by loss of

amylase activity (89). Sporulation was induced by growing cells in the absence of antibiotics

followed by the resuspension of cells in SM medium (89). At indicated times PS, samples (50

μL) were centrifuged ($12,000 \times g$ for 1 min), supernatants were removed, and cell pellets were

stored at -80°C. Whole-cell extracts were prepared as described for *E. coli* (37), except samples

were incubated at 50°C for 3 min instead of boiling for 3 min (56), and proteins were subjected

to immunoblot analysis (Supplementary File 3).

To image GFPΔ27BofA localization, samples collected at 3 h PS were examined by

fluorescence microscopy using an Olympus FluoView FV-1000 filter-based confocal

microscope. GFPΔ27BofA (ex/em ~488/507 nm) was excited using a 458 nM argon laser and

fluorescence was captured using a BA465-495 nm band pass filter. The lipophilic dye FM 4-64

(1 µg/mL) (AAT Bioquest) was used to stain membranes. FM 4-64 (ex/em ~515/640 nm) was

excited using a 515 nm argon laser and fluorescence was captured using a BA560IF band pass

filter (90).

Disulfide Cross-Linking. A method described previously (91) was used with slight

modifications (55). Briefly, as described above for Pro-σ^K(1-127) cleavage, *E. coli* BL21(DE3)

was transformed with a plasmid, grown in LB (10 mL), induced with IPTG, and equivalent

amounts of cells were collected. Cells were subjected to disulfide cross-linking, then lysed and

proteins were precipitated by addition of trichloroacetic acid (5%), and proteins were subjected

to immunoblot analysis (Supplementary File 3).

Co-Immunoprecipitation (FLAG₂ Pull-Down Assays). *E. coli* BL21(DE3) was transformed

with a plasmid, grown in LB (1 L), and induced with IPTG as described above. The culture was

split, cells were harvested, and cell pellets were stored at -80°C. Cell lysates were prepared as

described (65), except that each cell pellet was resuspended in 20 mL of lysis buffer containing

50 mM Tris-HCl pH 7.1 rather than PBS. Cell lysates were centrifuged (15,000 × *g* for 15 min

at 4 °C) to sediment cell debris and protein inclusion bodies. The supernatant was treated with

1% *n*-dodecyl-β-D-maltoside (DDM) (Anatrace) for 1 h at 4 °C to solubilize membrane proteins,

then centrifuged at 150,000 × *g* for 1 h at 4 °C. The supernatant was designated the input sample

and 1 mL was mixed with 50 uL anti-DYKDDDDK magnetic agarose (Pierce) that had been

equilibrated with buffer (50 mM Tris-HCl pH 7.1, 0.1% DDM, 5 mM 2-mercaptoethanol, 10%

glycerol) and the mixture was rotated for 1 h at 25°C. The magnetic agarose was removed with a

DynaMag-2 magnet (Invitrogen) and the supernatant was saved (unbound sample). The

magnetic agarose was washed three times by gently vortexing with 500 μL wash buffer (50 mM

Tris-HCl pH 7.1, 150 mM NaCl, 10% glycerol, 0.1% DDM), then washed once with 500 μL

water. The magnetic agarose was mixed with 50 μL of 2× sample buffer (50 mM Tris·HCl pH

6.8, 4% SDS, 20% glycerol, 200 mM DTT 0.03% bromophenol blue) and boiled for 3 min

(bound sample). A portion of the bound sample was diluted tenfold (1/10 bound sample) with

1× sample buffer to match the concentration of the input sample. Samples were subjected to

immunoblot analysis (Supplementary File 3).

Colbalt Affinity Purification (His$_6$ Pull-Down Assays). Input sample (15 mL) prepared as

described above was mixed with imidazole (5 mM) and 0.5 mL of Talon superflow metal affinity

resin (Clontech) that had been equilibrated with buffer (as above for magnetic agarose). The

mixture was rotated for 1 h at 4°C. The cobalt resin was sedimented by centrifugation at 708 × *g*

for 2 min at 4 °C and the supernatant was saved (unbound sample). The resin was washed three

times with 5 mL wash buffer (as above plus 5 mM imidazole), each time rotating the mixture for

10 min at 4 °C and sedimenting resin as above. The resin was mixed with 0.5 mL 2× sample

buffer and boiled for 3 min (bound sample). A portion of the bound sample was diluted

fifteenfold (1/15 bound sample) with $1\times$ sample buffer to match the concentration of the input

sample. Samples were subjected to immunoblot analysis (Supplementary File 3).

Modeling of Complexes Containing SpoIVFB, BofA, and Parts of SpoIVFA and Pro-σ^K.

The modeling proceeded through stages where initial monomeric models were assembled step-

by-step into multimeric complexes guided primarily by the restraints from cross-linking

experiments and predicted contacts from co-evolutionary coupling analysis. More specifically, a

SpoIVFB monomer was first assembled from the membrane and CBS domains. Two monomers

were combined into a dimer and the dimer was assembled into a plausible tetramer. Part of one

molecule of Pro-σ^K (residues 1-114) was subsequently added, which resulted in 'fb.sigk'. The

'fb.sigk.bofa.fa' model was developed by starting from 'fb.sigk', truncating the Pro-σ^K N-

terminus (residues 1-37), and subsequently adding first BofA and finally SpoIVFA (residues 65-

111). See Supplementary File 3 for additional description of the modeling.

Acknowledgments

We thank Daniel Parrell for assistance with fluorescence microscopy, Jon Kaguni for helpful

comments on the manuscript, and David Rudner, Simon Cutting, Ruanbao Zhou, and Yang

Zhang for plasmids. This research was supported by National Institutes of Health Grants R01

GM43585 (to L.K.) and R35 GM126948 (to M.F.), and by Michigan State University

AgBioResearch.

References

1.   Brown MS, Ye J, Rawson RB, & Goldstein JL (2000) Regulated intramembrane

proteolysis: a control mechanism conserved from bacteria to humans. *Cell* 100(4):391-398.

2.   Urban S (2013) Mechanisms and cellular functions of intramembrane proteases. *Biochim.*

Biophys. Acta - Biomembr. 1828:2797-2800.

3.   Kuhnle N, Dederer V, & Lemberg MK (2019) Intramembrane proteolysis at a glance: from

signalling to protein degradation. *J. Cell Sci.* 132(16):jcs217745.

4.   Manolaridis I*, et al.* (2013) Mechanism of farnesylated CAAX protein processing by the

intramembrane protease Rce1. *Nature* 504(7479):301-305.

5.   Hu J, Xue Y, Lee S, & Ha Y (2011) The crystal structure of GXGD membrane protease

FlaK. *Nature* 475(7357):528-531.

6.   Li X*, et al.* (2013) Structure of a presenilin family intramembrane aspartate protease. *Nature*

493(7430):56-61.

7.   Feng L*, et al.* (2007) Structure of a site-2 protease family intramembrane metalloprotease.

Science 318(5856):1608-1612.

8.   Wang Y, Zhang Y, & Ha Y (2006) Crystal structure of a rhomboid family intramembrane

protease. *Nature* 444(7116):179-183.

9.   Wu Z*, et al.* (2006) Structural analysis of a rhomboid family intramembrane protease

reveals a gating mechanism for substrate entry. *Nat. Struct. Mol. Biol.* 13(12):1084-1091.

10.   Cho S, Dickey SW, & Urban S (2016) Crystal structures and inhibition kinetics reveal a

two-stage catalytic mechanism with drug design implications for rhomboid proteolysis. *Mol.*

Cell 61(3):329-340.

11. Cho S, Baker RP, Ji M, & Urban S (2019) Ten catalytic snapshots of rhomboid

intramembrane proteolysis from gate opening to peptide release. *Nat. Struct. Mol. Biol.*

26(10):910-918.

12. Ticha A*, et al.* (2017) General and modular strategy for designing potent, selective, and

pharmacologically compliant inhibitors of rhomboid proteases. *Cell Chem. Biol.*

24(12):1523-1536 e1524.

13. Yang G*, et al.* (2019) Structural basis of Notch recognition by human γ-secretase. *Nature*

565(7738):192-197.

14. Zhou R*, et al.* (2019) Recognition of the amyloid precursor protein by human γ-secretase.

Science 363(6428):eaaw0930.

15. Rawson RB (2013) The site-2 protease. *Biochim. Biophys. Acta - Biomembr.*

1828(12):2801-2807.

16. Ye J (2013) Roles of regulated intramembrane proteolysis in virus infection and antiviral

immunity. *Biochim. Biophys. Acta - Biomembr.* 1828(12):2926-2932.

17. Adam Z (2013) Emerging roles for diverse intramembrane proteases in plant biology.

Biochim. Biophys. Acta - Biomembr. 1828(12):2933-2936.

18. Chang YC, Bien CM, Lee H, Espenshade PJ, & Kwon-Chung KJ (2007) Sre1p, a regulator

of oxygen sensing and sterol homeostasis, is required for virulence in *Cryptococcus*

neoformans. *Mol. Microbiol.* 64(3):614-629.

19. Urban S (2009) Making the cut: central roles of intramembrane proteolysis in pathogenic

microorganisms. *Nat. Rev. Microbiol.* 7:411-423.

20. Kroos L & Akiyama Y (2013) Biochemical and structural insights into intramembrane

metalloprotease mechanisms. *Biochim. Biophys. Acta - Biomembr.* 1828(12):2873-2885.

21. Schneider JS & Glickman MS (2013) Function of site-2 proteases in bacteria and bacterial

pathogens. *Biochim. Biophys. Acta - Biomembr.* 1828(12):2808-2814.

22. de Hoon MJ, Eichenberger P, & Vitkup D (2010) Hierarchical evolution of the bacterial

sporulation network. *Curr. Biol.* 20(17):R735-R745.

23. Galperin MY*, et al.* (2012) Genomic determinants of sporulation in *Bacilli* and *Clostridia*:

towards the minimal set of sporulation-specific genes. *Environ. Microbiol.* 14(11):2870-

2890.

24. Abecasis AB*, et al.* (2013) A genomic signature and the identification of new sporulation

genes. *J. Bacteriol.* 195(9):2101-2115.

25. Ramos-Silva P, Serrano M, & Henriques AO (2019) From root to tips: sporulation evolution

and specialization in *Bacillus subtilis* and the intestinal pathogen *Clostridioides difficile*.

Mol. Biol. Evol. 36(12):2714-2736.

26. McKenney PT, Driks A, & Eichenberger P (2013) The *Bacillus subtilis* endospore:

assembly and functions of the multilayered coat. *Nat. Rev. Microbiol.* 11(1):33-44.

27. Al-Hinai MA, Jones SW, & Papoutsakis ET (2015) The *Clostridium* sporulation programs:

diversity and preservation of endospore differentiation. *Microbiol. Mol. Biol. Rev.* 79(1):19-

37.

28. Checinska A, Paszczynski A, & Burbank M (2015) *Bacillus* and other spore-forming

genera: variations in responses and mechanisms for survival. *Annu. Rev. Food Sci. Technol.*

6:351-369.

29. Browne HP*, et al.* (2016) Culturing of 'unculturable' human microbiota reveals novel taxa

and extensive sporulation. *Nature* 533(7604):543-546.

30. Tan IS & Ramamurthi KS (2014) Spore formation in *Bacillus subtilis*. *Environ. Microbiol.*

Rep. 6(3):212-225.

31. Cutting S*, et al.* (1990) A forespore checkpoint for mother-cell gene expression during

development in *Bacillus subtilis*. *Cell* 62:239-250.

32. Cutting S, Roels S, & Losick R (1991) Sporulation operon *spoIVF* and the characterization

of mutations that uncouple mother-cell from forespore gene expression in *Bacillus subtilis*.

J. Mol. Biol. 221:1237-1256.

33. Ricca E, Cutting S, & Losick R (1992) Characterization of *bofA*, a gene involved in

intercompartmental regulation of pro-σ^K processing during sporulation in *Bacillus subtilis*.

J. Bacteriol. 174:3177-3184.

34. Resnekov O, Alper S, & Losick R (1996) Subcellular localization of proteins governing the

proteolytic activation of a developmental transcription factor in *Bacillus subtilis*. *Genes*

Cells 1(6):529-542.

35. Rudner DZ, Pan Q, & Losick RM (2002) Evidence that subcellular localization of a

bacterial membrane protein is achieved by diffusion and capture. *Proc. Natl. Acad. Sci. USA*

99(13):8701-8706.

36. Rudner DZ & Losick R (2002) A sporulation membrane protein tethers the pro-σ^K

processing enzyme to its inhibitor and dictates its subcellular localization. *Genes Dev.*

16(8):1007-1018.

37. Zhou R & Kroos L (2004) BofA protein inhibits intramembrane proteolysis of pro-σ^K in an

intercompartmental signaling pathway during *Bacillus subtilis* sporulation. *Proc. Natl.*

Acad. Sci. USA 101(17):6385-6390.

38. Kroos L, Yu YT, Mills D, & Ferguson-Miller S (2002) Forespore signaling is necessary for

pro-σ^K processing during *Bacillus subtilis* sporulation despite the loss of SpoIVFA upon

translational arrest. *J. Bacteriol.* 184(19):5393-5401.

39. Cutting S, Driks A, Schmidt R, Kunkel B, & Losick R (1991) Forespore-specific

transcription of a gene in the signal transduction pathway that governs pro-σ^K processing in

Bacillus subtilis. *Genes Dev.* 5:456-466.

40. Pan Q, Losick R, & Rudner DZ (2003) A second PDZ-containing serine protease

contributes to activation of the sporulation transcription factor σ^K in *Bacillus subtilis*. *J.*

Bacteriol. 185(20):6051-6056.

41. Dong TC & Cutting SM (2003) SpoIVB-mediated cleavage of SpoIVFA could provide the

intercellular signal to activate processing of Pro-σ^K in *Bacillus subtilis*. *Mol. Microbiol.*

49(5):1425-1434.

42. Campo N & Rudner DZ (2006) A branched pathway governing the activation of a

developmental transcription factor by regulated intramembrane proteolysis. *Mol. Cell*

23(1):25-35.

43. Campo N & Rudner DZ (2007) SpoIVB and CtpB are both forespore signals in the

activation of the sporulation transcription factor σ^K in *Bacillus subtilis*. *J. Bacteriol.*

189(16):6021-6027.

44. Mastny M*, et al.* (2013) CtpB assembles a gated protease tunnel regulating cell-cell

signaling during spore formation in *Bacillus subtilis*. *Cell* 155(3):647-658.

45. Zhou R & Kroos L (2005) Serine proteases from two cell types target different components

of a complex that governs regulated intramembrane proteolysis of pro-σ^K during *Bacillus*

subtilis development. *Mol. Microbiol.* 58(3):835-846.

46. Rudner D, Fawcett P, & Losick R (1999) A family of membrane-embedded

metalloproteases involved in regulated proteolysis of membrane-associated transcription

factors. *Proc. Natl. Acad. Sci. USA* 96:14765-14770.

47. Yu Y-TN & Kroos L (2000) Evidence that SpoIVFB is a novel type of membrane

metalloprotease governing intercompartmental communication during *Bacillus subtilis*

sporulation. *J. Bacteriol.* 182:3305-3309.

48. Zhou R, Cusumano C, Sui D, Garavito RM, & Kroos L (2009) Intramembrane proteolytic

cleavage of a membrane-tethered transcription factor by a metalloprotease depends on ATP.

Proc. Natl. Acad. Sci. USA 106:16174-16179.

49. Kroos L, Kunkel B, & Losick R (1989) Switch protein alters specificity of RNA polymerase

containing a compartment-specific sigma factor. *Science* 243:526-529.

50. Eichenberger P*, et al.* (2004) The program of gene transcription for a single differentiating

cell type during sporulation in *Bacillus subtilis*. *PLoS Biol.* 2(10):e328.

51. Morohashi Y & Tomita T (2013) Protein trafficking and maturation regulate intramembrane

proteolysis. *Biochim. Biophys. Acta* 1828(12):2855-2861.

52. Lichtenthaler SF, Lemberg MK, & Fluhrer R (2018) Proteolytic ectodomain shedding of

membrane proteins in mammals-hardware, concepts, and recent developments. *EMBO J.*

37(15):e99456.

53. Beard HA, Barniol-Xicota M, Yang J, & Verhelst SHL (2019) Discovery of cellular roles of

intramembrane proteases. *ACS Chem. Biol.* 14(11):2372-2388.

54. Kinch LN, Ginalski K, & Grishin NV (2006) Site-2 protease regulated intramembrane

proteolysis: sequence homologs suggest an ancient signaling cascade. *Protein Sci.* 15(1):84-

93.

55. Zhang Y, Luethy PM, Zhou R, & Kroos L (2013) Residues in conserved loops of

intramembrane metalloprotease SpoIVFB interact with residues near the cleavage site in

Pro-σ^K. *J. Bacteriol.* 195(21):4936-4946.

56. Halder S, Parrell D, Whitten D, Feig M, & Kroos L (2017) Interaction of intramembrane

metalloprotease SpoIVFB with substrate Pro-σ^K. *Proc. Natl. Acad. Sci. USA* 114:E10677-

E10686.

57. Saribas AS, Gruenke L, & Waskell L (2001) Overexpression and purification of the

membrane-bound cytochrome P450 2B4. *Protein Expr. Purif.* 21(2):303-309.

58. Prince H, Zhou R, & Kroos L (2005) Substrate requirements for regulated intramembrane

proteolysis of *Bacillus subtilis* pro-σ^K. *J. Bacteriol.* 187:961-971.

59. Ramirez-Guadiana FH, *et al.* (2018) Evidence that regulation of intramembrane proteolysis

is mediated by substrate gating during sporulation in *Bacillus subtilis*. *PLoS Genet.*

14(11):e1007753.

60. Varcamonti M, Marasco R, De Felice M, & Sacco M (1997) Membrane topology analysis

of the *Bacillus subtilis* BofA protein involved in pro-σ^K processing. *Microbiol.* 143(Pt

4):1053-1058.

61. Resnekov O (1999) Role of the sporulation protein BofA in regulating activation of the

Bacillus subtilis developmental transcription factor σ^K. *J. Bacteriol.* 181:5384-5388.

62. Arolas JL, Goulas T, Cuppari A, & Gomis-Ruth FX (2018) Multiple architectures and

mechanisms of latency in metallopeptidase zymogens. *Chem. Rev.* 118(11):5581-5597.

63. Van Wart HE & Birkedal-Hansen H (1990) The cysteine switch: a principle of regulation of

metalloproteinase activity with potential applicability to the entire matrix metalloproteinase

gene family. *Proc. Natl. Acad. Sci. USA* 87(14):5578-5582.

64. Odintsov SG, Sabala I, Marcyjaniak M, & Bochtler M (2004) Latent LytM at 1.3 Å

resolution. *J. Mol. Biol.* 335(3):775-785.

65. Zhang Y*, et al.* (2016) Complex formed between intramembrane metalloprotease SpoIVFB

and its substrate, Pro-σ^K. *J. Biol. Chem.* 291:10347-10362.

66. Akiyama K*, et al.* (2015) Roles of the membrane-reentrant β-hairpin-like loop of RseP

protease in selective substrate cleavage. *eLife* 4:e08928

67. Rawson RB (2003) The SREBP pathway--insights from Insigs and insects. *Nat. Rev. Mol.*

Cell Biol. 4(8):631-640.

68. Lima S, Guo MS, Chaba R, Gross CA, & Sauer RT (2013) Dual molecular signals mediate

the bacterial response to outer-membrane stress. *Science* 340(6134):837-841.

69. Hizukuri Y*, et al.* (2014) A structure-based model of substrate discrimination by a

noncanonical PDZ tandem in the intramembrane-cleaving protease RseP. *Structure*

22(2):326-336.

70. Bolduc DM, Montagna DR, Gu YL, Selkoe DJ, & Wolfe MS (2016) Nicastrin functions to

sterically hinder γ-secretase-substrate interactions driven by substrate transmembrane

domain. *Proc. Natl. Acad. Sci. USA* 113(5):E509-E518.

71. Fukumori A & Steiner H (2016) Substrate recruitment of γ-secretase and mechanism of

clinical presenilin mutations revealed by photoaffinity mapping. *EMBO J.* 35(15):1628-

1643.

72. Pereira FC*, et al.* (2013) The spore differentiation pathway in the enteric pathogen

Clostridium difficile. *PLoS Genet.* 9(10):e1003782.

73. Saujet L*, et al.* (2013) Genome-wide analysis of cell type-specific gene transcription during

spore formation in *Clostridium difficile*. *PLoS Genet.* 9(10):e1003756.

74. Haraldsen JD & Sonenshein AL (2003) Efficient sporulation in *Clostridium difficile*

requires disruption of the σ^K gene. *Mol. Microbiol.* 48(3):811-821.

75. Ticha A, Collis B, & Strisovsky K (2018) The rhomboid superfamily: structural

mechanisms and chemical biology opportunities. *Trends Biochem. Sci.* 43(9):726-739.

76. Wolfe MS (2019) Dysfunctional γ-secretase in familial Alzheimer's disease. *Neurochem Res*

44(1):5-11.

77. De Strooper B & Chavez Gutierrez L (2015) Learning by failing: ideas and concepts to

tackle γ-secretases in Alzheimer's disease and beyond. *Annu. Rev. Pharmacol. Toxicol.*

55:419-437.

78. Szaruga M, *et al.* (2017) Alzheimer's-causing mutations shift Aβ length by destabilizing γ-

secretase-Aβn interactions. *Cell* 170(3):443-456 e414.

79. Konovalova A, *et al.* (2018) Inhibitor of intramembrane protease RseP blocks the σ^E

response causing lethal accumulation of unfolded outer membrane proteins. *Proc. Natl.*

Acad. Sci. USA 115(28):E6614-E6621.

80. Jacobsen JA, Major Jourden JL, Miller MT, & Cohen SM (2010) To bind zinc or not to bind

zinc: an examination of innovative approaches to improved metalloproteinase inhibition.

Biochim. Biophys. Acta 1803(1):72-94.

81. Fields GB (2015) New strategies for targeting matrix metalloproteinases. *Matrix Biol.* 44-

46:239-246.

82. Gomis-Ruth FX (2017) Third time lucky? Getting a grip on matrix metalloproteinases. *J.*

Biol. Chem. 292(43):17975-17976.

83. Moss ML*, et al.* (2007) The ADAM10 prodomain is a specific inhibitor of ADAM10

proteolytic activity and inhibits cellular shedding events. *J. Biol. Chem.* 282(49):35712-

35721.

84. Wong E*, et al.* (2016) Harnessing the natural inhibitory domain to control TNFα Converting

Enzyme (TACE) activity *in vivo*. *Sci. Rep.* 6:35598.

85. Zorzi A, Deyle K, & Heinis C (2017) Cyclic peptide therapeutics: past, present and future.

Curr. Opin. Chem. Biol. 38:24-29.

86. Sohrabi C, Foster A, & Tavassoli A (2020) Methods for generating and screening libraries

of genetically encoded cyclic peptides in drug discovery. *Nat. Rev. Chem.* 4(2):90-101.

87. Hanahan D (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol.*

Biol. 166:557-580.

88. Notredame C, Higgins DG, & Heringa J (2000) T-Coffee: A novel method for fast and

accurate multiple sequence alignment. *J. Mol. Biol.* 302(1):205-217.

89. Harwood CR & Cutting SM (1990) *Molecular Biological Methods for Bacillus* (John Wiley

& Sons, Chichester, England) p 581.

90. Parrell D & Kroos L (2020) Channels modestly impact compartment-specific ATP levels

during *Bacillus subtilis* sporulation and a rise in the mother cell ATP level is not necessary

for Pro-σ^K cleavage. *Mol. Microbiol.* 114:563-581.

91. Koide K, Ito K, & Akiyama Y (2008) Substrate recognition and binding by RseP, an

Escherichia coli intramembrane protease. *J. Biol. Chem.* 283(15):9562-9570.

92. Ha Y (2009) Structure and mechanism of intramembrane protease. *Semin. Cell Dev. Biol.*

20(2):240-250.

93. Zelenski N, Rawson R, Brown M, & Goldstein J (1999) Membrane topology of S2P, a

protein required for intramembranous cleavage of sterol regulatory element-binding

proteins. *J. Biol. Chem.* 274:21973-21980.

94. Nohturfft A, DeBose-Boyd RA, Scheek S, Goldstein JL, & Brown MS (1999) Sterols

regulate cycling of SREBP cleavage-activating protein (SCAP) between endoplasmic

reticulum and Golgi. *Proc. Natl. Acad. Sci. USA* 96(20):11235-11240.

95. DeBose-Boyd RA*, et al.* (1999) Transport-dependent proteolysis of SREBP: relocation of

site-1 protease from Golgi to ER obviates the need for SREBP transport to Golgi. *Cell*

99(7):703-712.

96. Chen G & Zhang X (2010) New insights into S2P signaling cascades: regulation, variation,

and conservation. *Protein Sci.* 19(11):2015-2030.

97. Sakai J*, et al.* (1996) Sterol-regulated release of SREBP-2 from cell membrane requires two

sequential cleavages, one within a transmembrane domain. *Cell* 85:1037-1048.

98. Sakai J*, et al.* (1997) Identification of complexes between the COOH-terminal domains of

sterol regulatory element-binding proteins (SREBPs) and SREBP cleavage-activating

protein. *J. Biol. Chem.* 272(32):20213-20221.

99. Horton JD*, et al.* (2003) Combined analysis of oligonucleotide microarray data from

transgenic and knockout mice identifies direct SREBP target genes. *Proc. Natl. Acad. Sci.*

USA 100(21):12027-12032.

100. Kanehara K, Akiyama Y, & Ito K (2001) Characterization of the *yaeL* gene product and its

S2P-protease motifs in *Escherichia coli*. *Gene* 281(1-2):71-79.

101. Drew D*, et al.* (2002) Rapid topology mapping of *Escherichia coli* inner-membrane proteins

by prediction and PhoA/GFP fusion analysis. *Proc. Natl. Acad. Sci. USA* 99(5):2690-2695.

102. Ades SE, Connolly LE, Alba BM, & Gross CA (1999) The *Escherichia coli* σ^E-dependent

extracytoplasmic stress response is controlled by the regulated proteolysis of an anti-σ

factor. *Genes Dev.* 13(18):2449-2461.

103. Walsh NP, Alba BM, Bose B, Gross CA, & Sauer RT (2003) OMP peptide signals initiate

the envelope-stress response by activating DegS protease via relief of inhibition mediated by

its PDZ domain. *Cell* 113(1):61-71.

104. Kanehara K, Ito K, & Akiyama Y (2002) YaeL (EcfE) activates the σ^E pathway of stress

response through a site-2 cleavage of anti-σ^E, RseA. *Genes Dev.* 16(16):2147-2155.

105. Alba BM, Leeds JA, Onufryk C, Lu CZ, & Gross CA (2002) DegS and YaeL participate

sequentially in the cleavage of RseA to activate the σ^E-dependent extracytoplasmic stress

response. *Genes Dev.* 16(16):2156-2168.

106. Flynn JM, Neher SB, Kim YI, Sauer RT, & Baker TA (2003) Proteomic discovery of

cellular substrates of the ClpXP protease reveals five classes of ClpX-recognition signals.

Mol. Cell 11(3):671-683.

107. Chaba R, Grigorova IL, Flynn JM, Baker TA, & Gross CA (2007) Design principles of the

proteolytic cascade governing the σ^E-mediated envelope stress response in *Escherichia coli*:

keys to graded, buffered, and rapid signal transduction. *Genes Dev.* 21(1):124-136.

  Figure 1 with 4 supplements
952

953
954
Inhibition of Pro-σ^K cleavage.
(**A**) Diagram of SpoIVFB inhibition during *B. subtilis* endosporulation and upon heterologous
expression in *E. coli*. During endosporulation (*Left*), SpoIVFB and its inhibitory proteins BofA
and SpoIVFA are produced in the mother cell (MC) and localize to the outer forespore (FS)
membrane (OFM). Pro-σ^K is also produced in the MC and associates with membranes. When
these proteins are synthesized in *E. coli* (*Right*), they localize to the inner membrane (IM). The
expanded view of the membranes (*Center*) shows a SpoIVFB variant with an extra N-terminal
transmembrane segment (cytTM), and highlights several residues (E44, V70, P135) at or near the
active site in the membrane domain, which is connected to the CBS domain by an interdomain

linker. When produced in *E. coli*, cytTM-SpoIVFB cleaves Pro-σ^K(1-127), removing its N-
terminal Proregion [Note: Pro-σ^K(1-126) was renamed Pro-σ^K(1-127) as explained in (56)]. Co-
production of SpoIVFA and either full-length BofA or GFPΔ27BofA (lacking predicted TMS1)
inhibits Pro-σ^K(1-127) cleavage. The dashed double-headed arrows indicate that SpoIVFB,
BofA or GFPΔ27BofA, and SpoIVFA form a complex of unknown structure. (**B**) Cleavage
assays comparing inhibition by SpoIVFA and either GFPΔ27BofA or full-length BofA in *E. coli*.
Pro-σ^K(1-127) and cytTM-SpoIVFB were produced alone (lane 1, pYZ2) or in combination with
GFPΔ27BofA and SpoIVFA (lane 2, pYZ46) or full-length BofA and SpoIVFA (lane 3,
pSO212). Alternatively, "pET Quartet" plasmids were used to produce Pro-σ^K(1-127), cytTM-
SpoIVFB, SpoIVFA, and either GFPΔ27BofA (lane 4, pSO40) or full-length BofA (lane 5,
pSO213). Samples collected after 2 h of IPTG induction were subjected to immunoblot analysis
with SpoIVFA (*Top*), GFP (*Middle*), or penta-His antibodies (*Bottom*, 2 and 30 s exposures).
The single star (*) indicates cross-reacting proteins below SpoIVFA and the double star (**)
indicates breakdown species of GFPΔ27BofA. A breakdown species below SpoIVFA (not
indicated) is observed in some experiments. The graph shows quantification of the cleavage
ratio (cleavage product/[Pro-σ^K(1-127) + cleavage product]) for three biological replicates. Error
bars, 1 standard deviation. (**C**) Cleavage assays comparing inhibition by either GFPΔ27BofA or
SpoIVFA. pET Triplet plasmids were used to produce Pro-σ^K(1-127), cytTM-SpoIVFB, and
either GFPΔ27BofA (lane 2, pSO64) or SpoIVFA (lane 3, pSO65). Samples were subjected to
immunoblot analysis and quantification as in (**B**).
984
985   **Figure 1-source data 1**
986   **Immunoblot images (raw and annotated) and quantification of cleavage assays (Figure 1 B**
987   **and C).**

     Figure 2 with 2 supplements

990
Effects of alanine substitutions in GFPΔ27BofA on inhibition of Pro-σ^K(1-127) cleavage
in *E. coli*.
pET Quartet plasmids were used to produce Pro-σ^K(1-127), cytTM-SpoIVFB, SpoIVFA, and
GFPΔ27BofA (lane 1, pSO40) or Ala-substituted GFPΔ27BofA (lanes 2-20, pSO44-pSO58 and
pSO60-pSO63). Samples collected after 2 h of IPTG induction were subjected to immunoblot
analysis with SpoIVFA, GFP, and penta-His antibodies. Single (*) and double (**) stars are
explained in the Figure 1B legend, as is the graph.
998
999   **Figure 2-source data 1**
1000   **Immunoblot images (raw and annotated) and quantification of cleavage assays.**

Figure 3 with 1 supplement

Effects of alanine substitutions in GFPΔ27BofA during *B. subtilis* sporulation.
(**A**) Effects of GFPΔ27BofA variants (N48A, N61A, and T64A) on Pro-σ^K cleavage. Wild-type
strain PY79, a *spoIVB165* null mutant, a *spoIVB165 bofA::erm* double mutant, and the double
mutant with P*bofA*-*gfpΔ27bofA* integrated ectopically at *amyE* to express GFPΔ27BofA with no
substitution (none) or the indicated Ala substitution, were starved to induce sporulation.
Samples collected at 4 and 5 h poststarvation (PS) were subjected to immunoblot analysis with
antibodies against SpoIVFA, GFP, SpoIVFB, and Pro-σ^K. The graph shows quantification of the
cleavage ratio [σ^K/(Pro-σ^K + σ^K)] for three biological replicates. Error bars, 1 standard deviation.
Student's two-tailed *t* tests were performed to compare certain cleavage ratios (*p* values). (**B**)
Localization of GFPΔ27BofA and the three variants. Samples collected at 3 h PS were treated
with FM 4-64 to stain membranes. Confocal microscopy images of fluorescence from

1047 GFPΔ27BofA, membranes, and merged images are shown for representative sporangia with
1048 discrete (no substitution in GFPΔ27BofA, designated "none"), partial (N48A and T64A), or no
1049 forespore (FS) localization (N61A). Scale bar, 1 μm. The percentage of sporangia (44-93
1050 counted; non-sporulating cells were not counted) with each localization pattern is shown.
1051
1052 **Figure 3-source data 1**
1053 **Immunoblot images (annotated) and quantification of cleavage assays (Figure 3A), and**
1054 **confocal microscopy images (see the readme file) and a table of the numbers of sporangia**
1055 **in which FS localization of GFPΔ27BofA and the three variants was counted (Figure 3B).**
1056
1057 **Figure 3-source data 2**
1058 **Immunoblot images (raw) (Figure 3A).**

1059   **Figure 4** with 5 supplements

1060
BofA TMS2 is near the SpoIVFB active site.
(**A**) Disulfide cross-linking of E44C at the cytTM-SpoIVFB active site to C46 in TMS2 of
MBPΔ27BofA. pET Quartet plasmids were used to produce single-Cys E44C cytTM-SpoIVFB
in combination with MBPΔ27BofA C46 (pSO91) or Cys-less MBPΔ27BofA C46S as a negative
control (pSO110), and Cys-less variants of SpoIVFA and Pro-σ^K(1-127) in *E. coli*. Samples
collected after 2 h of IPTG induction were treated for 60 min with Cu^{2+}(phenanthroline)$_3$ (Cu +)
to promote disulfide bond formation or with 2-phenanthroline (Cu –) as a negative control, then
treated with TCA to precipitate proteins and resuspended in sample buffer with DTT (+) to
reverse cross-links or without (–) to preserve cross-links, and finally subjected to immunoblot
analysis with FLAG antibodies to visualize cytTM-SpoIVFB monomer, dimer, and complex
with MBPΔ27BofA. (**B**) Disulfide cross-linking of V70C or P135C near the cytTM-SpoIVFB
active site to C46 in TMS2 MBPΔ27BofA. pET Quartet plasmids were used to produce single-
Cys V70C or P135C cytTM-SpoIVFB E44Q variants in combination with MBPΔ27BofA
(pSO92, pSO93) or Cys-less MBPΔ27BofA C46S as a negative control (pSO111, pSO112), and
Cys-less variants of SpoIVFA and Pro-σ^K(1-127) in *E. coli*. Samples collected after 2 h of IPTG
induction were treated and subjected to immunoblot analysis as in (**A**). A representative result
from at least two biological replicates is shown in (**A**) and (**B**).
1078
1079   **Figure 4-source data 1**
1080   **Immunoblot images (raw and annotated).**
1081

1082 **Figure 5** with 3 supplements

1083
Inhibitory proteins do not prevent Pro-σ^K(1-127) from interacting with SpoIVFB.
(**A**) Pro-σ^K(1-127), SpoIVFA, and GFPΔ27BofA co-purify with cytTM-SpoIVFB. pET Quartet
plasmids were used to produce a catalytically-inactive E44C cytTM-SpoIVFB variant with a
FLAG$_2$ tag (pSO73), or a variant lacking FLAG$_2$ as a negative control (pSO149), in combination
with Pro-σ^K(1-127), SpoIVFA, and GFPΔ27BofA in *E. coli*. Samples collected after 2 h of
IPTG induction were subjected to co-immunoprecipitation with anti-FLAG antibody beads.
Input, unbound, 1/10 bound (diluted to match input), and (undiluted) bound samples were
subjected to immunoblot analysis with FLAG, penta-His, SpoIVFA, and GFP antibodies. The
single star (*) indicates cross-reacting proteins below SpoIVFA and the double star (**)
indicates a cross-reacting protein or breakdown species of GFPΔ27BofA that fail to co-purify. A
representative result from two biological replicates is shown. (**B**) A231C in the cytTM-SpoIVFB
CBS domain forms disulfide cross-links with A97C in the C-terminal region of Pro-σ^K(1-127) in
the absence or presence of inhibitory proteins. pET Duet plasmids were used to produce single-
Cys A231C cytTM-SpoIVFB E44Q in combination with single-Cys A97C Pro-σ^K(1-127)
(pSO130; squares and line with long dashes labeled "No BofA", although SpoIVFA is also
absent) or with Cys-less Pro-σ^K(1-127) as a negative control (pSO255; crosses and line labeled
"Cys-less") in *E. coli*. pET Quartet plasmids were used to produce single-Cys A231C cytTM-
SpoIVFB E44Q, single-Cys A97C Pro-σ^K(1-127), and Cys-less SpoIVFA in combination with
Cys-less MBPΔ27BofA (pSO133; circles and line labeled "MBPΔ27BofA") or with Cys-less
full-length BofA (pSO246; diamonds and line labeled "BofA") in *E. coli*. Samples collected
after 2 h of IPTG induction were treated for 15, 30, 45, or 60 min with Cu^{2+}(phenanthroline)$_3$
oxidant to promote disulfide bond formation and subjected to immunoblot analysis with FLAG
antibodies to visualize the cytTM-SpoIVFB monomer, dimer, and complex with Pro-σ^K(1-127)
(Figure 5-figure supplement 3C). Abundance of the complex was divided by the total amount of
cytTM-SpoIVFB monomer, dimer, and complex. The ratio over time was plotted (n=2) with a
best-fit trend line.
1110
1111 **Figure 5-source data 1**
1112 **Immunoblot images (raw and annotated) (Figure 5A) and quantification of cross-linking**
1113 **(Figure 5B).**

Figure 6 with 5 supplements

Inhibitory proteins block access of the Pro-σ^K(1-127) N-terminal region to the SpoIVFB active site and hinder interaction with the SpoIVFB membrane-reentrant loop and interdomain linker.

(**A**) E44C at the cytTM-SpoIVFB active site forms abundant disulfide cross-links with V20C or K24C in the N-terminal region of Pro-σ^K(1-127) in the absence of inhibitory proteins. pET Duet plasmids were used to produce single-Cys E44C cytTM-SpoIVFB in combination with single-Cys F18C (pSO167), V20C (pSO169), S21C (pSO170), or K24C (pSO128) Pro-σ^K(1-127), or with Cys-less Pro-σ^K(1-127) (pSO79) as a negative control, in *E. coli*. Samples collected after 2 h of IPTG induction were treated with Cu^{2+}(phenanthroline)$_3$ oxidant for 15, 30, 45, or 60 min to promote disulfide bond formation and subjected to immunoblot analysis with FLAG antibodies to visualize the cytTM-SpoIVFB monomer, dimer, and complex with Pro-σ^K(1-127) (Figure 6-figure supplement 1A). Abundance of the complex was divided by the total amount of cytTM-SpoIVFB monomer, dimer, and complex. The ratio over time was plotted (n=2) with a best-fit trend line. (**B**) Representative immunoblots of 60-min samples from the experiment described in (**A**). (**C**) MBPΔ27BofA and SpoIVFA decrease cross-linking between E44C cytTM-SpoIVFB and V20C or K24C Pro-σ^K(1-127). pET Quartet plasmids were used to produce single-Cys E44C cytTM-SpoIVFB in combination with single-Cys F18C (pSO163), V20C (pSO165), S21C (pSO166), or K24C (pSO131) Pro-σ^K(1-127), or with Cys-less Pro-σ^K(1-127) (pSO110) as a negative control, and Cys-less variants of MBPΔ27BofA and SpoIVFA in *E. coli*. Samples collected after 2 h of IPTG induction were treated and subjected to immunoblot analysis as in (**A**) (Figure 6-figure supplement 1B). The complex/total ratio was plotted as in (**A**). (**D**) Representative immunoblots of 60-min samples from the experiment described in (**C**). (**E** and **F**) Summaries of the effects of inhibitory proteins on cross-linking between E44C cytTM-SpoIVFB and V20C or K24C Pro-σ^K(1-127). Data from (**A**) (labeled "No BofA" in (**E**), although SpoIVFA is also absent), (**C**) (labeled "MBPΔ27BofA" in (**E**), although SpoIVFA is also present), and Figure 6-figure supplement 2 (labeled "BofA" in (**E**), although SpoIVFA is also present) are plotted along with Cys-less Pro-σ^K(1-127) as a negative control. In (**F**), symbols and lines are as in (**E**). (**G** and **H**) Summaries of the effects of inhibitory proteins on cross-linking between V70C in the cytTM-SpoIVFB membrane-reentrant loop and F18C or K24C in the Pro-σ^K(1-127) N-terminal region. Data from Figure 6-figure supplement 3 are plotted using symbols and lines as in (**E**). (**I**) Summary of the effects of inhibitory proteins on cross-linking between Y214C in the cytTM-SpoIVFB interdomain linker and L41C in the Pro-σ^K(1-127) N-terminal region. Data from Figure 6-figure supplement 5 are plotted using symbols and lines as in (**E**).

Figure 6-source data 1
Quantification of cross-linking (Figure 6 A, C, and E-I).

Figure 7 with 1 supplement

1155
Model of SpoIVFB monomer with BofA and parts of SpoIVFA and Pro-σ^K.
(**A**) Model of SpoIVFB, BofA TMS2, and the C-terminal part of Pro-σ^K(1-127). At *Left*, a side
view of the complex, showing the six TMSs of the SpoIVFB membrane domain with the active
site zinc ion (magenta), the interdomain linker, and the CBS domain (green), BofA TMS2
(cyan), and Pro-σ^K(38-114) (red). In the enlarged view of the active site cleft (*Center*), TMSs 1–
6 of SpoIVFB and TMS2 of BofA are numbered. At *Right*, a top view is shown. (**B**) Model of
SpoIVFB with full-length BofA and Pro-σ^K(38-114). Predicted TMSs 1 and 2 of BofA are
numbered and its C-terminal region is labeled "C" near the C-terminus in the views shown in the

Center and at *Right*. (**C**) Model of SpoIVFB with full-length BofA, SpoIVFA(65-111) (purple,
residue 111 is numbered), and Pro-σ^K(38-114).
1166
1167   **Figure 7-source data 1**
1168   **PyMOL session file used to produce the images and PDB file of the model of a SpoIVFB**
1169   **tetramer with BofA and parts of SpoIVFA and Pro-σ^K.**
1170

1171 **Figure 8**

1172

1173
Regulation of intramembrane proteolysis by SpoIVFB differs from that of S2P and RseP.
(**A**) S2P regulation. Human S2P localizes to membranes of the Golgi apparatus and has eight
predicted TMSs and a PDZ domain (20, 92, 93). SREBP substrates localize to endoplasmic
reticulum (ER) membranes *via* interaction of their regulatory domain (Reg) with SCAP, which is
bound by Insig proteins when the cholesterol level is high (15). Cholesterol depletion releases
SREBP-SCAP complexes from Insig and allows vesicular substrate transport to the Golgi
membranes (94-96). Cleavage (lightning bolt) by S1P at site-1 in the luminal loop of SREBPs
promotes separation of the TMSs and site-2 cleavage by S2P (97, 98). S2P cleavage releases the
basic helix-loop-helix (bHLH) domain of SREBP substrates into the cytosol, for subsequent
entry into the nucleus and transcriptional activation of genes required for fatty acid and
cholesterol synthesis and uptake (99). (**B**) RseP regulation. RseP localizes to the *E. coli* inner
membrane (IM) and has four predicted TMSs and tandem PDZ domains (20, 100, 101). The
substrate RseA also localizes to the inner membrane and sequesters σ^E in the absence of
envelope stress (102). The PDZ domains of RseP block access of the RseA/σ^E complex to the
RseP active site (69). Stress causes misfolding of outer membrane proteins and accumulation of
lipopolysaccharide intermediates, which trigger site-1 cleavage by DegS in the RseA periplasmic
domain (68, 103). Removal of the RseA extramembrane domain permits entry into the RseP
active site cleft for site-2 cleavage (104, 105). Release of RseA/σ^E into the cytosol allows ClpXP
protease to degrade RseA, liberating σ^E to direct transcription of stress-responsive genes (106,
107). (**C**) SpoIVFB regulation. SpoIVFB localizes to the outer forespore membrane (OFM)
during *B. subtilis* sporulation and has six predicted TMSs and a CBS domain (20, 32, 34, 54).

Our results support a novel regulatory mechanism in which the inhibitory proteins BofA and
SpoIVFA occupy the SpoIVFB active site cleft and block access of the Pro-σ^K Proregion. The
inhibitory proteins did not prevent interaction of Pro-σ^K with the SpoIVFB CBS domain in our *E.*
coli system. Proteases SpoIVB and CtpB are secreted from the forespore into the intermembrane
space and cleave the inhibitory proteins (41-43, 45), allowing the Pro-σ^K Proregion to enter the
SpoIVFB active site cleft for cleavage (46-48), which releases σ^K into the mother cell to direct
transcription of sporulation genes (49, 50).
1202

1203   **Figure Supplements**
1204
1205   **Figure 1-figure supplement 1**
1206   **Morphological changes during endosporulation and regulated intramembrane proteolysis**
1207   **of Pro-σ^K in *B. subtilis*.**
1208
1209   **Figure 1-figure supplement 2**
1210   **Full-length BofA alone fails to inhibit Pro-σ^K(1-127) cleavage in *E. coli* and full-length Pro-**
1211   **σ^K is similar to Pro-σ^K(1-127) in terms of requirements for cleavage inhibition.**
1212
1213   **Figure 1-figure supplement 3**
1214   **Cleavage assays examining the effects of SpoIVB production on BofA and SpoIVFA**
1215   **inhibition of SpoIVFB in *E. coli*.**
1216
1217   **Figure 1-figure supplement 4**
1218   **An F66A substitution in cytTM-SpoIVFB partially overcomes inhibition by GFPΔ27BofA**
1219   **and SpoIVFA in *E. coli*.**
1220
1221   **Figure 2-figure supplement 1**
1222   **Sequence alignments of BofA orthologs to determine conserved residues.**
1223
1224   **Figure 2-figure supplement 2**
1225   **Effects of BofA C-terminal residues and residues preceding predicted TMS2 on SpoIVFB**
1226   **inhibition in *E. coli*.**
1227
1228   **Figure 3-figure supplement 1**
1229   **GFPΔ27BofA variants are intact during *B. subtilis* sporulation.**
1230
1231   **Figure 4-figure supplement 1**
1232   **Models of SpoIVFB and BofA TMS2.**
1233
1234   **Figure 4-figure supplement 2**
1235   **Cys-less variants of SpoIVFA and MBPΔ27BofA inhibit cleavage of Pro-σ^K(1-127) by**
1236   **cytTM-SpoIVFB in *E. coli*.**
1237
1238   **Figure 4-figure supplement 3**
1239   **BofA TMS2 is in proximity to the active site of SpoIVFB.**
1240
1241   **Figure 4-figure supplement 4**
1242   **Full-length BofA is in proximity to the active site of SpoIVFB.**
1243
1244   **Figure 4-figure supplement 5**
1245   **BofA TMS2 has a preferred orientation in the active site cleft of SpoIVFB.**
1246
1247
1248

1249 **Figure 5-figure supplement 1**
1250 **Neither GFPΔ27BofA nor full-length BofA when co-produced with SpoIVFA prevent Pro-**
1251 **σ^K(1-127) from interacting with SpoIVFB.**
1252
1253 **Figure 5-figure supplement 2**
1254 **GFPΔ27BofA and SpoIVFA do not prevent full-length Pro-σ^K from interacting with**
1255 **SpoIVFB.**
1256
1257 **Figure 5-figure supplement 3**
1258 **Disulfide cross-linking between the cytTM-SpoIVFB CBS domain and the Pro-σ^K(1-127)**
1259 **C-terminal region.**
1260
1261 **Figure 6-figure supplement 1**
1262 **Disulfide cross-linking between E44C at the active site of cytTM-SpoIVFB and the**
1263 **Proregion of Pro-σ^K(1-127) in the absence of inhibitory proteins or in the presence of**
1264 **MBPΔ27BofA and SpoIVFA.**
1265
1266 **Figure 6-figure supplement 2**
1267 **Full-length BofA and SpoIVFA decrease cross-linking between E44C cytTM-SpoIVFB and**
1268 **V20C or K24C Pro-σ^K(1-127).**
1269
1270 **Figure 6-figure supplement 3**
1271 **Disulfide cross-linking between V70C in the cytTM-SpoIVFB membrane-reentrant loop**
1272 **and F18C or K24C in the Pro-σ^K(1-127) N-terminal region is decreased more by full-length**
1273 **BofA than by MBPΔ27BofA (lacking TMS1).**
1274
1275 **Figure 6-figure supplement 4**
1276 **Comparison of disulfide cross-linking between C46 in TMS2 of full-length BofA or**
1277 **MBPΔ27BofA (lacking TMS1) and E44C at or P135C near the active site of cytTM-**
1278 **SpoIVFB.**
1279
1280 **Figure 6-figure supplement 5**
1281 **Disulfide cross-linking between Y214C in the cytTM-SpoIVFB interdomain linker and**
1282 **L41C in the Pro-σ^K(1-127) N-terminal region is decreased more by full-length BofA than**
1283 **by MBPΔ27BofA (lacking TMS1).**
1284
1285 **Figure 7-figure supplement 1**
1286 **Model of SpoIVFB tetramer with part of one Pro-σ^K molecule.**
1287

1288 **Source Data Files**
1289
1290 **Figure 1-source data 1**
1291 **Immunoblot images (raw and annotated) and quantification of cleavage assays (Figure 1 B**
1292 **and C).**
1293
1294 **Figure 1-figure supplement 2-source data 1**
1295 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1296
1297 **Figure 1-figure supplement 3-source data 1**
1298 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1299
1300 **Figure 1-figure supplement 4-source data 1**
1301 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1302
1303 **Figure 2-source data 1**
1304 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1305
1306 **Figure 2-figure supplement 2-source data 1**
1307 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1308
1309 **Figure 3-source data 1**
1310 **Immunoblot images (annotated) and quantification of cleavage assays (Figure 3A), and**
1311 **confocal microscopy images (see the readme file) and a table of the numbers of sporangia**
1312 **in which FS localization of GFPΔ27BofA and the three variants was counted (Figure 3B).**
1313
1314 **Figure 3-source data 2**
1315 **Immunoblot images (raw) (Figure 3A).**
1316
1317 **Figure 3-figure supplement 1-source data 1**
1318 **Immunoblot images (raw and annotated).**
1319
1320 **Figure 4-source data 1**
1321 **Immunoblot images (raw and annotated).**
1322
1323 **Figure 4-figure supplement 1-source data 1**
1324 **PyMOL session file used to produce the images.**
1325
1326 **Figure 4-figure supplement 2-source data 1**
1327 **Immunoblot images (raw and annotated) and quantification of cleavage assays.**
1328
1329 **Figure 4-figure supplement 4-source data 1**
1330 **Immunoblot images (raw and annotated).**
1331
1332
1333

1334 **Figure 4-figure supplement 5-source data 1**
1335 **Immunoblot images (raw and annotated) (Figure 4-figure supplement 5 B and C) and**
1336 **quantification of cleavage assays (Figure 4-figure supplement 5B).**
1337
1338 **Figure 5-source data 1**
1339 **Immunoblot images (raw and annotated) (Figure 5A) and quantification of cross-linking**
1340 **(Figure 5B).**
1341
1342 **Figure 5-figure supplement 1-source data 1**
1343 **Immunoblot images (raw and annotated).**
1344
1345 **Figure 5-figure supplement 2-source data 1**
1346 **Immunoblot images (raw and annotated).**
1347
1348 **Figure 5-figure supplement 3-source data 1**
1349 **Immunoblot images (raw and annotated) and quantification of cleavage assays (Figure 5-**
1350 **figure supplement 3B).**
1351
1352 **Figure 6-source data 1**
1353 **Quantification of cross-linking (Figure 6 A, C, and E-I).**
1354
1355 **Figure 6-figure supplement 1-source data 1**
1356 **Immunoblot images (raw and annotated) (Figure 6-figure supplement 1A).**
1357
1358 **Figure 6-figure supplement 1-source data 2**
1359 **Immunoblot images (raw and annotated) (Figure 6-figure supplement 1B).**
1360
1361 **Figure 6-figure supplement 2-source data 1**
1362 **Immunoblot images (raw and annotated) (Figure 6-figure supplement 2A) and**
1363 **quantification of cross-linking (Figure 6-figure supplement 2B).**
1364
1365 **Figure 6-figure supplement 3-source data 1**
1366 **Immunoblot images (raw and annotated) (Figure 6-figure supplement 3 A, C, E, and G)**
1367 **and quantification of cross-linking (Figure 6-figure supplement 3 B, D, and F).**
1368
1369 **Figure 6-figure supplement 4-source data 1**
1370 **Immunoblot images (raw and annotated) (Figure 6-figure supplement 4 A, B, D, and E)**
1371 **and quantification of cross-linking (Figure 6-figure supplement 4 C and F).**
1372
1373 **Figure 6-figure supplement 5-source data 1**
1374 **Immunoblot images (raw and annotated).**
1375
1376 **Figure 7-source data 1**
1377 **PyMOL session file used to produce the images and PDB file of the model of a SpoIVFB**
1378 **tetramer with BofA and parts of SpoIVFA and Pro-σ^K.**
1379

1380 **Figure 7-figure supplement 1-source data 1**
1381 **PyMOL session file used to produce the images and PDB file of the model of a SpoIVFB**
1382 **tetramer with part of one molecule of Pro-σ^K.**
1383

1384 **Supplementary Files**
1385
1386 **Supplementary File 1**
1387 **Supplementary Results.**
1388
1389 **Supplementary File 2**
1390 **Supplementary Discussion.**
1391
1392 **Supplementary File 3**
1393 **Supplementary Materials and Methods.**
1394
1395 **Supplementary File 4**
1396 **Plasmids, primers, and *B. subtilis* strains used in this study.**
1397
1398 **Supplementary File 5**
1399 **Applied restraints from cross-linking, and distance evaluation, for the model of SpoIVFB**
1400 **in complex with part of Pro-σ^K.**
1401
1402 **Supplementary File 6**
1403 **Applied restraints from co-evolutionary coupling analysis, and distance evaluation, for the**
1404 **model of SpoIVFB in complex with part of Pro-σ^K.**
1405
1406 **Supplementary File 7**
1407 **Applied restraints from cross-linking, and distance evaluation, for the model of SpoIVFB**
1408 **in complex with BofA and parts of SpoIVFA and Pro-σ^K.**
1409
1410 **Supplementary File 8**
1411 **Applied restraints from co-evolutionary coupling analysis, and distance evaluation, for the**
1412 **model of SpoIVFB in complex with BofA and parts of SpoIVFA and Pro-σ^K.**