Tracking neural markers of template formation and implementation in attentional inhibition under different distractor consistency

Wen Wen[1], Zhibang Huang[1], Yin Hou[1], Sheng Li[*1,2,3,4]

1. School of Psychological and Cognitive Sciences, Peking University, Beijing, China
2. Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China
3. PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China
4. Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing, China

This work was supported by a grant from National Key R&D Program of China (2017YFB1002503).

Correspondence concerning this article should be addressed to Sheng Li, School of Psychological and Cognitive Sciences, Peking University, 5 Yiheyuan Road, Haidian, Beijing, 100871, China. E-mail: sli@pku.edu.cn

Word count: 5509

21   **Abstract**

To perform challenging visual search tasks, attention should be selectively allocated to

promote targets and inhibit distractors. Rejection templates based on the distractor's feature

can be built to constrain the search process by filtering-out matched stimuli. We measured

electroencephalography (EEG) of human participants when they performed a visual search

task under conditions in which the distractor cues were constant within a block (fixed-cueing)

or changed on a trial-by-trial basis (varied-cueing). Sustained representation of the cued

colors could be decoded from the whole-brain EEG signal during the retention interval in the

fixed-cueing condition, implying a formed rejection template. By contrast, transient

representation of the cued colors was observed in the varied-cueing condition only after cue

onset with significant but lower decoding accuracy than the fixed-cueing condition. The

differential neural representations of the to-be-ignored color in the two cueing conditions

agreed with the visual search performance that the fixed-cueing condition demonstrated a

larger inhibition benefit of the distractor cueing than the varied-cueing condition. During the

retention interval, we also observed stronger posterior alpha lateralization and mid-frontal

theta/beta power in the varied-cueing condition, indicating the cognitive costs in template

formation caused by the trialwisely change of distractor colors. The stable template formed in

the fixed-cueing condition was utilized to inhibit the matched distractors in the search array

and this was accompanied by larger mid-frontal theta activity. Taken together, our findings

highlighted the critical roles of distractor consistency in linking template formation to

successful distractor inhibition and the functional relevance of frontal theta activity in support

of template formation and implementation.

Keywords: rejection template, attentional suppression, visual search, cognitive control,

theta oscillation.

Introduction

Human beings can allocate limited attentional resources to task-relevant items whilst

filtering out irrelevant distractors. Extensive evidence has demonstrated that target templates

held in memory facilitate visual performance [1–5]. Importantly, the neural representation of

target templates during the preparation stage is positively correlated with subsequent task

performance [5–7]. By contrast, divergent findings have been reported regarding whether

foreknowledge of the to-be-ignored feature, referred to as a rejection template [8,9], can benefit

search efficiency. For those studies that showed the behavioral suppression benefits of the

rejection template, the emergence of such benefits seemed to be dependent on multiple

factors [10–16]. Of note, the consistency of the to-be-ignored feature across trials was

particularly prepotent [17–20]. These studies have demonstrated robust benefits of the distractor

cues when the cues remained unchanged within a block (fixed-cueing condition) and

minimum benefits if not absent when the cues changed on a trial-by-trial basis (varied-cueing

condition).

To reconcile the discrepancies of findings in the fixed and varied distractor-cueing

conditions, researchers proposed that a stable rejection template is essential in distractor

inhibition and can be created only under the fixed-cueing condition, whereas the varied-

cueing context obstructed its formation [8,17,21]. So far, there was limited neural evidence

shedding light on the existence of rejection templates. Reeder and colleagues [22] provided

their participants with three types of cues (positive cues, negative cues, and neutral cues) and

found that negative cues that indicated the distractor feature led to decreased responses in the

early visual cortex (V1-V3) prior to the search array. However, the authors later reported that

this decrement in the early visual cortex was not a feature-specific suppression and there

were no distinctive representations of the rejection templates [23]. Therefore, it is necessary to

examine whether selective representations of the rejection templates could be identified.

Recent studies come to reach the consensus that distractor inhibition could be learned

when statistical regularities of distractors' feature [24] or spatial location [25–27] are expected (see

Geng et al., 2019 for a review). By adopting a variant of Posner cueing paradigm, Noonan et

al. [20] found that benefits of distractor suppression, as indexed by a decreased amplitude of

early visual component P1, emerged only in the blocked condition where the distractor's

location was fixed. Similarly, by manipulating distractors' locations, van Moorelaar and

Slageter [29] showed that distractor-location-learning induced behavioral benefits appeared

after several repetitions and the learned expectation of distractors reduced stimulus-specific

processing as reflected by the decreased Pd component and location-decoding accuracy. On

the other hand, Vatterott and Vecera [19] examined the experience-dependent nature of feature-

based distractor suppression by manipulating the color contingency of a singleton distractor

in each trial within and between blocks. The capture effect of the singleton distractor

decreased within a block where the singleton's color remained unchanged but rebounded

when a new block started with another colored singleton. With all the evidence showing the

influence of distractor consistency on stimulus processing, yet it remained less understood

how distractor consistency modulates the process of template formation and subsequent

implementation of the template for distractor filtering.

Mid-frontal theta-band activities have been linked to goal-related prioritization in

working memory tasks [30,31] and conflict solving in the framework of cognitive control [32,33].

Specifically, stronger theta activity was evoked by distractor cues relative to target cues

during the delay period as a means of reducing interference from matched distractors [34].

Meanwhile, Dugué and colleagues [35] observed stronger post-stimulus theta activity in correct

trials than incorrect trials in a visual search task, revealing the association between post-

stimulus theta power and target selection. Although previous studies have indicated the tight

connection between template-based distractor inhibition and cognitive control [12,36], the exact

functional roles of theta activity in distractor inhibition, especially pertaining to template

formation and implementation, need more empirical investigations.

The current study aimed to reveal the feature-specific representation of rejection

template and the associated neural dynamics. Participants performed a cued visual search task

in which they were asked to locate the target and discriminate the orientation of the bar inside

of the target (Figure 1A). The participants were provided with distractor cues and were

informed that the search items in the same color as the distractor cues would never be the

target. The experiment contained the fixed-cueing and varied-cueing conditions that were

defined according to the consistency of the distractor cues within a block of trials. We

recorded electroencephalography (EEG) signals when the participants performed the task and

applied multivariate decoding method on the EEG signals to reveal the neural representations

of the to-be-ignored colors during the retention interval. We also examined the oscillatory

activities at different frequency bands to identify the neural markers that corresponded to

attentional selection, memory transformation, and cognitive control in the template formation

and implementation stages. Our results provide novel neural evidence in support of the

crucial roles of distractor consistency in linking template formation to successful distractor

inhibition.

Method

Participants.

We tested thirty-four participants in total (mean age = 21.3, SD = 2.2), ranging from 18

to 27 years old, 15 females, 1 left-handed). According to the G-power estimation [37], thirty

participants at least should be tested to guarantee a medium effect size (f = 0.25) and a high

statistic power (90%). Participants were compensated with monetary rewards. All

participants had normal or corrected-to-normal (color) vision and had no history of

psychiatric or neurological disorders. All participants provided written informed consent

prior to the experiment. The study was approved by the Committee for Protecting Human and

Animal Subjects at the School of Psychological and Cognitive Sciences at Peking University

(IRB protocol #2017-11-02).

Stimuli and procedure.

Participants were faced with a CRT monitor (refresh rate = 60 Hz) in a dark shielded

EEG booth and their heads were stabilized using a chin rest. The schematic procedure of a

trial is shown in Figure 1A. After a jittered blank (~1.5 s), a cue patch (1.5° × 1.5°) and an

equal-sized gray patch were presented (1.5° to the center) for 0.1 s. The gray patch was

presented to equalize the visual input in both hemispheres. The cue patch could appear at

either hemifield with equal probability. Participants were instructed that target in the

following search array would never be in the same color as the cued color. The color cue

remained the same within one block in the fixed-cueing condition while it changed

trialwisely in the varied-cueing condition. Two gray frames of the same size and locations as

the two patches were showed during the retention interval (1.5 s). The search array contained

eight items of different shapes (diamond/circle/hexagon, in the same square size of 2°× 2°)

and colors (four in target color and four in distractor color). Participants were instructed to

find the target, defined solely by its shape (diamond), and discriminate the orientation of the

bar inside of it. The target's color was randomly selected in each trial. In matched trials

(75%), the distractor's color was the same as the cued color. In neutral trials (25%), the

distractor's color was a new color different from the target color and the cued color. The

search array lasted for 0.2 s. A final memory test required participants to select the patch that

was in the same color as the cue from two presented patches. Both the search task and the

memory test had a response limit of 2 s.

Each cueing condition (fixed and varied) had 12 blocks with 64 trials per block. Six cue

colors (selected from L*a*b color space, L = 70, equally distributed in the a*b plane)

appeared in the first six blocks as well as the last six blocks in the fixed-cueing condition.

Participants came twice and completed the fixed-cueing and varied-cueing conditions on

separate days. The order of the conditions was counterbalanced between participants.

Participants were instructed to look at the fixation dot and their eye movements were

monitored using Eyelink 1000/1000Plus (1000 Hz, monocular).

EEG recording.

EEG data were acquired at a sampling rate of 1000 Hz using BrainAmps amplifier from a

64-channel EasyCap (Brain Products, Munich, Germany) with Ag/AgCl electrodes placed

according to the 10-20 system. An external electrode placed below the right eye was used to

measure eye movements. Electrode impedance was kept below 10 kΩ.

Data analysis.

Behavioral data: Only trials with correct responses in both the search task and the

memory task were included for search task's reaction times (RTs) analysis. Besides, we

recursively removed outlier trials where search RTs were beyond 2.5 standard deviations

from the mean. This resulted in 4.4% of the trials in the fixed-cueing condition and 3.7% of

the trials in the varied-cueing condition being discarded. There was a general practice effect

given that participants completed the two cueing conditions on separate days. Hence, we

calculated the benefit index for each cueing condition to represent the RT facilitation resulted

from the inhibition of matched distractors:

$$\text{RT benefit index} = 100 * \frac{RT_{neutral} - RT_{matched}}{(RT_{neutral} + RT_{matched})/2}$$

To examine the possible influence of testing order on the suppression benefit, we

performed a two-way mixed ANOVA on benefit index with testing order (fixed-cueing first

or varied-cueing first) as between-subject factor and cueing condition (fixed vs. varied) as

within-subject factor. We further split the trials into five bins based on the search RT

percentile for each trial type in each cueing condition. Then, we calculated the benefit index

for each percentile bin in each cueing condition and conducted one-sample t-tests (two-tailed,

alpha = .05) to compare the obtained benefit indices with zero. A two-way repeated ANOVA

with factors of cueing condition (fixed vs. varied) and trial type (matched vs. neutral) was

performed on the accuracies of the search task and the memory test. We reported the

corrected degrees of freedom (Greenhouse-Geisser correction) when the equal variances

assumption was violated.

EEG preprocessing: Offline preprocessing was performed with EEGLAB [38]. After down-

sampled to 500 Hz, the data was bandpass-filtered (1~30 Hz). Bad channels were spherically

interpolated. We then re-referenced the data to the averaged signal of mastoids and performed

independent component analysis (ICA) to correct ocular movements and other artifacts.

Finally, the continuous EEG was epoched for further analyses (the exact epoch length and

time-locked onset point were chosen depending on specific analyses). Besides behavioral

exclusion described above, epochs in which any channel showed above-threshold amplitude

(60 μV) at any time point of the epoch were removed. Since matched trials and neutral trials

differed from each other only after the onset of search array, we used both types of trials

when the analysis focused on the activities during the retention interval.

EEG decoding: To reveal the neural representations of the rejection templates, we

performed color-decoding based on the whole-brain electrodes' broad-band EEG data (EOG

excluded) during the retention interval. The decoding was separately performed on cue-left

and cue-right trials and the averaged results were used for statistical analysis. The continuous

EEG was segmented into -0.2~1.6 s relative to the cue onset and baseline-corrected to the

mean amplitude of the pre-stimulus period. We Gaussian-smoothed (window size = 16 ms)

the data along the time dimension to reduce temporal noises. At each time point, data was

partitioned into six training folds and one test fold. To create an unbiased classifier, the

number of trials for each cue color in the training set was equalized by subsampling. Trial-

averaged training set was generated by averaging the trials from the same cue color. We first

estimated the covariance matrix based on the trial-averaged training set. Next, we computed

the pair-wise Mahalanobis distances between test trials and the trial-averaged training set

using the covariance matrix [39]. A smaller distance toward the trial-averaged activation of one

specific color cue condition indicated a larger pattern similarity. Therefore, the decoding was

marked as a success when the distance between this given trial and the activation profile of

the same color was smaller than any of other colors' activation profile in the training data.

Cross-validation was carried out in a leave-one-fold-out manner. This routine was iterated

1000 times and trials were randomly assigned to the training and test set in each iteration.

The averaged decoding accuracy was taken for the statistical analysis. The accuracy time

series were Gaussian-smoothed (window size = 16 ms) and compared with the chance level

(i.e., 1/6). Cluster-based permutation was performed on the decoding accuracy time-series

over participants (alpha = .05, cluster-based nonparametric alpha = .05, cluster statistic = sum,

two-tail, permutation times = 10000 [40]).

Cross-temporal decoding: We followed the same decoding procedure as explained above

except that we trained the model using data at one time point and tested it on all time points.

We down-sampled the data into 50 Hz for computational efficiency. For the 2D result map

(as in Figure 2B), the diagonal line represents the accuracies when training and testing were

performed at the same time point while the off-diagonal points correspond to the temporal-

generalized decoding. We adopted the leave-one-fold-out cross-validation method throughout

the decoding process. Hence, the training-fold and the test-fold were never based on the same

trials no matter whether the training and test time points were the same or different. This

procedure was iterated 1000 times for each participant and the averaged result was used for

group statistical tests. Cluster-based permutation test was conducted on the 2D result map at

the participants' population level (alpha = .01, cluster-based nonparametric alpha = .01,

cluster statistic = sum, two-tail, permutation times = 10000 [40]).

Single-trial's averaged accuracy: Single-trial's averaged accuracy of the matched trials

was calculated to assess the relationship between the representation of the to-be-ignored color

and the following search performance. Instead of calculating the averaged decoding accuracy

(i.e., percentage of correct prediction) for each time point of the test-fold, we applied the

trained model to each trial in the test-fold and labeled the prediction of the trial using a binary

tag (1 for correct, 0 for incorrect). Once this was done for all time points of the trial, we took

the averaged binary tags (0~1600 ms) as the averaged accuracy for this specific trial. This

procedure was iterated for 1000 times and the mean accuracy was used for statistics. We

calculated the Pearson's correlation between trials' averaged accuracy and search RTs for

each participant. The populations' correlation coefficients in each cueing condition were

tested against zero using the one-sample bootstrapping t-test (alpha = 0.05, two-tail, bootstrap

samples = 10000).

Posterior alpha lateralization: A surface Laplacian filter (a 10th-order Legendre

polynomial and a lambda of 10^{-5}) was applied to increase topographical specificity and

reduce the effects of volume conduction (Cohen, 2019). To calculate the posterior lateral

alpha power, we segmented the continuous data into -0.4~1.6 s relative to the cue onset, and

the averaged ERP was subtracted from each trial to remove phase-locked activity evoked by

stimulus onset. Hilbert transform was used to extract the alpha-band (8~12 Hz) power of each

trial using posterior occipital electrodes (P5/6, P7/8, PO7/8). Power data was decibel

normalized to the averaged baseline (-0.4~-0.1 s). This process was separately performed on

left-side cue and right-side cue trials and the averaged results were used for the statistical

analysis. We calculated the alpha lateralization as the power difference between contralateral

and ipsilateral electrodes (contralateral minus ipsilateral).

Time-frequency analysis: We performed time-frequency decomposition on the

Laplacian-filtered data using Fieldtrip [42]. The continuous data were epoched -1.5~3 s relative

to cue onset. After the time-frequency decomposition, we selected data between -0.4~1.6 s

for further analysis. Each trial's signal was decomposed using Morlet wavelet convolution

every 50 ms. The frequencies ranged from 1~30 Hz in 30 linearly spaced steps and the width

of the wavelet kernel increased linearly from 1 to 10 along with frequency bins.

Normalization was done by decibel baselined to the pre-stimulus interval (-0.4~-0.1 s). Time-

frequency decomposition was separately done on the left- and right-cued trials. We further

combined the resulting data from the left- and right-cued trials so that the left-hemifield

electrodes were ipsilateral to the stimulus. To test the difference between conditions, cluster-

based permutation was performed at the group level. Adjacent time-frequency time points

exceeding the threshold (alpha = .01, two-tail) were grouped as a cluster. The cluster-level

statistic was calculated by taking the sum of the difference values within the cluster (cluster

alpha = .05). The number of random permutations using the Monte Carlo method was set to

10000.

We also examined the neural oscillations related to template implementation during

visual search. Continuous data were epoched -1~2 s relative to search array onset. We

adopted the same time-frequency decomposition method to obtain the oscillatory activities

during -0.4~1.2 s for all trials.

Results

Behavioral suppression benefits in matched trials

We observed larger response benefits in the fixed-cueing condition than the varied-

cueing condition (F(1,32) = 9.661, p = .004, η_p^2 = .232, Figure 1B) but no significant effect

of testing order (F(1,32) = 1.558, p = .221, η_p^2 = .046). There was no significant interaction

effect between the two factors (F(1,32) = .064, p = .802, η_p^2 = .002)[1].

As shown in Figure 1C, we analyzed the benefit index across five RT percentile bins.

The results showed that the suppression benefits were evident across all bins in the fixed-

cueing condition (all p < .001). However, in the varied-cueing condition, we observed

significant suppression benefit only in the percentile bin with longest RTs (p < .001). The

early and robust behavioral benefits in the fixed-cueing condition indicated the engagement

of proactive inhibition when the to-be-ignored color remained the same. By contrast, the

benefit of reactive inhibition was relatively late and emerged only in the group of trials with

longest RTs in the varied-cueing condition. These results were consistent with our previous

findings that cognitive control played a crucial role in distractor inhibition in the varied-

cueing condition [18].

We further analyzed the results of accuracy for visual search and memory test (Figure

---

[1] See Supplementary Materials Figure S1 for the search RTs of different conditions.

1D). For the visual search accuracy, there were no significant differences between conditions

(F(1,33) = 1.784, p = .191, η_p^2 = .051) or trial types (F(1,33) = 3.115, p = .087, η_p^2 = .086)

but their interaction effect was significant (F(1,33) = 8.983, p = .005, η_p^2 = .214). Simple

effect analysis showed that participants were more accurate in the matched trials than the

neutral trials in the fixed-cueing condition (F(1,33) = 11.31, p = .002). However, this

difference was not evident in the varied-cueing condition (F(1,33) = 0.19, p = .662). For the

memory test, participants were more accurate in the fixed-cueing condition (F(1, 33) =

100.318, p < .001, η_p^2 = .752) and the matched trials (F(1, 33) = 93.234, p < .001, η_p^2

= .739). The significant interaction effect (F(1, 33) = 46.202, p < .001, η_p^2 = .583) indicated

that when the cued color reappeared in the search array as the distractor color, it promoted

memory performance to a larger extent in the varied-cueing condition (5.8%, F(1, 33) =

83.26, p < .001) than in the fixed-cueing condition (1.3%, F(1, 33) = 19.28, p < .001). This

result might be explained by the refreshment strategy (or perceptual sampling) that using the

reappeared matched distractors to enhance the memory of the to-be-ignored distractor color

in the varied-cueing condition [9].

EEG decoding of the to-be-ignored color during retention interval

We performed decoding analysis on EEG data to reveal the representation of the

distractor color during the retention interval. As shown in Figure 2A, color information could

be decoded in both conditions (fixed-cueing: 0.078~1.6 s, cluster p < .001; varied-cueing:

0.26~0.806 s, cluster p < .001). Importantly, the decoding accuracy was higher in the fixed-

cueing condition compared with the varied-cueing condition, especially after the cue onset

(cluster p < .001, 0.08~0.438 s) and prior to search array onset (cluster p = .021, 1.096~1.272

s; cluster p = .005, 1.342~1.6 s).

Next, we focused on representation stability and performed cross-temporal decoding

separately for the two cueing conditions. Figure 2B shows the two-dimensional maps of

decoding accuracies. There was a wide range of generalization for color decoding between

time points in the fixed-cueing condition, indicating a more stable representation of the cued

color during the retention interval. In contrast, significant decoding performance was

observed only when training and testing were performed on the data from the same time

points in the varied-cueing condition, suggesting a highly dynamic representation of the cued

color. These results were in line with the larger behavioral suppression benefit in the fixed-

cueing condition as such benefit would highly rely on the stable representation of the

distractor's feature to form an effective rejection template.

To further unfold the relationship between the neural representation of the cued color and

distractor inhibition, we calculated the single-trial's averaged accuracy on the matched trials

in the two cueing conditions. We then computed the trialwise correlation between the single-

trials' mean accuracies and search RTs for each participant. We hypothesized that stable

rejection template would facilitate distractor filtering and therefore lead to faster response in

the matched trials. If the neural representation instead reflected initial sensory processing,

better representation of the distractor cue would lead to behavioral cost. Results in Figure 2C

showed that single-trial's averaged accuracy was positively correlated with search RT in the

varied-cueing condition (mean of correlation coefficients r = .014, p = .042) confirming that

better representation of the color would cause attraction. This effect was not evident in the

fixed-cueing condition (mean of correlation coefficients r = .004, p = .562) which might be

explained by the ceiling effect of behavioral suppression benefits. These discrepancies

suggested distinct neural representations of the to-be-ignored cue in the two cueing

contingencies. In fact, when we performed multivariate representation similarity analysis

(RSA) on the topographical data within and between two cueing conditions (see

Supplementary Materials Figure S2), the results showed that RSA scores of the within-

condition scheme were higher than the between-condition scheme. Thus, although based on

the same sensory information, brain signals during the retention interval manifested different

topographical distributions under different cueing conditions.

Posterior alpha lateralization during early stage of retention interval

An increase in posterior alpha-band (8~12 Hz) activity has been regarded as a neural

marker of attentional inhibition [43–45]. Correspondingly, the decrease of alpha power at the

contralateral side of an attended stimulus indicated the release from inhibition (i.e., increase

in attention). As shown in Figure 3A, we observed significant alpha lateralization

(contralateral minus ipsilateral) induced by the color cue in both cueing conditions (fixed-

cueing condition: 0.282~0.74 s, cluster p = .008; varied-cueing condition: 0.266~1.124 s,

cluster p = .001). Stronger alpha lateralization (herein, more negative) was found in the

varied-cueing condition relative to the fixed-cueing condition (0.338~1.066 s, cluster p

= .003), indicating a larger attentional bias towards the color cue at the initial stage of the

retention interval.

Mid-frontal theta and beta activities during the late stage of retention interval

Next, we focused on mid-frontal activities in the two cueing conditions (Figure 3B left).

The time-frequency difference map of mid-frontal electrodes showed that varied cues

induced stronger oscillatory activities in the range of theta-band and beta-band at the later

stage of the retention interval in comparison with the fixed cues (0.35~1.6 s, 2~19 Hz, cluster

p = .002). Frontal theta oscillation (3~7 Hz) has been regarded as a reflection of top-down

control in goal-directed cognitive processes [32]. In two recent studies, stronger mid-frontal

theta was observed during the retention interval if the cue indicated the distractor rather than

the target [34] or switching rather than repeating the task sets [46]. The enhanced mid-frontal theta

activity was also suggested to be associated with the prevention of interference from the

anticipated distractor [34] or the proactive control of task set switching [46]. In our experiment,

the varied-cueing condition shared these critical aspects of task structure with these two

studies. Therefore, we consider that both the prevention of interference from the anticipated

distractor and the proactive control of task set switching could contribute to the observed

mid-frontal theta effects during the retention interval.

On the other hand, frontal beta activity (13~30 Hz) has been proposed to reflect a short-

lived transition period between representations (e.g., from latent to active) late in the WM

delay in the service of task demands [47]. We speculated that the higher mid-frontal beta

activity in the varied-cueing condition (Figure 3B, right-top) reflected a stronger engagement

in transferring the cueing information from perceptual representation into rejection template

that can be used to filter out the matched distractors in the following search task. However,

given the lack of literature on the roles of frontal beta in attentional suppression, future

experiments with more specific designs are required to test this interpretation.

Mid-frontal theta activity during visual search period

Finally, time-frequency analysis also highlighted the dominant low-frequency oscillatory

activity in mid-frontal electrodes during the search process (1~9 Hz, 0.2-1.2 s, cluster p

< .001, Figure 4A). Specifically, as shown in Figure 4B, mid-frontal theta power during

visual search correlated with individuals' RTs in the matched (fixed-cueing: r = -.398, p

= .020; varied-cueing: r = -.500, p = .003) and neutral trials (fixed-cueing: r = -.445, p = .009;

varied-cueing: r = -.657, p < .001). The observed correlations demonstrated that mid-frontal

theta reflected a general process of cognitive control that filters out the distractors of different

colors and shapes. Participants who utilized stronger control process were faster in localizing

the target.

Of particular interest, we looked into the relationship between post-stimulus theta

activity and template implementation. If mid-frontal theta also signaled the implementation

of the rejection template on top of the general control process in visual search, we would

expect to observe stronger mid-frontal theta for the trials (e.g., matched trials) in which the

rejection templates were successfully utilized for search. To test this assumption, we

conducted a repeated-measures ANOVA on the mid-frontal theta power with cueing

condition (fixed vs. varied) and trial type (matched vs. neutral) as within-subject factors. We

did not find significant difference for either cueing condition ($F(1,33) = 1.928$, $p = .174$, $\eta^2 = $

0.055) or trial type ($F(1,33) = 0.006$, $p = .939$, $\eta^2 < .001$). However, there was a marginal

significant interaction effect between the two factors ($F(1,33) = 3.675$, $p = .064$, $\eta^2 = 0.100$).

Simple effect analysis revealed that the fixed-cueing condition had stronger mid-frontal theta

power than the varied-cueing condition in the matched trials ($F(1,33) = 6.94$, $p = .013$), but

not in the neutral trials ($F(1,33) = 0.09$, $p = .762$). These results were consistent with the

observed larger behavioral suppression benefits in the fixed-cueing condition, indicating that

mid-frontal theta served as a neural marker not only for the general control process of

distractor filtering but also for the implementation of the established rejection template.

Discussion

Distractor inhibition is challenging and context-dependent. Inconsistent results have been

reported concerning whether distractors matched with the to-be-ignored feature could be

inhibited. The present study identified the neural signature of the rejection template in the

fixed-cueing condition as characterized by the sustained decoding of the to-be-ignored color

and the stable cross-temporal decoding performance. By contrast, trial-wise updates of

distractor cues in the varied-cueing condition hampered the establishment of stable rejection

templates. This was reflected as the smaller behavioral suppression benefit and the transient

decoding performance that did not persist in the late stage of the retention interval. Moreover,

the results of single-trial's averaged accuracy implied that the representation of distractor

cues in the varied-cueing condition would conversely interfere with the following visual

search performance. These findings collectively suggested that creating rejection templates is

difficult when the distractor consistency is violated.

The specific processes related to template formation and implementation were further

revealed by oscillatory activities in different frequency bands. First, we observed stronger

posterior alpha lateralization toward the color cue in the varied-cueing condition than the

fixed-cueing condition, reflecting the higher necessity of paying attention to the color cue

when it changed on a trial-by-trial basis. Second, the posterior alpha lateralization was

followed by mid-frontal theta and beta activities in the late stage of the retention interval.

Based on previous literature, we suggest that the theta band activity reflected the control

processes such as the prevention of the anticipated distractor interference and the proactive

control of cue switching [34,46]. The beta band activity, on the other hand, may index the

memory transition process that transferred the color cue from perceptual representation to

rejection template [47]. Third, increased mid-frontal theta activity was found after the search

array onset and was behaviorally relevant with search performance. This post-stimulus mid-

frontal theta activity was stronger in the fixed-cueing condition than the varied-cueing

condition restrictively in the matched trials where participants could utilize the rejection

template. Overall, we suggest that the post-stimulus mid-frontal theta also reflected the

control processes of distractor filtering and implementation of rejection template in visual

search.

Capitalized on color decoding of EEG signal, we demonstrated selective representations

of the to-be-ignored feature in both the fixed-cueing and varied-cueing conditions. In Reeder

and colleagues' [23] fMRI study, they performed representational similarity analysis to

calculate the distance metric of five cued colors in target, distractor, and neutral cueing

conditions. Compared with the neutral condition, a more distinctive representation was

observed only when the color was cued to be the target but not when the color was a

distracting feature. The authors argued that the representations of general task-irrelevant

features in neutral condition might not be an ideal baseline because participants might inhibit

the neutral cue as well. Instead of comparing the representation distinctiveness across

different cueing contingencies, we emphasized selective representation of the to-be-ignored

feature within each condition. Meanwhile, the decoding accuracy time series provided us

with a temporally fine-scaled picture of the representations of rejection templates. In another

study, by repeating distractor locations, van Moorselaar and colleagues [29] showed how

learned expectations promoted distractor inhibition. Whereas distractor expectation reduced

distractor-specific processing after search array onset, meaningful preparatory neural tuning

to distractor locations was not observed in their study. Herein, we first provide the evidence

of color-selective representation of the distractor cue during the retention interval,

particularly in the fixed-cueing condition. Discrepancies in the contents of rejection template

as well as the experimental design might contribute to the different findings. Compared to the

location-based expectation in van Moorselaar's study which only excluded one item, taking

the advantage of the to-be-ignored color in the present study would help to filter out a half of

the items in a search array. Moreover, fixed cues that were presented within one block could

further facilitate the generation of stable rejection templates.

We found better color decoding performance in the fixed-cueing condition compared to

the varied-cueing condition right after cue onset. The constant cueing contingency in fixed-

cueing condition promoted the expectation of distractor color and therefore boosted the color

selectivity during the retention interval. This might account for the very early (<100 ms)

emergence of the template information extracted by decoding. In fact, previous work showed

that expectation could pre-activate representations of expected sensory stimuli [48] and

strategically bias gazes toward the foreknown distractor in the preparatory stage to reduce

stimulus-evoked attractions [49]. Behaviorally, fixed cues also facilitate distractor suppression

which echoed previous findings that participants can be more effective in excluding the

irrelevant items in a predictive context [14,17,20].

Given the current design, we could not elucidate how the matched distractors were

filtered out based on the rejection template. Previous work found that the target template was

pre-activated briefly before stimulus onset and served as a match filter to convert the sensory

input of stimulus into the decision-relevant representation [50]. However, it remains unknown

whether the rejection template functions the same way as the target template. It has been

proposed that the rejection template might reduce the activation of matched features and

therefore prevent those matched distractors from interfering with target processing in the first

place [20,51]. Spatial tuning to the distractor-location also decreased when participants learned

to ignore the distractor [29]. Our behavioral results showed that inhibition was achieved even in

the fastest RT bin in the fixed-cueing condition, indicating a proactive suppression

presumably through the reduction of the distractor-evoked response. On the contrary,

rejection templates could be utilized reactively [52] which fitted our finding that suppression

benefits emerged only at the longest RT bin in the varied-cueing condition. Future studies are

required to further investigate the factors that modulate the implementation process.

In conclusion, our results revealed the neural markers for the distinctive processes of

template formation and implementation between the fixed-cueing and varied-cueing

conditions. Distractor consistency in the fixed-cueing condition promoted the formation of

rejection template and corresponded to the significant behavioral suppression benefit. In the

varied-cueing condition, transient decoding performance as well as the stronger posterior

alpha lateralization and mid-frontal theta/beta power during the retention interval implied the

cognitive costs in template formation caused by the trial-by-trial changes of the distractor

features. Differences in template formation could contribute to the utilization phase where we

observed successful implementation of the rejection template under the fixed-cueing

condition. This behavioral advantage of fixed-cueing was accompanied by the stronger mid-

frontal theta power after search array onset in the matched trials. Together, these findings

provide an integrative view of template-guided distractor inhibition in visual search.

Author contributions

WW and SL designed the research. WW, ZH and YH collected data. WW analysed data

and interpreted findings under the supervision of SL. WW drafted the manuscript. SL revised

the manuscript. All authors have reviewed the content and approved the final version for

publication.

References

1.   Desimone, R. & Duncan, J. Neural mechanisms of selective visual attention. *Annu.*

Rev. Neurosci. **18**, 193–222 (1995).

2.   Soto, D., Heinke, D., Humphreys, G. W. & Blanco, M. J. Early, involuntary top-down

guidance of attention from working memory. *J. Exp. Psychol. Hum. Percept. Perform.*

31, 248–261 (2005).

3.   Soto, D., Hodsoll, J., Rotshtein, P. & Humphreys, G. W. Automatic guidance of

attention from working memory. *Trends Cogn. Sci.* **12**, 342–348 (2008).

4.   Reeder, R. R. & Peelen, M. V. The contents of the search template for category-level

search in natural scenes. *J. Vis.* **13**, 13 (2013).

5.   Eimer, M. The neural basis of attentional control in visual search. *Trends Cogn. Sci.* **18**,

526–535 (2014).

6.   Stokes, M., Thompson, R., Nobre, A. C. & Duncan, J. Shape-specific preparatory

activity mediates attention to targets in human visual cortex. *Proc. Natl. Acad. Sci. U.*

S. A. **106**, 19569–19574 (2009).

7.   Vogel, E. K., McCollough, A. W. & Machizawa, M. G. Neural measures reveal

individual differences in controlling access to working memory. *Nature* **438**, 500–503

(2005).

8.   Arita, J. T., Carlisle, N. B. & Woodman, G. F. Templates for rejection: configuring

attention to ignore task-irrelevant features. *J. Exp. Psychol. Hum. Percept. Perform.* **38**,

580 (2012).

9.   Woodman, G. F. & Luck, S. J. Do the Contents of Visual Working Memory

Automatically Influence Attentional Selection During Visual Search? *J. Exp. Psychol.*

Hum. Percept. Perform. **33**, 363–377 (2007).

10.   Beck, V. M., Luck, S. J. & Hollingworth, A. Whatever you do, don't look at the. .:

Evaluating guidance by an exclusionary attentional template. *J. Exp. Psychol. Hum.*

Percept. Perform. **44**, 645–662 (2018).

11.   Conci, M., Deichsel, C., Müller, H. J. & Töllner, T. Feature guidance by negative

attentional templates depends on search difficulty. *Vis. cogn.* **0**, 1–10 (2019).

12.   Han, S. W. & Kim, M. S. Do the Contents of Working Memory Capture Attention?

Yes, But Cognitive Control Matters. *J. Exp. Psychol. Hum. Percept. Perform.* **35**,

1292–1302 (2009).

13.   Tanda, T. & Kawahara, J. I. Association between cue lead time and template-for-

rejection effect. *Atten. Percept. Psychophys.* (2019).

14.   Töllner, T., Conci, M. & Müller, H. J. Predictive distractor context facilitates

attentional selection of high, but not intermediate and low, salience targets. *Hum.*

Brain Mapp. **36**, 935–944 (2015).

15.   Olivers, C. N. L. What drives memory-driven attentional capture? The effects of

memory type, display type, and search type. *J. Exp. Psychol. Hum. Percept. Perform.*

35, 1275 (2009).

16.   Zhang, Z., Gapelin, N. & Carlisle, N. B. Probing early attention following negative

and positive templates. *Attention, Perception, Psychophys.* **82**, 1166–1175 (2020).

17.   Cunningham, C. A. & Egeth, H. E. Taming the White Bear: Initial Costs and Eventual

Benefits of Distractor Inhibition. *Psychol. Sci.* **27**, 476–485 (2016).

18.   Wen, W., Hou, Y. & Li, S. Memory guidance in distractor suppression is governed by

the availability of cognitive control. *Attention, Perception, Psychophys.* **80**, 1157–1168

(2018).

19.   Vatterott, D. B. & Vecera, S. P. Experience-dependent attentional tuning of distractor

rejection. *Psychon. Bull. Rev.* **19**, 871–878 (2012).

20.   Noonan, M. A. P. *et al.* Distinct mechanisms for distractor suppression and target

facilitation. *J. Neurosci.* **36**, 1797–1807 (2016).

21.   Carlisle, N. B. Flexibility in attentional control: Multiple sources and suppression. *Yale*

J. Biol. Med. **92**, 103–113 (2019).

22.   Reeder, R. R., Olivers, C. N. L. & Pollmann, S. Cortical evidence for negative search

templates. *Vis. cogn.* **25**, 278–290 (2017).

23.   Reeder, R. R., Olivers, C. N. L., Hanke, M. & Pollmann, S. No evidence for enhanced

distractor template representation in early visual cortex. *Cortex* **108**, 279–282 (2018).

24.   Stilwell, B. T., Bahle, B. & Vecera, S. P. Feature-based statistical regularities of

distractors modulate attentional capture. *J. Exp. Psychol. Hum. Percept. Perform.* **45**,

419–433 (2019).

25.   Wang, B. & Theeuwes, J. Statistical regularities modulate attentional capture. *J. Exp.*

Psychol. Hum. Percept. Perform. **44**, 13–17 (2018).

26.   Ferrante, O. *et al.* Altering spatial priority maps via statistical learning of target

selection and distractor filtering. *Cortex* **102**, 67–95 (2018).

27.   Leber, A. B., Gwinn, R. E., Hong, Y. & O'Toole, R. J. Implicitly learned suppression

of irrelevant spatial locations. *Psychon. Bull. Rev.* **23**, 1873–1881 (2016).

28.   Geng, J. J., Won, B.-Y. Y. & Carlisle, N. B. Distractor Ignoring: Strategies, Learning,

and Passive Filtering. *Curr. Dir. Psychol. Sci.* **28**, 096372141986709 (2019).

29.   van Moorselaar, D. & Slagter, H. A. Learning What Is Irrelevant or Relevant:

Expectations Facilitate Distractor Inhibition and Target Facilitation through Distinct

Neural Mechanisms. *J. Neurosci.* **39**, 6953–6967 (2019).

30.   Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S. & D'Esposito, M. Causal Evidence

for a Role of Theta and Alpha Oscillations in the Control of Working Memory. *Curr.*

Biol. **30**, 1748-1754.e4 (2020).

31. de Vries, I. E. J., Slagter, H. A. & Olivers, C. N. L. Oscillatory Control over

Representational States in Working Memory. *Trends Cogn. Sci.* **24**, 150–162 (2020).

32. Cavanagh, J. F. & Frank, M. J. Frontal theta as a mechanism for cognitive control.

Trends Cogn. Sci. **18**, 414–421 (2014).

33. Cohen, M. X. A neural microcircuit for cognitive conflict detection and signaling.

Trends Neurosci. **37**, 480–490 (2014).

34. de Vries, I. E. J., Savran, E., van Driel, J. & Olivers, C. N. L. Oscillatory Mechanisms

of Preparing for Visual Distraction. *J. Cogn. Neurosci.* **31**, 1873–1894 (2019).

35. Dugué, L., Marque, P. & VanRullen, R. Theta Oscillations Modulate Attentional

Search Performance Periodically. *J. Cogn. Neurosci.* **27**, 945–958 (2014).

36. Kiyonaga, A., Egner, T. & Soto, D. Cognitive control over working memory biases of

selection. *Psychon. Bull. Rev.* **19**, 639–646 (2012).

37. Faul, F., Erdfelder, E., Lang, A. G. & Buchner, A. G*Power 3: A flexible statistical

power analysis program for the social, behavioral, and biomedical sciences. *Behav.*

Res. Methods **39**, 175–191 (2007).

38. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *J. Neurosci. Methods*

134, 9–21 (2004).

39. Wolff, M. J., Jochim, J., Akyürek, E. G. & Stokes, M. G. Dynamic hidden states

underlying working-memory-guided behavior. *Nat. Neurosci.* **20**, 864–871 (2017).

40. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG-and MEG-data. *J.*

Neurosci. Methods **164**, 177–190 (2007).

41.   Cohen, M. X. *Analyzing Neural Time Series Data*. *Analyzing Neural Time Series Data*

(MIT press, 2019). doi:10.7551/mitpress/9609.001.0001

42.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Comput. Intell. Neurosci. **2011**, (2011).

43.   Worden, M. S., Foxe, J. J., Wang, N. & Simpson, G. V. Anticipatory biasing of

visuospatial attention indexed by retinotopically specific alpha-band

electroencephalography increases over occipital cortex. *J. Neurosci.* **20**, RC63 (2000).

44.   Kelly, S. P., Lalor, E. C., Reilly, R. B. & Foxe, J. J. Increases in Alpha Oscillatory

Power Reflect an Active Retinotopic Mechanism for Distracter Suppression During

Sustained Visuospatial Attention. *J. Neurophysiol.* **95**, 3844–3851 (2006).

45.   Jensen, O. & Mazaheri, A. Shaping functional architecture by oscillatory alpha activity:

gating by inhibition. *Front. Hum. Neurosci.* **4**, 186 (2010).

46.   Cooper, P. S. *et al.* Frontal theta predicts specific cognitive control-induced

behavioural changes beyond general reaction time slowing. *Neuroimage* **189**, 130–140

(2019).

47.   Spitzer, B. & Haegens, S. Beyond the status quo: A role for beta oscillations in

endogenous content (RE)activation. *eNeuro* **4**, (2017).

48.   Kok, P., Mostert, P. & de Lange, F. P. Prior expectations induce prestimulus sensory

templates. *Proc. Natl. Acad. Sci.* **114**, 10473–10478 (2017).

49.   Wen, W., Zhang, Y. & Li, S. Gaze dynamics of feature-based distractor inhibition

under the prior-knowledge and expectation. *Atten. Percept. Psychophys.* (2021).

doi:10.1101/2020.07.19.210781

50.   Myers, N. E. *et al.* Testing sensory evidence against mnemonic templates. *Elife* **4**, 1–

25 (2015).

51.   Moher, J., Lakshmanan, B. M., Egeth, H. E. & Ewen, J. B. Inhibition Drives Early

Feature-Based Attention. *Psychol. Sci.* **25**, 315–324 (2014).

52.   Moher, J. & Egeth, H. E. The ignoring paradox: Cueing distractor features leads first

to selection, then to inhibition of to-be-ignored items. *Attention, Perception,*

Psychophys. **74**, 1590–1605 (2012).

Figure 1. Task and behavioral results. (A) Task. The color cue indicated the to-be-ignored color in the search array. Participants were asked to find the target (diamond) and judge whether the bar inside of it was clockwise or counter-clockwise tilted relative to the mid-vertical. In the memory test, participants were instructed to select the color patch that matched the cue. (B) Suppression benefit index. Red and blue dots represent the individuals' benefit indices in the fixed-cueing and varied-cueing conditions, respectively. Black lines represent the averaged benefit indices across participants. (C) Suppression benefit indices of RT percentile bins. Errorbars represent the standard deviations of 5000 times bootstrapped means. (D) Accuracy of visual search task and memory test. * $p < .05$, ** $p < .01$, *** $p < .001$, *̲ $p < .001$ (two-tailed).

Figure 2. Results of color-decoding during retention interval. (A) Time-series of decoding

accuracy in the fixed-cueing and varied-cueing conditions. The shaded areas represent the

standard deviation of 5000 times bootstrapped mean. The solid/dashed vertical line indicate

the cue onset/offset. Solid lines below the x-axis indicated significant time-clusters for each

condition and the black lines correspond to the significant time-clusters for condition

difference. (B) Cross-temporal decoding of cued color in the two cueing conditions. Alpha-

blending was performed on insignificant time-points for better illustration. (C) Histogram of

participants' correlation coefficients between single-trial's averaged accuracy and search RT

in the matched trials. Vertical dashed lines are the average correlation coefficients across

participants for the two cueing conditions. The x-axis represents the trialwise Pearson's

correlation coefficients. The y-axis represents the proportion of participants. P-values are

one-sample bootstrapping t-test of populations' correlation coefficients against zero.

Figure 3. Oscillatory activities during retention interval. (A) Top: Posterior lateralized alpha

power in the fixed-cueing and varied-cueing conditions. The shaded areas represent the

between-subject standard error. The solid/dashed vertical line indicated the cue onset/offset.

Solid lines below the x-axis indicated significant time-clusters for each condition with the

black line corresponding to their difference. Bottom: Topography of the averaged lateralized

alpha power during 300~800 ms. Topographical maps indicate the difference values of alpha

power between left-cue minus right-cue trials. Since contralateral alpha power would

decrease relative to ipsilateral alpha power when distractor cues are attended, left posterior

electrodes are positive in this plot. The black dots highlight the channels used for alpha

lateralization analysis. Black dots correspond to the channels used for alpha lateralization

analysis (P5/6, P7/8, PO7/8). (B) Left: Time-frequency map extracted from mid-frontal

electrodes (FCz, FC1, FC2, Fz, F1, F2). Dotted contours were adopted from the time-

frequency difference map on the right column. See Supplementary Materials Figure S3 for

the topography of alpha-band and beta-band activity during retention interval. Right-top:

Time-frequency difference map between the varied-cueing and fixed-cueing conditions

(varied minus fixed). Significant time-frequency cluster from the non-parametric permutation

test are out-lined by the dark contour. Data was extracted from mid-frontal electrodes. Right-

bottom: Topographical distribution of mid-frontal theta activity (3~7 Hz, 0.1~1.6 s) for the

fixed-cueing and varied-cueing conditions. Black dots correspond to the channels used for

time-frequency analysis.

Figure 4. Mid-frontal theta after search array onset. (A) Top: Time-frequency map after

search array onset. The significant time-frequency cluster from the non-parametric

permutation test was outlined by the dark contour. Bottom: Topography of averaged theta

power during 0.2~0.7s and the power difference between conditions. To avoid stimulus-

driven and response-related contaminates, we chose the time window to be 0.2~0.7s. The

averaged search RTs across conditions was 735 ms. The black dots indicated the channels

used for power analysis (FCz, FC1, FC2, Fz, F1, F2). (B) Spearman-rank correlations

between frontal theta power and RTs in matched trials and neutral trials of two cueing

conditions. The shaded area represents the 95% confidence interval for the fitted line. (C)

Mid-frontal theta power of different trial types and cueing conditions. Errorbars represent the

within-subject corrected standard error.