Tracking neural markers of template formation and implementation in attentional inhibition under different distractor consistency

Wen Wen [1], Zhibang Huang [1], Yin Hou [1], Sheng Li [*1,2,3,4]

1. School of Psychological and Cognitive Sciences, Peking University, Beijing, China

2. Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China

3. PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

4. Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing, China

Correspondence concerning this article should be addressed to Sheng Li, School of Psychological and Cognitive Sciences, Peking University, 5 Yiheyuan Road, Haidian, Beijing, 100871, China. E-mail: sli@pku.edu.cn

Declaration of Interests: The authors declare no competing financial interests.

Acknowledgments: This work was supported by a grant from National Key R&D Program of China (2017YFB1002503).

18   **Abstract**

Performing visual search tasks requires optimal attention deployment to promote targets

and inhibit distractors. Rejection templates based on the distractor's feature can be built to

constrain the search process. We measured electroencephalography (EEG) of human

participants of both sexes when they performed a visual search task in conditions where the

distractor cues were constant within a block (fixed-cueing) or changed on a trial-by-trial basis

(varied-cueing). In the fixed-cueing condition, sustained decoding of the cued colors could be

achieved during the retention interval and the participants with higher decoding accuracy

showed larger suppression benefit of the distractor cueing in the search period. In the varied-

cueing condition, the cued color could only be transiently decoded after its onset and the

higher decoding accuracy was observed from the participants who demonstrated lower

suppression benefit. The differential neural representations of the to-be-ignored color in the

two cueing conditions as well as their reverse associations with behavioral performance

implied that rejection templates were formed in the fixed-cueing condition but not in the

varied-cueing condition. Additionally, we observed stronger posterior alpha lateralization and

mid-frontal theta/beta power during the retention interval of the varied-cueing condition,

indicating the cognitive costs in template formation caused by the trialwise change of

distractor colors. Taken together, our findings revealed the neural markers associated with the

critical roles of distractor consistency in linking template formation to successful inhibition.

Keywords: rejection template, attentional suppression, cognitive control, theta

oscillation, visual search.

Introduction

Human beings can allocate limited attentional resources to task-relevant items whilst

filtering out irrelevant distractors. Extensive evidence has demonstrated that target templates

held in memory facilitate visual performance (Desimone & Duncan, 1995; Eimer, 2014;

Reeder & Peelen, 2013; Soto et al., 2005, 2008). By contrast, divergent findings have been

reported regarding whether foreknowledge of the to-be-ignored feature, referred to as

rejection templates (Arita et al., 2012; Woodman & Luck, 2007), can benefit search

efficiency. Studies showed that the behavioral benefit of the rejection template is dependent

on many factors (Beck et al., 2018; Conci et al., 2019; Han & Kim, 2009; Olivers, 2009;

Tanda & Kawahara, 2019; Töllner et al., 2015; Zhang et al., 2020). Notably, the consistency

of the to-be-ignored feature across trials was particularly prepotent (Cunningham & Egeth,

2016; Noonan et al., 2016; Vatterott & Vecera, 2012; Wen et al., 2018). These studies have

demonstrated robust inhibition benefits when distractor cues remained unchanged within a

block (fixed-cueing condition) and minimum benefits if not absent when the cues changed on

a trial-by-trial basis (varied-cueing condition).

Researchers proposed that a stable rejection template is essential in distractor inhibition

and can be created only under the fixed-cueing condition, whereas the varied-cueing context

obstructed its formation (Arita et al., 2012; Carlisle, 2019; Cunningham & Egeth, 2016). So

far, there was limited neural evidence shedding light on the existence of rejection templates.

Reeder and colleagues (2017) found that negative cues that indicated the distractor feature led

to decreased responses in the early visual cortex (V1-V3) prior to the search array. However,

they later reported that this decrement was not a feature-specific suppression and there were

no distinctive representations of the rejection templates (Reeder et al., 2018). Therefore, it is

necessary to examine whether selective representations of the rejection templates could be

identified.

Recent studies have reached the consensus that distractor inhibition could be learned

when statistical regularities of distractors' feature (Stilwell et al., 2019) or spatial location

(Ferrante et al., 2018; Leber et al., 2016; Wang & Theeuwes, 2018) are expected (Geng et al.,

2019). For example, Noonan et al. (2016) found decreased P1 component in the blocked

condition where the distractor's location was fixed. Similarly, van Moorselaar and Slagter

(2019) showed that distractor-location learning induced behavioral benefits appeared after

several repetitions and the learned expectation of distractors reduced stimulus-specific

processing as reflected by the decreased Pd component and location-decoding accuracy. With

all the evidence showing the influence of distractor consistency on stimulus processing, yet it

remained less understood how distractor consistency modulates the process of template

formation and subsequent implementation.

Mid-frontal theta-band activities have been linked to goal-related prioritization in

working memory tasks (de Vries et al., 2020; Riddle et al., 2020) and conflict solving in the

framework of cognitive control (Cavanagh & Frank, 2014; Cohen, 2014). Specifically,

stronger theta activity was evoked by distractor cues relative to target cues during the delay

period as a means of reducing interference from matched distractors (de Vries et al., 2019).

Meanwhile, Dugué and colleagues (2014) observed stronger post-stimulus theta activity in

correct trials than incorrect trials, revealing the association between post-stimulus theta

power and target selection. Although behavioral studies have indicated the tight connection

between template-based distractor inhibition and cognitive control (Han & Kim, 2009;

Kiyonaga et al., 2012), the exact functional roles of theta activity in distractor inhibition,

especially pertaining to template formation and implementation, need more empirical
investigations.

The current study aimed to reveal the feature-specific representation of rejection
template and the associated neural dynamics. Participants were provided with distractor cues
and were asked to perform a cued visual search task (Figure 1A). We recorded
electroencephalography (EEG) signals and applied multivariate decoding to reveal the neural
representations of the to-be-ignored colors. We also examined the oscillatory activities
corresponding to attentional selection, memory transformation, and cognitive control in the
template formation and implementation stages.

Materials and Methods

Participants.

We tested thirty-four participants in total (mean age = 21.3, SD = 2.2), ranging from 18
to 27 years old, 15 females, 1 left-handed). According to the G-power estimation (Faul et al.,
2007), thirty participants at least should be tested to guarantee a medium effect size (f = 0.25)
and a high statistic power (90%). Participants were compensated with monetary rewards. All
participants had normal or corrected-to-normal (color) vision and had no history of
psychiatric or neurological disorders. All participants provided written informed consent
prior to the experiment. The study was approved by the Committee for Protecting Human and
Animal Subjects at the School of Psychological and Cognitive Sciences at Peking University
(IRB protocol #2017-11-02).

Stimuli and procedure.

Participants were faced with a CRT monitor (refresh rate = 60 Hz) in a dark shielded

EEG booth and their heads were stabilized using a chin rest. The schematic procedure of a

trial is shown in Figure 1A. After a jittered blank (~1.5 s), a cue patch (1.5° × 1.5°) and an

equal-sized gray patch were presented (1.5° to the center) for 0.1 s. The gray patch was

presented to equalize the visual input in both hemispheres. The cue patch could appear at

either hemifield with equal probability. Participants were instructed that target in the

forthcoming search array would never be in the same color as the cued color. The color cue

remained the same within one block in the fixed-cueing condition but changed on a trial-by-

trial basis in the varied-cueing condition. Two gray frames of the same size and locations as

the two patches were shown during the retention interval (1.5 s). The search array contained

eight items of different shapes (diamond/circle/hexagon, in the same square size of 2°× 2°)

and colors (four in target color and four in distractor color). Participants were instructed to

find the target, defined solely by its shape (diamond), and discriminate the orientation of the

bar inside of it. The target's color was randomly selected in each trial. In matched trials

(75%), the distractor's color was the same as the cued color. In neutral trials (25%), the

distractor's color was a new color different from the target color and the cued color. The

search array lasted for 0.2 s and was masked by checkerboards following its offset. Masks

disappeared when participants made a response. A final memory test required participants to

select the patch that was in the same color as the cue from two presented patches. To

discourage participants from memorizing the cue color using semantic categorization, some

pairs (~30%) in the memory test were set to be visually similar. Both the search task and the

memory test had a response limit of 2 s.

Each cueing condition (fixed and varied) had 12 blocks with 72 trials per block. Six cue

colors (selected from L*a*b color space, L = 70, equally distributed in the a*b plane)

appeared in the first six blocks as well as the last six blocks in the fixed-cueing condition.

Participants came twice and completed the fixed-cueing and varied-cueing conditions on

separate days. The order of the conditions was counterbalanced between participants.

Participants were instructed to look at the fixation dot and their eye movements were

monitored using Eyelink 1000/1000Plus (1000 Hz, monocular).

Statistical analysis.

Only trials with correct responses in both the search task and the memory task were

included for search task's reaction times (RTs) analysis. Besides, we recursively removed

outlier trials where search RTs were beyond 2.5 standard deviations from the mean. This

resulted in 4.4% of the trials in the fixed-cueing condition and 3.7% of the trials in the varied-

cueing condition being discarded. There was a general practice effect given that participants

completed the two cueing conditions on separate days. Hence, we calculated the benefit index

for each cueing condition to represent the RT facilitation resulted from the inhibition of

matched distractors:

$$\text{RT benefit index} = 100 * \frac{RT_{neutral} - RT_{matched}}{(RT_{neutral} + RT_{matched})/2}$$

To examine the possible influence of testing order on the suppression benefit, we

performed a two-way mixed ANOVA on benefit index with testing order (fixed-cueing first

or varied-cueing first) as between-subject factor and cueing condition (fixed vs. varied) as

within-subject factor. We further split the trials into five bins based on the search RT

percentile for each trial type in each cueing condition. Then, we calculated the benefit index

for each percentile bin in each cueing condition and conducted one-sample t-tests (two-tailed,

alpha = .05) to compare the obtained benefit indices with zero. A two-way repeated ANOVA

with factors of cueing condition (fixed vs. varied) and trial type (matched vs. neutral) was

performed on the accuracies of the search task and the memory test. We reported the

corrected degrees of freedom (Greenhouse-Geisser correction) when the equal variances

assumption was violated.

EEG recording.

EEG data were acquired at a sampling rate of 1000 Hz using BrainAmps amplifier from a

64-channel EasyCap (Brain Products, Munich, Germany) with Ag/AgCl electrodes placed

according to the 10-20 system. An external electrode placed below the right eye was used to

measure eye movements. Electrode impedance was kept below 10 kΩ.

EEG data analysis.

EEG preprocessing: Offline preprocessing was performed with EEGLAB (Delorme &

Makeig, 2004). After down-sampled to 500 Hz, the data was bandpass-filtered (1~30 Hz).

Bad channels were spherically interpolated. We then re-referenced the data to the averaged

signal of mastoids and performed independent component analysis (ICA) to correct ocular

movements and other artifacts. On average, there were 3.8 (SD=1.4) and 4.1 (SD=1.5)

components removed in the fixed-cueing and varied-cueing conditions respectively. Finally,

the continuous EEG was epoched for further analyses (the exact epoch length and time-

locked onset point were chosen depending on specific analyses). Besides behavioral

exclusion described above, epochs in which any channel showed above-threshold amplitude

(60 µV) at any time point of the epoch were removed. Since matched trials and neutral trials

differed from each other only after the onset of search array, we used both types of trials

when the analysis focused on the activities during the retention interval.

EEG decoding: To reveal the neural representations of the rejection templates, we

performed color-decoding based on the whole-brain electrodes' broad-band EEG data (EOG

excluded) during the retention interval. The decoding was separately performed on cue-left

and cue-right trials and the averaged results were used for statistical analysis. The continuous

EEG was segmented into -0.2~1.6 s relative to the cue onset and baseline-corrected to the

mean amplitude of the pre-stimulus period. We Gaussian-smoothed (window size = 16 ms)

the data along the time dimension to reduce temporal noises. At each time point, data was

partitioned into six training folds and one test fold. To create an unbiased classifier, the

number of trials for each cue color in the training set was equalized by subsampling. Trial-

averaged training sets were generated by averaging the trials from the same cue color. We

first estimated the covariance matrix based on the trial-averaged training set. Next, we

computed the pair-wise Mahalanobis distances between test trials and the trial-averaged

training set using the covariance matrix (Wolff et al., 2017). A smaller distance toward the

trial-averaged activation of one specific color cue condition indicated a larger pattern

similarity. Therefore, the decoding was marked as a success when the distance between this

given trial and the activation profile of the same color was smaller than any of other colors'

activation profiles in the training data. Cross-validation was carried out in a leave-one-fold-

out manner. This routine was iterated 1000 times and trials were randomly assigned to the

training and test set in each iteration. The averaged decoding accuracy was taken for the

statistical analysis. The accuracy time series were Gaussian-smoothed (window size = 16 ms)

and compared with the chance level (i.e., 1/6). Cluster-based permutation was performed on

the decoding accuracy time-series over participants (alpha = .05, cluster-based nonparametric

alpha = .05, cluster statistic = sum, two-tail, permutation times = 10000 (Maris & Oostenveld,

2007)).

Cross-temporal decoding: We followed the same decoding procedure as explained above

except that we trained the model using data at one time point and tested it on all time points.

We down-sampled the data into 50 Hz for computational efficiency. For the 2D result map

(as in Figure 2B), the diagonal line represents the accuracies when training and testing were

performed at the same time point while the off-diagonal points correspond to the temporal-

generalized decoding. We adopted the leave-one-fold-out cross-validation method throughout

the decoding process. Hence, the training-fold and the test-fold were never based on the same

trials no matter whether the training and test time points were the same or different. This

procedure was iterated 1000 times for each participant and the averaged result was used for

group statistical tests. Cluster-based permutation test was conducted on the 2D result map at

the participants' population level (alpha = .01, cluster-based nonparametric alpha = .01,

cluster statistic = sum, two-tail, permutation times = 10000 (Maris & Oostenveld, 2007)).

Cross-condition decoding: To compare the representations of color cues in the two

cueing conditions, we performed cross-condition decoding by training the model using data

from one condition and testing it on the other one (i.e., trained on the fixed-cueing condition

and tested on the varied-cueing condition or vice versa). For each color of the training set, we

subsampled equal number of trials to create an unbiased classifier. This subsampling process

was iterated 1000 times. Bi-directional cross-condition decoding demonstrated similar results

and therefore we performed statistical analysis on the average data (cluster-based permutation,

alpha = .05, cluster-based nonparametric alpha = .05, cluster statistic = sum, two-tail,

permutation times = 10000).

Multivariate topographical pattern similarity: The topographical activation pattern was

defined as the whole-brain channels' activities at a certain time point. First, we reorganized

the data to integrate the left- and right-cued trials so that the left-hemifield electrodes were

ipsilateral to the cue. Topographical pattern similarity was calculated as the correlation

between the pattern of a given trial and the averaged pattern from the same cueing condition

(i.e., within-condition scheme) or different cueing condition (i.e., between-condition scheme)

(Figure 3A). The correlation was separately computed for each cued color to avoid the

influence of sensory difference. For the between-condition scheme, the averaged

topographical activation pattern was created by averaging trials of the same cued color from

the other cueing condition. For the within-condition scheme, the averaged topographical

activation pattern was created by averaging trials of the same cued color from the same

cueing condition but in different blocks. We excluded the trials in the same block to reduce

the temporal similarity effect. Considering that the number of trials that contributed to the

averaged topographical activation patterns might confound the results, we subsampled the

trials so that the number of trials used for the averaged topographical activation patterns

across colors in each cueing condition was equalized. The subsampling process was iterated

1000 times. We treated the topographical activation pattern at each time point as a vector and

calculated the Pearson's correlation coefficients between the vector of the given trial and the

vector of the averaged topographical activation pattern (Figure 3B). The averaged correlation

coefficients across all interactions and all time points during the retention interval were used

to represent each participant's similarity scores in the four conditions (2 cueing conditions $\times$

2 correlating schemes). The similarity scores were fisher-Z transformed when performing the

statistical test at the population level.

Posterior alpha lateralization: A surface Laplacian filter (a 10th-order Legendre

polynomial and a lambda of 10^{-5}) was applied to increase topographical specificity and

reduce the effects of volume conduction (Cohen, 2019). To calculate the posterior lateral

alpha power, we segmented the continuous data into -0.4~1.6 s relative to the cue onset, and

the averaged ERP was subtracted from each trial to remove phase-locked activity evoked by

stimulus onset. Hilbert transform was used to extract the alpha-band (8~12 Hz) power of each

trial using posterior occipital electrodes (P5/6, P7/8, PO7/8). Power data was decibel

normalized to the averaged baseline (-0.4~-0.1 s). This process was separately performed on

left-side cue and right-side cue trials and the averaged results were used for the statistical

analysis. We calculated the alpha lateralization as the power difference between contralateral

and ipsilateral electrodes (contralateral minus ipsilateral).

Time-frequency analysis: We performed time-frequency decomposition on the

Laplacian-filtered data using Fieldtrip (Oostenveld et al., 2011). The continuous data were

epoched -1.5~3 s relative to cue onset. After the time-frequency decomposition, we selected

data between -0.4~1.6 s for further analysis. Each trial's signal was decomposed using Morlet

wavelet convolution every 50 ms. The frequencies ranged from 1~30 Hz in 30 linearly

spaced steps and the width of the wavelet kernel increased linearly from 1 to 10 along with

frequency bins. Normalization was done by decibel baselined to the pre-stimulus interval (-

0.4~-0.1 s). Time-frequency decomposition was separately done on the left- and right-cued

trials. We further combined the resulting data from the left- and right-cued trials so that the

left-hemifield electrodes were ipsilateral to the stimulus. To test the difference between

conditions, cluster-based permutation was performed at the group level. Adjacent time-

frequency points exceeding the threshold (alpha = .01, two-tail) were grouped as a cluster.

The cluster-level statistic was calculated by taking the sum of the difference values within the

cluster (cluster alpha = .05). The number of random permutations using the Monte Carlo

method was set to 10000.

We also examined the neural oscillations related to template implementation during

visual search. Continuous data were epoched -1~2 s relative to search array onset. We

adopted the same time-frequency decomposition method to obtain the oscillatory activities

during -0.4~1.2 s for all trials.

Results

Behavioral suppression benefits in matched trials

As shown in Figure 1B, search RT had a significant main effect of trial type ($F(1,33) =$

$21.132, p < .001, \eta_p^2 = .390$) but not for cueing condition ($F(1,33) = .016, p = .899, \eta_p^2 < .001$),

indicating that participants were generally faster in matched trials than neutral trials (19.8 ms).

The significant interaction effect between cueing-condition and trial type ($F(1,33) = 8.558, p$

$= .006, \eta_p^2 = .206$) was driven by the larger suppression benefits in the fixed-cueing condition

(26.8 ms, $F(1,33) = 28.21, p < .001$) than the varied-cueing condition (12.9 ms, $F(1,33) =$

$7.17, p = .011$). Considering the practice effect across testing days, we calculated benefit

index which represents the suppression benefit of matched trials over neutral trials. Larger

suppression benefits were observed in the fixed-cueing condition than the varied-cueing

condition ($F(1,32) = 9.661, p = .004, \eta_p^2 = .232$, Figure 1C) and testing order had no

significant influence on suppression benefit ($F(1,32) = 1.558, p = .221, \eta_p^2 = .046$). There

was no significant interaction effect between the two factors ($F(1,32) = .064, p = .802, \eta_p^2$

$= .002$).

Figure 1D demonstrates the benefit index across five RT percentile bins. Suppression

benefits were evident across all bins in the fixed-cueing condition (all $p < .001$). However,

for the varied-cueing condition, we observed significant suppression benefit only in the

percentile bin with longest RTs ($p < .001$). The early and robust behavioral benefits in the

fixed-cueing condition indicated the engagement of proactive inhibition when the to-be-

ignored color remained the same. By contrast, the late suppression benefit of the longest RT

bin in the varied-cueing condition implies that participants adopted a reactive control which

required sufficient time to come into effect. These results were consistent with our previous

findings that cognitive control played a crucial role in distractor inhibition in the varied-

cueing condition (Wen et al., 2018).

We further analyzed the results of accuracy for visual search and memory test (Figure

1E). For the visual search accuracy, there were no significant differences between conditions

($F(1,33) = 1.784$, $p = .191$, $\eta_p^2 = .051$) or trial types ($F(1,33) = 3.115$, $p = .087$, $\eta_p^2 = .086$)

but their interaction effect was significant ($F(1,33) = 8.983$, $p = .005$, $\eta_p^2 = .214$). Simple

effect analysis showed that participants were more accurate in the matched trials than the

neutral trials in the fixed-cueing condition ($F(1,33) = 11.31$, $p = .002$). However, this

difference was not evident in the varied-cueing condition ($F(1,33) = 0.19$, $p = .662$). For the

memory test, participants were more accurate in the fixed-cueing condition ($F(1, 33) =$

100.318, $p < .001$, $\eta_p^2 = .752$) and the matched trials ($F(1, 33) = 93.234$, $p < .001$, $\eta_p^2 = .739$).

The significant interaction effect ($F(1, 33) = 46.202$, $p < .001$, $\eta_p^2 = .583$) indicated that when

the cued color reappeared in the search array as the distractor color, it promoted memory

performance to a larger extent in the varied-cueing condition (5.8%, $F(1, 33) = 83.26$, p

$< .001$) than in the fixed-cueing condition (1.3%, $F(1, 33) = 19.28$, $p < .001$). This result

might be explained by the refreshment strategy (or perceptual sampling) that using the

reappeared matched distractors to enhance the memory of the to-be-ignored distractor color

in the varied-cueing condition (Woodman & Luck, 2007).

EEG decoding of the to-be-ignored color during retention interval

We performed multivariate decoding analysis on EEG data to reveal the representation of

the distractor color during the retention interval. Color information could be decoded in both

conditions (Figure 2A, fixed-cueing: 0.078~1.6 s, cluster $p < .001$; varied-cueing: 0.26~0.806

s, cluster $p < .001$). Importantly, the decoding accuracy was higher in the fixed-cueing

condition compared with the varied-cueing condition, especially after the cue onset (cluster p

$< .001$, 0.08~0.438 s) and prior to search array onset (cluster $p = .021$, 1.096~1.272 s; cluster

$p = .005$, 1.342~1.6 s). To further rule out the possibility that the selective color

representation was driven by the memory test, we split the participants based on their overall

memory performance (across matched and neutral trials) and compared the decoding

accuracy of the two subgroups in each cueing condition. No significant difference in

decoding accuracy between the high and low performer groups was observed (fixed-cueing:

t(32) = 0.433, $p = .668$; varied-cueing: t(32) = 0.155, $p = .878$).

Next, we focused on representation stability and performed cross-temporal decoding

separately for the two cueing conditions. Figure 2B shows the two-dimensional maps of

decoding accuracies. There was a wide range of generalization for color decoding between

time points in the fixed-cueing condition, indicating a more stable representation of the cued

color during the retention interval. In contrast, significant decoding performance was

observed only when training and testing were performed on the data from the same time

points in the varied-cueing condition, suggesting a highly dynamic representation of the cued

color. Moreover, the difference map demonstrated that the fixed-cueing condition had higher

off-diagonal decoding accuracy than the varied-cueing condition (Figure 2C). These results

were in line with the larger behavioral suppression benefit in the fixed-cueing condition as

such benefit would highly rely on the stable representation of the distractor's feature to form

an effective rejection template.

Opposite effects in relating EEG decoding to behavior in two cueing conditions

To examine how the representations of distractor cues affect distractor filtering during

visual search, we half-split the participants into two groups based on their behavioral benefit

indices in each cueing condition. In the fixed-cueing condition, participants who showed

larger suppression benefit had higher decoding accuracy (Figure 2D, $t(32) = 2.119$, $p = .042$,

Cohen'd = 0.73). This pattern was reversed in the varied-cueing condition as higher decoding

accuracy was observed among the low suppression benefit participants (Figure 2D, $t(32) =$

2.092, $p = .044$, Cohen'd = 0.71). This difference suggested that distinct neural

representations of the to-be-ignored cue in the two cueing contingencies. To further examine

the neural representation difference, we performed cross-condition decoding. Classifiers

trained on trials from one cueing condition can only decode distractor cues from the other

condition during the early retention interval (Figure 2E, 134~442 ms, cluster $p < .001$). The

decoding accuracy was around chance level thereafter. This result indicated that the

representation of color cues underwent two stages where sensory signals were emphasized in

the initial stage. The unsuccessful cross-condition decoding suggested that different coding

patterns were adopted to represent the color cue.

Additionally, we conducted multivariate topographical pattern similarity analysis on the

basis of the within- and between-condition scheme correlations (Figure 3). If the rejection

template was represented similarly in the two cueing conditions, one would expect that

similarity scores for the within-condition and between-condition schemes were of no

significant difference. However, the results revealed that the similarity scores were

significantly different between the two cueing conditions ($F(1,33) = 19.305$, $p < .001$, $\eta_p^2 =$

0.369) and the two correlating schemes ($F(1,33) = 54.373$, $p < .001$, $\eta_p^2 = 0.622$). There was

also an interaction effect ($F(1,33) = 26.9$, $p < .001$, $\eta_p^2 = 0.449$). Simple effect analysis

revealed that the similarity score of the within-condition scheme was higher than that of the

between-condition scheme for the varied-cueing condition ($F(1,33) = 68.78$, $p < .001$). The

same trend was observed for the fixed-cueing condition with marginal significance ($F(1,33) =$

4.05, $p = .052$). These results suggest that, although exposed to the same sensory information,

the brain signals during the retention interval manifested different topographical

representations under the two cueing conditions.

Posterior alpha lateralization during the early stage of retention interval

An increase in posterior alpha-band (8~12 Hz) activity has been regarded as a neural

marker of attentional inhibition (Jensen & Mazaheri, 2010; Kelly et al., 2006; Worden et al.,

2000). Correspondingly, the decrease of alpha power at the contralateral side of an attended

stimulus indicated the release from inhibition (i.e., increase in attention). As shown in Figure

4A, we observed significant alpha lateralization (contralateral minus ipsilateral) induced by

the color cue in both cueing conditions (fixed-cueing condition: 0.282~0.74 s, cluster p

$= .008$; varied-cueing condition: 0.266~1.124 s, cluster $p = .001$). Stronger alpha

lateralization (herein, more negative) was found in the varied-cueing condition relative to the

fixed-cueing condition (0.338~1.066 s, cluster $p = .003$), indicating a larger attentional bias

towards the color cue at the initial stage of the retention interval. This attentional bias was

attenuated with the repetition of color cues in the fixed-cueing condition.

Mid-frontal theta and beta activities during the late stage of retention interval

Next, we focused on mid-frontal activities in the two cueing conditions (Figure 4B). The

time-frequency difference map of mid-frontal electrodes showed that varied cues induced

stronger oscillatory activities in the range of theta-band and beta-band at the later stage of the

retention interval in comparison with the fixed cues (0.35~1.6 s, 2~19 Hz, cluster $p = .002$).

Frontal theta oscillation (3~7 Hz) has been regarded as a reflection of top-down control in

goal-directed cognitive processes (Cavanagh & Frank, 2014). In two recent studies, stronger

mid-frontal theta was observed during the retention interval if the cue indicated the distractor

rather than the target (de Vries et al., 2019) or switching rather than repeating the task sets

(Cooper et al., 2019). The enhanced mid-frontal theta activity was also suggested to be

associated with the prevention of interference from the anticipated distractor (de Vries et al.,

2019) or the proactive control of task set switching (Cooper et al., 2019). In our experiment,

the varied-cueing condition shared these critical aspects of task structure with these two

studies. Therefore, we consider that both the prevention of interference from the anticipated

distractor and the proactive control of task set switching could contribute to the observed

mid-frontal theta effects during the retention interval.

On the other hand, frontal beta activity (13~30 Hz) has been proposed to reflect a short-

lived transition period between representations (e.g., from latent to active) late in the WM

delay in the service of task demands (Spitzer & Haegens, 2017). We speculated that the

higher mid-frontal beta activity in the varied-cueing condition (Figure 4B) reflected a

stronger engagement in transferring the cueing information from perceptual representation

into rejection template that can be used to filter out the matched distractors in the following

search task. However, given the lack of literature on the roles of frontal beta in attentional

suppression, future experiments with more specific designs are required to test this

interpretation.

Mid-frontal theta activity during visual search period

Finally, time-frequency analysis also highlighted the dominant low-frequency oscillatory

activity in mid-frontal electrodes during the search process (1~7 Hz, 0.2-1.2 s, cluster p

< .040, Figure 5A). There were no significant differences between conditions, trial types, or

their interaction (repeated measures ANOVA, all *ps* > .302). We then examined the

functional relevance of the post-stimulus theta power with search performance (Figure 5B).

Particularly, theta power predicted search RT in the varied-cueing condition (matched trials:

Spearman's ρ = -.564, *p* = .001; neutral trials: Spearman's ρ = -.680, *p* < .001) but not in the

fixed-cueing condition (matched trials: Spearman's ρ = -.174, *p* = .324; neutral trials:

Spearman's ρ = -.108, *p* = .542). These results indicate that mid-frontal theta at the post-

stimulus stage reflected a reactive control process that filtered out the distractors in the

absence of stable rejection templates.

Discussion

Distractor inhibition is challenging and context-dependent. Inconsistent results have been

reported concerning whether distractors matched with the to-be-ignored feature could be

inhibited. The present study identified the neural signature of the rejection template in the

fixed-cueing condition as characterized by the sustained decoding of the to-be-ignored color

and the stable cross-temporal decoding performance. By contrast, trialwise updates of

distractor cues in the varied-cueing condition hampered the establishment of stable rejection

templates. This was reflected as the smaller behavioral suppression benefit and the transient

decoding performance that did not persist in the late stage of the retention interval. Moreover,

we found that better neural representation of the distractor cues in the varied-cueing condition

led to smaller suppression benefits. These findings collectively suggested that creating

rejection templates is difficult when the distractor consistency is violated.

The specific processes related to template formation were unfolded by oscillatory

activities in different frequency bands. First, we observed stronger posterior alpha

lateralization toward the color cue in the varied-cueing condition than the fixed-cueing

condition, reflecting the higher necessity of paying attention to the color cue when it changed

on a trial-by-trial basis. The posterior alpha lateralization was followed by mid-frontal theta

and beta activities in the late stage of the retention interval. Based on previous literature, we

suggest that the theta band activity reflected the control processes such as the prevention of

the anticipated distractor interference and the proactive control of cue switching (Cooper et

al., 2019; de Vries et al., 2019). The beta band activity, on the other hand, may index the

memory transition process that transferred the color cue from perceptual representation to

rejection template (Spitzer & Haegens, 2017). However, this tentative explanation needs

future empirical investigations.

Capitalized on color decoding of EEG signal, we demonstrated selective representations

of the to-be-ignored feature in both the fixed-cueing and varied-cueing conditions. In Reeder

and colleagues' (2018) fMRI study, they performed representational similarity analysis to

calculate the distance metric of five cued colors in target, distractor, and neutral cueing

conditions. Compared with the neutral condition, a more distinctive representation was

observed only when the color was cued to be the target but not when the color was a

distracting feature. The authors argued that the representations of general task-irrelevant

features in neutral condition might not be an ideal baseline because participants might inhibit

the neutral cue as well. Instead of comparing the representation distinctiveness across

different cueing contingencies, we emphasized selective representation of the to-be-ignored

feature within each condition. Meanwhile, the decoding accuracy time series provided us

with a temporally fine-scaled picture of the representations of rejection templates. In another

study, by repeating distractor locations, van Moorselaar and colleagues (2019) showed how

learned expectations promoted distractor inhibition. Whereas distractor expectation reduced

distractor-specific processing after search array onset, meaningful preparatory neural tuning

to distractor locations was not observed in their study. Herein, we first provide the evidence

of color-selective representation of the distractor cue during the retention interval,

particularly in the fixed-cueing condition. Discrepancies in the contents of rejection template

as well as the experimental design might contribute to the different findings. Compared to the

location-based expectation in van Moorselaar's study which only excluded one item, taking

the advantage of the to-be-ignored color in the present study would help to filter out half of

the items in a search array. Moreover, fixed cues that were presented within one block could

further facilitate the generation of stable rejection templates.

The constant cueing contingency in fixed-cueing condition promoted the expectation of

distractor color and therefore boosted the color selectivity during the retention interval. This

might account for the very early (<100 ms) emergence of the template information extracted

by decoding. In fact, previous work showed that expectation could pre-activate

representations of expected sensory stimuli (Kok et al., 2017) and strategically bias gazes

toward the foreknown distractor in the preparatory stage to reduce stimulus-evoked

attractions (Wen et al., 2021). Behaviorally, fixed cues also facilitate distractor suppression

which echoed previous findings that participants can be more effective in excluding the

irrelevant items in a predictive context (Cunningham & Egeth, 2016; Noonan et al., 2016;

Töllner et al., 2015).

The opposite effects in relating decoding accuracy to suppression benefit in the two

cueing conditions suggest that neural representations of the distractor cues were different in

nature. Together with oscillatory activity in retention, this reverse pattern implied that the

formation of rejection template was dependent on distractor consistency. Since robust

rejection templates were built in the fixed-cueing condition, participants could implement the

rejection template in this condition to proactively suppress the matched distractors, but relied

on reactive control to filter out non-targets in the varied-cueing condition. This was

evidenced by the negative correlation between the post-stimulus theta power and search RT

that was only observed in the varied-cueing condition. In addition, behavioral suppression

benefits in the varied-cueing condition were smaller than the fixed-cueing condition and

appeared only in the longest RT bin. Hence, while proactive inhibition played a central role

in the fixed-cueing condition, distractor suppression in the varied-cueing condition was

primarily undertaken by reactive control. Notably, the negative correlation was observed in

both matched trials and neutral trials, implying that reactive control operated in a universal

manner and rejected all nontargets in the absence of a stable rejection template.

Given the current design, we could not elucidate how the matched distractors were

filtered out based on the rejection template especially in the fixed-cueing condition. Previous

work found that the target template was pre-activated briefly before stimulus onset and

served as a match filter to convert the sensory input of stimulus into the decision-relevant

representation (Myers et al., 2015). However, it remains unknown whether the rejection

template functions the same way as the target template. It has been proposed that the

rejection template might reduce the activation of matched features and therefore prevent

those matched distractors from interfering with target processing in the first place (Moher et

al., 2014; Noonan et al., 2016; van Zoest et al., 2021). Spatial tuning to the distractor location

also decreased when participants learned to ignore the distractor (van Moorselaar & Slagter,

2019). Our behavioral results showed that inhibition was achieved even in the fastest RT bin

in the fixed-cueing condition, indicating a proactive suppression presumably through the

reduction of the distractor-evoked response. On the contrary, rejection templates could be

utilized reactively (Moher & Egeth, 2012) and contribute to suppression benefits of the

longest RT bin in the varied-cueing condition. Future studies are required to further

investigate the factors that modulate the implementation process.

In conclusion, our results revealed the neural markers for the distinctive processes of

template formation and implementation between the fixed-cueing and varied-cueing

conditions. Distractor consistency in the fixed-cueing condition promotes the formation of

rejection template and leads to the significant behavioral suppression benefit. In the varied-

cueing condition, transient decoding performance as well as the stronger posterior alpha

lateralization and mid-frontal theta/beta power during the retention interval imply the

cognitive costs in template formation caused by the trial-by-trial changes of the distractor

features. In comparison to the fixed-cueing condition where rejection template can be

implemented to proactively suppress matched distractors, the varied-cueing condition relies

on reactive control to filter out non-targets. Together, these findings provide an integrative

view of template-guided distractor inhibition in visual search.

References

Arita, J. T., Carlisle, N. B., & Woodman, G. F. (2012). Templates for rejection: configuring

attention to ignore task-irrelevant features. *Journal of Experimental Psychology: Human*

Perception and Performance, *38*(3), 580.

Beck, V. M., Luck, S. J., & Hollingworth, A. (2018). Whatever you do, don't look at the. .:

Evaluating guidance by an exclusionary attentional template. *Journal of Experimental*

Psychology: Human Perception and Performance, *44*(4), 645–662.

https://doi.org/10.1037/xhp0000485

Carlisle, N. B. (2019). Flexibility in attentional control: Multiple sources and suppression.

Yale Journal of Biology and Medicine, *92*(1), 103–113.

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control.

Trends in Cognitive Sciences, *18*(8), 414–421. https://doi.org/10.1016/j.tics.2014.04.012

Cohen, M. X. (2014). A neural microcircuit for cognitive conflict detection and signaling.

Trends in Neurosciences, *37*(9), 480–490. https://doi.org/10.1016/j.tins.2014.06.004

Cohen, M. X. (2019). Analyzing Neural Time Series Data. In *Analyzing Neural Time Series*

Data. MIT press. https://doi.org/10.7551/mitpress/9609.001.0001

Conci, M., Deichsel, C., Müller, H. J., & Töllner, T. (2019). Feature guidance by negative

attentional templates depends on search difficulty. *Visual Cognition*, *0*(0), 1–10.

https://doi.org/10.1080/13506285.2019.1581316

Cooper, P. S., Karayanidis, F., McKewen, M., McLellan-Hall, S., Wong, A. S. W., Skippen,

P., & Cavanagh, J. F. (2019). Frontal theta predicts specific cognitive control-induced

behavioural changes beyond general reaction time slowing. *NeuroImage*, *189*(August

2018), 130–140. https://doi.org/10.1016/j.neuroimage.2019.01.022

Cunningham, C. A., & Egeth, H. E. (2016). Taming the White Bear: Initial Costs and

Eventual Benefits of Distractor Inhibition. *Psychological Science*, *27*(4), 476–485.

https://doi.org/10.1177/0956797615626564

de Vries, I. E. J., Savran, E., van Driel, J., & Olivers, C. N. L. (2019). Oscillatory

Mechanisms of Preparing for Visual Distraction. *Journal of Cognitive Neuroscience*,

31(12), 1873–1894. https://doi.org/10.1162/jocn_a_01460

de Vries, I. E. J., Slagter, H. A., & Olivers, C. N. L. (2020). Oscillatory Control over

Representational States in Working Memory. *Trends in Cognitive Sciences*, *24*(2), 150–

162. https://doi.org/10.1016/j.tics.2019.11.006

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*(1), 9–21.

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual*

Review of Neuroscience, *18*, 193–222.

Dugué, L., Marque, P., & VanRullen, R. (2014). Theta Oscillations Modulate Attentional

Search Performance Periodically. *Journal of Cognitive Neuroscience*, *27*(5), 945–958.

https://doi.org/10.1162/jocn_a_00755

Eimer, M. (2014). The neural basis of attentional control in visual search. *Trends in*

Cognitive Sciences, *18*(10), 526–535. https://doi.org/10.1016/j.tics.2014.05.005

Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: A flexible statistical

power analysis program for the social, behavioral, and biomedical sciences. *Behavior*

Research Methods, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Ferrante, O., Patacca, A., di Caro, V., della Libera, C., Santandrea, E., & Chelazzi, L. (2018).

Altering spatial priority maps via statistical learning of target selection and distractor

filtering. *Cortex*, *102*, 67–95. https://doi.org/10.1016/j.cortex.2017.09.027

Geng, J. J., Won, B.-Y. Y., & Carlisle, N. B. (2019). Distractor Ignoring: Strategies, Learning,

and Passive Filtering. *Current Directions in Psychological Science*, *28*(6),

096372141986709. https://doi.org/10.1177/0963721419867099

Han, S. W., & Kim, M. S. (2009). Do the Contents of Working Memory Capture Attention?

Yes, But Cognitive Control Matters. *Journal of Experimental Psychology: Human*

Perception and Performance, *35*(5), 1292–1302. https://doi.org/10.1037/a0016452

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha

activity: gating by inhibition. *Frontiers in Human Neuroscience*, *4*, 186.

Kelly, S. P., Lalor, E. C., Reilly, R. B., & Foxe, J. J. (2006). Increases in Alpha Oscillatory

Power Reflect an Active Retinotopic Mechanism for Distracter Suppression During

Sustained Visuospatial Attention. *Journal of Neurophysiology*, *95*(6), 3844–3851.

https://doi.org/10.1152/jn.01234.2005

Kiyonaga, A., Egner, T., & Soto, D. (2012). Cognitive control over working memory biases

of selection. *Psychonomic Bulletin and Review*, *19*(4), 639–646.

https://doi.org/10.3758/s13423-012-0253-7

Kok, P., Mostert, P., & de Lange, F. P. (2017). Prior expectations induce prestimulus sensory

templates. *Proceedings of the National Academy of Sciences*, *114*(39), 10473–10478.

https://doi.org/10.1073/pnas.1705652114

Leber, A. B., Gwinn, R. E., Hong, Y., & O'Toole, R. J. (2016). Implicitly learned

suppression of irrelevant spatial locations. *Psychonomic Bulletin & Review*, *23*(6),

1873–1881. https://doi.org/10.3758/s13423-016-1065-y

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data.

Journal of Neuroscience Methods, *164*(1), 177–190.

Moher, J., & Egeth, H. E. (2012). The ignoring paradox: Cueing distractor features leads first

to selection, then to inhibition of to-be-ignored items. *Attention, Perception, and*

Psychophysics, *74*(8), 1590–1605. https://doi.org/10.3758/s13414-012-0358-0

Moher, J., Lakshmanan, B. M., Egeth, H. E., & Ewen, J. B. (2014). Inhibition Drives Early

Feature-Based Attention. *Psychological Science*, *25*(2), 315–324.

https://doi.org/10.1177/0956797613511257

Myers, N. E., Rohenkohl, G., Wyart, V., Woolrich, M. W., Nobre, A. C., & Stokes, M. G.

(2015). Testing sensory evidence against mnemonic templates. *Elife*, *4*, e09000.

https://doi.org/10.7554/eLife.09000

Noonan, M. A. P., Adamian, N., Pike, A., Printzlau, F., Crittenden, B. M., & Stokes, M. G.

(2016). Distinct mechanisms for distractor suppression and target facilitation. *Journal of*

Neuroscience, *36*(6), 1797–1807. https://doi.org/10.1523/JNEUROSCI.2133-15.2016

Olivers, C. N. L. (2009). What drives memory-driven attentional capture? The effects of

memory type, display type, and search type. *Journal of Experimental Psychology:*

Human Perception and Performance, *35*(5), 1275.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Computational Intelligence and Neuroscience, *2011*.

Reeder, R. R., Olivers, C. N. L., Hanke, M., & Pollmann, S. (2018). No evidence for

enhanced distractor template representation in early visual cortex. *Cortex*, *108*, 279–282.

https://doi.org/10.1016/j.cortex.2018.08.005

Reeder, R. R., Olivers, C. N. L., & Pollmann, S. (2017). Cortical evidence for negative search

templates. *Visual Cognition*, *25*(1–3), 278–290.

https://doi.org/10.1080/13506285.2017.1339755

Reeder, R. R., & Peelen, M. v. (2013). The contents of the search template for category-level

search in natural scenes. *Journal of Vision*, *13*(3), 13. https://doi.org/10.1167/13.3.13

Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S., & D'Esposito, M. (2020). Causal

Evidence for a Role of Theta and Alpha Oscillations in the Control of Working Memory.

Current Biology, *30*(9), 1748-1754.e4. https://doi.org/10.1016/j.cub.2020.02.065

Soto, D., Heinke, D., Humphreys, G. W., & Blanco, M. J. (2005). Early, involuntary top-

down guidance of attention from working memory. *Journal of Experimental Psychology:*

Human Perception and Performance, *31*(2), 248–261. https://doi.org/10.1037/0096-

1523.31.2.248

Soto, D., Hodsoll, J., Rotshtein, P., & Humphreys, G. W. (2008). Automatic guidance of

attention from working memory. *Trends in Cognitive Sciences*, *12*(9), 342–348.

https://doi.org/10.1016/j.tics.2008.05.007

Spitzer, B., & Haegens, S. (2017). Beyond the status quo: A role for beta oscillations in

endogenous content (RE)activation. *Eneuro*, *4*(4).

https://doi.org/10.1523/ENEURO.0170-17.2017

Stilwell, B. T., Bahle, B., & Vecera, S. P. (2019). Feature-based statistical regularities of

distractors modulate attentional capture. *Journal of Experimental Psychology: Human*

Perception and Performance, *45*(3), 419–433. https://doi.org/10.1037/xhp0000613

Tanda, T., & Kawahara, J. I. (2019). Association between cue lead time and template-for-

rejection effect. *Attention, Perception & Psychophysics*.

Töllner, T., Conci, M., & Müller, H. J. (2015). Predictive distractor context facilitates

attentional selection of high, but not intermediate and low, salience targets. *Human*

Brain Mapping, *36*(3), 935–944.

Van Moorselaar, D., & Slagter, H. A. (2019). Learning What Is Irrelevant or Relevant:

Expectations Facilitate Distractor Inhibition and Target Facilitation through Distinct

Neural Mechanisms. *The Journal of Neuroscience : The Official Journal of the Society*

for Neuroscience, *39*(35), 6953–6967. https://doi.org/10.1523/JNEUROSCI.0593-

19.2019

van Zoest, W., Huber-Huber, C., Weaver, M., & Hickey, C. (2021). Strategic distractor

suppression improves selective control in human vision. *The Journal of Neuroscience*,

JN-RM-0553-21. https://doi.org/10.1523/JNEUROSCI.0553-21.2021

Vatterott, D. B., & Vecera, S. P. (2012). Experience-dependent attentional tuning of

distractor rejection. *Psychonomic Bulletin and Review*, *19*(5), 871–878.

https://doi.org/10.3758/s13423-012-0280-4

Wang, B., & Theeuwes, J. (2018). Statistical regularities modulate attentional capture.

Journal of Experimental Psychology. Human Perception and Performance, *44*(1), 13–

17. https://doi.org/10.1037/xhp0000472

Wen, W., Hou, Y., & Li, S. (2018). Memory guidance in distractor suppression is governed

by the availability of cognitive control. *Attention, Perception, & Psychophysics*, *80*(5),

1157–1168.

Wen, W., Zhang, Y., & Li, S. (2021). Gaze dynamics of feature-based distractor inhibition

under the prior-knowledge and expectation. *Attention, Perception & Psychophysics*.

https://doi.org/10.1101/2020.07.19.210781

Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic hidden states

underlying working-memory-guided behavior. *Nature Neuroscience*, *20*(6), 864–871.

https://doi.org/10.1038/nn.4546

Woodman, G. F., & Luck, S. J. (2007). Do the Contents of Visual Working Memory

Automatically Influence Attentional Selection During Visual Search? *Journal of*

Experimental Psychology: Human Perception and Performance, *33*(2), 363–377.

https://doi.org/10.1037/0096-1523.33.2.363

Worden, M. S., Foxe, J. J., Wang, N., & Simpson, G. v. (2000). Anticipatory biasing of

visuospatial attention indexed by retinotopically specific alpha-band

electroencephalography increases over occipital cortex. *The Journal of Neuroscience：*

The Official Journal of the Society for Neuroscience, *20*(6), RC63.

https://doi.org/10.1523/JNEUROSCI.20-06-j0002.2000

Zhang, Z., Gapelin, N., & Carlisle, N. B. (2020). Probing early attention following negative

and positive templates. *Attention, Perception, & Psychophysics*, *82*(3), 1166–1175.

https://doi.org/10.3758/s13414-019-01864-8

Figure 1. Task and behavioral results. (A) Task. The color cue indicated the to-be-ignored

color in the search array. Participants were asked to find the target (diamond) and judge

whether the bar inside of it was clockwise or counter-clockwise tilted relative to the mid-

vertical. In the memory test, participants were instructed to select the color patch that

matched the cue. (B) Search RT. Gray lines represent individuals' data. Errorbars represent

between-subject standard errors. (C) Suppression benefit index. Red and blue dots represent

the individuals' benefit indices in the fixed-cueing and varied-cueing conditions, respectively.

Black lines represent the averaged benefit indices across participants. (D) Suppression benefit

indices of RT percentile bins. Errorbars represent the standard deviations of 5000 times

bootstrapped means. (E) Accuracy of visual search task and memory test. Errorbars represent

the standard deviations of 5000 times bootstrapped means. $* p < .05$, $** p < .01$, $*** p < .001$,

$\underline{*} p < .001$ (two-tailed).

684
Figure 2. Results of color-decoding during retention interval. (A) Time-series of decoding

accuracy in the fixed-cueing and varied-cueing conditions. The shaded areas represent the

standard deviation of 5000 times bootstrapped mean. The solid/dashed vertical line indicate

the cue onset/offset. Lines below the x-axis indicated significant time-clusters for each

condition and the black lines correspond to the significant time-clusters for condition

difference. (B) Cross-temporal decoding of cued color in the two cueing conditions. Alpha-

blending was performed on insignificant time-points for better illustration. (C) Difference

map of cross-temporal decoding. Black outlined areas indicate significantly higher

decodability in the fixed-cueing condition compared to the varied-cueing condition as

revealed by cluster-based permutation (alpha = .05, cluster-based nonparametric alpha = .05,

cluster statistic = sum, two-tail, permutation times = 10000). (D) Decoding accuracies are

shown for the low and high suppression benefit groups in both cueing conditions. Gray

circles represent individual participants' decoding accuracy, which is the average of the time-

resolved decoding accuracies during retention interval (0~1600 ms). Errorbar represensts the

between-subject standard error. * $p <.05$. (E) Cross-condition decoding. F2V stands for the

decoding result when trained on data of the fixed-cueing condition and tested on data of the

varied-cueing condition. The black time series is the average of bi-directional cross-condition

decoding accuracy.

Figure 3. Multivariate topographical pattern similarity analysis. (A) Similarity matrix. C1F

refers color1 in the fixed-cueing condition. Each row represents a given individual trial while

each colume represents the averaged topographyical pattern of each color. Topographical

similarities of highlighted filled grids were used for statistical analysis. (B) Schematic

illustration of the topographical pattern similarity. The left panel is the correlation time series

of the example trial and the averaged pattern during retention interval. The right side are the

topographies of the example trial and the averaged pattern at three time points as indicated by

the black circles in the time series. (C) Similarity scores. Errorbars correspond to the within-

subject standard error.

714
Figure 4. Oscillatory activities during retention interval. (A) Top: Posterior lateralized alpha

power in the fixed-cueing and varied-cueing conditions. The shaded areas represent the

between-subject standard error. The solid/dashed vertical line indicated the cue onset/offset.

Lines below the x-axis indicated significant time-clusters for each condition with the black

line corresponding to their difference. Bottom: Topography of the averaged lateralized alpha

power during 300~800 ms. Topographical maps indicate the difference values of alpha power

between left-cue minus right-cue trials. Since contralateral alpha power would decrease

relative to ipsilateral alpha power when distractor cues are attended, left posterior electrodes

are positive in this plot. The black dots highlight the channels used for alpha lateralization

analysis (P5/6, P7/8, PO7/8). (B) Time-frequency map extracted from mid-frontal electrodes

(FCz, FC1, FC2, Fz, F1, F2). Dotted contours were adopted from the time-frequency

difference map (varied minus fixed). Significant time-frequency clusters from the non-

parametric permutation test are outlined by the dark contour. Data was extracted from mid-

frontal electrodes. Right-bottom: Topographical distribution of mid-frontal theta activity (3~7

Hz, 0.1~1.6 s) for the fixed-cueing and varied-cueing conditions. Black dots correspond to

the channels used for time-frequency analysis.

Figure 5. Mid-frontal theta after search array onset. (A) Top: Time-frequency map after

search array onset. The significant time-frequency cluster from the non-parametric

permutation test was outlined by the dark contour. Topography reflects averaged theta power

during 0.2~0.7s. The time window was chosen to avoid stimulus-driven and response-related

contaminates. The average search RTs across conditions was 735 ms. The black dots

indicated the channels used for power analysis (FCz, FC1, FC2, Fz, F1, F2). Bar plot shows

the average mid-frontal theta power within time window of interest. Errorbars represent the

between-subject corrected standard error. (B) Spearman-rank correlations between frontal

theta power and RTs in matched trials and neutral trials of two cueing conditions. The shaded

area represents the 95% confidence interval for the fitted line.