Tributyltin exposure leads to increased adiposity and reduced abundance of

leptogenic bacteria in the zebrafish intestine

Sol Gómez de la Torre Canny[1*], Olaf Mueller[1], Camil V. Craciunescu[1], Bruce

Blumberg[2], and John F. Rawls[1*]

[1] Department of Molecular Genetics and Microbiology, Duke Microbiome Center, Duke

University School of Medicine, Durham, NC 27710

[2] Department of Developmental and Cell Biology, and Pharmaceutical Sciences,

University of California, Irvine, CA 92697-2300

*correspondence to sgdltc@gmail.com and john.rawls@duke.edu

ABSTRACT

The chemical obesogen tributyltin (TBT) is known to promote fat storage in adipose

tissue through direct action on vertebrate cells. TBT also has direct toxic effects on

microorganisms, raising the possibility that TBT may also promote fat storage in

vertebrates by altering their microbiota. Here we show that exposure of conventionally-

reared post-embryonic zebrafish to TBT results in increased adiposity, reduced body

size, and altered intestinal microbiota composition including reduced relative abundance

of *Plesiomonas* bacteria. To test if those microbiota alterations affected host adiposity,

we exposed conventionally-reared zebrafish to intestinal bacterial strains representative

of TBT-altered taxa. We found that introduction of a *Plesiomonas* strain into

conventionally-reared zebrafish was sufficient to reduce adiposity and alter intestinal

microbiota composition. Using new long-term gnotobiotic zebrafish husbandry methods,

we found that colonization of germ-free zebrafish with *Plesiomonas* was sufficient to

reduce host adiposity. Together these results show the leptogenic activity of

Plesiomonas on zebrafish hosts, indicating that the ability of TBT to increase adiposity

in vivo may be due in part to TBT-mediated modification of the abundance of leptogenic

bacteria like *Plesiomonas*. These findings underscore how complex reciprocal

interactions between animals and their microbial and chemical environments can

influence energy balance and metabolic health.

IMPORTANCE

Obesogens are environmental chemicals that promote fat storage and are generally

thought to exert this effect directly on animal cells. Using zebrafish, we show that the

obesogen tributyltin can also promote fat storage by acting upon intestinal microbiota

via reduction of bacteria that are sufficient to reduce fat storage.

37
38
39
INTRODUCTION

Energy balance is a central aspect of animal physiology. When energy intake

surpasses energy expenditure (i.e., positive energy balance), excess energy is stored

as triglycerides in white adipocyte cells constituting adipose tissue (1, 2). Obesity is a

condition of excessive accumulation of adipose tissue relative to body size. Obesity is a

growing global public health challenge (3, 4) and a major risk factor for other non-

communicable diseases (5, 6). Identification of environmental factors promoting

sustained positive energy balance and expansion of adipose tissue is therefore an

important goal (7). The intestinal microbiota has emerged as a key environmental factor

modulating adipose tissue and body size. Mice reared germ-free (GF) in the absence of

microorganisms have reduced adipose tissue (8) and body size (9) compared to mice

colonized with their normal microbiota. Moreover, transplantation of fecal microbial

communities into GF recipient mice from identical twins discordant for obesity (10), or

from individuals with or without a form of malnutrition called kwashiorkor (11),

recapitulated the respective phenotypes of the human host. Thus, both the presence

and specific composition of gut microbiota have the potential to alter energy storage in

adipose tissue and body size in vertebrates.

Interaction between the gut microbiota and their vertebrate hosts can be further

impacted by xenobiotics (i.e., chemicals not produced by metabolisms naturally

occurring in organisms and the environment) (12). Some xenobiotics can alter the

composition of intestinal microbiotas and alter host physiology (13-16). Also, xenobiotics

can be chemically transformed by intestinal microbiota thus modifying their toxicity,

bioavailability, and efficacy when used as therapeutics (17-20).

One class of xenobiotic chemicals particularly relevant to energy balance are

obesogens, which are defined by their ability to promote adipose tissue accumulation

and alter metabolism (21, 22). The organotin compound tributyltin (TBT) is a

prototypical obesogen. TBT was once broadly used in antifouling paints to inhibit growth

of organisms on the submerged surfaces of ships, and consequently accumulated in

marine sediments and organisms prior to the ban on its use (23, 24). Although TBT

toxicity was first identified as causing imposex in neogastropods (25, 26), it was later

shown to act as an obesogen through *in vivo* studies in multiple tetrapod species (27-

31). The effects of TBT in fishes remain less understood, but include increased visceral

adipose tissue in the rare minnow (*Gobiocypris rarus*) (32) and inhibited fat mobilization

from adipose tissue in response to starvation in zebrafish (*Danio rerio*) (33, 34).

Importantly, these *in vivo* studies were conducted in conventionally-reared (CR) animals

colonized with their normal microbiota. Thus, it remains unclear whether TBT promotes

adiposity via direct effects on host tissues or indirectly via their associated microbiota.

Indeed, TBT directly promotes adipocyte differentiation in vertebrate cells *in vitro*, likely

acting upon peroxisome proliferator-activated (PPAR) γ and retinoid X (RXR) receptors

(27, 28, 35-40). The possibility that TBT might directly affect microbial physiology is

supported by studies in bacteria, fungi, and other environmental microbial communities

(41-49). Variable levels of microbial susceptibility to TBT have been linked to

differences in TBT adsorption, bioaccumulation, enzymatic transformation, efflux and

other adaptations (47, 50-58). It is therefore possible that TBT could also directly impact

the composition and activity of host-associated microbial communities. Indeed, recent

studies showed TBT exposure affects gut microbiota composition in rats and mice (59-

61). However, the specific role of TBT-modified microbes on host adiposity remains

untested, and such TBT-microbiota interactions in other vertebrates are yet unexplored.

In this study, we used the zebrafish to test if the host-associated microbiota and

TBT interact to modulate host body size and adiposity. We and others have previously

defined the composition of the conventional zebrafish gut microbiota (62-64) and

developed short-term gnotobiotic methods for larval zebrafish to reveal impacts of

specific microbiota members on host physiology (62, 65-67). We and others also

showed that zebrafish develop white adipose tissues during later post-embryonic stages

that display extensive homologies with those of mammals (68-71). Using TBT and

microbial exposures under conventional as well as new long-term gnotobiotic husbandry

we developed for this work, we tested the hypothesis that TBT affects vertebrate

adiposity by modifying microbiota composition.

RESULTS

Tributyltin exposure affects body size and adiposity

We conducted a repeated cross-sectional study to evaluate the impact of

tributyltin chloride (TBT) exposure on post-embryonic zebrafish (Fig. 1A). We raised

zebrafish embryos conventionally until 18 days post-fertilization (dpf) when the intestinal

microbiota had been established (64, 72). We then transferred the experimental cohort

to a modified flow through aquaculture system (mFTS, see Methods), where we initiated

exposure to TBT (0.01, 0.1, and 1 µg/l) or DMSO vehicle control in tank water at 22 dpf.

These TBT concentrations were selected based on ranges observed in the environment

and in animal tissues (73-76). Because adipose tissue development in zebrafish

depends on body size, particularly on standard length (SL) (69, 70), we size-matched

fish prior to exposure. We sampled fish from individual tanks at 7, 14, and 21 days post-

exposure (dpE) to measure SL (77) and adiposity (defined as the ratio of the total two-

dimensional area of adipose tissues per total body area excluding fins) (78), before

dissecting their intestines for 16S rRNA gene sequencing analysis of the intestinal

microbiota. We also sampled the tank water to evaluate concomitant effects of TBT on

the aquatic microbiota (Fig. 1A, see Methods).

We detected no significant differences in zebrafish SL or adiposity between

treatment groups (0 dpE; Fig. 2A and B) or replicate tanks (data not shown) prior to

exposure. Although no differences in adiposity between the control and all three TBT

concentrations were observed at 7 or 14 dpE, exposure to 1 µg/l TBT did cause an

increase in adiposity compared to controls at 21 dpE (P = 0.0120; Fig. 2B and D). In

addition to greater adiposity, we observed a reduction of SL in zebrafish exposed to 1

µg/l of TBT at 14 and 21 dpE (P = 0.0294 and P = 0.0105, respectively; Fig. 2A and C).

Other measurements of body size including height at anterior of anal fin (HAA) (77) and

total body area were only significantly reduced at 21dpE and only in the 1 µg/l TBT

treatment group (P = 0.0042 and P = 0.0083; Fig. S1). Therefore, 1 µg/l TBT inhibits

body growth and promotes adiposity in zebrafish after 21 days of exposure.

Since zebrafish display substantial inter-individual variation in SL even when

age-matched (Fig. 2A), we combined data from 7, 14, and 21 dpE to assess if TBT

treatment impacted the overall developmental relationship between SL and total

adipose tissue area (69, 70). A linear relationship between SL and adipose tissue area

was maintained in all treatment groups (all $P \geq$ 0.0853), but the slope of the 1 µg/l TBT

group was significantly greater than DMSO vehicle control (P = 0.0005; Figure 2E).

Together, this indicates that exposure to 1 µg/l TBT alters the developmental

relationship between SL and adipose tissue accumulation.

Exposure to TBT is not a major determinant of intestinal or aquatic microbiota

composition but does alter the relative abundance of specific bacterial taxa

Having observed the effects of TBT on adiposity and body size, we tested if

chemical exposure also affected zebrafish intestinal or tank water microbiotas. We

performed 16S rRNA gene sequencing of whole intestines and tank water sampled

during TBT exposure (Fig. 1A). Intestinal and aquatic microbiota composition were

significantly different from each other at all time points and TBT concentrations (all $P \le$

0.003; Table S1). Within each TBT concentration, the composition of intestinal

microbiota at each time point was significantly different from all other time points (all $P \le$

0.028; Table S1). Therefore, sample type and developmental age were major

determinants of intestinal microbiota composition. In contrast, TBT exposure had a

relatively minor effect on intestinal and aquatic microbial community composition. We

found only two instances during the study in which the composition of intestinal

communities from zebrafish treated with TBT significantly differed from those in vehicle

control tanks: at 7 dpE when compared to 0.1 μg/l TBT (P = 0.0030; Table S1) and at

14 dpE when compared to 1 μg/l TBT (P = 0.0430; Table S1). Notably, it was at 14 dpE

that we first observed SL differences in zebrafish treated with 1 μg/L TBT. However, a

significant microbiota composition difference in zebrafish treated with 1 μg/l TBT was no

longer detected at the end of the experiment at 21 dpE, when both adiposity and body

size phenotypes were evident (Table S1, Fig. 2, and Fig. S1). Also, we found that

aquatic communities from control tanks were only significantly different compared to 1

μg/l or DMSO at 7 and 21 dpE (P = 0.028 and P = 0.029, respectively; Table S1).

Comparison of indices and estimators of α-diversity between controls and all TBT

concentrations revealed no differences in the total number of OTUs, Chao1, Shannon or

Faith's phylogenetic indices. Together, these data indicate that TBT does not markedly

alter the overall composition or diversity of the microbial communities in zebrafish

intestine or their tank water.

Although TBT exposure did not influence the overall composition of microbial

communities, we reasoned that it may affect individual microbial taxa. We therefore

applied linear discriminant analysis effect size analysis using LEfSe (79). LEfSe

identified biomarker taxa through pairwise comparisons of the relative abundance of

OTUs in microbial communities from zebrafish treated with DMSO or with 1 µg/l TBT

(the only TBT concentration that rendered body size and adiposity phenotypes) in every

timepoint of the exposure. We discovered 51 biomarker taxa at the genus level in the

zebrafish intestine (Table S2) and 79 in water (Table S3) associated with either of the

selected chemical treatments at any of the time points sampled (7, 14, and 21 dpE).

Further, only 4/51 biomarkers in intestinal communities and 9/79 in aquatic communities

were found in more than one of the time points sampled (Tables S2 and S3), in accord

with our observation above that age is a major determinant of community composition

(Table S1). Furthermore, only 7/51 biomarkers in the intestine were also identified as

biomarkers in the aquatic communities, in accord with the intestinal and aquatic

microbial communities being distinct (Table S1). Finally, most biomarkers in the

intestine (47/51) and water (67/79) were only significantly altered at 14 or 21 dpE, the

same time points when host phenotypes were also altered (Tables S2 and S3; Fig. 2).

Together, these results identify bacterial genera in the zebrafish intestine and tank

water that were altered by treatment with 1 µg/l TBT.

Supercolonization of the zebrafish intestine with Plesiomonas ZOR0011 reduces

adiposity but does not affect body size

We hypothesized that the effects of TBT on adiposity and body size may be

mediated in part by the observed alterations in the relative abundance of specific

bacterial genera. To test that possibility, we exposed CR zebrafish to specific bacterial

isolates for 14 days (Figure 1B; see Methods). We reasoned that this supercolonization

approach would allow us to test the effects of increasing abundance of specific

community members in an otherwise normal intestinal microbial community, presumably

resembling the specific changes in microbiota composition induced by TBT exposure.

We referenced LEfSe-identified intestinal biomarker taxa associated with TBT

exposure and alteration in adiposity and body size phenotypes (Fig. 2; Table S2)

against a collection of bacterial strains previously isolated from the zebrafish intestine

(64, 80). We found strains representing biomarker genera *Plesiomonas* (Class

Gammaproteobacteria, Order Enterobacterales; strain *Plesiomonas* ZOR0011), *Vibrio*

(Class Gammaproteobacteria, Order Vibrionales; strain *Vibrio* ZWU0020), and

Chitinibacter (Class Betaproteobacteria, Order Neisseriales; strains *Chitinibacter*

ZOR0017 and *Chitinibacter* ZOR0013), all of which grow aerobically and were originally

isolated at similar ages to that of our experimental cohort (64)*. In accord with our LEfSe

results, the relative abundance of *Chitinibacter* OTU in the intestines of fish exposed to

1 µg/l TBT was higher compared to DMSO controls at 14 and 21 dpE (*P* = 0.0045 and *P*

= 0.0224, respectively; Fig. 3A). Further, the relative abundances of *Vibrio* and

Plesiomonas OTUs in the intestines of fish exposed to 1 µg/l TBT were lower compared

to DMSO controls at 14 dpE (*P* = 0.0335 and *P* = 0.0205, respectively; Fig. 3C and E).

In addition, *Plesiomonas* OTU was also more abundant in the intestines of fish treated

with 0.1 µg/l TBT at 7 dpE (*P* = 0.0098; Fig. 3E). The relative abundances of these three

OTUs remained unaltered in the tank water at all time points and all TBT concentrations

(Fig. 3B, D, and F). These results show that TBT altered the relative abundance of all

three selected bacterial genera (*Plesiomonas*, *Vibrio*, and *Chitinibacter)* in the intestine

at 14 dpE, the time point of TBT exposure when the earliest host phenotype was first

observed (Fig. 2), and one of the only two timepoints in which TBT was a major

determinant of microbiota composition (Table S1). Based on the directionality of the

TBT-induced changes in the relative abundance of these OTUs by 14 dpE, we

hypothesized that an increase in *Chitinibacter* would promote adiposity and reduce body

size; while an increase in *Plesiomonas* or *Vibrio* would reduce adiposity and increase

body size.

Zebrafish entering the supercolonization experiment were size-matched to those

that had entered the *in vivo* TBT exposure experiment to ensure they were at similar

developmental stages. Mock supercolonization using only autoclaved RAS water was

used as a negative control. We sampled fish from individual tanks to evaluate the

impact of the bacterial treatments on body size, adiposity, as well as intestinal and

aquatic microbial communities (Fig. 1B). Prior to exposure, measurements of zebrafish

adiposity and body size did not reveal differences between experimental groups (Fig.

4A and B). However, after 14 days of supercolonization, animals treated with

Plesiomonas ZOR0011 had a significant reduction in adiposity (P = 0.0334; Fig. 4B and

D) without effects on standard length or other body size measurements (Fig. 4A and C;

Fig. S1). The other three bacterial strains did not significantly affect either of these traits.

The observation of the predicted adiposity phenotype after supercolonization with

Plesiomonas ZOR0011 supports the possibility that the reduction in the relative

abundance of *Plesiomonas* bacteria following TBT exposure may mediate, at least

partially, the resulting host phenotypes.

We next used 16S rRNA gene sequencing to verify if supercolonization

effectively increased the relative abundance of the introduced bacterial strains in the

water or zebrafish intestinal microbial communities. We examined the relative

abundance of *Chitinibacter*, *Plesiomonas*, and *Vibrio* OTUs in the aquatic communities

after 14 days of supercolonization with the four different intestinal bacterial strains.

None of those bacterial treatments significantly increased the relative abundance of

Chitinibacter, *Plesiomonas* or *Vibrio* OTUs in the tank water when compared to negative

controls (Fig. 4E). Nevertheless, results from the more sensitive LEfSe analysis

revealed that the OTUs assigned to each introduced bacterial strain were in fact

biomarker genera of their corresponding treatment group (Table S4). Therefore,

supercolonization effectively increased the relative abundance of introduced bacterial

genera in the aquatic environment, thereby potentially allowing them to also colonize

the zebrafish intestine. However, changes in the relative abundance of members of the

intestinal microbial communities of supercolonized animals were unique to the strain

Plesiomonas ZOR0011. The relative abundance of OTU *Plesiomonas* in the intestine

was increased upon supercolonization with *Plesiomonas* ZOR0011 (P = 0.0022; Fig.

4E), a result confirmed by LEfSe analysis (Table S5). For the other three strains tested,

the respective OTUs were not altered nor identified as biomarkers by LEfSe analysis.

These data support the hypothesis that the host adiposity phenotype caused by *in vivo*

exposure to 1 µg/l TBT may be mediated in part by the chemically-induced alteration of

the relative abundance of *Plesiomonas* in the zebrafish intestine.

Supercolonization with Plesiomonas ZOR0011 alters zebrafish intestinal

microbiota

We next asked if supercolonization with *Plesiomonas* ZOR0011 or any of the

other selected bacterial strains, altered the composition of the intestinal or aquatic

microbiota. As expected, composition of intestinal and aquatic microbiotas remained

significantly different regardless of the treatment (all $P \leq 0.008$; Fig. S2; Table S6).

Supercolonization with the different bacterial strains did not alter α-diversity of aquatic or

intestinal communities (total number of OTUs, Chao1, Shannon or Faith's phylogenetic

indices). However, the composition of intestinal microbial communities in zebrafish

supercolonized with *Plesiomonas* ZOR0011 or *Chitinibacter* ZOR0017 displayed

significant differences compared to negative controls ($P = 0.001$ and $P = 0.027$

respectively; Fig. S2, Table S6). Also, the intestinal microbial communities in zebrafish

supercolonized with *Plesiomonas* ZOR0011 and *Chitinibacter* ZOR0017 are distinct (P

$= 0.001$; Fig. S2, Tables S6). These results show that supercolonization with

Plesiomonas ZOR0011 and *Chitinibacter* ZOR0017 had distinct effects on intestinal

microbiota composition.

Monoassociation of zebrafish with Plesiomonas ZOR0011 inhibits adiposity and

body size

Supercolonization with *Plesiomonas* ZOR0011 reduced adiposity, but also

altered the composition of the established intestinal microbiota. We therefore sought to

test the sufficiency of *Plesiomonas* ZOR0011 to directly influence adiposity and body

size in the absence of other microbes using a gnotobiotic approach. Although methods

for gnotobiotic husbandry of zebrafish embryos and larvae have been established, they

were not optimized to support growth of zebrafish into later stages when adipose

tissues develop (68-70). We therefore developed new methods for long-term gnotobiotic

zebrafish husbandry (see Methods) to raise germ-free (GF) zebrafish to 14 dpf before

conducting a monoassociation with *Plesiomonas* ZOR0011 or *Chitinibacter* ZOR0013.

Chitinibacter ZOR0013 was selected as a control because it did not elicit a phenotype

nor altered the composition of the intestinal microbiota after supercolonization. As an

additional reference control, we tested the impact of TBT on zebrafish body size and

adiposity in the absence of any microbes by exposing GF zebrafish to 1 µg/l TBT or

DMSO. We subjected GF zebrafish to these treatments for 21 days to match the length

of adiposity phenotype during the TBT exposure (Fig. 4B and D). We detected no

differences in mortality (all $P \geq 0.1986$ at 7 dpE/dpC; and $P \geq 0.5213$ at 21 dpE/dpC) or

the density (all $P \geq 0.8869$ at 7 dpE/dpC; and $P \geq 0.9474$ at 21 dpE/dpC) of the flasks

between the chemical or bacterial conditions tested (Methods). Whereas there were no

phenotypical alterations in fishes after 7 days of treatment, we observed a reduction in

adiposity of gnotobiotic zebrafish 21 days post-colonization (dpC) with *Plesiomonas*

ZOR0011 ($P = 0.0399$; Fig. 5A, B, and D), resembling the effect of supercolonization of

CR zebrafish with this bacterial strain (Fig. 4B and D). In contrast to its

supercolonization phenotype, monoassociation with *Plesiomonas* ZOR0011 also

reduced body size of gnotobiotic zebrafish ($P = 0.0256$; Fig. 5A and C). Finally, neither

monoassociation with *Chitinibacter* ZOR0013 or exposure to 1 µg/l TBT in the GF

background elicited body size or adiposity phenotypes after 21 days (Fig. 5). Together,

these results show that *Plesiomonas* ZOR0011 is sufficient to directly reduce adiposity

and body size, and suggest that obesogenic effects of TBT on zebrafish may require the

presence of microbiota.

TBT affects specific members of the intestinal microbiota directly in the absence

of the zebrafish host

The studies above did not resolve whether TBT affects *Plesiomonas* and other

intestinal bacteria *in vivo* directly, or indirectly through host-dependent processes. To

investigate the direct effect of TBT on zebrafish intestinal bacteria, we harvested

intestinal microbiota from CR zebrafish and cultured them *in vitro* in non-selective rich

media containing different concentrations of TBT similar to those tested in the *in vivo*

chemical exposure (0.01, 0.1, and 1 µg/l TBT) and two additional higher concentrations

(10 and 1000 µg/l TBT) anticipating the presence of highly-resistant bacteria observed

in some aquatic environments (42). Zebrafish used as microbiota donors were size-

matched to zebrafish from control tanks at the endpoint of the *in vivo* TBT exposure

(see Methods). We incubated the communities *in vitro* for 24 hours at room temperature

under aerobic and anaerobic conditions and used 16S rRNA gene sequencing to

interrogate their composition. Of a total of 272 OTUs detected across the different

treatments and the input community, only 62 were observed both in the input

community and at least one condition after *in vitro exposure to TBT*. Nevertheless, most

of the OTUs detected in the *in vitro* cultures (85/90) were also observed in the intestinal

microbiota of zebrafish at 21 dpE during the *in vivo* chemical exposure in at least one of

the chemical conditions. Therefore, although OTUs from the *in vitro* experiments

represent only the culturable fraction of an *in vivo* community, they provide a relatively

broad representation of OTUs found in the intestinal microbiota of zebrafish from the

same size that underwent TBT exposure *in vivo* (Table S7).

As in the *in vivo* chemical exposures, diversity analysis showed that TBT was not

a major determinant of the composition of the culturable zebrafish intestinal microbiota

at any of the concentrations tested *in vitro* (all $P \geq 0.05$, Table rename). In addition, α-

diversity was not altered by the exposure to TBT (total number of OTUs, Chao1,

Shannon or Faith's phylogenetic indices). LEfSe analysis identified 37 biomarker taxa of

in vitro TBT exposure at the genus level, in at least one of the pairwise comparisons

between the DMSO and the different TBT concentrations (Table S8). Of the 37

biomarker genera, 11 of them were also identified as biomarkers of the *in vivo* exposure

to 1 µg/l TBT at time points when zebrafish phenotypes were detected (14 and 21 dpE,

Table S8). These 11 biomarkers included genera *Vibrio and Plesiomonas*. *Plesiomonas*

showed an increased relative abundance (2.7-fold) only at the highest TBT

concentration of 1000 µg/l under aerobic conditions. In contrast, the lower TBT

concentrations used in the *in vivo* chemical exposure experiment (0.01, 0.1, and 1 µg/l

TBT) did not affect the relative abundance of *Plesiomonas* in microbiota cultured *in vitro*

(Tables S7 and S8). These results suggest that *Plesiomonas* sensitivity to TBT may be

determined in part by the host.

DISCUSSION

TBT is considered a prototypical obesogen (27-31). In zebrafish and other

teleosts, TBT promotes the accumulation of lipids in different tissues and organs (32,

81-88). Exposure of post-embryonic zebrafish to TBT impaired adipose tissue

mobilization in response to food restriction (33, 34), but its impact on adipose tissue

accumulation in fishes was unknown. Our results establish that a 21-day TBT exposure

promotes adiposity in zebrafish, indicating its obesogenic activity on teleost fishes. We

also found that the TBT-induced adiposity phenotype was accompanied by reduced

body size. This is in accord with reports that TBT compounds reduced body length in

zebrafish (89), other teleost species (90, 91) and in *Xenopus tropicalis* (92), as well as

body weight in other vertebrates (93-95), although these studies did not examine co-

occurring lipid or adipose phenotypes. In humans, stunted growth and other forms of

malnutrition coexist with obesity (96) and earlier manifestations of undernutrition may

predispose to obesity (97). We observed that the impacts of TBT on SL (by 14 dpE)

preceded overt impacts on overall body size and adiposity (by 21 dpE), but it remains

unclear if these adiposity and body size phenotypes are mechanistically linked. It is

possible that TBT-induced impairments in fat mobilization from adipose tissues in

zebrafish (34) reduce the energy available for growth. Identification of the direct and

indirect mechanisms by which TBT affects the relationship between adiposity and body

size is therefore warranted.

There is a growing appreciation that xenobiotics may exert their impacts on

animal health by altering the composition of their microbiotas (15, 16, 59-61, 98-100).

While TBT has well-documented direct actions on adipocyte differentiation and other

aspects of host physiology (27, 28, 35-40), it was only recently shown to alter the

composition of gut microbiota in rodents (59-61). In attempt to test if TBT-modified gut

microbiota is sufficient to affect host physiology, Zhan and coworkers transplanted TBT-

modified fecal microbiota into recipient mice pre-treated with antibiotics to deplete their

resident microbiota, and recapitulated adiposity and metabolic phenotypes observed in

donors (60). However, this experimental approach does not control for potential

confounding effects of antibiotic-resistant microbes, the direct effects of antibiotics on

the recipient host (101), or the possibility of bioaccumulation of TBT in fecal microbiota.

We found that TBT altered the relative abundance of a subset of intestinal bacterial

genera in CR zebrafish. Whereas the alterations in intestinal microbiota composition

was relatively modest, this information allowed us to identify TBT-affected bacterial

genera for further sufficiency tests. It also remains possible that, beyond these

compositional changes, TBT affects the functional potential of the microbiota as has

been demonstrated with other xenobiotics (102-104).

We developed multiple experimental approaches to test if microbes affected by TBT

exposure *in vivo* were themselves sufficient to evoke the phenotypes observed in their

hosts. Supercolonization is a strategy that introduces microbial strains into animals

already colonized with a microbiota, and has been effective for defining the impact of

the introduced microbes on the host and the intestinal ecosystem (105-107). By

supercolonizing CR zebrafish with available individual bacterial strains representing

TBT-affected genera, we found that *Plesiomonas* ZOR0011 was the only strain,

amongst those tested, sufficient to reduce adiposity in zebrafish. One advantage of the

supercolonization approach is that it inherently accommodates the potential for

microbial interactions. Indeed, *Plesiomonas* ZOR0011 supercolonization altered the

microbiota composition, making it unclear whether the impact of *Plesiomonas* ZOR0011

supercolonization on host adiposity was direct or indirect via alterations in the

abundance of other microbiota members. For these reasons we used a complementary

gnotobiotic approach to study the isolated effect of *Plesiomonas* ZOR0011.

Although methods for gnotobiotic zebrafish husbandry have been in use for over

17 years (62, 67, 108), almost all studies have focused on early free-feeding larval

stages. This postponed the need to develop nutrition and housing systems for raising

zebrafish into later life stages. The husbandry methods that we developed for this study

permit growth of zebrafish to standard lengths at which most adipose tissue depots

develop (70). We anticipate these methods will also be useful for testing the role of

microbial exposures on other aspects of post-embryonic physiology in the zebrafish.

Using these new methods, we showed that the exposure of GF zebrafish to a TBT

concentration that was phenotypical in CR zebrafish did not affect body size or

adiposity. This suggests that GF zebrafish may be resistant to TBT-induced

phenotypes, but additional studies are needed to determine if this is limited to specific

developmental stages or TBT dosages. Nevertheless, our combined approaches

showed that *Plesiomonas* ZOR0011 was sufficient to inhibit adiposity in zebrafish, both

in the absence and the presence of a conventional microbiota.

A major finding of this study is the discovery that a bacterial strain isolated from

the zebrafish intestine, *Plesiomonas* ZOR0011, has leptogenic effects on zebrafish

hosts. *Plesiomonas* bacteria are common members of the intestinal microbiota in

laboratory-reared zebrafish at all life stages and across multiple aquaculture facilities,

and have also been observed in adult zebrafish recently captured from their natural

habitat (63, 64). *Plesiomonas* abundance in the zebrafish intestine has also been

shown to be sensitive to other xenobiotics besides TBT (109-113), but this study is the

413 first to explore its direct physiologic effects on this host. Although the relative

414 abundance of *Plesiomonas* in the intestine was relatively low in our study, it had a

415 significant effect on host physiology and on microbiota composition. Other studies have

416 also underscored how members of the microbiota of relatively low abundance can have

417 significant effects on host-microbe commensalism (66, 114). The mechanisms by which

418 *Plesiomonas* ZOR0011 exerts its leptogenic effects on zebrafish and affects microbiota

419 composition remain unknown. The genus *Plesiomonas,* thought to consist of a single

420 species *P. shigelloides,* is common in aquatic ecosystems and inhabits diverse aquatic

421 vertebrates and invertebrates (115). *Plesiomonas* has also been detected in the

422 intestinal microbiota in healthy humans and some mammals, yet also frequently

423 associated with gastroenteritis in humans (115). We did not observe any overt

424 pathology or mortality in zebrafish supercolonized or monoassociated with *Plesiomonas*

425 ZOR0011, thus its sufficiency to reduce adiposity and body size may be consistent with

426 a role on digestive physiology or energy balance. Moreover, our recent finding that the

427 relative abundance of *Plesiomonas* members was markedly reduced by starvation in

428 adult zebrafish, suggests potential roles in host nutritional physiology (116). Further

429 studies are needed to discern the mechanisms by which *Plesiomonas* ZOR0011 exerts

430 its leptogenic effect.

431  Whereas the focus of this study was to assess impacts of TBT on gut microbiota

432 and its consequences *in vivo*, we also investigated the direct effect of TBT on zebrafish

433 microbiota *in vitro* in the absence of the host. Our results demonstrated that the impact

434 of xenobiotics like TBT on members of the zebrafish intestinal microbiota can differ

435 between *in vivo* and *in vitro* scenarios. Many genera including *Plesiomonas* were not

affected by phenotypical levels of TBT *in vivo* (0.1 and 1 µg/l TBT), and their relative

abundance *in vitro* was only significantly affected by much higher concentrations (1000

µg/l TBT). We were unable to test the effect of TBT exposure to ≥ 10 µg/l on intestinal

microbiota *in vivo* as this concentration caused >92% mortality as early as 1 dpE (see

Methods). The global increase in synthetic chemicals like TBT is recent in the shared

evolutionary history of vertebrate animals and their commensal bacteria with whom they

continue to co-evolve (117-119). The effect of chemical novelty on intestinal bacteria

like *Plesiomonas* is presumably an anthropogenic impact on organisms across the tree

of life, as individuals or as holobionts. Zebrafish, a well-established animal model for

human disease, can also serve as a model for wild and farmed fishes, vulnerable to

xenobiotic exposure and central to food security (120-122). Further understanding of the

specific effect of xenobiotics on intestinal microbiota can contribute to both planetary

sustainability and the improvement of human and animal health.

MATERIALS AND METHODS

Conventional zebrafish husbandry and obtention of experimental cohorts

All zebrafish experiments were approved by the Institutional Animal Care and

Use Committees of Duke University Medical Center under protocol A115-16-05 in

accordance with the Public Health Service Policy on the Human Care and Use of

Laboratory Animals under the United States of America National Institutes of Health

Office of Laboratory Animal Welfare (OLAW). Adult zebrafish (wild-type Ekkwill strain)

were raised under conventional conditions on a Pentair recirculating aquaculture system

(RAS) and set up for natural breeding to obtain the embryos constituting the different

experimental cohorts. Four adult fish (1:1 male to female ratio) from five families were

set in mating boxes for group spawning. The resulting clutches of embryos were

collected synchronously, transferred to Petri dishes containing RAS water, and

maintained at a density of ~70 fish per dish and incubated at 28 °C. Abnormal embryos

were removed during the first 24 hours post-fertilization (hpf). At ~36 hpf, embryos were

pooled and randomized into groups of 62 embryos which were transferred to clean 3-

liter tanks (30 tanks total). Each 3-liter tank, filled with 500 ml of RAS water, was kept

static until 5 dpf. At 5 dpf, the 3-liter tanks were connected to the RAS and maintained

according to our standard husbandry protocols (123) with one modification to feeding

before their transfer to the modular flow through aquaculture system (mFTS, see below)

where the experimental cohorts were housed during the experiments. The modification

was that, instead of receiving an exclusive powdered diet of Zeigler AP100 <50-micron

larval diet (Pentair, LD50-AQ) twice daily from 5 dpf until 14 dpf, larvae were

supplemented twice daily with 500 µl of *Artemia* (Brine Shrimp Direct, BSEACASE) that

was collected, rinsed and resuspended in an equal volume of RAS water.

The experimental cohort for the *in vivo* chemical exposure to TBT was

transferred to the mFTS on 18 dpf. On the day of transfer to the mFTS, zebrafish were

collected from the RAS in sterile 25 mm-deep Petri dishes, where they were

anesthetized using 0.2 g/L buffered Tricaine to measure their SL (77) for size-matching

individuals between 7 and 11 mm. The SL of the experimental cohort prior to TBT

exposure (0 days post-exposure, dpE) was of 8.39 ± 0.82 mm (mean ± SD), which was

measured at 22 dpf. A total of 20 pools of 30 individuals constituted the experimental

cohort during the *in vivo* chemical exposure study. For the supercolonization

experiments, the cohort was obtained in the same manner with the exception that

transfer to the mFTS occurred on 24 dpf. Zebrafish entering the supercolonization

experiment were size matched to those that had entered the TBT exposure to avoid

developmental differences between these two experiments. The SL of the experimental

cohort prior to bacterial supercolonization (0 days post-supercolonization, dpSC) was of

8.77 ± 0.84 (mean ± SD), which was measured at 27 dpf. 15 pools of 30 individuals

constituted the experimental cohort during the supercolonization study.

Modular flow through system (mFTS) construction and operation

Because TBT is likely degraded by filtration through activated carbon (124) and

UV exposure (125) which are used to maintain water quality on the RAS, we designed,

built and installed a modular flow-through system (mFTS) in one of the racks at our

RAS. The mFTS housed the experimental cohorts for both the *in vivo* TBT exposure

and the supercolonization experiments.

Each shelf in our facility fits five 10-liter polycarbonate tanks connected to the

RAS by five manifolds that deliver pre-conditioned RAS water into the experimental

tanks. The mFTS used the standard polycarbonate tanks from the RAS system which

we modified by drilling and installing polycarbonate food-grade beverage dispenser

faucets serving as spigots for draining the tanks during water exchanges (2712212,

Carlisle). Inside the tanks, the spigots were covered with fiberglass mesh (33105,

Lowe's) serving as baffles to prevent fish from escaping during water exchanges. The

spigots from the five tanks in each shelf were connected through fitted vinyl hoses to a

3/4-inch PVC pipe mounted with a slight angle of inclination with respect to the fish tank

shelf. This angle prevented backflow during the water exchanges. One PVC pipe was

used to collect the outflow of 5 tanks from a shelf. Four PVC-pipes were finally

connected, through a 90-degree 3/4-inch elbow, to a fitted 1-inch vinyl hose and

secured with an adjustable clamp (57188, Lowe's). The 1-inch vinyl hose ran

perpendicular to the fish tank shelves into a drain, to collect the outflow of the mFTS

system. Thus, the mFTS thus allowed us to drain several fish tanks simultaneously to

perform water exchange in a relatively automated and uniform manner. The outflow of

each fish tank was controlled by the location of the spigot along the height of the tank.

We installed the spigot to regulate an outflow of 5.5 L, with 2.5 L retained below the

spigot after draining. After draining, the tanks in the mFTS were refilled with the

conditioned water from the RAS system via the RAS manifold. The modified tanks in the

mFTS were initially filled with 7.5 L of conditioned RAS. A total of 1.6 volumes of water

were exchanged daily to maintain an adequate water quality.

The mFTS held twenty 10-liter modified polycarbonate tanks during the *in vivo*

chemical exposure to TBT, and fifteen during the supercolonization experiments. Water

quality parameters were assessed in the mFTS during the two studies. Conductivity was

measured in the conditioned RAS water before water exchange. pH, nitrate and nitrite,

and toxic ammonia were measured in the mFTS and in the conditioned water using test

kits according to the manufacturer's instructions (147008, 2745425, and 224100

respectively, HACH). Water quality parameters were measured on every tank in the

mFTS on the day before adding the experimental cohort to the tanks, and on every time

point sampled during the studies. The pH and conductivity in the conditioned water

used for the water exchange was recorded daily. The nitrite levels in the conditioned

water were 0 ppm, nitrate between 2 and 10 ppm, and toxic ammonia was 0 mg/l.

During the *in vivo* chemical exposure to TBT, the pH in the tanks ranged between 7 and

7.6, nitrite between 0 and 0.15 ppm, nitrate between 2 and 10 ppm, and toxic ammonia

between 0 and 0.00521 mg/l. The water pH and nitrate in the experimental tanks during

the chemical exposure was comparable to that of the conditions water, while the nitrite

and the toxic ammonia were higher. During the supercolonization experiments, water

pH in the experimental tanks ranged between 7.37 and 7.41, nitrite was 0, nitrate

ranged between 2 and 10 ppm, and toxic ammonia between 0 and 0.00106 mg/l. When

compared to the conditioned water, the pH was lower, the nitrite and nitrate were

comparable, and the toxic ammonia was slightly higher. Finally, the conductivity in the

RAS water used during the water exchanges during the *in vivo* exposure to TBT ranged

between 712 and 845; and during the supercolonization experiments, between 773 and

801. Overall, water quality parameters in the tanks were within the optimal ranges for

the zebrafish (126). The only substantial environmental difference between the mFTS

and the RAS was the temperature. In the RAS, the water temperature is maintained at

28.5 °C by an in-line heating element. In the mFTS, tanks were filled with conditioned

RAS water but then equilibrated to air temperature over time. Water temperature in the

tanks after the water exchange was recorded daily in the two studies. During the *in vivo*

exposure TBT, the average water temperature recorded in the afternoon was 22.3 °C;

and in the supercolonization experiments, 22.9°C. Notably, the water temperature

equilibrated with the room temperature during the evening, resulting in a lower water

temperature in the morning. During the *in vivo* exposure to TBT, the average water

temperature recorded in the morning was 19.1°C, and in the supercolonization

experiments 20.1°C. Because the mFTS was located inside our main zebrafish facility,

we were not able to change the room temperature or install space heaters.

During their housing in the mFTS, zebrafish received an exclusively live feed

diet. *Artemia* (Brine Shrimp Direct, BSEACASE) was offered twice a day. *Artemia* was

collected, rinsed and resuspended in RAS water. The volume and the concentration of

Artemia (volume collected from hatching cone/volume of RAS water used for

resuspension after rinsing) were adjusted to prevent the accumulation of uneaten food

at the bottom of the tanks and recorded daily. Each tank in the mFTS received the same

amount of food, and the order of feeding was alternated daily to prevent any feeding

bias that could affect growth. Uneaten food and debris accumulated at the bottom of the

tank were removed as required using a 10 ml sterile serological pipette during the water

exchange.

Repeated cross-sectional study of in vivo TBT exposure

Four stocks of 1 mg/ml, 0.1 mg/ml, 0.01 mg/ml, and 0.001 mg/ml of TBT (Sigma,

T50202) in DMSO (Fisher, D136-L) were prepared. All stocks were maintained in the

dark at 4° C until use. Small volumes of these stocks were aliquoted in sterile

borosilicate glass vials to avoid repeated freeze-thaw cycles. DMSO stocks were

aliquoted in the same manner.

Tanks in the mFTS were initially dosed with 75 µl of the corresponding stocks of

TBT and DMSO stocks to achieve final concentrations of TBT in water of 10 µg/l, 1 µg/l,

0.1 µg/l, and 0.01 µg/l, respectively; as well as DMSO vehicle control. By aliquoting the

same volume of stocks, we also maintained an equal volume of DMSO in all treatments.

Stocks were added before refilling tanks after the water exchange to facilitate mixing.

We assumed that TBT was not degraded over a 24-hour period, so we added a volume

of stock daily to adjust the concentration of TBT in the tanks for the total loss of

chemical during the daily water exchange.

Each TBT concentration and the control had four tank replicates and 30 fish per

tank (Fig. 1A). The TBT concentration of 10 μg/l resulted in a mortality rates between

92.31 and 100% after one day of exposure so this treatment was excluded from the

study and the schematic in Fig. 1A. The tank replicates for each treatment were

randomly distributed in our zebrafish facility racks to avoid feeding and water exchange

biases (not represented in Fig. 1A). Four time points were sampled during this repeated

cross-sectional study: one before starting the exposure to TBT (0 days post-exposure,

dpE); as well as 7, 14, and 21 dpE (Fig. 1A). The fish were not offered food on the day

of sampling and had received their last meal and chemical treatments 14 hours before

sampling. 200 ml of water from each tank were collected in sterile conical vials: 150 ml

were used for the aquatic microbial community analysis (see below), and the remainder

50 ml we used for measuring the water quality parameters (pH, nitrate, nitrite, and toxic

ammonia) and for counting colony forming units (CFU). Five individual fish per tank (20

fish per treatment) were sampled at each time point indicated in Fig. 1A. Body size (SL,

HAA, and body area), adipose tissue accumulation, and composition of the intestinal

microbial communities from each fish were also analyzed (see below, Fig. 1A).

Zebrafish body size and adipose tissue analysis

Nile Red (Sigma, N1142) staining was performed as described in (33, 70, 78)

except that the Nile red solution was prepared with filtered and autoclaved RAS water

(FASW). Zebrafish were euthanized in 1.34g/L buffered Tricaine and laid on 1.5%

methylcellulose in FASW on the right flank for imaging. We used a Leica M205

fluorescence stereomicroscope with a GFP bandpass filter to image Nile Red

fluorescence in adipose tissues and corresponding brightfield images of each animal.

All images per time point were acquired using the same exposure settings. Image

analysis was conducted in FIJI/ImageJ 1.5p (127). SL, HAA, and total body area were

measured to evaluate body size as described in (70, 77, 78).

Intestinal and aquatic sample collection

After imaging, euthanized zebrafish were transferred to a Petri dish coated with

sterile 3% agarose, onto which fish were immobilized by the tail using an insect pin

dipped in 70% ethanol. After decapitation with an autoclaved razorblade, the whole

gastrointestinal (GI) tract was removed using forceps. Individual whole intestines were

separated from the other organs of the GI and transferred to sterile screw cap tube

containing 100 µl of sterile 0.1-mm-diameter zirconia beads (BioSpec Products,

NC0268065) and 400 µl of sterile lysis buffer (20 mM Tris-HCl pH 8.0, 2 mM EDTA pH

8.0, 1% Triton X-100; modified from (72)). Tank water from the upper water column was

filtered as described in (72). Filters were then extracted using flame-sterilized forceps

and cut in half using a sterile scalpel. One half of the paper disc was rolled and inserted

into a sterile screw cap tube containing 200 µl of sterile 0.1-mm-diameter zirconia

beads, to which 200 µl of sterile lysis were immediately added. Intestinal and water

samples were flash frozen in a dry ice-ethanol bath and stored at −80°C for subsequent

16S rRNA gene sequencing.

Supercolonization with zebrafish intestinal bacterial strains

624   In the supercolonization experiments described here, we used four bacterial

strains isolated from the zebrafish intestine: *Chitinibacter* ZOR0017, *Chitinibacter*

ZOR0013, *Plesiomonas* ZOR0011; and *Vibrio* ZWU0020 (64, 80). Strains ZOR0017,

ZOR0013, and ZOR0011 were generously provided by Karen Guillemin (University of

Oregon). Single colonies grown on tryptic soy agar (TSA) plates were picked and grown

overnight in tryptic soy broth (TSB). 750 µl of the overnight cultures were collected to

extract DNA and confirm the identity of the strain by sequencing the 16S rRNA gene

using the universal 8F and 1492R primers. An aliquot of this overnight verified culture

was diluted and plated on TSA for growth of well-spread single colonies that were used

to inoculate fresh daily overnight cultures used for preparing the bacterial suspension

administered daily during the supercolonization experiments. The daily overnight

cultures of 10 ml of tryptic soy broth (TSB) were incubated aerobically at 30 $^{\circ}$C for 14 to

16 h in rotary mixer at 100 rpm. 300 µl of the daily overnight culture were collected for

16S rRNA sequencing-based verification of the bacterial strains.

638   A starter culture of 10 ml of fresh TSB was inoculated with the overnight culture

at a starting OD_{600} of 0.1 and incubated for 6 to 8 h at 30°C in a roller drum at 100 rpm.

The starter culture was used to inoculate 200 ml of fresh sterile TSB in an culture flaks

(i.e. supercolonization culture). Because of differences in the growth rate of the different

strains, and to obtain supercolonization cultures of approximately the same density at

the same time, the supercolonization culture for *Chitinibacter* ZOR0017 and

Chitinibacter ZOR0013 was inoculated at a starting OD_{600} of 0.05, whereas

Plesiomonas ZOR0011 and *Vibrio* ZWU0020 were inoculated at a starting OD_{600} of

0.025. The supercolonization culture was incubated at 30°C in a shaker at 100 rpm for

14 hours.

To prepare the bacterial suspension, we calculated the volume of the

supercolonization culture required to prepare 200 ml of bacterial suspension in fresh

TSB at an OD_{600} of 1. The diluted bacterial suspension was transferred to an autoclaved

Nalgene centrifuge bottle (B1033, Sigma), and spun in a Sorvall RC5B centrifuge at

5000rpm at a set temperature range of $10 - 25^{o}C$. The pellet was washed twice with

100 ml of autoclaved RAS water, and resuspended in 60 ml of autoclaved RAS water

(i.e., bacterial suspension for daily supercolonization). 750 µl of each bacterial

suspension were sampled daily to count CFU on TSA and 16S rRNA sequencing-based

verification of the bacteria strains. The CFU/ml in the tanks after supercolonization

(estimated from the CFU/ml in the bacterial suspension) was measured daily. The

average CFU/ml was not significantly different amongst the bacterial strains over the 14

days of supercolonization (One-way ANOVA and Tukey's multiple comparisons test; $P =$

0.7770). The bacterial suspension was divided into 20 ml aliquots used to supercolonize

each tank. Before adding the bacterial suspension, the tanks were drained to the 2.5 L

volume. To promote supercolonization of the intestine with this inoculum, zebrafish were

offered *Artemia* after adding the bacterial suspension. Zebrafish, bacteria, and *Artemia*

were incubated for 2.5 h, before continuing with the water exchange as described

above. Control tanks received a mock inoculation of 20 ml of autoclaved RAS.

Each bacterial treatment and the control had three tank replicates (15

experimental tanks total with ~30 fish per tank) randomized in order. We performed

daily supercolonization for 14 days, which was a period equivalent to the earliest time

point at which we observed body size and adiposity phenotypes during the *in vivo*

chemical exposure (Fig. 2). Five fish per tank were sampled at (0 dpSC) for intestinal

microbiota and body size analysis. Fifteen fish were sampled at the 14 dpSC (as there

was only one time point in this experiment) for body size and adipose tissue analysis.

Five of these fifteen fish were also sampled for intestinal microbiota analysis (Fig. 1B).

Tank water was collected from each tank for aquatic microbiota analysis at 0 and 14

dpSC as previously described for aquatic microbiota and water quality analysis.

Gnotobiotic zebrafish husbandry and obtention of experimental cohorts

The maintenance of the flasks containing GF zebrafish, diet administered and

monoassociation procedures were adapted from (108, 128) to maintain GF animals up

to 35 dpf. This substantial improvement of gnotobiotic husbandry was necessary for our

experiment because adipose tissue starts to develop when zebrafish are SL of ~4.2 mm

which in conventional husbandry is detectable around 12 dpf (69, 70). Moreover, we

wanted to mimic the phenotypical *in vivo* chemical exposure to 1 µg/l TBT (both body

size and adiposity at 21 dpE; Fig.2) and supercolonization with *Plesiomonas* ZOR0011

(at 14 dpSC; Fig.4) under conventional husbandry conditions.

Four conventionally reared adult fish (1:1 male to female ratio) from two different

families were set in mating boxes for group spawning. Zebrafish embryos were obtained

by natural mating. The collection of the embryos and the derivation procedure to obtain

GF zebrafish was conducted as described in (108). All procedures after GF derivation

were conducted using sterile technique, sterile materials wiped with ethanol and UV-

irradiated, inside a laminar flow cabinet. Sterility tests throughout this experiment were

conducted as described (108). Zebrafish were maintained in cell culture flasks with

sterile gnotobiotic zebrafish medium (GZM) in a 28.5°C incubator with a 14:10 hour

light: dark cycle. Flasks were filled with 50 ml of GZM from 0 to 21 dpf, and this volume

was increased to 100 ml GZM from 23 to 35 dpf to accommodate the growth of the

animals. 60% volume GZM exchange was conducted daily to maintain water quality

levels. pH, nitrate, nitrite, and toxic ammonia were measured in all flasks as previously

described. The pH in the flasks ranged between 6.6 and 7, nitrite between 0 and 0.15

ppm, nitrate between 0 and 10 ppm, and toxic ammonia between 0 and 0.0042 mg/l.

The pH in the flasks was lower than in the conditioned water in our RAS facility and the

in vivo chemical exposure, but comparable to the pH during the supercolonization

experiments. The nitrite in the flasks was higher than in the conditioned water in our

RAS zebrafish facility and the supercolonization experiments, but comparable to the *in*

vivo chemical exposure. The nitrate in the flasks was comparable to the conditioned

water in our RAS zebrafish facility, the *in vivo* chemical exposure and the

supercolonization experiments. The toxic ammonia in the flasks was higher than in our

RAS zebrafish facility, but comparable to the *in vivo* chemical exposure and the

supercolonization experiments. Overall, water quality parameters in the flasks were

within the optimal ranges for zebrafish (126).

Flasks of GF zebrafish were raised for 13 days post-fertilization before randomly

assigning flask to the chemical and microbial treatments (see below). These flasks also

underwent a thorough fish health check and verification of GF status. On 14 dpf, one

fish per flask was sampled to measure the SL. The average SL of the cohort was 4.21 ±

0.22 mm (n=29), similar to the SL expected at this age under conventional husbandry.

Exposure of GF zebrafish to TBT and their monoassociation with *Plesiomonas*

ZOR0011 or *Chitinibacter* ZOR0013 were initiated on 15 dpf. Flasks of GF zebrafish (5

flasks per treatment except for *Chitinibacter* ZOR0013 which had only four, 19 flasks

total) were randomly assigned to chemical and bacterial treatments. Two time points

were sampled: 7 and 21 days post-exposure (dpE) or monoassociation (days post-

colonization, dpC). Flasks were sampled terminally at both time points: two flasks per

treatment at 7 dpE, and 3 samples per treatment at 21 dpE (except for the *Chitinibacter*

ZOR0013 treatment from which only two flasks) were sampled at each time point. The

fish were not offered food on the day of sampling and had received their last meal and

treatments 14 hours before sampling. Body size and adipose tissue area from each fish

in the flask were analyzed as previously described. There were no significant

differences in the rearing density of the flasks assigned to the chemical/bacterial

treatments that could result in differences in body size (Kruskal-Wallis, Dunn's multiple

comparison test of flasks from each chemical/bacterial treatment against the germ-free

DMSO control; all $P \geq 0.8869$ at 7 dpE/dpC; and $P \geq 0.9474$ at 21 dpE/dpC). Similarly,

there were no differences in mortality in flasks sampled at 7 or 14 dpE/dpC (Kruskal-

Wallis, Dunn's multiple comparison test of flasks from each chemical/bacterial treatment

against the germ-free DMSO; all $P \geq 0.1986$ at 7 dpE/dpC; and $P \geq 0.5213$ at 21

dpE/dpC).

Culture and provision of germ-free Artemia-Tetrahymena diet

One of the main challenges of long-term gnotobiotic husbandry is the transition in

zebrafish larvae from reliance on yolk to free-feeding before they are capable of hunting

live *Artemia*. Also, we wanted to reproduce as closely as possible the diet zebrafish

received in the mFTS during the *in vivo* chemical exposure, and the supercolonization

experiments. However, previous pilot experiments showed that an *Artemia*-only diet

was not compatible with survival past 14 dpf (data not shown). Thus, we designed a GF

live diet of *Tetrahymena thermophila* strain CU428.2 (TSC_SD00178, Tetrahymena

Stock Center) and San Francisco strain *Artemia* (Brine Shrimp Direct). GF *Artemia* was

offered daily starting 5 dpf, and supplemented with GF *T. thermophila* on 5, 8, and 11

dpf. GF *Artemia* and *T. thermophila* cultures were prepared as described below.

T. thermophila stocks were maintained in 5 ml of Modified Neff's Media (MNM; 3

mg $FeCl_3.6H_20$, 10 g proteose peptone, 2.5 g yeast extract, and 5 g glucose) with 5 µl

Penicillin-Streptomycin Stabilized solution (Sigma-Aldrich, P4458) and 24 µl of

Amphotericin B solution (Sigma-Aldrich, A2942) at room temperature. These stocks

were renewed every two weeks by transferring 100 µl of stock to fresh media. 200 µl of

MNM stock were used to inoculate 14.8 ml of Proteose Peptone Yeast Agar (PPYE; 2,5

g proteose peptone; 2.5 g yeast extract) media with 10 µl of Penicillin-Streptomycin

Stabilized solution (Sigma-Aldrich, P4458). PPYE cultures were maintained at 28.5° C

for one week before its use for inoculation of milk yeast extract media (MYE; 10g/l yeast

extract, 10 g/l non-fat powdered milk). MYE is a rich media that is used for growing *T.*

thermophila for feeding zebrafish (129). 4 ml of a one-week old PPYE culture were used

to inoculate 40 ml of sterile MYE and was incubated at 28.5° C overnight in a 50 ml

conical vial. For food preparation, the conical vial was centrifuged at 5000 rcf for 3

minutes, 25 ml of the supernatant were decanted, and the conical vial was refilled to the

50 ml mark with sterile GZM to rinse the pellet. The pellet was rinsed two more times

with sterile GZM by decanting 35 ml of the supernatant, before its final resuspension in

40 ml of sterile GZM. This preparation had an average of ~30 000 cells/ml. 4 ml of the

suspension of *T. thermophila* were used to feed each fish flask. Sterility tests were

conducted on the *T. thermophila* MNM and PPYE cultures, as well as on prepared food

suspensions as described in (108).

To prepare GF Artemia, ~250 µl of Artemia cysts were rehydrated in 10 ml of

sterile saltwater (12.5g/l NaCl, 0.83g/l NaHCO3) for at least 2 hours with gentle shaking

in a nutator. The cysts were left to settle, and the saltwater was exchanged for 10 ml of

household bleach (8.25% sodium hypochlorite) for decapsulation. The Artemia-bleach

suspension was mixed on a nutator for 3 minutes, and then left to settle at the bottom of

the conical tube for 5 more minutes. After decapsulation, the bleach and floating cysts

were removed. The cysts were rinsed by adding saltwater, mixed by gentle rolling for

1.5 minutes, before letting the cysts settle for 30 seconds. The procedure was repeated

4 times, before resuspending the cysts in 10 ml of sterile saltwater, and incubation for at

least 24 h at room temperature with gentle shaking under a warm light for hatching. For

food preparation, the actively swimming *Artemia* were collected using a sterile

serological pipette, collected on a 100 µm cell strainer (431752, Corning) and rinsed

with 10 ml of sterile GZM. *Artemia* was transferred to a conical vial and resuspended in

GZM. The volume and the concentration (i.e. volume of collected from hatching

vial/volume of GZM for resuspension) of *Artemia* was adjusted to prevent the

accumulation of food debris in the flasks. Volume and concentration were recorded

daily. Each flask of fish received the same amount of food, and the order of feeding was

alternated daily to prevent to prevent any feeding bias that could affect growth. Uneaten

food and debris were removed daily during the GZM exchange.

Germ-free zebrafish exposure to TBT

Stocks of 1 mg/ml of TBT (Sigma, T50202) in DMSO (Fisher, D136-L) and of

DMSO vehicle were prepared and aliquoted using sterile technique. Although the

presence of bacteria in these chemicals was unlikely, the sterility of the stocks was

tested by plating 100 µl of the stock solutions on TSA and incubated for 1 week at

28.5°C under aerobic conditions before starting the experiment. The plates were

inspected daily, and no microbial growth was observed.

Flasks containing GF zebrafish were initially dosed with 5 µl of the TBT stock and

DMSO to achieve a final concentration of 1 µg/l TBT as well as an equal total volume of

DMSO vehicle in the control flasks. Stocks were added before fresh GZM during the

media exchange to facilitate mixing. We assumed that TBT was not degraded over a

24-hour period, so we added a volume of stock daily to adjust the concentration of TBT

in the flask for the total loss of chemical during the daily GZM exchange.

Monoassociation with Plesiomonas ZOR0011 and Chitinibacter ZOR0013

Monoassociation is the colonization of GF zebrafish with a single microbial strain.

A starter culture of each bacterial strain was prepared by inoculating 10 ml of sterile

TSB with a single colony, followed by overnight aerobic incubation in a roller drum at

100 rpm at 30° C. 300 µl of the starter culture were collected to confirm the identity of

the strain by sequencing as described for the supercolonization experiments. The

starter culture was used to inoculate 40 ml of sterile TSB in a sterile 50 ml conical via

(i.e. monoassociation culture). To obtain monoassociation cultures of approximately the

same density at the same time, *Chitinibacter* ZOR0013 was inoculated at a starting

OD_{600} of 0.05 and *Plesiomonas* ZOR0011 at 0.025 as *Plesiomonas* ZOR0011 grew

faster in TSB than *Chitinibacter* ZOR00R13. The monoassociation culture was

incubated at 30° C in a shaker at 100 rpm for 14 hours. To prepare the bacterial

suspension for the monoassociation, we calculated the volume of the culture needed to

prepare 40 ml of a bacterial suspension at an OD_{600} of 0.25. We chose the OD_{600} and

dilutions to ensure that the zebrafish fish received amounts of bacteria in the

monoassociation comparable to those in the supercolonization experiments. The diluted

bacterial suspension was spun at 5000 rpm for 10 minutes. The supernatant was

decanted, and the pellet was washed twice with 40 ml of sterile GZM. The washed

bacterial pellet was finally resuspended in 40 ml of autoclaved GZM, and this

suspension was used for the monoassociation. 750 µl of each bacterial suspension

were sampled to count CFU on TSA plates and to confirm the identity of the strain by

16S rRNA gene sequencing as described above. The CFU/ml in the flask after

inoculation were of 1.01E+08 CFU/ml for *Chitinibacter* ZOR0013 and 1.20E+07 CFU/ml

for *Plesiomonas* ZOR0011 (estimated from the CFU/ml of the bacterial suspension).

Nevertheless, the average CFU/ml counts from GZM from individual flasks sampled at 7

dpC and 21 dpC were not significantly different (t-test, $P = 0.7541$ and $P = 0652$

respectively) between *Chitinibacter* ZOR0013 and *Plesiomonas* ZOR0011, suggesting

that this initial difference was not maintained and likely did not affect the results of our

experiments. In addition, whereas the estimated CFU/ml in the fish flasks after

monoassociation was lower than the average estimated CFU/ml in the tanks after the

supercolonization experiments, these values were within the minimum to maximum

CFU/ml range in tanks after supercolonization measured during the 14 days of

supercolonization (data not shown).

GF flasks assigned to the bacterial treatments received 6 ml *Chitinibacter*

ZOR0013 or *Plesiomonas* ZOR0011 suspension. In addition, the monoassociated flasks

were initially dosed with 5 µl of DMSO since the GF DMSO control was shared with the

chemical exposure to TBT in the GF background. DMSO was added before fresh GZM

during the media exchange to facilitate mixing, and a volume was added daily to adjust

for the loss chemical during the daily GZM exchange. Bacteria were added only once

during the experiment at 15 dpf.

In vitro exposure of zebrafish intestinal microbial communities to TBT

Four stocks of 1000 µg/ml, 10 µg/ml, 1 µg/ml, 0.1 µg/ml, and 0.01 µg/ml of TBT

(Sigma, T50202) in DMSO (Fisher, D136-L) were prepared and aliquoted using sterile

technique. All stocks were maintained in the dark at 4°C until use. Borosilicate tubes

containing 15 ml of sterile tryptic soy broth (TSB) were used for the aerobic *in vitro*

exposure; and borosilicate tubes containing 14.25 ml of sterile TSB and 0.75 µl of 10 g/l

filter sterilized L-cysteine were used for the anaerobic *in vitro* chemical exposure. 15 µl

of the stocks were added to the growth media to achieve final TBT concentrations of 0

µg/l (DMSO vehicle control), 0.01 µg/l, 0.1 µg/l, 1 µg/l, and 10 µg/l, and 1000 µg/l TBT.

Each TBT concentration was prepared in triplicate.

An experimental cohort of zebrafish was obtained and conventionally reared. On

day 25 post-fertilization, zebrafish were size matched between 7 mm and 9 mm. Two

pools of 30 size-matched individuals were transferred to two clean and autoclaved fish

tanks and maintained in the RAS for 4 weeks before sampling to match the SL at 21

dpE *in vivo* chemical exposure to TBT. Two fish per tank were collected as donors (n =

4; average SL = 16.75 ± 1.5 mm). To reproduce the sampling of whole intestines during

the *in vivo* chemical exposure, zebrafish were incubated in 1 mM epinephrine in FASW

before euthanasia with 1.34 g/L filter sterilized tricaine. Whole gastrointestinal (GI) tracts

were dissected as previously described, and cut open using sterile technique before

transfer to a sterile screw cap tube containing 200 µl of sterile 1-mm-diameter zirconia

beads (NC9847287) and 500 µl of FASW. Four GI tracts were pooled in a single

beaded screw cap tube. Intestinal samples were vortexed for one minute twice to obtain

an intestinal slurry used to inoculate TSB media with TBT and DMSO prepared as

described above (10 µl of slurry in 5 ml of media). The remainder of the slurry

constituted the input intestinal microbial community and was transferred to a sterile

Eppendorf tube. The beads were rinsed with 500 µl of FASW, and after vortexing, this

rinsing volume was pooled with the slurry. The collected material was centrifuged,

before centrifugation at 7500 rcf for 15 minutes. After removing the supernatant, the

pellet was flash frozen in a dry ice-ethanol bath and stored at −80°C until DNA isolation

for 16S rRNA gene sequencing.

TSB cultures dosed with TBT or DMSO were incubated aerobically and

anaerobically for 24 hours without agitation at room temperature. Anaerobic incubation

was performed in a Coy anaerobic chamber. 750 µl of each culture tube was collected

in a sterile 1.5 ml Eppendorf tube and spun at 7500 rcf. The pellets were flash frozen in

a dry ice-ethanol bath and stored at −80°C until DNA isolation for 16S rRNA gene

sequencing for microbial community analysis.

DNA isolation

The Duke Microbiome Shared Resource extracted bacterial DNA from all

samples using Qiagen MO BIO PowerSoil DNA Kit that allows for the isolation of

samples in a 96 well plate format using the Retsch MM400 plate shaker. Quantity of

samples were assessed using the PerkinElmer Victor plate reader with the

ThermoFisher Scientific Qubit dsDNA HS assay kit.

Sequencing of 16S rRNA gene amplicons

Bacterial community composition in isolated DNA samples was characterized by

amplification of the V4 variable region of the 16S rRNA gene by polymerase chain

reaction using the forward primer 515 and reverse primer 806 16S rRNA following the

Earth Microbiome Project protocol (http://www.earthmicrobiome.org/) by the Duke

Microbiome Shared Resource. These primers (515F and 806R) carry unique barcodes

allowing for multiplexing and co-sequencing of hundreds of samples at a time.

Equimolar 16S rRNA gene PCR products were quantified and pooled for sequencing.

Sequencing was performed by the Duke Microbiome Center using an Illumina MiSeq

instrument configured for 150 bp paired-end sequencing runs.

Bioinformatic analysis

Data analysis conducted in QIIME v. 1.9.1 (131). Paired reads were joined with

join_paired_ends.py, using standard parameters, and joined reads were subsequently

demultiplexed with split_libraries_fastq.py, with the phred quality threshold set to Q30.

Demultiplexed amplicons were filtered for chimeric sequences with

identify_chimeric_seqs.py, using the usearch61 (132) detection method, and

substituting usearch v. 6.1 with vsearch v. 2.7.1 (133). Only amplicons predicted as

chimeric by both denovo and reference-based detection were removed. After

completion of base quality and chimera filtering, totals of 21,069,394 (*in vivo* TBT

exposure study), 1,283,416 (supercolonization study) and 5,801,110 (*in vitro* TBT

exposure study) amplicons were used in downstream analyses. Amplicons were

clustered into operational taxonomic units (OTUs), using QIIME's closed reference OTU

picking strategy, implemented in pick_closed_reference_otus.py. The script was

configured to cluster amplicons with uclust (132) and assign taxonomy against the 97%

clustered 16S reference sequence set of SILVA, v. 1.23 (134). A subsequent basic

statistical diversity analysis was performed, using QIIME's core diversity analysis

workflow script (core_diversity_analyses.py), calculating alpha- and beta-diversity, and

relative taxa abundances in sample groups. Relative taxa abundances were further

analyzed with LEfSe (Linear discriminant analysis effect size, (79) to identify differential

biomarkers in sample groups.

Data availability

16S rRNA sequence data are available at the NCBI Short Read Archive under

accessions PRJNA692784 (*in vivo* TBT exposure study), PRJNA693006

(supercolonization study), PRJNA693015 (*in vitro* TBT exposure study).

ACKNOWLEDGEMENTS

This work was supported by grants to J.F.R. from the National Institutes of Health

(R21-ES023369) and the Gordon and Betty Moore Foundation. The authors are grateful

to Karen Guillemin and Brendan Bohannan (University of Oregon) for providing bacterial

strains; José M. Abelleira-Pereira (Universidad de Cádiz) for assistance in design of the

mFTS; Philip Zandona for technical assistance in construction of the mFTS; Jim Burris

and Patrick Williams (Duke Zebrafish Core Facility) for accommodation and support of

our zebrafish experiments; Holly Dressman and Zhengzheng Wei for their assistance in

the optimization of extraction protocols and other helpful conversations; Colin Lickwar

and Jia Wen for helpful feedback on this manuscript; and the competent staff at Public

Hardware Inc. (Durham, NC).

REFERENCES

1.   Hill JO, Wyatt HR, Peters JC. 2012. Energy balance and obesity. Circulation 126:126-32.
2.   Rutkowski JM, Stern JH, Scherer PE. 2015. The cell biology of fat expansion. J Cell Biol 208:501-12.
3.   World Health Organization. 2020. Obesity and overweight https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight.
4.   Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, Mullany EC, Biryukov S, Abbafati C, Abera SF, Abraham JP, Abu-Rmeileh NM, Achoki T, AlBuhairan FS, Alemu ZA, Alfonso R, Ali MK, Ali R, Guzman NA, Ammar W, Anwari P, Banerjee A, Barquera S, Basu S, Bennett DA, Bhutta Z, Blore J, Cabral N, Nonato IC, Chang JC, Chowdhury R, Courville KJ, Criqui MH, Cundiff DK, Dabhadkar KC, Dandona L, Davis A, Dayama A, Dharmaratne SD, Ding EL, Durrani AM, Esteghamati A, Farzadfar F, Fay DF, Feigin VL, Flaxman A, Forouzanfar MH, Goto A, Green MA, Gupta R, et al. 2014. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet 384:766-81.
5.   Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. 2009. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. BMC Public Health 9:88.
6.   Nyberg ST, Batty GD, Pentti J, Virtanen M, Alfredsson L, Fransson EI, Goldberg M, Heikkila K, Jokela M, Knutsson A, Koskenvuo M, Lallukka T, Leineweber C, Lindbohm JV, Madsen IEH, Magnusson Hanson LL, Nordin M, Oksanen T, Pietilainen O, Rahkonen O, Rugulies R, Shipley MJ, Stenholm S, Suominen S, Theorell T, Vahtera J, Westerholm PJM, Westerlund H, Zins M, Hamer M, Singh-Manoux A, Bell JA, Ferrie JE, Kivimaki M. 2018. Obesity and loss of disease-free years owing to major non-communicable diseases: a multicohort study. Lancet Public Health 3:e490-e497.
7.   Schwartz MW, Seeley RJ, Zeltser LM, Drewnowski A, Ravussin E, Redman LM, Leibel RL. 2017. Obesity Pathogenesis: An Endocrine Society Scientific Statement. Endocr Rev 38:267-296.
8.   Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, Semenkovich CF, Gordon JI. 2004. The gut microbiota as an environmental factor that regulates fat storage. Proc Natl Acad Sci U S A 101:15718-23.
9.   Schwarzer M, Makki K, Storelli G, Machuca-Gayet I, Srutkova D, Hermanova P, Martino ME, Balmand S, Hudcovic T, Heddi A, Rieusset J, Kozakova H, Vidal H, Leulier F. 2016. Lactobacillus plantarum strain maintains growth of infant mice during chronic undernutrition. Science 351:854-7.

979   10.   Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, Griffin NW,
980   Lombard V, Henrissat B, Bain JR, Muehlbauer MJ, Ilkayeva O, Semenkovich CF,
981   Funai K, Hayashi DK, Lyle BJ, Martini MC, Ursell LK, Clemente JC, Van Treuren
982   W, Walters WA, Knight R, Newgard CB, Heath AC, Gordon JI. 2013. Gut
983   microbiota from twins discordant for obesity modulate metabolism in mice.
984   Science 341:1241214.

985   11.   Smith MI, Yatsunenko T, Manary MJ, Trehan I, Mkakosya R, Cheng J, Kau AL,
986   Rich SS, Concannon P, Mychaleckyj JC, Liu J, Houpt E, Li JV, Holmes E,
987   Nicholson J, Knights D, Ursell LK, Knight R, Gordon JI. 2013. Gut microbiomes
988   of Malawian twin pairs discordant for kwashiorkor. Science 339:548-54.

989   12.   Comittee on Advancing Understanding of the Implication of Environmental-
990   Chemical Interactions with the Human Microbiome, National Academies of
991   Sciences Engineering and Medicine. 2017. Environmental Chemicals, the
992   Human Microbiome, and Health Risk: A Research Strategy doi:10.17226/24960,
993   Washington (DC).

994   13.   Joly C, Gay-Queheillard J, Leke A, Chardon K, Delanaud S, Bach V, Khorsi-
995   Cauet H. 2013. Impact of chronic exposure to low doses of chlorpyrifos on the
996   intestinal microbiota in the Simulator of the Human Intestinal Microbial
997   Ecosystem (SHIME) and in the rat. Environ Sci Pollut Res Int 20:2726-34.

998   14.   Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. 2013.
999   Exercise attenuates PCB-induced changes in the mouse gut microbiome.
Environ Health Perspect 121:725-30.

15.   Lu K, Abo RP, Schlieper KA, Graffam ME, Levine S, Wishnok JS, Swenberg JA,
Tannenbaum SR, Fox JG. 2014. Arsenic exposure perturbs the gut microbiome
and its metabolic profile in mice: an integrated metagenomics and metabolomics
analysis. Environ Health Perspect 122:284-91.

16.   Suez J, Korem T, Zeevi D, Zilberman-Schapira G, Thaiss CA, Maza O, Israeli D,
Zmora N, Gilad S, Weinberger A, Kuperman Y, Harmelin A, Kolodkin-Gal I,
Shapiro H, Halpern Z, Segal E, Elinav E. 2014. Artificial sweeteners induce
glucose intolerance by altering the gut microbiota. Nature 514:181-6.

17.   Lindenbaum J, Rund DG, Butler VP, Jr., Tse-Eng D, Saha JR. 1981. Inactivation
of digoxin by the gut flora: reversal by antibiotic therapy. N Engl J Med 305:789-
94.

18.   Yan A, Culp E, Perry J, Lau JT, MacNeil LT, Surette MG, Wright GD. 2018.
Transformation of the Anticancer Drug Doxorubicin in the Human Gut
Microbiome. ACS Infect Dis 4:68-76.

19.   Maini Rekdal V, Bess EN, Bisanz JE, Turnbaugh PJ, Balskus EP. 2019.
Discovery and inhibition of an interspecies gut bacterial pathway for Levodopa
metabolism. Science 364.

20.   Wallace BD, Wang H, Lane KT, Scott JE, Orans J, Koo JS, Venkatesh M, Jobin
C, Yeh LA, Mani S, Redinbo MR. 2010. Alleviating cancer drug toxicity by
inhibiting a bacterial enzyme. Science 330:831-5.

21.   Egusquiza RJ, Blumberg B. 2020. Environmental Obesogens and Their Impact
on Susceptibility to Obesity: New Mechanisms and Chemicals. Endocrinology
161.

22. Lind L, Lind PM, Lejonklou MH, Dunder L, Bergman A, Guerrero-Bosagna C,
Lampa E, Lee HK, Legler J, Nadal A, Pak YK, Phipps RP, Vandenberg LN, Zalko
D, Agerstrand M, Oberg M, Blumberg B, Heindel JJ, Birnbaum LS. 2016.
Uppsala Consensus Statement on Environmental Contaminants and the Global
Obesity Epidemic. Environ Health Perspect 124:A81-3.
23. Copisarow M. 1945. Marine Fouling and Its Prevention. Science 101:406-7.
24. Evans SM, Birchenough AC, Brancato MS. 2000. The TBT Ban: Out of the
Frying Pan into the Fire? Marine Pollution Bulletin 40:204-211.
25. Gibbs PE, Pascoe PL, Burt GR. 1988. Sex Change in the Female Dog-Whelk,
Nucella Lapillus , Induced by Tributyltin from Antifouling Paints. Journal of the
Marine Biological Association of the United Kingdom 68:715-731.
26. Smith BS. 1981. Tributyltin compounds induce male characteristics on female
mud snails Nassarius obsoletus = Ilyanassa obsoleta. J Appl Toxicol 1:141-4.
27. Grun F, Watanabe H, Zamanian Z, Maeda L, Arima K, Cubacha R, Gardiner DM,
Kanno J, Iguchi T, Blumberg B. 2006. Endocrine-disrupting organotin compounds
are potent inducers of adipogenesis in vertebrates. Mol Endocrinol 20:2141-55.
28. Yao W, Wei X, Guo H, Cheng D, Li H, Sun L, Wang S, Guo D, Yang Y, Si J.
2020. Tributyltin reduces bone mineral density by reprograming bone marrow
mesenchymal stem cells in rat. Environ Toxicol Pharmacol 73:103271.
29. Bertuloso BD, Podratz PL, Merlo E, de Araujo JF, Lima LC, de Miguel EC, de
Souza LN, Gava AL, de Oliveira M, Miranda-Alves L, Carneiro MT, Nogueira CR,
Graceli JB. 2015. Tributyltin chloride leads to adiposity and impairs metabolic
functions in the rat liver and pancreas. Toxicol Lett 235:45-59.
30. Yan H, Guo H, Cheng D, Kou R, Zhang C, Si J. 2018. Tributyltin reduces the
levels of serum adiponectin and activity of AKT and induces metabolic syndrome
in male mice. Environ Toxicol 33:752-758.
31. Chamorro-Garcia R, Sahu M, Abbey RJ, Laude J, Pham N, Blumberg B. 2013.
Transgenerational inheritance of increased fat depot size, stem cell
reprogramming, and hepatic steatosis elicited by prenatal exposure to the
obesogen tributyltin in mice. Environ Health Perspect 121:359-66.
32. Zhang J, Zhang C, Ma D, Liu M, Huang S. 2017. Lipid accumulation, oxidative
stress and immune-related molecules affected by tributyltin exposure in muscle
tissues of rare minnow (Gobiocypris rarus). Fish Shellfish Immunol 71:10-18.
33. Tingaud-Sequeira A, Ouadah N, Babin PJ. 2011. Zebrafish obesogenic test: a
tool for screening molecules that target adiposity. J Lipid Res 52:1765-72.
34. Ouadah-Boussouf N, Babin PJ. 2016. Pharmacological evaluation of the
mechanisms involved in increased adiposity in zebrafish triggered by the
environmental contaminant tributyltin. Toxicol Appl Pharmacol 294:32-42.
35. Kanayama T, Kobayashi N, Mamiya S, Nakanishi T, Nishikawa J. 2005.
Organotin compounds promote adipocyte differentiation as agonists of the
peroxisome proliferator-activated receptor gamma/retinoid X receptor pathway.
Mol Pharmacol 67:766-74.
36. Regnier SM, El-Hashani E, Kamau W, Zhang X, Massad NL, Sargis RM. 2015.
Tributyltin differentially promotes development of a phenotypically distinct
adipocyte. Obesity (Silver Spring) 23:1864-71.

37.   Kirchner S, Kieu T, Chow C, Casey S, Blumberg B. 2010. Prenatal exposure to
the environmental obesogen tributyltin predisposes multipotent stem cells to
become adipocytes. Mol Endocrinol 24:526-39.
38.   Inadera H, Shimomura A. 2005. Environmental chemical tributyltin augments
adipocyte differentiation. Toxicol Lett 159:226-34.
39.   Carfi M, Croera C, Ferrario D, Campi V, Bowe G, Pieters R, Gribaldo L. 2008.
TBTC induces adipocyte differentiation in human bone marrow long term culture.
Toxicology 249:11-8.
40.   Lutfi E, Riera-Heredia N, Cordoba M, Porte C, Gutierrez J, Capilla E, Navarro I.
2017. Tributyltin and triphenyltin exposure promotes in vitro adipogenic
differentiation but alters the adipocyte phenotype in rainbow trout. Aquat Toxicol
188:148-158.
41.   Gadd GM. 2000. Microbial interactions with tributyltin compounds: detoxification,
accumulation, and environmental fate. Sci Total Environ 258:119-27.
42.   Cruz A, Caetano T, Suzuki S, Mendo S. 2007. Aeromonas veronii, a tributyltin
(TBT)-degrading bacterium isolated from an estuarine environment, Ria de
Aveiro in Portugal. Mar Environ Res 64:639-50.
43.   Cruz A, Oliveira V, Baptista I, Almeida A, Cunha A, Suzuki S, Mendo S. 2012.
Effect of tributyltin (TBT) in the metabolic activity of TBT-resistant and sensitive
estuarine bacteria. Environ Toxicol 27:11-7.
44.   Fukagawa T, Suzuki S, Fukunaga K, Suzuki T, Takama K. 1992. Isolation and
characterization of tributyltin chloride-resistant marine Vibrio. FEMS Microbiol
Lett 72:83-6.
45.   Cruz A, Henriques I, Sousa AC, Baptista I, Almeida A, Takahashi S, Tanabe S,
Correia A, Suzuki S, Anselmo AM, Mendo S. 2014. A microcosm approach to
evaluate the degradation of tributyltin (TBT) by Aeromonas molluscorum Av27 in
estuarine sediments. Environ Res 132:430-7.
46.   Pain A, Cooney JJ. 1998. Characterization of organotin-resistant bacteria from
boston harbor sediments. Arch Environ Contam Toxicol 35:412-6.
47.   Cruz A, Anselmo AM, Suzuki S, Mendo S. 2015. Tributyltin (TBT): A Review on
Microbial Resistance and Degradation. Critical Reviews in Environmental
Science and Technology 45:970-1006.
48.   Wuertz S, Miller CE, Pfister RM, Cooney JJ. 1991. Tributyltin-resistant bacteria
from estuarine and freshwater sediments. Appl Environ Microbiol 57:2783-9.
49.   Dowson PH, Bubb JM, Lester JN. 1996. Persistence and Degradation Pathways
of Tributyltin in Freshwater and Estuarine Sediments. Estuarine, Coastal and
Shelf Science 42:551-562.
50.   Bernat P, Siewiera P, Sobon A, Dlugonski J. 2014. Phospholipids and protein
adaptation of Pseudomonas sp. to the xenoestrogen tributyltin chloride (TBT).
World J Microbiol Biotechnol 30:2343-50.
51.   Jude F, Arpin C, Brachet-Castang C, Capdepuy M, Caumette P, Quentin C.
2004. TbtABM, a multidrug efflux pump associated with tributyltin resistance in
Pseudomonas stutzeri. FEMS Microbiol Lett 232:7-14.
52.   Inoue H, Takimura O, Kawaguchi K, Nitoda T, Fuse H, Murakami K, Yamaoka Y.
2003. Tin-carbon cleavage of organotin compounds by pyoverdine from
Pseudomonas chlororaphis. Appl Environ Microbiol 69:878-83.

53. Cruz A, Micaelo N, Felix V, Song JY, Kitamura S, Suzuki S, Mendo S. 2013.
sugE: A gene involved in tributyltin (TBT) resistance of Aeromonas molluscorum
Av27. J Gen Appl Microbiol 59:39-47.
54. Hernould M, Gagne S, Fournier M, Quentin C, Arpin C. 2008. Role of the
AheABC efflux pump in Aeromonas hydrophila intrinsic multidrug resistance.
Antimicrob Agents Chemother 52:1559-63.
55. Cruz A, Rodrigues R, Pinheiro M, Mendo S. 2015. Transcriptomes analysis of
Aeromonas molluscorum Av27 cells exposed to tributyltin (TBT): Unravelling the
effects from the molecular level to the organism. Mar Environ Res 109:132-9.
56. Khanolkar D, Dubey SK, Naik MM. 2015. Tributyltin chloride (TBTCl)-enhanced
exopolysaccharide and siderophore production in an estuarine Alcaligenes
faecalis strain. Arch Environ Contam Toxicol 68:612-21.
57. Dubey SK, Roy U. 2003. Review: Biodegradation of tributyltins (organotins) by
marine bacteria. Applied Organometallic Chemistry 17:3-8.
58. Mimura H, Sato R, Furuyama Y, Taniike A, Yagi M, Yoshida K, Kitamura A.
2008. Adsorption of tributyltin by tributyltin resistant marine Pseudoalteromonas
sp. cells. Mar Pollut Bull 57:877-82.
59. Yuan G-h, Zhang Z, Gao X-s, Zhu J, Guo W-h, Wang L, Ding P, Jiang P, Li L.
2020. Gut microbiota-mediated tributyltin-induced metabolic disorder in rats. RSC
Advances 10:43619-43628.
60. Zhan J, Ma X, Liu D, Liang Y, Li P, Cui J, Zhou Z, Wang P. 2020. Gut
microbiome alterations induced by tributyltin exposure are associated with
increased body weight, impaired glucose and insulin homeostasis and endocrine
disruption in mice. Environ Pollut 266:115276.
61. Guo H, Yan H, Cheng D, Wei X, Kou R, Si J. 2018. Tributyltin exposure induces
gut microbiome dysbiosis with increased body weight gain and dyslipidemia in
mice. Environ Toxicol Pharmacol 60:202-208.
62. Rawls JF, Samuel BS, Gordon JI. 2004. Gnotobiotic zebrafish reveal
evolutionarily conserved responses to the gut microbiota. Proc Natl Acad Sci U S
A 101:4596-601.
63. Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin
K, Rawls JF. 2011. Evidence for a core gut microbiota in the zebrafish. ISME J
5:1595-608.
64. Stephens WZ, Burns AR, Stagaman K, Wong S, Rawls JF, Guillemin K,
Bohannan BJ. 2016. The composition of the zebrafish intestinal microbial
community varies across development. ISME J 10:644-54.
65. Semova I, Carten JD, Stombaugh J, Mackey LC, Knight R, Farber SA, Rawls JF.
2012. Microbiota regulate intestinal absorption and metabolism of fatty acids in
the zebrafish. Cell Host Microbe 12:277-88.
66. Rolig AS, Parthasarathy R, Burns AR, Bohannan BJ, Guillemin K. 2015.
Individual Members of the Microbiota Disproportionately Modulate Host Innate
Immune Responses. Cell Host Microbe 18:613-20.
67. Bates JM, Mittge E, Kuhlman J, Baden KN, Cheesman SE, Guillemin K. 2006.
Distinct signals from the microbiota promote different aspects of zebrafish gut
differentiation. Dev Biol 297:374-86.

68. Flynn EJ, 3rd, Trent CM, Rawls JF. 2009. Ontogeny and nutritional control of
adipogenesis in zebrafish (Danio rerio). J Lipid Res 50:1641-52.
69. Imrie D, Sadler KC. 2010. White adipose tissue development in zebrafish is
regulated by both developmental time and fish size. Dev Dyn 239:3013-23.
70. Minchin JEN, Rawls JF. 2017. A classification system for zebrafish adipose
tissues. Dis Model Mech 10:797-809.
71. Minchin JE, Dahlman I, Harvey CJ, Mejhert N, Singh MK, Epstein JA, Arner P,
Torres-Vazquez J, Rawls JF. 2015. Plexin D1 determines body fat distribution by
regulating the type V collagen microenvironment in visceral adipose tissue. Proc
Natl Acad Sci U S A 112:4363-8.
72. Wong S, Stephens WZ, Burns AR, Stagaman K, David LA, Bohannan BJ,
Guillemin K, Rawls JF. 2015. Ontogenetic Differences in Dietary Fat Influence
Microbiota Assembly in the Zebrafish Gut. MBio 6:e00687-15.
73. Antizar-Ladislao B. 2008. Environmental levels, toxicity and human exposure to
tributyltin (TBT)-contaminated marine environment. a review.
b_antizar@hotmail.com. Environ Int 34:292-308.
74. Kannan K, Senthilkumar K, Giesy JP. 1999. Occurrence of Butyltin Compounds
in Human Blood. Environmental Science & Technology 33:1776-1779.
75. Schoyen M, Green NW, Hjermann DO, Tveiten L, Beylich B, Oxnevad S, Beyer
J. 2019. Levels and trends of tributyltin (TBT) and imposex in dogwhelk (Nucella
lapillus) along the Norwegian coastline from 1991 to 2017. Mar Environ Res
144:1-8.
76. Castro IB, Iannacone J, Santos S, Fillmann G. 2018. TBT is still a matter of
concern in Peru. Chemosphere 205:253-259.
77. Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. 2009. Normal
table of postembryonic zebrafish development: staging by externally visible
anatomy of the living fish. Dev Dyn 238:2975-3015.
78. Minchin JE, Rawls JF. 2017. In vivo imaging and quantification of regional
adiposity in zebrafish. Methods Cell Biol 138:3-27.
79. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower
C. 2011. Metagenomic biomarker discovery and explanation. Genome Biol
12:R60.
80. Rawls JF, Mahowald MA, Ley RE, Gordon JI. 2006. Reciprocal gut microbiota
transplants from zebrafish and mice to germ-free recipients reveal host habitat
selection. Cell 127:423-33.
81. Zhang J, Zuo Z, Xiong J, Sun P, Chen Y, Wang C. 2013. Tributyltin exposure
causes lipotoxicity responses in the ovaries of rockfish, Sebastiscus marmoratus.
Chemosphere 90:1294-9.
82. Zhang J, Zuo Z, He C, Cai J, Wang Y, Chen Y, Wang C. 2009. Effect of tributyltin
on testicular development in Sebastiscus marmoratus and the mechanism
involved. Environ Toxicol Chem 28:1528-35.
83. Crawford KA, Clark BW, Heiger-Bernays WJ, Karchner SI, Claus Henn BG,
Griffith KN, Howes BL, Schlezinger DR, Hahn ME, Nacci DE, Schlezinger JJ.
2019. Altered lipid homeostasis in a PCB-resistant Atlantic killifish (Fundulus
heteroclitus) population from New Bedford Harbor, MA, U.S.A. Aquat Toxicol
210:30-43.

84.   Meador JP, Sommers FC, Cooper KA, Yanagida G. 2011. Tributyltin and the
1207   obesogen metabolic syndrome in a salmonid. Environ Res 111:50-6.
1208   85.   Qiu X, Iwasaki N, Chen K, Shimasaki Y, Oshima Y. 2019. Tributyltin and
1209   perfluorooctane sulfonate play a synergistic role in promoting excess fat
1210   accumulation in Japanese medaka (Oryzias latipes) via in ovo exposure.
1211   Chemosphere 220:687-695.
1212   86.   Lyssimachou A, Santos JG, Andre A, Soares J, Lima D, Guimaraes L, Almeida
1213   CM, Teixeira C, Castro LF, Santos MM. 2015. The Mammalian "Obesogen"
1214   Tributyltin Targets Hepatic Triglyceride Accumulation and the Transcriptional
1215   Regulation of Lipid Metabolism in the Liver and Brain of Zebrafish. PLoS One
1216   10:e0143911.
1217   87.   Zhang J, Sun P, Kong T, Yang F, Guan W. 2016. Tributyltin promoted hepatic
1218   steatosis in zebrafish (Danio rerio) and the molecular pathogenesis involved.
1219   Aquat Toxicol 170:208-215.
1220   88.   Riu A, McCollum CW, Pinto CL, Grimaldi M, Hillenweck A, Perdu E, Zalko D,
1221   Bernard L, Laudet V, Balaguer P, Bondesson M, Gustafsson JA. 2014.
1222   Halogenated bisphenol-A analogs act as obesogens in zebrafish larvae (Danio
1223   rerio). Toxicol Sci 139:48-58.
1224   89.   Lima D, Castro LF, Coelho I, Lacerda R, Gesto M, Soares J, Andre A, Capela R,
1225   Torres T, Carvalho AP, Santos MM. 2015. Effects of Tributyltin and Other
1226   Retinoid Receptor Agonists in Reproductive-Related Endpoints in the Zebrafish
1227   (Danio rerio). J Toxicol Environ Health A 78:747-60.
1228   90.   Shimasaki Y, Kitano T, Oshima Y, Inoue S, Imada N, Honjo T. 2003. Tributyltin
1229   causes masculinization in fish. Environ Toxicol Chem 22:141-4.
1230   91.   Manning CS, Lytle TF, Walker WW, Lytle JS. 1999. Life-cycle toxicity of
1231   bis(tributyltin) oxide to the sheepshead minnow (Cyprinodon variegatus).
1232   Archives of Environmental Contamination and Toxicology 37:258-266.
1233   92.   Guo SZ, Qian LJ, Shi HH, Barry T, Cao QZ, Liu JQ. 2010. Effects of tributyltin
1234   (TBT) on Xenopus tropicalis embryos at environmentally relevant concentrations.
1235   Chemosphere 79:529-533.
1236   93.   Seinen W, Helder T, Vernij H, Penninks A, Leeuwangh P. 1981. Short-Term
1237   Toxicity of Tri-N-Butyltinchloride in Rainbow-Trout (Salmo-Gairdneri Richardson)
1238   Yolk-Sac Fry. Science of the Total Environment 19:155-166.
1239   94.   Omura M, Ogata R, Kubo K, Shimasaki Y, Aou S, Oshima Y, Tanaka A, Hirata
1240   M, Makita Y, Inoue N. 2001. Two-generation reproductive toxicity study of
1241   tributyltin chloride in male rats. Toxicological Sciences 64:224-232.
1242   95.   Adeeko A, Li D, Forsyth DS, Casey V, Cooke GM, Barthelemy J, Cyr DG, Trasler
1243   JM, Robaire B, Hales BF. 2003. Effects of in utero tributyltin chloride exposure in
1244   the rat on pregnancy outcome. Toxicol Sci 74:407-15.
1245   96.   Popkin BM, Corvalan C, Grummer-Strawn LM. 2020. Dynamics of the double
1246   burden of malnutrition and the changing nutrition reality. Lancet 395:65-74.
1247   97.   Wells JC, Sawaya AL, Wibaek R, Mwangome M, Poullas MS, Yajnik CS, Demaio
1248   A. 2020. The double burden of malnutrition: aetiological pathways and
1249   consequences for health. Lancet 395:75-88.

98. Gaulke CA, Barton CL, Proffitt S, Tanguay RL, Sharpton TJ. 2016. Triclosan
Exposure Is Associated with Rapid Restructuring of the Microbiome in Adult
Zebrafish. PLoS One 11:e0154632.
99. Zhou L, Limbu SM, Shen M, Zhai W, Qiao F, He A, Du ZY, Zhang M. 2018.
Environmental concentrations of antibiotics impair zebrafish gut health. Environ
Pollut 235:245-254.
100. Catron TR, Keely SP, Brinkman NE, Zurlinden TJ, Wood CE, Wright JR, Phelps
D, Wheaton E, Kvasnicka A, Gaballah S, Lamendella R, Tal T. 2019. Host
Developmental Toxicity of BPA and BPA Alternatives Is Inversely Related to
Microbiota Disruption in Zebrafish. Toxicol Sci 167:468-483.
101. Morgun A, Dzutsev A, Dong X, Greer RL, Sexton DJ, Ravel J, Schuster M, Hsiao
W, Matzinger P, Shulzhenko N. 2015. Uncovering effects of antibiotics on the
host and microbiota using transkingdom gene networks. Gut 64:1732-43.
102. Lin XB, Farhangfar A, Valcheva R, Sawyer MB, Dieleman L, Schieber A, Ganzle
MG, Baracos V. 2014. The role of intestinal microbiota in development of
irinotecan toxicity and in toxicity reduction through dietary fibres in rats. PLoS
One 9:e83644.
103. Kang DJ, Kakiyama G, Betrapally NS, Herzog J, Nittono H, Hylemon PB, Zhou
H, Carroll I, Yang J, Gillevet PM, Jiao C, Takei H, Pandak WM, Iida T, Heuman
DM, Fan S, Fiehn O, Kurosawa T, Sikaroodi M, Sartor RB, Bajaj JS. 2016.
Rifaximin Exerts Beneficial Effects Independent of its Ability to Alter Microbiota
Composition. Clin Transl Gastroenterol 7:e187.
104. Vila AV, Collij V, Sanna S, Sinha T, Imhann F, Bourgonje AR, Mujagic Z, Jonkers
DMAE, Masclee AAM, Fu JY, Kurilshikov A, Wijmenga C, Zhernakova A,
Weersma RK. 2020. Impact of commonly used drugs on the composition and
metabolic function of the gut microbiota. Nature Communications 11.
105. Lee SM, Donaldson GP, Mikulski Z, Boyajian S, Ley K, Mazmanian SK. 2013.
Bacterial colonization factors control specificity and stability of the gut microbiota.
Nature 501:426-9.
106. Lam LH, Monack DM. 2014. Intraspecies competition for niches in the distal gut
dictate transmission during persistent Salmonella infection. PLoS Pathog
10:e1004527.
107. Tai N, Peng J, Liu F, Gulden E, Hu Y, Zhang X, Chen L, Wong FS, Wen L. 2016.
Microbial antigen mimics activate diabetogenic CD8 T cells in NOD mice. J Exp
Med 213:2129-46.
108. Pham LN, Kanther M, Semova I, Rawls JF. 2008. Methods for generating and
colonizing gnotobiotic zebrafish. Nat Protoc 3:1862-75.
109. Wang CY, Zhao Y, Jin YX. 2020. The emerging PFOS alternative OBS exposure
induced gut microbiota dysbiosis and hepatic metabolism disorder in adult
zebrafish. Comparative Biochemistry and Physiology C-Toxicology &
Pharmacology 230.
110. Gonzalez-Penagos CE, Zamora-Briseno JA, Cerqueda-Garcia D, Amendola-
Pimenta M, Perez-Vega JA, Hernandez-Nunez E, Rodriguez-Canul R. 2020.
Alterations in the Gut Microbiota of Zebrafish (Danio rerio) in Response to Water-
Soluble Crude Oil Components and Its Mixture With a Chemical Dispersant.
Front Public Health 8:584953.

111. Brugman S, Liu KY, Lindenbergh-Kortleve D, Sansom JN, Furuta GT, Renshaw
SA, Willemsen R, Nieuwenhuis EES. 2009. Oxazolone-Induced Enterocolitis in
Zebrafish Depends on the Composition of the Intestinal Microbiota.
Gastroenterology 137:1757-1767.
112. Chen LG, Hu CY, Lai NLS, Zhang WP, Hua JH, Lam PKS, Lam JCW, Zhou BS.
2018. Acute exposure to PBDEs at an environmentally realistic concentration
causes abrupt changes in the gut microbiota and host health of zebrafish.
Environmental Pollution 240:17-26.
113. Jin Y, Xia J, Pan Z, Yang J, Wang W, Fu Z. 2018. Polystyrene microplastics
induce microbiota dysbiosis and inflammation in the gut of adult zebrafish.
Environ Pollut 235:322-329.
114. Hajishengallis G, Liang S, Payne MA, Hashim A, Jotwani R, Eskan MA, McIntosh
ML, Alsam A, Kirkwood KL, Lambris JD, Darveau RP, Curtis MA. 2011. Low-
Abundance Biofilm Species Orchestrates Inflammatory Periodontal Disease
through the Commensal Microbiota and Complement. Cell Host & Microbe
10:497-506.
115. Janda JM, Abbott SL, McIver CJ. 2016. Plesiomonas shigelloides Revisited. Clin
Microbiol Rev 29:349-74.
116. Jawahar J, McCumber A, Lickwar C, Amoroso C, Canny SGdlT, Wong S,
Bohannan B, Guillemin K, Rawls J. 2020. Starvation causes changes in the
intestinal transcriptome and microbiome that are reversed upon refeeding
doi:10.21203/rs.3.rs-120052/v1. Research Square.
117. Groussin M, Mazel F, Alm EJ. 2020. Co-evolution and Co-speciation of Host-Gut
Bacteria Systems. Cell Host Microbe 28:12-22.
118. Baillie-Hamilton PF. 2002. Chemical toxins: a hypothesis to explain the global
obesity epidemic. J Altern Complement Med 8:185-92.
119. Anonymous. 1979. Toxic Substances Control Act (TSCA) : Chemical Substance
inventory / United States Environmental Protection Agency, Office of Toxic
Substances. U.S. Environmental Protection Agency, Office of Toxic Substances,
Washington.
120. FAO. 2020. The State of World Fisheries and Aquaculture 2020. In brief.
Sustainability in action. Rome, Italy.
121. Belton B, Thilsted SH. 2014. Fisheries in transition: Food and nutrition security
implications for the global South. Global Food Security-Agriculture Policy
Economics and Environment 3:59-66.
122. Bene C, Arthur R, Norbury H, Allison EH, Beveridge M, Bush S, Campling L,
Leschen W, Little D, Squires D, Thilsted SH, Troell M, Williams M. 2016.
Contribution of Fisheries and Aquaculture to Food Security and Poverty
Reduction: Assessing the Current Evidence. World Development 79:177-196.
123. Murdoch CC, Espenschied ST, Matty MA, Mueller O, Tobin DM, Rawls JF. 2019.
Intestinal Serum amyloid A suppresses systemic neutrophil activation and
bactericidal activity in response to microbiota colonization. PLoS Pathog
15:e1007381.
124. Brandli RC, Breedveld GD, Cornelissen G. 2009. Tributyltin sorption to marine
sedimentary black carbon and to amended activated carbon. Environ Toxicol
Chem 28:503-8.

125. Brosillon S, Bancon-Montigny C, Mendret J. 2014. Study of photocatalytic
degradation of tributyltin, dibutylin and monobutyltin in water and marine
sediments. Chemosphere 109:173-9.
126. Lawrence C. 2007. The husbandry of zebrafish (*Danio rerio*): A review.
Aquaculture 269:1-20.
127. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein
V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for
biological-image analysis. Nat Methods 9:676-82.
128. Melancon E, Gomez De La Torre Canny S, Sichel S, Kelly M, Wiles TJ, Rawls
JF, Eisen JS, Guillemin K. 2017. Best practices for germ-free derivation and
gnotobiotic zebrafish husbandry. Methods Cell Biol 138:61-100.
129. Cassidy-Hanley DM. 2012. Tetrahymena in the laboratory: strain resources,
methods for culture, maintenance, and storage. Methods Cell Biol 109:237-76.

1356

1357

1358 **FIGURES**

1359

Figure 1: Schematic representation of *in vivo* experiments using conventionally-

reared zebrafish. (A) Experimental design for exposure to tributyltin chloride (TBT). We

prepared four tank replicates per chemical treatment. The concentrations of TBT in tank

water are indicated in the schematic. DMSO was used as vehicle control. (B)

Experimental design for supercolonization with individual bacterial strains. We prepared

three tank replicates per bacterial treatment, including the four indicated bacterial

strains and a mock supercolonization negative control of water from the recirculating

aquaculture system (RAS). For both experiments, the timeline represents the fish age in

days post-fertilization (black ticks, dpf) in relation to the days of chemical exposure (red

ticks, dpE) or supercolonization (red ticks, dpSC). The different bar colors represent the

aquaculture systems used for housing the zebrafish and are listed in the box below

panels A and B. We sampled 5 fish per tank to measure body size and adiposity, prior

to dissecting their whole intestine for 16S rRNA gene sequencing analysis of the

intestinal microbial communities. We also sampled 150 ml of tank water for 16S rRNA

gene sequencing analysis of the aquatic microbial communities. (*) The same sampling

strategy was used in all timepoints of both experiments, except at 14 dpSC when we

sampled all the remainder fish as this experiment had only two timepoints.

1377

Figure 2: Exposure to 1 µg/l TBT increases adiposity and inhibits body size *in vivo* after 21 days. (A) Standard length (SL, the distance between the snout and the caudal peduncle) measurements. (B) Adiposity measurements (the total two-dimensional area of adipose tissues per body area excluding fins). Bars represent the mean and error bars the standard deviation. Means were compared to the DMSO control using one-way ANOVA and post-hoc Dunnett's multiple comparison tests. When

significant, the multiplicity adjusted *P*-value is reported for the comparison. (C)

Brightfield images of whole fish at 21 dpE. Selected fish are representative of the mean

SL per chemical treatment. (D) Nile red fluorescence images reveal adipose tissues in

whole fish at 21 dpE. Selected fish are representative of the mean adiposity per

treatment. Their corresponding SL is reported. (E) Linear regression analysis between

SL (x-axis) and total adipose tissue area (y-axis). All samples across all timepoints (7

dpE, 14 dpE, and 21 dpE) and treatments are included. Deviation from linearity per

chemical treatment was tested and found not significant (all $P \geq 0.0853$). Equations and

goodness of fit values per chemical treatment are reported on the panel. The best fit

values of the slopes per treatment were compared to the DMSO control using one-way

ANOVA and post-hoc Dunnett's multiple comparison tests. The only significant adjusted

P-value corresponded to the comparison between the DMSO and 1 µg/l TBT treatments

($P < 0.0005$).

1399

Figure 3: TBT exposure alters the relative abundance of specific bacterial taxa in the zebrafish intestine but not in the water. Relative abundance of selected members of intestinal microbial communities during in vivo exposure to the TBT concentrations and at the time points indicated. Selected OTUs from the intestinal (A, C, E) and aquatic (B, D, F) microbial communities assigned to genera *Chitinibacter* (A, B), *Vibrio* (C, D), and *Plesiomonas* (E, F). Box and whisker plots represent the interquartile

1406   range, with the line in the box defining the median. Whiskers represent the minimum

1407   and maximum value. The mean of each bacterial treatment was compared to the control

1408   using the Mann-Whitney test per time point and sample types. When significant, the P-

1409   values are reported for the comparisons.

1410

1411

1412

Figure 4: Supercolonization with *Plesiomonas* ZOR0011 is sufficient to inhibit

adiposity. Zebrafish were supercolonized daily for 14 days with one of four bacterial

strains: *Chitinibacter* ZOR0017, C17; *Chitinibacter* ZOR0013, C13; *Vibrio* ZWU0020,

V20; or *Plesiomonas* ZOR0011, P11. (A) Standard length (SL) and (B) adiposity

measurements before initiating the bacterial treatment (0 days) and after 14 days of

daily supercolonization (dpSC). Bars represent the mean and error bars the standard

deviation. Means were compared to the negative control using unpaired t-test. When

significant, the *P*-value is reported for the comparison. (C) Brightfield images of whole

fish representative of the mean SL per treatment at 14 dpSC. (D) Nile red fluorescence

images reveal adipose tissues in whole fish representative of the mean adiposity per

treatment at 14 dpSC. Their corresponding SL is reported. (E) Relative abundances of

OTUs assigned to genera *Plesiomonas*, *Chitinibacter*, and *Vibrio* in aquatic and

intestinal microbial communities at 14 dpSC. Box and whisker plots represent the

interquartile range, with the line in the box defining the median. Whiskers represent the

minimum and maximum value. The mean of each bacterial treatment was compared to

the control using the Mann-Whitney test. When significant, the *P*-values are reported for

the comparisons.

1430

1431

Figure 5: Monoassociation reveals that *Plesiomonas* ZOR0011 is sufficient to

reduce body size and adiposity in zebrafish without an established microbiota.

(A) Standard length and (B) adiposity after exposure of germ-free zebrafish to TBT; or

monoassociation (colonized) with *Chitinibacter* ZOR0013 (*C*13) and *Plesiomonas*

ZOR0011 (*P*11). Measurements were taken 7 and 14 days post-exposure (dpE) or post-

colonization (dpC). Bars represent the mean and error bars the standard deviation.

Means of each treatment were compared to the DMSO control using one-way ANOVA

1439   and the post-hoc Dunnett's multiple comparison test. When significant, the multiplicity

1440   adjusted *P*-value is reported for the comparison. (C) Brightfield images of whole fish

1441   representative of the mean SL per treatment measured after 14 days. (D) Nile red

1442   fluorescence images reveals adipose tissues in whole fish representative of the mean

1443   adiposity per treatment measured after 14 days. Their corresponding SL are reported.

1444