1

Title:

Remodeling of organelles and microtubules during spermiogenesis in the liverwort
Marchantia polymorpha

Running title:

Spermiogenesis in *M. polymorpha*

Authors:

Naoki Minamino[1], Takuya Norizuki[1], Shoji Mano[2, 3], Kazuo Ebine[1, 3], Takashi Ueda[1, 3]

[1]Division of Cellular Dynamics, National Institute for Basic Biology, Nishigonaka 38,
Myodaiji, Okazaki, Aichi, 444-8585, Japan

[2]Laboratory of Organelle Regulation, National Institute for Basic Biology, Nishigonaka 38,
Myodaiji, Okazaki, Aichi, 444-8585, Japan

[3]Department of Basic Biology, SOKENDAI (The Graduate University for Advanced Studies),
Nishigonaka 38, Myodaiji, Okazaki, Aichi, 444-8585, Japan

corresponding author's email address

tueda@nibb.ac.jp

Keywords

Spermiogenesis, microtubule, organelle, *Marchantia polymorpha*

Summary statement

 We classified developmental processes of spermatozoids into 1 + 5 stages and characterized
remodeling of organelles and microtubules during spermiogenesis in the liverwort
Marchantia polymorpha.

Abstract

Gametogenesis is an essential biological event for sexual reproduction in various organisms.

Bryophytes and some other plants employ motile sperms (spermatozoids) as male gametes,

which self-locomote to the egg cells to accomplish fertilization. Spermatozoids of bryophytes

harbor distinctive morphological characteristics, including the cell body with a helical slender

shape and two motile flagella at the anterior edge. During transformation from a spermatid to

spermatozoid (spermiogenesis), the shape and cellular contents of spermatids are dynamically

reorganized. However, how each organelle is reorganized during plant spermiogenesis

remains obscure. In this study, we classified the developmental processes during

spermiogenesis in the liverwort *Marchantia polymorpha* according to cellular and nuclear

shapes and flagella development. We then examined the remodeling of microtubules and

reorganization of endomembrane organelles during spermiogenesis. The results indicate that

the state of post-translational modification of tubulin is dynamically changed during the

formation of the flagella and spline, and the plasma membrane and endomembrane organelles

are drastically reorganized in a precisely regulated manner during spermiogenesis. These

findings are expected to provide useful indexes to classify developmental and subcellular

processes of spermiogenesis in bryophytes.

Introduction

Sexual reproduction, one of the most crucial events in the lifecycle of multicellular organisms,

is necessary for genetic variation and thus adaptation to environments. To accomplish sexual

reproduction, reproductive cells, which are generally classified into male and female gametes,

are generated through precisely regulated differentiation and maturation processes. A

majority of seed plants generate non-motile male gametes, which are transported to the egg

cells via pollen tubes for fertilization. Conversely, some species in charophycean algae as

well as bryophytes, lycophytes, ferns, and some groups of gymnosperms, such as, Ginkgo

and cycads, generate motile sperm cells (spermatozoids) as male gametes, which harbor two

or more motile flagella and move to egg cells in water to accomplish fertilization.

Gametogenesis in angiosperms has been intensively studied using several species including

Arabidopsis, and its mechanisms are well documented at the genetic and molecular levels

(Berger and Twell, 2011; Hackenberg and Twell, 2019; Russell and Jones, 2015; Twell,

2011). Conversely, molecular genetic studies of gametogenesis in basal land plants producing

spermatozoids have only recently begun, using a few bryophyte models such as

Physcomitrium patens and *Marchantia polymorpha* (Higo et al., 2018; Koi et al., 2016;

Koshimizu et al., 2018; Minamino et al., 2017; Sanchez-Vera et al., 2017).

Spermatozoids of bryophytes have distinctive morphological characteristics, some of which

are also shared with spermatozoids of other plant lineages. The cell body exhibits a helical

shape and consists mostly of an elongated nucleus. The anterior region of the cell body

contains one mitochondrion, which is associated with the multilayered structure (MLS), a

characteristic of plant spermatids. The uppermost layer of the MLS (the opposite side to the

mitochondrion) is a bundle of microtubules, termed the spline, which extends along the

helical nucleus through the cell body. Two flagella elongate from two basal bodies, which are

attached to the spline of the MLS. The posterior side of the cell body contains the other

mitochondrion and a plastid, and other cytoplasmic components are not appreciably observed

in mature spermatozoids (Fig. 1A; Shimamura, 2016). During the transformation from

spermatids to spermatozoids, which is termed spermiogenesis, the cell shape and cellular

contents of spermatids are drastically reorganized. The MLS and flagella are formed *de novo*

during spermiogenesis, the process that has been investigated mainly using electron

microscopy (Carothers and Kreitner, 1967; Carothers and Kreitner, 1968; Kreitner, 1977a;

Kreitner and Carothers, 1976; Renzaglia and Duckett, 1987; Renzaglia and Garbary, 2001). It

is expected that most of the cytoplasm, including various organelles other than the nucleus,

anterior and posterior mitochondria, and a plastid, is eliminated during spermiogenesis.

However, when and how organelles are removed from spermatids during spermiogenesis

remains unknown, mainly due to the lack of proper methods to observe spatiotemporal

organelle dynamics during plant spermiogenesis.

Marchantia polymorpha is a model of liverworts, a group of bryophytes whose genomic

information is available (Bowman et al., 2017; Montgomery et al., 2020), and various tools

for molecular genetic analyses have been established (Kohchi et al., 2021). Transcriptome

data and a wide variety of organelle markers have also been developed for *M. polymorpha*

(Bowman et al., 2017; Higo et al., 2016; Kanazawa et al., 2016; Minamino et al., 2017;

Minamino et al., 2018). Employing this relatively new model plant for analyzing the cellular

dynamics during plant spermiogenesis, we previously reported that subcellular localizations

of some organelle markers drastically change in spermatogenous tissue (Minamino et al.,

2017). However, the temporal resolution was not sufficient because of the lack of defined

developmental stages in spermiogenesis in *M. polymorpha*. In this study, we first classified

the developmental processes of spermiogenesis in *M. polymorpha* according to cellular and

nuclear morphology and flagellar formation. We then examined microtubule organization in

each developmental stage during spermiogenesis and found that the state of post-translational

tubulin modification is significantly changed during the development of microtubule-

containing structures. Furthermore, we monitored markers of endomembrane organelles in

each developmental stage and found that each organelle/organelle protein is degraded at a

distinctive stage during spermiogenesis. These results demonstrate a drastic but highly

organized rearrangement of cellular components during spermiogenesis in *M. polymorpha*

and provide a unified standard for classification of developmental stages of spermatids

undergoing spermiogenesis in bryophytes.

Results

Determining developmental stages of spermiogenesis in *M. polymorpha*

An antheridium of *M. polymorpha* consists of outer jacket cells and inner reproductive cells.

Spermatogenous cells proliferate by continuous transverse and vertical cell division, and the

resultant spermatid mother cells divide diagonally to generate spermatids (Fig. 1B;

Shimamura, 2016). Spermatids then transform into mature spermatozoids through

spermiogenesis. Spermiogenesis of *M. polymorpha* comprises dynamic cellular

reorganization events, including condensation and elongation of the nucleus, *de novo*

synthesis of the locomotory apparatus, and elimination of the cytoplasm (Fig. 1B; Shimamura,

2016). Based on the morphology of the cell body, nuclear shape observed with Hoechst

33342, and formation of flagella as yardsticks, we first attempted to classify spermatid cells

undergoing spermiogenesis, which were fixed with paraformaldehyde and separated from

each other by treating with cell-wall digesting enzymes, into 1 + 5 developmental stages (Fig.

1C-H). The cells harboring the round nucleus without visible flagella were classified as stage

0. These cells presumably contained spermatogenous cells, spermatid mother cells, and early-

stage spermatids, which were not distinguishable from each other based on their morphology

after cell wall digestion (Fig. 1C). At stage 1, protruding or elongating flagella were observed,

whereas the nucleus remained spherical (Fig. 1D). A change in the nuclear shape was

detected beginning from stage 2. At this stage, spermatids were equipped with fully elongated

flagella, and projection of the anterior side of the nucleus was also visible (Fig. 1E). At stage

3, the nucleus became crescent-shaped (Fig. 1F), and a cylindrical nucleus was observed at

stage 4 (Fig. 1G). Mature spermatozoids were defined as stage 5 (Fig. 1H).

Altered modification of microtubules during spermiogenesis in *M. polymorpha*

The spermatozoid of *M. polymorpha* comprises two distinctive microtubule-containing

structures, the flagella and spline. To examine how microtubules are organized during

spermiogenesis to form these structures, we performed immunostaining of tubulin in

spermatids at distinct developmental stages. In addition to immunostaining using an anti-α-

tubulin antibody, we also performed staining with an anti-poly-glutamate (polyE) antibody,

which recognizes the carboxyl-terminally located linear alpha-glutamate chain comprising

four or more glutamate residues. This method was chosen knowing that post-translational

modifications, including glutamylation, are detected in persistent microtubules in the

axoneme in animals and in Chlamydomonas (Janke and Magiera, 2020; Wloga et al., 2017).

In some cells at stage 0, only the anti-α-tubulin antibody-stained fibrous structures were

observed near the cell surface (Fig. 2A). In other cells, rod-like forming flagella, which had

not yet protruded from the cell body, were stained by both the anti-α-tubulin and anti-polyE

antibodies (Fig. 2B). In stage 1 spermatids, elongating flagella were observed, stained by

both antibodies (Fig. 2C). Notably, microtubule fibers extending radially from or toward the

basal region of the forming flagella were frequently observed at stages 0 and 1 (Fig. 2B,C).

The radial fibers were not observed in stage 2 spermatids, whereas the spline microtubule

was stained by the anti-α-tubulin antibody in stage 2 and 3 spermatids (Fig. 2D,E). The radial

structure and spline were not stained by polyE antibody (Fig. 2B-E). Intriguingly, the signal

intensity from α-tubulin on flagella became weaker except for that from the distal-most

region at stages 2 to 3; however, the signal from polyE was uniformly detected at the

corresponding region (Fig. 2D,E). Similar results were also obtained when we used another

anti-α-tubulin and anti-β-tubulin antibodies (Fig. S2A-D). In stage 4, the polyE signal was

detected on flagella, but α-tubulin was not detected on either the flagella or spline (Fig. 2F).

At stage 5 (mature spermatozoids), the α-tubulin signal was detected again on both flagella

and spline. The polyE signal was also detected on flagella and spline, although the intensity

of the signal from the spline was weak compared with that from the flagella (Fig. 2G; Fig.

S1). We then performed immunoblotting to examine whether the anti-polyE antibody

recognized the same tubulin population as anti-α- and anti-β-tubulin antibodies. The anti-α-

and β-tubulin antibodies detected tubulin molecules at approximately 50 kDa, whereas the

anti-polyE antibody reacted with the product at approximately 58 kDa in cell lysates prepared

from antheridia (Fig. S3). The anti-α- and -β-tubulin antibodies did not react with the 58 kDa

product detected with the anti-polyE antibody, and vice versa. Given that the shift of the

molecular weight would reflect the covalently added poly-glutamate chain, as reported in

previous studies (Kubo et al., 2010; Kubo et al., 2014; Vu et al., 2016), and that both the

epitopes of the anti-tubulin antibodies used in this study and polyglutamylation sites are

mapped to the C-terminal region of tubulin, the anti-tubulin antibodies and the anti-polyE

antibody react exclusively with distinct populations of tubulin molecules. Thus, our results

strongly suggest that tubulin molecules in the axoneme and spline undergo progressive

alteration in post-translational modification during spermiogenesis.

Selection of promoters for expression of organelle markers in spermatogenous tissue of

M. polymorpha

With the goal of observing the behavior of endomembrane organelles during spermiogenesis

without interference from surrounding tissues, we expressed organelle markers under the

regulation of promoters predominantly active in spermatids. We selected three candidate

genes, which were reported to express strongly during spermiogenesis in *M. polymorpha*.

Mp*DUO1* is a MYB transcription factor, which is required for sperm differentiation of *M.*

polymorpha (Higo et al., 2018). Mp*CEN1* is a homolog of CENTRIN essential for the

formation of locomotory apparatus in the fern *Marsilea vestita* and is strongly expressed in

spermatogenous tissue independent of Mp*DUO1* in *M. polymorpha* (Higo et al., 2018; Klink

and Wolniak, 2001). IFT52 is one of the components of the intraflagellar transport complex

essential for flagellar formation in Chlamydomonas (Brazelton et al., 2001; Deane et al.,

2001), whose homolog, Mp*IFT52*, in *M. polymorpha*, is abundantly expressed in

antheridiophores according to the transcriptome data deposited in MarpolBase

(https://marchantia.info/). Transgenic lines expressing the fluorescent protein Citrine driven

by the promoters of these genes exhibited strong expression in spermatogenous tissue (Fig.

S4A-C), suggesting that these promoters may be useful for expression of organelle markers

during spermatogenesis. We thus used these promoters for further experiments in this study

(Fig. S4D).

Reorganization of vacuoles during spermiogenesis in *M. polymorpha*

We previously reported that plasma membrane (PM) proteins and several endomembrane

organelle proteins are transported into the luminal space of spherical vacuoles to be degraded

during spermiogenesis (Minamino et al., 2017). However, it remains unknown when the

vacuole transforms to spherical shape, and it also remains unclear how transport of organelle

proteins to the vacuole is organized. To determine this, we first examined alteration of the

vacuolar morphology during spermiogenesis using a vacuole-residing soluble *N*-

ethylmaleimide sensitive factor attachment protein receptor (SNARE), MpVAMP71, tagged

with monomeric Citrine (mCitrine-MpVAMP71), expressed under the regulation of the

Mp*DUO1* promoter (Kanazawa et al., 2016). In cells at stages 0 and 1, small and fragmented

vacuoles with intricate morphology were observed around the nucleus (Fig. 3A,B). However,

only one spherical vacuole was observed in each spermatid at stage 2 (Fig. 3C), which

persisted through stages 3 and 4 (Fig. 3D,E). In mature spermatozoids (stage 5),

compartments with mCitrine-MpVAMP71 were not detected (Fig. 3F). Thus, the vacuole

drastically changes its morphology and number before stage 2 to form a spherical vacuole,

which is eliminated after stage 4 to accomplish spermiogenesis.

Dynamic reorganization of the plasma and endo-membranes during spermiogenesis

A plasma membrane SNARE, MpSYP12A (Kanazawa et al., 2016), tagged with mCitrine

(mCitrine-MpSYP12A), was shown to be removed from the PM during spermiogenesis,

suggesting that drastic remodeling also takes place in the PM in spermatids undergoing

spermiogenesis (Minamino et al., 2017). To elucidate the precise timing of this event, we

observed spermatids expressing mCitrine-MpSYP12A at each developmental stage. At stage

0, mCitrine-MpSYP12A was detected on the PM and punctate compartments (Fig. 4A,B). At

stage 1, the fluorescent signal on the PM was diminished, and bright fluorescence was

observed in punctate compartments (Fig. 4C), which probably represents small vacuoles

observed in spermatids at stage 1 (Fig. 3B). The signal from mCitrine was still detectable in

the lumen of the larger spherical vacuole in some spermatids at stage 2 (Fig. 4D), whereas

only faint or almost no fluorescence was detected in the majority of spermatids at this stage

(Fig. 4E). We did not detect any fluorescence after stage 3 (Fig. 4F-H). These results indicate

that endocytosis was highly activated between stages 0 and 1, and at least a set of PM

proteins including MpSYP12A was removed almost completely from the PM and degraded in

the vacuole during stage 2, suggesting rapid remodeling of the PM at early stages during

spermiogenesis.

The Golgi apparatus also exhibited drastic changes in morphology and number during

spermiogenesis. We observed the Golgi apparatus during spermiogenesis using the Venus-

fused transmembrane domain of sialyltransferase (ST) derived from rat, which is localized to

the Golgi apparatus in *M. polymorpha* (Kanazawa et al., 2016). While several discrete Golgi

apparatuses were observed at stage 0, only one Golgi apparatus, which was enlarged

compared with the Golgi at earlier stages, was observed in cells at stage 2 (Fig. 5A-C,G,H).

Appearance of the large Golgi apparatus in spermatids undergoing spermiogenesis is also

reported in electron microscopic observations of another liverwort, *Blasia pusilla* (Renzaglia

and Duckett, 1987). The fluorescence from ST-Venus was decreased in spermatids at stage 3

and was not detectable in cells at stages 4 and 5 (Fig. 5D-F). We also observed similar

behavior of the Golgi apparatus using other Golgi markers, mCitrine-fused MpGOS11, and

MpSFT1 (Kanazawa et al., 2016); these markers were hardly detected at as early as stage 3

(Fig. S5). Thus, one large Golgi apparatus is formed during early stages of spermiogenesis,

which is degraded during later stages with distinct temporal regulation depending on proteins.

We then observed changes in the endoplasmic reticulum (ER). As an ER membrane marker,

we used mCitrine-fused MpSEC22 (mCitrine-MpSEC22, Kanazawa et al., 2016). We also

monitored a soluble ER protein marker SP-mCitrine-HDEL, which consists of mCitrine

tagged with the signal peptide (SP) at its N-terminus and four amino acids (His-Asp-Glu-

Leu) sufficient for ER localization at the C-terminus (Mano et al., 2018). These ER markers

illuminated the nuclear envelope (NE) and the ER until stage 2 (Fig. 6A-F). Intriguingly, the

fluorescence at the NE was hardly detected at stage 3, whereas the signal on the ER remained

detectable at this stage (Fig. 6G-J), suggesting distinct regulation of remodeling between the

NE and ER during spermiogenesis. Furthermore, we found that these two ER markers

behaved distinctly at later stages; SP-mCitrine-HDEL remained at the anterior region of

spermatozoids, and a faint signal was also detected in other regions (Fig. 6L), whereas the

signal from mCitrine-MpSEC22 was not detected in spermatozoids (Fig. 6K). Thus,

remodeling of the ER also takes place in a highly organized manner, with distinct temporal

regulation depending on domains and proteins.

Discussion

The unified criterion for staging developmental processes is useful for developmental studies

of any organisms, as exemplified by human embryogenesis or spermatogenesis staging, or

development of the flower or anther in Arabidopsis (Bowman et al., 1991; Meistrich and

Hess, 2013; O'Rahilly and Müller, 1987; Oakberg, 1956; Sanders et al., 1999). A shared

criterion for staging developmental processes would also be quite useful in studies of plant

spermatozoid development. In this study, we propose 1 + 5 stages of spermiogenesis in *M.*

polymorpha, according to the shape of the cell body and nucleus and flagellar formation (Fig.

7). The continuous morphological changes of the nucleus in fixed spermatids were roughly

classified as spherical, drop-shaped, crescent-shaped, or cylinder-shaped (Fig. 1A-F), which

is also consistent with previous observations using electron microscopy (Kreitner, 1977b).

The change in nuclear morphology during spermiogenesis is accompanied by chromatin

condensation, which is associated with conversion of DNA-binding nuclear basic proteins

from histones to protamine-like proteins (D'Ippolito et al., 2019; Higo et al., 2016).

Chromatin condensation also occurs during spermiogenesis in animal systems, and

transcription ceases completely around mid-spermiogenesis as chromatin condensation

progresses (Kierszenbaum and Tres, 1975; Sassone-Corsi, 2002). Further progression of

spermiogenesis is mediated by translating mRNA that is accumulated before spermiogenesis

at appropriate time points (Chalmel and Rolland, 2015; Dai et al., 2019). Similar cessation of

transcription may occur during spermatogenesis in plants; therefore, morphological

characteristics, rather than transcriptome information, would be useful indexes for

distinguishing developmental stages of plant spermiogenesis.

After dividing the process of transformation from spermatids to spermatozoids into 1 + 5

stages, we analyzed microtubule organization in spermatids undergoing spermiogenesis and

spermatozoids by immunostaining with antibodies for tubulin or a post-translational

modification of tubulin (polyE). At stage 1, microtubules radially extending from the basal

region of elongating flagella were observed (Fig. 2B,C). This structure was a transient

structure and disappeared before stage 2, at which point flagellar elongation seemed to be

nearly complete (Fig. 2D). Therefore, the radial microtubules might function as rails for

transporting components of flagella to the bases of the flagella. A similar structure, termed

rootlet microtubules, has been observed in Chlamydomonas and which extends from the

basal bodies and have been involved in arrangement of the eyespot (Boyd et al., 2011).

Although spermatids of *M. polymorpha* do not contain the eye spot, the radial structure might

be involved in the arrangement of organelles, such as mitochondria, during spermatogenesis.

Tubulin of the basal bodies and axonemes undergoes several post-translational modifications

including glutamylation (Janke and Magiera, 2020; Wloga et al., 2017). We found that the

flagella and spline in spermatozoids of *M. polymorpha* were recognized by the anti-polyE

antibody (Fig. 2; Fig. S1, S2), indicating that axonemal and spline microtubules are

glutamylated in the basal land plant. Glutamylation is reported to be involved in regulation of

flagellar motility in Chlamydomonas and Tetrahymena (Kubo et al., 2010; Suryavanshi et al.,

2010), assembly of doublet microtubules in mouse (Konno et al., 2016; Lee et al., 2013), and

stability of microtubules in *Caenorhabditis elegans* (O'Hagan and Barr, 2012; O'Hagan et al.,

2011). The function of the glutamylation of axonemal and spline microtubules in plant

spermatozoids should be verified in future studies.

We found that the signals of anti-tubulin antibodies used for immunostaining in this study

became weaker as spermiogenesis progressed and then recovered in mature spermatozoids

(Fig. 2; Fig. S2). One plausible explanation of this phenomenon is that the status of post-

translational modification of tubulin, including glutamylation, changes as spermiogenesis

progresses. The α- and β-tubulin molecules are poly-glutamylated at the C-terminal region

(Wloga et al., 2017), and epitopes of anti-tubulin antibodies used in this study are also located

in the C-terminal region of tubulin molecules. Thus, glutamylation could hinder access of the

anti-tubulin antibodies to their epitopes, resulting in diminished signals in immunostaining.

This is consistent with the result of immunoblotting; anti-α- and -β-tubulin antibodies did not

recognize poly-glutamylated tubulin, which was detected by the anti-polyE antibody (Fig.

S3). Thus, our results indicate that an extent of glutamylation of tubulin molecules may

change during spermiogenesis; the degree of glutamylation increases along formation of the

axoneme, and after its completion, microtubules might be de-glutamylated. An effect of post-

translational modification of tubulin on microtubule properties in spermatids and

spermatozoids and its physiological significance would be interesting topics for further study.

We investigated the reorganization of endomembrane organelles with the progression of

spermiogenesis; the results are summarized in Figure 7. The vacuole changed its morphology

and number from a complicated and fragmented shape to a spherical structure at stage 3,

when the nucleus assumed a crescent-like shape (Fig. 3C,D). This morphological transition

suggests that vacuoles actively fuse with each other as spermiogenesis proceeds. It would be

interesting to see whether evolutionarily conserved machinery components of vacuole

biogenesis, such as RAB7/RABG GTPase and the HOPS complex (Brillada et al., 2018; Rojo

et al., 2001; Takemoto et al., 2018), are involved in vacuole remodeling during

spermiogenesis in *M. polymorpha*, which would also provide useful information for

understanding its biological significance. Stage 4 spermatids contained one spherical vacuole

in the cell body, whereas the mature spermatozoid had no discernable vacuoles or remnants

of the vacuole (Fig. 3E,F). Thus, it appears that the vacuole is removed from the spermatid at

the final step of spermiogenesis. In spermiogenesis of Drosophila, the individualization

complex comprising actin cones is required for elimination of unneeded organelles and the

cytosol. The complex is initially formed near the nucleus and moves from the head to the tail

of the spermatid. During this movement, majority of the cytoplasm is removed to form the

cystic bulge, which is finally detached from the tip of the tail (Fabian and Brill, 2012).

Similarly in plants, actin seems to play an important role in the cytoplasm removal during

spermiogenesis, although a structure corresponding to the individual complex has not been

reported (Renzaglia and Garbary, 2001).

Our observation of mCitrine-MpSYP12A revealed that endocytosis of this protein is highly

active between stages 0 and 1 (Fig. 4). This result suggests that the protein composition of the

PM drastically changes at the early stage of spermiogenesis. The spermatozoid PM must have

competence that is distinct from that of the spermatid, such as the ability to respond to an

attractant from egg cells and resistance to low osmolarity to prevent rupture in fresh water

without the rigid cell wall. Therefore, the cell surface of the spermatozoid is reorganized

during spermiogenesis, and the high endocytic activity observed in early spermiogenesis

might contribute to the thorough remodeling of the PM.

The morphology of the Golgi apparatus has diverged among organisms and even among

tissues, probably reflecting diverse and tissue-specific Golgi functions (Ito et al., 2014;

Sengupta and Linstedt, 2011). We found that the number and size of the Golgi apparatus

changed during spermiogenesis (Fig. 5, Fig. S5), which suggests that the function of the

Golgi apparatus changes during spermiogenesis. Furthermore, we found that the signal from

Golgi markers disappeared by stage 4, which strongly suggests that the Golgi apparatus (or a

major set of Golgi proteins) is eliminated before removal of the cytoplasm. Although the

molecular mechanism of the Golgi removal remains ambiguous, we previously found that

autophagy is highly activated during spermiogenesis and a Golgi-like membrane structure is

transported into the vacuole in *M. polymorpha* (Minamino et al., 2017). Therefore, autophagy

could be a key mechanism for Golgi clearance during spermiogenesis, which would be

verified in future studies.

Drastic deformation of the nucleus during spermiogenesis is generally observed in land plants

(Renzaglia and Garbary, 2001). However, the mechanisms of the nuclear remodeling remain

unknown. We found that the signal from SP-mCitrine-HDEL and mCitrine-MpSEC22 at the

NE was markedly decreased during spermiogenesis, suggesting that the composition of the

nuclear envelope proteins drastically changes during spermiogenesis. Because NE-localized

proteins are involved in the regulation of the nuclear shape in Arabidopsis (Goswami et al.,

2020; Meier et al., 2016), the alteration in composition of the NE might be related to the

nuclear reshaping during spermiogenesis in *M. polymorpha*. To our knowledge, the existence

of the ER in bryophyte spermatozoids has not been reported (Renzaglia and Garbary, 2001),

which infers that the ER is largely eliminated from spermatozoids. Surprisingly, however, we

found that the soluble ER marker remained in the cell body of the spermatozoid (Fig. 6L). It

is reported that the anterior tip of the nucleus is embedded between the spline and anterior

mitochondrion during spermiogenesis in *M. polymorpha* (Kreitner, 1977a). The accumulated

ER marker at the anterior region of the cell body could represent this anterior tip of the

nucleus; soluble luminal ER proteins in this region might escape from degradation. Further

analyses would be needed to conclusively demonstrate how the nuclear shaping and ER

remodeling are coordinated during spermiogenesis.

Material and Methods

Plant materials and transformation

Male accession of *Marchantia polymorpha*, Takaragaike-1 (Tak-1), was used throughout this

study. Plants were grown on $1/2 \times$ Gamborg's B5 medium containing 1.4% (w/v) agar at 22

$^{\circ}$C under continuous white light. Transformation was performed according to a previously

described method (Kubota et al., 2013). Transgenic lines were selected with 10 mg·L^{-1}

hygromycin B and 100 mg·L^{-1} cefotaxime for plants transformed with pMpGWB101-based

binary vectors (see below), and 0.5 µM chlorsulfuron and 100 mg·L^{-1} cefotaxime for plants

transformed with pMpGWB301-based binary vectors (see below). Induction of sexual organs

by far-red irradiation was performed as described previously (Chiyoda et al., 2008).

Constructions

Open reading frames (ORFs) and genomic sequences of *M. polymorpha* genes were

amplified by PCR from cDNA and genomic DNA prepared from Tak-1. Amplified fragments

were subcloned into the pENTR^{TM}/D-TOPO vector (Thermo Fisher Scientific) according to

the manufacturer's instructions. The primer sequences and sizes of amplified products are

listed in Table S1. To construct pENTR $_{pro}$Mp*DUO1* and $_{pro}$Mp*IFT52*, the promoter region of

each gene (5.0 and 5.4 kb, respectively) was amplified using Tak-1 genomic DNA, which

was subcloned into the pENTR^{TM}/D-TOPO vector. To construct pENTR $_{pro}$Mp*CEN1*, the

promoter region of Mp*CEN1* (5.0 kb) was amplified with a *Sma*I site at the 3$^{'}$ end and

subcloned into the pENTR^{TM}/D-TOPO vector. The resultant sequence was introduced into

pMpGWB307 (Ishizaki et al., 2015) using Gateway LR Clonase^{TM} II Enzyme Mix (Thermo

Fisher Scientific).

To construct pENTR $_{pro}$Mp*DUO1:mCitrine*, $_{pro}$Mp*CEN1:mCitrine*, and $_{pro}$Mp*IFT52:mCitrine*,

Mp*DUO1*, Mp*CEN1*, and Mp*IFT52* promoter sequences were amplified with a *Sma*I site at

the 3$^{'}$ end and subcloned into the pENTR^{TM}/D-TOPO vector. The cDNA of mCitrine was

amplified and inserted into the *Sma*I site. To construct pMpGWB101-based and

pMpGWB301-based Gateway vectors (Fig. S4D), the promoter regions followed by cDNA

for mCitrine or only the promoter regions were amplified from plasmids described above,

which were inserted at the *Hin*dIII site of pMpGWB101 or pMpGWB301 using the In-fusion

HD Cloning System (Clontech). To construct $_{pro}$Mp*DUO1*:*ST-Venus*, pENTR *ST-Venus*

(Kanazawa et al., 2016; Uemura et al., 2012) was subjected to LR reaction with

pMpGWB301 $_{pro}$Mp*DUO1*. To construct $_{pro}$Mp*DUO1: mCitrine*-Mp*SYP12A*, $_{pro}$Mp*DUO1:*

mCitrine-Mp*VAMP71,* $_{pro}$Mp*DUO1:mCitrine*-Mp*GOS11*, $_{pro}$Mp*DUO1:mCitrine*-Mp*SFT1*,

and $_{pro}$Mp*DUO1:mCitrine*-Mp*SEC22*, pENTR Mp*SYP12A*, pENTR Mp*VAMP71*, pENTR

Mp*GOS11*, pENTR Mp*SFT1*, and pENTR Mp*SEC22* (Kanazawa et al., 2016) were subjected

to LR recombination with pMpGWB301 $_{pro}$Mp*DUO1:mCitrine*, respectively. To construct

$_{pro}$Mp*EF1α*:*SP-mCitrine-HDEL* fusion genes, pDONR $_{pro}$Mp*EF1α* (Mano et al., 2018) was

subjected to LR recombination with R4pMpGWB394 (Mano et al., 2018).

Microscopy

To prepare the antheridial cells of *M. polymorpha* for observation, the antheridial receptacles

between stages 3 and 5 (Higo et al., 2016) were sliced manually with a razor blade, placed on

a glass slide (Matsunami), and then covered with a coverslip. To prepare the spermatids of *M.*

polymorpha, antheridia were fixed for over 60 min with 4% (w/v) paraformaldehyde (PFA)

in PME buffer (50 mM PIPES-NaOH, 5 mM EGTA, and 1 mM MgSO$_4$, pH 6.8), and treated

for 30 min with cell wall digestion buffer (1% (w/v) cellulase Onozuka RS (SERVA), 0.25%

(w/v) pectolyase Y-23 (KYOWA CHEMICALPRODUCTS), 1% (w/v) BSA, 0.1% (w/v)

NP-40, 1% glucose, and 1 x cOmplete^{TM} EDTA-free protease inhibitor cocktail (Roche

Applied Science) in PME buffer). The samples were placed on a glass slide and mounted

with PBS containing 0.1% Hoechst 33342. To prepare the mature spermatozoids of *M.*

polymorpha, collected spermatozoids were fixed for over 60 min with 4% (w/v) PFA in PBS

buffer, placed on a glass slide, washed with PBS buffer three times, and mounted with PBS

containing 0.1% Hoechst 33342. The prepared samples were observed under LSM780 (Carl

Zeiss) with an oil immersion lens (x63).

For immunostaining of spermatids of *M. polymorpha*, we slightly modified the method

described in Shimamura (2015). Antheridia were fixed for over 60 min with 4% (w/v) PFA in

PME buffer and treated for 30 min with the cell wall digestion buffer. Cells were then treated

with permeabilization buffer (0.01% (v/v) Triton X-100 and 1% (w/v) BSA in PME buffer)

for 10 min. After washing with PME buffer three times, cells were placed on a MAS-coated

glass slide, and incubated for 30 min at room temperature with blocking solution (1% (w/v)

BSA in PBS buffer). After removing the blocking solution, cells were incubated with the

primary antibody in PBS buffer at 4 $^{\circ}$C overnight. After washing with PBS buffer three times,

the samples were incubated for 60 min at 37 $^{\circ}$C with the secondary antibody and 0.1% (v/v)

Hoechst 33342 in PBS buffer. After washing with PBS buffer three times, slides were

mounted using the ProLong Diamond Antifade reagent (Thermo Fisher Scientific).

Immunostaining of mature spermatozoids was performed almost similar to that of

spermatozoids, with additional centrifugation to collect spermatozoids. Samples were

observed under LSM780 (Carl Zeiss) with an oil immersion lens (x63). The reactivities of the

antibodies were examined using immunoblotting with cell lysates prepared from antheridia of

Tak-1.

Antibodies

Anti α-tubulin antibody (DM1A) and anti α-tubulin antibody (B-5-1-2) were purchased from

Sigma-Aldrich and used at ×1,000 dilution for immunostaining (DM1A and B-5-1-2) and

immunblotting (B-5-1-2). Anti β-tubulin antibody (AA2) was purchased from Abcam and

used at ×1,000 dilution for immunostaining and immunoblotting. Anti polyE antibody was

purchased from AdipoGen Life Sciences (AG25B-0030-C050) and used at ×5,000 dilution

for immunoblotting and ×10,000 dilution for immunostaining. Alexa Fluor 488 plus goat

anti-mouse IgG (A32723) and Alexa Fluor 594 plus goat anti-rabbit IgG (A32740) were

purchased from Thermo Fisher Scientific and used at ×1,000 dilution for immunostaining.

Gene accession numbers

Mp*SYP12A* (Mapoly0163s0015, Mp6g00050), Mp*VAMP71* (Mapoly0064s0109,

Mp8g00880), Mp*SYP2* (Mapoly0187s0013, Mp8g15260), Mp*DUO1* (Mapoly0019s0071,

Mp1g13010), Mp*CEN1* (Mapoly0103s0016, Mp1g00710), Mp*IFT52* (Mapoly0059s0029,

Mp6g13200), Mp*GOS11* (Mapoly0001s0245, Mp1g19070), Mp*SFT1* (Mapoly0006s0109,

Mp3g06390), Mp*SEC22* (Mapoly0023s0071, Mp2g11050)

Acknowledgments

We thank Dr. Takehiko Kanazawa (NIBB) for sharing plasmids used in this study. We also

thank Dr. Masaki Shimamura (Hiroshima University) for sharing the protocol for

immunostaining. We thank the Model Plant Research Facility of the NIBB Bioresource

Center for technical support.

Competing interests

No competing interests declared.

Funding

This work was supported by the following grants from the KAKENHI Program of the Japan Society for the Promotion of Science (JSPS): Grant No. 20K15824 to N.M., Grants No. 17H05850, 19H04872, and 21K06222 to K.E., and Grants No. 19H05675, 19H05670, and 21H02515 to T.U., as well as a Grant-in-Aid for JSPS fellows (to TN, Grant No. 19J13751).

References

Berger, F. and Twell, D. (2011). Germline specification and function in plants. *Annu. Rev.*
Plant Biol. **62**, 461–484, doi:10.1146/annurev-arplant-042110-103824.

Bowman, J. L., Drews, G. N. and Meyerowitz, E. M. (1991). Expression of the
Arabidopsis floral homeotic gene AGAMOUS is restricted to specific cell types late in
flower development. *Plant Cell* **3**, 749–758, doi:10.1105/tpc.3.8.749.

Bowman, J. L., Kohchi, T., Yamato, K. T., Jenkins, J., Shu, S., Ishizaki, K., Yamaoka,
S., Nishihama, R., Nakamura, Y., Berger, F., et al. (2017). Insights into land plant
evolution garnered from the *Marchantia polymorpha* genome. *Cell* **171**, 287-304.e15,
doi:10.1016/j.cell.2017.09.030.

Boyd, J. S., Gray, M. M., Thompson, M. D., Horst, C. J. and Dieckmann, C. L. (2011).
The daughter four-membered microtubule rootlet determines anterior-posterior
positioning of the eyespot in *Chlamydomonas reinhardtii*. *Cytoskeleton* **68**, 459–469,
doi:10.1002/cm.20524.

Brazelton, W. J., Amundsen, C. D., Silflow, C. D. and Lefebvre, P. A. (2001). The *bld1*
mutation identifies the *Chlamydomonas osm-6* homolog as a gene required for flagellar
assembly. *Curr. Biol.* **11**, 1591–1594, doi:10.1016/S0960-9822(01)00485-7.

Brillada, C., Zheng, J., Krüger, F., Rovira-Diaz, E., Askani, J. C., Schumacher, K. and
Rojas-Pierce, M. (2018). Phosphoinositides control the localization of HOPS subunit
VPS41, which together with VPS33 mediates vacuole fusion in plants. *Proc. Natl. Acad.*
Sci. USA. **115**, 8305–8314, doi:10.1073/pnas.1807763115.

Carothers, Z. B. and Kreitner, G. L. (1967). Studies of spermatogenesis in the hepaticae. I.
Ultrastructure of the *vierergruppe* in *Marchantia*. *J. Cell Biol.* **33**, 43–51,
doi:10.1083/jcb.33.1.43.

Carothers, Z. B. and Kreitner, G. L. (1968). Studies of spermatogenesis in the Hepaticae.
II. Blepharoplast structure in the spermatid of *Marchantia*. *J. Cell Biol.* **36**, 603–616,
doi:10.1083/jcb.36.3.603.

Chalmel, F. and Rolland, A. D. (2015). Linking transcriptomics and proteomics in
spermatogenesis. *Reproduction* **150**, 149–157, doi:10.1530/REP-15-0073.

Chiyoda, S., Ishizaki, K., Kataoka, H., Yamato, K. T. and Kohchi, T. (2008). Direct
transformation of the liverwort *Marchantia polymorpha* L. by particle bombardment
using immature thalli developing from spores. *Plant Cell Rep.* **27**, 1467–1473,
doi:10.1007/s00299-008-0570-5.

D'Ippolito, R. A., Minamino, N., Rivera-Casas, C., Cheema, M. S., Bai, D. L., Kasinsky, H. E., Shabanowitz, J., Eirin-Lopez, J. M., Ueda, T., Hunt, D. F., et al. (2019). Protamines from liverwort are produced by post-translational cleavage and C-terminal di-aminopropanelation of several male germ-specific H1 histones. *J. Biol. Chem.* **294**, 16364–16373, doi:10.1074/jbc.RA119.010316.

Dai, P., Wang, X., Gou, L. T., Li, Z. T., Wen, Z., Chen, Z. G., Hua, M. M., Zhong, A., Wang, L., Su, H., et al. (2019). A translation-activating function of MIWI/piRNA during mouse spermiogenesis. *Cell* **179**, 1566–1581, doi:10.1016/j.cell.2019.11.022.

Deane, J. A., Cole, D. G., Seeley, E. S., Diener, D. R. and Rosenbaum, J. L. (2001). Localization of intraflagellar transport protein IFT52 identifies basal body transitional fibers as the docking site for IFT particles. *Curr. Biol.* **11**, 1586–1590, doi:10.1016/S0960-9822(01)00484-5.

Fabian, L. and Brill, J. A. (2012). Drosophila spermiogenesis. *Spermatogenesis* **2**, 197–212, doi:10.4161/spmg.21798.

Goswami, R., Asnacios, A., Hamant, O. and Chabouté, M. E. (2020). Is the plant nucleus a mechanical rheostat? *Curr. Opin. Plant Biol.* **57**, 155–163, doi:10.1016/j.pbi.2020.09.001.

Hackenberg, D. and Twell, D. (2019). The evolution and patterning of male gametophyte development. *Curr. Top. Dev. Biol.* **131**, 257–298, doi:10.1016/bs.ctdb.2018.10.008.

Higo, A., Niwa, M., Yamato, K. T., Yamada, L., Sawada, H., Sakamoto, T., Kurata, T., Shirakawa, M., Endo, M., Shigenobu, S., et al. (2016). Transcriptional framework of male gametogenesis in the liverwort *Marchantia polymorpha* L. *Plant Cell Physiol.* **57**, 325–338, doi:10.1093/pcp/pcw005.

Higo, A., Kawashima, T., Borg, M., Zhao, M., López-Vidriero, I., Sakayama, H., Montgomery, S. A., Sekimoto, H., Hackenberg, D., Shimamura, M., et al. (2018). Transcription factor DUO1 generated by neo-functionalization is associated with evolution of sperm differentiation in plants. *Nat. Commun.* **9**, 1–13, doi:10.1038/s41467-018-07728-3.

Ishizaki, K., Nishihama, R., Ueda, M., Inoue, K., Ishida, S., Nishimura, Y., Shikanai, T. and Kohchi, T. (2015). Development of gateway binary vector series with four different selection markers for the liverwort *Marchantia polymorpha*. *PLoS One* **10**, 1–13, doi:10.1371/journal.pone.0138876.

Ito, Y., Uemura, T. and Nakano, A. (2014). Formation and maintenance of the Golgi

apparatus in plant cells. *Int. Rev. Cell Mol. Biol.* **310**, 221–287, doi:10.1016/B978-0-12-

800180-6.00006-2.

Janke, C. and Magiera, M. M. (2020). The tubulin code and its role in controlling

microtubule properties and functions. *Nat. Rev. Mol. Cell Biol.* **21**, 307–326,

doi:10.1038/s41580-020-0214-3.

Kanazawa, T., Era, A., Minamino, N., Shikano, Y., Fujimoto, M., Uemura, T.,

Nishihama, R., Yamato, K. T., Ishizaki, K., Nishiyama, T., et al. (2016). SNARE

molecules in *Marchantia polymorpha*: Unique and conserved features of the membrane

fusion machinery. *Plant Cell Physiol.* **57**, 307–324, doi:10.1093/pcp/pcv076.

Kierszenbaum, A. L. and Tres, L. L. (1975). Structural and transcriptional features of the

mouse spermatid genome. *J. Cell Biol.* **65**, 258–270, doi:10.1083/jcb.65.2.258.

Klink, V. P. and Wolniak, S. M. (2001). Centrin is necessary for the formation of the motile

apparatus in spermatids of *Marsilea*. *Mol. Biol. Cell* **12**, 761–776,

doi:10.1091/mbc.12.3.761.

Kohchi, T., Yamato, K. T., Ishizaki, K., Yamaoka, S. and Nishihama, R. (2021).

Development and molecular genetics of *Marchantia polymorpha*. *Annu. Rev. Plant Biol.*

72, 19–20, doi:10.1146/annurev-arplant-082520-094256.

Koi, S., Hisanaga, T., Sato, K., Shimamura, M., Yamato, K. T., Ishizaki, K., Kohchi, T.

and Nakajima, K. (2016). An evolutionarily conserved plant RKD factor controls germ

cell differentiation. *Curr. Biol.* **26**, 1775–1781, doi:10.1016/j.cub.2016.05.013.

Koshimizu, S., Kofuji, R., Sasaki-Sekimoto, Y., Kikkawa, M., Shimojima, M., Ohta, H.,

Shigenobu, S., Kabeya, Y., Hiwatashi, Y., Tamada, Y., et al. (2018). Physcomitrella

MADS-box genes regulate water supply and sperm movement for fertilization. *Nat.*

Plants **4**, 36–45, doi:10.1038/s41477-017-0082-9.

Kreitner, G. L. (1977a). Influence of the multilayered structure on the morphogenesis of

Marchantia spermatids. *Am. J. Bot.* **64**, 57–64, doi:10.1002/j.1537-2197.1977.tb07605.x.

Kreitner, G. L. (1977b). Transformation of the nucleus in Marchantia spermatids:

morphogenesis. *Am. J. Bot.* **64**, 464–475, doi:10.2307/2441777.

Kreitner, G. L. and Carothers, Z. B. (1976). Studies of spermatogenesis in the Hepaticae V.

Blepharoplast development in *Marchantia polymorpha*. *Am. J. Bot.* **63**, 545–557,

doi:10.2307/2441818.

Kubo, T., Yanagisawa, H. aki, Yagi, T., Hirono, M. and Kamiya, R. (2010). Tubulin

polyglutamylation regulates axonemal motility by modulating activities of inner-arm

dyneins. *Curr. Biol.* **20**, 441–445, doi:10.1016/j.cub.2009.12.058.

Kubo, T., Yanagisawa, H. A., Liu, Z., Shibuya, R., Hirono, M. and Kamiya, R. (2014). A

conserved flagella-associated protein in *Chlamydomonas*, FAP234, is essential for

axonemal localization of tubulin polyglutamylase TTLL9. *Mol. Biol. Cell* **25**, 107–117,

doi:10.1091/mbc.E13-07-0424.

Kubota, A., Ishizaki, K., Hosaka, M. and Kohchi, T. (2013). Efficient agrobacterium-

mediated transformation of the liverwort *Marchantia polymorpha* using regenerating

thalli. *Biosci. Biotechnol. Biochem.* **77**, 167–172, doi:10.1271/bbb.120700.

Mano, S., Nishihama, R., Ishida, S., Hikino, K., Kondo, M., Nishimura, M., Yamato, K.

T., Kohchi, T. and Nakagawa, T. (2018). Novel gateway binary vectors for rapid

tripartite DNA assembly and promoter analysis with various reporters and tags in the

liverwort *Marchantia polymorpha. PLoS One* **13**, e0204964,

doi:10.1371/journal.pone.0204964.

Meier, I., Griffis, A. H. N., Groves, N. R. and Wagner, A. (2016). Regulation of nuclear

shape and size in plants. *Curr. Opin. Cell Biol.* **40**, 114–123,

doi:10.1016/j.ceb.2016.03.005.

Meistrich, M. L. and Hess, R. A. (2013). Assessment of spermatogenesis through staging of

seminiferous tubules. *Methods Mol. Biol.* **927**, 299–307, doi:10.1007/978-1-62703-038-

0_27.

Minamino, N., Kanazawa, T., Nishihama, R., Yamato, K. T., Ishizaki, K., Kohchi, T.,

Nakano, A. and Ueda, T. (2017). Dynamic reorganization of the endomembrane

system during spermatogenesis in *Marchantia polymorpha. J. Plant Res.* **130**, 433–441,

doi:10.1007/s10265-017-0909-5.

Minamino, N., Kanazawa, T., Era, A., Ebine, K., Nakano, A. and Ueda, T. (2018). RAB

GTPases in the basal land plant *Marchantia polymorpha. Plant Cell Physiol.* **59**, 845–

856, doi:10.1093/pcp/pcy027.

Montgomery, S. A., Tanizawa, Y., Galik, B., Wang, N., Ito, T., Mochizuki, T.,

Akimcheva, S., Bowman, J. L., Cognat, V., Maréchal-Drouard, L., et al. (2020).

Chromatin organization in early land plants reveals an ancestral association between

H3K27me3, transposons, and constitutive heterochromatin. *Curr. Biol.* **30**, 573-588.e7,

doi:10.1016/j.cub.2019.12.015.

O'Hagan, R. and Barr, M. M. (2012). Regulation of tubulin glutamylation plays cell-

specific roles in the function and stability of sensory cilia. *Worm* **1**, 155–159,

doi:10.4161/worm.19539.

O'Hagan, R., Piasecki, B. P., Silva, M., Phirke, P., Nguyen, K. C. Q., Hall, D. H.,

Swoboda, P. and Barr, M. M. (2011). The tubulin deglutamylase CCPP-1 regulates the

function and stability of sensory cilia in *C. elegans*. *Curr. Biol.* **21**, 1685–1694,

doi:10.1016/j.cub.2011.08.049.

O'Rahilly, R. and Müller, F. (1987). *Developmental Stages in Human Embryos, Including a*

Revision of Streeter's 'Horizons' and a Survey of the Carnegie Collection. Washington,

Carnegie Institution of Washington.

Oakberg, E. F. (1956). A description of spermiogenesis in the mouse and its use in analysis

of the cycle of the seminiferous epithelium and germ cell renewal. *Am. J. Anat.* **99**, 391–

413, doi:10.1002/aja.1000990303.

Renzaglia, K. S. and Duckett, J. G. (1987). Spermatogenesis in *Blasia pusilla*: from young

antheridium through mature spermatozoid. *Bryologist* **90**, 419, doi:10.2307/3243109.

Renzaglia, K. S. and Garbary, D. J. (2001). Motile gametes of land plants: diversity,

development, and evolution. *CRC. Crit. Rev. Plant Sci.* **20**, 107–213,

doi:10.1080/20013591099209.

Rojo, E., Gillmor, C. S., Kovaleva, V., Somerville, C. R. and Raikhel, N. V. (2001).

VACUOLELESS1 is an essential gene required for vacuole formation and

morphogenesis in *Arabidopsis*. *Dev. Cell* **1**, 303–310, doi:10.1016/S1534-

5807(01)00024-7.

Russell, S. D. and Jones, D. S. (2015). The male germline of angiosperms: repertoire of an

inconspicuous but important cell lineage. *Front. Plant Sci.* **6**, 1–10,

doi:10.3389/fpls.2015.00173.

Sanchez-Vera, V., Kenchappa, C. S., Landberg, K., Bressendorff, S., Schwarzbach, S.,

Martin, T., Mundy, J., Petersen, M., Thelander, M. and Sundberg, E. (2017).

Autophagy is required for gamete differentiation in the moss *Physcomitrella patens*.

Autophagy **13**, 1939–1951, doi:10.1080/15548627.2017.1366406.

Sanders, P. M., Bui, A. Q., Weterings, K., McIntire, K. N., Hsu, Y. C., Lee, P. Y.,

Truong, M. T., Beals, T. P. and Goldberg, R. B. (1999). Anther developmental defects

in *Arabidopsis thaliana* male-sterile mutants. *Sex. Plant Reprod.* **11**, 297–322,

doi:10.1007/s004970050158.

Sassone-Corsi, P. (2002). Unique chromatin remodeling and transcriptional regulation in
spermatogenesis. *Science (80-.).* **296**, 2176–2178, doi:10.1126/science.1070963.

Sengupta, D. and Linstedt, A. D. (2011). Control of organelle size: the Golgi complex.
Annu. Rev. Cell Dev. Biol. **27**, 57–77, doi:10.1146/annurev-cellbio-100109-104003.

Shimamura, M. (2015). Whole-mount immunofluorescence staining of plant cells and
tissues. In Plant Microtechniques and Protocols (ed. Yeung, E.), pp181-196.
Switzerland: Springer.

Shimamura, M. (2016). *Marchantia polymorpha*: taxonomy, phylogeny and morphology of
a model system. *Plant Cell Physiol.* **57**, 230–256, doi:10.1093/pcp/pcv192.

Takemoto, K., Ebine, K., Askani, J. C., Krüger, F., Gonzalez, Z. A., Ito, E., Goh, T.,
Schumacher, K., Nakano, A. and Ueda, T. (2018). Distinct sets of tethering
complexes, SNARE complexes, and Rab GTPases mediate membrane fusion at the
vacuole in Arabidopsis. *Proc. Natl. Acad. Sci. USA.* **115**, 2457–2466,
doi:10.1073/pnas.1717839115.

Twell, D. (2011). Male gametogenesis and germline specification in flowering plants. *Sex.*
Plant Reprod. **24**, 149–160, doi:10.1007/s00497-010-0157-5.

Uemura, T., Kim, H., Saito, C., Ebine, K., Ueda, T., Schulze-Lefert, P. and Nakano, A.
(2012). Qa-SNAREs localized to the *trans*-Golgi network regulate multiple transport
pathways and extracellular disease resistance in plants. *Proc. Natl. Acad. Sci. USA.* **109**,
1784–1789, doi:10.1073/pnas.1115146109.

Vu, H. T., Akatsu, H., Hashizume, Y., Setou, M. and Ikegami, K. (2016). Increase in α-
tubulin modifications in the neuronal processes of hippocampal neurons in both kainic
acid-induced epileptic seizure and Alzheimer's disease. *Sci. Rep.* **7**, 40205,
doi:10.1038/srep40205.

Wloga, D., Joachimiak, E., Louka, P. and Gaertig, J. (2017). Posttranslational
modifications of Tubulin and cilia. *Cold Spring Harb. Perspect. Biol.* **9**,
doi:10.1101/cshperspect.a028159.

 Figure legends

Figure 1. Developmental stages of spermiogenesis in *M. polymorpha*

(A-B) Schematic illustration of spermatozoids (A) and male gamete development (B). (C-H)

Maximum intensity projection images of spermatids and a spermatozoid stained with Hoechst

33342. Left panels, merged images of differential interference contrast (DIC) and fluorescent

images; middle panels, higher magnification images of left panels; right panels, fluorescent

images of nuclei in middle panels. The blue pseudo-color indicates fluorescence from

Hoechst 33342. Scale bars = 5 μm.

Figure 2. Microtubule dynamics during spermatogenesis

(A-G) Maximum intensity projection images of spermatids and a spermatozoid

immunostained with anti-α-tubulin and anti-polyE antibodies. Left panels, differential

interference contrast (DIC) images; middle panels, fluorescent images; right panels, higher

magnification images of middle panels. Green, magenta, and blue pseudo-colors indicate

fluorescence from Alexa 488 (α-tubulin), Alexa594 (polyE), and Hoechst 33342, respectively.

Scale bars = 5 μm.

Figure 3. Changes in vacuolar morphology during spermiogenesis

(A-F) Maximum intensity projection images of spermatids and a spermatozoid expressing

mCitrine-MpVAMP71 driven by the Mp*DUO1* promoter. Left panels, fluorescent images;

right panels, differential interference contrast (DIC) images. Green and blue pseudo-colors

indicate fluorescence from mCitrine and Hoechst 33342, respectively. Scale bars = 5 μm.

Figure 4. Remodeling of the plasma membrane during spermiogenesis

(A-H) Maximum intensity projection images of spermatids and a spermatozoid expressing

mCitrine-MpSYP12A driven by the Mp*DUO1* promoter. Left panels, fluorescence images;

right panels, differential interference contrast (DIC) images. Green and blue pseudo-colors

indicate fluorescence from mCitrine and Hoechst 33342, respectively. Scale bars = 5 μm.

Figure 5. Reorganization of the Golgi apparatus during spermiogenesis

(A-F) Maximum intensity projection images of spermatids and a spermatozoid expressing

ST-Venus driven by the Mp*DUO1* promoter. Left panels, fluorescence images; right panels,

differential interference contrast (DIC) images. Green and blue pseudo-colors indicate

fluorescence from mCitrine and Hoechst 33342, respectively. Scale bars = 5 μm. (G) The size

of the Golgi apparatus labeled by ST-Venus. n = 137 (stage 0), 91 (stage 1), and 45 puncta

(stage 2). The boxes and solid lines in the boxes indicate the first quartile and third quartile,

and median values, respectively. The whiskers indicate $1.5 \times$ interquartile ranges. Different

letters denote significant differences based on Tukey's test ($p < 0.05$). (H) The number of the

Golgi apparatus labeled by ST-Venus. The difference in the number of the Golgi apparatus

among stages was statistically significant based on Tukey's test ($p < 0.05$). See also Fig. S5.

Figure 6. Reorganization of the ER and the nuclear envelope during spermiogenesis

Maximum intensity projection images of spermatids and spermatozoids expressing mCitrine-

MpSEC22 driven by the Mp*DUO1* promoter (A, C, E, G, I, K) or SP-mCitrine-HDEL driven

by the Mp*EF1α* promoter (B, D, F, H, J, L). Arrowheads indicate SP-mCitrine-HDEL

remaining in spermatozoids. Left panels, fluorescence images of mCitrine; middle panel,

merged images of mCitrine and Hoechst 33342; right panels, differential interference contrast

(DIC) images. Green and blue pseudo-colors indicate fluorescence from mCitrine and

Hoechst 33342, respectively. Scale bars = 5 μm.

Figure 7. Schematic diagram of spermiogenesis progression in *M. polymorpha*.

At stage 0, flagellar formation begins in a cell body, and endocytosis is highly activated. At

stage 1, flagella protrude and elongate. Radially extending microtubules are formed

transiently from or to the basal region of the flagella. Plasma membrane (PM) proteins

including mCitrine-MpSYP12A are transported to vacuoles. At stage 2, flagella elongation is

mostly completed, and morphological change of the nucleus begins. Spline microtubules are

visible. Vacuoles fuse with each other to form a large spherical vacuole. PM proteins are

degraded in the spherical vacuoles. One large Golgi apparatus is observed. At stage 3, Golgi

proteins are degraded. The nuclear envelope is remodeled. At stage 4, most of endoplasmic

reticulum (ER) proteins are eliminated. The spherical vacuole persists until stage 4, which is

removed to transform to the mature spermatozoid. Glutamylation of tubulin in axonemal and

spline microtubules is dynamically regulated during spermiogenesis.