Application of stable isotopes to study movement ecology and diet

variation in a migratory songbird

Andrea Contina[1, 2,†] Allison K. Pierce[1,*], Scott W. Yanco[1,3,4,*], Eli S. Bridge[5], Jeffrey F. Kelly[5,6,7],

Michael B. Wunder[1]

[1] University of Colorado Denver, Department of Integrative Biology, Denver, CO, USA.

[2] Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA.

[3] Center for Biodiversity and Global Change, Yale University, New Haven, CT, USA.

[4] Department of Ecology and Evolutionary Biology, Yale University, CT, USA.

[5] Oklahoma Biological Survey, University of Oklahoma, Norman, OK, USA.
10
[6] Corix Plains Institute and Department of Biology, University of Oklahoma, Norman, OK, USA.

[7] Department of Biology, University of Oklahoma, Norman, OK, USA.

* Equal contribution

† Corresponding author's email: acontina@utexas.edu

Current author's address: Department of Integrative Biology, University of Texas at Austin,

Austin, TX, USA.

Authors' contributions

AC conceptualized the project, analyzed the data, and wrote the manuscript. AKP analyzed the

data, reviewed, and edited the manuscript. SWY provided key feedback on data analysis,

reviewed, and edited the manuscript. ESB collected the data and reviewed the manuscript. JFK

collected the data and reviewed the manuscript. MBW supervised data analysis and reviewed the

manuscript.

Introduction

Animal migration is a multifaceted life-history trait that evolved across taxa through complex

behavioral and physiological adaptations (Alerstam and Bäckman, 2018). Migratory species

maintain energetic balance between fuel consumption (e.g., fatty acid catabolism) and

replenishing resources to effectively perform a long-distance migration (Guglielmo, 2018).

However, research into how migrants sustain themselves during migration remains particularly

challenging due to the small body size of many migratory songbirds and the fact that they travel,

in some cases, thousands of kilometers per day (Rani et al., 2017; DeLuca et al., 2015). Species

have developed unique behavioral strategies associated with seasonal movements (Horton et al.,

2016), which depend on resource availability and seasonal environmental stochasticity. Thus, an

understanding of these strategies requires examination of fluctuations in body mass and fat

deposition in conjunction with behavior across the spatial extent of the species' distribution. This

is relevant because variation in migration and foraging behavior at the intra-species level can

place different evolutionary pressures on individual physiology and act as selective forces across

populations. That is, there might be a trade-off between longer and more energetically costly

migratory routes, potentially resulting in decreased body condition, and alternative strategies

involving diet changes or morphological adaptations (e.g., longer or pointed wings) to

accommodate the increased energetic demands of their migratory journey (Kaboli et al, 2007;

Reif et al., 2016).

Many species of songbirds are too small to be fitted with tracking devices (e.g., satellite

transmitters) and information on their migratory patterns remain unknown. However, indirect

molecular approaches can be implemented to study large-scale movements without the use of

extrinsic tracking devices. (Rundel et al., 2013). In particular, movement ecology studies have

exploited the latitudinal gradient in stable hydrogen isotopes occurring across North America to

address questions related to avian migratory patterns and connectivity between breeding and

wintering grounds (Rubenstein and Hobson, 2004; Vander Zanden et al. 2018; Besozzi et al.

2021). Stable isotope ratios of hydrogen obtained from inert animal tissues (e.g., claws, hair, and

feathers), provide insight on the environmental conditions and latitudinal gradient where the

keratin-based tissues were grown and offer a valid molecular tool to infer large-scale animal

movements in species too small for direct movement observations (Hobson, 2005). Importantly,

most songbirds replace their flight feathers at the breeding ground and before starting fall

migration to minimize the energetic costs of the journey (Barta et al., 2008). By replacing old

wing feathers with new ones, presumably in better conditions, migratory birds gain aerodynamic

efficiency (Bowlin and Wikelski, 2008). Thus, feather samples collected at the wintering

grounds, sometimes thousands of kilometers away from the breeding locations, can be used to

infer where the feathers were grown based on stable isotope similarities between environmental

and tissue hydrogen values (Wunder, 2010).

A suitable study system for disentangling different migratory strategies from other key

evolutionary adaptations is offered by the Dark-eyed Junco (*Junco hyemalis*), a migratory

songbird comprising numerous subspecies broadly distributed across North America. It is found

breeding and molting in the northern regions of Canada and Alaska, and mountainous regions of

the United States (U.S.) and wintering in the lower latitudes of Canada and the majority of the

continental U.S. and northern Mexico (Nolan et al., 2002). This species has shown negative

population trends of over 40% in some areas in the last decade (PIF 2017; Rockwell et al., 2019)

and its physiology and migratory behaviors have been the focus of numerous studies (Nolan and

Ketterson, 1983; Rogers et al., 1994.; Fudickar et al., 2016; Liebgold et al., 2019).

Bridge et al. (2010) analyzed hydrogen isotope ratios from secondary feathers in a

population of Dark-eyed Juncos, all from the same *hyemalis* subspecies group (slate-colored),

sampled during the winter of 2009 in central Oklahoma (U.S.) and showed isotopic variation

ranging from -175.6‰ to -118.6‰ from Vienna Standard Mean Ocean Water (VSMOW). This

large range of variation (~60‰) in hydrogen stable isotope values prompted the authors to

speculate that individuals from a wide geographic range of breeding, and consequently molting,

locations wintered in Oklahoma, although the study did not include explicit probabilistic

geographic assignments of wintering individuals, nor did it consider their variation in trophic

level.

Although the subspecies of Dark-eyed Junco's are primarily differentiated by

morphology and/or geographic distribution, considerable behavioral diversity between and even

within subspecies presents a unique opportunity to address large-scale spatial and ecological

questions. We used hydrogen stable isotope data to estimate breeding location origins of a

population of Dark-eyed Juncos wintering in Oklahoma, and we collected a suite of

morphological measurements after the birds completed fall migration. Moreover, we expect that

Oklahoma wintering birds originating from northern latitudes would have a higher carbohydrate

intake, but lower protein consumption compared to wintering birds from southern latitudes due

to their need to fuel a longer migratory journey. To test this hypothesis, we assessed pre-

migratory diets by examining carbon and nitrogen stable isotope ratios in flight feathers. In

addition to quantifying geographic connectivity in the Dark-eyed Junco population wintering in

Oklahoma, this study offers new insights into how this species negotiates tradeoffs between

migration distance and diet. Importantly, we present a framework to identify possible

confounding effects in explanatory models for different migration strategies across populations

(Yanco et al. 2020).

Methods

Study Area

We studied Dark-eyed Juncos wintering in central Oklahoma where the species forages in

open woodland habitats and feeds on grass sprouts, *Taraxacum* sp. and various species of insects

(Bridge et al. 2010). The weather conditions during our study (January and February 2009) as

recorded by the Oklahoma Climatological Survey and the Oklahoma Mesonet

(www.mesonet.org) at the Norman (NRMN) weather station indicated monthly average

temperature ranging from 2.7 to 8.9 °C, monthly average precipitation ranging from 22.8 to 24.6

mm, and monthly average wind speed ranging from 15.6 to 18.1 kph.

Sample collection and stable isotope analysis

We used published stable hydrogen isotope data from 80 Dark-eyed Juncos sampled in

central Oklahoma, USA (35°10' N, 97°26' W) during winter 2009 (Bridge et al. 2010). After

removing individuals with missing capture information, our data set comprised stable hydrogen

isotope values of secondary feathers from 76 migrants that we used to infer geographic breeding

and molting origin. We also present novel carbon and nitrogen stable isotope data obtained from

a subset (N = 74) of those migrants sampled in Oklahoma that we used to study habitat use and

trophic level. However, after filtering out individuals with missing morphometric measurements

we had data from 62 migrants (39 adult males and 23 adult females) that we used for

morphological models. All samples were collected between January and February of the same

year. Feathers were prepared for analysis of stable isotope ratios of hydrogen (δ^2H), carbon

($\delta^{13}C$), and nitrogen ($\delta^{15}N$) by following standard washing protocol in a 2:1 solution of

chloroform-methanol to remove debris and oil contaminants (Paritte and Kelly 2009; Chew et al.

2019). Samples for hydrogen stable isotope ratios analysis were processed through a Thermo-

Finnigan Delta V isotope ratio mass spectrometer interfaced with a high-temperature pyrolysis

elemental analyzer (TC/EA, Thermo-Finnigan, Bremen, Germany). Feather δ^2H values (δ^2H$_f$) are

reported as mean ± SD in delta notation of parts per mil (‰) from the standards ($\delta D_{sample} =$

$[(R_{sample}/R_{standard}) - 1]$) compared to the Vienna Standard Mean Ocean Water (VSMOW). We

used chicken feather standard (CFS; -147.4‰), cow hooves (CHS; -187‰), and bowhead whale

baleen (BWB; -108‰) as keratin which are widely adopted for sample comparative equilibration

(Wassenaar and Hobson, 2003). The details of hydrogen sample measurement are in Kelly et al.

2009. We conducted carbon and nitrogen stable isotope ratios analysis on a Thermo-Finnigan

DeltaPlus isotope ratio mass spectrometer connected to a Carlo Erba elemental analyzer. We

normalized the δ^{13}C and δ^{15}N values to the Vienna Pee Dee Belemnite (VPDB) and AIR scales,

respectively, and used Brown-headed cowbird (BHCO; *Molothrus ater*) feather standards. The

values of δ^{13}C and δ^{15}N standards were -15.7 ± 0.1‰ (δ^{13}C BHCO), and 7.6 ± 0.1‰ (δ^{15}N

BHCO).

Feather Calibration and Geographical Assignments

We implemented a Bayesian analytical framework in the R package assignR to spatially

reconstruct juncos' breeding and/or molting areas (Ma et al., 2020; R Core Team, 2020). First,

we rescaled stable hydrogen isotope values in precipitation (δ^2H$_p$) from a North America

geostatistical model (Bowen et al. 2005) to 22 known-origin Song Sparrow (*Melospiza melodia*)

samples provided by Hobson et al. (2012) and Hobson and Koehler (2015). To ensure a more

accurate isotopic calibration (Wunder 2010), we used linear regression to model mean tissue

isotope values as a function of δ^2H$_p$ using the *calRaster* function in assignR. We recognize that a

species-specific calibration provides higher degree of accuracy. However, when species-specific

known-origin tissues are not available, calibration data from closely related species can be used

(Hobson et al., 2012). Our data set comprised entirely unknown-origin feather samples from

Dark-eyed Juncos; therefore, we used published data of *M. melodia*, a species of comparable

size, diet, and spatial distribution (Arcese et al., 2002; Hobson et al., 2012).

We determined the molting locations of juncos wintering in Oklahoma by computing

posterior probability density maps through the *pdRaster* function in assignR (Wunder 2010; Ma

et al., 2020). We present normalized posterior probability maps representing both the individual

and cumulative group probabilities of origin from each grid cell restricted to the breeding and

year-round distribution ranges (BirdLife International 2016). To visualize the cumulative group

probabilities of origin, we clustered the samples into four isotopic groups based on the reported

stable isotope feather values (δ^2H_f) across the junco data set arranged from the lowest values

(G1) to the highest values (G4): the first group (G1) included δ^2H_f values ranging from -175.6‰

to -160.2‰ (M = -167.1‰, SD = 5.02‰, N = 18); the second group (G2) included values

ranging from -159.3‰ to -150.4‰ (M = -154.9‰, SD = 3.17‰, N = 23); the third group (G3)

included values ranging from -149.8‰ to -140.1‰ (M = -145.4‰, SD = 2.88‰, N = 19); and

the fourth group (G4) included values ranging from -139.3‰ to -118.6‰ (M = -130.4‰, SD =

6.39‰, N = 16). Even though this clustering approach is *ad hoc*, it still represents a useful

subdivision based on δ^2H_f variation across birds in our dataset. Wassenaar and Hobson (2006)

estimated that ±3‰ variation in hydrogen stable isotope extracted from feathers translates

approximately to 1 degree of latitude (or ~111 km). Therefore, the differential range of about

10‰ or more that we used to cluster birds into four groups corresponds to ~370 km of

geographic separation (Hobson and Wassenaar, 1996; Wassenaar and Hobson, 2006). This

distance is presumably large enough to consider our samples from different breeding

populations.

Modeling morphometric parameters

We investigated the effect and relative predictive importance of five different

morphometrics on mean hydrogen stable isotope ratios in Dark-eyed Junco feathers. Our

explanatory variables were: wing chord (mm), tarsus length (mm), tail length (mm), mass (g),

and fat depositions scored on a six-point ordinal scale (Helms and Drury 1960). We did not

include body mass index normalized by a structural measure because this would be collinear

with other included variables and additive model combinations should account for size

normalized mass (Green, 2001). We controlled for morphological variance across males and

females by including 'sex' as a random effect in all our models and to standardize evaluation of

predictive influence between the morphometric variables measured on different scales we Z-

scored values for all variables. We modeled a complete combinatorial set of linear models in the

R package MuMIn (Barton and Barton, 2019) and computed the relative importance of

individual morphometric variables through a variable importance analysis using Akaike

Information Criterion corrected for small sample sizes (AICc) and summed model weights for

each model containing a given variable. We then compared the sum of all the AICc weights and

odds ratios to evaluate the relative importance of each variable.

Trophic Level Analysis

We analyzed stable carbon ($\delta^{13}C$) and nitrogen ($\delta^{15}N$) isotopes in feather samples to

explore trophic level partition among birds migrating from northern and southern latitudes as

revealed by hydrogen (δ^2H) data. We exploited the occurrence of distinct plant photosynthetic

pathways, knowing that trees and shrubs in temperate climates (e.g., C3 plants) are depleted in

carbon compared to plants inhabiting arid environments and adopting C4 or CAM

photosynthesis (Keeley and Rundel, 2003). These differences help to assess the proportion of

proteins deriving from C3 and C4 plants, while higher nitrogen values indicate a shift towards a

diet less dependent on seeds and richer in insects. Thus, we used the *corr.test* function in the R

package psych (Revelle and Revelle, 2015) to perform a Pearson correlation test between carbon

(δ^{13}C) and hydrogen (δ^2H) and between nitrogen (δ^{15}N) and hydrogen (δ^2H) stable isotopes. This

test highlights diet trends in relation to latitudinal origin as determined by δ^2H data. We then

used the *levelplot* function in the R package lattice (Sarkar 2008) to create a visual representation

of δ^2H values from feathers collected in Oklahoma and their distribution within the sampling

space.

Results

Rescaling model and posterior probability density maps

The rescaling model provided a robust relationship between precipitation isotope values

($\delta^2 H_p$) and known-origin samples of the species *M. melodia* (y = -35.08 + 0.91x; R^2 = 0.79, $CI_{95\%}$

[0.74, 0.95], n = 22) and allowed us to calibrate our isoscape to determine junco's posterior

probability of origin (Fig 1 – panel A). We obtained a rescaled environmental isoscape raster of

52,884 cells at 0.33km resolution and a range of hydrogen stable isotope values spanning -

193.3‰ and -41.3‰ (M = -122.9‰; SD = 37.2‰; Fig 1 – panel B and C).

Once we performed isoscape tissue calibration, we inferred the spatial extent of the

breeding and molting origin of 76 Dark-eyed Juncos wintering in Oklahoma through a Bayesian

probabilistic framework. Our posterior probability density maps, show distinct latitudinal

variation across the breeding and molting range of migratory juncos (Fig 2). Overall, the

northernmost migrants were assigned to either Alaska and western Canada, including Yukon

Territory and British Columbia, or central Northwestern Territories, while southern migrants

were assigned to either Northern Saskatchewan or southern Manitoba and northwestern United

States near the Rocky Mountains. However, even though the probability assignment surfaces for

the southern migrants (group G3 and G4) were largely distributed at latitudes lower than 60°N

(Fig 2), we also found weak probabilities of origin along the western coast of Alaska due to the

underlying structure of the precipitation isoscape used for rescaling (Fig 1, panel B-C). Taken

together, these results suggest that different Dark-eyed Junco populations admix into the same

wintering area in central Oklahoma.

249
250

Variable importance analysis

The analysis of the importance of variables revealed no clear predictive support for any

variable considered in our models. We fit a full set of 32 linear mixed-effect models for mean

δ^2H_f including single variable models and all the possible combinations of the five predictors,

with a random effect of sex and ranked them using Akaike Information Criterion corrected for

small sample size (AICc). All 5 variables we considered had relatively high summed AICc

evidence weights suggesting models containing each respective variable outperformed those that

did not (Table 1 and Table 2). Fat score had the highest summed AICc weight value and odds

ratios, indicating that models containing the fat score variable accounted for all of the weight of

AICc values for the model set (Table 1). However, the estimated effect of fat score on mean δ^2H_f

was indistinguishable from 0 in all models weighted by AICc (Fig 3A) suggesting that despite

being the most comparatively informative variable, it had no discernible effect on δ^2H_f values in

our data set. Similarly, estimates for other variables we considered also lacked support for non-

zero effects Fig 3B).

Migrant Diet Preferences and Trophic Level

The Pearson test did not identify a strong correlation between carbon ($\delta^{13}C$) and

hydrogen (δ^2H) isotopes values ($r = -0.17$, $p = 0.22$), but it suggested a moderately positive

correlation between nitrogen ($\delta^{15}N$) and hydrogen (δ^2H) isotopes values ($r = 0.33$, $p = 0.01$).

These results suggest that a trophic level shift towards insect consumption might occur in

migrants originating from lower latitudes (Fig 4 panel A and B). Interestingly, when samples

were grouped by wing length (Fig 4 panel C and D), it appeared that birds with longer wings had

a broader range of $\delta^{13}C$ values, indicating variability in their habitat use. Moreover, when

plotting $\delta^{13}C$ against $\delta^{15}N$ over δ^2H feather isoscape within the sampling space, the changes in

dietary niche across latitudinal origin appeared more evident (Fig 5 panel A-C; supplementary

material; S1-S2). Northern migrants showed lower protein consumption (δ^{15}N: 1-4‰) and C3-

shrub habitat use (δ^{13}C: -22 − -24‰), while southern migrants were characterized by a higher

trophic level (δ^{15}N: 4-8‰) and δ^{13}C towards C3-vegetation compatible with temperate forests

(δ^{13}C: -24 − -26‰). Analysis of diet preferences between sexes did not show marked differences,

although males seem to have a slightly larger dietary niche compared to females.

Discussion

Bridge et al. (2010) conducted a captive experiment in which wintering Dark-eyed

Juncos were fed *ad libitum* to demonstrate that fat depositions allow them to initiate migration

earlier in the spring compared to the control group. We extended previous work by providing a

geostatistical approach illustrating the explicit spatial distribution of breeding and molting origin

of fall migrants and related that distribution to morphological characteristics and trophic level

differentiation. Our geographical assignments clearly suggested that migratory juncos wintering

in central Oklahoma originated from different breeding and molting areas (Fig 2). We identified

northern and southern migrants while each group also showed intra-cohort spatial variation

ranging from Alaska and western Canada to central Northwestern Territories, and Northern

Saskatchewan to southern Manitoba and northwestern United States, respectively. However,

despite the posterior probability surfaces showing population admixture at the wintering area in

Oklahoma, our modeling effort showed no association between winter body condition and

breeding or molting latitude.

Stable isotope ratios of hydrogen, carbon, and nitrogen, extracted from inert animal

tissues, such as claws and feathers, reflect the isotope ratios in the environment where those

tissues were grown and can provide inferences about large-scale animal movements and diet

preferences in species with small body size (Rubenstein and Hobson, 2004; Zenzal et al., 2018;

Vander Zanden et al., 2018). However, a clear advantage of adopting a multi-element stable

isotope approach is that migrants can be assigned to different breeding and molting grounds of

origin while their diet preferences are also revealed and clustered in trophic guilds (Herrera et al.,

2003). The positive correlation between nitrogen ($\delta^{15}N$) and hydrogen (δ^2H) values, although

moderate, corroborated our hypothesis that wintering birds originating from northern latitudes

would have lower protein intake relative to southern migrants and prefer a diet rich in

carbohydrates before migrating. That is, increased carbohydrate anabolism and lower protein

consumption is necessary to quickly synthesize enough subcutaneous lipid reserves to sustain a

long-distance migration (Cecere, et al., 2011; Domer et al., 2018). Alternatively, higher δ^{15}N

values in southern migrants might be explained by environmental stressors. An early study in

captive quails (*Coturnix japonica*) and geese (*Anser rossii*) showed that increased δ^{15}N values

can be obtained from growing tissues of individuals maintained under severe nutritional stress

conditions (e.g., fasting) compared to control groups (Hobson et al. 1993). However, subsequent

analysis in sparrows (*Melospiza melodia*) found no differences in δ^{15}N between birds kept under

a restricted diet versus birds fed *ad libitum* (Kempster et al., 2007) suggesting a less

straightforward relationship between physiological stress conditions and δ^{15}N values. Therefore,

even though we cannot pinpoint the exact causes of increased δ^{15}N values in growing feathers

(e.g., baseline difference) and that other species have shown variation in feather δ^{15}N values

between boreal and more southern food webs based on land use (Hobson 1999), it is also

possible that migratory juncos originating from southern latitudes were foraging at a higher level

in the trophic system and consuming more insects compared to northern migrants. We do not

exclude that such diet preference might be linked to an opportunistic foraging strategy due to an

outburst of insect availability (Yang et al., 2008). However, analysis of isotopic niche shifts in

avian migratory populations have shown sharp changes during different life cycle stages (Hahn,

et al., 2013), confirming that variations in diet compositions have a defined role in maintaining

fluctuating energy needs (Marshall et al., 2016). Moreover, our δ^{13}C results indicated that

northern juncos might exploit food resources from a broader range of habitats compared to

southern juncos, ranging from pine forest to deciduous forest (e.g., oak) and C3 shrubs. Bearhop

et al. (2004) studied the Thick-billed Vireos (*Vireo crassirostris*) in the Bahamas and based on

mean claw $\delta^{13}C$ values found differences among two groups of birds occupying environments

with either prevalent xerophytic shrubs or pine forest. While our feather $\delta^{13}C$ results showed that

northern juncos seed consumption fell within the spectrum of plants characterized by a C3

photosynthetic pathway, and we found enough variation to highlight habitat occupancy

differentiations across latitudinal origins (Fig 4 and Fig 5), we note that a robust environmental

isotopic baseline is necessary to determine dietary and habitat preferences with confidence (Post,

2002).

The interactions between avian body mass fluctuations and migratory behaviors within

and between seasons remain poorly understood (Brown and Sherry, 2006; Tsvey et al., 2007;

Marra et al. 2015). Importantly, body condition appears to both influence and be influenced by

migratory behavior. For example, body condition may affect migratory decisions (Schaub and

Jenni 2000); several species of migrating shorebirds underwent longer migratory flights when

departing with higher body condition scores (Anderson et al. 2019). Similarly, spring arrival

timing of red knots (*Calidris canutus*) was associated with better body condition at a major

stopover site (Duijns et al. 2017) and conditions experienced during winter months by American

Redstarts (*Setophaga ruticilla*) lead to variation in body conditions that predicted spring

departure dates (Marra et al., 1998). In our investigation, individuals that migrated minimum

distances did non exhibit larger body masses or higher fat scores over the early winter period.

However, our study was limited because individual migrants were not captured immediately

upon arrival in Oklahoma. While challenging, an earlier capturing effort might reveal mass as a

predictor of latitude of origin because the body condition of those migrants would provide a

stronger indication of the migration strategy performed. Moreover, there may be other

mechanisms that could explain the absence of a relationship between body conditions and

migration strategies in our study population. For example, the taxonomy of the genus *Junco* has

been challenging for decades and is far from being resolved (Miller 1941; Nolan et al., 2002;

Aleixandre et al., 2013). The species *Junco hyemalis* has been recently proposed to be

phylogenetically reassessed based on extensive genomic data to determine if it needs to be split

into four new species that also account for distinct phenotypic variation (Friis et al., 2016). Thus,

in a complex migratory system, it is possible that in some populations different morphs of

migrants characterized by a larger and smaller than average body size could come into contact at

the winter ground and therefore confounding the effects of energy consumption during

migration. Alternatively, long distance migrants could also improve their body conditions (e.g.,

increase mass at stopover sites) before arriving at their final wintering destination and reduce the

difference in body mass compared to short distance migrants. However, if the stopover site is

only used to refuel without growing new feathers, the stable isotope analysis will still detect the

northern origin (e.g., molting ground) and correctly identify the migration latitudinal origin

without revealing any geographical information on the stopover location.

383
384   **Conclusions**

385   By implementing a Bayesian assignment framework, we offered a probabilistic

geographic explanation of the junco's feather isotopic variation across the species' breeding and

molting areas in the northern hemisphere and attempted to explain latitudinal origin in relation to

morphological traits (Wunder, 2010; Alerstam, 2011; Maggini et al., 2016). In our system, none

of the variables considered had a clear effect on feather hydrogen stable isotope values (used as a

proxy for latitudinal origin) although we found indication of a possible higher trophic level shift

in individuals migrating from southern latitudes. These findings promote the concurrent use of

different stable isotope markers as a means of studying interconnected behavioral traits. Our

results also highlight the need to carefully examine confounding effects in explanatory models in

relation to the species' natural history and possible differential migratory behaviors across

populations.

Abbreviations

NRMN: Norman; AICc: Akaike Information Criterion corrected for small sample sizes; G1:

Group 1; G2: Group 2; G3: Group 3; G4: Group 4.

References

Alerstam, T., 2011. Optimal bird migration revisited. *Journal of Ornithology*, *152*(1), pp.5-23.

Alerstam, T. and Bäckman, J., 2018. Ecology of animal migration. *Current Biology*, *28*(17), pp.R968-R972.

Anderson, A. M., S. Duijns, P. A. Smith, C. Friis, and E. Nol. 2019. Migration Distance and Body Condition Influence Shorebird Migration Strategies and Stopover Decisions During Southbound Migration. Frontiers Ecol Evol 7:251.

Arcese, Peter, Mark K. Sogge, Amy B. Marr and Michael A. Patten. (2002). Song Sparrow (*Melospiza melodia*), version 2.0. In The Birds of North America (P. G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, New York, USA.

Barta, Z., McNamara, J.M., Houston, A.I., Weber, T.P., Hedenström, A. and Fero, O., 2008. Optimal moult strategies in migratory birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1490), pp.211-229.

Barton, K. and Barton, M.K., 2019. Package 'MuMIn'. *Multi-model inference. version*, *1*(6).

Bearhop, S., Hilton, G.M., Votier, S.C. and Waldron, S., 2004. Stable isotope ratios indicate that body condition in migrating passerines is influenced by winter habitat. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *271*(suppl_4), pp.S215-S218.

Besozzi, E., Chew, B., Allen, D.C. and Contina, A., 2021. Stable isotope analysis of an aberrant Painted Bunting (*Passerina ciris*) feather suggests post-molt movements. *The Wilson Journal of Ornithology*, *133*(1), pp.151-158.

BirdLife International and Handbook of the Birds of the World (2016) Bird species distribution maps of the world". Version 6.0. Available at http://datazone.birdlife.org/species/requestdis.

Bowlin, M.S. and Wikelski, M., 2008. Pointed wings, low wingloading and calm air reduce migratory flight costs in songbirds. *PloS one*, *3*(5).

Bridge, E.S., Kelly, J.F., Bjornen, P.E., Curry, C.M., Crawford, P.H. and Paritte, J.M., 2010. Effects of nutritional condition on spring migration: do migrants use resource availability to keep pace with a changing world? *Journal of Experimental Biology*, *213*(14), pp.2424-2429.

Brown, D.R. and Sherry, T.W., 2006. Food supply controls the body condition of a migrant bird wintering in the tropics. *Oecologia*, *149*(1), pp.22-32.

457   Cecere, J.G., Spina, F., Jenni-Eiermann, S. and Boitani, L., 2011. Nectar: an energy drink used
458   by European songbirds during spring migration. *Journal of Ornithology*, *152*(4), pp.923-931.
459

460   Chew B, Kelly JF, Contina A. 2019. Stable isotopes in avian research: a step by step protocol to
461   feather sample preparation for stable isotope analysis of carbon ($\delta 13C$), nitrogen ($\delta 15N$), and
462   hydrogen ($\delta 2H$). Version 1.1. protocols.io
463   https://dx.doi.org/10.17504/protocols.io.z2uf8ew
464

465   DeLuca, W.V., Woodworth, B.K., Rimmer, C.C., Marra, P.P., Taylor, P.D., McFarland, K.P.,
466   Mackenzie, S.A. and Norris, D.R., 2015. Transoceanic migration by a 12 g songbird. *Biology*
467   *letters*, *11*(4), p.20141045.
468

469   Domer, A., Ovadia, O. and Shochat, E., 2018. Energy for the road: Influence of carbohydrate and
470   water availability on fueling processes in autumn-migrating passerines. The Auk: Ornithological
471   Advances, 135(3), pp.534-546.
472

473   Duijns, S., L. J. Niles, A. Dey, Y. Aubry, C. Friis, S. Koch, A. M. Anderson, and P. A. Smith.
474   2017. Body condition explains migratory performance of a long-distance migrant. Proc Royal
475   Soc B 284:20171374.
476

477   Fudickar, A.M., Greives, T.J., Atwell, J.W., Stricker, C.A. and Ketterson, E.D., 2016.
478   Reproductive allochrony in seasonally sympatric populations maintained by differential response
479   to photoperiod: implications for population divergence and response to climate change. *The*
480   *American Naturalist*, *187*(4), pp.436-446.
481

482   Green, A.J., 2001. Mass/length residuals: measures of body condition or generators of spurious
483   results? *Ecology*, *82*(5), pp.1473-1483.
484

485   Guglielmo, C.G., 2018. Obese super athletes: fat-fueled migration in birds and bats. *Journal of*
486   *Experimental Biology*, *221*(Suppl 1), p.jeb165753.
487

488   Hahn, S., Amrhein, V., Zehtindijev, P. and Liechti, F., 2013. Strong migratory connectivity and
489   seasonally shifting isotopic niches in geographically separated populations of a long-distance
490   migrating songbird. Oecologia, 173(4), pp.1217-1225.
491

492   Helms, C.W. and Drury, W.H., 1960. Winter and migratory weight and fat field studies on some
493   North American buntings. *Bird-banding*, *31*(1), pp.1-40.
494

495   Herrera, L.G., Hobson, K.A., Rodríguez, M. and Hernandez, P., 2003. Trophic partitioning in
496   tropical rain forest birds: insights from stable isotope analysis. Oecologia, 136(3), pp.439-444.
497

Hobson, K.A., Alisauskas, R.T. and Clark, R.G., 1993. Stable-nitrogen isotope enrichment in
avian tissues due to fasting and nutritional stress: implications for isotopic analyses of diet. *The*
Condor, *95*(2), pp.388-394.

Hobson, K.A. and Wassenaar, L.I., 1996. Linking breeding and wintering grounds of neotropical
migrant songbirds using stable hydrogen isotopic analysis of feathers. *Oecologia*, *109*(1),
pp.142-148.

Hobson, K.A., 1999. Stable-carbon and nitrogen isotope ratios of songbird feathers grown in two
terrestrial biomes: implications for evaluating trophic relationships and breeding origins. The
Condor, 101(4), pp.799-805.

Hobson, K.A., 2005. Using stable isotopes to trace long-distance dispersal in birds and other
taxa. *Diversity and Distributions*, *11*(2), pp.157-164.

Hobson, K.A., Van Wilgenburg, S.L., Wassenaar, L.I. and Larson, K., 2012. Linking hydrogen
(δ^2H) isotopes in feathers and precipitation: sources of variance and consequences for
assignment to isoscapes. *PloS one*, *7*(4), p.e35137.

Hobson, K.A. and Koehler, G., 2015. On the use of stable oxygen isotope (δ18O) measurements
for tracking avian movements in North America. *Ecology and Evolution*, *5*(3), pp.799-806.

Horton, K.G., Van Doren, B.M., Stepanian, P.M., Farnsworth, A. and Kelly, J.F., 2016. Seasonal
differences in landbird migration strategies. *The Auk: Ornithological Advances*, *133*(4), pp.761-
769.

Kaboli, M., Aliabadian, M., Guillaumet, A., Roselaar, C.S. and Prodon, R., 2007.
Ecomorphology of the wheatears (genus Oenanthe). *Ibis*, *149*(4), pp.792-805.

Keeley, J.E. and Rundel, P.W., 2003. Evolution of CAM and C4 carbon-concentrating
mechanisms. *International Journal of Plant Sciences*, *164*(S3), pp.S55-S77.

Kelly, J.F., Bridge, E.S., Fudickar, A.M. and Wassenaar, L.I., 2009. A test of comparative
equilibration for determining non-exchangeable stable hydrogen isotope values in complex
organic materials. Rapid Communications in Mass Spectrometry: An International Journal
Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry,
23(15), pp.2316-2320.

Kempster, B., Zanette, L., Longstaffe, F.J., MacDougall-Shackleton, S.A., Wingfield, J.C. and
Clinchy, M., 2007. Do stable isotopes reflect nutritional stress? Results from a laboratory
experiment on song sparrows. *Oecologia*, *151*(3), pp.365-371.
539

540   Liebgold, E.B., Gerlach, N.M. and Ketterson, E.D., 2019. Density-dependent fitness, not
541   dispersal movements, drives temporal variation in spatial genetic structure in dark-eyed juncos
542   (*Junco hyemalis*). *Molecular ecology*, *28*(5), pp.968-979.
543

544   Ma, C., Vander Zanden, H.B., Wunder, M.B. and Bowen, G.J., 2020. assignR: An R package for
545   isotope-based geographic assignment. *Methods in Ecology and Evolution*.
546

547   Maggini, I., Metzger, B., Voss, M., Voigt, C.C. and Bairlein, F., 2016. Morphometrics and stable
548   isotopes differentiate wintering populations of a migratory bird. *Movement ecology*, *4*(1), p.20.
549

550   Marra, P. P., E. B. Cohen, S. R. Loss, J. E. Rutter, and C. M. Tonra. 2015. A call for full annual
551   cycle research in animal ecology. Biol Letters 11:20150552.
552

553   Marra, P. P., K. A. Hobson, and R. T. Holmes. 1998. Linking winter and summer events in a
554   migratory bird by using stable-carbon isotopes. Science 282:1884–1886.
555

556   Marshall, T.J., Dick, M.F. and Guglielmo, C.G., 2016. Seasonal dietary shifting in yellow-
557   rumped warblers is unrelated to macronutrient targets. *Comparative Biochemistry and*
558   *Physiology Part A: Molecular & Integrative Physiology*, *192*, pp.57-63.
559

560   Nolan, V., Jr, and E. D. Ketterson. 1983. An analysis of body mass, wing length, and visible fat
561   deposits of Dark-eyed Juncos wintering at different latitudes. The Wilson bulletin.
562

563   Nolan, V., Jr, Ketterson, E. D., Cristol, D. A., Rogers, C. M., Clotfelter, E. D., Titus, R. C.,
564   Schoech, S. J., Snajdr, E, 2002. Dark-eyed Junco (*Junco hyemalis*), version 2.0. In The Birds of
565   North America (A. F. Poole and F. B. Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY,
566   USA. https://doi.org/10.2173/bna.716
567

568   Paritte, J.M. and Kelly, J.F., 2009. Effect of cleaning regime on stable-isotope ratios of feathers
569   in Japanese Quail (*Coturnix japonica*). *The Auk*, *126*(1), pp.165-174.
570

571   PIF (Partners in Flight), 2017. Avian Conservation Assessment Database, version 2017.
572   http://pif.birdconservancy.org.
573

574   Post, D.M., 2002. Using stable isotopes to estimate trophic position: models, methods, and
575   assumptions. Ecology, 83(3), pp.703-718.
576

577   R Core Team (2020) R: A Language and Environment for Statistical Computing. R Foundation
578   for Statistical Computing, Vienna, Austria. Available: http://www.r-project.org.

579   Rani, S., Singh, S., Malik, S. and Kumar, V., 2017. Insights into the Regulation of Spring
580   Migration in Songbirds. In *Biological Timekeeping: Clocks, Rhythms and Behaviour* (pp. 625-
581   642). Springer, New Delhi.
582
583   Reif, J., Hořák, D., Krištín, A., Kopsová, L. and Devictor, V., 2016. Linking habitat
584   specialization with species' traits in European birds. *Oikos*, *125*(3), pp.405-413.
585
586   Revelle, W. and Revelle, M.W., 2015. Package 'psych'. *The comprehensive R archive network*.
587
588   Rogers, C.M., Nolan, V. and Ketterson, E.D., 1994. Winter fattening in the dark-eyed junco:
589   plasticity and possible interaction with migration trade-offs. *Oecologia*, *97*(4), pp.526-532.
590
591   Rubenstein, D. R. and K. A. Hobson. 2004. From birds to butterflies: animal movement patterns
592   and stable isotopes. *Trends in Ecology and Evolution* 19(5): 256-263.
593
594   Rundel, C.W., Wunder, M.B., Alvarado, A.H., Ruegg, K.C., Harrigan, R., Schuh, A., Kelly, J.F.,
595   Siegel, R.B., DeSante, D.F., Smith, T.B. and Novembre, J., 2013. Novel statistical methods for
596   integrating genetic and stable isotope data to infer individual-level migratory connectivity.
597   *Molecular ecology*, *22*(16), pp.4163-4176.
598
599   Tsvey, A., Bulyuk, V.N. and Kosarev, V., 2007. Influence of body condition and weather on
600   departures of first-year European robins, Erithacus rubecula, from an autumn migratory stopover
601   site. *Behavioral Ecology and Sociobiology*, *61*(11), pp.1665-1674.
602
603   Sarkar, D., 2008. The Lattice Package: Lattice Graphics. R package Version 0.17-15. *URL:*
604   *http://cran. r-project. org/web/packages/lattice/index. html*.
605
606   Schaub, M., and L. Jenni. 2000. Body mass of six long-distance migrant passerine species along
607   the autumn migration route. J Für Ornithol 141:441–460.
608
609   Vander Zanden, H.B., Nelson, D.M., Wunder, M.B., Conkling, T.J. and Katzner, T., 2018.
610   Application of isoscapes to determine geographic origin of terrestrial wildlife for conservation
611   and management. *Biological Conservation*, *228*, pp.268-280.
612
613   Wassenaar, L.I. and Hobson, K.A., 2003. Comparative equilibration and online technique for
614   determination of non-exchangeable hydrogen of keratins for use in animal migration studies.
615   Isotopes in environmental and health studies, 39(3), pp.211-217.
616
617   Wassenaar, L.I. and Hobson, K.A., 2006. Stable-hydrogen isotope heterogeneity in keratinous
618   materials: mass spectrometry and migratory wildlife tissue subsampling strategies. *Rapid*

619 *Communications in Mass Spectrometry: An International Journal Devoted to the Rapid*
620 *Dissemination of Up‑to‑the‑Minute Research in Mass Spectrometry*, *20*(16), pp.2505-2510.
621
622 Wunder, M.B., 2010. Using isoscapes to model probability surfaces for determining geographic
623 origins. In *Isoscapes* (pp. 251-270). Springer, Dordrecht.
624
625 Yanco, S.W., McDevitt, A., Trueman, C.N., Hartley, L. and Wunder, M.B., 2020. A modern
626 method of multiple working hypotheses to improve inference in ecology. *Royal Society Open*
627 *Science*, *7*(6), p.200231.
628
629 Yang, L.H., Bastow, J.L., Spence, K.O. and Wright, A.N., 2008. What can we learn from
630 resource pulses. *Ecology*, *89*(3), pp.621-634.
631
632 Zenzal, T.J., Contina, A.J., Kelly, J.F. and Moore, F.R., 2018. Temporal migration patterns
633 between natal locations of ruby-throated hummingbirds (*Archilochus colubris*) and their Gulf
634 Coast stopover site. *Movement ecology*, *6*(1), p.2.

 Figure Legends:

Figure 1. Linear regression model between environmental precipitation isotope values ($\delta^2 H_p$)

and 22 known-origin feather samples of *M. melodia* samples (panel A) used to rescale the North

America hydrogen isoscape (panel B and C) for assigning migratory juncos wintering in

Oklahoma to their molting locations at the breeding grounds across northern United States and

Canada. The green and orange bars along the horizontal and vertical axis represent the

distributions of the sampling data across environmental and tissue isotope values, respectively

(panel A). The calibrated hydrogen isoscape mean values and standard deviations are presented

in panel B and C, respectively.

Figure 2. Cumulative assignment posterior probability density maps across four isotopic groups

of 76 Dark-eyed Juncos wintering in Oklahoma (red dot) and breeding at northern latitudes in the

United States and Canada. The bright yellow color represents high probabilities of molting

origin. The northern migrants (groups G1 and G2; panel A and B) were assigned to either Alaska

and western Canada, including Yukon and British Columbia, or central Northwestern Territories.

The southern migrants (groups G3 and G4; panel C and D) were prevalently assigned to either

Northern Saskatchewan or southern Manitoba and northern western United States. The hydrogen

stable isotope approach suggests that Dark-eyed Junco migrants from different breeding

populations converge into the same location in Oklahoma over the winter.

Figure 3. Model averaged estimates of mean $\delta^2 H_f$ values with 95% CI for levels of fat scores

with centered and scaled morphometric metric measures held at 0, their mean value (A) and

model averaged coefficient estimates of centered and scaled morphometric parameters with 95%

CI with level 1 of fat score as the reference for the intercept (B). Both plots indicate effects for

fat scores and morphometric variables on mean δ^2H_f are insignificant for our data set as

evidenced by CI overlap between fat score levels (A) and with zero (B).

Figure 4. Correlation between $\delta^{13}C$ and hydrogen δ^2H (panel A) and nitrogen $\delta^{15}N$ and hydrogen

δ^2H values (panel B). Migrants exhibiting longer wings show a broader range of $\delta^{13}C$ values

compared to medium- and short-length wings birds suggesting more flexibility in their habitat

use (panel C). We did not find noticeable trends between $\delta^{15}N$ and wing length (panel D).

Figure 5. Isotopic diet composition of $\delta^{13}C$ and $\delta^{15}N$ plotted over δ^2H spatial data distribution

(panel A). Blue areas represent lower latitudes of origin (δ^2H = from -110‰ to -145‰) while red

areas represent higher latitudes of origin (δ^2H = from -145‰ to -180‰). Each open circle

represents a single individual. Northern migrants show a moderate trend towards lower protein

consumption ($\delta^{15}N$: 1-4‰) and C3-plant habitat use ($\delta^{13}C$: -22 − -24‰). Southern migrants show

higher trophic level ($\delta^{15}N$: 4-8‰) and temperate forests habitat use ($\delta^{13}C$: -24 − -26‰). Panels B

and C show the distributions of $\delta^{13}C$ and $\delta^{15}N$ values across migrants originating from extreme

northern and extreme southern latitudes. Dashed red lines represent isotope mean values for each

group.

Table Legends:

Table 1. Variable importance results. Although the variable 'body fat' emerged as the

comparatively best predictor of hydrogen variation in feathers, the effect for this and all other

variables could not be differentiated from zero.

Table 2. Akaike Information Criterion corrected for small sample size (AICc) model selection

results. The most parsimonious model was the full model including all five morphological

variables but was equally competitive with a model omitting tarsus based on <2 unit difference

in ΔAICc values. Simpler single term or intercept only models were generally outperformed by

more complex models.

Supplementary Material

724
725   **Figure S1.** Distributions of $\delta^{13}C$ values across migrants originating across latitudinal groups.

726   Dashed red lines represent stable isotope mean values for each group.

728   **Figure S2.** Distributions of $\delta^{15}N$ values across migrants originating across latitudinal groups.

729   Dashed red lines represent stable isotope mean values for each group.

Figure 1

Figure 2

A

B

Figure 3

Figure 4

Figure 5

812
813
814 **Figure S1**
815
816
817
818

Figure S2

Variable	Sum of AICc model weights	Odds ratio
Fat	1	inf
Mass	0.87	6.69
Wing	0.86	6.14
Tail	0.72	2.57
Tarsus	0.54	1.17

Table 1

Model ID	Model Variables	logLik	AICc	delta	weight
32	h ~ Fat + Mass + Tail + Tarsus + Wing + (1|sex)	-222.1	471.48	0	0.3
24	h ~ Fat + Mass + Tail + Wing + (1|sex)	-223.74	471.8	0.33	0.25
28	h ~ Fat + Mass + Tarsus + Wing + (1|sex)	-224.61	473.54	2.06	0.11
20	h ~ Fat + Mass + Wing + (1|sex)	-226.25	473.95	2.47	0.09
30	h ~ Fat + Tail + Tarsus + Wing + (1|sex)	-225.43	475.16	3.69	0.05
22	h ~ Fat + Tail + Wing + (1|sex)	-226.94	475.33	3.86	0.04
16	h ~ Fat + Mass + Tail + Tarsus + (1|sex)	-225.65	475.61	4.13	0.04
8	h ~ Fat + Mass + Tail + (1|sex)	-227.18	475.83	4.35	0.03
12	h ~ Fat + Mass + Tarsus + (1|sex)	-227.46	476.38	4.91	0.03
4	h ~ Fat + Mass + (1|sex)	-228.98	476.68	5.21	0.02
26	h ~ Fat + Tarsus + Wing + (1|sex)	-227.69	476.85	5.37	0.02
18	h ~ Fat + Wing + (1|sex)	-229.2	477.11	5.63	0.02
14	h ~ Fat + Tail + Tarsus + (1|sex)	-229.25	479.96	8.48	0
6	h ~ Fat + Tail + (1|sex)	-230.84	480.39	8.91	0
10	h ~ Fat + Tarsus + (1|sex)	-232.01	482.74	11.26	0
31	h ~ Mass + Tail + Tarsus + Wing + (1|sex)	-233.91	483.9	12.42	0
2	h ~ Fat + (1|sex)	-233.92	483.91	12.44	0
23	h ~ Mass + Tail + Wing + (1|sex)	-235.4	484.32	12.85	0
27	h ~ Mass + Tarsus + Wing + (1|sex)	-236.3	486.13	14.66	0
19	h ~ Mass + Wing + (1|sex)	-237.79	486.64	15.16	0
15	h ~ Mass + Tail + Tarsus + (1|sex)	-237.14	487.81	16.33	0
29	h ~ Tail + Tarsus + Wing + (1|sex)	-237.26	488.04	16.56	0
7	h ~ Mass + Tail + (1|sex)	-238.62	488.32	16.84	0
21	h ~ Tail + Wing + (1|sex)	-238.76	488.6	17.12	0
11	h ~ Mass + Tarsus + (1|sex)	-238.93	488.93	17.45	0
3	h ~ Mass + (1|sex)	-240.4	489.51	18.03	0
25	h ~ Tarsus + Wing + (1|sex)	-239.41	489.89	18.41	0
17	h ~ Wing + (1|sex)	-240.9	490.51	19.03	0
13	h ~ Tail + Tarsus + (1|sex)	-240.75	492.58	21.1	0
5	h ~ Tail + (1|sex)	-242.43	493.56	22.09	0
9	h ~ Tarsus + (1|sex)	-243.45	495.6	24.13	0
1	h ~ (1|sex)	-245.58	497.58	26.1	0

860
861
862 **Table 2**