1

Comparative transcriptomics reveals unique patterns of convergence in the evolution of eusociality

9
10
11

Velasque, M.[1,2], Tan, Y.[1], Liu, A.W.[1], Luscombe, N.M.[1] & Denton, J.A.[1,3, *]

[1] Genomics & Regulatory Systems Unit, Okinawa Institute of Science & Technology, 1919-1

Tancha, Okinawa, Japan

[2] The Experimental Evolutionary Biology Lab, School of Biological Sciences, Monash University,

Clayton, Victoria, Australia

[3] The World Mosquito Program, Monash University, Clayton, Victoria, Australia

* Corresponding author: jai.denton@worldmosquitoprogram.org

Abstract

Eusociality represents a major evolutionary transition that arose independently in at least 12 insect lineages. Despite this prevalence, there remains considerable uncertainty surrounding the catalysing event and underlying genomic changes that enable such modifications. Commonly associated with this evolutionary transition is establishing and maintaining the reproductive division of labour (e.g. a reproductive queen and no-reproductive workers). This division is, at least in part, induced and maintained by highly species-specific pheromones. However, genomic analysis remains conflicted on the role of pheromones in this evolutionary transition. Specifically, if there was co-option of a common pheromone-sensitive genetic pathway present in all progenitor species or strong lineage-specific selection converging on similar transcriptomic signatures.

Using a solitary insect model, we sought to determine if various species-specific pheromones induced similar transcriptomic responses, thus activating similar pathways. We measured the transcriptomic and physiological response of a solitary insect, *Drosophila melanogaster*, to pheromones from bumblebees, honey bees, and termites. Each treatment induced the same strong physiological response - a decreased ovary size. However, employing several methods of transcriptomic analysis, we did not observe conservation in pheromone-mediated gene/pathway regulation.

Thus, despite a conserved phenotypic response, the underpinning transcriptome was vastly different. This suggests that pheromone-mediated eusociality is the result of convergent evolution. We propose that mechanisms maintaining eusociality (i.e. proto-pheromone) in early stages of eusocial evolution in each group, thus, acting as a primer for eusociality. This early state is then refined through strong selective pressure, resulting in a converging eusocial phenotype.

48   Visual Abstract

Introduction

The transition from solitary to group living is a major evolutionary jump (1). However, it requires complex communication mechanisms to convey essential information to the group, such as the presence of predators and resource location (2, 3). Although there are multiple methods to transmit and receive information (e.g. visual, auditory, electric, olfactory), chemical cues are one of the most important and earliest to evolve (2, 4).

Insects use chemical components for recruitment, defence, predator recognition, and mate attraction, as well as to establish and shape social interactions (5–7). However, some eusocial organisms have evolved beyond context-dependent messaging. Acting as primer pheromones, they induce long-term behavioural and physiological effects by regulating colony structure and dynamics (8–11). This ranges from modulating aggression and attraction to monopolising colony reproduction through regulating female fecundity (12, 13). This monopolisation of reproduction by a few or one groups (queens/kings) creates a reproductive division of labour (8, 9). This is the cornerstone of eusociality (14–17).

Pheromone-mediated eusociality is not unique to honey bees as it has independently evolved at least 12 times in arthropods (18). The diverse evolutionary history of eusociality is also reflected in the production and structure of pheromones controlling the process (19–21). For instance, in several species, cuticular hydrocarbons (CHCs), a component ubiquitously produced in insects to prevent desiccation and infections, can act as queen pheromones (22–24). Such as in stingless and bumble bees, *Bombus terrestris,* where CHCs act as a queen pheromone (22, 25). In other species, such as the subterranean termite *Reticulitermes speratus,* queen pheromone is a mixture of CHCs, n-butyl-n-butyrate and 2-methyl-1butanol. However, the use of CHCs is not conserved in other eusocial animals. In *Apis mellifera,* queen pheromone comprises a complex blend of fatty acids secreted by the (26–28) mandibular gland (26–28). Moreover, pheromones produced by the brood (brood pheromone – Brood) were also found to regulate behaviour and worker reproduction in several species, such as *A. mellifera* (29–31).

In eusocial animals, the reproductive division of labour usually initiates during
early larval development and is maintained throughout adulthood via exposure to queen
pheromone (32, 33). Hence, in the absence of such pheromones, the division of labour
breaks down as young workers activate their ovaries and engage in more selfish
behaviour (34–38).

In eusocial animals, the division of labour and reproduction are suggested to be
intertwined, arising from modifications of the reproductive life cycle of solitary insects
(39, 40). Comparative genomics of multiple eusocial insect species corroborates this
hypothesis and suggests that eusocial traits evolved from the co-option of pathways
present in solitary ancestors (41–45). Moreover, there is strong evidence that specific
queen pheromones evolved from conserved signals of solitary ancestors (22, 45, 46). It
has been proposed that eusociality requires a highly conserved group of core genes,
present in the solitary ancestor, that serves as a ground plan for eusociality (43).
However, clear unambiguous evidence of this co-option hypothesis remains elusive (42,
47).

Further expansion of comparative genomics has added to this ambiguity.
Genome analysis revealed branch-specific positive selection in five bee species from
the *Apis* genus (48). Specifically, a greater proportion of species-specific genes were
under positive selection than *Apis*-specific (45, 48). Furthermore, despite similar
eusocial behaviour, three eusocial termite species demonstrated vastly different
underpinning genetic mechanisms when contrasted against a wide diversity of non-
eusocial and eusocial insect species (49). Thus, eusociality may be induced and
maintained through diverse genetic means.

Pheromone-induced transcriptomic changes reflect extensive evolutionary
optimisation of affected networks. Thus, cross-species transcriptomic comparisons are
complicated due to the evolutionary distances between eusocial insect species, as inter-
species gene orthologs may have vastly different functions (44, 50–52). The birth of
novel genes and the expansion or contraction of gene families can make clear
orthologue identification impossible and can be conflated with evolutionary forces (52–
54). Therefore, differences or similarities in expression patterns between two highly
divergent species can be highly misleading (55). One approach to overcome this

problem is by using a single species to investigate transcriptomic changes induced by

various pheromones (56–58). This permits the study of conserved and co-opted

transcriptomic changes and identifies molecular signatures of these rare evolutionary

events.

Here, we investigate pheromone-mediated gene expression in a solitary species.

Using *Drosophila melanogaster*, we examine if primer pheromones, regulating

eusociality, evolved from the co-option of similar pathways present in a solitary

ancestor. Specifically, whether distinct pheromones have a similar transcriptomic and

physiological profile across multiple instances on which they evolved, as previously

suggested. We present a comprehensive transcriptome dataset set investigating gene

expression associated with the reproductive division of labour. We show the variation in

differential gene expression induced termite (*Reticulitermes speratus*), bumblebee

(*Bombus terrestris*) and honey bee (*Apis mellifera*) queen pheromone and honey bee

brood pheromone. We leverage this data set to perform an up to date and

comprehensive comparison of differentially vs distinct (i.e. non-homologous) expressed

genes associated with the maintenance of eusociality.

Results

Pheromone-induced Ovary Changes

We identified a significant change in ovary size across all pheromone-treated *D.*

melanogaster (Repeated measures ANOVA $F_{7,180}$, $p < 0.0001$; Figure 1; Tables S1 &

S2). Although all pheromone treatments strongly inhibited ovary size, their effects varied

in strength. Bumblebee and Termite pheromone had similar ($p = 1$) severe pheromone-

mediated ovary reductions of 0.32 mm^2 and 0.35 mm^2 (mean; n=10), respectively

(Figure 1; Table S1). Pheromone treatments from *A. mellifera* queen mandibular

pheromone (QMP), Brood and the QMP and Brood mix pheromone induced similar

reductions in ovary size (QMP and Brood vs QMP $p =1$; QMP and Brood vs Brood $p =1$;

QMP vs Brood $p = 0.064$) of 0.51 mm^2, 0.65 mm^2 and 0.59 mm^2 (mean; n=10)

respectively (Figure 1; Table S1).

Figure 1. Pheromone treatment resulted in reduced *Drosophila melanogaster* ovary size.
Treatments were queen pheromone produced by termites (*Reticulitermes speratus*), bumblebees
(*Bombus terrestris*) and honey bees (*Apis mellifera*) (QMP), brood pheromone produced by honey bee
larvae and a combination of both brood and QMP from honey bees. The pheromones were introduced to
D. melanogaster from egg to adult via supplementation in food. Letters, either a, b or c, denote statistical
significance with varying letters being significantly different (p-value < 0.05; Multiple Comparisons of
Means: Tukey Contrasts; Table S2). Ten biological replicates were performed for each treatment or
control with each biological replicate. Ovaries were measured from three flies per each replicate.

Pheromone-induced Transcriptomic Changes

Differentially expressed genes and networks

Despite phenotypic similarities, we identified no transcriptomic overlap between
pheromone treatment groups. Within each treatment, we identified similar numbers of
differentially expressed genes with brood pheromone treatment being elevated: QMP
(180 genes), Brood (640 genes), QMP and Brood (293 genes), Termites (207 genes)
and Bumblebees (102 genes). There was no shared core of pheromone response
genes and negligible inter-treatment overlap (Figure 2A; Figure S1). Due to their

potential in eusociality regulation, differential expression of the Notch signalling pathway

and the yellow gene family was specifically examined (59, 60). No differential

expression of any genes encoding the 12 Notch signalling pathway core components or

any of the 14 genes in the yellow family was detected in any treatment.

As we were unable to identify a core of differentially expressed genes, we

investigated if pheromone-mediated changes in gene expression affected reproductive

physiology through similar cellular processes. We compared gene enrichment of

pheromone-induced changes in the transcriptome using GO (gene ontology) and KEGG

(Kyoto Encyclopedia of Genes and Genomes) annotations. Although we identified some

inter-treatment pathway enrichment, again, no core pathways were identified (Figure

2B).

In the absence of a conserved transcriptomic core, a broader general correlation

between treatments was undertaken. There was a high degree correlation between the

$\log_2$-fold-change (Log_2FC) of all significantly expressed genes in each treatment (Figure

2C). To further investigate pheromone-mediated changes in genetic pathways and

transcriptomic networks, we employed two different network analysis approaches.

Figure 2. Differential gene expression of pheromone-sensitive pathways and genes in *Drosophila melanogaster* does not show a shared response. The DeSeq package was employed to determine differentially expressed genes in each of the pheromone treatments. A) Venn diagram showing the number of significantly differentially expressed genes per pheromone treatment. B) Results of signalling pathway analysis of differentially expressed genes across different pheromone treatments. C) Pearson's correlation coefficient between the Log_2-FC of significantly expressed transcripts in each treatment. Treatments were queen pheromone produced by termites (*Reticulitermes speratus*), bumblebees (*Bombus terrestris*) and honey bees (*Apis mellifera*), brood pheromone produced by honey bee larvae and a combination of both brood and queen pheromone from honey bees.

Network Analysis

Employing network-based transcriptomic analysis, and weighted gene co-

expression network analysis (WGCNA), no shared regulatory network common among

all pheromone treatments was identified (Figure 3). We used WGCNA to screen and

build a gene co-expression network, screen important modules, and identify candidate

genes associated with each pheromone treatment (hub genes). We identified candidate

genes by estimating the correlation between each gene and the pheromone treatment

using Gene Significance analysis (61–63). WGCNA identified statistically significant

differentially co-expressed genes between control groups and *D. melanogaster* treated

with QMP (219 genes), Brood (219 genes), the mix of QMP and Brood (396 genes),

Termites (307 genes) and Bumblebees (276 genes) (Figure 3). Once again, each

treatment had limited overlap (Figure 3A; Figure S2). Employing GO enrichment and

gene set enrichment analysis (GSEA) on the WGCNA differentially co-expressed genes

also failed to identify high levels of overlap between functions (Figure 2B).

Once again, there was a high degree correlation between the $\log_2$-fold-change

(Log_2FC) of all significantly expressed genes in each treatment (Figure 2C). However,

Brood pheromone treatment resulted in a negative correlation for the WGCNA-identified

transcripts. This was not observed in the mix of QMP and Brood treatment.

Figure 3. Network analysis showing pheromone-sensitive pathways and genes in *Drosophila melanogaster* does not show a shared response. Weighted gene co-expression analysis was used to screen and build a gene co-expression network, screen important modules, and identify candidate genes associated with each pheromone treatment. A) Venn diagram showing the number of significantly differentially expressed genes per pheromone treatment. B) Results of signalling pathway analysis of differentially expressed genes across different pheromone treatments. C) Pearson's correlation coefficient between the Log$_2$-FC of significantly expressed transcripts in each treatment. Treatments were queen pheromone produced by termites (*Reticulitermes speratus*), bumblebees (*Bombus terrestris*) and honey bees (*Apis mellifera*), brood pheromone produced by honey bee larvae and a combination of both brood and queen pheromone from honey bees.

Cross-species comparison

We found that pheromone-mediated transcriptomic changes between eusocial

insects and *D. melanogaster* were preserved*.* Although *D. melanogaster* has been

previously used as a model organism in the study of eusociality (64), there are no direct

comparisons between pheromone-mediated changes in gene expression for flies

compared to eusocial insects. Holman et al. studied the transcriptomic response of

several species, including *A. mellifera* and *B. terrestris,* treated with their own species'

pheromone (44). Using WGCNA module preservation, we determined the conservation

of pheromone-mediated network signatures in our *D. melanogaster* experiments to their

corresponding equivalents from Holman et al. Specifically, between our *D.*

melanogaster exposed to QMP (*A. mellifera*) and Holman et al. *A. mellifera* exposed to

QMP (*A. mellifera*), and our *D. melanogaster* exposed to bumblebee QMP (*B. terrestris*)

and Holman et al. *B. terrestris* exposed to bumblebee QMP.

Comparing these datasets, we identified a high degree of WGCNA module

preservation (Table 1; Figure 4; Figure S3). Overall, the *A. mellifera* comparison had a

higher number of preserved modules and, thus, a greater number of shared genes in

these modules. However, *A. mellifera* preservation also had a greater proportion of

preserved modules with strong and moderate preservation (Figure 4A; Figure S3). The

contrast between these two comparisons is likely down to the greater number of

identified orthologues in *A. mellifera*. Regardless of the number, the preservation of

these modules does demonstrate, in a broad sense, a similar transcriptomic response,

and those serve as an external control for our analysis.

Table 1. Cross-species model preservation of pheromone-response. The WGCNA module-
preservation function facilitated comparisons between *Drosophila melanogaster* and *Apis mellifera*
pheromone response to *A. mellifera* queen pheromone and *D. melanogaster* and *Bombus terrestris*
pheromone-response to *B. terrestris* queen pheromone. Number of shared modules between the two.
Total number of genes identified.

Species	Number of Modules	Number of Genes
Strong		
A. mellifera	30	2102
B. terrestris	9	721
Moderate		
A. mellifera	41	2636
B. terrestris	15	1310
Low		
A. mellifera	16	769
B. terrestris	23	713
Dissimilar		
A. mellifera	1	43
B. terrestris	4	115

244
Figure 4. Pheromone-responsive module preservation between *Drosophila melanogaster* and *Apis*
***mellifera* or *Bombus terrestris*.** Comparing *D. melanogaster* pheromone-response data from this work
and *A. mellifera* and *B. terrestris* pheromone-response data from Holman et al. Gene modules were
determined using WGCNA module preservation. *A)* Proportion conserved pheromone sensitive genes in
D. melanogaster, *A. mellifera* and *B. terrestris*. B) Summary of preservation evidence for pheromone
response. Pheromonal sensitivity in *A. mellifera* and *B. terrestris* were compared with *D. melanogaster*.
Shared genes indicate those highly conserved between the two comparison groups.

Origins of Pheromone Response Genes

The responsive genes of each pheromone treatment do not have a conserved

evolutionary origin, and thus eusociality is not co-oping recently evolved genes. The

absence of overlapping differentially expressed pheromone-sensitive genes between

treatments led to querying the evolutionary origins of these responsive pathways. We

employed phylostratigraphy and calculated the transcriptome age index (TAI) to infer

the evolutionary origins of pheromone-activated networks and genes. When separated

by treatment, each group of differentially expressed pheromone-responsive genes lack

a statistically significant conversed age (Figure 4A). If each treatment was further

separated into up- and down-regulated differentially expressed transcripts, up-regulated

transcripts from both termite and QMP treatments were statistically significant (Figure

4B).

A phylogenetic tree was created to determine the evolutionary relationship

between target species homologues of the pheromone-responsive genes (Figure 4C).

The tree was built using pylostratigraphic gene ages and phylogenetic relationships

amongst a diverse collection of 182 species, with most species collapsed into

systematic order or higher classification.

Figure 4. The evolutionary origins of pheromone responsive genes. The transcriptome age index

(TAI) was used to infer the evolutionary origins of pheromone-responsive genes. A) Summary of all

pheromone-sensitive genes per strata. B) Proportion of pheromone-sensitive genes in each treatment per

strata. C) A phylogenetic tree with pylostratigraphic gene ages and phylogenetic relationships amongst

182 species. Dots indicate the evolutionary age of different pheromone-sensitive genes. Asterisks

indicate phylogenetic levels (phylostrata) with significant changes in the expression of pheromone-

sensitive genes.

Discussion

Although all pheromone treatments suppressed ovary development in *D.*
melanogaster, we found no evidence of a shared evolutionary pathway or conserved
mechanism underpinning this phenotype. In social insects, chemical communication has
been independently co-opted in numerous species to facilitate the inhibition of worker
reproduction by dominant females (4, 5). The prevailing hypothesis is that this induced
sterility evolved from modifications in conserved molecular pathways regulating the
physiology and behaviour of solitary species (44, 45, 59, 65, 66). Therefore, it was
reasonable to expect similar pheromone-induced responses between different insect
lineages and, thus, similar physiological and regulatory pathway responses. We
removed the complexities of cross-species analysis and millions of years of convergent
evolution by applying pheromones that inhibit worker reproduction from honeybees,
bumblebees, and termites to a single species.

In contextualising our work, we are mindful of conflating factors introduced
through cross-species analysis, especially when the species are separated by large
evolutionary distances (18, 20, 48, 67). Furthermore, transcriptomic enrichment
analyses, employed widely for cross-species studies, can potentially introduce biases
(68, 69). However, we employed relaxed differential gene filtering (p<0.05, irrespective
of expression fold-change) and diverse enrichment methods. Our view was that
subsequent manual curation would allow us to identify candidate genes acting as
universal eusocial catalysts. This proved unnecessary as we could not identify a core of
differentially expressed genes and minimal inter-treatment overlap (Figure 2). Even GO
enrichment and pathway analysis, highly permissive methods, only revealed minimal
inter-group overlap.

Unable to identify a core of pheromone-responsive transcripts, we employed a
candidate-based approach. Pheromone exposure has previously been shown to
activate pathways related to lipid biosynthesis, oogenesis and olfaction (44, 45, 59). For
instance, Notch appears to be an essential pathway regulating ovary development in
the presence of queen pheromone (59) and it is generally assumed to be a co-opted
pathway in solitary animals that facilitates eusociality (59). Although we found Notch to
be differentially co-expressed in three of the five treatments, termite, QMP and QMP +

brood, differing Notch signalling genes were affected in each treatment. An alternative

candidate pathway is the yellow gene family. In Hymenoptera, this gene family is also

responsible for regulating aspects of eusociality and includes a clade of genes referred

to as the major royal jelly proteins (60, 70). In *D. melanogaster,* there are 14 members

of this gene family (60, 71). The yellow gene family (*yellow-h* and *yellow-g2 in D.*

melanogaster) was differentially co-expressed across all groups except for the brood

treatment. Thus, these pathways seemingly provide a potential co-option target but do

not appear to be a universal driver of pheromone response. Rather, given the

widespread regulatory changes associated with eusociality (72–74), these gene families

likely serve as candidate targets in refining eusocial traits.

Despite differences in the unpinning transcriptomic response, each treatment

yielded smaller ovary size (Figure 1). As previous work demonstrated, pheromone

exposure had no effect, even at much higher doses, on adult *D. melanogaster* (75). The

ineffectiveness of pheromone-mediated reproductive inhibition on adults is also

observed in other eusocial species (75–79). Insect oogenesis is a highly sensitive

process disrupted by numerous environmental, developmental and gene-regulatory

factors (80, 81). This includes the fact that cell death in the ovary is distinct from other

tissues and mediated independently (80, 81). Furthermore, pheromone-mediated

reproductive inhibition is not observed in primitively eusocial species like *Polistes satan*

(paper wasp) or species with high reproductive plasticity like *Euglossa dilemma* (orchid

bee) (41, 82). However, orchid bees form small 2–3 individual social groups wherein an

abundance of specific cuticular hydrocarbons correlated with social dominance and

ovary size (83). Therefore, oogenesis is a key candidate for pheromone-mediated

disruption. Moreover, influencing oogenesis provides a potential mechanism for the

monopolisation of colony reproduction and the evolution of eusociality.

Despite arising independently, several eusocial species share phenotypic and

transcriptomic similarities (41–43). However, consistent with a growing body of

comparative genomics (20, 84), we suggest this is not due to the co-option of identical

or similar pathways in solitary ancestors. Rather, positive selection drives the

convergence of characteristically eusocial phenotypes with diverse genomic origins. An

example of this arises in the evolution of intra-colony communication. Insects rely on a

host of olfactory signals and cues with the majority of these being detected by

chemosensory receptors in the odorant receptor (OR) or ionotropic receptor (IR) gene

families (85). These chemosensory pathways provide the core mechanism for intra-

colony communication in several eusocial species (19, 20, 86, 87). Genomic analysis

revealed that although termites and honey bees have expanded receptor gene families,

termites had greater IRs and bees greater ORs (49, 86, 88–91). This means that

although high-level analysis, such as GO, may detect these two gene family expansions

as similar, closer examination demonstrated their independence.

D. melanogaster accurately depicts the effect of queen pheromone in eusocial

species. When comparing our results with the transcriptome of honey bees and

bumblebees exposed to queen pheromone (44), we found strong conservation of

differentially expressed genes in pheromone treatments. Demonstrating the reliability of

our results and the power of comparing the effect of distinct components (i.e.

pheromones) in a single species.

Our work demonstrates the potential origins of eusociality evolution. The first

steps of eusocial evolution likely began with a localised reproductive impediment that,

over time, resulted in quasi-eusocial living. The absence of a core pheromone-induced

transcriptomic response and yet conserved phenotypic response, suggests female

fertility suppression can rapidly evolve using numerous compounds that induce

regulatory changes. This is evident as eusociality has evolved at least eight times

independently with limited overlap between the specific pheromones meditating this (67,

92).

The primer pheromonal signal in insects originated at least 100 Mya and evolved

independently in multiple species. Their evolution and signal can be well traced across

three groups: Blatoidea, Hymenoptera and Coleoptera, causing virtually a similar

phenotype in all groups. The independent evolution of pheromones, the high

conservation with their eusocial counterpart, combined with our results (i.e. pheromone

treatment causing a similar ovary reaction in flies with no overlapping gene network and

pathways) suggests pheromones are not a product of gene co-option, but the result of

convergent phenotypic evolution. Producing a similar phenotype (inhibition of ovary

development) through a different pathway in the species on which it evolved.

Materials & Methods

We used the fruit fly *D. melanogaster* to investigate the hypothesis that eusociality evolved multiple times by co-opting existing regulatory pathways present in a solitary ancestor. We achieved this by studying the pheromone genetic signature of fruit flies exposed to pheromones from three different eusocial species: honey bee queen mandibular pheromone (QMP), bumblebee queen pheromone (Bumblebee), termite queen pheromone (Termite) and a pheromone produced by honey bee larvae, the brood pheromone (Brood). To simulate the conditions present in honey bee colonies, when they are exposed to both brood and queen pheromone, we also added a group with a mix of queen and brood pheromone (QMP + Brood).

Fly husbandry

Flies were raised on standard Bloomington media at 24 °C under a 16h light / 8h dark cycle in 50 mL vials. The *D. melanogaster* strain Canton-S, a generous gift from the Van Vactor Unit, was used in all experiments. Each experimental vial contained ten males and females, of approximately 5-day-old. Each treatment group consisted of 10 vial replicates.

Pheromone treatment

Bloomington fly food was made and cooled to >50°C before mixing in each phenomenon treatment (Table 2) at a 1 in 150 ml concentration. Ten millilitres of supplemented food were added to each vial and left to dry for 24 hours. Vials were kept at 25°C and under a powerful extractor to prevent cross-contamination due to pheromone volatility. Adult flies were left on the vial for four days to lay eggs and then removed and discarded. Virgin females were collected and allowed to mature for five days in vials containing the aforementioned food with the corresponding treatment pheromone.

Table 2 – Pheromone Treatments, suppliers, and dilution.

Treatment	Pheromone Formula	Supplier (ID)	Dilution
Control	-	-	50 µl isopropanol
QMP (*Apis mellifera*)	9-ODA, two isomers of 9-HDA, HOB and HVA	TempQueen (DC-705)	20 µl of synthetic QMP mix diluted in 30 µl isopropanol
Brood pheromone (*Apis mellifera*)	Methyl palmitate, Ethyl palmitate, Methyl stearate, Ethyl stearate, Methyl oleate, Ethyl oleate, Methyl linoleate, Ethyl linoleate, Methyl linolenate, Ethyl linolenate	Contech (SuperBoost)	10 µl SuperBoost diluted in 40 µl isopropanol
QMP and Brood pheromone mix (*Apis mellifera*)	9-ODA, two isomers of 9-HDA, HOB and HVA; Methyl palmitate, Ethyl palmitate, Methyl stearate, Ethyl stearate, Methyl oleate, Ethyl oleate, Methyl linoleate, Ethyl linoleate, Methyl linolenate, Ethyl linolenate	TempQueen (DC-705) and Contech (SuperBoost)	- 20µL of synthetic QMP, 10µL Supperboost and 20µL isopropanol
Bumblebee queen pheromone (*Bombus terrestris*)	CHC pentacosane (C25)	Sigma-Aldrich (286931)	1 µg C25 diluted in 50 µl isopropanol
Termite queen pheromone (*Reticulitermes speratus*)	n-butyl-n-butyrate and 2-methyl-1-butanol	n-butyl-n-butyrate Sigma-Aldrich (281964); 2-methyl-1-butanol Wako Chemicals (133-08385)	15 µl n-butyl-n-butyrate and 7.5 µl 2-methyl-1-butanol diluted in 27.5 µl isopropanol

Ovary measurements

Ten biological replicates were used for each treatment or control. For each replicate, ovary collection for measurement was performed in triplicate and on the same day to prevent temporal changes in physiological measurements of ovaries. Ovary image identities were encoded such that they could be measured blind to reduce bias. Measurements were performed using ImageJ2 (93). We used normal QQplot and Shapiro-Wilk test to identify deviations from normality. Although termites had two

outliers, all the points fall approximately along the reference line, and thus we assumed

normality. We compared differences in ovary size between different treatment groups

using One-way repeated-measures ANOVA.

RNA Sequencing

For RNA sequencing, we randomly selected four vials and pulled five ovaries per

vial. RNA extraction was performed with an in-house modified protocol (Supporting

Text). Libraries were prepared using NEBNext® Ultra™ II Directional RNA Library Prep

Kit (E7760L) for Illumina as instructed by the manufacturer.

Data Analysis

The complete analysis pipeline and associated results are publicly available at

https://github.com/marivelasque/eusociality_evolution. This pipeline is a modified

version of our previously developed pipeline (94). RNA data were trimmed with

Trimmomatic 0.38 (95) and quality was assessed using FASTQC (96). Transcript

abundance was calculated with RSEM/bowtie2 (97, 98). Mapping and abundance

calculations were performed against the *D. melanogaster* genome assembly BDGP6

(release 89). Differential expression analysis was performed using edgeR (99). We

adjusted p-values to control the false discovery rate using the Benjamini-Hochberg

method. The threshold of significance was defined by $p < 0.05$. Visualisation relied on

packages venn (100), ggplot2 (101), igraph (102) and ggraph (103). Analysis, figures

and HTML files were generated using R in the RStudio environment.

Gene co-expression network analysis

To investigate changes in gene networks mediated by eusocial pheromones we

used Weighted Gene Co-Expression Network Analysis WGCNA (62). We used the

WGCNA R package to construct the co-expression network using gene counts obtained

from RSEM. Prior to the analysis, we removed gene outliers using the

"goodSamplesGenes" (62) function and accessed overall sample clustering in the

expression matrix using "hclust" (104). We used the lowest threshold power that

resulted in approximate scale-free topology to construct the gene co-expression

network using the blockwiseModules function (62). Then, we estimated the gene

expression profile, module eigengene, and the module-treatment relationship,

calculating the correlation between the module eigengene and treatment group. We

implemented the analysis in two steps (i.e. lowest threshold power and differential co-

expression) using the functions SoftPower and wgcna.est from (94). The threshold of

significance was defined as $p<0.05$.

Enrichment and pathway analysis

We obtained Gene Ontology (GO) and KEGG (Kyoto Encyclopedia of Genes and

Genomes) annotations using the package biomaRt (105) clusterProfiler (104)

respectively. We used Fisher's exact test to identify significantly enriched genes using

the package topGO (106) with the threshold of significance defined by $P<0.05$. We

identified enriched pathways using Pathway-Express, a package part of the Onto-Tools

(107). The significant pathways for predicting target genes were identified according to

the Kyoto Encyclopedia of Genes and Genomes (KEGG) database obtained using the

package clusterProfiler (104, 108). A minimal size of 40 genes of each geneSet was

used, and the significance threshold was defined by $p<0.05$. Although there was an

overlap on "Ribosome biogenesis in eukaryotes", the enrichment showed a different

directionality, overrepresented in Bumblebee, Termites, QMP, a mix of QMP and Brood

over-represented in Brood pheromone.

The pathway analysis was implemented using the Pathway-Express package

part of the Onto-Tools (107). Signalling pathways were identified using only differentially

expressed genes. The relationship between each gene was inferred based on their

KEGG pathway and magnitude of expression change (logFC,) and the visualisation of

significantly enriched pathways was performed using Igraph (102).

Cross-species comparison

To ensure pheromonal-induced changes in the fruit fly transcriptome are

comparable to eusocial insects, we compared the network conservation of eusocial

insects and fruit flies exposed to pheromone. Holman et al. (44) compared the whole

transcriptome of several species, including *Apis mellifera* and *Bombus terrestris,* treated

with either solvent-only control or their own species' queen pheromone. To determine

whether the QMP network signature was well conserved between *D. melanogaster*, *A.*

mellifera and *B. terrestris,* we used a WGCNA integrated function (modulePreservation)

(62) to calculate module preservation statistics. Orthologs between these two groups

were obtained using BLAST-based on the *D. melanogaster* genome. Holman et al. data

were obtained from BioProject DRP004516 (44). A total of 5507 unique *D.*

melanogaster orthologs were identified in *A. mellifera* and 2744 in *B. terrestris*. To

ensure that the conserved genes between the two data sets correspond to pheromone-

sensitive genes, we compared the differential expression between each QMP treatment

and control using exactTest function in edgeR (99). The threshold of significance was

defined by P<0.05. No publically available pheromone termite treatment data were

identified.

Evolutionary Origins

We performed a protein-based homology search across the tree of life to

determine the evolutionary age of pheromone-sensitive genes in *D.melanogaster* using

Phylostratr (109). We compared *D. melanogaster* gene sequences against 182 spp.

(NCBI proteins database) by the blastp algorithm at the e-value cutoff of 1e-03 and

selected the best match for pheromone-sensitive ortholog in each species. Following,

we compared the phylogenetic age of the pheromone-sensitive genes within each

treatment using the Transcriptome age index (TAI)(110, 111). To estimate TAI, we

retrieved the phylostrata (15 out of 27 phylostrata) of all genes that were differentially

expressed across all pheromone treatments (all pheromone-sensitive genes) and their

respective normalised expression. We used the pheromone-sensitive phylostrata to

reconstruct phylogeny using ape and ggtree (112–115).

Declarations

Acknowledgements

The Okinawa Institute of Science & Technology Sequencing Centre kindly

generated the sequencing data. Resources and expertise from the Okinawa Institute of

Science & Technology High-Performance Computing Section were employed for data

analysis. This work was funded by the Japanese Society for the Promotion of Science

KAKENHI grant scheme (grant numbers 19K16205, 19K06795 awarded to MV and

JAD, respectively) and the Okinawan Institute of Science & Technology. All authors

were supported by the Okinawa Institute of Science & Technology. MV and JAD were

also supported by Monash University and the World Mosquito Program Ltd,

respectively.

Availability of Data & Materials

All code and compiled markdown file showing will be available on GitHub prior to

publication https://github.com/marivelasque/eusociality_evolution. A permanent copy of

the final analysis markdown file is available at XXX <DOI provided prior to final

publication>. All raw sequencing data are available under the BioProject XXX <provided

prior to final publication>.

Author Contributions

MV and JAD conceived the project, designed and performed the experiments,

analysed the results and wrote the paper. YT, MV and AL prepared the sequencing

libraries. All authors edited the manuscript and interpreted the results.

References

1. J. M. Smith, E. Szathmary, *The Major Transitions in Evolution* (Oxford University Press, 1997).

2. T. D. Wyatt, *Pheromones and Animal Behavior: Chemical Signals and Signatures* (Cambridge University Press, 2014).

3. F. G. Barth, A. Schmid, *Ecology of Sensing* (Springer Science & Business Media, 2013).

4. Bradbury, Jack W., and Sandra L. Vehrencamp, *Principles of Animal Communication* (Sunderland, 1998).

5. N. M. A. El-Ghany, N. M. Abd El-Ghany, "Pheromones and Chemical Communication in Insects" in *Pests, Weeds and Diseases in Agricultural Crop and Animal Husbandry Production*, Dimitrios Kontogiannatos, Anna Kourti, Kassio Ferreira Mendes, Ed. (IntechOpen, 2020), pp. 16–30.

6. J. F. A. Traniello, S. K. Robson, Trail and Territorial Communication in Social Insects. *Chemical Ecology of Insects 2*, 241–286 (1995).

7. C. G. Butler, "Chemical Communication in Insects: Behavioral and Ecologic Aspects" in *Communication by Chemical Signals*, Johnston Jr., J. W., Moulton, D. G. and Turk, A., Ed. (D. Appleton, 1970), pp. 35–78.

8. M. L. Winston, K. N. Slessor, Honey bee primer pheromones and colony organization: gaps in our knowledge. *Apidologie (Celle)* **29**, 81–95 (1998).

9. S. D. Leonhardt, F. Menzel, V. Nehring, T. Schmitt, Ecology and evolution of communication in social insects. *Cell* **164**, 1277–1287 (2016).

10. J. S. van Zweden, The evolution of honest queen pheromones in insect societies. *Commun. Integr. Biol.* **3**, 50–52 (2010).

11. A. Ward, M. Webster, "The Evolution of Group Living" in *Sociality: The Behaviour of Group-Living Animals*, A. Ward, M. Webster, Eds. (Springer International Publishing, 2016), pp. 191–216.

12. J. Olejarz, C. Veller, M. A. Nowak, The evolution of queen control over worker reproduction in the social Hymenoptera. *Ecol. Evol.* **7**, 8427–8441 (2017).

13. G. C. Varley, E. O. Wilson, The insect societies. *J. Anim. Ecol.* **43**, 261 (1974).

14. Y. L. Conte, Y. Le Conte, A. Hefetz, Primer Pheromones in Social Hymenoptera. *Annual Review of Entomology* **53**, 523–542 (2008).

15. T. I. Szabo, Behavioural studies on queen introduction in the honeybee 3. Relationship between queen attractiveness to workers and worker aggressiveness towards a queen. *J. Apic. Res.* **13**, 161–171 (1974).

16. J. Beetsma, The process of queen-worker differentiation in the honeybee. *Bee World* **60**, 24–39 (1979).

17. S. Dobata, Arms race between selfishness and policing: two-trait quantitative genetic model
for caste fate conflict in eusocial Hymenoptera. *Evolution* **66**, 3754–3764 (2012).

18. E. O. Wilson, B. Hölldobler, Eusociality: origin and consequences. *Proc. Natl. Acad. Sci. U.*
S. A. **102**, 13367–13371 (2005).

19. B. J. Fischman, S. H. Woodard, G. E. Robinson, Molecular evolutionary analyses of insect
societies. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 2**, 10847–10854 (2011).

20. H. Yan, J. Liebig, Genetic basis of chemical communication in eusocial insects. *Genes Dev.*
35, 470–482 (2021).

21. J. Billen, J. Šobotník, Insect exocrine glands. *Arthropod Struct. Dev.* **44**, 399–400 (2015).

22. A. Van Oystaeyen, *et al.*, Conserved Class of Queen Pheromones Stops Social Insect
Workers from Reproducing. *Science* **343**, 287–290 (2014).

23. C. A. Oi, *et al.*, Dual effect of wasp queen pheromone in regulating insect sociality. *Curr.*
Biol. **25**, 1638–1640 (2015).

24. C. A. Oi, J. G. Millar, J. S. van Zweden, T. Wenseleers, Conservation of Queen
Pheromones Across Two Species of Vespine Wasps. *J. Chem. Ecol.* **42**, 1175–1180
(2016).

25. T. M. Nunes, *et al.*, Queen signals in a stingless bee: suppression of worker ovary
activation and spatial distribution of active compounds. *Sci. Rep.* **4**, 7449 (2014).

26. T. Pankiw, *et al.*, Mandibular gland components of european and africanized honey bee
queens (Apis mellifera L.). *J. Chem. Ecol.* **22**, 605–615 (1996).

27. E. Plettner, K. N. Slessor, M. L. Winston, J. E. Oliver, Caste-Selective Pheromone
Biosynthesis in Honeybees. *Science* **271**, 1851–1853 (1996).

28. E. Plettner, K. N. Slessor, M. L. Winston, Biosynthesis of Mandibular Acids in Honey Bees
(Apis mellifera): De novo Synthesis, Route of Fatty Acid Hydroxylation and Caste Selective
β-Oxidation. *Insect Biochem. Mol. Biol.* **28**, 31–42 (1998).

29. Y. Le Conte, A. Mohammedi, G. E. Robinson, Primer effects of a brood pheromone on
honeybee behavioural development. *Proc. Biol. Sci.* **268**, 163–168 (2001).

30. H. B. G. S. Consortium, Pheromone communication in the honeybee (Apis mellifera L.). *J.*
Chem. Ecol. **31**, 2731–2745 (2005).

31. S. A. Princen, *et al.*, Honeybees possess a structurally diverse and functionally redundant
set of queen pheromones. *Proc. Biol. Sci.* **286**, 20190517 (2019).

32. K. Matsuura, *et al.*, Identification of a pheromone regulating caste differentiation in termites.
Proc. Natl. Acad. Sci. U. S. A. **107**, 12963–12968 (2010).

33. D. H. Collins, *et al.*, Gene expression during larval caste determination and differentiation in
intermediately eusocial bumblebees, and a comparative analysis with advanced eusocial
honeybees. *Mol. Ecol.* **30**, 718–735 (2021).

34. M. Orlova, J. Starkey, E. Amsalem, A small family business: synergistic and additive effects
of the queen and the brood on worker reproduction in a primitively eusocial bee. *J. Exp.*
Biol. **223** (2020).

35. J. Starkey, A. Brown, E. Amsalem, The road to sociality: brood regulation of worker
reproduction in the simple eusocial bee Bombus impatiens. *Anim. Behav.* **154**, 57–65
(2019).

36. Y. Ulrich, D. Burns, R. Libbrecht, D. J. C. Kronauer, Ant larvae regulate worker foraging
behavior and ovarian activity in a dose-dependent manner. *Behav. Ecol. Sociobiol.* **70**,
1011–1018 (2016).

37. A. Maisonnasse, *et al.*, New insights into honey bee (Apis mellifera) pheromone
communication. Is the queen mandibular pheromone alone in colony regulation? *Front.*
Zool. **7**, 18 (2010).

38. J. R. Dampney, A. B. Barron, B. P. Oldroyd, Policing of adult honey bees with activated
ovaries is error prone. *Insectes Soc.* **49**, 270–274 (2002).

39. M. J. West-Eberhard, Phenotypic Plasticity and the Origins of Diversity. *Annu. Rev. Ecol.*
Syst. **20**, 249–278 (1989).

40. R. Gadagkar, R. Gadagkar, *The Social Biology of Ropalidia marginata: toward*
understanding the evolution of eusociality (Harvard University Press, 2009).

41. S. H. Woodard, *et al.*, Genes involved in convergent evolution of eusociality in bees. *Proc.*
Natl. Acad. Sci. U. S. A. **108**, 7472–7477 (2011).

42. A. J. Berens, J. H. Hunt, A. L. Toth, Comparative transcriptomics of convergent evolution:
different genes but conserved pathways underlie caste phenotypes across lineages of
eusocial insects. *Mol. Biol. Evol.* **32**, 690–703 (2015).

43. M. R. Warner, L. Qiu, M. J. Holmes, A. S. Mikheyev, T. A. Linksvayer, Convergent eusocial
evolution is based on a shared reproductive groundplan plus lineage-specific plastic genes.
Nat. Commun. **10**, 2651 (2019).

44. L. Holman, H. Helanterä, K. Trontti, A. S. Mikheyev, Comparative transcriptomics of social
insect queen pheromones. *Nat. Commun.* **10**, 1593 (2019).

45. L. Holman, Queen pheromones and reproductive division of labor: a meta-analysis.
Behavioral Ecology **29**, 1199–1209 (2018).

46. C. F. Funaro, K. Böröczky, E. L. Vargo, C. Schal, Identification of a queen and king
recognition pheromone in the subterranean termite Reticulitermes flavipes. *Proceedings of*
the National Academy of Sciences **115**, 3888–3893 (2018).

47. K. M. Kapheim, *et al.*, Genomic signatures of evolutionary transitions from solitary to group
living. *Science* **348**, 1139–1143 (2015).

48. B. Fouks, *et al.*, The genomic basis of evolutionary differentiation among honey bees.
Genome Res. **31**, 1203–1215 (2021).

49. M. C. Harrison, *et al.*, Hemimetabolous genomes reveal molecular basis of termite

eusociality. *Nat Ecol Evol* **2**, 557–566 (2018).

50. M. Lynch, J. S. Conery, The evolutionary demography of duplicate genes. *J. Struct. Funct. Genomics* **3**, 35–44 (2003).

51. E. V. Koonin, Orthologs, paralogs, and evolutionary genomics. *Annu. Rev. Genet.* **39**, 309–338 (2005).

52. A. Kuzniar, R. C. H. J. van Ham, S. Pongor, J. A. M. Leunissen, The quest for orthologs: finding the corresponding gene across genomes. *Trends Genet.* **24**, 539–551 (2008).

53. T. Gabaldón, E. V. Koonin, Functional and evolutionary implications of gene orthology. *Nat. Rev. Genet.* **14**, 360–366 (2013).

54. P. Natsidis, P. Kapli, P. H. Schiffer, M. J. Telford, Systematic errors in orthology inference and their effects on evolutionary analyses. *iScience* **24**, 102110 (2021).

55. B. Misof, *et al.*, Phylogenomics resolves the timing and pattern of insect evolution. *Science* **346**, 763–767 (2014).

56. J. M. Ranz, J. Parsch, Newly evolved genes: moving from comparative genomics to functional studies in model systems. How important is genetic novelty for species adaptation and diversification? *Bioessays* **34**, 477–483 (2012).

57. A. L. Camiletti, G. J. Thompson, Drosophila As a Genetically Tractable Model for Social Insect Behavior. *Frontiers in Ecology and Evolution* **4** (2016).

58. K. R. Sieber, T. Dorman, N. Newell, H. Yan, (Epi)Genetic Mechanisms Underlying the Evolutionary Success of Eusocial Insects. *Insects* **12** (2021).

59. E. J. Duncan, O. Hyink, P. K. Dearden, Notch signalling mediates reproductive constraint in the adult worker honeybee. *Nat. Commun.* **7**, 12427 (2016).

60. M. D. Drapeau, S. Albert, R. Kucharski, C. Prusko, R. Maleszka, Evolution of the Yellow/Major Royal Jelly Protein family and the emergence of social behavior in honey bees. *Genome Res.* **16**, 1385–1394 (2006).

61. J. Li, *et al.*, Application of Weighted Gene Co-expression Network Analysis for Data from Paired Design. *Sci. Rep.* **8**, 622 (2018).

62. P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).

63. X. Qu, *et al.*, Weighted gene co expression network analysis (WGCNA) with key pathways and hub-genes related to micro RNAs in ischemic stroke. *IET Syst. Biol.* **15**, 93–100 (2021).

64. K. C. Galang, J. R. Croft, G. J. Thompson, A. Percival-Smith, Analysis of the Drosophila melanogaster anti-ovarian response to honey bee queen mandibular pheromone. *Insect Mol. Biol.* **28**, 99–111 (2019).

65. G. V. Amdam, K. Norberg, M. K. Fondrk, R. E. Page Jr, Reproductive ground plan may mediate colony-level selection effects on individual foraging behavior in honey bees. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 11350–11355 (2004).

655 66. K. M. Weiss, A. V. Buchanan, *Genetics and the Logic of Evolution* (John Wiley & Sons,
656   2004).

657 67. M. A. Nowak, C. E. Tarnita, E. O. Wilson, The evolution of eusociality. *Nature* **466**, 1057–
658   1062 (2010).

659 68. J. A. Timmons, K. J. Szkop, I. J. Gallagher, Multiple sources of bias confound functional
660   enrichment analysis of global -omics data. *Genome Biol.* **16**, 186 (2015).

661 69. F. Maleki, K. Ovens, D. J. Hogan, A. J. Kusalik, Gene Set Analysis: Challenges,
662   Opportunities, and Future Research. *Front. Genet.* **11**, 654 (2020).

663 70. J. Schmitzová, *et al.*, A family of major royal jelly proteins of the honeybee Apis mellifera L.
664   *Cell. Mol. Life Sci.* **54**, 1020–1030 (1998).

665 71. L. C. Ferguson, J. Green, A. Surridge, C. D. Jiggins, Evolution of the insect yellow gene
666   family. *Mol. Biol. Evol.* **28**, 257–272 (2011).

667 72. P. K. Dearden, Evo-Devo Lessons Learned from Honeybees. *Evolutionary Developmental*
668   *Biology* (2021) https:/doi.org/10.1007/978-3-319-32979-6.pdf#page=795.

669 73. K. M. Glastad, L. Ju, S. L. Berger, Tramtrack acts during late pupal development to direct
670   ant caste identity. *PLoS Genet.* **17**, e1009801 (2021).

671 74. J. Ge, Z. Ge, D. Zhu, X. Wang, Pheromonal Regulation of the Reproductive Division of
672   Labor in Social Insects. *Front Cell Dev Biol* **8**, 837 (2020).

673 75. M. R. Lovegrove, P. K. Dearden, E. J. Duncan, Honeybee queen mandibular pheromone
674   induces a starvation response in Drosophila melanogaster. *Insect Biochem. Mol. Biol.* **154**,
675   103908 (2023).

676 76. M. H. Pham-Delègue, J. Trouiller, C. M. Caillaud, B. Roger, C. Masson, Effect of queen
677   pheromone on worker bees of different ages: behavioural and electrophysiological
678   responses. *Apidologie* **24**, 267–281 (1993).

679 77. E. M. Litsey, S. Chung, J. D. Fine, The Behavioral Toxicity of Insect Growth Disruptors on
680   Apis mellifera Queen Care. *Frontiers in Ecology and Evolution* **9** (2021).

681 78. C. M. Grozinger, G. E. Robinson, Endocrine modulation of a pheromone-responsive gene
682   in the honey bee brain. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **193**,
683   461–470 (2007).

684 79. V. Vergoz, *et al.*, Peripheral modulation of worker bee responses to queen mandibular
685   pheromone. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 20930–20935 (2009).

686 80. T. L. Pritchett, E. A. Tanner, K. McCall, Cracking open cell death in the Drosophila ovary.
687   *Apoptosis* **14**, 969–979 (2009).

688 81. M. Buszczak, L. Cooley, Eggs to die for: cell death during Drosophila oogenesis. *Cell Death*
689   *Differ.* **7**, 1071–1074 (2000).

690 82. C. A. Oi, *et al.*, Do primitively eusocial wasps use queen pheromones to regulate
691   reproduction? A case study of the paper wasp Polistes satan. *Front. Ecol. Evol.* **7** (2019).

83. N. W. Saleh, *et al.*, Social Behavior, Ovary Size, and Population of Origin Influence
Cuticular Hydrocarbons in the Orchid Bee Euglossa dilemma. *Am. Nat.* **198**, E136–E151
(2021).

84. E. Treanore, N. Derstine, E. Amsalem, What Can Mechanisms Underlying Derived Traits
Tell Us About the Evolution of Social Behavior? *Ann. Entomol. Soc. Am.* **114**, 547–561
(2020).

85. C. Gomez-Diaz, F. Martin, J. M. Garcia-Fernandez, E. Alcorta, The Two Main Olfactory
Receptor Families in Drosophila, ORs and IRs: A Comparative Approach. *Front. Cell.*
Neurosci. **12**, 253 (2018).

86. P. Brand, S. R. Ramírez, The Evolutionary Dynamics of the Odorant Receptor Gene Family
in Corbiculate Bees. *Genome Biol. Evol.* **9**, 2023–2036 (2017).

87. S. K. McKenzie, I. Fetter-Pruneda, V. Ruta, D. J. C. Kronauer, Transcriptomics and
neuroanatomy of the clonal raider ant implicate an expanded clade of odorant receptors in
chemical communication. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 14091–14096 (2016).

88. N. Terrapon, *et al.*, Molecular traces of alternative social organization in a termite genome.
Nat. Commun. **5**, 3636 (2014).

89. S. Shigenobu, *et al.*, Genomic and transcriptomic analyses of the subterranean termite
Reticulitermes speratus: Gene duplication facilitates social evolution. *Proc. Natl. Acad. Sci.*
U. S. A. **119** (2022).

90. H. M. Robertson, K. W. Wanner, The chemoreceptor superfamily in the honey bee, Apis
mellifera: expansion of the odorant, but not gustatory, receptor family. *Genome Res.* **16**,
1395–1403 (2006).

91. M. Paoli, G. C. Galizia, Olfactory coding in honeybees. *Cell Tissue Res.* **383**, 35–58 (2021).

92. W. O. H. Hughes, B. P. Oldroyd, M. Beekman, F. L. W. Ratnieks, Ancestral monogamy
shows kin selection is key to the evolution of eusociality. *Science* **320**, 1213–1216 (2008).

93. C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data. *BMC*
Bioinformatics **18**, 529 (2017).

94. J. A. Denton, M. Velasque, F. A. Reed, Cross-species transcriptomic network analysis
reveals links between ribosomal protein mutation and cancer. *bioRxiv* (2020)
https:/doi.org/10.1101/2020.12.11.420604.

95. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).

96. S. Andrews, FastQC: a quality control tool for high throughput sequence data. *Babraham*
Bioinformatics (2010) (2018).

97. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

98. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).

99. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).

100.   A. Dusa, Venn: draw venn diagrams. *R package version* (2018).

101.   L. Wilkinson, *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2016).

102.   G. Csardi, T. Nepusz, Others, The igraph software package for complex network
research. *InterJournal, complex systems* **1695**, 1–9 (2006).

103.   B. Si, *et al.*, GGraph: An Efficient Structure-aware Approach for Iterative Graph
Processing. *IEEE Transactions on Big Data*, 1–1 (2020).

104.   G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. *OMICS* **16**, 284–287 (2012).

105.   S. Durinck, *et al.*, BioMart and Bioconductor: a powerful link between biological
databases and microarray data analysis. *Bioinformatics* **21**, 3439–3440 (2005).

106.   A. Alexa, J. Rahnenführer, Gene set enrichment analysis with topGO. *Bioconductor*
Improv **27**, 1–26 (2009).

107.   S. Draghici, *et al.*, Onto-Tools, the toolkit of the modern biologist: Onto-Express, Onto-
Compare, Onto-Design and Onto-Translate. *Nucleic Acids Res.* **31**, 3775–3781 (2003).

108.   T. Wu, *et al.*, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innovation (Camb) **2**, 100141 (2021).

109.   Z. Arendsee, *et al.*, Phylostratr: A framework for phylostratigraphy. *Bioinformatics* **35**,
3617–3627 (2019).

110.   M. Quint, *et al.*, A transcriptomic hourglass in plant embryogenesis. *Nature* **490**, 98–101
(2012).

111.   T. Domazet-Lošo, D. Tautz, A phylogenetically based transcriptome age index mirrors
ontogenetic divergence patterns. *Nature* **468**, 815–818 (2010).

112.   G. Yu, *Data Integration, Manipulation and Visualization of Phylogenetic Trees* (Chapman
& Hall/CRC Biostatistics, 2022).

113.   G. Yu, Using ggtree to Visualize Data on Tree-Like Structures. *Curr. Protoc.*
Bioinformatics **69**, e96 (2020).

114.   G. Yu, T. T.-Y. Lam, H. Zhu, Y. Guan, Two methods for mapping and visualizing
associated data on phylogeny using ggtree. *Mol. Biol. Evol.* **35**, 3041–3043 (2018).

115.   G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, Ggtree : An r package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

Supporting Text & Materials

Materials & Methods
RNA Extraction.
A modified TRIzol RNA extraction protocol used for this work.

Materials
- ● 1 ml TRIzol
- ● 200 µl RNase-free chloroform
- ● 1 µl Glycogen
- ● 1 ml Isopropanol 100% w/w – chilled to -80°C
- ● 3 ml Ethanol 75% w/w – prepared fresh
- ● 500 µl Back Extraction Buffer (BEB)
- ● 20 µl Nuclease-free water

Sample homogenization and RNA extraction
1. Flash freeze samples (samples must be kept at -80oC) and transfer to dry ice (for processing) or -80oC freezer.
2. Add 500µL of TRIzol to each Eppendorf tube (keep samples in dry ice)
3. Homogenise sample with a pestle
4. Add 500µL of TRIzol
5. Mix by pipetting samples
6. Let it sit at room temperature for 5 min
7. Add 200µL of RNase-free chloroform (make sure it is a new vial or that it is only used on RNA work)
8. Shake vigorously by hand for 15s DO NOT VORTEX SAMPLES (vortex degrades the DNA)
9. Let it sit at room temperature for 3 min
10. Centrifuge at 4oC < 12,000g for 15 min
11. Transfer the upper aqueous phase to a new tube (leave 8-10µL in the bottom of the Eppendorf tube to prevent TRIzol contamination from getting into the next phase
12. Transfer the upper aqueous phase to a new tube (leave 8-10µL in the bottom of the tube to prevent TRIzol contamination from getting into the next phase. DO NOT DISCARD LOWER PHASE
13. Keep both upper and lower phases in separate tubes and proceed to 1- RNA and siRNA purification (upper aqueous phase) or 2- DNA purification (lower phase)

RNA and siRNA purification (upper aqueous phase)
1. Add 1µL of glycogen (can be skipped)
2. Add 500µL of RNase-free 100% isopropanol/ethanol (chilled to -80oC)
3. Invert by hand 10-20 times to mix (or very gently with the pipet)
4. Move samples to the fridge and let them rest overnight at -80 fridge
5. Centrifuge at 4oC < 12,000g for 10 min to precipitate the RNA (move samples very carefully not to agitate the mix, if disturbed, leave them overnight again).
6. Remove the supernatant and discard the rest.
7. Add 1µL of 75% Ethanol to pellet

8. Centrifuge at <7,500g for 5 min at 4oC
9. Remove supernatant and discard
10. Add 1µL of 75% Ethanol to pellet
11. Centrifuge at <7,500g for 5 min at 4oC
12. Remove supernatant and discard
13. Pulse spin samples at room temperature
14. Carefully remove the remaining supernatant with a pipette without disturbing the RNA pellet (use
gel loading tips)
15. Air-dry the pellet at room temperature for 2-5 min to evaporate Ethanol
16. If necessary, heat tubes at 42°C to evaporate remaining Ethanol
17. Add 20µL of nuclease-free water to the pellet (over-dried RNA might take longer to dissolve)
18. Gentle mix the sample to solubilise the RNA
19. Place the solubilised RNA on ice immediately
20. Aliquot 8µL of the sample for quantification and cDNA synthesis (keep the remaining 12µL in -
80oC)
21. Quantify RNA concentration and purity.
830
831
832

833
834 **Figure S1. Log$_2$–fold change heatmap of the top 25 differentially expressed genes.** Differential gene
835 expression was determined using deseq. Both up- and down-regulated genes were considered. The
836 colour scale reflects the fold change in gene expression in pheromone compared to the controls, ranging
837 from down-regulated (blue) to up-regulated (red). Threshold of significance was defined as p<0.05.
838

Figure S2. Log$_2$–fold change heatmap of the top 25 differentially expressed genes. Differential gene expression was determined using WGCNA. Both up- and down-regulated genes were considered. The colour scale reflects the fold change in gene expression in pheromone compared to the controls, ranging from down-regulated (blue) to up-regulated (red). Threshold of significance was defined as p<0.05.

Figure S3. All pheromone-responsive modules preserved in cross-species comparison between ***Drosophila melanogaster*** **and** ***Apis mellifera*** **or** ***Bombus terrestris.*** The WGCNA module-preservation function facilitated comparisons between *Drosophila melanogaster* and *Apis mellifera* pheromone-response to *A. mellifera* queen pheromone, and *D. melanogaster* and *Bombus terrestris* pheromone-response to *B. terrestris* queen pheromone. Number indicates the number of genes associated with each module.

853 **Table S1**. *Drosophila melanogaster* **ovary size by pheromone treatment.** Flies were exposed to
854 treatment pheromone from egg until collection. Virgin female flies were collected and dissected five days
855 after emergence. SD denotes standard deviation.

Treatment	Replicates	Mean size (mm^2)	SD
Bumblebee	10	0.32	0.064
Termite	10	0.35	0.145
QMP	10	0.51	0.085
QMP and Brood	10	0.59	0.156
Brood	10	0.65	0.123
Control (2nd)	10	0.80	0.146
Control (1st)	10	0.88	0.136

856
857

Table S2. **Multiple Comparisons of Means: Tukey Contrasts for ovary sizes of each pheromone treatment.** Highlighted comparisons are statistically significantly different. All treatments are significantly different to the controls. The controls are not significantly different.

Group 1	Group 2	Estimate	p.Val
Termite	Bumblebee	0.027	1.000000
QMP	Bumblebee	0.189	0.014997
QMP and Brood	Bumblebee	0.270	0.000026
Brood	Bumblebee	0.328	0.000000
Control (2nd)	Bumblebee	0.483	0.000000
QMP	Termite	0.162	0.076808
QMP and Brood	Termite	0.244	0.000262
Brood	Termite	0.301	0.000002
Control (2nd)	Termite	0.456	0.000000
QMP and Brood	QMP	0.082	1.000000
Brood	QMP	0.139	0.262967
Control (1st)	QMP	0.371	0.000000
Brood	QMP and Brood	0.058	1.000000
Control (1st)	QMP and Brood	0.289	0.000005
Control (1st)	Brood	0.231	0.000823
Control (1st)	Control (2nd)	0.076	1.000000

862
863