Evolution with private resources reverses some changes

from long-term evolution with public resources

Katrina van Raay[1,2], Sergey Stolyar[2,3], Jordana Sevigny[1,2], Jeremy A. Draghi[4],

Richard E. Lenski[2,5], Christopher J. Marx[2,3,6,7], Benjamin Kerr[1,2], Luis Zaman[2,8]

1. Department of Biology, University of Washington, Seattle, WA;

2. BEACON Center for the Study of Evolution in Action, Michigan State University, East Lansing, MI;

3. Department of Biological Sciences, University of Idaho, Moscow, ID;

4. Department of Biological Sciences, Virginia Tech, Blacksburg, VA;

5. Department of Microbiology and Molecular Sciences; Michigan State University, East Lansing, MI;

6. Institute for Modeling Collaboration and Innovation, University of Idaho, Moscow, ID;

7. Institute for Bioinformatics and Evolutionary Studies, University of Idaho, Moscow, ID;

8. Department of Ecology and Evolutionary Biology, and Center for the Study of Complex Systems,

University of Michigan, Ann Arbor, MI.

Abstract

A population under selection to improve one trait may evolve a sub-optimal state for another trait due to tradeoffs and other evolutionary constraints. How this evolution affects the capacity of a population to adapt when conditions change to favor the second trait is an open question. We investigated this question using isolates from a lineage spanning 60,000 generations of the Long-Term Evolution Experiment (LTEE) with *Escherichia coli*, where cells have access to a shared pool of resources, and have evolved increased competitive ability and a concomitant reduction in numerical yield. Using media-in oil emulsions we shifted the focus of selection to numerical yield, where cells grew in isolated patches with private resources. We found that the time spent evolving under shared resources did not affect the ability to re-evolve toward higher numerical yield. The evolution of numerical yield commonly occurred through mutations in the phosphoenolpyruvate phosphotransferase system. These mutants exhibit slower uptake of glucose, making them poorer competitors for public resources, and produce smaller cells that release less carbon as overflow metabolites. Our results demonstrate that mutations that were not part of adaptation under one selective regime may enable access to ancestral phenotypes when selection changes to favor evolutionary reversion.

Keywords: evolution, tradeoffs, constraints, metabolism

Introduction

The history of life has been inextricably shaped by evolutionary constraints. Given phenotypic trade-offs, an organism cannot simultaneously optimize two traits with respect to fitness (Agrawal et al. 2010; Stearns 1989). Thus, a population under strong selection to improve one trait may evolve a sub-optimal state for another trait. More generally, pleiotropy means that mutations in one gene affect multiple traits simultaneously (Pavličev & Cheverud, 2015), and antagonistic pleiotropy occurs when mutations are beneficial for one trait but detrimental for another. While there is a rich literature on evolutionary tradeoffs (Beardmore et al. 2011; Gounand et al. 2016; Guillaume & Otto, 2012) and antagonistic pleiotropy (Rose, 1982; Rose, 1985), there has been less attention given to how the evolutionary history of a lineage under selection for one trait affects its subsequent trajectory when selection shifts to favor a second trait that trades off with the first (but see Ostrowski et al. 2015; Teotónio & Rose, 2000; Travisano & Lenski, 1996; Velicer, 1999).

To understand how previous adaptation might affect future evolution, let us imagine a population under long-term selection for one trait that trades off with a second trait (Figure 1a). For simplicity, we assume higher values of each trait correspond to higher fitness when under selection. Long-term improvement in the favored trait (trait 1 in Figure 1a) occurs along with concomitant decreases in the second trait (trait 2). If selection shifted to favoring trait 2, how would the duration of history evolving under selection for trait 1 affect future evolution (Figure 1b)? There are at least two plausible hypotheses about how that duration would influence the rate of adaptation. The "Mutational Access" hypothesis (Figure 1c) posits that populations that spent more time under selection for trait 1 have fewer available mutations that would improve fitness by increasing trait 2 (i.e., the "late descendant" compared to the "intermediate descendant" in Figure

1a). If so, then this would lead to a *lower* rate of adaptation for the late descendant than for the

intermediate one when trait 2 is favored. This mutational limitation could result from the necessity

of compensating for (or reverting) more mutational steps; alternatively, it might occur if mutations

increasing trait 2 become increasingly deleterious over time due to the entrenchment of trait 1

(Schank & Wimsatt, 2011; Shah et al. 2015). The "Strength of Selection" hypothesis (Figure 1d),

by contrast, suggests that the intermediate and late descendants have equal access to mutations

improving trait 2, but the selective benefits for the late descendant are greater due to the relatively

larger changes in fitness upon improvement (Barrick et al. 2010; Chou et al. 2011). Under this

hypothesis, the rate of adaptation is predicted to be *higher* for the late descendant.

If adaptation is possible under selection for trait 2, then we expect a return towards the

progenitor's value of that trait. Does this phenotypic return occur by "rediscovering" the ancestral

genetic and mechanistic underpinnings of the second trait (Levin et al. 1997; Pennings et al. 2020)?

This rediscovery could occur by reverting key mutations, or by otherwise returning to the

regulatory, physiological, or metabolic states that enabled the progenitor's phenotype. Indeed, the

mechanisms that underly a return to the ancestral state (e.g., Figures 1b-d) might differ depending

on whether evolution is initiated with intermediate or late descendants.

83
Figure 1: Evolutionary trajectories and alternative hypotheses. a) A population evolves under direct selection for
one trait (trait 1, bolded axis). A higher value along either axis indicates an improvement in that trait under selection.
Each circle represents a genotype in an evolutionary line of descent. We focus on three genotypes along an
evolutionary trajectory: the progenitor, an intermediate descendent, and a late descendant. b) Selection is relaxed for
trait 1 and strengthened for trait 2 (bolded axis). The curved arrow represents the transition of a population with
descendent traits towards traits that resemble the progenitor when selection acts on trait 2. Whether, how fast, and by
what mechanisms this transition occurs are the main questions addressed in this paper. c) Mutational access
hypothesis. Intermediate (light green) and late (dark green) descendants are shown after selection is relaxed on trait 1
and strengthened on trait 2. The width of arrows pointing from each descendent towards the progenitor indicates the
rate of change. The thick arrow from the intermediate descendent denotes a faster return to the progenitor phenotype
than the dashed arrow from the late descendent. Under this hypothesis, the late descendent is either mutationally
further from the progenitor state or the mutational options to revert to the progenitor phenotype are more costly due
to entrenchment. d) Strength of selection hypothesis. Intermediate (light green) and late (dark green) descendants are
shown when selection flips from favoring trait 1 to trait 2. The width of arrows pointing from each descendent towards
the progenitor indicates the relative strength of selection. The thick arrow from the late descendent indicates stronger
selection favoring a mutation that restores the ancestral phenotype, and thus a faster response, than for the same
mutation in the intermediate descendant (dashed arrow). Under this hypothesis, both descendants have the same
mutational access to various positions in phenotype space.
102
103

Here we present a realization of the thought experiment in Figure 1, in which we test how

the length of time that populations of *Escherichia coli* had been selected for faster growth affects

their future adaptation in a novel environment that favors increased numerical yield (i.e., number

of cells per concentration of substrate added). To do so, we take advantage of the Long-Term

Evolution Experiment (LTEE), in which 12 replicate populations of *E. coli* have been evolving for

over 70,000 generations in a relatively simple environment (Lenski 2017; Vasi et al. 1994; Wiser

et al. 2013). Periodic samples from each line have been frozen in suspended animation, so that the

"progenitor," "intermediate descendant," and "late descendant" are available by reviving samples

from this living fossil record. The well-stirred, unstructured environment maintains a single, public

pool of resources for the population and thus favors rapid growth, above all else (Vasi et al. 1994,

Novak et al. 2006). Furthermore, there is no expectation of direct selection for either numerical

yield or metabolic yield (proportion of carbon in the substrate that is incorporated into biomass).

In an unstructured environment, a mutant that consumes resources more slowly but efficiently is

disadvantaged, because a faster-growing but less-efficient competitor can deplete the shared

resources in the meantime. Instead, a mutant that grows faster has an advantage in an unstructured

environment, even if it leads to a reduction in the population's numerical yield, metabolic yield,

or both.

The outcomes in the LTEE are slightly more complicated, and they are mediated in part by

changes to a third trait: cell size (Mongold & Lenski 1996). All of the LTEE lineages show

increases in competitive ability relative to their common ancestor, while simultaneously having

lower numerical yields (Vasi et al. 1994, Wiser et al. 2013). The reductions in numerical yield

result from the evolution of much larger individual cells (Lenski & Travisano 1994, Vasi et al.

1994, Grant et al. 2021). In many bacteria, increased cell size is a physiological response to

increased growth rate (Johnston et al. 1979; Pierucci, 1978). In addition, some mutations in

bacteria affect both growth and cell size or shape (Monds et al. 2014, Yulo & Hendrickson 2019).

Therefore, it is reasonable to expect that selection for faster growth has led to larger cells and, as

a pleiotropic consequence, decreased numerical yield.

These evolutionary changes also extend to metabolic yield—the efficiency with which

carbon sources are converted into biomass. During exponential growth on glucose, when most of

the doublings occur during the LTEE daily cycle, direct measurements of metabolic yield through

central metabolism show a small (but significant) decrease in the evolved populations. The

excretion of acetate, an overflow metabolite, also increased by about 50% in the LTEE (Harcombe

et al. 2013). Similar to the case with cell size, some loci with beneficial mutations in the LTEE

lineages are known to affect acetate metabolism (Quandt et al. 2015); it is also known that rapid

growth *per se* leads to increased overflow metabolite production in *E. coli* (Barrick & Lenski 2009;

Farmer & Jones 1976). This situation is further complicated, however, by the finding that the LTEE

populations have evolved increased total biomass when measured after a full 24-hour growth cycle

(Lenski & Mongold 2000, Novak et al. 2006). While the increase might be caused in part by a

change in biomass composition, it also reflects an increased ability to use excreted metabolites

after the glucose has been depleted. Indeed, the evolved LTEE strains have increased their ability

to grow on acetate, as well as to grow on other overflow metabolites that the LTEE ancestor could

not use (Leiby & Marx, 2014). Although there was no direct advantage to increased biomass in a

well-mixed environment, nonetheless biomass increased alongside faster growth in the LTEE. This

constellation of changes can be understood by considering that, in the novel environment of the

LTEE, the ancestral strain was likely far from the trade-off front between growth rate and total

biomass, leaving room for improvements in both trats (Novak et al. 2006).

Now consider what might happen if the bacteria that previously evolved in the LTEE

moved to and evolved in a structured environment that was otherwise similar to the unstructured

LTEE environment. By distributing a transplanted population into many isolated 'patches,' each

seeded by a single cell, any resources saved by growing more efficiently would accrue

disproportionately to cells with the same genotype, which would effectively eliminate resource

competition. Now imagine that the cells in these patches were periodically pooled, diluted

distributed as single founders over a new set of empty patches. In that case, selection would favor

increased numerical yield since each individual cell is a potential propagule able to establish a new

subpopulation. Increased numerical yield could be achieved by increasing the metabolic yield, by

producing smaller individual cells, or perhaps by some combination of these changes. In this way,

selection shifts to favor numerical yield over competitive ability for shared resources

(accomplishing the transformation from Figure 1a to Figure 1b in our thought experiment).

In our study, we achieve this shift by propagating cells in water-in-oil emulsions comprised

of millions of aqueous media-filled droplets surrounded by an oil phase (Figure 2) (Bachmann et

al. 2013). If a population of cells is diluted sufficiently before creation of the emulsion, then each

cell will usually be the sole occupant of a droplet. In such a case, barring mutation, there is no

competition between genotypes for resources, as each cell (and its progeny) has access to a private

resource supply demarcated by the droplet. If growth of isolated cells inside droplets proceeds for

sufficient time to exhaust the substrate and is followed by demulsification, dilution, and

redistribution into droplets across successive transfers (as in Figure 2a), then those genotypes that

produce *more* cells (rather than the fastest-growing cells) will have an advantage. Indeed, prior

experiments have demonstrated that various forms of isolation can favor numerical yield at the

expense of competitive ability in several different biological systems (Bachmann et al. 2013;

Eshelman et al. 2010; Kerr et al. 2006; van Tatenhove-Pel et al. 2021).

Here, we instantiate the thought experiment of Figure 1 by considering two bacterial traits:

competitive ability (the relative increase in proportion of a focal strain when competing with

another strain for shared resources) and numerical yield (the absolute number of cells produced by

a focal strain when resources are not shared). We focus on a single LTEE lineage in order to

explore how previous selection for one trait (competitive ability) affects the tempo and mode of

evolution under selection for a second trait (numerical yield) in the presence of a phenotypic

tradeoff. We probe how the length of time adapting to the well-mixed LTEE regime affects the

ability of a population to adapt under the new emulsion condition. We examine whether increases

in numerical yield during growth with private resources come with associated decreases in

competitive ability for shared resources, and whether they are achieved through changes in

metabolic yield and associated overflow metabolite production, decreases in cell size, or by both

factors. Finally, by determining the nature and number of mutations that occurred, we can

investigate whether any evolutionary reversions to the ancestral phenotypes were caused by

changes in the same loci/processes that were implicated in the evolution of fast growth in the

original LTEE environment.

Methods

Strains

All experiments were founded with *E. coli* B isolates that came from the Ara–5 lineage in the

LTEE. The isolates used in our experiment came from 0 generations (the founder of the LTEE),

20,000 generations, and 60,000 generations of evolution in the LTEE (Table S1; hereafter called

0K, 20K, and 60K, respectively). The 0K isolate is the ultimate progenitor of the 20K and 60K

isolates in the context of the LTEE. However, because we conduct evolution experiments in the

emulsion system with these three strains, we will refer to these LTEE isolates as the "0K emulsion

ancestor", the "20K emulsion ancestor", and the "60K emulsion ancestor".

Emulsion evolution experiment

Following the pioneering work of Bachmann et al. (2013), we propagated three replicate

cultures founded by each of the three LTEE isolates (the emulsion ancestors) under selection for

increased numerical yield by using water-in-oil emulsions for 100 transfers. This method (see

below) creates millions of media-filled, picoliter-sized droplets surrounded by an oil phase, in

which bacterial cells in one droplet are isolated from those in others (Figure 2). Emulsions were

set up in 1.5-ml Eppendorf conical microcentrifuge tubes, and they were incubated at 37 °C

without shaking.

In order to select for increased numerical yield, we diluted the bacterial culture before

setting up the emulsions, such that 90% of occupied droplets would contain only 1 cell at the

beginning of each transfer cycle. We subsequently showed that occupied droplets had an average

of 1.12 cells (see Supplementary section *Calculating Droplet Statistics*). Bacteria were allowed to

grow and divide in droplets for 24 hours, after which the droplets were broken with 1*H*,1*H*,2*H*,2*H*-

perfluoro-1-octanol, releasing all the cells into a common pool. A fraction of the cells from this

pool were immediately used to initiate the next transfer (see Figure 2a). Because of the level of

dilution before emulsification, lineages from this treatment have evolved in "private droplets" (see

Figure 2c). This treatment was continued for 100 transfers.

Figure 2: a) Schematic of transfer protocol: 1. A mixture of oil and surfactant is added to a diluted culture of bacteria in growth media and vortexed to form millions of droplets, 2. Bacteria in the emulsion are incubated to allow subpopulation growth within the droplets, 3. After overnight growth, a demulsifier is added to break the emulsion and resuspend the bacteria, 4. The free cells are diluted into fresh media and the cycle is repeated, **b)** Micrograph of bacteria growing in emulsion droplets, **c)** Schematic of inoculation and growth for private droplets, **d)** Schematic of inoculation and growth for shared droplets. For parts c and d, the two colors of bacterial cells represent distinct genotypes.

To control for any effects of growing in emulsions, we also propagated three replicate cultures founded by each LTEE isolate for 100 transfers under higher starting-density conditions, in which each occupied droplet had almost 3 cells on average (see Supplementary section

Calculating Droplet Statistics). We predicted that we would not observe evolution towards

increased numerical yield in this "shared-droplet" treatment (see Figure 2d).

During the evolution experiment, the cell density of freshly broken emulsions was

measured with a FilterMax F5 Multi-Mode Microplate Reader (Molecular Devices) after 24 hours.

Briefly, we pipetted 150 µl of overnight culture from a broken emulsion into a flat-bottom 96-well

microtiter plate, and we recorded an OD_{595} reading. This reading was then used to estimate cell

density using a calibration curve giving the relationship between OD_{595} and CFU/ml, which was

established in prior experiments. Cultures were then diluted to 2×10^6 CFU/ml for the private

droplet treatment and 5×10^7 CFU/ml for the shared droplet treatment. Emulsions were created

by adding 200 µl of a mixture of 9 parts Novec 7500 HFE to 1 part Pico-surf (5% (w/w)) to 300

µl of diluted culture in the growth medium DM1000 in 1.5 ml Eppendorf tubes, and vortexing at

maximum speed for 2 minutes. DM1000 contains 1000 mg/l of glucose, and it was used instead

of the DM25 (25 mg/l glucose) used in the LTEE because DM1000 exhibited clearer differences

between the 0K and 60K samples in terms of numerical yield and competitive ability than DM25

in pilot experiments.

Single-colony isolates were picked from each population at the end of our evolution

experiment (after transfer 100) and frozen at -80 °C as 20% v/v glycerol stocks. We refer to these

evolved isolates as "emulsion descendants." Two loci previously shown to distinguish the 0K,

20K, and 60K ancestors were Sanger sequenced for each isolate to confirm the intended derivation

of the isolates. We detected within-experiment contamination via Sanger and subsequent whole-

genome sequencing in 4 of the 18 emulsion descendants, so these populations were dropped from

our analysis. The complete genomes of the remaining 14 strains (8 private droplet, 6 shared

droplet) were sequenced and analyzed, and the 6 isolates of private-droplet descendants that had

mutations were used for further analysis. For the shared droplet treatment, one descendent was

randomly chosen from each lineage to be assessed in the phenotypic assays (3 isolates).

Competition assays

257   Using the frozen samples of the emulsion ancestors (0K, 20K, and 60K) and their emulsion

258   descendants from the private and shared droplet treatments, we assessed the relative competitive

259   abilities of all of our strains. The emulsion descendants competed against a common marked strain

260   (equivalent to our 0K emulsion ancestor) in shared droplet conditions. We used a neutral marker

261   (a point mutation in the *araA* gene) to distinguish the focal and common competitor strains on

262   tetrazolium-arabinose indicator agar plates.

263   Competitions ran for 3 transfer cycles in emulsions. Competitions were initiated with a 1:1

264   ratio of the focal strain (denoted F) to the common competitor (denoted C). The competitive ability

265   of the focal strain relative to the common competitor ($w(F, C)$) was assessed by calculating a ratio

266   of their realized Malthusian growth rates (Lenski et al. 1991):

267
$$w(F, C) = \frac{\ln \frac{F_{\text{final}}}{F_{\text{initial}}} + \sum_{i=2}^{3} \ln z_i}{\ln \frac{C_{\text{final}}}{C_{\text{initial}}} + \sum_{i=2}^{3} \ln z_i},$$

268   where X_{initial} and X_{final} are the initial (beginning of the first transfer) and final (end of the third

269   transfer) densities, respectively, of strain X (with $X \in \{F, C\}$) and z_i is the factor by which the

270   population is diluted in order to initiate the i^{th} transfer.

272   *Genome Sequencing*

273   Single-colony isolates from 14 evolved lines and their 3 ancestors were grown in DM1000

and frozen as 20% v/v glycerol stocks at -80 °C. For genome sequencing, cells from these frozen

stocks were reanimated in 5 ml of LB (lysogeny broth). Genomic DNA was extracted using a

Qiagen DNEasy kit with 1 ml of overnight culture, following the manufacture's protocol. Library

preparation and sequencing for 16 strains was done at the Microbial Genome Sequencing Center

(MiGS) at the University of Pittsburgh. One strain (the 0K emulsion ancestor) was sequenced at

the University of Washington on Illumina's NextSeq platform and prepped with Illumina Nextera

barcodes. Genome sequences had an average of 131X coverage.

Mutations (predicted mutations, unassigned missing coverage, and unassigned new

junctions) were called using breseq (Version 0.33.2) (Deatherage & Barrick, 2014) with default

parameters. All mutations were confirmed using Integrative Genomics Viewer (Robinson et al.

2011), and four point mutations identified in strains that differed phenotypically from their

ancestors in cell size or numerical yield were Sanger sequenced for confirmation (Table S4).

Numerical yield and cell size assays

Populations of each emulsion ancestor and descendant were grown under the shared-

droplet conditions over a standard 24-hour growth cycle. We used the shared-droplet emulsions

instead of the well-stirred conditions to control for the microenvironmental effects of growing in

an emulsion (e.g., oxygen depletion during growth). The final cell number in the 24-hour sample

was used as our measure of numerical yield. Cell size was measured as the median cell volume

(μm^3) of several thousand cells using a Beckman Coulter MSE 4 instrument.

Metabolic analysis

Supernatants from the yield assays were filtered and stored at -80 °C prior to metabolic

analysis. After thawing, samples were analyzed using High Performance Liquid Chromatography

(HPLC) on a 20A chromatographic system (Shimadzu) equipped with diode array and refractive

index detectors. Samples were eluted with 6 mM H_2SO_4 at a flow rate of 0.6 ml/min from an

Aminex HPX 87H organic acid analysis column (300 by 7.8 mm) (BioRad). We recorded the areas

for all noticeable peaks associated with detected metabolites (glucose, citrate, acetate, formate, and

lactate), and these values were converted to metabolite concentrations (in mM) using organic acid

and sugar standards (from BioRad or made in-house). Citrate, an iron chelator present in DM

medium, is neither produced nor consumed by these bacterial strains, and it has been shown not to

affect the growth of the Ara–5 lineage (Leiby et al. 2012). Thus, citrate was used as an internal

control.

Analysis of metabolic data

The rates of production for each metabolite are difficult to compare because of differences

in the growth rates of the strains. To minimize this potentially confounding factor, we performed

linear regression of the concentration of each metabolite against the remaining concentration of

glucose. The magnitude of the slope of this relationship estimates the ratio of metabolite

produced for each unit of glucose consumed, assuming an approximate equilibrium between the

uptake of glucose and the release of metabolites into the medium. As glucose levels drop, however,

there is increased uncertainty in the glucose measurements; moreover, as the glucose becomes

scarce, the cells may begin to consume the excreted metabolites, which would bias our estimates

of the production rates. To avoid these problems, we excluded time-points in the linear regressions

for which the mean glucose concentration was below 0.8 mM. We chose this threshold because,

at lower concentrations, growth curves showed clear signs of departure from exponential growth.

We also excluded the measurements made at t=0 and, instead, we constrained the x-intercept of

the regressions to reflect the known initial concentrations in the media (i.e., 6 mM glucose and

none of the overflow metabolites). To estimate how quickly the glucose was consumed, we

performed regressions of glucose concentrations against time, using the same criteria for data

exclusion. This approach provided a measure of growth rate that was insensitive to differences in

cell size across strains.

Results & Discussion

Private-droplet treatment selected for high numerical yield

Our private droplet treatment was designed to minimize local competition between

genotypes, thereby focusing selection upon increased numerical yield. To assess whether the

emulsion-evolved populations underwent changes in numerical yield, we measured their cell

densities, along with those of their ancestors, at the end of 24 hours of growth in the emulsion

system. Providing some reassurance that salient features of the LTEE environment are maintained

in our emulsion cultures, the decreases in numerical yield seen during the LTEE were also detected

in emulsion -- the 20K and 60K ancestors had decreased numerical yield compared to the 0K

ancestor (Figure 3, Table S3; $p < 0.01$, unpaired one-tailed t-tests). All 6 private droplet

descendants had significantly increased numerical yield (cell number) compared with their direct

ancestor (Figure 3, Table S2; BK 433: $p < 0.05$; BK 431, BK 437, BK 438, BK 441, BK 442: $p <$

0.01, unpaired two-tailed t-tests). Our shared-droplet treatment served as control for the private-

droplet treatment, allowing local competition between genotypes, while controlling for growth in

the emulsion environment. Two of the three shared-droplet descendants had increased numerical

yield and one had decreased yield; however, the gains were less than those in the private-droplet

treatment, and all three changes were non-significant. Combining the 0K, 20K, and 60K lines, the

private-droplet descendants were significantly more productive than the shared-droplet

descendants (p=0.039, paired one-tailed t-test; see Supplementary section *Statistical Analysis*).

The two evolutionary treatments had different effective population sizes. However, the weaker

numerical-yield response in the shared-droplet treatment relative to the private-droplet treatment

cannot be explained by the lack of mutational opportunity, because the number of cell divisions

per transfer is actually greater in the shared-droplet treatment (see Supplementary section

Calculating Number of Cell Divisions in Droplets). Taken together, these data confirm our

prediction that the private-droplet treatment exerts stronger selection for increased numerical yield

than does the shared-droplet treatment.

Figure 3: Numerical yield (cell density after the standard 24-hour growth cycle) compared to relative competitive ability, both measured in the emulsion system. Each panel shows strains originating from a single emulsion ancestor (in grey, at right) from the Ara−5 lineage of the LTEE (0, 20,000, and 60,000 generations). Colors indicate emulsion evolution treatments (labels at top). The coordinate positions of each point are the averages of three replicate assays for each phenotypic trait. Error bars show SEM; when they are not visible along either axis, the corresponding SEM is smaller than the symbol itself. Dashed lines show the mean competitive ability and numerical yield of the relevant ancestor.

In particular, the shared-droplet results demonstrate that the increase in numerical yield in

the private-droplet treatment was not simply an evolutionary response to propagation under the

emulsion conditions. Instead, these results imply that the initial density within a droplet affects the

evolutionary outcome. By reducing competition between genotypes within the private droplets,

selection has favored numerical yield.

Shared droplet treatment favored competitive ability

When resources are shared, selection for greater competitive ability is expected. To assess

competitive ability, a focal strain was paired with a marked version of the 0K ancestor and

proportions were tracked under high density emulsion conditions such that resources were shared

between competitors. Again, emulsion populations showed consistent results with the batch culture

environment used during the LTEE -- the competitive ability of the 20K ancestor was elevated

compared to the 0K ancestor, and the 60K ancestor had a further increase over the 20K ancestor

(Figure 3,Table S3; $p < 0.05$ unpaired, one-tailed t-tests). Competitive ability increased in all three

shared-droplet descendants that we tested (Figure 3). However, the change was significant relative

to the corresponding emulsion ancestor only in the case of the line derived from the 0K ancestor

(Table S2). Competitive ability also increased in three of the six private-droplet descendants, but

none of these increases were significant, and the magnitudes of the increases were smaller than

those observed in the shared-droplet treatment. Competitive ability decreased in the other three

private-droplet descendants, and the decline was significant in one case (BK 441: $p < 0.01$,

unpaired two-tailed t-test). Combining the results from the 0K, 20K, and 60K lines, the shared-

droplet descendants were significantly more competitive than the private-droplet descendants ($p = $

0.038, paired one-tailed t-test; see Supplementary section *Statistical Analysis*). Overall, these

results support our prediction that starting each droplet with multiple cells favors the evolution of

increased competitive ability.

Tradeoff between competitive ability and numerical yield

If competitive ability and numerical yield strictly traded off, then competitive ability would

decrease in the private-droplet treatment as numerical yield increased, and numerical yield would

decrease in the shared-droplet treatment as competitive ability increased. However, we did not see

such a strict tradeoff. Although the private-droplet descendants had larger gains in numerical yield,

some shared-droplet descendants also experienced moderate gains in numerical yield. Similarly,

although the shared-droplet descendants exhibited larger gains in competitive ability, some

private-droplet descendants also showed moderate improvements in their competitive ability.

Thus, these two traits do not strictly tradeoff with one another, and instead there seems to be some

misalignment, in which larger gains in one trait sometimes occur with more modest gains in the

other trait.

To develop this notion of misalignment further, we can contrast the case of a strict tradeoff,

or pure antagonistic pleiotropy (Figure 4a), with two different scenarios of synergistic pleiotropy.

The first synergistic scenario involves strict alignment, such that a mutation that is most beneficial

for trait 1 is also the most beneficial for trait 2 (Figure 4b). In essence, this scenario is the opposite

of strict antagonistic pleiotropy. The second form of synergism involves misalignment. In this

scenario, all mutations simultaneously improve both traits, but the mutations that are the strongest

for trait 1 are the weakest for trait 2, and *vice versa* (Figure 4c). Misalignment occurs when the

relative effect of a mutation on one trait is the opposite of its relative effect on another trait.

Therefore, antagonistic pleiotropy (Figure 4a) is also a case of misalignment, although the term is

broader than the traditional definition of "antagonism" (as illustrated in Figure 4c). In the

Supplementary section, we develop a statistical test for misalignment. Using this test, we find that

the increases in numerical yield in the private-droplet treatment came with concomitant weaker

performance in competitive ability, while the improvements in competitive ability in the shared-

droplet treatment came with weaker performance in numerical yield. Despite the absence of a strict

tradeoff, these traits exhibited misalignment in our experiment. When two traits are misaligned,

independent evolution in environments that favor one trait or the other can produce a pattern that

supports a canonical tradeoff curve when the descendants are pooled across environments and the

ancestors are ignored (i.e., considering only the colored points, while ignoring the grey points, in

Figures 3 and 4c).

418
Figure 4: Three scenarios for pleiotropy with respect to two traits. The grey circle is the focal, ancestral genotype and
the colored circles show the phenotypes of evolved genotypes. Here, fitness improvement with regard to a trait is
taken to mean a *positive* change relative to the ancestor. (a) Antagonistic pleiotropy under a "strict tradeoff" situation,
in which an improvement in one trait leads to a decline in the other, and the magnitude of enhancement in one
dimension correlates with the magnitude of deterioration in the other. (b) Aligned synergistic pleiotropy describes a
situation where the magnitude of improvement in one trait correlates with the magnitude of improvement in the other.
(c) Misaligned synergistic pleiotropy describes a situation in which evolution improves both traits, but those genotypes
with the strongest improvement in one trait have the weakest improvement in the other.

Private-droplet treatment favored smaller cells

429    While performing microscopy to assess droplet diameter (Supplemental Figures S8 & S9),

we noticed variation in cell size across the emulsion treatments. The cells that had evolved in

private droplets appeared smaller than cells that had evolved in shared droplets. We therefore

decided to investigate this trait systematically. Figure 5 shows the relationship between cell size

and numerical yield.

Both cell size and total biomass increased in all 12 LTEE populations during selection for

increased competitive ability for shared resources despite the fact that neither is directly a target

of selection under those conditions (Grant et al. 2021, Lenski & Mongold, 2000). This increase in

cell size can be seen in our data for the three time points that we sampled from the Ara–5

population, by comparing the values of the three grey points along the x-axis in Figure 5. For the

emulsion evolved lines, the median cell volume decreased in all of the isolates from the private-

droplet treatment, and the changes were significant in three cases (BK 437, BK 438, BK 441: $p <$

0.01, unpaired, two-tailed t-test). Using an emulsion protocol similar to our private-droplet

treatment, Bachman *et al*. (2013) also reported observing an evolved decrease in cell size in

Lactococcus lactis. They noted that propagation in an emulsion system could favor such a change

because any mutant that distributed the same total amount of biomass into a greater number of

smaller cells would be able to colonize more droplets each transfer. Our shared-droplet treatment

provides an appropriate control for this inference. Indeed, the median cell size actually increased

slightly in all three replicates of this treatment, and the change was significant in one case (BK

443: $p < 0.01$, unpaired, two-tailed t-test). The opposing directional changes in our two treatments

indicates that neither the emulsion environment nor the transfer protocol (pooling all droplets and

transferring only a fraction of cells to the next growth cycle, as shown in Figure 2a) was responsible

for the consistently smaller cells seen in the private-droplet descendants. Instead, these findings

support the hypothesis that smaller cells evolved in response to the reduced inter-genotype

competition in the private-droplet treatment. When the 0K, 20K, and 60K lines are pooled, the

difference in cell size between the private-droplet and shared-droplet descendants is marginally

significant (p=0.051, paired one-tailed t-test; see Supplementary section *Statistical Analysis)*.

Figure 5: Median cell volume compared to numerical yield. Each panel shows strains originating from a single

emulsion ancestor (in grey, at right) from the Ara−5 lineage of the LTEE (0, 20,000, and 60,000 generations). Colors

indicate emulsion evolution treatments (labels at top). The coordinate positions of each point show the average of

three replicate assays for each phenotypic trait. Error bars show SEM; when they are not visible along either axis, the

corresponding SEM is smaller than the symbol itself. Dashed lines show the mean numerical yield and cell volume

of the relevant ancestor.

Genome sequencing reveals mutations in phosphoenolpyruvate phosphotransferase system

Given the evolution of smaller cells in the private-droplet treatment, we wondered if these

populations were reverting to the ancestral state of the LTEE. After all, the progenitor of the LTEE

(the 0K ancestor in our emulsion experiment) had very small cells. In particular, would the same

genes and metabolic pathways that changed in the LTEE and led to larger cells in that experiment

undergo reversion? Or would novel genetic and metabolic changes lead to a similar phenotypic

endpoint? Also, would these changes depend on the particular LTEE-derived ancestor used in the

emulsion experiment (e.g., 20K versus 60K)? To find out, we sequenced the complete genomes of

14 emulsion-evolved descendants and their three corresponding ancestors.

This whole-genome sequencing revealed a total number of 13 mutations that distinguish

the evolved descendants from their ancestors (Table S4). We saw no cases of identical mutations

in multiple descendants, nor was any gene mutated in descendants that evolved in the two different

treatments. However, three of the six descendants from the private-droplet treatment had point

mutations in genes encoding proteins in the phosphoenolpyruvate (PEP) phosphotransferase

system (PTS) (Table 1), which is involved in glucose uptake (Carmona et al. 2015; Escalante et

al. 2012; Nam et al. 2001; New et al. 2014; Notley-Mcrobb et al. 2006; Xia et al. 2017). All three

of these descendants also had increased numerical yield and reduced cell size (Table S2), and all

three were derived from the 20K and 60K emulsion ancestors. Of the three private-droplet

descendants without mutations in PTS genes, two derived from the 0K ancestor, which already

had high numerical yield in the emulsion conditions. The third private-droplet descendant without

mutations in PTS genes derived from the 60K ancestor, and it exhibited much smaller changes in

numerical yield and cell size than its counterpart with a PTS-associated mutation. The changes in

numerical yield and cell size were also small for both of the private-droplet descendants derived

from the 0K ancestor, suggesting that the high-yield phenotype of the LTEE progenitor was largely

maintained. Bachmann et al. (2013) also found a point mutation in a PTS gene in one strain that

evolved increased numerical yield in emulsions, suggesting that PTS might have some role in

modulating numerical yield.

The PTS-associated mutations occurred in *ptsH*, which encodes a phosphocarrier protein;

ptsG, which encodes a subunit of the glucose transporter; and *crp*, which encodes a global

transcriptional regulator that affects some steps in glucose uptake (Kimata et al. 1997). Given that

these mutations include an early stop codon and non-conservative changes in amino acids, it is

likely that some or all of them cause reductions or even losses of function. Mutations in *pykF,*

which encodes pyruvate kinase, were among the earliest and most repeatable genetic changes in

the LTEE (Woods et al. 2006; Tenaillon et al. 2016; Peng et al. 2018). These mutations presumably

reduce the conversion of PEP to pyruvate in central metabolism, which would leave more PEP

available to power the transport of PTS-dependent sugars. Indeed, distinct patterns of pleiotropy

were observed in the LTEE for various substrates that differed in terms of whether their uptake

requires the PTS (Travisano & Lenski, 1996). Thus, although the mutations selected in the private

droplet-treatment were not in the same genes as those responsible for phenotypic changes in the

LTEE, they appear to affect the same system.

STRAIN (BK#)	LTEE GEN	POSITION	REF	ALT	ANNOTATION	GENE	DESCRIPTION
437	20K	2462747	A	C	*86Y (TAA→TAC)	*ptsH*	phosphocarrier protein Hpr
437	20K	3414187	G	A	E55K (GAA→AAA)	*crp*	cAMP-activated global transcriptional regulator CRP
438	20K	1173797	G	A	G454S (GGT→AGT)	*ptsG*	PTS glucose transporter subunit IIBC
441	60K	2462629	T	C	L47P (CTG→CCG)	*ptsH*	phosphocarrier protein Hpr

Table 1: Mutations in private-droplet descendants associated with PTS.

Slower glucose uptake and decreased overflow metabolism in private-droplet descendants

Perhaps the private-droplet descendants achieved their higher numerical yield and smaller

cell size in a physiologically similar way to the LTEE progenitor, despite genetic changes that did

not merely reverse those that occurred during the LTEE. Because the mutations we found in the

PTS system would be predicted to affect glucose transport, we decided to directly examine whether

the private-droplet descendants had evolved toward a metabolic state, at least in terms of glucose

use and overflow metabolite production, that resembled the LTEE progenitor.

We examined the concentrations of both glucose and various excreted metabolites in spent

medium as the various bacterial strains grew in emulsions (Figure 6a). We observed that the

glucose concentration for the 20K and 60K private-droplet descendants decreased more slowly

than for the corresponding shared-droplet descendants, their ancestors, and the 0K descendants of

either the shared or private droplets (Figure 6a, glucose panel). The slower glucose consumption

was most pronounced in the private-droplet descendants that had mutations in PTS, indicating that

these mutations likely simultaneously reduce the rate of glucose utilization while increasing

numerical yield.

The primary metabolite that is excreted by *E. coli* growing on glucose, and by the LTEE

lines in particular, is acetate (Harcombe et al. 2013). Acetate is a byproduct of both glycolysis and

fermentation, which is a less efficient form of metabolism than respiration (Szenk et al. 2017). The

0K emulsion evolution decedents did not markedly differ in metabolic signatures from the LTEE

ancestor. However, we saw differences in acetate production between the private-droplet

descendants from 20K and 60K and their emulsion ancestors. Specifically, the private-droplet

descendants produced less acetate, which suggests they were more efficient at glucose metabolism

under these conditions than their ancestor. In contrast, some of the shared-droplet descendants

accumulated higher levels of acetate (Fig. 6a). In both cases, the acetate concentration eventually

declines as it is consumed by the cells after they have depleted the available glucose. Interestingly,

in emulsion-evolved populations of a different strain of *E. coli* (MG1655), Rabbers et al. (2021)

found an increased numerical yield was achieved with no change in acetate production. Those

results are consistent with our observations for the 0K descendants but differ from what we observe

in the 20K and 60K descendants, which suggests a history of rate adaptation may leave its mark

on future evolution.

In addition to acetate, other organic acids such as formate and lactate can be generated by

E. coli during growth on glucose. For all of the isolates in our study, either formate or lactate was

generated, but not both. For the 0K lineages, formate was produced by all types (ancestor, private-

droplet descendants, and shared-droplet descendant). For the 20K and 60K lineages, only the

shared-droplet descendant produced formate, whereas lactate was produced by the 20K and 60K

ancestors and their private-droplet descendants.

Differences between strains in the accumulated concentrations of excreted metabolites

might reflect different production rates, but they might also be influenced by differences in growth

rate, as well as subsequent consumption of those products. We therefore standardized these

measurements by computing the rate of production of each excreted metabolite per unit of glucose

consumed, and by excluding time points after the remaining glucose fell below a threshold

concentration (see *Methods—Analysis of metabolic data*). After so doing, Figure 6b shows a

positive relationship between the initial rate of glucose utilization and the fraction of this carbon

that is "wasted" as excreted metabolites in the 20K and 60K emulsion lines. Carbon waste was

calculated as a sum of the concentrations across all excreted metabolites concentrations, weighted

by the number of carbon atoms in each metabolite. The data indicate that the shared-droplet

descendants use the glucose rapidly, but in a relatively wasteful way, whereas the private-droplet

descendants consume glucose more slowly and efficiently.

We used these standardized rates (Table S5) to perform a principal components analysis

(PCA; Figure 6c) in order to visualize the differences between the emulsion ancestors and their

descendants. The metabolic changes between the 0K ancestor and its descendants were modest,

but the changes in the 20K and 60K lines were much greater. For the 20K and 60K ancestors, the

metabolic shifts in the private-droplet treatment were distinct from those in the shared-droplet

treatment, while the overall patterns of change were nearly identical for the 20K and 60K lines.

Thus, in addition to PTS-related mutations conferring high numerical yield for three of the four

private-droplet descendants of these two ancestors, the magnitude and nature of the metabolic

changes appear similar. One private-droplet descendant (derived from the 60K ancestor) exhibited

a smaller decrease in cell size and a more moderate increase in numerical yield (Figure 5). This

isolate also did not have mutations associated with PTS, and it differed somewhat less in its

metabolic profile from its ancestor. On balance, therefore, we lack clear support for either the

Mutational Access Hypothesis or the Strength of Selection Hypothesis (Figure 1). Nonetheless, it

is clear that just one or a few novel genetic changes can restore, at least approximately, the

ancestral phenotypes of slow growth, small cells, and high numerical yield, even after tens of

thousands of generations that led to faster growth, larger cells, and lower numerical yield.

Figure 6: Glucose consumption and metabolite production. a) The concentrations of glucose, acetate, formate and lactate over 10 hours. Each trajectory shows one strain, and each point along a trajectory is the average of up to three replicates per strain. Colors indicate emulsion evolution treatments (labels at top). b) Glucose utilization versus the fraction of carbon wasted. c) Principal components analysis (PCA) based on the strain-specific rates of excreted metabolites per unit glucose consumed. The 0K strains are shown as empty points, the 20K strains as filled points without borders, and the 60K strains as filled points with borders. Lines, colored by treatment, connect emulsion ancestors (grey points) to their descendants.

Summary

Three of the four private-droplet descendants of the 20K and 60K ancestors achieved or even surpassed the numerical yield of the ultimate LTEE progenitor (the 0K emulsion ancestor). When compared to their ancestors, these high-yield descendants showed several parallel changes: smaller cells, more efficient glucose metabolism, mutations in genes encoding the PTS, and lower levels of acetate production. None of these changes occurred in the descendants that evolved from the same ancestors under the shared-droplet treatment. While the increased numerical yield and

smaller cell size appeared to be reversions to the phenotypic state of the ultimate progenitor, the

genetic and metabolic underpinnings of the descendants' phenotypes were distinct from those of

the ultimate progenitor. Specifically, mutations in genes encoding the PTS were not responsible

for evolutionary improvement in growth rate during the LTEE, and the metabolic profiles of the

20K and 60K private-droplet descendants were distinct from that of the 0K emulsion ancestor (Fig.

6c). Therefore, these data indicate a novel mechanistic basis for phenotypic reversion in this

system. Specifically, mutations in the PTS likely involved a reduction in glucose uptake, thereby

lowering the growth rate (and leading to a reduced competitive ability). However, these same

mutations appear to lead to both smaller size and limited loss of carbon via overflow metabolites,

thereby improving numerical yield. Notably, the same kind of mutations produced the phenotypic

reversion for both the 20K and 60K emulsion lines.

Dollo's law of irreversibility states that "An organism never returns exactly to a former

state, even if it finds itself placed in conditions of existence identical to those in which it has

previously lived. But by virtue of the indestructibility of the past [...] it always keeps some trace

of the intermediate stages through which it has passed" (Gould, 1970). Our experiment certainly

did not return these organisms to their ancestral conditions. However, the private-droplet treatment

was designed to place populations under selection for an ancestral phenotype, namely higher yield.

Even as the private-droplet lineages derived from later generations of the LTEE recovered the high

numerical yield of the LTEE ancestor, their phenotypic "return" did not involve a straightforward

reversal of the steps in their evolutionary history. Rather, the mutational targets and metabolic

patterns underlying their return were distinct from those evolutionary paths explored in the LTEE.

By contrast, when under the same selection for increased numerical yield, the private-droplet

descendants of the 0K emulsion ancestor exhibited genetic and metabolic changes distinct from

the 20K and 60K emulsion lines, suggesting that changes during the first 20,000 generations of

the LTEE had altered the potential for subsequent adaptation to the private-droplet regime Such

contingency is a recurring feature of adaptation in biological systems (Blount et al. 2008; Card et

al. 2019). However, we found no evidence of contingency in the later generations, as both the

intermediate (20K) and late (60K) isolates from the LTEE readily recovered their ancestral form,

seemingly by taking similar steps.   Therefore, past evolution need not invariably alter future

evolution, even when the future involves a return to the past.

References

Agrawal, A. A., Conner, J. K., & Rasmann, S. (2010). Tradeoffs and Negative Correlationsin Evolutionary Ecology BT - Evolution Since Darwin. *Evolution Since Darwin*, (10), 243–268.

Bachmann, H., M. Fischlechner, I. Rabbers, N. Barfa, F. Branco dos Santos, D. Molenaar, and B. Teusink. 2013. "Availability of Public Goods Shapes the Evolution of Competing Metabolic Strategies." *Proceedings of the National Academy of Sciences* 110(35):14302–7. doi: 10.1073/pnas.1308523110.

Barrick, J E, and Lenski, R. E. "Genome-wide mutational diversity in an evolving population of Escherichia coli." *Cold Spring Harbor symposia on quantitative biology* vol. 74 (2009): 119-29. doi:10.1101/sqb.2009.74.018

Barrick, J. E., Kauth, M. R., Strelioff, C. C., & Lenski, R. E. (2010). Escherichia coli rpoB mutants have increased evolvability in proportion to their fitness defects. *Molecular Biology and Evolution*, *27*(6), 1338–1347. https://doi.org/10.1093/molbev/msq024

Beardmore, R. E., Gudelj, I., Lipson, D. A., & Hurst, L. D. (2011). Metabolic trade-offs and the maintenance of the fittest and the flattest. *Nature*, *472*(7343), 342–346. https://doi.org/10.1038/nature09905

Blount, Z. D., C. Z. Borland, and Lenski, R. E. 2008. Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proceedings of the National Academy of Sciences, USA* 105: 7899-7906.

Card, K. J., T. LaBar, J. B. Gomez, and Lenski, R. E. 2019. Historical contingency in the evolution of antibiotic resistance after decades of relaxed selection. *PLOS Biology* 17: e3000397.

Carmona, S. B., Moreno, F., Bolívar, F., Gosset, G., & Escalante, A. (2015). Inactivation of the PTS as a Strategy to Engineer the Production of Aromatic Metabolites in Escherichia coli. *Journal of Molecular Microbiology and Biotechnology*, *25*(2–3), 195–208. https://doi.org/10.1159/000380854

Chou, H. H., Chiu, H. C., Delaney, N. F., Segrè, D., & Marx, C. J. (2011). Diminishing returns epistasis among beneficial mutations decelerates adaptation. *Science*, *332*(6034), 1190–1192. https://doi.org/10.1126/science.1203799

Deatherage D.E., Barrick J. E. 2014. "Identification of Mutations in Laboratory-Evolved Microbes from Next-Generation Sequencing Data Using Breseq. In: Sun L., Shou W. (Eds) Engineering and Analyzing Multicellular Systems. Methods in Molecular Biology (Methods and Protocols)." 1151:1–22. doi: 10.1007/978-1-4939-0554-6.

Escalante, A., Cervantes, A. S., Gosset, G., & Bolívar, F. (2012). Current knowledge of the
Escherichia coli phosphoenolpyruvate-carbohydrate phosphotransferase system:
Peculiarities of regulation and impact on growth and product formation. *Applied*
Microbiology and Biotechnology, *94*(6), 1483–1494. https://doi.org/10.1007/s00253-012-
4101-5

Eshelman, C. M., Vouk, R., Stewart, J. L., Halsne, E., Lindsey, H. a, Schneider, S., … Kerr, B.
(2010). Unrestricted migration favours virulent pathogens in experimental metapopulations:
evolutionary genetics of a rapacious life history. *Philosophical Transactions of the Royal*
Society of London. Series B, Biological Sciences, *365*, 2503–2513.
https://doi.org/10.1098/rstb.2010.0066

Farmer, I. S., & Jones, C. W. (1976). The Energetics of Escherichia coli during Aerobic Growth
in Continuous Culture. *European Journal of Biochemistry*, *67*(1), 115–122.
https://doi.org/10.1111/j.1432-1033.1976.tb10639.

Gould, S. J. (1970). Dollo on Dollo ' s Law : Irreversibility and the Status of Evolutionary Laws
Author (s): Stephen Jay Gould Source : Journal of the History of Biology , Vol . 3 , No . 2
(Autumn , 1970), pp . 189-212 Published by : Springer Stable URL : http://www.jsto. *J*
Hist Biol., *3*(2), 189–212.
690
691   Gounand, I., Daufresne, T., Gravel, D., Bouvier, C., Bouvier, T., Combe, M., … Mouquet, N.
692   (2016). Size evolution in microorganisms masks trade-offs predicted by the growth rate
693   hypothesis. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1845), 20162272.
694   https://doi.org/10.1098/rspb.2016.2272
695
696   Grant, N. A., Abdel Magid, A., Franklin, J., Dufour, Y., & Lenski, R. E. (2021). Changes in Cell
697   Size and Shape During 50,000 Generations of Experimental Evolution with Escherichia
698   coli. *Journal of Bacteriology*, (March). https://doi.org/10.1128/jb.00469-20
699
700   Guillaume, F., & Otto, S. P. (2012). Gene functional trade-offs and the evolution of pleiotropy.
701   *Genetics*, *192*(4), 1389–1409. https://doi.org/10.1534/genetics.112.143214
702
Harcombe, W. R., Delaney, N. F., Leiby, N., Klitgord, N., & Marx, C. J. (2013). The Ability of
Flux Balance Analysis to Predict Evolution of Central Metabolism Scales with the Initial
Distance to the Optimum. *PLoS Computational Biology*, *9*(6).
https://doi.org/10.1371/journal.pcbi.1003091

Johnston, G. C., Ehrhardt, C. W., Lorincz, A., & Carter, B. L. A. (1979). Regulation of cell size
in the yeast Saccharomyces cerevisiae. *Journal of Bacteriology*, *137*(1), 1–5.
https://doi.org/10.1128/jb.137.1.1-5.1979

Kerr, B., Neuhauser, C., Bohannan, B. J. M., & Dean, A. M. (2006). Local migration promotes
competitive restraint in a host-pathogen "tragedy of the commons." *Nature*, *442*(7098), 75–
78. https://doi.org/10.1038/nature04864
713

714 Kimata, K., Takahashi, H., Inada, T., Postma, P., & Aiba, H. (1997). cAMP receptor protein-
715   cAMP plays a crucial role in glucose-lactose diauxie by activating the major glucose
716   transporter gene in Escherichia coli. *Proceedings of the National Academy of Sciences of*
717   *the United States of America*, *94*(24), 12914–12919.
718   https://doi.org/10.1073/pnas.94.24.12914
719
Leiby, N., Harcombe, W. R., & Marx, C. J. (2012). Multiple long-term, experimentally-evolved
populations of Escherichia coli acquire dependence upon citrate as an iron chelator for
optimal growth on glucose. *BMC Evolutionary Biology*, *12*(1).
https://doi.org/10.1186/1471-2148-12-151
Leiby, N., & Marx, C. J. (2014). Metabolic Erosion Primarily Through Mutation Accumulation,
and Not Tradeoffs, Drives Limited Evolution of Substrate Specificity in Escherichia coli.
PLoS Biology, *12*(2). https://doi.org/10.1371/journal.pbio.1001789
Lenski, R. E. 2017. "Experimental Evolution and the Dynamics of Adaptation and Genome
Evolution in Microbial Populations." *The ISME Journal* 1–14. doi: 10.1038/ismej.2017.69.
Lenski, R. E., and Travisano, M. 1994. "Dynamics of Adaptation and Diversification: A 10,000-
Generation Experiment with Bacterial Populations." *Proceedings of the National Academy*
of Sciences of the United States of America 91(15):6808–14. doi: 10.1073/pnas.91.15.6808.
732
733 Levin, B. R., Lipsitch, M., Perrot, V., Schrag, S., Antia, R., Simonsen, L., … Stewart, F. M.
734   (1997). The population genetics of antibiotic resistance. *Clinical Infectious Diseases : An*
735   *Official Publication of the Infectious Diseases Society of America*, *24 Suppl 1*, S9-16.
736   Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8994776
737
Manhart, M., Adkar, B. V., & Shakhnovich, E. I. (2016). *Tradeoffs between microbial growth*
phases lead to frequency-dependent and non-transitive selection.
https://doi.org/http://dx.doi.org/10.1101/096453
Monds, R. D., Lee, T. K., Colavin, A., Ursell, T., Quan, S., Cooper, T. F., & Huang, K. C.
(2014). Systematic Perturbation of Cytoskeletal Function Reveals a Linear Scaling
Relationship between Cell Geometry and Fitness. *Cell Reports*, *9*(4), 1528–1537.
https://doi.org/10.1016/j.celrep.2014.10.040
Mongold, J. A., and Lenski, R. E. 1996. Experimental rejection of a nonadaptive explanation for
increased cell size in *Escherichia coli*. *Journal of Bacteriology* 178: 5333-5334.
Nam, T. W., Cho, S. H., Shin, D., Kim, J. H., Jeong, J. Y., Lee, J. H., … Seok, Y. J. (2001). The
Escherichia coli glucose transporter enzyme IICBGlc recruits the global repressor Mlc.
EMBO Journal, *20*(3), 491–498. https://doi.org/10.1093/emboj/20.3.491
New, A. M., Cerulus, B., Govers, S. K., Perez-Samper, G., Zhu, B., Boogmans, S., …
Verstrepen, K. J. (2014). Different Levels of Catabolite Repression Optimize Growth in

Stable and Variable Environments. *PLoS Biology*, *12*(1), 17–20. https://doi.org/10.1371/journal.pbio.1001764

Notley-McRobb, L., Death, A., & Ferenci, T. (2006). The relationship between external glucose concentration and cAMP levels inside. *Biochemical Journal*, (1997), 1909–1918. https://doi.org/10.1099/00221287-143-6-1909

Novak, M., Pfeiffer, T., Lenski, R. E., Sauer, U., & Bonhoeffer, S. (2006). Experimental tests for an evolutionary trade-off between growth rate and yield in E. coli. *The American Naturalist*, *168*(2), 242–251. https://doi.org/10.1086/506527

Pavličev, M., & Cheverud, J. M. (2015). Constraints Evolve: Context Dependency of Gene Effects Allows Evolution of Pleiotropy. *Annual Review of Ecology, Evolution, and Systematics*, *46*(1), 413–434. https://doi.org/10.1146/annurev-ecolsys-120213-091721

Peng, F., Widmann, S., Wünsche, A., Duan, K., Donovan, K. A., Dobson, R. C. J., … Cooper, T. F. (2018). Effects of Beneficial Mutations in pykF Gene Vary over Time and across Replicate Populations in a Long-Term Experiment with Bacteria. *Molecular Biology and Evolution*, *35*(1), 202–210. https://doi.org/10.1093/molbev/msx279

Pennings, P. S., Ogbunugafor, C. B., & Hershberg, R. (2020). Reversion is most likely under high mutation supply, when compensatory mutations don't fully restore fitness costs. *BioRxiv*. https://doi.org/10.1101/2020.12.28.424568

Pfeiffer, T., Schuster, S., & Bonhoeffer, S. (2001). *Cooperation and Competition in the Evolution of ATP-Producing Pathways*. *292*(April).

Pierucci, O. (1978). Dimensions of Escherichia coli at various growth rates: model for envelope growth. *Journal of Bacteriology*, *135*(2), 559–574.

Ostrowski, E. A., Ofria, C., & Lenski, R. E. (2015). Genetically integrated traits and rugged adaptive landscapes in digital organisms Experimental evolution. *BMC Evolutionary Biology*, *15*(1), 1–14. https://doi.org/10.1186/s12862-015-0361-x

Quandt, E. M., Gollihar, J., Blount, Z. D., Ellington, A. D., Georgiou, G., & Barrick, J. E. (2015). Fine-tuning citrate synthase flux potentiates and refines metabolic innovation in the lenski evolution experiment. *ELife*, *4*(OCTOBER2015), 1–22. https://doi.org/10.7554/eLife.09696

Rabbers, I., Gottstein, W., Feist, A., Teusink, B., & Bruggeman, F. J. (2021). Selection for cell yield does not reduce overflow metabolism in E. coli. *BioRxiv*, 2021.05.24.445453. Retrieved from https://doi.org/10.1101/2021.05.24.445453

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., & Mesirov, J. P. (2011). Integrative Genomics Viewer. *Nature Biotechnology*, *29*(1), 24–26. https://doi.org/10.1038/nbt0111-24

789   Rose, M. (1982). *Antagonistic pleiotropy, dominance, and genetic variation*. 48*(2501).
790
791   Rose, M. R. (1985). Life history evolution with antagonistic pleiotropy and overlapping
792   generations. *Theoretical Population Biology*, *28*(3), 342–358. https://doi.org/10.1016/0040-
793   5809(85)90034-6
794
795   Schank, J., & Wimsatt, W. (2011). *Generative Entrenchment and Evolution Author (s): Jeffrey
796   C . Schank and William C . Wimsatt Source : PSA : Proceedings of the Biennial Meeting of
797   the Philosophy of Science Association , Vol . 1986 , Volume Two : Symposia and Invited
798   Papers (1986), pp. 1986*(1986), 33–60.
799
800   Shah, P., McCandlish, D. M., & Plotkin, J. B. (2015). Contingency and entrenchment in protein
801   evolution under purifying selection. *Proceedings of the National Academy of Sciences of the
802   United States of America*, *112*(25), E3226–E3235. https://doi.org/10.1073/pnas.1412933112
803
804   Stearns, S. C. 1989. "Trade-Offs in Life-History Evolution Published by: British Ecological
805   Society Stable URL: Https://Www.Jstor.Org/Stable/2389364 Trade-Offs in Life-History
806   Evolution." *Functional Ecology* 3(3):259–68.
807
Stevenson, K., McVey, A. F., Clark, I. B. N., Swain, P. S., & Pilizota, T. (2016). General
calibration of microbial growth in microplate readers. *Scientific Reports*, *6*(December), 4–
10. https://doi.org/10.1038/srep38828
Szenk, M., Dill, K. A., & de Graff, A. M. R. (2017). Why Do Fast-Growing Bacteria Enter
Overflow Metabolism? Testing the Membrane Real Estate Hypothesis. *Cell Systems*, *5*(2),
95–104. https://doi.org/10.1016/j.cels.2017.06.005
Tenaillon, O., J. E. Barrick, N. Ribeck, D. E. Deatherage, J. L. Blanchard, A. Dasgupta, G. C.
Wu, S. Wielgoss, S. Cruveiller, C. Médigue, D. Schneider, and R. E. Lenski. 2016. Tempo
and mode of genome evolution in a 50,000-generation experiment. *Nature* 536: 165-170.
Teotónio, H., & Rose, M. R. (2000). Variation in the reversibility of evolution. *Nature*,
408(6811), 463–466. https://doi.org/10.1038/35044070
Travisano, M., and R. E. Lenski. 1996. "Long-Term Experimental Evolution in Escherichia Coli.
IV. Targets of Selection and the Specificity of Adaptation." *Genetics* 143(1):15–26. doi:
10.1007/s000360050052.
822
823   van Tatenhove-Pel, R. J., Zwering, E., Boreel, D. F., Falk, M., van Heerden, J. H., Kes, M. B. M.
824   J., … Bachmann, H. (2021). Serial propagation in water-in-oil emulsions selects for
825   Saccharomyces cerevisiae strains with a reduced cell size or an increased biomass yield on
826   glucose. *Metabolic Engineering*, *64*(November 2020), 1–14.
827   https://doi.org/10.1016/j.ymben.2020.12.005
828

829 Vasi, F., Travisano, M., & Lenski, R. E. (1994). Long-Term Experimental Evolution in
830   Escherichia coli . II . Changes in Life-History Traits During Adaptation to a Seasonal
831   Environment. *The University of Chicago Press 144*(3), 432–456.
832

833 Velicer, G. J. (1999). Pleiotropic effects of adaptation to a single carbon source for growth on
834   alternative substrates. *Applied and Environmental Microbiology*, *65*(1), 264–269.
835   https://doi.org/10.1128/aem.65.1.264-269.1999
836

837 Wiser, M. J., Ribeck, N., and Lenski, R. E. 2013. "Long-Term Dynamics of Adaptation in
838   Asexual Populations." *Science (New York, N.Y.)* 342(6164):1364–67. doi:
839   10.1126/science.1243357.
840

841 Woods, R. J., Barrick, J. E., Cooper, T. F., Shrestha, U., Kauth, M. R., & Lenski, R. E. (2011).
842   *Second-Order Selection for Evolvability in a Large Escherichia coli Population*. (March),
843   1433–1437.
844

845 Woods, R., D. Schneider, C. L. Winkworth, M. A. Riley, and Lenski, R. E. (2006). Tests of
846   parallel molecular evolution in a long-term experiment with *Escherichia coli*. *Proceedings*
847   *of the National Academy of Sciences, USA* 103:9107-9112.
848

849 Xia, T., Sriram, N., Lee, S. A., Altman, R., Urbauer, J. L., Altman, E., and Eiteman, M. A. 2017.
850   "Glucose Consumption in Carbohydrate Mixtures by Phosphotransferase-System Mutants
851   of Escherichia Coli." *Microbiology (United Kingdom)* 163(6):866–77. doi:
852   10.1099/mic.0.000480.
853

854 Yulo, P. R. J., & Hendrickson, H. L. (2019). The evolution of spherical cell shape; progress and
855   perspective. *Biochemical Society Transactions*, *47*(6), 1621–1634.
856   https://doi.org/10.1042/BST20180634

875
876
877
878