1   **ERK-mediated Curvature Feedback Regulates Branching Morphogenesis in Lung**
2   **Epithelial Tissue**
3

4   *Tsuyoshi Hirashima[1,2,3] and Michiyuki Matsuda[2,4,5]
5
1. The Hakubi Center, Kyoto University, Kyoto, Japan
2. Laboratory of Bioimaging and Cell Signaling, Graduate School of Biostudies, Kyoto
University, Kyoto, Japan
3. Japan Science and Technology Agency, PRESTO, Kawaguchi, Japan
4. Department of Pathology and Biology of Diseases, Graduate School of Medicine, Kyoto
University, Kyoto, Japan
5. Institute for Integrated Cell-Material Sciences, Kyoto University, Kyoto, Japan
13
14
15   ORCID
16   TH: 0000-0001-7323-9627
17   MM: 0000-0002-5876-9969
18
19
20   *Corresponding author: Tsuyoshi Hirashima, Ph.D
21
22   The Hakubi Center/Graduate School of Biostudies, Kyoto University
23   Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan
24   Phone: +81-75-753-4422
25   hirashima.tsuyoshi2m@kyoto-u.ac.jp
26

Abstract

Intricate branching patterns emerge in internal organs because of the repetitive presence of simple deformations in epithelial tissues. During murine lung development, epithelial cells in distal tips of a single tube require fibroblast growth factor (FGF) signals generated by their surrounding mesenchyme to form repetitive tip bifurcations. However, it remains unknown how the cells employ FGF signaling to convert their behaviors to achieve the recursive branching processes. Here we show a self-sustained epithelial regulatory system during the murine lung branching morphogenesis, mediated by extracellular signal-regulated kinase (ERK), which acts as a downstream driver of FGF signaling. We found that tissue-scale curvature regulates ERK activity in the lung epithelium using two-photon live cell imaging and mechanical perturbations. ERK is activated specifically in epithelial tissues with a positive curvature, regardless of whether the change in curvature was attributable to morphogenesis or artificial perturbations. Moreover, we found that ERK activation accelerates actin polymerization specifically at the apical side of cells, and mechanically contributes to the extension of the apical membrane, leading to a decrease in epithelial tissue curvature. These results indicate the existence of a negative feedback loop between tissue curvature and ERK activity beyond scale. We confirmed that this regulation was sufficient to generate the recursive branching processes by a mathematical model. Taken together, we propose that ERK mediates the curvature feedback loop underlying the process of branching morphogenesis in developing lungs.

Keywords: Branching morphogenesis, ERK, FRET cell imaging, lung development, mathematical model, self-organization

Introduction

Characteristic patterns are generated in developing tubular organs through the repeated occurrence of tissue morphogenetic modes, such as elongation, bending, and bifurcation [1–4]. During murine lung development, the complex branched architecture is stereotypically created from a simple epithelial tube, by assembling morphogenetic modes as building motifs with a spatio-temporal order [2]. Previous genetic studies have revealed the essential molecules and signaling pathways underlying lung epithelial morphogenesis [5–7]. Moreover, previous studies have proposed how cellular behaviors and mechanical forces determine the tube shape [8–10]. Thus, the interplay between chemical signals and cell mechanics in constituent cells determines the organized periodic branched pattern, though it is not completely understood yet.

Several pharmacological and genetic studies have demonstrated that fibroblast growth factor (FGF) signaling is a key regulator of lung branching morphogenesis during murine development [11]. The treatment of an isolated epithelium in which the mesenchyme is removed with different FGF ligands has clarified the distinct roles of FGFs in lung epithelial morphology. For example, FGF1 and FGF10 induce bud formation, while FGF7 gives rise to a balloon-like shape, and any of these ligands could promote epithelial proliferation [12–14]. An investigation of the spatio-temporal expression pattern uncovered the fact that *Fgf10* is expressed locally in the mesenchyme surrounding the distal bud of the lung epithelium [14], and its main receptor, *Fgfr2b*, is homogeneously expressed in the lung epithelium [12]. These results led to hypothesize that the *Fgf10* prepattern initiates branching during lung development [14]; this was partially supported by results of previous studies on mice in which *Fgf10* or its main receptor *Fgfr2b* were knocked out and showed the complete absence of lungs [15–17]. However, it was demonstrated that the ubiquitous overexpression of *Fgf10* in its knocked out background could restore the branching process [18], indicating that the *Fgf10* expression is required but the precise prepattern is dispensable for the murine lung branching morphogenesis. Thus, we explored how FGF signals are converted to cellular behaviors underlying tissue morphogenesis.

A main FGF downstream signaling pathway, the extracellular signal-regulated kinase ERK/MAP kinase pathway, includes a three-tiered kinase cascade, i.e., RAF - MEK – ERK [11]. The genetic inactivation of either *Mek1/2* or *Erk1/2* specifically in the airway epithelium results in lung agenesis [19], implying that ERK and its activation were essential for lung development [20]. Previously, immunofluorescence staining results suggested that the phosphorylation of ERK was most likely to occur in the distal tips of the lung epithelium, and may thus be involved in the budding process during lung development [21–23]. FGFR2-mediated ERK activation reportedly upregulates the expression of *Anxa4*, whose products are able to bind with F-actin

bundles, and promote cell migration within epithelial tissues [22]. These reports revealed the role

of ERK activity in cellular behavior during lung development. However, it is still unclear how

the epithelial cells collectively behave to generate the periodic branching morphogenesis

mediated via ERK signaling.

In this study, we show a physical basis for self-sustained branching morphogenesis by

combining two-photon live cell imaging of a Förster resonance energy transfer (FRET)-based

biosensor, mechanical perturbation, and mathematical modeling. We found that ERK

activation occurs specifically in epithelial tissues with a positive curvature, and promotes cell

extension through actin polymerization at the apical edges, resulting in a decreased tissue

curvature. Thus, our study clarifies that the negative feedback loop between the tissue curvature

and ERK activity underlies the process of repetitive branching morphogenesis in developing

murine lung epithelial tissues.

Results

1. ERK activation occurs in distal tips of lung epithelial tissue

We first examined the impact of ERK activity on lung epithelial morphology during murine

development. We cultured developing murine lungs dissected at embryonic day (E) 12.5 in the

presence or absence of PD0325901, an inhibitor of the ERK activator MEK, under explant

culture conditions. The growth of 1-day cultured lung epithelial tissues was remarkably

impaired by the inhibition of ERK activity (Figure 1A, 1B), indicating the importance of ERK

activation for epithelial morphogenesis during this stage.

Then, we visualized the spatial distribution of ERK activity in lung lobes dissected from

murine embryos that ubiquitously expressed the FRET biosensor for the ERK activity [24–26] via

whole tissue two-photon live imaging (Figure 1C, Movie 1). FRET images showed that the

ERK activity in the epithelium was highly dependent on tissue curvature; the ERK activity was

higher in the curved sheet protruding the mesenchyme (convex), while it was lower in the sheet

bent towards the lumen (concave) (Figure 1C). Although endothelial cells exhibited relatively

high levels of ERK activity, they were less involved in epithelial branching morphogenesis [27];

thus, we focused on ERK activity in epithelial tissues. The basal surface of epithelial tissue

was extracted for quantification, and positions at nodes of generated meshes were embedded

on the surface, enabling us to map chemical and geometrical quantities, such as ERK activity,

epithelial tissue curvature, and tissue thickness (Figure 1D, S1A). As expected, there was a

positive correlation between the ERK activity and tissue curvature ($\rho=0.51$, where ρ is the

Spearman's rank correlation coefficient; Figure 1E, 1G, S1B), which indicates that ERK

activation would occur mainly in convex regions. Moreover, there was a negative correlation

between the ERK activity and tissue thickness ($\rho=-0.48$; Figure 1F, 1G). As observed in distal

buds, the epithelial tissue was thinner in the tip region, where ERK was activated, than in the

stalk region, where ERK was inactivated (Figure 1H). These results suggest the existence of

interplay between the ERK activity and cell morphology in developing murine lungs.

Figure 1 Relationship between ERK activity and tissue geometry in the lung epithelium
(A) Immunofluorescence images of anti-E-cadherin staining in an ex vivo culture of lungs for
1 day from E12.5, after treatment with dimethyl sulfoxide (DMSO) (left) and 1 µM PD0325901
(right). Scale bar, 200 µm. (B) Area and perimeter of epithelial tissues from the projected
images. Data represent mean and standard deviation (SD) values. N=6, two-sample t-test,
p<0.001.
(C) Representative ERK activity map in a developing lung at E12.5. Scale bar, 100 µm.
(D) Gaussian curvature map on the basal side of lung epithelial cells.
(E, F) Scatter plot of ERK activity versus Gaussian curvature (E) and that of ERK activity
versus sheet thickness (F). Color indicates frequency. n=427, N=3.
(G) Correlation coefficient of (E) and (F). N=3, one-sample t-test, p=0.001 (E), and p=0.004
(F).
(H) ERK activity map in the distal region of a lung dissected at E12.5. Arrows facing each
other indicate the sheet thickness at a tip (magenta) and distal stalk (blue). Scale bar, 50 µm.

2. ERK activity follows the morphological change of isolated epithelial tissues

To analyze how ERK activity is regulated in the lung epithelium, we isolated epithelial tissues from mesenchymal tissues, and embedded them in Matrigel with culture media containing FGFs for ex vivo culture (Figure 2A). This simplified experimental system allows us to explore responses to external chemical and mechanical stimuli in epithelial cells under regulated conditions. As reported previously [12,13], treatment with 500 ng mL^{-1} FGF1 for 1 day induces branched epithelial buds, while that with 100 ng mL^{-1} FGF7 gives rise to a cyst-like structure with an enlarged lumen (Figure 2B). PD0325901 treatment with FGF1 results in the collapse of the lumen and absence of large deformations; this was also observed with the absence of growth factors with 1-day-old culture (Figure 2B). We then examined the spatial distribution of ERK activity in epithelial tissues, 1 hour after chemical treatment. FGF1 activates ERK exclusively at the distal buds, in a manner similar to that observed with the mesenchyme (Figure 1C), through ligand endocytosis from the basal side of cells (Figure S2, Movie 2), despite the homogeneous expression of four FGF receptors, i.e., FGFR1, FGFR2, FGFR3, and FGFR4 in the distal epithelial tissues [12,18,28,29]. In contrast, FGF7 activates the ERK almost homogeneously in the epithelial tissues (Figure 2C). As expected, ERK activity was homogeneously low in epithelial cells treated with PD0325901 or cultured without any growth factors (Figure 2C). These results suggest that localized ERK activation at the distal tips is key to the process of branching morphogenesis.

To examine the spatio-temporal dynamics of ERK activity during the branching of isolated epithelial tissues, we performed two-photon live imaging under FGF1 supplemented conditions. We found that the ERK activity increased during the protrusion of epithelial tips, and decreased when the tip apex became flat during the tip bifurcation process (Figure 2D, Movie 3). Notably, no mesenchymal cells were observed around the distal tip of the epithelium (Figure 2D), suggesting that tip bifurcation could occur in a self-sustaining manner without mesenchymal cells. The quantification of the ERK activity and tissue curvature at the tip apex shows that the decrease in ERK activity follows that in curvature, with an average delay of 97 min (Figure 2E, 2F). This observation suggests that the ERK activity is controlled by the curvature of the epithelial cells.

To clarify the morphological differences in cells between the convex and concave regions of epithelial tissues, we measured the basal and apical areas of epithelial cells through whole-tissue staining using E-cadherin. The basal cell area of isolated epithelial tissues in the convex region is significantly larger than that in the concave region (Figure 2G, left), while the apical cell area in the convex region is almost equivalent to that in the concave region (Figure 2G,

176 right). The epithelial basal edges act as an interface for cells to receive growth factor signals

177 from external environments that activate ERK (Figure S2). Therefore, we speculate that the

178 extension of basal membranes would trigger ERK activation in the convex regions of epithelial

179 tissues.

Figure 2 ERK activity and tissue curvature in isolated epithelial cells

(A) Schematics of the culture system for isolated lung epithelial tissues.

(B) Bright field images of isolated epithelial tissues subjected chemical treatments for 1 day

from E12.5. Concentrations: 500 ng mL^{-1} FGF1, 100 ng mL^{-1} FGF7, 1 μM PD0325901 (MEKi).

Scale bar, 100 μm.

(C) ERK activity of isolated epithelial tissues subjected to chemical treatments shown in (B)

for 1 hour from E12.5. Scale bar, 50 μm.

(D) Time-lapse images of ERK activity (upper) and bright field (lower) during terminal

bifurcation of lung epithelial tissues. Scale bar, 30 μm.

(E) Tissue curvature (blue) and ERK activity (red) at the tip apex over time (E). Vertical dotted

lines indicate the peak time for the tissue curvature and ERK activity, respectively.

(F) Lag time from the peak for tissue curvature to the peak for ERK activity. N=3.

(G) Basal area (left) and apical area (right) of individual cells in the convex and concave

regions of isolated epithelial tissues. Bold and error bars represent the mean and SD values,

respectively. n=60, N=3. Welch's t test, p<0.001 (left) and p=0.18 (right).

3. The shape of epithelial tissues controls the ERK activity

To test whether the shape of epithelial tissues controls the ERK activity, we performed a physical perturbation assay using isolated epithelial tissues. We embedded isolated epithelial tissues within the FGF1-containing Matrigel filled inside a polydimethylsiloxane (PDMS) chamber, and uniaxially compressed it by 33% either parallelly or vertically to the distal-proximal axis of the lung epithelium (Figure 3A). Time-lapse images obtained during this process allow us to examine the ERK activity response to perturbations in different axes (Figure 3B, 3C). For parallel compression, the curved epithelial tissues at the tip apex became flattened (Figure 3D) and ERK activity decreased exclusively in the flat region (Figure 3E, Movie 4). However, for vertical compression, the tip apex of the epithelium became sharp (Figure 3D) and ERK activation followed the change in tissue curvature (Figure 3E, Movie 5).

We further examined the ERK activity response to cell deformation. The manipulation of cell shape in curved epithelial tissues remains unfeasible under 3D culture conditions; thus, we performed a strain perturbation assay under planar culture conditions using dissociated lung epithelial cells (Figure 3F). Under planar culture conditions, the stretching of cells led to ERK activation, while their compression led to ERK inactivation (Figure 3G, 3H), indicating that the strain on the cell area is key to curvature-dependent ERK activity regulation. Notably, the surface area of the basal side of a cell is significantly larger in the convex region with high ERK activity than in the concave region with lower ERK activity (Figure 2G). Together, epithelial tissue curvature controls the ERK activity in cells, most likely through the strained state of their basal membranes.

Figure 3 Response of ERK activity to mechanical perturbation

(A) Schematics of uniaxial compression assay performed on an isolated epithelium under 3D culture conditions. The *x* and *y* axes indicate directions parallel and vertical to the distal and proximal axes of the epithelium, respectively.

(B, C) Time-lapse images of the ERK activity for parallel compression (B) and vertical compression (C). Time origin indicates the timing of mechanical compression. Scale bars, 50 μm.

(D, E) Tissue curvature at the tip apex (D) and normalized ERK activity (E) over time. Colors represent the axes of compression, as shown in (A). Data represent mean and SD values. N=3.

(F) Schematics of uniaxial mechanical perturbations under 2D culture conditions.

(G) Time-lapse images of the ERK activity in the isolated lung epithelium for the extension (upper) and compression (lower) processes. Scale bar, 50 μm.

(H) Normalized ERK activity for the extension (red) and the compression (blue) processes over time. Time origin indicates the timing of mechanical perturbations. Data represent mean and SD values. N=3.

4. ERK activation modulates cell shape via actin polymerization at the apical side

Next, we investigate how the ERK activity contributes to epithelial morphology and mechanics. First, we observed the change in the cell shape of isolated epithelial tissues treated with PD0325901. Drug treatment led to an inhibition in the ERK activity within 30 min after administration (Figure S3A), and the apico-basal length of epithelial cells became longer than that observed in the control within 2 hours (Figure 4A, Movie 6). Hereafter, we refer to the apico-basal length of cells as the cell height, and the cell length orthogonal to the apico-basal axis as the cell width (Figure 4B). ERK inactivation in cells treated with PD0325901 lengthens the cell height to 124% and shortens the cell width to 92% on an average, despite no significant changes in the cell size, compared to mock-treated cells (Figure 4C). This trend is consistent with that observed in the dissected murine lungs (Figure 1E–G). These results indicate that the ERK activity regulates the cell height, while maintaining the cell size.

Next, to explore what causes the cell shape to change mechanically, we performed inhibitor assays on cytoskeletons, such as actin, microtubule, and non-muscle myosin, which allowed us to determine the mechanical contributions of targeted cytoskeletons to the cell morphology. Given that the cell size is maintained even if the cell shape varies (Figure 4C), two representative geometrical parameters, cell height and cell width, are in an inverse relationship (see Materials and Methods IIa). Thus, we focused only on the change in the cell height caused by cytoskeletal perturbations. The cell height was increased markedly upon treatment with Latrunculin A or Cytochalasin D, each an inhibitor of actin polymerization, as observed in inhibiting ERK activity with PD0325901; conversely, the cell height was decreased by treatment with Jasplakinolide, which promotes actin polymerization (Figure 4D, 4E). Because F-actin remarkably accumulates at the apical side of cells (Figure S3B), we speculated that actin polymerization at the apical edge would contribute to cell width extension, leading to the shortening of the cell height. The cell height was also shortened by interfering with microtubule polymerization with Nocodazole; combined treatment with Nocodazole and Latrunculin A resulted in significant cell height extension via the rupture of lateral cell edges (Figure S3D). Microtubules within cells are distributed along the apico-basal axis (Figure S3C), and would thus stabilize cytoskeletal architectures sustaining cell height with F-actin. The inhibition of the activity of non-muscle myosin II with Blebbistatin resulted in no significant difference in cell height, compared to that observed with DMSO control (Figure 4E). This was consistent with the fact that the activity of myosin is insignificant in the tips of lung epithelial tissue (Figure S3E-E"). To further analyze the mode of actin polymerization, we used CK-666, an Arp2/3 complex inhibitor CK-666, and SMIFH2, an inhibitor of the formin homology 2

domain. We found that the cell height was lengthened for CK-666 treatment, but slightly
shortened for SMIFH2 treatment (Figure 4E), suggesting that actin polymerization would
depend on the Arp2/3-mediated nucleation of branched filaments, rather than formin-mediated
actin nucleation.

Because the inhibition of ERK activity and actin polymerization resulted in similar effects
on the cell height, we speculated that ERK activation promoted F-actin accumulation within
cells. Simultaneous imaging for the two factors revealed that F-actin accumulation at the apical
side of cells was observed exclusively in the tips where ERK activation occurs, under FGF1-
treated conditions (Figure 4F). Mapping between the ERK activity and F-actin accumulation
level via 3D whole-tissue imaging clearly showed a positive correlation with FGF1 (ρ=0.51),
although the inhibition of the ERK activity led to a significant decrease in F-actin accumulation
at the apical side of cells throughout the tissue, and relatively uniform distribution (Figure 4G).
In addition, treatment with FGF7, which activates ERK globally in the isolated epithelial
tissues, regardless of the tissue curvature (Figure 2C), led to an increase in the apical F-actin
accumulation overall throughout the epithelial tissues (Figure 4G). Thus, we found that ERK
activation promotes actin polymerization at the apical edges of cells, and ultimately controls
the cellular shape in the epithelial monolayer.

Figure 4 Response of cell morphology to chemical perturbations
(A) Change in the cell shape of an isolated epithelium upon treatment with either 500 ng mL^{-1}
FGF1 (left) or 500 ng mL^{-1} FGF1 and 1 μM PD0325901. Cell membranes were visualized

using the whole-mount immunofluorescence of anti-E-cadherin. The inset windows represent
a magnified 3D rendering focusing on a representative single cell marked in yellow. Scale bar,
30 μm.
(B) Schematics for cellular morphological measurement.
(C) Quantification of cell morphological parameters, including cell height, width, and size.
n=60, N=3. Welch's t test, $p<0.001$ (left), $p=0.013$ (middle), and $p=0.4$ (right).
(D) Change in the cell morphology of an isolated epithelium upon treatment with either DMSO
as a control, 1 μM Latrunculin A, or 1 μM Jasplakinolide. The cell shape was visualized using
SiR-Actin, a F-actin labeling probe. Scale bar, 20 μm.
(E) Relative change in cell height upon inhibitor treatment for cytoskeletons. Concentrations
of inhibitors: 1 μM Latrunculin A, 1 μM Cytochalasin D, 1 μM Nocodazole, 30 μM
Blebbistatin, 1 μM Jasplakinolide, 100 μM CK-666, and 10 μM SMIFH2. The bar graph and
error bars represent mean and SD values. N≥3 for each treatment.
(F) Spatial map of ERK activity and F-actin accumulation level visualized by SiR-Actin
labeling upon treatment with 500 ng mL^{-1} FGF1. Inset windows show magnified views that
focus on their subcellular localization. Scale bar, 50 μm.
(G) Scatter plot and contour map of ERK activity in the cytosol and F-actin intensity at the
apical edge. Colors indicate the administration of different chemicals; red: 500 ng mL^{-1} FGF1,
n=1204, N=4, blue: 500 ng mL^{-1} FGF1 and 1 μM PD0325901 (MEKi), n=687, N=3, green:
100 ng mL^{-1} FGF7, n=787, N=3.
316

5. ERK activation flattens epithelial tissues via apical actin polymerization

How does the ERK-mediated actin polymerization at the apical side of cells result in morphological changes in monolayer tissues? To answer this, we used a simple geometrical model of a two-dimensional monolayer tissue, composed of cells with identical morphologies. In this model, individual cells are depicted by tetragons, the edges of which each correspond to the apical, basal, and lateral sides of cells. Then, the sheet curvature κ can be represented as a function of the length of these edges; especially, a change in the curvature with respect to the apical edge length a can be derived as $\partial\kappa/\partial a < 0$. Thus, the sheet curvature decreases with an expansion in the apical edge (see Materials and Methods IIb). This theoretical analysis predicts that ERK activation would decrease the tissue curvature via actin polymerization at the apical edges.

To confirm this prediction, we examined the change in epithelial tissue curvature in response to acute ERK activation. For this purpose, an isolated lung epithelium was incubated in the culture medium without any growth factors. Then, FGF1 was added into the medium to simulate ERK. Acute ERK activation exclusively promoted F-actin accumulation at the apical side of cells in epithelial distal tips; this was followed by the flattening of the epithelial sheet (Figure 5A–D, Movie 7). Moreover, the simultaneous administration of FGF1 and PD0325901 resulted in neither F-actin accumulation nor tip flattening; this was also observed for PBS control treatment (Figure 5B–D). These results indicate that ERK activation flattens the epithelial tissues in the tips via actin polymerization at the apical sides of cells.

Figure 5 Simultaneous imaging of FGF-triggered ERK activation, actin polymerization, and tissue flattening

(A) Time-lapse images of ERK activity (upper) and F-actin intensity (lower). SiR-Actin was pretreated for F-actin visualization, 1 hour before the addition of FGF1. The time origin indicates the timing of FGF1 administration. Scale bar, 30 μm.

(B–D) Normalized ERK activity in the tips (B), normalized F-actin intensity in the tips (C), and tissue curvature at the tip apex (D) over time. Colors indicate different chemical treatments, including with 500 ng mL^{-1} FGF1 and 1 μM PD0325901 (MEKi). The time origin indicates the timing of chemical administration. Data represent mean and SD values. N=4.

6. Feedback between ERK activity and tissue curvature explains repetitive branching

To examine whether the curvature-dependent apical extension would be able to produce the periodic branching pattern, we performed numerical analysis using vertex dynamics model, a cell-based mechanical model [30–32]. The model represents epithelial cells as polygons with vertices and edges, and the edges on the apical side actively deform via F-actin dynamics that depend on the tissue-scale curvature and its decay (as described in Materials and Methods IIc). We conducted numerical simulations with an initial shape and bud diameters obtained from the experiment (Figure 6A), and examined the impact of curvature on the change in apical edge length through a model parameter α, using the standardized parameter set (Figure S4A). The apical edge extends/shrinks in accordance with the positive/negative values of α in a curvature-dependent manner (Figure 6B). Clearly, the curvature and cell height at the tip apex decreased with an increase in α at equilibrium (Figure 2B, S5B). Moreover, during curvature-dependent apical extension ($\alpha = 1$), the curvature decreases at the tip apex (arrowhead, Figure 2C), but increases at both sides over time (arrows, Figure 2C), indicating that the curvature-dependent apical extension can create new buds for terminal bifurcation. This does not happen in the opposite case ($\alpha = -1$), suggesting that curvature-dependent apical constriction is not likely to drive repetitive branching, contradictory to a previous report [33].

We then enabled cell proliferation to occur in the model system, in order to perform in silico experiments in virtual growing monolayer tissues. The simulation of the regime of curvature-dependent apical extension and cell proliferation clearly demonstrates the formation of repetitive branching patterns (Figure 6D, Movie 8). In contrast, the repetitive branching pattern is not generated in counterpart regimes, such as no regulation on the apical edge, and global apical extension as observed for the FGF7 treatment – apical extension happens in all cells, irrespective of curvature (Figure 6E). In these counterpart regimes, there are significant differences in both the averaged curvature, a measure of curvature averaged over the entire tissue, and the correlation coefficient between the curvature and monolayer thickness (Figure 6F, 6G). Moreover, the virtual experiments show that luminal pressure, which has been proposed to be a physical factor for lung morphogenesis [34,35], is not essential for branching pattern formation; rather, it suppresses its constant large magnitude value (Figure S4C). These results support the fact that that the exclusive occurrence of apical edge extension in cells with high curvatures drives repetitive terminal bifurcation in the growing epithelial monolayer in a self-sustaining manner.

Furthermore, numerical investigations have clarified that the retention time of curvature memory in F-actin, i.e., the inverse of the F-actin decay rate γ, is crucial for generating the

branching pattern (Figure 6H, Movie 9–11). The averaged curvature becomes larger for a longer curvature memory, and the correlation between the curvature and F-actin becomes higher for a shorter curvature memory. Thus, there is an optimal range for the retention time of the curvature memory in F-actin (Figure 6I). This result suggests the importance of the tissue curvature hysteresis of cellular force generation via F-actin, for the emergence of repetitive branch patterns.

Figure 6 Simulation and analysis of the mathematical model
(A) Distal tip of a virtual epithelial monolayer tissue during the initial configuration of the simulation. Colors indicate the tissue curvature. D represents the diameter of the tip. Scale bar, 30 µm.
(B) Tissue curvature at the tip apex over the parameter α. Colors indicate the tip diameter.
(C) Time evolution of the basal sides of cells in different α. Colors indicate the time periods in simulations. The arrowhead indicates the tip apex and arrows indicate the regions with a growing curvature. Scale bars, 50 µm.
(D) Time evolution of the basal sides of cells in the simulation with cell proliferation at $\alpha=1$.
(E) Generated morphology of epithelial monolayer tissues in different regimes. Curvature-dependent apical extension: $\alpha=1$, $\beta=0$; no regulation: $\alpha=0$, $\beta=0$; global apical extension: $\alpha=0$, $\beta=0.2$.

(F, G) Averaged curvature (F) and correlation coefficient between the tissue curvature and
tissue thickness (G) for the 3 regulatory regimes. Bars represent mean values. N=5. Welch's t-
test, p<0.001.
(H) Spatial maps of the curvature (upper) and F-actin level (lower) for different curvature
memory time periods γ^{-1}. left: $\gamma=1$, middle: $\gamma= 10^{-3}$, right: $\gamma= 10^{-6}$.
(I) Tissue curvature (green) and correlation coefficient between the curvature and the F-actin
level. Data represent mean and SD values. N=5. The gray shaded region indicates the region
in which both the values are high.
413

Discussion

Here, we have demonstrated a regulatory mechanism by which epithelial monolayer cells undergo repetitive branching morphogenesis through curvature sensing and ERK-mediated mechanical force generation in a self-sustained manner. Our two-photon live imaging analysis has shown that the ERK activity changes following a dynamic change in the tissue curvature, and ERK activation occurs preferentially in curved epithelial tissues. Moreover, we found that ERK-mediated actin polymerization contributes to the extension of the apical side of cells, and demonstrated its importance for achieving a decrease in tissue curvature during terminal bifurcation via mathematical model analyses. Thus, there is a closed negative feedback loop between the tissue curvature and ERK activity across multiple scales in developing lung epithelial tissues (Figure 7).

The results presented in this study provide a better understanding of lung epithelial branching, in conjunction with previously determined theory-based mechanisms, based on mechanical and chemical aspects. From a mechanical viewpoint, it has been proposed that the spatial patterning of the isolated lung epithelium arises from the proliferation-induced physical instability of the epithelial layer, known as mechanical buckling [36], in various developing epithelial tissues [1,37,38]. This purely mechanical model may explain the formation of an initial branching pattern, under the assumption that the epithelial tissue acts as an elastic layer, but may require the physical conditions under which mechanical buckling occurs. However, our results have demonstrated that repetitive branching can be generated even at a range where proliferation-induced buckling did not occur (Figure 6E–G). We emphasize that the regime presented here is not mutually exclusive to buckling-induced patterning. Thus, ERK-mediated tissue curvature feedback results in a robust regulatory system underlying branching morphogenesis, together with the buckling regime. In addition, a simple reaction–diffusion model of substrate–depletion has been proposed for integrating chemical signal and tissue growth [39,40]; the model assumes that lung epithelial cells consume FGF ligands that have slow diffusion rates, and thereby predicts that a concentration gradient of FGF is generated in the extracellular matrix. However, live imaging analysis results have shown that no FGF gradient was formed in the Matrigel, while FGF was internalized in the epithelial cells through endocytosis (Figure S2). Other reaction-diffusion systems have focused on genetic regulatory networks that can achieve the localization of *Fgf10* expression observed in vivo [41–43]. Although the spatial distribution of the chemoattractant is dispensable for epithelial branching [18] (Figure 2D), the incorporation of such fundamental biochemical signals into our model would be useful for studying detailed systems for lung branching and patterning in vivo.

Our findings highlight the curvature-sensitive ERK activation of lung epithelial cells and
raise an important question about how cells sense the tissue curvature and transmit FGF–ERK
signals. Because the ERK activity response occurs within 10 min after the perturbation in the
tissue curvature (Figure 3B–E), the chemical feedbacks via gene expression, e.g., the interplay
between the FGF and Sonic hedgehog expressed in the epithelium, are unlikely but rather via
unknown protein–protein interactions should be involved in this process. We determined from
the results (Figure 2G, 3G–H) that the phenomenon of curvature-sensitive ERK activation in
lung epithelial tissues was attributable to a strain exhibiting a certain cell membrane or level of
membrane tension, as shown previously for other tissues [44–46]. Moreover, the state of the
membrane would trigger receptor-mediated endocytosis, as FGF internalization occurs
exclusively in epithelial tissues with a positive curvature (Figure S2). Multiple intracellular
events, including the endocytic trafficking of FGFR and molecular sensing of plasma
membrane curvature, would be involved in this process [47–49].

This study has focused on the core mechanism to minimally generate the self-sustained
repetitive branching patterns in epithelium; however, there are other additional contributions
that could achieve the robust tissue morphogenesis in vivo. For example, smooth muscle cells
are not necessary for the initiation of cleft formation (Figure 2D)[50], but should mechanically
assist the cleft formation from the outside of an epithelium, realizing the stable terminal
bifurcation [27,50]. Luminal fluid is another potentially important factor that can mechanically
affect epithelial tissue deformation through shear stress and hydrostatic pressure [51]. It has been
previously reported that the manipulation of luminal pressure in murine developing lungs under
ex vivo culture conditions altered branch formation [34]. As oscillatory contractions of airway
smooth muscles are likely to produce dynamic changes in the luminal pressure, it has been
suggested that rhythmic luminal pressure would control the global synchronization of
branching events in the murine lung [35], although it does not contribute to the tissue flattening
during the terminal bifurcation.

The ERK-mediated curvature feedback mechanism proposed here advances the concept
of mechanochemical regulation of the epithelial monolayer [32,44,52,53], in terms of taking the
impact of tissue geometry into account. In various tissues, the epithelial cells collectively form
their shape in the tissue scale and the constituent cells respond to tissue curvature at different
levels, including the transcription and cell activity levels [54–57]. We believe that the proposed
system works as a regulatory principle underlying the mechanisms of self-organized tissue
morphogenesis.

Figure 7 ERK-mediated curvature feedback regulates branching morphogenesis in murine lung epithelial tissue

The epithelial cells activate ERK specifically in tissues with a positive curvature via reception of extracellular FGF ligands (red arrow). ERK-mediated actin polymerization contributes to the extension of the apical side of cells, which decreases the tissue curvature (blue arrow). Thus, there is a closed negative feedback loop between tissue curvature and ERK activity across multiple scales in developing epithelial lung tissues.

Acknowledgements

This work was supported by the JST PRESTO grant JPMJPR1949, by the JSPS KAKENHI 19H00993, by the SPIRITS 2018 of Kyoto University, and by the Medical Research Support Center of Kyoto University. We would like to thank Akane Kusumi for technical assistance, Edouard Hannezo, Naoya Hino, Mamoru Ishii, and Takuya Yoshida for fruitful discussion.

Author Contributions

Conceptualization, T.H.; Methodology, T.H.; Software, T.H.; Validation, T.H.; Formal analysis, T.H.; Investigation, T.H.; Resources, T.H., M.M.; Data curation, T.H.; Writing - original draft, T.H.; Writing - review & editing, T.H., M.M.; Visualization, T.H.; Supervision, T.H.; Project administration, T.H., M.M.; Funding acquisition, T.H., M.M.

Competing interests

The authors declare no competing interests.

References

1.   Hirashima, T. Pattern Formation of an Epithelial Tubule by Mechanical Instability
during Epididymal Development. *Cell Rep.* **9**, 866–73 (2014).

2.   Metzger, R. J., Klein, O. D., Martin, G. R. & Krasnow, M. A. The branching
programme of mouse lung development. *Nature* **453**, 745–750 (2008).

3.   Hannezo, E. *et al.* A Unifying Theory of Branching Morphogenesis. *Cell* (2017).
doi:10.1016/j.cell.2017.08.026

4.   Lu, P. & Werb, Z. Patterning mechanisms of branched organs. *Science (80-.).* **322**,
1506–1509 (2008).

5.   Cardoso, W. V. & Lü, J. Regulation of early lung morphogenesis: questions, facts and
controversies. *Development* **133**, 1611–1624 (2006).

6.   Warburton, D. *et al.* Lung organogenesis. in *Current Topics in Developmental Biology*
(2010). doi:10.1016/S0070-2153(10)90003-3

7.   Herriges, M. & Morrisey, E. E. Lung development: Orchestrating the generation and
regeneration of a complex organ. *Development (Cambridge)* (2014).
doi:10.1242/dev.098186

8.   Varner, V. D. & Nelson, C. M. Cellular and physical mechanisms of branching
morphogenesis. *Development* **141**, 2750–9 (2014).

9.   Ingber, D. E. Mechanical control of tissue morphogenesis during embryological
development. *Int. J. Dev. Biol.* (2006). doi:10.1387/ijdb.052044di

10.   Anlaş, A. A. & Nelson, C. M. Tissue mechanics regulates form, function, and
dysfunction. *Current Opinion in Cell Biology* (2018). doi:10.1016/j.ceb.2018.05.012

11.   Ornitz, D. M. & Itoh, N. The Fibroblast Growth Factor signaling pathway. *Wiley*
Interdiscip. Rev. Dev. Biol. **4**, 215–266 (2015).

12.   Cardoso, W. V., Itoh, A., Nogawa, H., Mason, I. & Brody, J. S. FGF-1 and FGF-7
induce distinct patterns of growth and differentiation in embryonic lung epithelium.
Dev. Dyn. (1997). doi:10.1002/(SICI)1097-0177(199703)208:3<398::AID-
AJA10>3.0.CO;2-X

13.   Nogawa, H. & Ito, T. Branching morphogenesis of embryonic mouse lung epithelium
in mesenchyme-free culture. *Development* (1995).

14.   Bellusci, S., Grindley, J., Emoto, H., Itoh, N. & Hogan, B. L. M. Fibroblast Growth
Factor 10 (FGF10) and branching morphogenesis in the embryonic mouse lung.
Development (1997).

15.   Min, H. *et al.* Fgf-10 is required for both limb and lung development and exhibits

striking functional similarity to Drosophila branchless. *Genes Dev.* (1998).

doi:10.1101/gad.12.20.3156

16. Sekine, K. *et al.* Fgf10 is essential for limb and lung formation. *Nat. Genet.* (1999).

doi:10.1038/5096

17. Arman, E., Haffner-Krausz, R., Gorivodsky, M. & Lonai, P. Fgfr2 is required for limb

outgrowth and lung-branching morphogenesis. *Proc. Natl. Acad. Sci. U. S. A.* (1999).

doi:10.1073/pnas.96.21.11895

18. Volckaert, T. *et al.* Localized Fgf10 expression is not required for lung branching

morphogenesis but prevents differentiation of epithelial progenitors. *Dev.* (2013).

doi:10.1242/dev.096560

19. Boucherat, O., Nadeau, V., Berube-Simard, F.-A., Charron, J. & Jeannotte, L. Crucial

requirement of ERK/MAPK signaling in respiratory tract development. *Development*

142, 3801–3801 (2015).

20. Yoshida, T., Matsuda, M. & Hirashima, T. Incoherent Feedforward Regulation via

Sox9 and ERK Underpins Mouse Tracheal Cartilage Development. *Front. Cell Dev.*

Biol. **8**, (2020).

21. Liu, Y. *et al.* Novel role for netrins in regulating epithelial behavior during lung

branching morphogenesis. *Curr. Biol.* (2004). doi:10.1016/j.cub.2004.05.020

22. Jiang, K., Tang, Z., Li, J., Wang, F. & Tang, N. Anxa4 mediated airway progenitor

cell migration promotes distal epithelial cell fate specification. *Sci. Rep.* (2018).

doi:10.1038/s41598-018-32494-z

23. Tang, N., Marshall, W. F., McMahon, M., Metzger, R. J. & Martin, G. R. Control of

Mitotic Spindle Angle by the RAS-Regulated ERK1/2 Pathway Determines Lung

Tube Shape. *Science (80-.).* **333**, 342–345 (2011).

24. Komatsu, N. *et al.* A platform of BRET-FRET hybrid biosensors for optogenetics,

chemical screening, and in vivo imaging. *Sci. Rep.* **8**, 8984 (2018).

25. Komatsu, N. *et al.* Development of an optimized backbone of FRET biosensors for

kinases and GTPases. *Mol. Biol. Cell* **22**, 4647–56 (2011).

26. Harvey, C. D. *et al.* A genetically encoded fluorescent sensor of ERK activity. *Proc.*

Natl. Acad. Sci. U. S. A. **105**, 19264–9 (2008).

27. Kim, H. Y. *et al.* Localized Smooth Muscle Differentiation Is Essential for Epithelial

Bifurcation during Branching Morphogenesis of the Mammalian Lung. *Dev. Cell*

(2015). doi:10.1016/j.devcel.2015.08.012

28. Yuan, T. *et al.* Temporospatial Expression of Fgfr1 and 2 During Lung Development,

Homeostasis, and Regeneration. *Front. Pharmacol.* **11**, (2020).

29.   De Moerlooze, L. *et al.* An important role for the IIIb isoform of fibroblast growth

factor receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse

organogenesis. *Development* **127**, 483–492 (2000).

30.   Fletcher, A. G., Osterfield, M., Baker, R. E. & Shvartsman, S. Y. Vertex models of

epithelial morphogenesis. *Biophys. J.* **106**, 2291–2304 (2014).

31.   Nagai, T. & Honda, H. A dynamic cell model for the formation of epithelial tissues.

Philos. Mag. B **81**, 699–719 (2001).

32.   Hirashima, T. & Adachi, T. Polarized cellular mechano-response system for

maintaining radial size in developing epithelial tubes. *Development* **146**, dev181206

(2019).

33.   Fumoto, K., Takigawa-Imamura, H., Sumiyama, K., Kaneiwa, T. & Kikuchi, A.

Modulation of apical constriction by Wnt signaling is required for lung epithelial

shape transition. *Development* **144**, 151–162 (2017).

34.   Unbekandt, M. *et al.* Tracheal occlusion increases the rate of epithelial branching of

embryonic mouse lung via the FGF10-FGFR2b-Sprouty2 pathway. *Mech. Dev.* **125**,

314–324 (2008).

35.   Nelson, C. M. *et al.* Microfluidic chest cavities reveal that transmural pressure controls

the rate of lung development. *Development* (2017). doi:10.1242/dev.154823

36.   Varner, V. D., Gleghorn, J. P., Miller, E., Radisky, D. C. & Nelson, C. M.

Mechanically patterning the embryonic airway epithelium. *Proc. Natl. Acad. Sci.* **112**,

9230–9235 (2015).

37.   Kourouklis, A. P. & Nelson, C. M. Modeling branching morphogenesis using

materials with programmable mechanical instabilities. *Curr. Opin. Biomed. Eng.* **6**,

66–73 (2018).

38.   Savin, T. *et al.* On the growth and form of the gut. *Nature* **476**, 57–62 (2011).

39.   Miura, T. & Shiota, K. Depletion of FGF acts as a lateral inhibitory factor in lung

branching morphogenesis in vitro. *Mech. Dev.* **116**, 29–38 (2002).

40.   Hartmann, D. & Miura, T. Modelling in vitro lung branching morphogenesis during

development. *J. Theor. Biol.* **242**, 862–872 (2006).

41.   Hirashima, T., Iwasa, Y. & Morishita, Y. Mechanisms for split localization of Fgf10

expression in early lung development. *Dev. Dyn.* **238**, 2813–2822 (2009).

42.   Menshykau, D., Kraemer, C. & Iber, D. Branch Mode Selection during Early Lung

Development. *PLoS Comput. Biol.* **8**, e1002377 (2012).

43. Menshykau, D., Blanc, P., Unal, E., Sapin, V. & Iber, D. An interplay of geometry and
signaling enables robust lung branching morphogenesis. *Development* **141**, 4526–4536
(2014).

44. Hino, N. *et al.* ERK-Mediated Mechanochemical Waves Direct Collective Cell
Polarization. *Dev. Cell* **53**, 646-660.e8 (2020).

45. Kinoshita, N. *et al.* Mechanical Stress Regulates Epithelial Tissue Integrity and
Stiffness through the FGFR/Erk2 Signaling Pathway during Embryogenesis. *Cell Rep.*
30, 3875-3888.e3 (2020).

46. Sitarska, E. & Diz-Muñoz, A. Pay attention to membrane tension: Mechanobiology of
the cell surface. *Curr. Opin. Cell Biol.* **66**, 11–18 (2020).

47. Francavilla, C. *et al.* Functional Proteomics Defines the Molecular Switch Underlying
FGF Receptor Trafficking and Cellular Outputs. *Mol. Cell* **51**, 707–722 (2013).

48. Belleudi, F. *et al.* Keratinocyte Growth Factor Receptor Ligands Target the Receptor
to Different Intracellular Pathways. *Traffic* **8**, 1854–1872 (2007).

49. Antonny, B. Mechanisms of Membrane Curvature Sensing. *Annu. Rev. Biochem.* **80**,
101–123 (2011).

50. Young, R. E. *et al.* Airway Smooth Muscle Differentiation is Essential for Promoting
Airway Size, Tracheal Cartilage Segmentation, But Dispensable for Epithelial
Branching. *SSRN Electron. J.* (2019). doi:10.2139/ssrn.3454659

51. Chan, C. J. & Hiiragi, T. Integration of luminal pressure and signalling in tissue self-
organization. *Development* **147**, dev181297 (2020).

52. Boocock, D., Hino, N., Ruzickova, N., Hirashima, T. & Hannezo, E. Theory of
mechanochemical patterning and optimal migration in cell monolayers. *Nat. Phys.* **17**,
267–274 (2021).

53. Ishii, M., Tateya, T., Matsuda, M. & Hirashima, T. Retrograde ERK activation waves
drive base-to-apex multicellular flow in murine cochlear duct morphogenesis. *Elife* **10**,
(2021).

54. Nikolaev, M. *et al.* Homeostatic mini-intestines through scaffold-guided organoid
morphogenesis. *Nature* **585**, 574–578 (2020).

55. Nelson, C. M., VanDuijn, M. M., Inman, J. L., Fletcher, D. A. & Bissell, M. J. Tissue
Geometry Determines Sites of Mammary Branching Morphogenesis in Organotypic
Cultures. *Science (80-.).* **314**, 298–300 (2006).

56. Luciano, M. *et al.* Large-scale curvature sensing by epithelial monolayers depends on
active cell mechanics and nuclear mechanoadaptation. *bioRxiv* (2020).

doi:doi.org/10.1101/2020.07.04.187468

57. Hughes, A. J. *et al.* Engineered Tissue Folding by Mechanical Compaction of the
Mesenchyme. *Dev. Cell* **44**, 165-178.e6 (2018).

58. Hirashima, T. & Adachi, T. Procedures for the Quantification of Whole-Tissue
Immunofluorescence Images Obtained at Single-Cell Resolution during Murine
Tubular Organ Development. *PLoS One* **10**, e0135343 (2015).

59. Hirashima, T. Mechanical Tissue Compression and Whole-mount Imaging at Single
Cell Resolution for Developing Murine Epididymal Tubules. *BIO-PROTOCOL* **10**,
(2020).

60. Lukinavičius, G. *et al.* Fluorogenic probes for live-cell imaging of the cytoskeleton.
Nat. Methods (2014). doi:10.1038/nmeth.2972

61. Milroy, L. G. *et al.* Selective chemical imaging of static actin in live cells. *J. Am.*
Chem. Soc. (2012). doi:10.1021/ja211708z

62. Tran, A. P., Yan, S. & Fang, Q. Improving model-based functional near-infrared
spectroscopy analysis using mesh-based anatomical and light-transport models.
Neurophotonics **7**, 1 (2020).

63. Meyer, M., Desbrun, M., Schr, P. & Barr, A. H. Discrete Differential-Geometry
Operators for Triangulated 2-Manifolds. *Vis. Math. III* 113–134 (2003).
doi:10.1016/j.cagd.2007.07.005

64. Hannezo, E., Prost, J. & Joanny, J.-F. Theory of epithelial sheet morphology in three
dimensions. *Proc. Natl. Acad. Sci.* **111**, 27–32 (2014).

65. Wen, F. L., Wang, Y. C. & Shibata, T. Epithelial Folding Driven by Apical or Basal-
Lateral Modulation: Geometric Features, Mechanical Inference, and Boundary Effects.
Biophys. J. (2017). doi:10.1016/j.bpj.2017.05.012

66. Boucherat, O., Nadeau, V., Berube-Simard, F.-A., Charron, J. & Jeannotte, L. Crucial
requirement of ERK/MAPK signaling in respiratory tract development. *Development*
142, 3197–3211 (2014).

Materials and Methods

(I) Experiments

Animals

It has been previously reported that transgenic mice expressed hyBRET-ERK-NES, a FRET

biosensor [24-26]. hyBRET-ERK-NES is comprised of cyan fluorescent protein (CFP) and yellow

fluorescent protein (YFP), which act as the donor and acceptor of FRET, respectively.

Otherwise, we used ICR mice purchased from Japan SLC, Inc. We designated the midnight

preceding the discovery of the vaginal plug as embryonic day 0.0 (E0.0), and all mice were

sacrificed via cervical dislocation to minimize suffering. All animal experiments were

approved by the local ethical committee for animal experimentation (MedKyo 18086, 19090,

and 20081), and were performed in compliance with the guidelines for the care and use of

laboratory animals, at Kyoto University.

Antibodies, small molecules, and recombinant proteins

We used the following primary antibodies at a dilution ratio of 1:200 unless stated otherwise:

rat monoclonal anti-E-cadherin (Cell Signaling Technology, #3195), rabbit polyclonal anti-

phospho-myosin light chain (pMLC) (Abcam, #ab2480), Alexa Fluor 546-conjugated goat

anti-rat IgG (H+L) antibody (Thermo Fisher Scientific, #A11081, 1:1000), Alexa Fluor 647-

conjugated goat anti-rabbit IgG (H+L) antibody (Abcam, #ab150079, or Thermo Fisher

Scientific, #A21247, 1:1000), Blebbistatin (Merck Millipore, #203391), CK-666 (Sigma-

Aldrich, #SML0006), Cytochalasin D (Merck Millipore, #250255), FGF1 (R&D Systems,

#232-FA), FGF7 (R&D Systems, #251-KG), Jasplakinolide (Abcam, #ab141409), Latrunculin

A (Cayman Chemical, #10010630), Nocodazole (Merck Millipore, #487928), PD0325901

(FUJIFILM Wako Pure Chemical Corporation, #162-25291), SMIFH2 (Sigma-Aldrich,

#S4826), Y-27632 (Merck Millipore, #SCM075).

Ex vivo culture

Dissected lung lobes were placed onto a hydrophilic polytetrafluoroethylene organ culture

insert with a pore size of 0.4 μm (Merck Millipore, #PICM01250), which was preset in a 35

mm petri dish filled with 800 μL of culture medium. Samples were cultured in an air-liquid

interface at 37°C under 5% CO_2 conditions. To culture isolated epithelial tissues, dissected

lobes were digested with 10 PU/mL of Dispase I (FUJIFILM Wako Pure Chemical Corporation,

#386-02271) in phosphate buffered saline (PBS) for 15 min on ice, and immersed into fetal

bovine serum (FBS) and kept on ice for 5 min, to stop the protease-mediated reaction. Then,

samples were washed in PBS twice on ice, and the distal tips of epithelial tissues were manually

excised using fine needles. Isolated epithelial tissues were cultured within growth factor-

reduced Matrigel (Corning, #356231) and covered with 500 µL of culture medium added to

each well of glass bottom dishes (Greiner, #627870) at 37°C under 5% CO_2 conditions. We

used FluoroBrite DMEM Media (Thermo Fischer Scientific, #A1896701) containing 1%

GlutaMAX (Thermo Fischer Scientific, #35050061) and 0.5% FBS as the culture medium.

Whole-tissue immune-fluorescence staining

Staining and optical clearing of dissected lung were performed according to previous studies

[58,59]. Briefly, samples were fixed with 4% paraformaldehyde (PFA) in PBS overnight at 4°C.

For pMLC staining, samples were fixed with 2% trichloroacetic acid in PBS containing a 1%

phosphatase inhibitor cocktail (1:100, Nacalai Tesque, #07575-51) for 15 min at 4°C. Then,

samples were blocked by incubation in 10% normal goat serum (Abcam, #ab156046) diluted

in 0.1% Triton X-100/PBS (PBT) for 3 h at 37°C. Samples were treated with primary

antibodies overnight at 4°C, washed in 0.1% PBT, and subsequently treated with secondary

antibodies conjugated to either Alexa Fluor 546 or Alexa Fluor 647 overnight at 4°C. For

nuclear counterstaining, we used Hoechst33342 (5 µg/ml, Dojindo Molecular Technologies,

#H342) or DAPI (Dojindo Molecular Technologies, #D523-10, 1:200). The samples were

mounted with 10 µL of 1% agarose gel onto a glass-based dish (Greiner Bio-One, #627871)

for obtaining stable images. Then, samples were immersed with CUBIC-R+ (Tokyo Chemical

Industry Co., # T3741) solution for 1 hour at 37°C or BABB solution (benzyl-alcohol and

benzyl-benzoate, 1:2) for 30 min at 23°C.

F-actin and tubulin labeling

To visualize F-actin and microtubules, we used SiR-Actin and SiR-tubulin, respectively [60,61],

according to the manufacturer's instructions (Cytoskeleton, Inc. #CY-SC001 and #CY-SC002).

We added 1 µM of far-red fluorogenic probes to the culture media, 1 hour before imaging.

Although SiR-Actin was derived from Jasplakinolide, which activated ERK in isolated lung

epithelial tissues in our experiments, ERK activation did not occur after treatment with SiR-

Actin in isolated lung epithelial tissues.

Fluorescence labeling of FGF

To visualizes internalized FGFs, we used an amine-reactive pH-sensitive pHrodo iFL Red STP

ester dye, according to the manufacturer's instructions (Thermo Fischer Scientific, #P36014).

For the imaging of FGF1, regardless of its localization inside or outside cells, we used the
Alexa Fluor 488 Microscale Protein Labeling Kit, according to the manufacturer's instructions
(Thermo Fischer Scientific, #A30006).
745

Microscopy
For live imaging, we used the incubator-integrated multiphoton fluorescence microscope
system (LCV-MPE, Olympus) with a 25× water-immersion lens (NA=1.05, WD=2 mm,
XLPLN25XWMP2, Olympus) or an inverted microscope (FV1200MPE-IX83, Olympus) with
a 30× silicone-immersion lens (NA=1.05, WD=0.8 mm, UPLSAPO30XS, Olympus). The
excitation wavelengths were set to 840 nm for the CFP of the ERK FRET biosensor, 930 nm
for Alexa Fluor 488-labeled FGF1, and 1040 nm for the pHrodo iFL Red-labeled FGFs and
the SiR probes (InSight DeepSee, Spectra-Physics). We used RDM690 IR cut filters (Olympus),
and the DM505 and DM570 dichroic mirrors. We used BA460-500, BA495-540, BA520-560,
BA575-630, and BA647/57 emission filters (Olympus) for CFP, Alexa Fluor 488, YFP or
FRET, iFL Red probes, and SiR probes, respectively. For fixed samples, images were obtained
using the Leica TCS SP8 confocal laser scanning platform equipped with the Leica HyD hybrid
detector, using the 20× objective lens (NA=0.75, WD=680 μm, HC PL APO CS2, Leica) and
40× objective lens (NA = 1.3, WD = 240 μm, HC PL APO CS2, Leica).
760

FRET image analysis
The median filter of a 3 × 3 window was processed to remove shot noise, and the background
signal was subtracted from each FRET and CFP channel. Then, the ratio of FRET intensity to
CFP intensity was calculated using a custom-made MATLAB (MathWorks) script. In the scale
bar, the color represents the FRET/CFP ratio and brightness represents the fluorescence
intensity of the FRET channel.
767

Three-dimensional mapping of different quantities
To determine the correlation between different quantities, such as ERK activity, F-actin signal,
tissue curvature, and tissue thickness (Figure S1A), we obtained 3D stacked images of lung
epithelial tissues with 1-μm-long z-intervals, and manually traced either the basal or apical side
of the cells. Next, we generated triangle meshes on the traced surface using "Iso2Mesh," a free
mesh generation toolbox [62]. Then, we calculated a discrete Gaussian curvature and a discrete
curvature on each node of generated meshes using a custom-made code according to the
definition proposed earlier [63]. Finally, we obtained all values except those for the curvatures

by averaging values around the nodes, and mapped them at each node. All processing was
performed using MATLAB.

Curvature measurement in 2D

We used a spline curve-based method described elsewhere [53]. Briefly, we manually traced the

basal side of epithelial cells from images, and determined discrete sampling points (x_i, y_i)

along traced curves at regular intervals. Upon fitting the 4 sampling points from i-1 to i+2 with

a cubic spline function, the function S_i at an interval $[x_i, x_{i+1}]$ was denoted as $S_i(x) =$

$a_i(x - x_i)^3 + b_i(x - x_i)^2 + c_i(x - x_i) + d_i$. Due to the definition of the curvature using the

formula $\kappa(x) = S''(1 + S'^2)^{-3/2}$, the curvature from the spline function was calculated as

follows:

$$\kappa_i(x) = \frac{6a(x - x_i) + 2b}{(1 + \{3a(x - x_i)^2 + 2b(x - x_i) + c\}^2)^{3/2}}.$$

The convex/concave surface of the lumen was assigned a positive/negative value, namely κ.

We defined tissue thickness as the length from the apical to the basal edge, which is vertical to

the traced curve at the sampling points.

Mechanical strain assay

We used a mechanical device (STREX, #STB-10) with a polydimethylsiloxane (PDMS)

chamber (STREX, #STB-CH-04). To compress isolated epithelium, we placed isolated lung

epithelial tissues in a 50% stretched state in the PDMS chamber, and added 1 mL of growth

factor-reduced Matrigel (Corning, #356231), and performed gelation for 15 min at 37°C. Then,

we added 2 mL of the culture medium supplemented with 500 ng/mL FGF1 into the chamber.

We conducted experiments involving epithelial tissues under 2D culture conditions in the

following manner. To obtain cell suspensions from primary tissues, lung tissues obtained via

dissection at E12.5 were treated with 1 mg/mL Collagenase/Dispase (Sigma-Aldrich,

#10269638001) in PBS for 50 min at 37°C while agitating the solution. Lung cells were then

dispersed into the solution by gentle pipetting, and the suspended solution was filtered using a

40 μm pore size mesh (Corning, #352340). After centrifugation at 200G for 5 min at 25°C, the

cell pellet was resuspended in a culture medium containing 10 μM Y-27632 and 10%

KnockOut Serum Replacement SR (Thermo Fischer Scientific, # 10828010) and the solution

was placed in a PDMS chamber pretreated with a collagen coating (Cellmatrix Type I-C, Nitta

Gelatin, #631-00771). After 2 days, the culture medium was exchanged with a medium not

containing Y-27632 and KSR, 1 hour before imaging. Notably, the PDMS chamber gradually

droops by becoming attached to the silicone oil used for objective lenses, because of which the

focus has to be adjusted manually every time an image needs to be obtained.

Statistical hypothesis testing

The number of cells or regions of interest analyzed (n) and the number of biological replicates

(N) are indicated in the figure legends. No particular statistical method was used to

predetermine the sample size. A minimum of N=3 independent experiments were performed,

based on previous studies in the field. No inclusion/exclusion criteria were used and all

analyzed samples were included in the analysis. No randomization was performed. Statistical

tests, sample sizes, test statistics, and *P*-values have been described in the main text. We

considered P-values < 0.05 to be statistically significant in two-tailed tests, and classified them

into the following 4 categories: * (p<0.05), ** (p<0.01), *** (p<0.001), and n.s. (not significant,

i.e., p $\geq$ 0.05).

Software

We used MATLAB (MathWorks) and Image J (National Institute of Health) software for

digital image processing. For graphics, MATLAB (MathWorks), Imaris (Bitplane), and Image

J (National Institute of Health) were used. MATLAB (MathWorks) was used for statistical

analysis.

Graph

In the boxplot, the central mark indicates the median, and the bottom and top edges of the box

indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme

data points not considered to be outliers, and the outliers are plotted individually using the '+'

symbol. All the graphs were drawn using MATLAB.

(II) Model analysis

(IIa) Apical length vs cell height at equilibrium

Here, we consider cells in a flat epithelial sheet to be identical, and assume that the basal edge

length is almost the same as the apical edge length for the simplicity of analysis. Then, potential

energy can be defined as follows:

$$U_{cell} = \frac{k_A}{2}(A - A^*)^2 + \frac{k_a}{2}(a - a^*)^2 + k_l l. \tag{1}$$

The first term represents the cell size constraint and its coefficient k_A, current cell area A, and

the target cell area A^*. The second and third terms each represent the regulation of cell edge

length at the apical and lateral side of cells a and l with controlling parameters k_a and k_l.

We set the second term as a function that converges to the variable target apical length a^*, as

explained later. Since the cell size was maintained even after a change in cell shape, caused by

the inhibitor assay, we set $A = A^*$. Then, upon considering that $A = rh$, we can redefine

equation 1 as follows:

$$U_{cell}^* = \frac{k_a}{2}(a - a^*)^2 + k_l \frac{A^*}{a}, \tag{2}$$

Now, we have two variables, a and a^*, and obtained equilibrium values at the minimum

potential energy value, i.e., $a_{eq} = (a|\partial U_{cell}^*/\partial a = 0)$ and $a_{eq}^* = (a^*|\partial U_{cell}^*/\partial a^* = 0)$ each,

as $a_{eq} = \sqrt{A^* k_l/k_b}$ and $a_{eq}^* = a_{eq}$. From these equations, it was noted that $a_{eq}^* \sim l_{eq}^{-1}$,

which indicates the existence of an inverse relationship between the apical length and cell

height at equilibrium. The analysis in this section is based on that of a previous study [64].

(IIb) Apical length vs epithelial sheet curvature

The analysis in this section is based on that of a previous study [65]. Here, we considered the

geometrical aspects of an epithelial monolayer, composed of smoothly-connected identical

tetragons along a line. The curvature κ at the midplane of the epithelial sheet is represented as

$$\kappa = \frac{2}{a + b}\left(\pi - 2\cos^{-1}\frac{b - a}{2l}\right), \tag{3}$$

869 where a, b, and l denote the apical, basal, and lateral edge lengths, respectively. Then, the

870 curvature with respect to the apical edge length is determined as follows:

$$\frac{\partial \kappa}{\partial a} = -\frac{2}{l(a + b)\sqrt{1 - \{(a - b)/4l\}^2}} - \frac{2}{(a + b)^2}\left(\pi - 2\cos^{-1}\frac{b - a}{2l}\right) < 0. \tag{4}$$

This indicates that the sheet curvature becomes small with an increase in the apical edge length.

(IIc) Mathematical modeling for multicellular dynamics

We modeled an epithelial monolayer using the vertex model framework [30–32]. In our model, a

single cell is represented as a polygon with four vertices and edges, which are shared by

neighboring cells. The dynamics of vertex position $\mathbf{r}_i$, where i is an index of the vertex, obeys

the equation of motion, based on the principle of least potential energy U, with a drag

coefficient η and luminal pressure p as follows:

884
$$\eta \dot{\mathbf{r}}_i = -\nabla_i U + f(p). \tag{5}$$

886 To determine the potential energy in the first term, we defined a minimal expression:

888
$$U = \sum_j \left\{ \frac{k_A}{2} \left(A_j - A_j^* \right)^2 + \frac{k_a}{2} \left(a_j - a_j^* \right)^2 + k_l l_j \right\} + \sum_i \frac{k_\theta}{2} \left\{ (\theta_{i_+} - \pi/2)^2 + (\theta_{i_-} - \pi/2)^2 \right\}. \tag{6}$$

The first term represents the cell size constraint with its coefficient k_A, current cell area A_j,

and the target cell area A_j^*. The second and third terms each represent the regulation of the cell

edge length at the apical and the lateral sides of cells a_j and l_j, with controlling parameters

k_a and k_l. As the apical edge is controlled by the ERK activation level, we set the second term

as a function that converges to the variable target apical length a_j^*; this has been explained

later. The fourth and fifth terms represent the bending energy of cells at the apical and the basal

sides, attributed to each vertex i; k_θ denotes the bending rigidity, and θ_{i_+} and θ_{i_-} are angles

at the vertex i across the cell edge.

The second term of Equation 5 represents the mechanical effect of luminal pressure [34,35,51]

on the apical edge of epithelial cells. The function f was defined as

901
$$f(p) = p\mathbf{n}_i, \tag{7}$$

where p is strength of pressure, and $\mathbf{n}_i$ is the normal vector at the apical vertex i. Note that this

function is effective only at vertices in the apical sides of cells.

To implement cell proliferation, we ensured that each cell was set with the cell timer τ_j,

and all cells had the same cell cycle length τ_{div}. We assumed that cell cycle progression

depended on the curvature; i.e., the cell cycle progresses when the cell curvature is not less

than zero $(\kappa_j \geq 0)$, and the cell cycle pauses when the cell curvature is less than zero $(\kappa_j < 0)$,

according to the previous report [36]. This is also attributable to the fact that cell proliferation in

the developing murine lung requires ERK activation, which can be observed in the distal tip

[23,66]. Cell division occurs and cells with almost equal sizes are produced when the cell timer

reaches the cell cycle length, $\tau_j = \tau_{div}$. Once cell division occurs, the τ_j value in one of the

daughter cells resets to zero and that in another is set at a value chosen stochastically from 0 to

$0.05\tau_{div}$ with a uniform distribution, to avoid perfect synchronization between neighboring

cells.

In our model, the target apical edge length depends on the cytoskeletal level c, observed

with a level of F-actin accumulation, as follows:

919
$$a_j^* = \zeta g(c_j) + a_0 \tag{8}$$

921
$$g(c_j) = \begin{cases} -1, & c_j \leq -1 \\ c_j, & -1 < c_j < 1 \\ 1, & 1 \leq c_j \end{cases} \tag{9}$$

923   Here, ζ denotes the maximum length after edge extension, and a_0 is the baseline for the apical

924   edge length. The saturation function g represents the physical limit for the change in edge

925   length. We modeled the cytoskeletal level c as follows:

927
$$\dot{c}_j = \alpha \tanh(\lambda \kappa_j) H(\kappa_j) + \beta - \gamma c_j. \tag{10}$$

The first term $\dot{c}_j$ represents the curvature-dependent contribution of FGF-ERK, along with the

magnitude parameter α and the sensitivity parameter λ, to the curvature κ. H is the step

function of the curvature, i.e., $H=1$ for $\kappa \geq 0$ and $H=0$ for $\kappa < 0$. The curvature κ_j at cell j

is calculated using the spline function-based method described above. As the cytoskeleton level

controls the apical target length observed with an increase in the monotonic function, as shown

in Equation 9, curvature-dependent apical extension/shrinkage is represented as a

positive/negative α value. The second term represents the curvature-independent contribution

of FGF-ERK, along with the parameter β. The third term represents the decay of cytoskeletons,

including F-actin depolymerization and severing. The rate γ controls the characteristic decay

time; the inverse value of γ denotes the retention time for cell curvature memory. The level of

curvature-dependent FGF-ERK activation was determined in Equation 10 because of its rapid

response.

(IId) Numerical simulations

We considered a manually traced line along the basal sides of cells in the image shown in

Figure 1 as the initial shape for the simulation, when the diameter of the distal tip D was in the

measured range. Upon an analysis of the tissue morphology, the number of cells was

determined according to the measured cell size, i.e., the average cell width and cell height,

which were 5 μm and 20 μm, respectively. The boundary in the proximal side was fixed in the

simulations shown in Figures 6B and 6C. For the simulations in Figure 6D–I, we used the

symmetrical image created based on the Figure 6A. Ordinary differential equations were

numerically solved using the forward Euler method, with a time step of 0.01, using MATLAB

(MathWorks). The simulations were terminated when the number of cells was more than 300.

Standard parameters were set as follows: $\eta = 1$, $k_A = 0.01$, $k_a = 1$, $k_l = 0.2$, $k_\theta =$

20, $\tau_{div} = 300$, $\alpha = 1$, $\beta = 0$, $\gamma = 0.001$, $\zeta = 3\,\mu m$, $\lambda = 100$, $p = 0.05$, and $D =$

$70\,\mu m$, unless otherwise noted. The k_A, k_a, k_l, k_θ, λ, and p values were determined by

numerical analyses using several criteria, under which the simulated curvature imitates the

experimental observations and became insensitive to the change in the values, as shown in the

supplementary figure S4. The maximum length of edge extension, i.e., the ζ value, was

determined from our measurements. A_j^* and a_0 values were set according to the initial

conditions, e.g., $A_j^* = 93\,\mu m$ and $a_0 = 4\,\mu m$ when $D = 70\,\mu m$.

 Supplementary figure legends

 Figure S1 3D mapping of ERK activity and tissue geometry

 (A) Mapping procedure for linking different quantities, including the ERK activity, epithelial

 sheet curvature, and sheet thickness. Details of the procedure are described in the Materials

 and Methods section. (B) The relationship between the ERK activity and discrete mean

 curvature. $\rho=0.61\pm0.04$, where ρ is Spearman's rank correlation coefficient. Data represent

 mean and SD values. n=427, N=3.

 Figure S2 FGF1 endocytosis from the basal side of an isolated lung epithelium

 (A) Time-lapse images of the bright field (upper), FGF1-pHrodo (middle), and FGF1-

 Alexa488 (lower). Red and blue circles in the bright field images indicate the tip apex and the

 cleft bottom, respectively. The time origin indicates the time of administration of labeled FGF1.

 FGF1 is distributed only in the Matrigel, not in the lumen, indicating that FGF1 internalization-

 based endocytosis occurs from the basal side of the epithelium. Notably, FGF1 distribution is

 almost homogeneous in the Matrigel (lower), indicating that no clear gradient was formed near

 the epithelium, even after ligand internalization. Scale bar, 50 μm.

 (B) FGF1-pHrodo intensity at the tip apex (red) and the cleft bottom (blue) over time. The

 response time of FGF1-pHrodo at the tip apex is 86 min. N=3.

 (C) FGF1-Alexa488 intensity in the Matrigel over time. The response time of FGF1-Alexa488

 in the Matrigel is 26 min. N=3.

 Figure S3 Morphological changes owing to chemical perturbations and cytoskeletal

 architecture

 (A) Morphological change in isolated epithelial cells after treatment with 1 μM PD0325901.

 Colors indicate the ERK activity level. Scale bar, 30 μm.

 (B, C) Isolated lung epithelium stained with SiR-Actin (B) and with SiR-Tubulin (C). Scale

 bar, 30 μm.

 (D) Relative change in cell height after treatment with cytoskeletal inhibitors, such as 1 μM

 Latrunculin A, 1 μM Nocodazole, and 30 μM Blebbistatin. N≥3 for each treatment.

 (E–E") Immunofluorescence images of cells interacting with anti-E-cadherin (blue), anti-

 phospho myosin light chain (pMLC, magenta) and the Hoechst (white) nuclear counterstain.

 Notably, the pMLC signal is significantly weaker at the tip of the epithelium than in smooth

 muscle cells around the stalk. In the magnified view (E"), the pMLC signal is clear at the

microtubule organizing centers (arrowhead), and is localized at the apical side of cells despite

its weak intensity. Scale bars, 100 μm (E, E') and 20 μm (E").

Figure S4 Simulation of the mathematical model

(A) Parameter dependency of the monolayer curvature in the simulation.

(B) Relative cell height, normalized by one at $\alpha=0$, as a function of α. Representative

morphologies of the virtual monolayer sheet are shown in the upper image.

(C) Tissue curvature as a function of luminal pressure p. Representative morphologies of

virtual monolayer tissues are shown in the upper image. N=5.

1004

Movie legends

Movie 1 Volumetric imaging of ERK activity in the murine lung

The 3D ERK activity map of a murine lung dissected at E12.5. Colors represent the ERK activity and the color scale corresponds to that in Figure 1B. Scale bar, 50 μm.

Movie 2 Endocytosis of FGF1 in the isolated lung epithelium

Time-lapse video of bright field (left), FGF1-Alexa488 (middle), and FGF1-pHrodo (right) in the isolated lung epithelium embedded within the Matrigel. The color scales correspond to those in Figure S2A. The time origin indicates the time of administration. Scale bars, 50 μm.

Movie 3 Dynamics of ERK activity in the growing isolated lung epithelium

Time-lapse video of bright field (left) and ERK activity (right) during the terminal bifurcation of an isolated lung epithelium. The color scale corresponds to that in Figure 2D. Scale bars, 30 μm.

Movie 4 ERK activity response in the isolated lung epithelium to compression parallel to the distal-proximal axis

Time-lapse video of ERK activity at the tip of the isolated lung epithelium against compression parallel to the distal-proximal axis. The color scale corresponds to that in Figure 3B. The time origin indicates the timing of compression. Scale bar, 30 μm.

Movie 5 ERK activity response of the isolated lung epithelium to compression vertical to the distal-proximal axis

Time-lapse video of ERK activity at the tip of an isolated lung epithelium against the compression vertical to the distal-proximal axis. The color scale corresponds to that in Figure 3C. The time origin indicates the timing of compression. Scale bar, 30 μm.

Movie 6 Morphological change in the isolated lung epithelium by ERK inactivation

Time-lapse video of ERK activity at the tip of the isolated lung epithelium after treatment with 1 μM PD0325901. The color scale corresponds to that in Figure S4A. Time origin indicates the timing of inhibitor administration. Scale bar, 30 μm.

Movie 7 Response of ERK activity and F-actin accumulation to acute simulation with FGF1

Time-lapse video of ERK activity (left) and SiR-Actin signal (right) at the tip of the isolated lung epithelium after treatment with 500 ng mL^{-1} FGF1. The color scales correspond to those in Figure 5A. The time origin indicates the timing of FGF1 administration. Scale bars, 30 μm.

Movie 8 Simulation for the morphogenesis of the proliferating epithelial monolayer in the regime of curvature-dependent apical extension

Colors represent the sheet curvature. The color scale corresponds to that in Figure 6E. Scale interval of the graph is 20 μm.

Movie 9 Simulation of virtual epithelial monolayer with a short retention time for curvature memory, γ=1

Colors represent the cellular F-actin levels. The color scale corresponds to that in Figure 6H. Scale interval of the graph is 20 μm.

Movie 10 Simulation of virtual epithelial monolayer with a medium retention time for curvature memory, γ= 10^{-3}

Colors represent cellular F-actin levels. The color scale corresponds to that in Figure 6H. Scale interval of the graph is 20 μm.

Movie 11 Simulation of virtual epithelial monolayer with a long retention time for curvature memory, γ= 10^{-6}

Colors represent cellular F-actin levels in. The color scale corresponds to that in Figure 6H. Scale interval of the graph is 20 μm.

Figure 1

A DMSO   PD0325901
              (MEK inhibitor)

200 µm

B DMSO
   PD03

Area [mm²]

Perimeter [mm]

C ERK activity (FRET/CFP)

1.6

1.0

100 µm

D Gaussian curvature [µm⁻²]

x10⁻⁴

-4

E High

Frequency

Low

Gaussian curvature [µm⁻²]

x10⁻⁴

-2

-4

0.9   1.25   1.6
ERK activity (FRET/CFP)

F Sheet thickness [µm]

0.9   1.25   1.6
ERK activity (FRET/CFP)

G Correlation coefficient

0.7

0.0

-0.7

ERK–Cunv   ERK–Thick

H ERK activity

1.0   1.6

50 µm

Figure 2

A Dispase | Matrigel | FGF

B FGF1 | FGF7 | FGF1+MEKi | w/o GF
100 μm

C ERK activity (FRET/CFP)
1.8 — 1.0
FGF1 | FGF7 | FGF1+MEKi | w/o GF
50 μm

D ERK activity (FRET/CFP)
1.0 — 1.8
ERK activity: 0 min | 200 min | 600 min | 1000 min
30 μm
Bright field: 0 min | 200 min | 600 min | 1000 min
30 μm

E Curvature [μm⁻¹] — ERK activity (FRET/CFP)
x10⁻²
Time [min]

F Lag time [min]

G Basal: *** | Apical: n.s.
Area [μm²]
Convex | Concave

Figure 3

Figure 4

A   FGF1   FGF1 + MEK inhibitor
30 µm

B   Basal
     Apical
     Cell height
     Cell width

C   Cell height [µm]   Cell width [µm]   Cell size [µm²]
     ***   *   n.s.
     FGF1 FGF1   FGF1 FGF1   FGF1 FGF1
           + MEKi   + MEKi   + MEKi

D   Before   After
     DMSO
     Latrunculin A
     Jasplakinolide
     20 µm

E   Relative change in cell height
     DMSO LatA CytoD Jasp Noco Bleb CK-666 SMIFH2

F   ERK activity (FRET/CFP) 1.8 ... 1.0
     50 µm
     F-actin intensity 1.0 ... 0.0

G   FGF1
     FGF1 + MEKi
     FGF7
     F-actin intensity
     ERK activity (FRET/CFP)

Figure 5

A

ERK activity (FRET/CFP)   0 min   30 min   60 min   90 min   120 min
1.8 — 1.0
30 μm

F-actin intensity   0 min   30 min   60 min   90 min   120 min
1.0 — 0.0

B
— FGF1
— FGF1 + MEKi
— PBS

Normalized ERK activity

0.8 — 1.4
Time [min]   0 — 60 — 120

C

Normalized F-actin intensity

1.0 — 3.0
Time [min]   0 — 60 — 120

D

Curvature [μm^{-1}]   x10^{-2}

0 — 3
Time [min]   0 — 60 — 120

Figure 6

A Curvature [μm⁻¹]
-0.1   0.1

30 μm   D

B (×10⁻²)
Curvature [μm⁻¹]

— D = 80 μm
— D = 100 μm
— D = 120 μm

Shrinkage   α   Extension

C
Curvature-dependent apical shrinkage | No regulation | Curvature-dependent apical extension
Time
α=−1   50 μm | α=0   50 μm | α=1   50 μm

D Time

E Curvature [μm⁻¹]
0.1 / −0.1

Curvature-dependent apical extension | No regulation | Global apical extension

F x10⁻²
Ave. curvature [μm⁻¹]
*** ***
Curv-dep | No reg | Global

G Correlation coefficient (Curvature vs Thickness)
*** ***
Curv-dep | No reg | Global

H Curvature memory time (γ⁻¹)
Short   Long

Curvature [μm⁻¹]
0.1 / −0.1

F-actin
1.0 / 0.0

I x10⁻²
Ave. curvature [μm⁻¹]
Correlation coefficient (Curvature vs F-actin)
Curvature memory time (γ⁻¹)

Figure 7

Curvature-sensitive ERK activation

Lung epithelium

Basal

Apical

Actin
polymerization

FGF

Basal

ERK

ERK-driven tissue bending