Isonicotinamide extends yeast chronological lifespan through a mechanism that diminishes nucleotides

Agata Kalita[1], Elisa Enriquez Hesles[1], Lindsey N. Power[1], Dezhen Wang[2], Pankaj K. Singh[2], and Jeffrey S. Smith[1*]

[1]Department of Biochemistry and Molecular Genetics, University of Virginia School of Medicine, Charlottesville, VA 22908; [2]Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, Omaha, NE 68198

*Corresponding Author
Jeffrey S. Smith
Department of Biochemistry and Molecular Genetics
University of Virginia School of Medicine
1340 Jefferson Park Ave.
Pinn Hall, Room 6229
Charlottesville, VA 22908

Phone: 434-243-5864
Email: jss5y@virginia.edu

ABSTRACT

Chronological lifespan (CLS) of budding yeast, *Saccharomyces cerevisiae,* is a commonly utilized model for cellular aging of non-dividing cells such as neurons. CLS is strongly extended by isonicotinamide (INAM), a non-metabolized isomer of the NAD^+ precursor nicotinamide (NAM), but the underlying mechanisms of lifespan extension remain uncharacterized. To identify potential biochemical INAM targets, we performed a chemical genetic screen with the yeast gene knockout (YKO) strain collection for INAM-hypersensitive mutants. Significantly enriched Gene Ontology terms that emerged included SWR1 and other transcription elongation factors, as well as metabolic pathways converging on one-carbon metabolism and contributing to nucleotide biosynthesis, together suggesting that INAM perturbs nucleotide pools. In line with this model, INAM effects on cell growth were synergistic with mycophenolic acid (MPA), which extends lifespan by reducing guanine nucleotide pools. Direct measurements of nucleotides and precursors by mass spectrometry indicated that INAM reduced nucleotides, including cAMP, at 24- and 96-hour time points post-inoculation. Taken together, we conclude that INAM extends CLS by perturbing nucleotide metabolism, which may be a common functional feature of multiple anti-aging interventions.

Keywords:
chronological aging/isonicotinamide/nucleotides/one-carbon metabolism/yeast

INTRODUCTION

Aging is a multifaceted process leading to decreased functioning over time, and ultimately, death of the cell or organism. Aging is also a major risk factor for many incapacitating conditions such as cardiovascular disease, cancer, and neurodegeneration (Niccoli & Partridge, 2012). Despite extraordinary variance in organism lifespans, many of the proteins and pathways involved in aging are evolutionary conserved (Smith *et al*, 2008; Bitto *et al*, 2015). This creates the possibility of identifying lifespan-extending mechanisms and interventions in model organisms that could be effective in understanding and combating human aging and associated disease.

Environmental interventions are promising in terms of potential application in human aging. Some of them, however, such as caloric restriction (CR), are difficult to implement in humans and can reduce the subjective quality of life (Dirks & Leeuwenburgh, 2006). Highly desired are interventions that do not require major changes in lifestyle, such as beneficial nutraceuticals or pharmaceuticals. Ideally, these types of molecules should be small, easily absorbed, and have limited negative side effects.

The budding yeast system has been effective in the identification and characterization of several such small molecule effectors of longevity, including the TOR inhibitor rapamycin (Powers *et al*, 2006). This unicellular eukaryote is highly suitable for testing environmental interventions using high throughput screening approaches due to its unicellular organization, short lifespan, and genetic malleability (Gershon & Gershon, 2000). Two types of aging can be studied in *S. cerevisiae*: chronological (Longo *et al*, 2012) and replicative (Mortimer & Johnston, 1959). Their corresponding longevity metrics are defined, respectively, as the survival of non-proliferating cells in stationary phase (CLS, chronological life span) (Longo *et al*, 2012) and the number of divisions that mother cells undergo before senescing (RLS, replicative life

span) (Mortimer & Johnston, 1959). Out of the two paradigms, chronological aging is more akin

to the aging of post-mitotic cells in mammalian organisms (such as neurons) (MacLean *et al*,

2001), whereas replicative aging is a better model for the aging of rapidly dividing, mitotically

active cells (e.g. skin stem cells) (Janssens & Veenhoff, 2016). CLS is more accurately described

as the survival of non-proliferating yeast cells in stationary phase, where they eventually lose the

ability to re-enter the cell cycle, even when provided with fresh nutrients (Postnikoff &

Harkness, 2014). The CLS system is therefore also used as a model for quiescence, since it

directly tests regrowth capacity of non-dividing cells (Mohammad *et al*, 2020). Importantly, CLS

is highly dependent on environmental factors, such as media composition and carbon source

levels, as well as the genetic strain background (Santos *et al*, 2020; Smith *et al*, 2016). Herein,

we take advantage of these features of CLS to study the mechanism of another longevity

promoting molecule, isonicotinamide (INAM).

Nicotinamide (NAM) is a water-soluble form of vitamin B3, a byproduct of the reaction

catalysed by NAD^+-dependent protein deacetylases (Sirtuins), and inhibits the deacetylation

reaction for these enzymes through a feedback mechanism (Sauve & Schramm, 2003; Bitterman

et al, 2002). INAM is an isomer of NAM commonly used for material synthesis and

pharmaceutical development due to its excellent cocrystallization properties that improve drug

bioavailability (Sánchez-Férez *et al*, 2019; Báthori *et al*, 2011). INAM does not occur in cells

naturally, but due to its structural similarity to NAM, has the potential for targeting NAM-

producing enzymes. Indeed, NAM and INAM both act as PARP-1 inhibitors in mammalian cells

(Sims *et al*, 1982). Of the two molecules, NAM function has been better characterized in yeast

and other systems, with relatively little known about INAM.

INAM, although structurally very similar to NAM, appears to elicit opposite effects to NAM in aging studies. In *S. cerevisiae*, NAM inhibition of Sir2 causes transcriptional silencing defects and destabilizes the ribosomal DNA (rDNA) locus, which drastically reduces RLS (Bitterman *et al*, 2002). INAM, however, functions as a Sir2 activating compound *in vitro* by relieving NAM inhibition (Sauve *et al*, 2005). INAM also promotes Sir2 function and RLS *in vivo* by indirectly maintaining elevated intracellular NAD^+ levels under conditions where NAD^+ is normally low, such as media lacking nicotinic acid (McClure *et al*, 2012). While INAM extension of RLS requires Sir2, INAM extends CLS in a sirtuin-independent manner (Wierman *et al*, 2015). Furthermore, unlike CLS extension induced by restriction of glucose or methionine, the INAM-induced longevity phenotype is not dependent on Snf1 (AMPK) activity (Wierman *et al*, 2017). The biochemical mechanisms of CLS extension by INAM treatment therefore remain uncertain, though INAM-treated cells do have increased resistance to acetic acid compared to controls (Wierman *et al*, 2015), which can be beneficial for CLS (Burtner *et al*, 2009). Consistent with this finding, transcriptome analysis previously revealed that INAM upregulates genes involved in cell wall synthesis (Wierman *et al*, 2015). However, because this RNA expression data was obtained very late in stationary phase (day 8), these changes in the transcriptional landscape might not reflect the primary mechanism and target of INAM, but rather the long-term consequences of its actions.

To better understand the primary mechanisms of INAM-induced lifespan extension, in this study we utilized an unbiased chemical genomics screening approach with the yeast knockout (YKO) strain collection to identify gene deletions that confer hypersensitivity to sublethal concentrations of INAM that normally extend CLS. The idea was that such genes may point toward biochemical pathways or cellular processes that INAM perturbs. The screen

unexpectedly implicated genes involved in transcriptional elongation and metabolic pathways related to one-carbon metabolism and nucleotide synthesis, leading to the conclusion that INAM-mediated extension of CLS involves reduction of nucleotide pools.

MATERIALS and METHODS

Yeast strains and media

The 'wild-type' laboratory strain in this study was BY4741 (*MAT*a *his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) (Brachmann *et al*, 1998), and most gene deletion mutants were obtained from the isogenic yeast knockout (YKO) collection (Winzeler *et al*, 1999). Detailed strain information is provided in Supplemental Table S1. Synthetic complete (SC) media with 2% glucose as a carbon source was used for most assays, with YPD used for cell propagation and selection prior to the experiments. For deletion strains with a KanMX cassette replacing the gene of interest, G418 was added to YPD at a final concentration of 200 mg/L. INAM and mycophenolic acid (MPA) were supplemented into SC media at the indicated concentrations. All liquid cultures and agar plates were grown at 30°C.

CLS assays

10 ml of SC media with 2% glucose and relevant supplements was inoculated with 100 µl of overnight culture. Liquid cultures were incubated on a rotating drum (TC-7, New Brunswick Scientific) in glass tubes with metal caps allowing for gas exchange. After 72 hours, the first measurement of colony forming units on YPD agar plates was made and this was treated as day 3 for the experiment (100% starting viability), to which all the other data was normalized. Measurements were taken every 3-4 days as previously described (Enriquez-Hesles *et al*, 2021; Wierman *et al*, 2017). At each time point, 20 µl of the cell suspension was removed from each tube and 10-fold serially diluted three times with sterile water. Next, 2.5 µl of each dilution

(1:10, 1:100, 1:1000) was spotted onto a YPD plate. After 16 hours, images of the spots were taken under a Nikon Eclipse E400 brightfield microscope at 30x magnification. Microcolonies within the spots were automatically counted from the digital images using OrganoSeg (Borten *et al*, 2018), with the parameters adjusted for yeast colony counting (Enriquez-Hesles *et al*, 2021). After accounting for the dilution factor, colony numbers from each day were divided by the number of colonies from the first time point (day 3) to give the viability metrics.

SGA screen

The *MAT*a haploid YKO collection, consisting of 4648 strains organized into 18 plates in 384-well format, was spotted onto Nunc Omnitray single-well plates containing YPD-agar with 200 µg/ml G418 using a floating pin manual replicator (VP 384FS, V&P Scientific). After 3 days growth, the replicator was used to transfer cells from colonies on the YPD plates to SC-agar Omnitrays supplemented with 0, 25, 50, 75, or 125 mM INAM. Colony growth for the 0, 25, 50, and 75 mM INAM plates was imaged with a Fluor Chem Q gel documentation system (ProteinSimple) after 2 days, while the 125 mM plates were imaged after 3 days incubation. The screen was performed in duplicate. SGAtools (http://sgatools.ccbr.utoronto.ca/) was then used to quantify colony growth from jpeg images, comparing the INAM-containing plates to control plates without INAM (Wagih *et al*, 2013). The score value of -0.3 was used as a cutoff for INAM sensitivity, and to limit false positives we required that both replicates reach this threshold for at least the 75- and 125-mM conditions. Most mutants with deletions of dubious open reading frames and genes without a confirmed function were omitted in further analysis, as their sensitivity would not provide any valuable information about the mechanism of action of INAM. Mutants with a score greater than +0.8 in both experiments at 125 mM INAM were considered for the identification of resistant strains.

Growth assays

For qualitative spots test growth assays, strains were patched onto YPD agar plates and grown overnight. The cell patches were resuspended in water at an OD_{600} of 1.0, then 10-fold serially diluted. Next, 2.5µl of each dilution was spotted onto SC agar plates containing 0, 25, 50, 75, or 125 mM INAM, and photos taken after 2 days of growth.

For IC_{50} assays, 10 ml SC cultures were incubated overnight in the roller drum, then added to wells of a 96-well plate at a starting OD_{600} of 0.1 in SC media containing a range of INAM concentrations (0-240 mM; 2-fold dilutions) and other supplements as indicated. The initial OD_{600} was measured with a Spectra Max M2 plate reader (Molecular Devices). Plates were then sealed with a sterile, gas-permeable membrane and incubated in a heated microplate shaker (Southwest Science, SBT1500-H) at 30°C for 8 hr at 900 RPM. Final OD_{600} was measured and the initial OD_{600} subtracted. Values were then normalized to fit the range 0-100%, where the control condition (no INAM) was treated as 100% and the highest INAM concentration as 0. These normalized growth values were then plotted on a graph using Origin 2018 (OriginLab, Northampton, MA, USA) and the sigmoidal fit was used to model a dose-response curve fitting the obtained data. IC_{50} was calculated as the concentration at which there was 50% viability. The 95% confidence intervals were also calculated by the Origin software from the 95% confidence bands. An additional benefit of this IC_{50} approach is precisely determining if mutant strains with intrinsically slow growth are sensitive to INAM, due to the assay being internally normalized to growth of a given strain in the control condition (no INAM), thus allowing direct comparison of strains with different growth rates.

Synergy assessment was performed similarly to the IC_{50} assays, with a combination of increasing MPA and INAM concentrations added in a matrix to a 96-well plate. Methanol

vehicle concentration was adjusted to be the same at all MPA concentrations used.

SynergyFinder (Ianevski *et al*, 2020) was used to analyze the synergy between compounds using baseline correction and a zero interaction potency (ZIP) model of synergy determination (Yadav *et al*, 2015).

Measurement of metabolite abundance by mass spectrometry

Cells were grown in 10 ml SC 2% glucose with or without 25 mM INAM on a roller drum. Cultures were started from overnight liquid cultures at OD_{600} adjusted to 0.05. Cell densities of cultures were measured before collection for normalization of metabolites. Cells were collected at log phase (~5 hr), 24 hr, and 96 hr. Intracellular metabolites were isolated as previously described (Shukla *et al*, 2015), with the extraction protocol modified for yeast. From these cultures, 9 ml was added to 20 ml of 100% cold (-80°C) LC/MS-grade methanol kept on dry ice (final methanol concentration: ~70% v/v) (Canelas *et al*, 2008). Samples were then kept on dry ice unless indicated otherwise. The quenched cell suspensions were spun down at 2000 × g in an Eppendorf 5810R centrifuge cooled to -9°C for 5 min. The cell pellets were resuspended in 1 ml 80% methanol (LC/MS grade) and transferred to screw-top microfuge tubes containing 0.5 mm acid-washed glass beads. A Mini-Beadbeater (Biospec products) was used to vigorously shake the microfuge tubes at 4°C for 3 cycles of 45 sec with 30-second pauses. Each tube was then punctured at the bottom with a needle, placed into a glass tube and spun down at -9°C for 5 minutes at 2000 RPM in the Eppendorf 5810R centrifuge to collect cell extracts. Beads were washed twice with 1.5 ml 80% methanol and spun down the same way (4 ml total pooled volume). The pellet was resuspended by vortexing and kept on ice for 15 minutes, followed by centrifugation at 3100 × g. Supernatants were transferred to two 2-ml microfuge tubes, and

evaporated in a Vacufuge™ Concentrator 5301 (Eppendorf, Hamburg, Germany) at 30°C until

20% volume remained. Remaining water was then removed by lyophilization.

Mass spectrometry

Metabolites were analyzed as described previously (Moshfegh *et al*, 2019; Gunda *et al*, 2016)

with a selected reaction monitoring (SRM) LC-MS/MS method with positive/negative ion

polarity switching using a Xevo TQ-S mass spectrometer. MassLynx 4.1 software was used to

calculate the peak areas for each metabolite. Peak areas were normalized to respective OD_{600} of

each sample. These normalized peaks were subsequently scaled relative to the mean peak area in

the control condition at each time point.

RNA isolation and quantitative RT-PCR

Total RNA was isolated from yeast cells harvested at log phase, 24 hr, or 96 hr using the acidic

phenol method (Collart & Oliviero, 2001). cDNA was then synthesized from 1 μg total RNA

using a Verso cDNA Synthesis Kit (Thermo Scientific), and quantitated by real-time PCR

reactions using SensiMix SYBR Hi-ROX Mastermix (Bioline) run on a StepOne Plus

thermocycler (Applied Biosystems). Transcript levels of tested genes were normalized to levels

of *TAF10* mRNA (Wierman *et al*, 2015). Sequences of primers used for qPCR are provided in

Table S2.

RESULTS

INAM-induced CLS extension is titratable and independent of auxotrophies

INAM is highly similar in chemical structure to NAM (Figure 1A), but not directly

utilized as an NAD^+ precursor (McClure *et al*, 2012). To begin characterizing the mechanism of

INAM-induced CLS extension, we first tested for dosage effects on the designer deletion strain BY4741, which contains several auxotrophies. Previous *in vivo* experiments with INAM were performed at a concentration of 25 mM due to its beneficial effect on Sir2-dependent transcriptional silencing (Sauve *et al*, 2005). In Figure 1B, INAM was titrated into CLS cultures at concentrations ranging from 2.5 to 50 mM, inducing moderate lifespan extension at 2.5 mM and maximal extension at 25 mM. INAM at 50 mM did not provide added benefit and actually caused mild toxicity at early time points. Since auxotrophic mutations in BY4741 significantly impacted CLS in previous studies (Gomes *et al*, 2007; Smith *et al*, 2016), we also confirmed that INAM at the optimal 25 mM concentration extended CLS of the related prototrophic strain, FY4 (Figure 1C). Furthermore, 25 mM INAM extended CLS of BY4741 when added to cultures at day 4, after stationary phase entry (Figure 1D). The effect was not as strong as adding INAM at the time of inoculation, but still suggested the underlying mechanism impacts both proliferating and quiescent cells.

INAM extends CLS independently of *SIR2* and *HST1*

Our group previously reported that Sir2 was not required for CLS extension by INAM (Wierman *et al*, 2015), which was unexpected given the positive effect of INAM on NAD$^+$ levels as cells enter the diauxic shift (McClure *et al*, 2012). Deleting *SIR2* slightly extended CLS in SC media (Fabrizio *et al*, 2005; Smith *et al*, 2007), while deleting its paralog, *HST1,* slightly shortened it (Smith *et al*, 2007). Given that significant functional redundancy exists between Sir2 and Hst1 (Fine *et al*, 2019; Hickman & Rusche, 2007; Li *et al*, 2013; Lamming *et al*, 2005), we hypothesized that deleting both sirtuins may block the INAM effect on CLS. As shown in Figure 1E, CLS of a *sir2Δ hst1Δ* double mutant without INAM addition was strongly reduced compared

to WT (BY4741), indicating that *SIR2* and *HST1* are indeed redundant in promoting CLS. However, 25 mM INAM still significantly improved lifespan of the *sir2Δ hst1Δ* strain by a similar percentage as observed for the WT strain (Figure 1E; statistics for all CLS assays are provided in Table S3), confirming that INAM works independently of Sir2 and Hst1 in the CLS system, while leaving the actual mechanism of INAM extension still unknown.

Chemical genomic screen reveals INAM sensitive mutant strains

In order to identify pathways affected by INAM, and thus provide insights about mechanisms of CLS extension, we took a non-targeted synthetic lethality screening approach using the yeast knockout (YKO) strain collection. We noticed that 50 mM INAM reduced the initial viability of BY4741 in CLS assays (Figure 1B), consistent with slightly reduced colony size in spot tests (Figure 2A). Higher INAM concentrations slowed colony growth even further (Figure 2A), suggesting the compound was perturbing one or more pathways important for cellular growth. We therefore hypothesized that applying INAM to the KO collection strains would identify clusters of hypersensitive mutants lacking genes that buffer INAM-impaired essential pathways or processes. Such an approach was successfully utilized in a number of screens for deletion mutants hypersensitive to specific chemicals, including the CLS-extending compound rapamycin (Xie *et al*, 2005). An analogous screen was also performed with nicotinamide (NAM) to identify sirtuin-dependent processes, implicating pathways that prevent genomic instability (Choy *et al*, 2016).

The *MAT*a haploid YKO strain collection was arrayed as colonies onto Nunc Omnitray plates containing YPD agar, then replicated onto SC 2% glucose agar plates containing 0, 25, 50, 75, or 125 mM INAM using a 384-pin applicator tool (Figure 2B). After 2 days, the growth of

each mutant on INAM-containing plates was quantitated relative to SC control plates using

SGAtools (Wagih *et al*, 2013). An example of growth inhibition is shown for *tho2Δ* in Figure 2C

(center colony). Note that growth of all strains was strongly reduced at 125 mM, similar to spot

assay results in Figure 2A. The full screen was independently performed twice, showing

significant correlation in SGAtools-generated scores, as shown for 75 mM INAM in Figure 2D

(R =0.42). We focused on mutants with sensitivity scores lower than -0.3 in both screen

replicates (Tables S4 and S5), as shown for the 240 such mutants from the 75 mM dataset in

Figure 2D (inset), fewer mutants met this stringent score criteria at 50 mM (173), and 25 mM

(150), which was expected as the degree of pathway disruption by INAM should be weaker at

lower concentrations. Only 39 genes were identified from all four INAM concentrations, though

there were greater overlaps for any two concentrations (Figure 2E). The 125 mM INAM

concentration caused a strong growth defect with the BY4741 control and most mutants (Figures

2A and C), and ~50% of the 326 mutants sensitive to this concentration did not overlap with

mutants from the other INAM concentrations (Figure 2E), suggesting there could be numerous

false positive hits in this portion of the screen. We therefore focused Gene Ontology (GO)

enrichment analysis on a collection of 341 genes identified from at least one of the 25, 50, or 75

mM INAM conditions (Table S4). Several 'Biological Process' GO terms with significant p-

values were related to transcriptional regulation, vacuolar transport, inositol phosphate

biosynthesis, and chromatin remodelling (Table S6). Significant 'Cellular Component' GO terms

covered the INO80 and SWR1 chromatin remodelling complexes, transcriptional elongation

factors, endosomes and the ESCRT complex (Table S6). With this functional information in

hand, we chose 61 candidate mutants for verification of INAM sensitivity using individual spot

test growth assays at 25, 50, and 75 mM (Figures S1 and S2). The mutants selected at this point

all had SGAtools scores lower than -0.3. INAM sensitivity was confirmed for 57 of 61 re-tested

deletion mutants (Figure S1 and S2), yielding a high positive predictive value (PPV) of 93.4%.

This high PPV also indicated the sensitivity cutoff score of -0.3 was highly stringent, suggesting

we likely missed some false-negative mutants. Therefore, based on the enriched GO terms in

Table S6 and the literature, we were able to identify 22 additional INAM-sensitive deletion

mutants (Figure S3).

Cells lacking *BNA7* are resistant to high INAM concentrations

Colony growth was strongly reduced by 125 mM INAM (Figure 2A and C), potentially

allowing for identification of INAM-resistant-mutants in the screen. While there were no

significantly enriched GO-terms identified among the mutants picked as resistant to 125 mM

INAM (Table S5; 0.8 cutoff score), we still tested 20 of these candidates for INAM-resistance in

spot test growth assays. The only mutant confirmed as resistant was *bna7Δ* (Figure S4A). We

verified this result by remaking the *bna7Δ::kanMX4* mutant from BY4741 and then retesting the

INAM-resistant phenotype. *BNA7* encodes formylkynurenine formamidase, an enzyme that

carries out the second step of *de novo* NAD$^+$ biosynthesis from tryptophan, releasing kynurenine

and formate as products (Figure S4B). This was intriguing because we previously isolated *bna2Δ*

in a screen for chronologically long-lived mutants (Matecic *et al*, 2010). However, deleting

BNA2, or other *BNA* genes in this pathway other than *BNA7*, had no effect on INAM sensitivity,

indicating a unique feature of lacking the Bna7 enzyme. The *bna7Δ* mutant was also long-lived

compared to BY4741, and 25 mM INAM significantly extended CLS even further (Figure S4C),

suggesting that INAM does not primarily target the Bna7 enzyme. Therefore, the mechanism of

how loss of Bna7 confers INAM resistance remains unclear, but could involve alterations in

enzyme substrate or product levels. Interestingly, this mutant was also resistant to the phenolic

compound, ferulic acid (Fletcher *et al*, 2019). Noteworthy, the Bna7 product, formate, is an input

to the one-carbon metabolism pathway.

SWR1 complex subunit mutants are sensitive to INAM

One of the strongest enriched GO terms for INAM hypersensitive mutants was the SWR1

complex (Table S6), a conserved ATP-dependent chromatin remodeler that exchanges H2A core

histones in nucleosome octamers with the H2A.Z variant encoded by *HTZ1* (Figure 3A;

(Santisteban *et al*, 2000; Mizuguchi *et al*, 2004)). Five SWR1 complex mutants (*bdf1Δ, vps72Δ,*

swc5Δ, swr1Δ, yaf9Δ) were identified from the screen at 25, 50, or 75 mM INAM concentrations,

and all were confirmed by spot test growth assays (Figures S1-S3). We also tested three

additional annotated subunit mutants found in the haploid YKO collection (*arp6Δ, swc3Δ,*

swc7Δ), but only *arp6Δ* and *swc3Δ* were confirmed as INAM sensitive. This actually made sense

because despite being part of the SWR1 complex, Swc7 was not required for H2A.Z binding or

histone replacement, and had minimal contributions to complex stability (Wu *et al*, 2009).

Importantly, the *htz1Δ* mutant strain itself was isolated from the screen at the 50, 75, and 125

mM INAM concentrations, and also confirmed as sensitive at 25 mM (Figure 3B). We note that

htz1Δ and SWR1 complex mutants were identified as sensitive to ferulic acid (Fletcher *et al*,

2019), highlighting another similarity between INAM and this phenolic compound.

Defects in the SWR1 complex were previously shown to extend CLS of a prototrophic

strain in nitrogen (glutamine)-rich media conditions, but not nitrogen (GABA)-poor media

(Garay *et al*, 2014). We were therefore curious if *swr1Δ* or *htz1Δ* mutants were also long-lived in

SC media with ammonium sulfate nitrogen conditions used for the INAM sensitivity screen. As

shown in Figure 3C, there were no significant differences in CLS between the mutants and WT

BY4741 without INAM, and their lifespans were fully extended by 25 mM INAM. Despite the requirement of SWR1 activity for resistance to high INAM concentrations, it was not required for CLS extension at the optimal 25 mM concentration. We concluded that deletion mutants isolated from the INAM-sensitivity screen will not necessarily be in pathways that directly regulate CLS, or its responsiveness to INAM.

Disruption of serine, cysteine and threonine metabolism confers sensitivity to INAM

Intriguing metabolism-related GO terms enriched in the 25, 50, or 75 mM INAM-sensitive mutant lists were *de novo* IMP biosynthesis (*ADE1*, *ADE4*, *ADE6*, *ADE5,7*), threonine metabolic process (*HOM2*, *HOM3*, *THR4*, and *GLY1*), and serine family amino acid biosynthetic process (*CYS4*, *GLY1*, *HOM2*, *HOM3*, *SER1*, *SER2*) (Table S6). *CYS3* was also identified from the 125 mM INAM concentration (Table S5). Figure 4A depicts connections between these enzymes, with the points of convergence at serine and glycine. Both of these amino acids are donors of one-carbon units in the folate/one-carbon metabolism pathway, whereby one-carbon moieties linked with tetrahydrofolate are major sources of building blocks for *de novo* purine synthesis (Appling, 1991). We previously reported that deletion of one-carbon metabolism and the *ADE* genes extended CLS in SC media (Matecic *et al*, 2010). Moreover, serine metabolism was strongly linked to CLS regulation, as serine supplementation extended CLS (Enriquez-Hesles *et al*, 2021a; Maruyama *et al*, 2016), and inactivation of *SER1* was identified as a major QTL for genetic CLS variation among strain backgrounds (Jung *et al*, 2018). We were able to recapitulate moderate CLS extension caused by impaired serine biosynthesis with a *ser2Δ* mutant (Figure 4B), but similar to the *swr1Δ* mutant, INAM fully extended CLS of the *ser2Δ* mutant (Figure

4B). Intriguingly, one-carbon moieties from the mitochondria are transported to the cytoplasm as formate, which is also a product of the Bna7 enzymatic reaction (Figure S4B).

The *ser2Δ* mutant visually showed strong sensitivity to 50 and 75 mM INAM on SC plates, but not 25 mM (Figure 4C). To more precisely quantify differences in sensitivity between strains, we switched from qualitative spot test growth assays to quantitative measurements of IC_{50} values in small liquid cultures (Figure 4D, see materials and methods for details). For BY4741, the IC_{50} with INAM in liquid cultures was between 60 and 70 mM depending on the experiment (Figure 4E). The IC_{50} of INAM for a *ser2Δ* mutant was only 38.5 mM (95% confidence intervals: 36.6 – 40.6 mM) (Figure 4E), consistent with the spot assay sensitivity results.

Having established a working IC_{50} assay, we employed it to test if the sensitivity of amino acid synthesis mutants such as *ser2Δ* and *thr1Δ* could be rescued by providing the respective amino acids. To this end, we tested INAM IC_{50} for BY4741 and deletion mutants when supplemented with 25 mM serine or threonine (Figure S5). As predicted, serine rescued low IC_{50} of the *ser2Δ* mutant without affecting BY4741. Similarly, threonine rescued low IC_{50} of *thr1Δ* and *hom3Δ* mutants defective in threonine synthesis, without affecting BY4741. These results suggested that abundant serine and threonine help cells tolerate the negative impact of high INAM concentrations.

INAM acts synergistically with MPA

Enriched GO terms for INAM sensitivity pointed to the SWR1 complex, transcriptional elongation complexes, and amino acid metabolic pathways clustering around one-carbon metabolism. Each of these terms is directly related to the availability of nucleotide pools. For

example, cells partially impaired for transcriptional elongation, including *swr1Δ* and *htz1Δ* mutants (Santisteban *et al*, 2011), are sensitive to mycophenolic acid (MPA) or 6-azauracil (6AU) (Desmoucelles *et al*, 2002), inhibitors of inosine 5'-monophosphate dehydrogenase (IMPDH) that deplete guanine nucleotide pools (Escobar-Henriques *et al*, 2001; Sweeney *et al*, 1972; Qiu *et al*, 2000). We therefore compared INAM-sensitive mutants identified in our screen to previously identified MPA-sensitive strains (Desmoucelles *et al*, 2002). As shown in Figure 5A, there was significant overlap between the datasets, with 45.1% (46/102) of MPA-sensitive mutants also identified as sensitive to at least one INAM concentration in the primary screen, but only 9.1% (46/504) of all candidate INAM-sensitive mutants, including at 125 mM, were also sensitive to MPA. To determine if such overlap might indicate a functional similarity between the compounds, perhaps even inhibiting the same target, we tested for synergy between MPA and INAM by measuring effects of combining the compounds at increasing concentrations on cellular growth in the IC_{50} assay. Visual representation of the synergistic parameter δ for each of the concentration combinations revealed clear synergistic effects at relatively low concentrations where they did not inhibit growth individually (Figure 5B). The strong synergistic interaction indicated that INAM and MPA likely target related pathways within an interaction network (Yin *et al*, 2014), but not the same enzyme, as this would lead to an additive effect.

MPA was known to extend RLS of *S. cerevisiae* (Liu *et al*, 2020; Sarnoski *et al*, 2017), and CLS of another yeast species, *Schizosaccharomyces pombe* (Stephan *et al*, 2013). As shown in Figure 5C, MPA also extended CLS of *S. cerevisiae* at 1 μM, and even more effectively at 10 μM. This effect was completely abolished by 0.1 mM guanine supplementation. Guanine is imported and then salvaged into GMP by hypoxanthine guanine phosphoribosyltransferase (Hpt1), thus bypassing the need for IMPDH, the target of MPA inhibition. This result confirmed

that MPA extended CLS through depletion of guanylic nucleotides. In contrast, CLS extension

by 25 mM INAM was not prevented by guanine supplementation (Figure 5D), indicating that

IMPDH was likely not a primary target of INAM, and thus consistent with the synergistic effects

of MPA and INAM on cell growth.

INAM modifies nucleotide metabolism

Since CLS extension by INAM was not prevented by guanine addition, and given the

involvement of serine and threonine, both of which support nucleotide pools (Kastanos *et al*,

1997; Hartman, 2007), we decided to test the impact of 25 mM INAM on a wider range of

nucleotides, nucleosides, and nucleobases. Levels of such metabolites in extracts from cells at

log phase, late diauxic shift (24 hr), or stationary phase (96 hr) were quantified by mass

spectrometry. We reasoned that nucleotide changes occurring at log phase and/or 24 hr of liquid

growth would reflect changes occurring in colonies growing on agar plates, specifically inner

cells of the colony are exposed to nutrient stress (Váchová *et al*, 2012). As shown in Figure 6A,

INAM significantly reduced several bases and nucleosides during log phase, which then

extended to nucleotides, including cAMP and cGMP, at 24 and 96 hrs (Figure 6A and B). By 96

hrs, some nucleotides (not the triphosphates) were elevated in INAM-treated cells compared to

the control (Figure 6A). This effect was most prominent with uracil, uridine, UMP, and UDP at

96 hrs (Figure S7A). We hypothesized this result could be related to the *ura3Δ* mutation in

BY4741, which blocks UMP synthesis from orotidine-5'-phosphate and makes cells auxotrophic

for uracil (Figure S7A). While supplementing uracil at 4x the normal concentration in SC media

extended CLS, it was less effective than 25 mM INAM (Figure 6C). Furthermore, INAM still

fully extended CLS in combination with the supplemented uracil without an additive effect. The

ura3Δ0 mutation in BY4741 applies pressure on cells to salvage uracil from the recycling of RNA, and INAM may augment this activity, but may also function elsewhere.

To complement the nucleotide quantification, quantitative RT-PCR was used to assay expression of several genes that function in nucleotide metabolism. As shown in Figures S6 and S7A, most tested genes showed reduced expression at 24 and 96 hrs, which was expected as cells enter stationary phase. However, INAM increased expression of *RNR3* and *IMD2* at 96 hrs. *RNR3* encodes a minor large ribonucleotide reductase subunit that is nearly undetectable under normal growth conditions, but strongly induced by DNA damaging agents, including hydroxyurea (HU) (Domkin *et al*, 2002). *IMD2* encodes a major isoform of inosine monophosphate dehydrogenase, the target of MPA. Expression of *IMD2*, but not the other *IMD* genes, is also induced by MPA (Escobar-Henriques & Daignan-Fornier, 2001). Both of these changes were consistent with impacts of INAM on nucleotide pools, but not indicative of a single mechanism. Interestingly, almost all the genes we tested by RT-PCR were elevated at day 8 in INAM compared to the control (Figure S7B), including *ADO1*, *AAH1*, *IMD4*, and *RNR1*, each of which was identified as upregulated by INAM in an earlier microarray transcriptomics study (Wierman *et al*, 2015). Taken all together, we conclude that INAM extends CLS through a mechanism that perturbs nucleotide metabolism.

DISCUSSION

CLS and Sir2

INAM first came to our attention because this synthetic product prevented feedback inhibition of Sir2 by the NAM product of its enzymatic reaction (Sauve *et al*, 2005). Unexpectedly, it extended RLS by indirectly maintaining elevated NAD^+ levels, most likely by

increasing flux through the NAD$^+$ salvage pathway (McClure *et al*, 2012), potentially through a

hormesis effect. When we examined the impact of INAM on CLS, it was found to strongly

extend lifespan independently of Sir2 (Wierman *et al*, 2015). The impact of Sir2 on CLS

depends on media composition and genetic strain background (Wierman & Smith, 2014), with

most lab media conditions resulting in CLS extension when Sir2 was deleted or inhibited

(Casatta *et al*, 2013; Fabrizio *et al*, 2005; Orlandi *et al*, 2017; Smith *et al*, 2007). Here, we found

that CLS was drastically reduced in cells lacking both *SIR2* and *HST1*, consistent with several

instances of these two paralogs compensating for each other in other assay systems (Hickman &

Rusche, 2007; Li *et al*, 2013). This new result unifies the positive role of Sir2 in yeast cell

survival, both replicatively and chronologically, as well as with sirtuins in higher eukaryotes.

Given the ability of INAM to maintain NAD$^+$ levels in cells entering stationary phase (McClure

et al, 2012), we were surprised that INAM still extended CLS of the *sir2Δ hst1Δ* double mutant,

although this pointed to INAM impacting CLS through a sirtuin-independent mechanism, thus

leading to the screen in this study.

INAM effects compared to NAM

The sensitivity selection criteria chosen in this study were very strict in order to minimize

the number of false positives that can occur with YKO screens. We know that INAM sensitive

mutants were missed because out of the deletion strains selected for testing based on follow-up

to primary screen hits and literature searching, 22 were sensitive to INAM, even though they did

not meet the original screening criteria. Such mutants turned out to be, on average, less INAM-

sensitive than those identified by the original criteria (score <-0.3). This result suggests that the

screen likely identified the most sensitive strains to INAM, but not all. Even at this point of the

screen, the candidates identified were sufficient to implicate nucleotide metabolism as a likely target for perturbation by INAM.

An earlier screen performed with the YKO strain collection for deletion mutants sensitive to 120 mM nicotinamide (NAM) identified genes that function in sister chromatid cohesion and maintenance of genome stability (Choy *et al*, 2016), cellular processes that were not significantly enriched in our screen for INAM-sensitive mutants. However, 21 of these 59 NAM-sensitive mutants overlapped with our list of candidate INAM-sensitive mutants from Table S4, including *htz1Δ*. From the supplemental tables (Choy *et al*, 2016), we also extracted NAM-sensitive mutants with normalized colony growth ratios of <0.5 in both screen replicates (107 genes), revealing significantly enriched GO terms of DNA repair (as expected), but also *de novo* IMP biosynthesis (*ADE1*, *ADE2*, *ADE4*, *ADE6*, *ADE8*). Several *ADE* gene mutants were also identified from our INAM-sensitivity screen, suggesting that NAM and INAM share some level of overlap in their physiological effects on yeast cells, as might be expected for compounds with such similar chemical structures (Figure 1A). There are also significant differences, including that NAM appears to be less toxic than INAM to yeast cells at high concentrations. Additionally, relatively low INAM concentrations (2.5 or 10 mM) extend CLS, while low NAM (1 or 5 mM) shortens CLS, most likely due to sirtuin inhibition (Smith *et al*, 2007). It remains possible that NAM at higher concentrations could also extend CLS similarly to INAM, thus overriding the inhibitory effect on sirtuins.

Interpretation and limitations of INAM sensitivity screening

It is important to note that the sensitivity of deletion strains to a given compound does not necessarily result from direct participation of the deleted gene in the compound-impaired pathway. For example, there are multidrug-resistance conferring genes such as non-specific

efflux membrane pumps that make cells sensitive to a wide array of toxic compounds when deleted. In *S. cerevisiae*, strains with deletions of genes relevant for vacuolar structure and function, follicular transport, cell structure, and lipid metabolism are also susceptible to many compounds with diverse, unconnected modes of action (Parsons *et al*, 2004). Only five of all strains that we confirmed as sensitive to INAM were also sensitive to the wider range of other compounds: *gal11Δ, ies6Δ, spc72Δ, vrp1Δ* and *def1Δ* (Parsons *et al*, 2004). Interestingly, several deletion strains previously identified as MPA-sensitive were also sensitive to rapamycin (Desmoucelles *et al*, 2002; Chan *et al*, 2000), three of which were sensitive to INAM in our study *(ser2Δ, hom2Δ,* and *cdc10Δ)*. Additionally, *hom3Δ, rnr1Δ, ser1Δ*, and *aat2Δ* mutants involved in serine, aspartate/threonine synthesis or ribonucleotide reductase activity were all sensitive to rapamycin (Chan *et al*, 2000). Importantly, such mutants are not generally sensitive to a broader spectrum of compounds (Parsons *et al*, 2004), implying there could be functional overlap between these CLS extending compounds at the level of nucleotide metabolism.

INAM effects on nucleotide metabolism

Our chemical genetic screen indicated that perturbing the pathways for serine, threonine, aspartate or cysteine synthesis conferred INAM sensitivity. Each of these pathways converge on the reaction of serine conversion to glycine, in which the methylene group of serine is incorporated into the folate cycle/one-carbon metabolism pathway by serine hydroxymethyltransferases (Kastanos *et al*, 1997). This pathway is the main source of one-carbon moieties in the cell, which in turn, together with glycine, provides key substrates for *de novo* purine nucleotide biosynthesis (Kastanos *et al*, 1997). The one-carbon pathway reactions can occur either in the mitochondria or in the cytoplasm, depending on the availability of nutrients (Kastanos *et al*, 1997). *De novo* purine synthesis occurs in the cytosol, so single carbon

groups formed in the mitochondria must be exported there. The trifunctional enzyme methylenetetrahydrofolate dehydrogenase (Mis1 in yeast, MTHD2 in mammals) releases the one-carbon moiety from mitochondria in the form of formic acid, where it can be re-incorporated into the cytosolic branch of the one-carbon pathway for use in purine biosynthesis (Kastanos *et al*, 1997). Formic acid is also a product of the reaction catalyzed by Bna7 (Wogulis *et al*, 2008) – a protein whose absence conferred INAM resistance. Formic acid might therefore be a metabolite that connects the observations made in this study between INAM and potential negative regulation of one-carbon metabolism. This is especially intriguing given that one-carbon metabolism is rapidly emerging as an important regulatory hub for longevity pathways (Maitra *et al*, 2020; Annibal *et al*, 2021; Enriquez-Hesles *et al*, 2021). Further work is necessary to explain in detail the effect of INAM on formic acid levels and one-carbon metabolism. In particular, identification of the molecular target of INAM will contribute to a better understanding of the action of this compound and its potential utility.

Unlike the effects of MPA in previous studies (Hyle *et al*, 2003), 25 mM INAM did not have a dramatic effect on nucleotide pools in exponentially growing cells, but the reduction in trinucleotides became noticeable in cells grown for 24 or 96 hours. We hypothesize that INAM perturbs the flux of nucleotide availability further upstream of MPA, which directly targets the IMP-dehydrogenase enzyme to produce an acute impact on guanine nucleotides. The effect of INAM on nucleotides becomes more noticeable when pools are naturally being depleted as cells enter stationary phase (Osório *et al*, 2005). We anticipate that the negative effect of INAM on trinucleotide pools would be stronger in proliferating cells if the concentration was sufficiently increased to slow growth rates, such as 50, 75, or 125 mM. The partial effect on nucleotides at 25 mM and lower could be inducing a hormesis response that is beneficial for longevity, similar to

that proposed for MPA (Liu *et al*, 2020). Evidence for this idea comes from earlier findings that 25 mM INAM prevents depletion of NAD^+ as cells enter stationary phase, increases expression of NAD^+ biosynthesis and salvage pathway genes (McClure *et al*, 2012), and incurs cell resistance to high exogenous acetic acid levels (Wierman *et al*, 2015). In this current study we also found that genes involved in the salvage of other nucleotides are induced by INAM relative to the control condition, especially at day 8 of the aging assays. Additionally, we observed that pyrimidine salvage intermediates were generally reduced in log phase cells, but became elevated in cells grown for 96 hr, suggesting upregulation of pyrimidine nucleotide salvage pathways.

cAMP and chronological aging

The strongest negative effects of INAM on nucleotides were for cAMP and cGMP at the 96 hr timepoint. Extracellular glucose is sensed by the G-protein coupled receptor (Gpr1/Gpa2) to activate cAMP synthesis by adenylyl cyclase (Cyr1). cAMP then activates protein kinase A (PKA), which regulates cell growth, metabolism, and stress resistance. For example, PKA directly represses the key transcription factor Rim15, preventing it from entering the nucleus (Broach, 2012). Glucose restriction reduces cAMP/PKA signalling, allowing Rim15 to enter the nucleus and activate genes with Post-Diauxic Shift (PDS) or STress Response Element (STRE) motifs, which mediates RLS extension (Wei *et al*, 2008). cAMP levels are typically high in proliferating yeast cells and then reduced as cells enter stationary phase, controlled by the balance between the activities of adenylyl cyclase and the cAMP phosphodiesterases, Pde1 and Pde2 (Park *et al*, 2005). Little is known about the role of cGMP in yeast cells, but it has been proposed to help fine tune cAMP levels (Cardarelli *et al*, 2019). The observed cAMP reduction in INAM-treated cells at 96 hr is consistent with lifespan extension, and suggests stimulation of Pde2 and/or reduced Cyr1 activity. Either way, the reduction in cAMP caused by INAM could

be enhancing stationary phase entry and quiescence maintenance, thus contributing to CLS extension.

INAM effects in mouse and humans

Could INAM be used as an intervention in humans? INAM actually inhibits the activity of PARP-1 in mammalian cells, similar to the effect of NAM (Ueda *et al*, 1995). No protein similar to PARP-1 in sequence or structure is encoded in the yeast genome (Citarelli *et al*, 2010). Due to inhibition of PARP-1, INAM has been studied in the past in mice. INAM exhibited anti-inflammatory and analgesic effects (Godin *et al*, 2011), also protected β-cells of the pancreas from streptozotocin-induced damage and apoptosis, and from diabetes caused by such damage (Fukaya *et al*, 2013). Such studies were not performed in PARP-1 knockout mice as a control, so it cannot be excluded that some of these positive effects are not due to PARP-1 inhibition by INAM, but occur through a path similar to the one described in this paper.

MPA is used in immunosuppressive therapy because it inhibits the proliferation of lymphocytes (Mitsui & Suzuki, 1969). Therefore, it is possible that the observed anti-inflammatory effect of INAM is due to a decrease in nucleotide levels by this compound. There are also data relevant to safety of INAM. For example, it has been shown that administration of this compound throughout the lifespan of mice does not cause increased cancer incidence (Toth, 1983). Unfortunately, this 1983 publication did not measure life expectancy data for the control group, which makes it impossible to assess the impact of INAM on lifespan in this experiment. Future INAM studies in metazoan organisms therefore appear warranted.

ACKNOWLEDGMENTS

We thank members of the Smith lab, David Auble, and Marty Mayo for helpful discussions. E.E.H was supported by the Medical Scientist Training Program (MSTP) training grant T32GM007267, the Cell and Molecular Biology (CMB) training grant T32GM008136, and an individual National Research Service Award from the NIH, F30AG067760. The study was supported by NIH grants RO1GM075240 and RO1GM127394 to J.S.S., as well as RO1CA216853 and RO1CA163649 to P.K.S.

REFERENCES

Annibal A, Tharyan RG, Schonewolff MF, Tam H, Latza C, Auler MMK & Antebi A (2021) Regulation of the one carbon folate cycle as a shared metabolic signature of longevity. *Nat Commun* 12: 3486

Appling DR (1991) Compartmentation of folate-mediated one-carbon metabolism in eukaryotes. *FASEB J* 5: 2645–2651

Báthori NB, Lemmerer A, Venter GA, Bourne SA & Caira MR (2011) Pharmaceutical Co-crystals with isonicotinamide-vitamin B3, clofibric acid, and diclofenac-and two isonicotinamide hydrates. *Cryst Growth Des* 11: 75-87

Bitterman KJ, Anderson RM, Cohen HY, Latorre-Esteves M & Sinclair DA (2002) Inhibition of silencing and accelerated aging by nicotinamide, a putative negative regulator of yeast Sir2 and human SIRT1. *J Biol Chem* 277: 45099–45107

Bitto A, Wang AM, Bennett CF & Kaeberlein M (2015) Biochemical Genetic Pathways that

Modulate Aging in Multiple Species: Figure 1. *Cold Spring Harb Perspect Med* 5: a025114

Borten MA, Bajikar SS, Sasaki N, Clevers H & Janes KA (2018) Automated brightfield morphometry of 3D organoid populations by OrganoSeg. *Sci Rep* 8: 1–10

Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P & Boeke JD (1998) Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: A useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* 14: 115–132

Broach JR (2012) Nutritional control of growth and development in yeast. *Genetics* 192: 73–105

Burtner CR, Murakami CJ, Kennedy BK & Kaeberlein M (2009) A molecular mechanism of chronological aging in yeast. *Cell Cycle* 8: 1256-1270

Canelas AB, Ras C, ten Pierick A, van Dam JC, Heijnen JJ & van Gulik WM (2008) Leakage-free rapid quenching technique for yeast metabolomics. *Metabolomics* 4: 226–239

Cardarelli S, Giorgi M, Poiana G, Biagioni S & Saliola M (2019) Metabolic role of cGMP in S. cerevisiae: the murine phosphodiesterase-5 activity affects yeast cell proliferation by altering the cAMP/cGMP equilibrium. *FEMS Yeast Res* 19: foz016

Casatta N, Porro A, Orlandi I, Brambilla L & Vai M (2013) Lack of Sir2 increases acetate consumption and decreases extracellular pro-aging factors. *Biochim Biophys Acta - Mol Cell Res* 1833: 593–601

Chan TF, Carvalho J, Riles L & Zheng XFS (2000) A chemical genomics approach toward understanding the global functions of the target of rapamycin protein (TOR). *Proc Natl Acad Sci U S A* 97: 13227–13232

Choy JS, Qadri B, Henry L, Shroff K, Bifarin O & Basrai MA (2016) A genome-wide screen

with nicotinamide to identify sirtuin-dependent pathways in *Saccharomyces cerevisiae*. *G3 Genes, Genomes, Genet* 6: 485–494

Citarelli M, Teotia S & Lamb RS (2010) Evolutionary history of the poly(ADP-ribose) polymerase gene family in eukaryotes. *BMC Evol Biol* 10: 1–26

Collart MA & Oliviero S (2001) Preparation of yeast RNA. *Curr Protoc Mol Biol* Chapter 13: Unit13.12

Desmoucelles C, Pinson B, Saint-Marc C & Daignan-Fornier B (2002) Screening the yeast 'Disruptome' for mutants affecting resistance to the immunosuppressive drug, mycophenolic acid. *J Biol Chem* 277: 27036–27044

Dirks AJ & Leeuwenburgh C (2006) Caloric restriction in humans: Potential pitfalls and health concerns. *Mech Ageing Dev* 127: 1–7

Domkin V, Thelander L & Chabes A (2002) Yeast DNA damage-inducible Rnr3 has a very low catalytic activity strongly stimulated after the formation of a cross-talking Rnr1/Rnr3 complex. *J Biol Chem* 277: 18574–18578

Enriquez-Hesles E, Smith Jr DL, Maqani N, Wierman MB, Sutcliffe MD, Fine RD, Kalita A, Santos SM, Muehlbauer MJ, Bain JR, *et al* (2021) A cell-nonautonomous mechanism of yeast chronological aging regulated by caloric restriction and one-carbon metabolism. *J Biol Chem* 296: 100125

Escobar-Henriques M, Balguerie A, Monribot C, Boucherie H & Daignan-Fornier B (2001) Proteome analysis and morphological studies reveal multiple effects of the immunosuppressive drug mycophenolic acid specifically resulting from guanylic nucleotide

depletion. *J Biol Chem* 49: 46237-46242

Escobar-Henriques M & Daignan-Fornier B (2001) Transcriptional regulation of the yeast GMP
synthesis pathway by its end products. *J Biol Chem* 276: 1523–1530

Fabrizio P, Gattazzo C, Battistella L, Wei M, Cheng C, McGrew K & Longo VD (2005) Sir2
blocks extreme life-span extension. *Cell* 123: 655–667

Fine RD, Maqani N, Li M, Franck E & Smith JS (2019) Depletion of limiting rDNA structural
complexes triggers chromosomal instability and replicative aging of Saccharomyces
cerevisiae. *Genetics* 212: 75-91

Fletcher E, Gao K, Mercurio K, Ali M & Baetz K (2019) Yeast chemogenomic screen identifies
distinct metabolic pathways required to tolerate exposure to phenolic fermentation
inhibitors ferulic acid, 4-hydroxybenzoic acid and coniferyl aldehyde. *Metab Eng* 52: 98–
109

Fukaya M, Tamura Y, Chiba Y, Tanioka T, Mao J, Inoue Y, Yamada M, Waeber C, Ido-
Kitamura Y, Kitamura T, *et al* (2013) Protective effects of a nicotinamide derivative,
isonicotinamide, against streptozotocin-induced β-cell damage and diabetes in mice.
Biochem Biophys Res Commun 442: 92–98

Garay E, Campos SE, González de la Cruz J, Gaspar AP, Jinich A & DeLuna A (2014) High-
resolution profiling of stationary-phase survival reveals yeast longevity factors and their
genetic interactions. *PLoS Genet* 10: e1004168

Gershon H & Gershon D (2000) The budding yeast, *Saccharomyces cerevisiae*, as a model for
aging research: A critical review. *Mech Ageing Dev* 120: 1–22

Godin AM, Ferreira WC, Rocha LTS, Seniuk JGT, Paiva ALL, Merlo LA, Nascimento EB, Bastos LFS & Coelho MM (2011) Antinociceptive and anti-inflammatory activities of nicotinamide and its isomers in different experimental models. *Pharmacol Biochem Behav* 99: 782–788

Gomes P, Sampaio-Marques B, Ludovico P, Rodrigues F & Leão C (2007) Low auxotrophy-complementing amino acid concentrations reduce yeast chronological life span. *Mech Ageing Dev* 128: 383–391

Gunda V, Yu F & Singh PK (2016) Validation of metabolic alterations in microscale cell culture lysates using hydrophilic interaction liquid chromatography (HILIC)-tandem mass spectrometry-based metabolomics. *PLoS One* 11: e0154416

Hartman 4th JL (2007) Buffering of deoxyribonucleotide pool homeostasis by threonine metabolism. *Proc Natl Acad Sci* U S A104: 11700–11705

Hickman MA & Rusche LN (2007) Substitution as a mechanism for genetic robustness: The duplicated deacetylases Hst1p and Sir2p in *Saccharomyces cerevisiae*. *PLoS Genet* 8: e126

Hyle JW, Shaw RJ & Reines D (2003) Functional distinctions between IMP dehydrogenase genes in providing mycophenolate resistance and guanine prototrophy to yeast. *J Biol Chem* 278: 28470–28478

Ianevski A, Giri AK & Aittokallio T (2020) SynergyFinder 2.0: visual analytics of multi-drug combination synergies. *Nucleic Acids Res* 48: W488–W493

Janssens GE & Veenhoff LM (2016) Evidence for the hallmarks of human aging in replicatively aging yeast. *Microb Cell* 3: 263–274

Jung PP, Zhang Z, Paczia N, Jaeger C, Ignac T, May P & Linster CL (2018) Natural variation of chronological aging in the *Saccharomyces cerevisia*e species reveals diet-dependent mechanisms of life span control. *NJP Aging Mech Dis* 12: 3

Kastanos EK, Woldman YY & Appling DR (1997) Role of mitochondrial and cytoplasmic serine hydroxymethyltransferase isozymes in *de novo* purine synthesis in *Saccharomyces cerevisiae*. *Biochemistry* 36: 14956–14964

Lamming DW, Latorre-Esteves M, Medvedik O, Wong SN, Tsang FA, Wang C, Lin SJ & Sinclair DA (2005) Cell biology: HST2 mediates SIR2-independent life-span extension by calorie restriction. *Science* 309: 1861-1864

Li M, Valsakumar V, Poorey K, Bekiranov S & Smith JS (2013) Genome-wide analysis of functional sirtuin chromatin targets in yeast. *Genome Biol* 14: R48

Liu P, Sarnoski EA, Olmez TT, Young TZ & Murat Acar · (2020) Characterization of the impact of GMP/GDP synthesis inhibition on replicative lifespan extension in yeast. *Curr Genet* 1: 3

Longo VD, Shadel GS, Kaeberlein M & Kennedy B (2012) Replicative and chronological aging in *Saccharomyces cerevisiae*. *Cell Metab* 16: 18–31

MacLean M, Harris N & Piper PW (2001) Chronological lifespan of stationary phase yeast cells; a model for investigating the factors that might influence the ageing of postmitotic tissues in higher organisms. *Yeast* 18: 499–509

Maitra N, He C, Blank HM, Tsuchiya M, Schilling B, Kaeberlein M, Aramayo R, Kennedy BK & Polymenis M (2020) Translational control of one-carbon metabolism underpins ribosomal protein phenotypes in cell division and longevity. *Elife* 9: e53127

Maruyama Y, Ito T, Kodama H & Matsuura A (2016) Availability of amino acids extends chronological lifespan by suppressing hyper-acidification of the environment in *Saccharomyces cerevisiae*. *PLoS One* 11: e0151894

Matecic M, Smith DL, Pan X, Maqani N, Bekiranov S, Boeke JD & Smith JS (2010) A microarray-based genetic screen for yeast chronological aging factors. *PLoS Genet* 6: e1000921

McClure JM, Wierman MB, Maqani N & Smith JS (2012) Isonicotinamide enhances Sir2 protein-mediated silencing and longevity in yeast by raising intracellular NAD^+ concentration. *J Biol Chem* 287: 20957–20966

Mitsui A & Suzuki S (1969) Immunosuppressive effect of Mycophenolic acid. *J Antibiot (Tokyo)* 22: 358–363

Mizuguchi G, Shen X, Landry J, Wu WH, Sen S & Wu C (2004) ATP-driven exchange of histone H2AZ variant catalyzed by SWR1 chromatin remodeling complex. *Science* 303: 343–348

Mohammad K, Junio JAB, Tafakori T, Orfanos E & Titorenko VI (2020) Mechanisms that link chronological aging to cellular quiescence in budding yeast. *Int J Mol Sci* 21: 1–14

Mortimer RK & Johnston JR (1959) Life span of individual yeast cells. *Nature* 183: 1751–1752

Moshfegh CM, Collins CW, Gunda V, Vasanthakumar A, Cao JZ, Singh PK, Godley LA & Case AJ (2019) Mitochondrial superoxide disrupts the metabolic and epigenetic landscape of $CD4^+$ and $CD8^+$ T-lymphocytes. *Redox Biol* 27:101141

Niccoli T & Partridge L (2012) Ageing as a risk factor for disease. *Curr Biol* 22: R741–R752

Orlandi I, Pellegrino Coppola D, Strippoli M, Ronzulli R & Vai M (2017) Nicotinamide

supplementation phenocopies SIR2 inactivation by modulating carbon metabolism and

respiration during yeast chronological aging. *Mech Ageing Dev* 161: 277–287

Osório H, Silles E, Maia R, Peleteiro B, Moradas-Ferreira P, Günther Sillero MA & Sillero A

(2005) Influence of chronological aging on the survival and nucleotide content of

Saccharomyces cerevisiae cells grown in different conditions: occurrence of a high

concentration of UDP-N-acetylglucosamine in stationary cells grown in 2% glucose. *FEMS

Yeast Res* 5: 387–398

Park J-I, Grant CM & Dawes IW (2005) The high-affinity cAMP phosphodiesterase of

Saccharomyces cerevisiae is the major determinant of cAMP levels in stationary phase:

involvement of different branches of the Ras–cyclic AMP pathway in stress responses.

Biochem Biophys Res Commun 327: 311–319

Parsons AB, Brost RL, Ding H, Li Z, Zhang C, Sheikh B, Brown GW, Kane PM, Hughes TR &

Boone C (2004) Integration of chemical-genetic and genetic interaction data links bioactive

compounds to cellular target pathways. *Nat Biotechnol* 22: 62–69

Postnikoff SDL & Harkness TAA (2014) Replicative and chronological life-span assays.

Methods Mol Biol 1163: 223–227

Powers RW, Kaeberlein M, Caldwell SD, Kennedy BK & Fields S (2006) Extension of

chronological life span in yeast by decreased TOR pathway signaling. *Genes Dev* 20: 174–

Qiu Y, Fairbanks LD, Rückemann K, Hawrylowicz CM, Richards DF, Kirschbaum B &

Simmonds HA (2000) Mycophenolic acid-induced GTP depletion also affects ATP and

pyrimidine synthesis in mitogen-stimulated primary human T-lymphocytes. *Transplantation* 69: 890-897

Sánchez-Férez F, Ejarque D, Calvet T, Font-Bardia M & Pons J (2019) Isonicotinamide-based compounds: From cocrystal to polymer. *Molecules* 24: 4169

Santisteban MS, Hang M & Smith MM (2011) Histone variant H2A.Z and RNA polymerase II transcription elongation. *Mol Cell Biol* 31: 1848–1860

Santisteban MS, Kalashnikova T & Smith MM (2000) Histone H2A.Z regulates transcription and is partially redundant with nucleosome remodeling complexes. *Cell* 103: 411–422

Santos SM, Laflin S, Broadway A, Burnet C, Hartheimer J, Rodgers J, Smith DL & Hartman IV JL (2020) High-resolution yeast quiescence profiling in human-like media reveals complex influences of auxotrophy and nutrient availability. *GeroScience* 43: 941–964

Sarnoski EA, Liu P & Acar M (2017) A high-throughput screen for yeast replicative lifespan identifies lifespan-extending compounds. *Cell Rep* 21: 2639–2646

Sauve AA, Moir RD, Schramm VL & Willis IM (2005) Chemical activation of Sir2-dependent silencing by relief of nicotinamide inhibition. *Mol Cell* 17: 595–601

Sauve AA & Schramm VL (2003) Sir2 regulation by nicotinamide results from switching between base exchange and deacetylation chemistry. *Biochemistry* 42: 9249–9256

Shukla SK, Gunda V, Abrego J, Haridas D, Mishra A, Souchek J, Chaika N V., Yu F, Sasson AR, Lazenby AJ, *et al* (2015) MUC16-mediated activation of mTOR and c-MYC reprograms pancreatic cancer metabolism. *Oncotarget* 6: 19118–19131

Sims JL, Sikorski GW, Catino DM, Berger SJ & Berger NA (1982)

Poly(adenosinediphosphoribose) Polymerase inhibitors stimulate unscheduled deoxyribonucleic acid synthesis in normal human lymphocytes. *Biochemistry* 21: 1813–1821

Smith Jr DL, Maharrey CH, Carey CR, White RA & Hartman IV JL (2016) Gene-nutrient interaction markedly influences yeast chronological lifespan. *Exp Gerontol* 86: 113–123

Smith Jr DL, McClure JM, Matecic M & Smith JS (2007) Calorie restriction extends the chronological lifespan of *Saccharomyces cerevisiae* independently of the Sirtuins. *Aging Cell* 6: 649–662

Smith ED, Tsuchiya M, Fox LA, Dang N, Hu D, Kerr EO, Johnston ED, Tchao BN, Pak DN, Welton KL, *et al* (2008) Quantitative evidence for conserved longevity pathways between divergent eukaryotic species. *Genome Res* 18: 564–570

Stephan J, Franke J & Ehrenhofer-Murray AE (2013) Chemical genetic screen in fission yeast reveals roles for vacuolar acidification, mitochondrial fission, and cellular GMP levels in lifespan extension. *Aging Cell* 12: 574–583

Sweeney MJ, Hoffman DH & Esterman MA (1972) Metabolism and biochemistry of mycophenolic acid. *Cancer Res* 32: 1803-1809

Toth B (1983) Lack of carcinogenicity of nicotinamide and isonicotinamide following lifelong administration to mice. *Oncology* 40: 72–75

Ueda K, Banasik M, Nakajima S, Yook HY & Kido T (1995) Cell differentiation induced by poly(ADP-ribose) synthetase inhibitors. *Biochimie* 77: 368–373

Váchová L, Cáp M & Palková Z (2012) Yeast colonies: a model for studies of aging,

environmental adaptation, and longevity. *Oxid Med Cell Longev* 2012: 601836

Wagih O, Usaj M, Baryshnikova A, VanderSluis B, Kuzmin E, Costanzo M, Myers CL, Andrews BJ, Boone CM & Parts L (2013) SGAtools: one-stop analysis and visualization of array-based genetic interaction screens. *Nucleic Acids Res* 41: W591–W596

Wei M, Fabrizio P, Hu J, Ge H, Cheng C, Li L & Longo VD (2008) Life span extension by calorie restriction depends on Rim15 and transcription factors downstream of Ras/PKA, Tor, and Sch9. *PLoS Genet* 4: e13

Wierman MB, Maqani N, Strickler E, Li M & Smith JS (2017) Caloric restriction extends yeast chronological life span by optimizing the Snf1 (AMPK) Signaling Pathway. *Mol Cell Biol* 37: e00562-16

Wierman MB, Matecic M, Valsakumar V, Li M, Smith Jr DL, Bekiranov S & Smith JS (2015a) Functional genomic analysis reveals overlapping and distinct features of chronologically long-lived yeast populations. *Aging (Albany NY)* 7: 177–194

Wierman MB & Smith JS (2014) Yeast sirtuins and the regulation of aging. *FEMS Yeast Res* 14: 73–88

Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, Bangham R, Benito R, Boeke JD, Bussey H, *et al* (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 285: 901-906

Wogulis M, Chew ER, Donohoue PD & Wilson DK (2008) Identification of formyl kynurenine formamidase and kynurenine aminotransferase from *Saccharomyces cerevisiae* using crystallographic, bioinformatic and biochemical evidence. *Biochemistry* 47: 1608–1621

Wu WH, Wu CH, Ladurner A, Mizuguchi G, Wei D, Xiao H, Luk E, Ranjant A & Wu C (2009) N terminus of Swr1 binds to histone H2AZ and provides a platform for subunit assembly in the chromatin remodeling complex. *J Biol Chem* 284: 6200–6207

Xie MW, Jin F, Hwang H, Hwang S, Anand V, Duncan MC & Huang J (2005) Insights into TOR function and rapamycin response: Chemical genomic profiling by using a high-density cell array method. *Proc Natl Acad Sci U S A* 102: 7215–7220

Yadav B, Wennerberg K, Aittokallio T & Tang J (2015) Searching for drug synergy in complex dose-response landscapes using an interaction potency model. *Comput Struct Biotechnol J* 13: 504–513

Yin N, Ma W, Pei J, Ouyang Q, Tang C & Lai L (2014) Synergistic and antagonistic drug combinations depend on network topology. *PLoS One* 9: e93960

FIGURE LEGENDS

Figure 1. INAM extends chronological lifespan. **A)** Chemical structures of nicotinamide and isonicotinamide. **B)** CLS assays with increasing dosage of INAM added to BY4741 at the time of inoculation. **C)** CLS assay comparing the effect of 25 mM INAM on the auxotrophic strain BY4741 and prototrophic strain FY4, added at inoculation. **D)** CLS of BY4741 when 25 mM INAM is added 96 hr after inoculation. **E)** WT (BY4741) or *sir2Δ hst1Δ* double mutant CLS with or without 25 mM INAM supplemented at time of inoculation. Viability for each CLS assay is calculated as the percentage of colony forming units at each time point compared to the stationary phase day 3 time point, which was collected 72 hours after inoculation. Error bars indicate standard deviations (n=3 biological replicates). Statistics are provided in Table S3.

Figure 2. Chemical genetic screen for INAM sensitive gene deletion mutants in the yeast knockout (YKO) strain collection. **A)** Spot test growth assay of BY4741 on SC agar plates containing 0, 25, 50, 75, or 125 mM INAM. **B)** The haploid yeast knockout collection was pinned onto control SC plates or SC plates with increasing concentrations of INAM using a manual pinning tool. **C)** Zoomed in portion of plate 7 showing INAM sensitivity of the *tho2Δ* strain (yellow arrow). **D)** Scatter plot of scores from each deletion strain obtained in experiment 1 (x-axis) and experiment 2 (y-axis). Strains with scores lower than -0.3 in both replicates were considered sensitive to a given INAM concentration (representative graph for 75 mM INAM). **E)** Venn diagram showing the overlap of strains scored as sensitive between different INAM concentrations.

Figure 3. Strains defective for the H2A.Z deposition by SWR1 are sensitive to INAM. **A)** SWR1 complex schematic diagram illustrating subunits and their sensitivity to INAM. **B)** Spot test growth assay showing confirmation of INAM sensitivity of *htz1Δ* and *swr1Δ* strains from YKO collection. The WT strain is BY4741. **C)** CLS assay of BY4741 and *swr1Δ* mutant strains supplemented with 25 mM INAM. **D)** CLS assay of BY4741 and *htz1Δ* strains supplemented with 25 mM INAM. Controls were supplemented with equal volume of sterile water instead of INAM stock solution.

Figure 4. Serine biosynthesis mutants are sensitive to INAM. **A)** Schematic of serine, aspartate, threonine, and cysteine synthesis pathways and their conversion at the one carbon metabolism pathway and purine metabolism. Deleted genes conferring INAM sensitivity are in bold. **B)** CLS assay showing moderate lifespan extension of *ser2Δ* mutant, and full extension by 25 mM INAM. **C)** Spot test growth assay for WT (BY4741) and *ser2Δ* strains on SC plates supplemented with 25, 50, or 75 mM INAM. **D)** Quantitative dose-response curve illustrating the

relative viability of BY4741 (WT) and the *ser2Δ* mutant at a range of INAM concentrations. Dots represent values obtained from three biological replicates with sigmoid curves fitted to the data using a dose response function and the Levenberg Marquardt iteration algorithm. Shaded areas around the curve are 95% confidence bands. **E)** Bar graph depicting IC_{50} of BY4741 (WT) and the *ser2Δ* mutant calculated from the equations of fitted curves in panel D. Error bars represent 95% confidence intervals.

Figure 5. Mycophenolic acid and INAM operate through related but distinct mechanisms. **A)** Venn diagram depicting a partial overlap between INAM- and MPA-sensitive deletion strains identified in a chemogenomic screen and confirmed by spot assays. **B)** Synergy map depicting interaction between INAM and MPA in reducing cell growth. Graph is representative of three independent experiments. **C)** CLS assay with BY4741 showing CLS extension by MPA supplementation (1 or 10 µg/ml), and suppression of the effect by 0.1 mM guanine. **D)** CLS assay with BY4741 showing INAM extension of CLS is that is not suppressed by guanine.

Figure 6. Alterations of intracellular nucleotide metabolite levels by 25 mM INAM. **A)** Summary of quantitative mass spectrometry analysis for a panel of purine and pyrimidine nucleotides and precursors. Shaded blocks (green-up, red-down) indicate metabolites with significant differences ($p<0.05$) in INAM treated cells compared to control. Darker shading indicates more significant p-values. **B)** Examples of significantly reduced purine nucleosides or nucleotides reduced by INAM at each time point. *$p<0.05$, **$p<0.01$. **C)** CLS assay showing partial lifespan extension when BY4741 (uracil auxotroph) is supplemented with 4x (0.8 mM) the normal concentration of uracil (0.2 mM) in SC medium. INAM (25 mM) fully extends CLS in combination with 4x uracil.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Confirmation of candidate mutants from the INAM sensitivity screen using spot test growth assays. The normalized growth scores from SGAtools are indicated for the two independent screens (performed sequentially). Scores for 25, 50, and 75 mM plates were from day 2 images, while scores for 125 mM plates were from day 3 images. Scores below the -0.3 cutoff are shaded in green.

Figure S2. Confirmation of additional candidate mutants from INAM sensitivity screen. **A)** Mutants that confirmed as INAM sensitive. Each experiment has its own BY4741 WT control. **B)** Mutants that did not confirm as INAM sensitive. SGAtools scores for the two screens are indicated to the right, as in Figure S1.

Figure S3. Strains not identified by the screen but confirmed as sensitive in the spot test growth assays. These additional strains were chosen based on identity of confirmed strains from the screen and the literature.

Figure S4. Loss of *BNA7* confers INAM resistance and extends CLS. **A)** Spot test growth assay of candidate INAM-resistant mutants chosen from the screen on 125 mM INAM plates. The only mutant confirmed as resistant was *bna7Δ*. **B)** Schematic diagram of *de novo* NAD$^+$ synthesis and salvage pathways in *S. cerevisiae*. The Bna7 reaction releases formate from L-formylkynurenine as products. **C)** CLS assay showing late stage lifespan extension of the *bna7Δ* mutant, and further extension by 25 mM INAM.

Figure S5. INAM sensitivity of serine and threonine synthesis mutants (*ser2Δ*, *thr1Δ* and *hom3Δ*) is rescued by the addition of respective amino acids. Error bars represent 95% confidence intervals derived from the dose-response curves.

Figure S6. Summary of purine nucleotide and metabolite levels and related gene expression changes induced by INAM in cells collected at log phase, after 24 hr, or 96 hr of liquid culture. Diagram was produced using VANTED.

Figure S7. Pyrimidine metabolite levels and gene expression changes at day 8. **A)** Summary of pyrimidine nucleotide and metabolite levels and related gene expression from log phase to 96 hr cultures. **B)** INAM-induced expression changes of select genes involved in nucleotide metabolism. Expression of each gene at day 8 was normalized to *TAF10*. Three biological replicates were tested.

A

B

C

D

A

B

ZIP synergy score: 1.359

Most synergistic area
Score: 9.86 (ZIP)

C

D

Figure 6