Title: The retrotransposon R2 maintains Drosophila ribosomal DNA repeats

One Sentence Summary: The retrotransposon R2 initiates restoration of ribosomal DNA copies

to trans-generationally maintain essential locus.

Authors: Jonathan O. Nelson[1], Alyssa Slicko[1,3], and Yukiko M. Yamashita[1,2,3*]

Affiliations: [1] Whitehead Institute for Biomedical Research,

[2]Department of Biology, MIT,

[3]Howard Hughes Medical Institute,

455 Main Street, Cambridge, MA 02142

*Corresponding Author

Abstract: Ribosomal DNA (rDNA) loci contain the hundreds of tandemly repeated copies of ribosomal RNA genes needed to support cellular viability. This repetitiveness makes it highly susceptible to copy loss, threatening multi-generational maintenance of rDNA. How this threat is counteracted to avoid extinction of the lineage has remained unclear. Here, we show that the rDNA-specific retrotransposon R2 is essential for rDNA copy number (CN) maintenance in the Drosophila male germline, despite the perceived disruptive nature of transposable elements. Depletion of R2 led to defective rDNA CN maintenance, causing a decline in fecundity over generations and eventual extinction. This study reveals that retrotransposons can provide a benefit to their hosts, contrary to their reputation as genomic parasitic, which may contribute to their widespread success throughout taxa.

Main Text:

Ribosomal RNAs (rRNAs) account for 80-90% of all transcripts in eukaryotic cells(*1*). To meet this demand, the ribosomal DNA (rDNA) gene that codes for rRNA is tandemly repeated hundreds of times, comprising rDNA loci on eukaryotic chromosomes. Ironically, this repetitive structure is susceptible to intra-chromatid recombination that causes rDNA copy number (CN) loss **(Fig. 1A)**, which is a major cause of replicative senescence in budding yeast(*2*). Evidence of similar rDNA CN instability has been noted in some tissues from aged dogs and humans(*3, 4*). Critically, age-associated rDNA CN loss also occurs in the Drosophila male germline and is inherited by the next generation(*5*). The essential yet unstable nature of rDNA raises the question as to how the degeneration of rDNA loci over successive generations is prevented to avoid the extinction of the lineage. Intensive studies have revealed that sister chromatid recombination mediates rDNA CN expansion in yeast, thereby maintaining rDNA repeat abundance over generations(*2, 6*). rDNA CN is variable between individuals of most species but maintained within a consistent range throughout the population(*7*), implying that trans-generational dynamic rDNA CN changes (loss and restoration) are a common feature of rDNA maintenance. However, the mechanism of trans-

generational rDNA CN maintenance in multicellular organisms has remained a mystery. Over 50 years ago, the phenomenon of 'rDNA magnification' in Drosophila was first described as the process wherein aberrant rDNA loci bearing minimal rDNA repeats recover to a normal rDNA CN(*8*). rDNA magnification was shown to require genes involved in homologous recombination-mediated repair, similar to yeast, which can create duplications of tandemly repeated elements(*9*). Despite the robust rDNA CN expansion activity, it remained unclear if the mechanisms of rDNA magnification served a physiological function in natural populations. We recently demonstrated that offspring who inherited reduced rDNA CN from old fathers could also recover normal rDNA CN in their germline(*5*). This recovery depends on the same set of the genes as rDNA magnification, leading us to propose that rDNA magnification is a manifestation of the physiological mechanisms to maintain rDNA CN across generations(*7*). However, the underlying factors responsible for rDNA magnification and their normal physiological functions remain poorly understood.

Metazoan rDNA genes are frequently inserted by rDNA-specific transposable elements (TEs), such as the retrotransposon *R2* in Drosophila. *R2* is found throughout arthropods(*10*) and *R2*-like elements are widely present across taxa, including Cnidaria(*11*), Planaria(*12*), nematodes(*13*), fish(*14, 15*), birds, and reptiles(*15*). These TEs use their sequence-specific nuclease to mobilize specifically within rDNA loci, inserting in rDNA genes and disrupting their function(*16*) **(Fig. S1A)**. Surprisingly, we found that RNAi-mediated knockdown of *R2* in the Drosophila male germline (nos-gal4 driving *R2* RNAi expression) **(Fig. S1A-B)** resulted in premature loss of germline stem cells (GSCs) during aging **(Fig. 1B-F)**. GSCs continuously produce differentiating germ cells to sustain sperm production throughout adulthood and thus are the source of the genome passed to the next generation(*17*). Whereas newly eclosed *R2* RNAi males contained similar numbers of GSCs to controls, GSC number more rapidly declined during aging in *R2* knockdown males compared to controls **(Fig. 1B-F)**. Given that *R2* is specifically

inserted at rDNA, we also examined the effect of *R2* knockdown on rDNA stability. Using highly

quantitative droplet digital PCR (ddPCR), we found that RNAi-mediated knockdown of *R2*

enhanced rDNA CN loss in the testis during aging compared to controls **(Fig. 1G)**. rDNA CN

insufficiency is likely the primary cause of GSC loss in *R2* RNAi animals, because increasing total

rDNA CN via introduction of a mini-chromosome harboring an rDNA locus(*18*) suppressed the

premature GSC loss caused by *R2* knockdown **(Fig. 1H)**. These results revealed that *R2*

contributes to sustaining GSC population during aging by supporting rDNA CN maintenance,

uncovering an unanticipated benefit of the R2 retrotransposon to the host, despite the widely-held

view of TEs being genetic parasites.

Why is *R2* necessary for rDNA copy number maintenance? We found that *R2* is required

for rDNA magnification, a phenomenon of rDNA CN expansion classically assayed by the

restoration of normal cuticular development in animals inheriting rDNA loci that had insufficient

rDNA CN in the previous generation **(Fig. 2A)**. The *Drosophila melanogaster* genome has two

rDNA loci, one on each X and Y chromosome(*19*). rDNA magnification almost exclusively occurs

to X chromosome rDNA loci harboring the minimal viable amount of rDNA (bb^{Z9}, **Fig. S2A)** when

combined with a Y chromosome completely lacking rDNA (bb^{Z9}/Ybb^0, 'magnifying males'

hereafter)(*8*) **(Fig. S2B)**. Importantly, rDNA magnification never occurs in males with a normal Y

chromosome containing intact rDNA (bb^{Z9}/Y$^+$, 'non-magnifying males' hereafter)(*20*), indicating

the presence of mechanisms to monitor rDNA CN to activate expansion. We found that RNAi-

mediated knockdown of *R2* eliminated rDNA magnification observed in magnifying males from

13.73% to 0% (n = 414, p = 5.6x10^{-7}) **(Fig. 2B)**. Moreover, quantification of rDNA CN by ddPCR

revealed that 87.5% of bb^{Z9} chromosomes increased rDNA CN in magnifying males compared to

the non-magnifying males (n = 96, p = 1.8x10^{-4}), with an average increase of 18.29 rDNA copies

across all samples (n = 96, p = 3.1x10^{-12}) **(Fig. 2C)**. This widespread rDNA CN increase indicates

that rDNA magnification broadly increases rDNA CN throughout the germline, despite only

13.73% of bb^{Z9} chromosomes having enough CN increase to support normal cuticle development.

We found this rDNA CN increase is also eliminated upon *R2* knockdown in magnifying males (n

= 96, p = 4.7×10^{-10}) **(Fig. 2C)**. These results reveal that *R2* is required for rDNA CN expansion

during rDNA magnification. Interestingly, we found that *R2* knockdown only disrupted rDNA

magnification when the *R2* RNAi construct was expressed by the *nanos*-gal4 driver in early germ

cells (including GSCs)(*21*), but not when expressed in later germ cells (excluding GSCs) by the

bam-gal4 driver(*22*) **(Fig. 2B)**. These results indicate that *R2* primarily functions in the earliest

stages of germ cells (including GSCs) to support rDNA magnification.

We further found that *R2* is sufficient for rDNA CN expansion. Ectopic expression of *R2* in

the germline induced rDNA magnification of the bb^{Z9} locus in non-magnifying males (bb^{Z9}/Y$^+$) **(Fig.**

2D-E). While assessment of dorsal cuticle phenotype never detects rDNA magnification from

control males, we found 3.3% of bb^{Z9} rDNA loci from non-magnifying males expressing a

transgenic copy of *R2* were phenotypically magnified **(Fig. 2D,** n = 877, p = 3.2×10^{-5}).

Quantification of rDNA CN by ddPCR revealed that ectopic overexpression of *R2* in non-

magnifying males (bb^{Z9}/Y$^+$) also increases the average rDNA CN at bb^{Z9} rDNA loci among all

offspring compared to controls **(Fig. 2E**, n = 94, p = 0.0256), revealing *R2* is sufficient to induce

rDNA CN expansion. Critically, expression of a nuclease dead R2 transgene (NucDeadR2) in

non-magnifying males **(Fig. S3A)**(*23*) failed to induce phenotypic detection of rDNA magnification

(Fig. 2D), suggesting that the nuclease activity of R2 is essential for its ability to induce rDNA CN

expansion.

How does R2's nuclease activity contribute to rDNA magnification? In yeast, rDNA CN

expansion is initiated by double-stranded breaks (DSBs) at the rDNA intergenic sequence, which

induces sister chromatid recombination that results in rDNA gene duplication(*24*). While the

mechanism of rDNA magnification remains unclear, all proposed models of Drosophila rDNA CN

expansion require an initiating DSB at the rDNA locus, the most prominent model being sister

chromatid recombination resulting in unequal sister chromatid recombination(*25*) **(Fig. 3A; Fig.**

S4A-B for additional models**)**. Indeed, artificial introduction of DSBs at rDNA loci by I-CreI

endonuclease expression has been reported to induce rDNA magnification(*26*), but the

endogenous factor that induces rDNA magnification remained unclear. R2 is capable of creating

DSBs through sequential nicking of opposite DNA strands during retrotransposition(*27*). It has

been speculated that DSBs created during R2 retrotranspostion may be an initiating event of

rDNA magnification(*28*), although this possibility has yet to be empirically tested. We first

confirmed that ectopic overexpression of R2, but not NucDeadR2, indeed induces chromosomal

breaks at rDNA loci identified by chromosome spreads **(Fig. S3B-D)**. R2 overexpression (but not

NucDeadR2) in the germline also increased the frequency of GSCs with DSBs, identified by

γH2Av expression **(Fig. S3E)**. Next, we found rDNA magnification is associated with an elevation

in DSBs in GSCs: the frequency of γH2Av-positive GSCs is increased in magnifying males

(bb^{Z9}/Ybb^0) compared to non-magnifying males (bb^{Z9}/Y^+) (**Fig. 3B-C, E**; n = 519, p = 8.8×10^{-4}).

Strikingly, we observed that knockdown of *R2* in magnifying males reduced the frequency of

γH2Av-positive GSCs to levels comparable to non-magnifying males **(Fig. 3D-E;** n = 537, p =

7.1×10^{-4}**)**, indicating that R2 is responsible for the DSBs formed in GSCs during rDNA

magnification. Taken together, these results suggest that rDNA-specific endonuclease activity of

R2, which functions during R2 retrotransposition, creates DSBs at the rDNA loci that may in turn

induce rDNA CN expansion.

Given the threat *R2* mobilization poses to the host genome, both by disruption of rRNA

function and causing excessive DSB formation(*27*), how is the potential benefit of *R2* to rDNA CN

maintenance balanced with the detriment of *R2* retrotransposition? We found *R2* expression in

the germline is specifically de-repressed under conditions of reduced rDNA CN, potentially

explaining how the conflicting consequences of *R2* expression are resolved. Using RNA

fluorescence *in situ* hybridization (RNA FISH) to examine *R2* expression at a single cell resolution,

we found that the frequency of GSCs expressing *R2* was significantly increased in magnifying

males (bb^{Z9}/Ybb^0), whereas non-magnifying males (bb^{Z9}/Y$^+$) rarely expressed *R2* **(Fig. 3F-H**; n =

231, p = 1.7x10^{-10}). Moreover, we found that GSCs from aged animals and the sons of old fathers,

which inherit reduced rDNA CN(*5*)**,** also exhibited a higher frequency of *R2* expression compared

to GSCs from young, wild type flies **(Fig. S5A-B, D**; n = 1,247, p = 8.3x10^{-4} for old animals; n =

1,107, p = 1.5x10^{-4} for offspring). Importantly, the frequency of *R2* expression among GSCs in the

sons of old fathers returned to the basal level after 20 days of age, when rDNA CN was shown to

have recovered(*5*) **(Fig. S5C-D**; n = 617, p = 0.036**)**. These results indicate that *R2* expression is

dynamically regulated in response to changing rDNA CN. Taken together, we propose that *R2*

expression is finely tuned to function when most beneficial to the host while minimizing

unnecessary exposure to the harmful effects of transposition.

Based on the finding that *R2* plays a critical role in maintaining germline rDNA CN, we

postulated that *R2* is essential to prevent continuous multi-generational rDNA loss capable of

causing the extinction of the lineage. In *C. elegans*, the loss of genome integrity is known to cause

gradual loss of fertility, a phenotype known as <u>mor</u>tal <u>g</u>ermline (morg)(*29*). To test whether *R2*-

mediated rDNA maintenance is required to maintain fertility through generations, we established

multiple independent lines expressing *R2* RNAi in their germline and tracked their fecundity at

each generation through the ability of each line to produce sufficient offspring to establish a new

generation **(Fig. S6)**. We found many *R2* RNAi lines failed to produce sufficient progeny, with

over half failing by the fourth generation, while nearly all control lines survived throughout the

duration of the experiments **(Fig. 4A)**, indicating that *R2* is essential for continuity of the germline

lineage. Males of extinguishing *R2* RNAi lineages had ~20% reduction in rDNA CN compared to

control lines (n = 22, p = 0.031) **(Fig. 4B)**. These results suggest that *R2*-mediated maintenance

of rDNA contributes to germline immortality.

Together, our findings reveal an unanticipated 'function' of retrotransposons to benefit the

host genome through a role in rDNA CN maintenance. Repetitive DNA sequences such as rDNA

are among the most vulnerable elements of the eukaryotic genome(*30, 31*). We propose that

DSBs generated by *R2* in GSCs with diminished rDNA loci stimulate sister chromatid

recombination that results in rDNA CN expansion **(Fig. 4C)**. Our proposed 'function' for *R2* in

rDNA maintenance may represent a novel mutualistic host-TE relationship, which are rarely

described in eukaryotes(*32*). *R2*'s de-repression when it can be beneficial (i.e. decreased rDNA

CN) may be the key to this mutualistic host-TE relationship. It awaits future investigation to

understand how the mechanisms that repress *R2*(*33–35*) may tune this activity for host's benefit.

The widespread presence of *R2* and other rDNA-specific TEs in both vertebrates and

invertebrates(*15*) suggests that similar host-TE mutualism may support rDNA CN maintenance

throughout Metazoa. Interestingly, many of these other *R2*-like elements have little sequence

similarity to R2, instead appearing to be derived from other non-specific TEs(*15*), suggesting this

host-TE mutualism may have evolved multiple times over the course of evolution. In summary,

our study provides an example of beneficial TEs in the maintenance of eukaryotic genomes, and

we propose that more beneficial TEs are yet to be discovered.

References:

1. A. F. Palazzo, E. S. Lee, Non-coding RNA: what is functional and what is junk? *Frontiers Genetics*. **6**,
2 (2015).

2. A. R. D. Ganley, T. Kobayashi, Ribosomal DNA and cellular senescence: new evidence supporting the
connection between rDNA and aging. *Fems Yeast Res*. **14**, 49–59 (2014).

3. B. L. Strehler, M.-P. Chang, L. K. Johnson, Loss of hybridizable ribosomal DNA from human post-
mitotic tissues during aging: I. Age-dependent loss in human myocardium. *Mech Ageing Dev*. **11**, 371–
378 (1979).

4. R. Johnson, B. L. Strehler, Loss of Genes coding for Ribosomal RNA in Ageing Brain Cells. *Nature*.
240, 412–414 (1972).

5. K. L. Lu, J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita, Transgenerational
dynamics of rDNA copy number in Drosophila male germline stem cells. *Elife*. **7**, e32421 (2018).

6. T. Kobayashi, Ribosomal RNA gene repeats, their stability and cellular senescence. *Proc Jpn Acad Ser*
B. **90**, 119–129 (2014).

7. J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita, Mechanisms of rDNA Copy Number
Maintenance. *Trends Genet*. **35**, 734–742 (2019).

8. F. M. Ritossa, Unstable redundancy of genes for ribosomal RNA. *Proc National Acad Sci*. **60**, 509–516
(1968).

9. A. B. Reams, J. R. Roth, Mechanisms of Gene Duplication and Amplification. *Csh Perspect Biol*. **7**,
a016592 (2015).

10. W. D. Burke, H. S. Malik, J. P. Jones, T. H. Eickbush, The domain structure and retrotransposition
mechanism of R2 elements are conserved throughout arthropods. *Mol Biol Evol*. **16**, 502–511 (1999).

11. K. K. Kojima, K. Kuma, H. Toh, H. Fujiwara, Identification of rDNA-Specific Non-LTR
Retrotransposons in Cnidaria. *Mol Biol Evol*. **23**, 1984–1993 (2006).

12. W. D. Burke, D. Singh, T. H. Eickbush, R5 Retrotransposons Insert into a Family of Infrequently
Transcribed 28S rRNA Genes of Planaria. *Mol Biol Evol*. **20**, 1260–1270 (2003).

13. W. D. Burke, F. Müller, T. H. Eickbush, R4, a non-LTR retrotransponson specific to the large subunit
rRNA genes of nematodes. *Nucleic Acids Res*. **23**, 4628–4634 (1995).

14. K. K. Kojima, H. Fujiwara, Cross-Genome Screening of Novel Sequence-Specific Non-LTR
Retrotransposons: Various Multicopy RNA Genes and Microsatellites Are Selected as Targets. *Mol Biol*
Evol. **21**, 207–217 (2004).

15. K. K. Kojima, H. Fujiwara, Long-Term Inheritance of the 28S rDNA-Specific Retrotransposon R2.
Mol Biol Evol. **22**, 2157–2165 (2005).

16. J. Yang, H. S. Malik, T. H. Eickbush, Identification of the endonuclease domain encoded by R2 and
other site-specific, non-long terminal repeat retrotransposable elements. *Proc National Acad Sci*. **96**,
7847–7852 (1999).

17. M. T. Fuller, A. C. Spradling, Male and Female Drosophila Germline Stem Cells: Two Versions of
Immortality. *Science*. **316**, 402–404 (2007).

18. D. L. Lindsley, L. Sandler, The Meiotic Behavior of Grossly Deleted X Chromosomes in Drosophila
Melanogaster. *Genetics*. **43**, 547–63 (1958).

19. F. M. Ritossa, K. C. Atwood, S. Spiegelman, A molecular explanation of the bobbed mutants of
Drosophila as partial deficiencies of "ribosomal" DNA. *Genetics*. **54**, 819–34 (1966).

20. K. D. Tartof, Regulation of ribosomal RNA gene multiplicity in Drosophila melanogaster. *Genetics*.
73, 57–71 (1973).

21. M. V. Doren, A. L. Williamson, R. Lehmann, Regulation of zygotic gene expression in Drosophila
primordial germ cells. *Curr Biol*. **8**, 243–246 (1998).

22. D. Chen, D. M. McKearin, A discrete transcriptional silencer in the bam gene determines asymmetric
division of the Drosophila germline stem cell. *Development*. **130**, 1159–1170 (2003).

23. A. Govindaraju, J. D. Cortez, B. Reveal, S. M. Christensen, Endonuclease domain of non-LTR
retrotransposons: loss-of-function mutants and modeling of the R2Bm endonuclease. *Nucleic Acids Res*.
44, 3276–3287 (2016).

24. S. Gangloff, H. Zou, R. Rothstein, Gene conversion plays the major role in controlling the stability of
large tandem repeats in yeast. *Embo J*. **15**, 1715–1725 (1996).

25. K. D. Tartof, Unequal Mitotic Sister Chromatid Exchange as the Mechanism of Ribosomal RNA
Gene Magnification. *Proc National Acad Sci*. **71**, 1272–1276 (1974).

26. S. Paredes, K. A. Maggert, Expression of I-CreI Endonuclease Generates Deletions Within the rDNA
of Drosophila. *Genetics*. **181**, 1661–1671 (2009).

27. T. H. Eickbush, D. G. Eickbush, Integration, regulation, and long-term stability of R2
retrotransposons. *Microbiol Spectr*. **3**, 1127–1146 (2015).

28. R. S. Hawley, C. H. Marcus, Recombinational Controls of rDNA Redundancy in Drosophila. *Annu
Rev Genet*. **23**, 87–120 (1989).

29. C. Smelick, S. Ahmed, Achieving immortality in the C. elegans germline. *Ageing Res Rev*. **4**, 67–82
(2005).

30. R. E. Brown, C. H. Freudenreich, Structure-forming repeats and their impact on genome stability.
Curr Opin Genet Dev. **67**, 41–51 (2021).

31. F. Pâques, W.-Y. Leung, J. E. Haber, Expansions and Contractions in a Tandem Repeat Induced by
Double-Strand Break Repair. *Mol Cell Biol*. **18**, 2045–2054 (1998).

32. R. L. Cosby, N.-C. Chang, C. Feschotte, Host–transposon interactions: conflict, cooperation, and
cooption. *Gene Dev*. **33**, 1098–1116 (2019).

33. E. Quénerch'du, A. Anand, T. Kai, The piRNA pathway is developmentally regulated during
spermatogenesis in Drosophila. *Rna*. **22**, 1044–1054 (2016).

34. F. He, A. James, H. Raje, H. Ghaffari, P. DiMario, Deletion of Drosophila Nopp140 induces
subcellular ribosomopathies. *Chromosoma*. **124**, 191–208 (2015).

35. P. A. Guerrero, K. A. Maggert, The CCCTC-Binding Factor (CTCF) of Drosophila Contributes to the
Regulation of the Ribosomal DNA and Nucleolar Stability. *Plos One*. **6**, e16401 (2011).

262
263

Acknowledgements

We thank the Bloomington Drosophila Stock Center, Kyoto Drosophila Stock Center and Developmental Studies Hybridoma Bank for reagents. We thank the Yamashita lab members and Dr. Andy Clark for discussion and comments on the manuscript. We thank ATCC for design of the 28S and RpL ddPCR assays. **Funding:** This research was supported by the Howard Hughes Medical Institute. Jonathan Nelson is supported by a Postdoctoral Fellowship (133949-PF-19-133-01-DMC) from the American Cancer Society. **Authors contributions:** Jonathan O. Nelson: Conceptualization, Methodology, Verification, Formal analysis, Investigation, Resources, Data curation, Writing – Original draft, Writing – Review & editing, Visualization, Funding acquisition. Alyssa Slicko: Validation, Investigation, Resources, Data curation, Writing – Review & editing. Yukiko M. Yamashita: Conceptualization, Methodology, Resources, Writing – Original draft, Writing – Review & editing, Visualization, Supervision, Project administration, Funding. **Competing interests:** The authors declare no competing interests. **Data materials availability:** All data is available in the manuscript or the supplementary materials.

List of Supplementary Materials

Supplementary Materials

Materials and Methods

Fig S1 – S6

Table S1 – S2

References 36 - 38

Figure 1: R2 is required for GSC maintenance via rDNA CN maintenance during

Drosophila male germline aging. (A) Tandem rDNA copy repeats are lost through

circularization and extrusion caused by intra-chromatid recombination between rDNA copies.

(B-E) GSCs in 0- and 40-day old control (*nos-gal4* driver alone) (B-C) and R2 RNAi (*nos-gal4*

driving *UAS-R2 RNAi*) testes (D-E). Yellow dotted circle = GSC. The major component of the

GSC signaling niche, the hub, is indicated by *. Green = Vasa, White = Fascillin III, Blue =

DAPI. Scale bar = 7.5 µM **(F)** Average number of GSCs per testis in control and R2 RNAi

animals during aging. * indicates $p < 10^{-5}$ determined by Student's t-test. Error bars = 95%

confidence interval (CI). **(G)** rDNA CN determined by ddPCR in testes of 0- and 40-day old

control and R2 RNAi animals. P-value determined by Student's t-test. Error bars = 95% CI. **(H)**

Average number of GSCs per testis during aging in Control and R2 RNAi animals with mini-X

chromosome. Error bars = 95% CI.

Figure 2: R2 is necessary and sufficient for rDNA magnification. (A) Diagram of rDNA

magnification. Animals with insufficient rDNA CN due to having the bb^{Z9} locus as their only

source of rDNA have distinct developmental defects resulting in gaps in the dorsal cuticle (red

arrow). rDNA CN expansion at the bb^{Z9} locus due to rDNA magnification is detected by

reversion of this phenotype to normal cuticles. **(B)** Percent magnified offspring determined by

cuticular phenotype in offspring from magnifying males of the indicated genotypes. P-value

determined by chi-squared test. Error bars = 95% CI. **(C)** bb^{Z9} locus rDNA CN determined by

ddPCR in daughters from males of indicated genotypes. P-value determined by Student's t-test.

Error bars = 95% CI. Non-magnifiying condition is the same data as panel (C). **(D)** Percent

magnified offspring from non-magnifying males of the indicated genotypes. P-value determined

by chi-squared test. Error bars = 95% CI. **(E)** bb^{Z9} locus rDNA CN determined by ddPCR in

daughters from non-magnifying males of the indicated genotype. P-value determined by

Student's t-test. Error bars = 95% CI.

Fig 3: Derepressed R2 creates DSBs in GSCs during rDNA magnification. (A) Diagram of

rDNA CN expansion by unequal sister chromatid exchange during DSB repair at rDNA loci.

Recombination between misaligned rDNA copies during DSB repair result in crossovers that

create unequal sister chromatid exchange that increases rDNA CN on one chromatid. **(B-D)**

Detection of DSBs in the early adult male germline by anti-γH2Av staining. R2 RNAi expressed

under the *nos-gal4* driver. Non-RNAi conditions contain the *nos-gal4* driver alone. GSCs

indicated by yellow dotted circle. Blue = DAPI, Green = vasa, Magenta = γH2Av, white = FasIII.

The hub is indicated by *. Scale bar = 10 µM. **(E)** Percentage of γH2Av positive GSCs in the

indicated conditions. P-value determined by chi-squared test. Error bars = 95% CI. **(F-G)** R2

expression in GSCs (yellow dotted circle). Blue = DAPI, Green = R2 mRNA. Isolated R2

channel in F'-G'. The hub is indicated by *. R2 positive cells GSCs are marked by yellow

arrowhead. **(H)** Percentage of R2 positive GSCs. P-value determined by chi-squared test. Error

bars = 95% CI.

Fig 4: R2 is required to maintain rDNA CN and fertility over successive generations. (A)
Kaplan-Meier curve of lineage survival in R2 RNAi and control lineages. Each lineage
constitutes an individual data point. P-value determined by log rank test. Shade represents 95%
CI. **(B)** rDNA CN determined by ddPCR in males of control (*nos-gal4* driver alone) animals at
the 6[th] generation or R2 RNAi (RNAi expressed under the *nos-gal4* driver) lineages at their
terminating generation. P-value determined by Student's t-test. Error = 95% CI. **(C)** Model of the
role of R2 in germline rDNA CN maintenance. R2 derepression in cells with reduced rDNA CN
leads to DSBs formed by rDNA-specific R2 endonuclease that can induce rDNA CN expansion
through sister chromatid recombination during their repair.

Supplementary Materials

Materials and Methods

Immunofluorescence

Immunofluorescence staining of testes was performed as previously described(*36*). Briefly, testes

were dissected in PBS, fixed in 4% formaldehyde in PBS for 30 minutes, then briefly washed two

times in PBS containing 0.1% Triton-X (PBS-T), followed by washing in PBS-T for 30 minutes.

After washes, samples were incubated at 4°C overnight with primary antibody in 3% bovine serum

albumin (BSA) in PBS-T. Samples were washed three consecutive times for 20 minutes in PBS-

T, then incubated at 4°C overnight with secondary antibody in 3% BSA in PBS-T, washed three

times again in PBS-T for 20 minutes, and mounted in VECTASHIELD with DAPI (Vector Labs).

The following primary antibodies were used: rat anti-vasa (1:20; DSHB; developed by A.

Spradling), mouse anti-Fascillin III (1:200; DSHB; developed by C. Goodman), and rabbit anti-γ-

H2AvD pS137 (1:200; Rockland). Images were taken with a Leica Stellaris 8 confocal microscope

with 63x oil-immersion objectives and processed using Fiji (ImageJ) software.

RNA FISH

RNA FISH samples were prepared as previously described(*5*). In short, dissected testes were

fixed in 4% formaldehyde in PBS for 30 minutes, briefly washed in PBS, and permeabilized in

70% ethanol overnight at 4º. Samples were then briefly rinsed in 2xSSC with 10% formamide

prior to hybridization with 50nM Stellaris probes overnight at 37º. Samples were washed twice in

2xSSC with 10% formamide for 30 minutes and mounted in VECTASHIELD with DAPI (Vector

Labs). Samples were imaged using a Leica Stellaris 8 confocal microscope with 63x oil-immersion

objectives and processed using Fiji (ImageJ) software.

DNA isolation

Testis DNA was isolated from 50 pooled dissected testes frozen in liquid N2. DNA isolation was

performed according to previously described methods for isolation from Drosophila tissues(*37*).

DNA was isolated from individual Drosophila animals using a modified protocol of the DNeasy

Blood and Tissue DNA extraction kit (Qiagen). In short, individual animals were homogenized in

200 µL Buffer ATL containing proteinase K using a pipette tip in Eppendorf tubes, vortexed for 15

seconds, and incubated for 1.5 hours at 56⁰. Samples were then prepared following the

manufacturer's protocol after incubation. All DNA samples were quantified and checked for purity

by Nanodrop One spectrophotometer (ThermoFisher).

rDNA copy number measurement by Droplet Digital PCR (ddPCR)

30 ng of DNA sample were used per 20 µL ddPCR reaction for control gene reactions (RpL and

Upf1), and 0.3 ng of DNA per 20 µL ddPCR reaction for 28S rDNA reactions. Primers and probes

for reactions are listed at **Table S1**. ddPCR reactions were carried out according to the

manufacturer's (Bio Rad) protocol. In short, master mixes containing ddPCR Supermix for Probes

(No dUTP) (Bio Rad), DNA samples, primer / probe mixes, and HindIII-HF restriction enzyme

(New England Biolabs) for 28S rDNA reactions (no restriction enzyme for control gene reactions)

were prepared in 0.2 mL Eppendorf tubes, and incubated at room temperature for 15 minutes to

allow for restriction enzyme digestion. ddPCR droplets were generated from samples using

QX200 Droplet Generator (Bio Rad) and underwent complete PCR cycling on a C100 deep-well

thermocycler (Bio Rad). Droplet fluorescence was read using the QX200 Droplet Reader (Bio

Rad). Sample copy number was determined using Quantasoft software (Bio Rad). rDNA copy

number per genome was determined by 28S sample copy number multiplied by 100 (due to the

100x dilution of sample in the 28S reaction compared to control reaction) divided by control gene

copy number multiplied by the expected number of control gene copies per genome (2 for RpL in

all samples; 2 for Upf1 in female samples; 1 for Upf1 in male samples). The 28S copy number

values determined by each control gene was averaged to determine 28S copy number for each

sample.

Mitotic chromosome spread and DNA FISH

Larval brain dissections, mitotic chromosome spreads, DNA FISH, and imaging were all done as

previously(*38*). In short, brains were dissected from male third instar larvae in PBS and fixed in

25μ μL of acetid acid and 4% formaldehyde in PBS. Samples were applied to Superfrost plus

slides and manually squashed under a coverslip, then immediately frozen in liquid N2. After

freezing slides were removed from N2, coverslip removed, and slides were dehydrated in 100%

ethanol and dried at room temperature. DNA FISH hybridization was performed in 20 μL of 50%

formamide, 10% dextran sulfate, 2X SSC buffer and 0.5 μM each probe applied directly to the

sample on the slide and covered with a cover slip. Samples were incubated at 95° for 5 minutes,

cooled and wrapped in parafilm, then incubated overnight at room temperature in a dark humid

chamber. Coverslips were removed and slides were washed three times for 15 minutes in 0.1x

SSC, dried, and then mounted in VECTASHIELD with DAPI (Vector Labs). Samples were imaged

using a Leica Stellaris 8 confocal microscope with 63x oil-immersion objectives and processed

using Fiji (ImageJ) software. Probes used for this study are as follows: 359, 5'-

AGGATTTAGGGAAATTAATTTTTGGATCAATTTTCGCATTTTTTGTAAG-3'-Cy5; $(TAGA)_6$-

Cy5;   IGS,   5'-

AGTGAAAAATGTTGAAATATTCCCATATTCTCTAAGTATTATAGAGAAAAGCCATTTTAGTGA

ATGGA-3'-Alexa488; $(AATAC)_6$-Cy3; and $(AATAAAC)_6$-Cy3.

Generation of rDNA deletion animals

rDNA copy number loss was induced during larval development in *yw* / Y ;; *HS-I-CreI*, *Sb* / TM6B

males with a *y, w* X chromosome by I-CreI expression as previously described(*26*). In brief,

parental animals mated and laid eggs for three days, then removed from food. After one additional

day of larval development, animals were exposed to 37° C heat shock for 45 minutes on two

consecutive days. identify X chromosomes with significant rDNA copy number reduction (bb),

adult males that experienced I-CreI expression were mated to bb^{158} / FM6 females, and virgin

non-*FM6* daughters (bb / bb^{158}) were screened for the *bobbed* phenotype. 28 out of 946 non-*FM6*

daughters screened were *bobbed*. To isolate potentially reduced rDNA loci and remove HS-I-CreI

from the background, TM6B containing *bobbed* females were individually mated to wild-type

males. Male offspring from each individual female candidates were subsequently individually

mated to bb^{158} / FM6 females. Any mating that failed to produce non-FM6 daughters were

eliminated (due to having the bb^{158} and not the candidate bb chromosome). All viable non-FM6

daughters were double checked for the *bobbed* phenotype, and stocks with all *bobbed* non-FM6

daughters had FM6 containing siblings collected and used to establish a bb / FM6 stock. This

method isolated the novel rDNA deletion allele, bb^{Z9}, used in this study.

Drosophila genetics

All Drosophila lines used in this study are found in **Table S2**. All animals we reared on standard

Bloomington medium at 25⁰. All aging was done in roughly 1:1 mixed presence of males and

females, provided fresh food every four to six days. UAS:R2 RNAi strains were designed using

SNAP-DRAGON shRNA target software, and oligos containing target hairpin sequence were

cloned into the WALLIUM20 vector for phiC31 site-directed integration into the Drosophila

genome. All UAS:R2 transgenes were synthesized into pUAST:attB by VetcorBuilder (Chicago,

IL) for site-directed integration. All injections and selection of animals containing integrated

transgenes were performed by BestGene, Inc (Chino Hills, CA).

Lineage survival assay

Independent lineages of nos:Gal4 / CyO; UAS:R2 RNAi / Tm6B or nos:Gal4 / CyO; TM2 / TM6B

animals were established by collecting siblings of the indicated genotypes from nos:Gal4 / CyO;

TM2 / TM6B males mated to Sp / CyO; UAS:R2 RNAi / TM6B females. At each generation in

each lineage, 3 males were mated with 5 females for 5 days, and offspring were collected 10 days

after mated ended. Any lineages that did not have at least 3 males and 5 females at collection

time were terminated due to insufficient animals.

Statistics

For all comparisons of percentage of samples with categorical values (percent γH2Av or R2

positive cells), significance was determined by chi-squared test. For all comparisons of samples

with independent values (number of GSCs; rDNA copy number), significance was determined by

Student's t-test between experimental and control conditions, unless otherwise indicated.

Figure S1: Design and efficiency of R2 RNAi. (A) Diagram the 45S rRNA cistron constituting an

'rDNA copy' and the sequence-specific R2 insertion site in the 28S rRNA gene. **(B)** Confirmation of

R2 expression knockdown by R2 RNAi. Precent of GSCs expressing R2 detected by RNA FISH in

nos-gal4 driver only condition (nos>) and R2 RNAi expressed by the *nos-gal4* driver (nos>R2 RNAi).

P-value determined by chi-squared test. Error bars = 95% CI.

Figure S2: Diagram of phenotypic assessment to detect rDNA magnification at the bb^{Z9}

allele. (A) rDNA CN quantification of the bb^{Z9} allele determined by ddPCR. All samples from

individual females with either the bb^{Z9} allele or X chromosome from which the allele was isolated

heterozygous with an X chromosome completely lacking rDNA (bb^{158}). P-value determined by

Student's t-test. Error bars = 95% CI. **(B)** Scheme to detect rDNA magnification of bb^{Z9} allele.

Tested magnifying males (bb^{Z9} / Ybb^{0}) are crossed to females heterozygous for the bb^{158} rDNA

deletion allele. Resultant bb^{Z9} / bb^{158} daughters completely rely on the bb^{Z9} locus for their rDNA.

(Left) bb^{Z9} / bb^{158} offspring with the 'bobbed' cuticular defects associated with rDNA insufficiency

(red arrow) are determined to have inherited a bb^{Z9} allele that did not undergo rDNA

magnification. (Right) bb^{Z9} / bb^{158} offspring with wild-type cuticles are determined to have

inherited a bb^{Z9} allele that restored normal rDNA content through rDNA magnification (bb^{Z9-Mag}).

Thus, the portion of bb^{Z9} / bb^{158} offspring with wild-type cuticles represents the portion of bb^{Z9}

alleles that underwent rDNA magnification. bb^{Z9} / bb^{158} animals are also collected for rDNA CN

quantification, and total rDNA CN in these animals represents the rDNA from the bb^{Z9} locus

alone.

Figure S3: Transgenic R2 expression induces double stranded breaks at rDNA loci. (A)

Alignment of 44 amino acids in the endonuclease domain of *Bombyx mori* R2 (Bm) and

Drosophila melanogaster R2 (Dm). Residues shown to be essential for Bm R2 endonuclease

activity(*23*) labeled in blue and respective Dm R2 residues in red. Mutations found to render Bm

R2 endonuclease dead(*23*) and their respective amino acid changes in Dm R2 are indicated

above highlighted residues. **(B-C)** DNA FISH in mitotic chromosome spreads of larval

neuroblasts in animals with the neuroblast driver *wor-gal4* (wor>, B) or transgenic R2 expressed

by the *wor-gal4* driver (wor>R2, C) animals. rDNA is labelled by probes targeting the rDNA

intergenic sequence (IGS) (Green), Y- chromosome is labelled by probes targeting AATAC and

AATAAAC satellite sequences (Cyan) and X-chromosome is labelled by probes against 359 and

TAGA satellite sequences (Red). Y* indicates Y-chromosome fragments broken at the rDNA

locus. **(D)** Percentage of neuroblasts with DSBs at rDNA loci determined by DNA FISH in mitotic

spreads in indicated genotypes. P-values determined by chi-squared test. Error bars = 95% CI.

(E) Percentage of γH2Av positive GSCs in the indicated genotypes. Transgenic R2 constructs

expressed in the germline by the *nos-gal4* driver (nos>). P-value determined by chi-squared

test. Error bars = 95% CI.

Figure S4: Models of rDNA CN expansion initiated by DSBs at rDNA locus. (A) rDNA CN

expansion by strand-dependent strand annealing (SDSA). DSBs at rDNA loci can be repaired

via recombination of a single strand with sister chromatid and synthesis. Synthesis can run on to

replicate multiple rDNA copies before re-annealing with broken strand. Upon repair after

synthesis, rDNA CN is expanded on one sister chromatid while CN of the other sister remains

intact. **(B)** DSBs formed at rDNA loci induce homology-dependent repair that causes

recombination with existing extrachromosomal rDNA circles (ERCs), allowing for reintegration of

the ERC rDNA copies into the genomic rDNA locus.

Figure S5: R2 expression in GSCs is regulated in response to changing rDNA copy

number during aging. (A-C) R2 RNA FISH in the male germline of young (0-days old), (A) old

(40-days old) (B), and recovered offspring of old animals (C). Isolated R2 channel in A'-C'.

GSCs indicated by yellow dotted circle. * indicates GSC niche signal hub. R2 positive cells

indicated by yellow arrowhead. **(D)** Percentage of R2 positive GSCs at indicated ages in the

offspring of 0- or 40-day old fathers. P-value determined by chi-squared test. Error bars = 95%

CI.

Figure S6: Scheme for testing lineage survival in multi-generational R2 RNAi experiment.
Each lineage is established using 3 males and 5 females of the given genotype. Males and
females are mated for 5 days and removed from the vial. 10 days later, all offspring are
collected. If total offspring from the mating contains at least 3 males and 5 females, then 3
males and 5 females are randomly selected to establish the next generation of the lineage. If
the total offspring contained fewer than 3 males or 5 females, then the lineage was terminated
and future generations were not established. This procedure was repeated at each generation
for each lineage for 6 generations or until termination.

184   **Table S1: List of oligomers used in this study for ddPCR assays.**

Oligo Name	Type	Target	Sequence	Modifications
dd-RpL32 F	Primer	RpL32	GCTTCAAGGGACAGTATCTG	
dd-RpL32 R	Primer	RpL32	AACGCGGTTCTGCATGAG	
dd-RpL32 Probe	Probe	RpL32	ATGCCCAACATCGGTTAC	5' HEX AND 3' Iowa Black FQ
dd-28S F	Primer	28S	GAGCTGCCATTGGTACAG	
dd-28S R	Primer	28S	GCTTTCGCCTTGAACTTAG	
dd-28S Probe	Probe	28S	TGGTGGATAGTAGCAAATAATCG	5' 6-FAM AND 3' Iowa Black FQ
dd-Upf1 F	Primer	Upf1	CACACTTTATGTCCACCATTATTG	
dd-Upf1 R	Primer	Upf1	GAGTTTCCGTAGGGACCAC	
dd-Upf1 Probe	Probe	Upf1	CCG TAA CCG CCA CTG CGG T	5' 6-FAM AND 3' Iowa Black FQ

187   **Table S2: List of Drosophila stocks used in this study.**

Genotype	Source	Use
Dp(1;f)122; C(1)RM, y^1 v^1 f^1 / C(1;Y)6, Df(1)259, w^1	Kyoto Drosophila Genomics and Genetic Resource Center (DGRC) #107274	Mini-x Chromosome (Dp(1;f))
yw	Bloomington Stock Center (BSC) #1495	Source for isolating novel rDNA deletion strain
w[1118]; P{v[+t1.8]=hs-I-CreI.R}1A Sb[1]/TM6	BSC #6937	I-CreI endonucelase
nos-gal4	PMID: 9501989	Early germ cell driver
wor-gal4	BSC #56553	Neuroblast driver
bam-gal4	PMID: 12571107	Germ cell driver
y[1] eq[1]/Df(YS)bb[-]	DGRC #101260	Y chromosome complete rDNA deletion
Df(1)bb158, y[1] / Dp(1;Y)y[+] / C(1)*; ca[1] awd[K]	DGRC #106876	X chromosome complete rDNA deletion

36. Cheng, J. *et al.* Centrosome misorientation reduces stem cell division during ageing. *Nature* **456**, 599–

604 (2008).

37. Huang, A. M., Rehm, E. J. & Rubin, G. M. Quick Preparation of Genomic DNA from Drosophila.

Cold Spring Harb Protoc **2009**, pdb.prot5198-pdb.prot5198 (2009).

38. Jagannathan, M., Warsinger-Pepe, N., Watase, G. J. & Yamashita, Y. M. Comparative Analysis of

Satellite DNA in the Drosophila melanogaster Species Complex. *G3 Genes Genomes Genetics* **7**, 693–

704 (2017).