Title: The retrotransposon *R2* maintains *Drosophila* ribosomal DNA repeats

One Sentence Summary: The retrotransposon *R2* initiates restoration of ribosomal DNA copies

to trans-generationally maintain essential locus.

Authors: Jonathan O. Nelson[1]*, Alyssa Slicko[1,3], and Yukiko M. Yamashita[1,2,3]*

Affiliations: [1]Whitehead Institute for Biomedical Research,

[2]Department of Biology, MIT,

[3]Howard Hughes Medical Institute,

455 Main Street, Cambridge, MA 02142

*Corresponding Authors: jonnels@wi.mit.edu, yukikomy@wi.mit.edu

Abstract: Ribosomal DNA (rDNA) loci contain the hundreds of tandemly repeated copies of ribosomal RNA genes needed to support cellular viability. This repetitiveness makes it highly susceptible to copy number (CN) loss, threatening multi-generational maintenance of rDNA. How this threat is counteracted to avoid extinction of the lineage has remained unclear. Here, we show that the rDNA-specific retrotransposon *R2* is essential for rDNA CN maintenance in the *Drosophila* male germline, despite the perceived disruptive nature of transposable elements. Depletion of *R2* led to defective rDNA CN maintenance, causing a decline in fecundity over generations and eventual extinction. This study reveals that active retrotransposons can provide a benefit to their hosts, contrary to their reputation as genomic parasites, which may contribute to their widespread success throughout taxa.

Main Text:

Ribosomal RNAs (rRNAs) account for 80-90% of all transcripts in eukaryotic cells(*1*). To meet this demand, the ribosomal DNA (rDNA) gene that codes for rRNA is tandemly repeated hundreds of times, comprising rDNA loci on eukaryotic chromosomes. Ironically, this repetitive structure is susceptible to intra-chromatid recombination that causes rDNA copy number (CN) loss **(Fig. 1A)**, which is a major cause of replicative senescence in budding yeast (*2*). Evidence of similar rDNA CN instability has been noted in some tissues from aged dogs and humans(*3*, *4*). Critically, age-associated rDNA CN loss also occurs in the *Drosophila* male germline and is inherited by the next generation(*5*). The essential yet unstable nature of rDNA raises the question as to how the degeneration of rDNA loci over successive generations is prevented to avoid the extinction of the lineage. Intensive studies have revealed that sister chromatid recombination mediates rDNA CN expansion in yeast, thereby maintaining rDNA repeat abundance over generations(*2*). rDNA CN is variable between individuals of most species but maintained within a consistent range throughout the population(*6*), implying that transgenerational dynamic CN changes (loss and restoration) are a common feature of rDNA maintenance. However, the mechanism of

transgenerational rDNA CN maintenance in multicellular organisms has remained a mystery.

Over 50 years ago, the phenomenon of 'rDNA magnification' in *Drosophila* was first described as

the process wherein aberrant rDNA loci bearing minimal rDNA repeats recover to a normal rDNA

CN(*7*, *8*). rDNA magnification requires genes involved in homologous recombination-mediated

repair, similar to yeast, which can duplicate tandemly repeated elements(*9*). Despite the robust

rDNA CN expansion activity, it remained unclear if the mechanisms of rDNA magnification served

a physiological function in natural populations. We recently demonstrated that offspring who

inherited reduced rDNA CN from old fathers could also recover rDNA CN in their germline(*5*). This

recovery depends on the same set of the genes as rDNA magnification, leading us to propose

that rDNA magnification is a manifestation of the physiological mechanisms to maintain rDNA CN

across generations(*6*). However, the underlying factors responsible for rDNA magnification

remains poorly understood.

Metazoan rDNA genes are frequently inserted by rDNA-specific transposable elements

(TEs), such as the retrotransposon *R2* in *Drosophila*. *R2* is found throughout arthropods and *R2*-

like elements are widely present across taxa, including Cnidaria, Planaria, nematodes, fish, birds,

and reptiles(*10*, *11*). These TEs use their sequence-specific nuclease to mobilize specifically

within rDNA loci(*12*), inserting into rDNA genes and likely disrupting 28S rRNA function(*13*) **(Fig.**

S1A). Surprisingly, we found that RNAi-mediated knockdown of *R2* in the *Drosophila* male

germline (*nos-gal4*-driven expression of RNAi lines, *nos>R2i-1* or *R2i-2,* hereafter) **(Fig. S1A-C)**

resulted in premature loss of germline stem cells (GSCs) during aging **(Fig. 1B-F)**. GSCs

continuously produce differentiating germ cells to sustain sperm production throughout adulthood

and thus are the source of the genome passed to the next generation(*14*). Whereas newly eclosed

R2 RNAi males contained similar numbers of GSCs to controls, GSC number more rapidly

declined during aging in *R2* knockdown males compared to controls **(Fig. 1F)**. Given that *R2* is

specifically inserted into rDNA, we also examined the effect of *R2* knockdown on rDNA stability.

Using highly quantitative droplet digital PCR (ddPCR), we found that RNAi-mediated knockdown

of *R2* enhanced rDNA CN loss in the testis during aging **(Fig. 1G-H)**. Interestingly, one of the

RNAi constructs (*R2i-2*) suffered rapid rDNA CN loss within the first 10 days of adulthood, but

recovered by 20 days of age **(Fig. 1G)**. As GSC number was also initially more drastically affected

with this RNAi line **(Fig. 1F)**, we speculate that severe rDNA loss caused by this RNAi construct

(*R2i-2*) may select for GSCs that are less sensitive to *R2* RNAi activity. rDNA CN loss in the

germline was further confirmed by DNA FISH on the meiotic chromosomes **(Fig S2A-F)**. rDNA

CN insufficiency is likely the primary cause of GSC loss in *R2* RNAi animals, because increasing

total rDNA CN via introduction of a mini-chromosome harboring an rDNA locus(*15*) suppressed

the premature GSC loss caused by *R2* knockdown **(Fig. 1I)**. These results revealed that *R2*

contributes to sustaining GSC population during aging through rDNA CN maintenance,

uncovering an unanticipated benefit of the *R2* retrotransposon to the host, despite the widely-held

view of TEs being genetic parasites.

Why is *R2* necessary for rDNA copy number maintenance? We found that *R2* is required

for rDNA magnification. rDNA magnification is detected as the emergence of offspring with normal

cuticle from fathers with abnormal ('bobbed') cuticle caused by insufficient rDNA CN(*7*) **(Fig. 2A)**.

Drosophila melanogaster rDNA loci reside on the sex chromosomes (X and Y)(*16*), and rDNA

magnification almost exclusively occurs to X chromosome rDNA loci harboring the minimal viable

amount of rDNA (bb^{Z9}, **Fig. S3A**) when combined with a Y chromosome lacking rDNA (bb^{Z9}/Ybb^{0},

'magnifying males' hereafter)(*7*) **(Fig. S3B)**. Importantly, rDNA magnification never occurs in

males with a normal Y chromosome containing intact rDNA (bb^{Z9}/Y^{+}, 'non-magnifying males'

hereafter)(*8*), indicating the presence of mechanisms to monitor rDNA CN to activate expansion.

We found that *R2* knockdown reduces rDNA magnification from 13.73% (control, n = 233) to 0%

(*R2i-1*, n = 181, p = 5.6×10^{-7}) and 2.36% (*R2i-2*, n = 127, p = 9.9×10^{-4}) **(Fig. 2B)**. Moreover,

quantification of rDNA CN by ddPCR revealed that 87.5% of bb^{Z9} chromosomes increased rDNA

CN in magnifying males (n = 96, p = 1.8×10^{-4}), with an average increase of 18.29 rDNA copies

across all samples (n = 96, p = 3.1×10^{-12}) **(Fig. 2C)**. This widespread rDNA CN increase indicates

that rDNA magnification broadly increases rDNA CN throughout the germline, despite only

13.73% of bb^{Z9} chromosomes recovering enough CN to support normal cuticle development. This

rDNA CN increase in magnifying males is also eliminated upon *R2* knockdown **(Fig. 2C)**. These

results reveal that *R2* is required for rDNA CN expansion during rDNA magnification. Interestingly,

we found that rDNA magnification was blocked only when the *R2* RNAi constructs were expressed

by the *nos-gal4* driver in early germ cells (including GSCs), but not when expressed in later germ

cells by the *bam-gal4* driver (see Methods) **(Fig. 2B)**. These results indicate that *R2* primarily

functions in the earliest stages of germ cells (including GSCs) to support rDNA magnification.

We further found that *R2* is sufficient for rDNA CN expansion. Ectopic expression of

transgenic *R2* in the germline **(Fig. S4A-F)** induced rDNA magnification of the bb^{Z9} locus in non-

magnifying males (bb^{Z9}/Y^+) **(Fig. 2D-E)**. While rDNA magnification assessed by dorsal cuticle is

never detected from control males (bb^{Z9}/Y^+ without *R2* expression), we found 3.3% of male

offspring exhibited magnification (normal cuticle) upon expression of transgenic *R2* **(Fig. 2D,** n =

877, p = 3.2×10^{-5}). Importantly, reversion of the cuticle phenotype was heritable to the subsequent

F2 generation throughout our experiments, confirming that CN restoration occurred in the

germline **(Fig. S5A-C)**. Quantification of rDNA CN by ddPCR revealed that ectopic

overexpression of *R2* in non-magnifying males (bb^{Z9}/Y^+) also increases the average rDNA CN at

bb^{Z9} rDNA loci among all offspring **(Fig. 2E**, n = 94, p = 0.0256), revealing *R2* is sufficient to

induce rDNA CN expansion. Critically, expression of a nuclease dead *R2* transgene (NucDead*R2*)

in non-magnifying males **(Fig. S4A-E)** failed to induce rDNA magnification **(Fig. 2D)**, suggesting

that the nuclease activity of *R2* is essential for its ability to induce rDNA CN expansion.

How does *R2*'s nuclease activity contribute to rDNA magnification? In yeast, rDNA CN

expansion is initiated by double-stranded breaks (DSBs) at the rDNA intergenic sequence, which

induces sister chromatid recombination that results in rDNA gene duplication(*17*). All proposed

models of *Drosophila* rDNA CN expansion (the most prominent model being unequal sister

chromatid recombination(*18*)) require an initiating DSB at the rDNA locus, **(Fig. 3A; Fig. S6A-B)**.

Indeed, artificial introduction of DSBs at rDNA loci by I-CreI endonuclease expression has been

reported to induce rDNA magnification(*19*), but the endogenous factor that induces rDNA

magnification remained unclear. *R2* is capable of creating DSBs through sequential nicking of

opposite DNA strands during retrotransposition(*10*). It has been speculated that DSBs created

during *R2* retrotranspostion may be an initiating event of rDNA magnification(*9*), although this

possibility has yet to be empirically tested. We confirmed that ectopic overexpression of *R2*, but

not NucDead*R2*, indeed induces chromosomal breaks at rDNA loci identified by chromosome

spreads **(Fig. S4B-D)**. *R2* overexpression (but not NucDead*R2*) in the germline also increased

the frequency of GSCs with DSBs, identified by γH2Av expression **(Fig. S4E)**. Next, we found

rDNA magnification is associated with an elevation in DSBs in GSCs: the frequency of γH2Av-

positive GSCs is increased in magnifying males (bb^{Z9}/Ybb^0) compared to non-magnifying males

(bb^{Z9}/Y^+) **(Fig. 3B-C, E**; n = 519, p = $8.8x10^{-4}$). Strikingly, we observed that knockdown of *R2* in

magnifying males reduced the frequency of γH2Av-positive GSCs to levels comparable to non-

magnifying males **(Fig. 3D-E;** n = 537, p = $7.1x10^{-4}$ for *R2i-1*; n = 521, p = $7.9x10^{-4}$ for *R2i-2***)**,

indicating that *R2* is responsible for the DSBs formed in GSCs during rDNA magnification. Taken

together, these results suggest that rDNA-specific endonuclease activity of *R2* creates DSBs at

the rDNA loci that may in turn induce rDNA CN expansion.

Given the threat *R2* mobilization poses to the host genome, both by disruption of rRNA

function and causing excessive DSB formation(*10*), how is the potential benefit of *R2* to rDNA CN

maintenance balanced with the detriment of *R2* retrotransposition? We found *R2* expression in

the germline is specifically de-repressed under conditions of reduced rDNA CN, potentially

explaining how the conflicting consequences of *R2* expression are resolved. Using RNA

fluorescence *in situ* hybridization (RNA FISH) to examine *R2* expression at a single cell resolution,

we found that the frequency of GSCs expressing *R2* was significantly increased in magnifying

males (bb^{Z9}/Ybb^0), whereas non-magnifying males (bb^{Z9}/Y^+) rarely expressed *R2* **(Fig. 3F-H**; n =

231, p = 1.7×10^{-10}). Moreover, we found that GSCs from aged animals and the sons of old fathers,

which inherit reduced rDNA CN(*5*)**,** also exhibited a higher frequency of *R2* expression compared

to GSCs from young flies **(Fig. S7A-B, D**; n = 1,247, p = 8.3×10^{-4} for old animals; n = 1,107, p =

1.5×10^{-4} for offspring). Importantly, the frequency of *R2* expression among GSCs in the sons of

old fathers returned to the basal level after 20 days of age, when rDNA CN was shown to have

recovered(*5*) **(Fig. S7C-D**; n = 617, p = 0.036**)**. These results indicate that *R2* expression is

dynamically regulated in response to changing rDNA CN. Taken together, we propose that *R2*

expression is finely tuned to function when most beneficial to the host while minimizing

unnecessary exposure to the harmful effects of transposition.

Based on the finding that *R2* plays a critical role in maintaining germline rDNA CN, we

postulated that *R2* is essential to prevent continuous multi-generational rDNA loss capable of

causing the extinction of the lineage. In *C. elegans*, the loss of genome integrity is known to cause

gradual loss of fertility, a phenotype known as <u>mor</u>tal <u>g</u>ermline (morg)(*20*). To test whether *R2*-

mediated rDNA maintenance is required to maintain fertility through generations, we established

multiple independent lines expressing *R2* RNAi in their germline and tracked their fecundity at

each generation through the ability of each line to produce sufficient offspring to establish a new

generation **(Fig. S8)**. While nearly all control lines survived throughout the duration of the

experiments, we found that lines expressing the *R2i-1* RNAi construct failed to consistently

produce sufficient progeny, with over half failing by the fourth generation **(Fig. 4A)** (n = 43, p =

3×10^{-6}), indicating that *R2* is essential for continuity of the germline lineage. Surviving males of

extinguishing *R2i-1* lineages had ~20% reduction in rDNA CN compared to control lines (n = 22,

p = 0.031) **(Fig. 4B)**. With the *R2i-2* RNAi, the lineage was maintained relatively well, after initial

sharp drop **(Fig. 4A)**: Considering that *R2* knockdown by the *R2i-2* construct exhibits only

transient rDNA CN decrease at day 10 (**Fig S2A**), we speculate that this RNAi construct quickly

selects for the germ cells and lineages insensitive to *R2* knockdown. Taken together, these results

suggest that *R2*-mediated maintenance of rDNA contributes to germline immortality.

Together, our findings reveal an unanticipated 'function' of retrotransposon activity to

benefit the host genome through a role in rDNA CN maintenance. Repetitive DNA sequences are

among the most vulnerable elements of the eukaryotic genome(*21*). We propose that DSBs

generated by *R2* in GSCs with reduced rDNA CN stimulate sister chromatid exchange that results

in rDNA CN expansion **(Fig. 4C)**. Our proposed 'function' for *R2* in rDNA maintenance may

represent a novel mutualistic host-TE relationship, which are rarely described in eukaryotes(*22*).

R2's de-repression when it can be beneficial (i.e. decreased rDNA CN) may be the key to this

mutualistic host-TE relationship. It awaits future investigation to understand how the mechanisms

that repress *R2*(*10*) may tune this activity for host's benefit. The widespread presence of *R2* and

other rDNA-specific TEs in both vertebrates and invertebrates(*11*) suggests that similar host-TE

mutualism may support rDNA CN maintenance throughout Metazoa. Interestingly, many of rDNA-

specific TEs have little sequence similarity to *R2*, instead appearing to be derived from other non-

specific TEs(*11*), suggesting this host-TE mutualism may have evolved multiple times over the

course of evolution. In summary, our study provides an example of mutualistic retrotransposons

in the maintenance of eukaryotic genomes, and we propose that more mutualistic TEs are yet to

be discovered.

Acknowledgements

We thank the Bloomington *Drosophila* Stock Center, Kyoto *Drosophila* Stock Center and

Developmental Studies Hybridoma Bank for reagents. We thank the Yamashita lab members

and Dr. Andy Clark for discussion and comments on the manuscript. We thank ATCC for design

of the 28S and RpL ddPCR assays. **Funding:** This research was supported by the Howard

Hughes Medical Institute. Jonathan Nelson was supported by an American Cancer Society

Postdoctoral Fellowship (133949-PF-19-133-01-DMC). **Authors contributions:** JON:

Conceptualization, Methodology, Verification, Formal analysis, Investigation, Resources, Data

curation, Writing – Original draft, Writing – Review & editing, Visualization, Funding acquisition.

AS: Validation, Investigation, Resources, Data curation, Writing – Review & editing. YMY

Conceptualization, Methodology, Resources, Writing – Original draft, Writing – Review &

editing, Visualization, Supervision, Project administration, Funding. **Competing interests:** The

authors declare no competing interests. **Data materials availability:** All data is available in the

manuscript or the supplementary materials.

References:

1. A. F. Palazzo, E. S. Lee, Non-coding RNA: what is functional and what is junk? *Frontiers Genetics*. **6**,
2 (2015).

2. T. Kobayashi, Ribosomal RNA gene repeats, their stability and cellular senescence. *Proc Jpn Acad Ser*
B. **90**, 119–129 (2014).

3. B. L. Strehler, M.-P. Chang, L. K. Johnson, Loss of hybridizable ribosomal DNA from human post-
mitotic tissues during aging: I. Age-dependent loss in human myocardium. *Mech Ageing Dev*. **11**, 371–
378 (1979).

4. R. Johnson, B. L. Strehler, Loss of Genes coding for Ribosomal RNA in Ageing Brain Cells. *Nature*.
240, 412–414 (1972).

5. K. L. Lu, J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita, Transgenerational
dynamics of rDNA copy number in Drosophila male germline stem cells. *Elife*. **7**, e32421 (2018).

6. J. O. Nelson, G. J. Watase, N. Warsinger-Pepe, Y. M. Yamashita, Mechanisms of rDNA Copy Number
Maintenance. *Trends Genet*. **35**, 734–742 (2019).

7. F. M. Ritossa, Unstable redundancy of genes for ribosomal RNA. *Proc National Acad Sci*. **60**, 509–516
(1968).

8. K. D. Tartof, Regulation of ribosomal RNA gene multiplicity in Drosophila melanogaster. *Genetics*.
73, 57–71 (1973).

9. R. S. Hawley, C. H. Marcus, Recombinational Controls of rDNA Redundancy in Drosophila. *Annu Rev*
Genet. **23**, 87–120 (1989).

10. T. H. Eickbush, D. G. Eickbush, Integration, regulation, and long-term stability of R2
retrotransposons. *Microbiol Spectr*. **3**, 1127–1146 (2015).

11. K. K. Kojima, H. Fujiwara, Long-Term Inheritance of the 28S rDNA-Specific Retrotransposon R2.
Mol Biol Evol. **22**, 2157–2165 (2005).

12. J. Yang, H. S. Malik, T. H. Eickbush, Identification of the endonuclease domain encoded by R2 and
other site-specific, non-long terminal repeat retrotransposable elements. *Proc National Acad Sci*. **96**,
7847–7852 (1999).

13. D. G. Eickbush, T. H. Eickbush, R2 Retrotransposons Encode a Self-Cleaving Ribozyme for
Processing from an rRNA Cotranscript ▽ . *Mol Cell Biol*. **30**, 3142–3150 (2010).

14. M. T. Fuller, A. C. Spradling, Male and Female Drosophila Germline Stem Cells: Two Versions of
Immortality. *Science*. **316**, 402–404 (2007).

15. D. L. Lindsley, L. Sandler, The Meiotic Behavior of Grossly Deleted X Chromosomes in Drosophila
Melanogaster. *Genetics*. **43**, 547–63 (1958).

16. F. M. Ritossa, K. C. Atwood, S. Spiegelman, A molecular explanation of the bobbed mutants of
Drosophila as partial deficiencies of "ribosomal" DNA. *Genetics*. **54**, 819–34 (1966).

17. S. Gangloff, H. Zou, R. Rothstein, Gene conversion plays the major role in controlling the stability of
large tandem repeats in yeast. *Embo J*. **15**, 1715–1725 (1996).

18. K. D. Tartof, Unequal Mitotic Sister Chromatid Exchange as the Mechanism of Ribosomal RNA
Gene Magnification. *Proc National Acad Sci*. **71**, 1272–1276 (1974).

19. S. Paredes, K. A. Maggert, Expression of I-CreI Endonuclease Generates Deletions Within the rDNA
of Drosophila. *Genetics*. **181**, 1661–1671 (2009).

20. C. Smelick, S. Ahmed, Achieving immortality in the C. elegans germline. *Ageing Res Rev*. **4**, 67–82
(2005).

21. R. E. Brown, C. H. Freudenreich, Structure-forming repeats and their impact on genome stability.
Curr Opin Genet Dev. **67**, 41–51 (2021).

22. R. L. Cosby, N.-C. Chang, C. Feschotte, Host–transposon interactions: conflict, cooperation, and
cooption. *Gene Dev*. **33**, 1098–1116 (2019).

255
256

List of Supplementary Materials

Supplementary Materials

Materials and Methods

Fig S1 – S8

Table S1 – S2

References 23 - 30

Figure 1: *R2* is required for GSC maintenance via rDNA CN maintenance during

***Drosophila* male germline aging. (A)** Model of rDNA repeat instability. **(B-E)** GSCs in 0- and

40-day old control (B-C) and *R2* RNAi testes (D-E). Yellow dotted circle = GSC. GSC signaling

niche indicated by *. Green = Vasa, White = Fascillin III, Blue = DAPI. Scale bar = 7.5 μM **(F)**

Average GSCs per testis in control and two *R2* RNAi lines during aging. * indicates $p < 10^{-3}$

determined by Student's t-test. Error = 95% confidence interval (CI). **(G-H)** Testis rDNA CN

determined by ddPCR. P-value by Student's t-test. Error = 95% CI. **(I)** Average GSCs per testis

during aging in Control and two *R2* RNAi lines containing mini-X chromosome. Error = 95% CI.

Figure 2: *R2* is necessary and sufficient for rDNA magnification. (A) Diagram of rDNA

magnification at the *bb^Z9* rDNA locus, during which dorsal cuticle defect (red arrow) revert to

normal cuticle. **(B)** Percent magnified offspring determined by cuticular phenotype in offspring

from magnifying males. P-value determined by chi-squared test. Error = 95% CI. **(C)** Mean *bb^Z9*

locus rDNA CN determined by ddPCR in daughters from males. P-value determined by

Student's t-test. Error = 95% CI. **(D)** Percent magnified offspring from non-magnifying males. P-

282   value determined by chi-squared test. Error = 95% CI. **(E)** Mean bb^{Z9} locus rDNA CN

283   determined by ddPCR in daughters from non-magnifying males. Non-magnifying condition is the

284   same data as panel (C). P-value determined by Student's t-test. Error = 95% CI.

Fig 3: Derepressed *R2* creates DSBs in GSCs during rDNA magnification. (A) Diagram of

rDNA CN expansion by unequal sister chromatid exchange during DSB repair at rDNA loci.

Recombination between misaligned rDNA copies during DSB repair result in crossovers that

create unequal sister chromatid exchange that increases rDNA CN on one chromatid. **(B-D)**

Detection of DSBs in the early adult male germline by anti-γH2Av staining. *R2* RNAi expressed

under the *nos-gal4* driver. Non-RNAi conditions contain the *nos-gal4* driver alone. GSCs

indicated by yellow dotted circle. Blue = DAPI, Green = vasa, Magenta = γH2Av, white = FasIII.

The hub is indicated by *. Scale bar = 10 μM. **(E)** Percentage of γH2Av positive GSCs. P-value

294 determined by chi-squared test. Error = 95% CI. **(F-G)** *R2* expression in GSCs (yellow dotted

295 circle). Blue = DAPI, Green = *R2* mRNA. Isolated *R2* channel in F'-G'. The hub is indicated by *.

296 *R2* positive cells GSCs are marked by yellow arrowhead. **(H)** Percentage of *R2* positive GSCs.

297 P-value determined by chi-squared test. Error = 95% CI.

Fig 4: *R2* is required to maintain rDNA CN and fertility over successive generations. (A)
Kaplan-Meier curve of lineage survival in control (*nos-gal4* driver alone and two *R2* RNAi
expressing (via the *nos-gal4* driver) lineages. Each lineage constitutes an individual data point.
P-values determined by log rank test. **(B)** rDNA CN determined by ddPCR in males of control
animals at the 6th generation or *R2* RNAi lineages at their terminating generation. P-value
determined by Student's t-test. Error = 95% CI. **(C)** Model of the role of *R2* in germline rDNA CN
maintenance.