Pparα and fatty acid oxidation coordinate hepatic transcriptional architecture.

Running title: Fatty acids alter the epigenome.

Kyle S. Cavagnini,[1] and Michael J. Wolfgang[1,2,3*]

[1]Department of Biological Chemistry, [2]Pharmacology and Molecular Sciences,

The Johns Hopkins University School of Medicine, Baltimore, Maryland.

[3]Lead Contact

*Address correspondence to**: Michael J. Wolfgang, Ph.D.,** Department of Biological Chemistry,

Johns Hopkins University School of Medicine, 855 N. Wolfe St., 475 Rangos Building, Baltimore,

MD 21205, Tel: 443-287-7680, Email: mwolfga1@jhmi.edu

HIGHLIGHTS

• Fasting-induced transcription and histone acetylation are largely independent of acetyl-

CoA concentration.

• Deficits in fatty acid oxidation prompt epigenetic changes and Pparα-sensitive

transcription.

• Fasting prompts enhancer priming and acetylation proximal to Pparα binding sites

independent of Pparα.

• Patterns of Pparα target genes can be distinguished by epigenetic marks at promoters

and enhancers.

ABSTRACT

Fasting requires tight coordination between the metabolism and transcriptional output of hepatocytes to maintain systemic glucose and lipid homeostasis. Genetically-defined deficits in hepatic fatty acid oxidation result in dramatic fasting-induced hepatocyte lipid accumulation and induction of genes for oxidative metabolism, thereby providing a mouse model to interrogate the mechanisms by which the liver senses and transcriptionally responds to fluctuations in lipid levels. While fatty acid oxidation is required for a rise in acetyl-CoA and subsequent lysine acetylation following a fast, changes in histone acetylation (total, H3K9ac, and H3K27ac) associated with transcription do not require fatty acid oxidation. Instead, excess fatty acids prompt induction of lipid catabolic genes largely via ligand-activated Pparα. We observe that active enhancers in fasting mice are enriched for Pparα binding motifs, and that inhibition of hepatic fatty acid oxidation results in elevated enhancer priming and acetylation proximal to Pparα binding sites within regulatory elements largely associated with genes in lipid metabolism. Also, a greater number of Pparα-associated H3K27ac signal changes occur at active enhancers compared to promoters, suggesting a genomic mechanism for Pparα to tune target gene expression levels. Overall, these data demonstrate the requirement for Pparα activation in maintaining transcriptionally permissive hepatic genomic architecture particularly when fatty acid oxidation is limiting.

INTRODUCTION

The liver is a principle regulator of systemic lipid physiology. In this role the hepatocyte requires signaling mechanisms by which to sense and respond to fluctuations in lipid availability. This is especially important during periods of nutrient deprivation, such as fasting (George, 2006). Fasting stimulates fatty acid mobilization from adipose, whereby they are taken up and broken down in the liver via mitochondrial β-oxidation to provide hepatocytes with ATP, NADH, and acetyl-CoA (Stern et al., 2016; The and Schulz, 1991). Errors in these fundamental catabolic

processes result in metabolic aberrations, such as fasting hypoketotic-hypoglycemia, and can

also contribute to the pathogenesis and progression of conditions such as diabetes, obesity,

and chronic liver disease (Asrani et al., 2019; Gong et al., 2017; Houten et al., 2016; Ponziani et

al., 2015).

The nuclear hormone receptor peroxisome proliferator-activated receptor alpha (Pparα)

plays a governing role in regulating hepatic lipid homeostasis and is thus a key component of

the fasting response (Kersten, 2014; Kersten et al., 1999; Leone et al., 1999). Upon activation

by lipid ligands, such as long chain fatty acids, Pparα and its heterodimer partner retinoid X

receptor alpha (RXRα) will bind to DNA and effect transcription of target genes (Bardot et al.,

1993; Boergesen et al., 2012; Evans and Mangelsdorf, 2014; Gearing et al., 1993). This

transcriptional program includes genes for mitochondrial and peroxisomal β-oxidation,

microsomal ω-oxidation, and ketogenesis (Aoyama et al., 1998; Mandard et al., 2004;

Rakhshandehroo et al., 2009). It prompts induction of regulators for mitochondrial metabolism

such as carnitine palmityltransferase 1a (*Cpt1a*) and pyruvate dehydrogenase kinase 4 (*Pdk4*),

which respectively promote mitochondrial fatty acid import and inhibit pyruvate oxidation,

(Huang et al., 2002; Song et al., 2010; Wu et al., 2001). Pparα also activates transcription of

pro-catabolic hepatokines such as fibroblast growth factor 21 (*Fgf21*), which help mediate the

adaptive starvation response systemically (Inagaki et al., 2007; Iroz et al., 2017). Pparα

knockout mice display diminished transcription of these and other genes for oxidative

metabolism, which over time contributes to hepatic lipid accumulation and steatosis (Aoyama et

al., 1998; Kersten et al., 1999; Leone et al., 1999; Reddy, 2001; Ruppert et al., 2019).

Transcription factor binding hubs, such as promoters and enhancer elements, play a

fundamental role in regulating hepatic transcription (Goldstein and Hager, 2015; Jump et al.,

2013; Karagianni and Talianidis, 2015; Qin et al., 2020). Enhancers are cis-regulatory genetic

elements that facilitate transcription initiation and processivity; they are classified as silenced,

poised, or active (Creyghton et al., 2010). Poised enhancers are marked by enrichment of

epigenetic modifications including H3K4me1 and H3K27me3 (Calo and Wysocka, 2013; Heinz

et al., 2015). Active enhancers are marked by a coincidence of H3K4me1 and H3K27ac, as

well as increased chromatin accessibility and the presence of histone acetyltransferases such

as p300 (Andersson and Sandelin, 2020; Creyghton et al., 2010; Raisner et al., 2018). The

hepatic fasting response requires careful balance between gluconeogenesis and ketogenesis.

The interplay between each process is in part governed by transcription factor binding patterns,

including changes in the active enhancer landscape. Glucocorticoid receptor (GR)-assisted

loading of cAMP responsive element binding protein I promotes activation of gluconeogenic

enhancers (Goldstein et al., 2017). In contrast, ketogenic enhancers are proposed to have a

more gradual enhancer maturation which is correlated to GR-stimulated expression of Pparα

and increased chromatin accessibility in proximity to Pparα binding motifs (Goldstein et al.,

2017). Moreover, Pparα binding has been directly detected at fasting-induced enhancers using

ChIP-seq (Lee et al., 2014; Sommars et al., 2019). Pparα-null mice have suppressed indicators

for active enhancers, including decreased enhancer RNA levels and H3K27ac ChIP-seq signal

(Guan et al., 2018; Sommars et al., 2019).

In this study we take advantage of mice with a genetically-defined deficit in hepatic fatty

acid oxidation to further investigate the contribution of Pparα in regulating the hepatic fasting

transcriptional landscape, including a close inspection of its activity at active enhancers.

Carnitine palmityltransferase 2 (Cpt2) is a required enzyme for the transport of long chain fatty

acids into the mitochondria (Houten et al., 2016). Its loss deprives the mitochondria of fatty acid

substrate, and it is thus an obligate enzyme for mitochondrial β-oxidation. During a fast, mice

with a liver-specific deletion of Cpt2 (Cpt2$^{L-/-}$) are unable to clear excess lipid (Lee et al., 2016).

This results in drastic fatty acid accumulation accompanied by augmented transcription of many

Pparα target genes (Lee et al., 2016). These mice are therefore a model of Pparα activation by

build-up of endogenous lipid ligands. We use this system to further explore the effect of hepatic

lipid sensing on gene activation, and the role for Pparα activation in maintaining transcriptionally

permissive chromatin. In particular we describe distinct patterns of Pparα target gene

transcription that are distinguished by epigenetic marks at promoters and enhancers. We

further observe that Pparα-sensitive enhancers are largely associated with lipid metabolism, and

deficits in fatty acid oxidation prompt elevated priming and acetylation at these loci. Altogether

our findings provide genomic insight into how fatty acids alter the epigenome to affect

transcription during periods of nutrient deprivation.

RESULTS

Impaired hepatic mitochondrial fatty acid oxidation elicits putative Pparα target genes

despite a suppression in lysine acetylation.

Mice lacking hepatic Cpt2 (Cpt2$^{L-/-}$) are unable to utilize long chain fatty acids for mitochondrial

β-oxidation. Following a fast, this results in fatty liver and dramatic induction of pro-catabolic

hepatic genes, thereby providing a useful genetic model to interrogate fatty-acid stimulated

transcription (Bowman et al., 2019; Lee et al., 2016). To fully characterize the transcriptional

landscape of the fasting Cpt2$^{L-/-}$ liver, we performed RNA-seq on livers harvested from 9-week

old wildtype (WT) and Cpt2$^{L-/-}$ mice fasted for 24 hours (**Fig. 1a, Table S1**). This expanded

upon our previously published observations that impaired hepatic β-oxidation is associated with

augmented fasting-induced transcription of genes for regulation of mitochondrial metabolism

(*Pdk4*), regulation of peroxisomal metabolism (*Ehhadh*), and pro-catabolic hepatokines (*Fgf21*)

(Lee et al., 2016). Indeed, gene ontology for all significantly upregulated transcripts (Cpt2$^{L-/-}$/WT

fold change ≥ 2, padj < 0.05) returned several terms for fatty acid metabolism (**Fig. S1a**). We

next performed TMT-based quantitative proteomics to confirm the induction of these genes at

the protein level (**Table S2**). We found that, in parallel to the RNA-seq data, Cpt2$^{L-/-}$ upregulated

peptides were enriched for gene ontology terms related to fatty acid biology (**Fig. S1b, S1c**).

These results provide additional evidence demonstrating that impaired β-oxidation triggers a

Fig 1. Fasted Cpt2$^{L-/-}$ liver exhibits deficits in lysine-acetylation along with metabolic gene induction.

  A. Volcano plot displaying differentially expressed transcripts (fold change ≥ |2|, padj < 0.05) between 24hr fasted WT and Cpt2$^{L-/-}$ liver as determined by RNA-seq (n=4).

  B. Tissue acetyl-CoA concentration from 24hr fasted WT and Cpt2$^{L-/-}$ liver (n=6, mean ± SEM). Significance determined using Student two-tailed t test, ***p < 0.001.

  C. Western blot for acetyl-lysine and HSC70 loading control in fed and fasted WT and Cpt2$^{L-/-}$ livers.

  D. Volcano plots showing magnitude and significance for fasted liver acetyl-proteome, measured by TMT-based quantitative mass spectrometry (n=5).

  E. Volcano plots showing magnitude and significance for mitochondrial acetyl-proteome and nuclear acetyl-proteome.

  F. Representative Western blot of acid-precipitated histones from fed and fasted WT and Cpt2$^{L-/-}$ liver tissue.

(legend continued on next page)

Fig 1. Fasted Cpt2$^{L-/-}$ liver exhibits deficits in lysine-acetylation along with metabolic gene induction.
  G. Aggregation plot depicting liver H3K9ac ChIP-seq mean tag density for (*left*) all H3K9ac ChIP-seq peak centers with ±5kb flanking regions and (*right*) across all gene bodies with ±2kb flanking regions. TSS = transcription start site, TTS = transcription termination site. n=1
  H. H3K9ac ChIP-seq genome browser tracks for representative genes that are upregulated in Cpt2$^{L-/-}$ RNA-seq, showing increased ChIP-seq signal in the fasted Cpt2$^{L-/-}$ liver.

compensatory transcriptional program in an attempt to relieve the lipid burden, a process which

includes shuttling fatty acids into other metabolic pathways such as peroxisomal oxidation.

Others have suggested that fatty acid oxidation promotes transcription for lipid

catabolism via histone acetylation from fatty acid-derived acetyl-CoA (McDonnell et al., 2016).

We observed that Cpt2$^{L-/-}$ mice exhibited suppressed hepatic acetyl-CoA concentration and

failed to induce protein acetylation following a fast, which therefore allowed us to examine this

hypothesis *in vivo* (**Fig. 1b, 1c**). To gain a more granular view of the proteins that exhibited

hypoacetylation, we mapped and quantified protein acetylation via TMT-based mass

spectrometry (**Fig. 1d, Table S3**). We found that mitochondrial protein acetylation was

suppressed in Cpt2$^{L-/-}$ mice as expected due to a lack of mitochondrial β-oxidation (**Fig. 1e,**

S1e). In fact, lysine acetylation was globally reduced in Cpt2$^{L-/-}$ mice with the noted exception of

nuclear peptides, which retained comparable levels of acetylation between WT and Cpt2$^{L-/-}$ mice

(**Fig. 1e, S1d**). These data indicate that the deficit in mitochondrial β-oxidation limits liver

acetyl-CoA substrate for lysine acetylation, with the exception of the nucleus where acetyl-lysine

levels appear buffered.

Analysis of acetyl-histone peptides provides evidence against bulk histone acetylation as

a primary driver of transcriptional activation. 91 unique acetyl-histone peptides were detected in

the acetyl-proteome, of which none were hyperacetylated, 38.5% were hypoacetylated and the

remaining 61.5% showed no significant change to acetylation state between Cpt2$^{L-/-}$ and WT

animals (**Table S3**). Consistent with those data, bulk changes in histone acetylation were not

observed for histone marks H3K9ac or H3K14ac (**Fig. 1f**). To gain detailed insight into fatty

acid oxidation-dependent histone acetylation, we next turned to ChIP-seq for H3K9ac. Global

H3K9ac levels were higher in Cpt2$^{L-/-}$ liver, including at gene promoters, consistent with the

augmented transcriptional profile of Cpt2$^{L-/-}$ mice (**Fig. 1g**). Local patterns of H3K9ac gene body

occupancy trended with RNA-seq fold-change (**Fig. 1h, S1f**). These data are not consistent

with the notion that the fatty acid oxidation-dependent generation of acetyl-CoA is a primary

driver of gene expression via an epigenetic mechanism, but rather point towards another lipid

sensing mechanism for transcriptional activation.

To better understand how fatty acids affect the chromatin landscape, including promotor

accessibility and binding motifs for transcription factors induced in fasting Cpt2$^{L-/-}$ liver, we next

turned to Assay for Transposase Accessible Chromatin Sequencing (ATAC-seq). ATAC-seq

profiles of fasted Cpt2$^{L-/-}$ and WT liver tissues were overall similar. A total of 69,485 ATAC-seq

sites were measured across all samples, of which 75.6% are shared by both fasted Cpt2$^{L-/-}$ and

WT animals (**Fig. 2a**). Cpt2$^{L-/-}$ livers had a notable increase in chromatin accessibility near

transcription start sites (**Fig. 2b).** Of the 9273 differentially upregulated ATAC sites (fold change

Cpt2$^{L-/-}$/WT ≥ +2), 16% were detected in gene promotors. Differential accessibility aligned with

RNA-seq trends (**Fig. 2c, S2a**). Motif analysis on ATAC peaks found at gene promotors for

upregulated Cpt2$^{L-/-}$ transcripts revealed Pparα and its heterodimer binding partner Rxr as the

top hits (**Fig. 2d**). Pparα targets are often defined by their response to a Pparα agonist, such as

WY-14643 (Brocker et al., 2020; Janssen et al., 2015; Lee et al., 1995; Li et al., 2018;

Rakhshandehroo et al., 2007). We determined that 52% of genes upregulated in Cpt2$^{L-/-}$ animals

are sensitive to WY-14643 (**Fig. 2e**) (Naiman et al., 2019). Together, the proteomic, acetyl-

proteomic, RNA-seq, ATAC-seq and H3K9ac ChIP-seq data suggest that deficits in hepatic fatty

acid oxidation by the loss of Cpt2 elevates endogenous lipid ligands to induce Pparα at

promoters of genes important for lipid catabolism despite a suppression in acetyl-CoA.

Pparα, Cpt2$^{L-/-}$ double knockout mice demonstrate Pparα dependent and independent

transcription.

Given the abundance of putative Pparα target genes and enrichment for Pparα promoter

binding motifs in fasted Cpt2$^{L-/-}$ livers, we decided to define the requirement for Pparα in

Fig 2. Fasting-induced genes in Cpt2$^{L-/-}$ liver are Pparα-responsive

A. Venn diagram showing unique and shared ATAC-seq peaks detected in fasted WT and Cpt2$^{L-/-}$ liver (n=2).

B. Aggregation plot depicting liver ATAC-seq mean tag density for all gene bodies with ±2kb flanking regions. TSS = transcription start site, TTS = transcription termination site.

C. ATAC-seq genome browser tracks for representative genes that were upregulated in Cpt2$^{L-/-}$ RNA-seq.

D. Top results from HOMER enrichment analysis for known motifs in ATAC-seq promoter peaks for genes induced in Cpt2$^{L-/-}$ liver.

E. Pie chart depicting Cpt2$^{L-/-}$ induced genes previously shown to respond to the Pparα agonist WY-14643 (GSE140063, Naiman et al., 2019) .

the transcriptional response of Cpt2$^{L-/-}$ mice by generating double knockout (DKO) mice.

Cpt2$^{L-/-}$;Ppar$\alpha^{-/-}$ DKO animals were generated using mice harboring a whole-body deletion of

Pparα (Ppar$\alpha^{-/-}$) bred to Cpt2$^{L-/-}$ mice. Subsequent experiments in this study were conducted

using 24hr fasted 9-week old WT , Cpt2$^{L-/-}$, Ppar$\alpha^{-/-}$, and DKO mice. All genotypes were viable

and fertile and were not associated with changes in body weight (**Fig. 3a**). We have previously

shown that fasted Cpt2$^{L-/-}$ mice have significantly increased liver mass due to excess fatty acids;

the DKO mice share this phenotype (**Fig. 3b**). While Cpt2$^{L-/-}$ mice did not exhibit differences in

fasting glycemia compared to controls, Ppar$\alpha^{-/-}$ and DKO mice exhibited mild fasting

hypoglycemia (**Fig. 3c**). This phenotype was previously described for fasted Ppar$\alpha^{-/-}$ animals

(Kersten et al., 1999). Ppar$\alpha^{-/-}$ mice had approximately half of the circulating fasting ketone

bodies such as β-hydroxybutyrate (βHB) compared to WT animals (**Fig. 3d**). However, Cpt2$^{L-/-}$

and DKO mice did not produce βHB following a 24hr fast, demonstrating a requirement for

hepatic fatty acid oxidation for ketone body generation.

To better understand the physiological impact of impaired hepatic β-oxidation and Pparα

loss, we performed global untargeted metabolomics on serum (**Fig. 3e, Table S4**). Principle

component analysis (PCA) showed that loss of Pparα caused broad perturbations to the serum

metabolome, more so than loss of Cpt2$^{L-/-}$ alone (**Fig. 3e, S3a**). This emphasizes the

importance of Pparα for systemic fasting physiology. These perturbations were compounded in

the DKO animals, which showed the greatest variation from WT controls. We've previously

reported that fasted Cpt2$^{L-/-}$ mice show depleted serum short- and medium-chain acyl-carnitines

(Lee et al., 2016). DKO mice phenocopy Cpt2$^{L-/-}$ serum levels of medium- and long-chain acyl

carnitines, indicating that this is likely a shared metabolic sink for the excess fatty acid in Cpt2$^{L-/-}$

livers (**Fig. S3b**). These data overall demonstrate that the loss of Pparα and/or the loss of

hepatic fatty acid oxidation result in systemic metabolic perturbations.

Fig 3. Pparα, Cpt2$^{L-/-}$ double knockout mice retain transcription of Pparα target genes

A. Body weight of male fasted WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and Cpt2$^{L-/-}$;Pparα$^{-/-}$ (DKO) animals (n=6-10 mean ± SEM).
B. Wet liver weight from fasted animals (n=6-10, mean ± SEM).
C. Blood glucose from fasted animals (n=6-10, mean ± SEM).
D. Serum levels of β-hydroxybutyrate from fasted animals determined by median-scaled untargeted metabolomics (n=6, mean ± SEM).
E. Heatmap of top 100 differentially regulated serum metabolites from fasted animals (n=6).
F. Principle component analysis of RNA-seq on fasted WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and DKO liver (n=4).
G. Volcano plot displaying RNA-seq differentially expressed transcripts between 24hr fasted WT and Pparα$^{-/-}$ livers.
H. Volcano plot displaying RNA-seq differentially expressed transcripts between 24hr fasted WT and DKO livers.
I. (*left*) Venn diagram depicting overlap between genes upregulated in Cpt2$^{L-/-}$and DKO liver compared to WT. (*right*) Pie chart depicting DKO fasting-induced genes known to respond to the Pparα agonist WY14643.

RNA-seq significance cutoff is fold change ≥ |2|, padj < 0.05. One-way ANOVA followed by Tukey's post-hoc test was performed as appropriate. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001; ns, not significant.

We next sought to determine the extent to which Pparα is implicated in the fasting-

induced Cpt2$^{L-/-}$ transcriptional response by carrying out RNA-seq on WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and

DKO livers (**Table S1**). PCA of RNA-seq data revealed that the transcriptional signature for

Pparα$^{-/-}$ mice overlapped with that of DKO mice, while those for WT and Cpt2$^{L-/-}$ animals were

distinct from the other genotypes (**Fig. 3f**). There were 330 differentially expressed transcripts

shared between the three knockout genotypes compared to WT animals (KO/WT fold change ≥

|2|, padj < 0.05) (**Fig. S3c**). In line with overall transcriptional signatures, a greater number of

differentially expressed genes were shared between the Pparα$^{-/-}$ and DKO mice compared to

Cpt2$^{L-/-}$ mice. These findings are not unexpected given the central role for Pparα in regulating

the hepatic genes necessary for fasting oxidative metabolism.

The loss of Pparα alone suppressed transcription of genes for fatty acid catabolism and

ketogenesis, including *Ehhadh*, *Hmgcs2*, and *Fgf21* (**Fig. 3g**). Many of these genes displayed

blunted expression in DKO livers as well (**Fig. 3h**). Curiously, several canonical Pparα target

genes, such as *Plin2* and *Pdk4*, retained induction in DKO liver compared to control animals

(**Fig. 3h**). Further analysis of the RNA-seq data revealed that Cpt2$^{L-/-}$ and DKO animals shared

257 upregulated transcripts, respectively representing 42% and 40% of induced genes for those

genotypes (**Fig. 3i**). This suggested that there may be a transcriptional program triggered by

the shared metabolic perturbations presented by loss of hepatic β-oxidation. Further, 49% of

induced transcripts in DKO are responsive to WY-14643 (**Fig. 3i**) (Naiman et al., 2019). We

therefore hypothesized that a subset of genes, which are typically thought to rely upon Pparα for

their induction, have in addition a secondary regulatory mechanism. The limited transcription of

target genes that do not require Pparα is likely prompted by excess lipid burden via an

independent mechanism driven by Err or Hnf4α, for example.

Fig 4. Unbiased clustering reveals patterns for Pparα-dependent and Pparα-independent transcription of target genes

A. *(left)* Tracing diagram of cluster 1 genes (Pparα-dependent) from fasted RNA-seq transcript levels. Blue line indicates mean expression. *(right)* Bar graph quantifying scaled transcript values. Presented as Z-score scaled transcript levels.

B. Normalized transcript read counts for select cluster 1 genes.

C. Gene ontology for cluster 1 genes ranked by significance.

D. *(left)* Tracing diagram of cluster 3 genes (Pparα-independent) from fasted RNA-seq transcript levels. Blue line indicates mean expression. *(right)* Bar graph quantifying scaled transcript values. Presented as Z-score scaled transcript levels.

E. Normalized transcript read counts for select cluster 3 genes.

F. Gene ontology for cluster 3 genes ranked by significance.

One-way ANOVA followed by Tukey's post-hoc test was performed as appropriate. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$; ns, not significant. Bar graphs represent 2%-98% percentile.

Impaired hepatic fatty acid oxidation prompts distinct modes of gene regulation.

We subsequently sought further resolution into the role for Pparα in regulating hepatic

transcription in response to impaired hepatic fatty acid oxidation, including the possibility of

Pparα-independent regulatory mechanisms. Unbiased k-means clustering was used to identify

unique patterns of gene transcription across all differentially expressed transcripts (**Fig. S4a**,

Table S5). Of the five clusters identified, two in particular (1 & 3) were associated with the

Cpt2$^{L-/-}$ gene induction signature (**Fig. 4a, 4d**). Notably, these two groups of transcripts display

significantly divergent modes of Pparα-sensitive transcription.

Cluster 1 genes were wholly dependent upon Pparα for their transcription, as indicated

by their repressed expression in Pparα$^{-/-}$ and DKO fasted liver (**Fig. 4a**). Cluster 1 genes

included multiple canonical Pparα targets, including *Cyp4a10*, *Ehhadh*, and *Hmgcs2* (**Fig. 4b,**

S3b). These genes are respectively implicated in Pparα regulation of ω-oxidation, peroxisomal

oxidation, and ketogenesis (Kersten et al., 2010). Indeed, gene ontology and KEGG pathway

analysis on all Cluster 1 genes returned terms related to fatty acid metabolism and Pparα

signaling (**Fig. 4c**). These data provide further evidence that impaired hepatic fatty acid

oxidation activates Pparα-dependent transcription upon fasting.

Cluster 3 genes were likewise induced in Cpt2$^{L-/-}$ mice in a Pparα-dependent manner.

However, this group differs from the previous cluster in that both Pparα-null backgrounds

retained baseline expression levels (**Fig. 4d**). Cluster 3 genes thus sustained transcriptional

activity in the absence of Pparα, yet still required the transcription factor for induction in

response to metabolic perturbations such as a buildup of Pparα ligands. This group included

canonical Pparα target genes *Acot2* and *Pdk4*, whose expression in Pparα-null animals

displayed no significant variation from control animals (**Fig. 4e**). KEGG pathway analysis

returned Ppar Signaling Pathway as a top result (**Fig. 4f**). Curiously, Cluster 3 contained

multiple genes associated with lipid droplets, including four members of the perilipin family

(*Plin1-4*) and *Cidec* (Greenberg et al., 2011). *Plin2*, *Plin4*, and *Cidec* are known Pparα targets

(Francque et al., 2015; Kersten et al., 2010). This suggests that modules of the Pparα

transcription program involve building upon pre-established Pparα-independent gene

expression patterns to respond to specific hepatocyte metabolic states.

Distinct Pparα target genes retain promotor accessibility despite loss of Pparα.

We next asked how these unique modes of Pparα target gene transcription were reflected at the

level of chromatin architecture by assessing ATAC-seq from all four genotypes. Loss of Pparα

drastically diminishes chromatin accessibility (**Fig. S5a**). Chromatin landscape remodeling for

the Pparα-null mice was expected given the transcription factor's crucial role in coordinating

fasting metabolism, yet the severe ATAC-seq signal depletion in gene promoter regions was

particularly striking (**Fig. 5a**). PCA emphasized this disparity, showing that Pparα and DKO

animals clustered uniquely, while WT and Cpt2$^{L-/-}$ livers overlapped substantially (**Fig. S5b**). We

further observed there are appreciably fewer changes in discrete peak intensities between

Pparα$^{-/-}$ and DKO animals compared to Cpt2$^{L-/-}$ and DKO animals (**Fig. 5b**). Together these data

indicate that Pparα is required for proper upkeep of fasting chromatin dynamics.

Cluster 1 genes, those dependent upon Pparα for their expression, exhibited a common

pattern of promotor chromatin accessibility that is represented by *Fgf21* (**Fig. 5c**). ATAC-seq

genome browser tracks indicated a higher promoter peak signal in the Cpt2$^{L-/-}$ livers compared

to WT; accessibility was lost under a Pparα-null background. Indeed, motif analysis on ATAC-

seq promotor peaks for this cluster returned Pparα and its heterodimer partner Rxrα as the top

two hits (**Fig 5d**). Suppressed chromatin accessibility at loci enriched for Pparα binding motifs

indicates the need for Pparα for proper promoter dynamics.

Fig 5. Loss of Pparα remodels genome-wide accessibility

A. Aggregation plot depicting liver ATAC-seq mean tag density for all gene bodies with ±2kb flanking regions in fasted WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and DKO liver (n=2). TSS = transcription start site, TTS = transcription termination site.

B. MA plot depicting ATAC-seq differential binding analysis results between DKO/Cpt2$^{L-/-}$ and DKO/Pparα$^{-/-}$ animals. The x-axis gives the mean signal and the y-axis gives the signal fold change for a given peak. Points in pink are significant (fold change ≥ |2|, FDR < 0.05).

C. ATAC-seq genome browser tracks for cluster 1 gene *Fgf21*.

D. HOMER enrichment analysis for known motifs in ATAC-seq promoter peaks for cluster 1 genes.

E. ATAC-seq genome browser tracks for cluster 3 gene *Pdk4*. Pparα-sensitive peak interval is indicated by green bar.

F. HOMER enrichment analysis for known motifs in ATAC-seq promoter peaks for cluster 3 genes.

Promoter accessibility for Cluster 3 genes, those with Pparα-independent baseline

expression, likewise shared a common promoter chromatin architecture as illustrated by *Pdk4*

(**Fig. 5e**). The Cpt2$^{L-/-}$ induction pattern was once again evident in the ATAC-seq signal.

However, in contrast to Cluster 1 genes, this group retained a degree of promotor accessibility

in Pparα$^{-/-}$ and DKO mice. Interestingly, 13% of Cluster 3 genes contained an interval of open

promotor chromatin that was lost in both Pparα-null animals, including at the promoters for

Pparα targets *Acot2*, *Fabp3*, *Pdk4*, and *Plin2* (**Fig. 5e** green bar). These likely indicate binding

sites for Pparα directly or for another DNA regulatory protein that depends on Pparα for an

aspect of its activity. Cluster 3 ATAC-seq promoter peaks are enriched for Pparα and Rxrα

binding motifs, albeit at a lower rate than for Cluster 1 genes (**Fig. 5f**). These data together

indicate that Cluster 3 promoters do not require Pparα to maintain accessibility during a fast,

though Pparα can interact with these loci to augment target gene transcription in response to a

metabolic perturbation.

Pparα-dependent and independent promotor acetylation.

Differential promotor accessibility prompted questions as to how the loss of Pparα affected

promotor dynamics *vis a vis* histone acetylation. We therefore performed ChIP-seq for

H3K27ac, a histone mark generally found at active chromatin (Andersson and Sandelin, 2020).

H3K27ac signal was comparable between genotypes, however Cpt2$^{L-/-}$ and DKO livers

displayed higher H3K27ac signal near transcription start sites compared to the WT and Pparα$^{-/-}$

animals (**Fig. 6a, S6a**). PCA showed that WT and Cpt2$^{L-/-}$ H3K27ac distributions were distinct

from one another, while Pparα$^{-/-}$ and DKO animals clustered together (**Fig. 6b**). This is the

inverse of the ATAC-seq PCA, in which WT and Cpt2$^{L-/-}$ animals were grouped and Pparα$^{-/-}$ and

DKO animals were separated (**Fig. S5b**). These data indicate that the metabolic perturbations

in Cpt2$^{L-/-}$ and DKO livers stimulate a common configuration of H3K27ac occupancy regardless

Fig 6. Differential H3K27ac occupancy patterns at Pparα target genes

A. Aggregation plot depicting liver H3K27ac ChIP-seq mean tag density for all gene bodies with ±2kb flanking regions in fasted WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and DKO liver (n=2). TSS = transcription start site, TTS = transcription termination site.

B. PCA of H3K27ac ChIP-seq data from fasted WT, Cpt2$^{L-/-}$, Pparα$^{-/-}$, and DKO liver.

C. Genome browser track for cluster 1 genes *Fgf21* and *Cyp4a10*. ATAC-seq is presented in red, H3K27ac ChIP-seq is presented in purple. Green bar indicates downstream regulatory region.

D. (*left*) Aggregation plot for H3K27ac ChIP-seq mean tag density for cluster 1 gene promoters. (*right*) Bar graph quantifying mean values of H3K27ac ChIP-seq peaks found in cluster 1 gene promoters.

E. Genome browser track for cluster 3 genes *Pdk4* and *Plin2*. ATAC-seq is presented in red, H3K27ac ChIP-seq is presented in purple.

F. (*left*) Aggregation plot for H3K27ac ChIP-seq mean tag density for cluster 3 gene promoters. (*right*) Bar graph quantifying mean values of H3K27ac ChIP-seq peaks found in cluster 3 gene promoters.

Significance determined by Kruskal-Wallis test with Dunn's post hoc correction. *p < 0.0001; ns, not significant. Bar graphs represent 2%-98% percentile.

of overall chromatin accessibility. This further demonstrates lipid-driven selective histone

acetylation despite absence of mitochondrial β-oxidation. Pparα is not required for this

H3K27ac deposition, however Pparα is still needed to promote chromatin accessibility and

ultimately transcription of lipid catabolic genes. Comparing the WT and Cpt2$^{L-/-}$ ATAC-seq and

H3K27ac ChIP-seq peaksets provides additional evidence that the Cpt2$^{L-/-}$ epigenetic program

builds upon the baseline fasting chromatin architecture to exaggerate an established response.

Clusters 1 and 3, together comprising the genes induced in Cpt2$^{L-/-}$ liver, provided a

useful contrast to understand variations in active chromatin acetylation and how they relate to

Pparα. H3K27ac occupancy at Cluster 1 gene promotors paralleled ATAC-seq trends in that

loss of Pparα was associated with significantly decreased promoter H3K27ac tag density, as

indicated by *Fgf21* and *Cyp4a10* (**Fig. 6c**). There was no statistical difference in H3K27ac

signal between WT and Cpt2$^{L-/-}$ animals (**Fig. 6d**). At Cluster 3 gene promoters, in contrast to

Cluster 1, Pparα loss was not associated with depleted promoter H3K27ac signal. H3K27ac

occupancy was instead distinctly independent of Pparα, as indicated by its canonical target

genes *Pdk4* and *Plin2* (**Fig 6e**). Moreover, the Cpt2$^{L-/-}$ background is associated with

significantly elevated H3K27ac promoter tag density compared to WT and Pparα$^{-/-}$ animals (**Fig.**

6f). These observations continue to demonstrate that the lipid-associated changes in chromatin

accessibility are linked to site-specific epigenetic modifications in gene promoters.

At Cluster 1 gene *Fgf21* we noticed an acetylation pattern indicative of a regulatory

region downstream from the transcription termination site (**Fig. 6c**, green bar). The H3K27ac

signal at this site was comparable between the four genotypes, yet accessibility was only

observed in the WT and Cpt2$^{L-/-}$ animals. Both ATAC-seq peaks contained Pparα binding

motifs. This chromatin pattern parallels that observed at cluster 3 gene promoters. We

speculate that at these loci Pparα is able to detect antecedent chromatin acetylation as part of

its DNA binding mechanism, or that an epigenetic reader detects that mark and subsequently

loads Pparα onto chromatin, thereby promoting a transcriptionally permissive state.

Loss of Pparα selectively alters the hepatic fasting enhancer landscape.

Fasting stimulates a cascade of transcription factor loading and binding onto hepatic enhancers

(Goldstein et al., 2017). Enhancers can be categorized as either poised or active. The former

are indicated by the histone modification H3K4me1, and the latter are defined by co-occupancy

of H3K4me1 and H3K27ac (Creyghton et al., 2010). We decided to use the Cpt2$^{L-/-}$ model as a

handle by which to investigate lipid-responsive enhancer elements, and the extent to which

Pparα regulates both their poising and activation. A total of 8998 active enhancers were

detected across all genotypes (**Fig. 7a, Table S6**). Strikingly, Pparα-null animals had

diminished chromatin accessibility over active enhancer elements (**Fig. S7a**). Loss of either

Cpt2 or Pparα alone did not impact H3K4me1 density at active enhancers, but their combined

deletion in DKO liver reduced global enhancer priming (**Fig. 7a**). Enhancer H3K27ac signal did

not differ between Pparα$^{-/-}$ and WT animals. However, loss of hepatic Cpt2 was associated with

overall decreased H3K27ac occupancy at active enhancers. This was an unanticipated contrast

to increased acetylation at gene promoters under the Cpt2$^{L-/-}$ background (**Fig. 1g, 6a**). Overall

these data imply that while loss of Pparα alone does not negatively impact the enhancer

epigenetic landscape during the latter stages of a fast, global enhancer accessibility is

significantly hindered without the transcription factor.

349   Active enhancers were highly enriched for Hnf4α and Pparα binding motifs (**Fig. 7b**).

350   Surprisingly, Pparα ChIP-seq sites located within enhancers showed elevated H3K4me1 and

351   H3K27ac signals in all knockout genotypes (**Fig. 7c**) (Sommars et al., 2019). There was no

352   difference in enhancer H3K4me1 signal between Cpt2$^{L-/-}$ and Pparα$^{-/-}$ liver, though loss of both

353   genes in DKO animals caused significantly increased methylation near Pparα binding sites.

Fig 7. Lipid-sensitive enhancer elements are primed for Pparα.

A. Bar graph depicting H3K4me1 and H3K27ac tag counts at active enhancer elements.

B. HOMER motif analysis for all active enhancers as defined by H3K4me1 and H3K27ac co-occupancy.

C. (*top*) Bar graphs depicting H3K4me1 and H3K27ac mean tag density proximal to Pparα ChIP-seq binding sites in all active enhancers (±500bp from Pparα ChIP peak center). (*bottom*) Aggregation plot for H3K4me1 and H3K27ac tag density within 1kb flanking regions of Pparα ChIP peak center. (Pparα ChIP-seq: GSE118788, Sommars et al., 2019)

D. (*top*) Pie chart showing genomic distribution (promoters, enhancers, other) of H3K4me1 and H3K27ac differential peak analysis at enhancers with repressed H3K27ac signal in Pparα liver compared to WT. Promoters were not considered for H3K4me1 ChIP-seq. (*bottom*) Percent overlap between peak genomic annotation and Pparα ChIP-seq coordinates.

(legend continued on next page)

E. Aggregation plots for H3K4me1 and H3K27ac tag density within 1kb flanking regions of Pparα ChIP peak center at enhancers with repressed H3K27ac signal in Pparα liver.

F. H3K27ac GREAT ontology for enhancers with repressed H3K27ac signal in Pparα liver.

G. (*top*) Pie chart showing genomic distribution of H3K4me1 and H3K27ac differential peak analysis for enhancers with increased H3K27ac signal in Cpt2$^{L-/-}$ liver compared to WT. (*bottom*) Percent overlap between peak genomic annotation and Pparα ChIP-seq coordinates.

H. Heatmap of normalized H3K27ac ChIP-seq (purple), H3K4me1 ChIP-seq (green), and ATAC-seq (red) signal at active enhancers classified according to DKO/Cpt2$^{L-/-}$ differential peak analysis. Coverage is calculated from peak center with ±5kb flanking regions.

I. (*top*) Pie chart showing genomic distribution of H3K27ac differential peak analysis for enhancers with increased, repressed, and unchanged H3K27ac signal in DKO liver compared to Cpt2$^{L-/-}$. (*bottom*) Percent overlap between peak genomic annotation and Pparα ChIP-seq coordinates.

J. H3K27ac GREAT ontology for Class B enhancers (DKO < Cpt2$^{L-/-}$).

K. Bar graphs quantifying normalized H3K27ac and H3K4me1 ChIP-seq signal at Class A and Class B active enhancers

Significance determined by Kruskal-Wallis test with Dunn's post hoc correction. *p < 0.0001; ns, not significant. Bar graphs represent 2%-98% percentile.

H3K27ac tag density flanking Pparα binding sites was increased in Cpt2$^{L-/-}$ animals compared to

Pparα$^{-/-}$. Cpt2$^{L-/-}$ animals also had increased ATAC-seq signal over Pparα enhancer sites

compared to WT animals (**Fig. S7b**). As might be expected, chromatin accessibility over Pparα

sites was decreased in Pparα-null livers, though this was despite increased enhancer priming

and acetylation over WT baselines at those loci (**Fig. 7c, S7b**). This indicates that the active

epigenetic state is maintained independent of Pparα at these enhancers, but Pparα is required

to promote chromatin accessibility. Together these data show that perturbations to fatty acid

catabolism are associated with amplified enhancer priming and acetylation specifically near

Pparα binding sites even in the absence of the transcription factor. We propose this is an

epigenetic mechanism for the hepatocyte to prime a genomic response to impaired lipid

catabolism.

We next examined the specific changes to enhancer state at loci that are differentially

regulated in Pparα$^{-/-}$ liver compared to WT. Loss of Pparα was associated with targeted

repression of H3K27ac signal, and to a lesser extent H3K4me1 signal (**Fig. 7d**). Only 169

enhancers had suppressed priming in Pparα$^{-/-}$ animals in contrast to the 612 enhancers with

suppressed H3K27ac signal in Pparα$^{-/-}$ animals. As expected, these downregulated regions

contained a high frequency of Pparα ChIP-seq binding sites. Curiously, 33% of H3K27ac peaks

repressed in Ppar$\alpha^{-/-}$ liver were located within enhancer elements, while only 19% of H3K27ac

signal changes were found in gene promoters (**Fig. 7d**). In other words, loss of Pparα

diminished chromatin acetylation at a greater number of enhancers than promoters. Enhancers

with blunted acetylation in Ppar$\alpha^{-/-}$ liver displayed highly enriched H3K4me1 and H3K27ac

signals in Cpt2$^{L-/-}$ animals (**Fig. 7e**). The Genomic Regions Enrichment of Annotations Tool

(GREAT), which provides ontologies for enhancer function using a binomial test specific for

long-range genomic regulatory domains, showed that enhancers repressed by loss of Pparα

were associated with ontology terms for fatty acid metabolism (**Fig. 7f**) (McLean et al., 2010).

These data are consistent with a model in which Cpt2 loss drives Pparα signaling, and that

Pparα target gene transcription is associated with heightened activation of enhancers

associated with lipid metabolism.

Following up on these observations, we next examined how a deficit in β-oxidation

affected the hepatic enhancer landscape, and the extent to which Pparα is implicated in

genomic maintenance under these metabolic conditions. Loss of Cpt2 had minimal effects on

overall enhancer priming (**Fig. 7g**). In contrast, Cpt2$^{L-/-}$ mice had 915 enhancer regions with

increased H3K27ac signal. 55% of these regions contain Pparα binding sites. Indeed, Cpt2$^{L-/-}$

animals had a prominent increase in both enhancer priming and acetylation over Pparα ChIP

sites (**Fig. S7c**). At these Cpt2$^{L-/-}$-sensitive loci we also observed increased H3K4me1 and

H3K27ac ChIP signals over Pparα binding sites in Pparα-null animals compared to WT controls,

indicating that enhancers with epigenetic sensitivity to impaired β-oxidation likewise respond to

impaired Pparα signaling, perhaps designating a response to drive augmented transcription

such as we see in the Cpt2$^{L-/-}$ mice (**Fig. S7c**).

The DKO mice clarify the role for Pparα at these enhancer elements. These mice had a

striking change to their active enhancer landscape when compared to their Cpt2$^{L-/-}$counterparts.

We initially noticed there were far fewer changes in enhancer priming compared to changes in

enhancer acetylation (**Fig. 7i, S7d**). A total of 1487 differentially regulated enhancers were

detected between the two genotypes, which were broken down into three groups according to

differential H3K27ac ChIP signal (**Fig. 7h**). Class B enhancers (n=876) were defined by

statistically significant H3K27ac depletion in DKO mice compared to Cpt2$^{L-/-}$ (**Fig. 7h, 7k**). We

consider these to be the enhancer elements at which the ligand-activated Pparα signaling

program specifically impacts acetylation in response to lipid sensing. The high incidence of

Pparα binding sites within these enhancers suggests that Pparα itself plays a direct role in local

H3K27ac deposition (**Fig. 7i**). Cpt2$^{L-/-}$ animals had increased H3K27ac signal at these regions

compared to WT (**Fig. 7k**). We further noted that loss of Pparα on the Cpt2$^{L-/-}$ background had

an outsized effect on enhancer acetylation compared to promoter acetylation (**Fig. 7i**). Finally,

GREAT analysis indicates that proximal genes for Class B enhancers were associated with

multiple lipid gene ontology terms (**Fig. 7j**). Together these data demonstrate the substantial

role for enhancer elements in the Pparα transcriptional response, and that lipid-amplified Pparα

signaling is perhaps mediated more substantially by epigenetic changes to enhancer regions

than at promoters.

Class A enhancers (n=611) were defined by a greater than two-fold increase in H3K27ac

signal in DKO mice compared to Cpt2$^{L-/-}$ liver. DKO Class A enhancer acetylation was similarly

increased over WT animals (**Fig. 7k**). This augmented H3K27ac signal was shared by Pparα$^{-/-}$

animals. Likewise, both Pparα$^{-/-}$ and DKO animals displayed elevated H3K4me1 priming levels

at Class A enhancers (**Fig. 7k**). This implies Pparα loss is associated with enhancer activation

that may be linked to other metabolic processes. Class C enhancers (n=7511), which showed

no change in acetylation state between Cpt2$^{L-/-}$ and DKO animals, represent 83% of the active

enhancer landscape (**Fig. S7e**), emphasizing that the genomic perturbations presented by loss

of Cpt2 or Pparα result in specific, targeted changes to enhancer epigenetic state.

Altogether these data demonstrate that metabolic perturbations such as inhibiting

hepatic fatty acid oxidation or loss of Pparα result in higher levels of enhancer priming and

acetylation proximal to Pparα sites that are largely associated with lipid metabolism. Pparα

DNA binding is proceeded by enhancer priming and acetylation. This is clearly illustrated at

Pparα-bound enhancers found near *Fgf21* and *Pdk4* (**Fig. S7f**). The acetylation may even be

the signal for Pparα to engage these regulatory elements. Furthermore, while changes to

enhancer epigenetic state are more localized near Pparα binding sites, the transcription factor is

so crucial to maintaining permissive chromatin that loss of Pparα repressed chromatin

accessibility across the global enhancer network. We also surmise that changes to the

enhancer landscape comprise a significant component of ligand-activated Pparα signaling, with

quantitatively more Pparα-associated changes in acetylation occurring at enhancers compared

to promoters. Overall, these data show the requirement for Pparα activation in maintaining

transcriptionally permissive hepatic genomic architecture, particularly when the hepatocyte must

sense and respond to elevated lipid.

DISCUSSION

A foundational question in liver metabolism is how the hepatocyte senses and responds to

fluctuations in lipid availability. This ultimately requires tight coordination between the

hepatocyte's metabolic state, genome architecture, and transcriptional output. Fasting Cpt2$^{L-/-}$

mice prompts accumulation of fatty acids, including putative endogenous Pparα ligands (Lee et

al., 2016; Selen et al., 2021). Using this mouse genetic model and a combination of next-

generation sequencing platforms we clarify the role for hepatic lipid content in metabolic gene

transcription, and in particular how Pparα, activated by natural ligands, maintains the

transcriptional architecture necessary for target gene expression (**Fig. 8**).

Fig 8. Pparα is required for lipid-responsive enhancer and promoter activation
Deficit in mitochondrial β-oxidation and/or excess fatty acids prompts H3K4me1 and H3K27ac deposition at lipid-responsive enhancer elements. This signals ligand-activated Pparα recruitment, and subsequent enhancer accessibility and activity. Pparα then binds to target gene promoters to drive chromatin acetylation, thereby inducing transcription of fatty acid catabolic genes to alleviate heightened lipid burden.

Our system benefits from use of *in vivo* metabolic perturbations to assess the role for

excess fatty acids in driving transcription. It has been stipulated that fatty acid oxidation

promotes transcription of metabolic genes via bulk increase to histone acetylation by lipid-

derived acetyl-CoA (McDonnell et al., 2016). Our data instead reveal that changes in hepatic

lipid levels can induce metabolic gene transcription even when mitochondrial β-oxidation, and

subsequent acetyl-CoA production, is inhibited. Indeed we detected no changes in bulk histone

acetylation in Cpt2$^{L-/-}$ liver. We instead observed that promoter acetylation patterns for many

lipid catabolic genes are Pparα-dependent. Enhancers displayed Pparα-independent

acetylation proximal to Pparα binding sites, but enhancer chromatin remained inaccessible in

the absence of the transcription factor. This indicates that fatty acid-induced transcription is

stimulated not by epigenetic changes, but instead by ligand-induced Pparα activation and the

associated chromatin remodeling. Studies in mice given a high fat diet (HFD) provide

supplemental evidence for our observations on fatty acid-driven transcription. In parallel with

our results, HFD-fed mice show increased hepatic lipid content as well as Pparα-dependent

hepatic gene expression (Patsouris et al., 2006). HFD-fed mice likewise exhibit suppressed

liver acetyl-CoA content but no difference in bulk histone acetylation compared to chow-fed

controls (Carrer et al., 2017).

A common method to stimulate Pparα and define target genes is to use a

pharmacological agonist (Brocker et al., 2020; Janssen et al., 2015; Lee et al., 1995;

Rakhshandehroo et al., 2007). Here we instead use fasting-induced buildup of endogenous

ligand to activate Pparα. We observed two distinct patterns of target gene transcription. In the

first group (Cluster 1), target genes require Pparα for promoter accessibility and acetylation

under conditions of high lipid availability. In the second group (Cluster 3), target genes do not

require Pparα for promoter acetylation. Promoter accessibility for the latter cluster is maintained

in Pparα$^{-/-}$ animals, though to a lesser extent than measured in WT controls. We therefore

hypothesize that a secondary mechanism maintains transcriptionally permissive H3K27ac

occupancy at these loci independent of Pparα, thereby promoting baseline gene expression.

The additional requirement of Pparα for gene induction above baseline is reflected in both the

RNA-seq and ATAC-seq datasets.

Cluster 3 gene promoters are enriched for the Hnf4α biding motif, suggesting the nuclear

hormone receptor may be involved in Pparα-independent promoter acetylation. It should be

noted that Hnf4α can bind directly to promoter sequences of *Fgf21*, *Hmgcs2*, and *Cpt1* to

obstruct Pparα binding and gene transcription (Martinez-Jimenez et al., 2010). However, these

three genes were assigned to Cluster 1, leaving open the possibility of Hnf4α involvement in

Cluster 3 promoter activation. Hnf4α is highly expressed in the three knockout genotypes,

providing additional evidence for this suggestion. We similarly observe high incidence of Hnf4α

motifs at active enhancers, in line with previous work demonstrating that Hnf4α maintains

H3K27ac occupancy at active enhancers in mouse liver (Thakur et al., 2019).

Previous *in vivo* work has advanced a role for Pparα localization to hepatic enhancers.

In a model of diet induced obesity, Pparα-null mice had suppressed circadian transcription for

eRNA proximal to Pparα binding motifs (Guan et al., 2018). In another study, hepatic fed state

Pparα ChIP-seq binding sites were observed to overlap with fasting-induced enhancers

(Goldstein et al., 2017). Our research builds upon those observations by demonstrating that

Pparα-sensitive enhancer elements in fasting liver are associated with lipid signaling. Elevated

enhancer accessibility in Cpt2$^{L-/-}$ animals, including particularly high ATAC-seq signal over

Pparα binding sites, further indicates that lipids drive Pparα engagement with active chromatin.

Moreover, depleted chromatin accessibility over all active enhancers in Pparα-null animals

indicates the requirement of Pparα to properly maintain the hepatic fasting enhancer landscape.

It's notable that Pparα loss does not significantly impact global enhancer epigenetic

state. Indeed, enhancer priming proximal to Pparα binding sites is independent of Pparα itself.

Increased H3K4me1 signal in DKO animals compared to other genotypes likely indicates a

compensatory signal to engage lipid-sensitive enhancers in response to the hepatocyte's need

for oxidative clearance of excess fatty acid. Enhancer H3K27ac signal adjacent to Pparα

binding sites is likewise globally independent of Pparα. Both Cpt2$^{L-/-}$ and DKO animals display

the highest levels of enhancer acetylation near Pparα binding sites, and yet these two

genotypes also display the lowest global enhancer H3K27ac signal. In other words, the Cpt2$^{L-/-}$

background drives enhancer acetylation specifically in proximity to Pparα binding sites,

providing evidence for a mechanism in which high lipid levels prompt H3K4me1 and H3K27ac

deposition to prime Pparα-sensitive enhancers, and credibly acting as a signal for Pparα DNA

binding.

While overall enhancer maintenance near Pparα binding sites is independent of Pparα,

insight into the Pparα enhancer program is gained by examining differential comparisons

between genotypes. One striking observation was that Pparα-associated H3K27ac signal

changes occurred more frequently in enhancer elements compared to promoters. Cpt2$^{L-/-}$-

associated lipid prompts an H3K27ac ChIP-seq signal increase at 5.5x more enhancers than

promoters. Similarly, loss of Pparα in Class B enhancers (DKO < Cpt2$^{L-/-}$) is associated with

H3K27ac signal depletion at 3.7x more enhancers than promoters. Finally, Ppar$\alpha^{-/-}$ animals

show H3K27ac depletion at 1.7x more enhancers than promoters. This notable degree of signal

change at enhancer regions, combined with the high incidence of Pparα binding sites and

association with GREAT lipid metabolism ontology terms, suggests that Pparα effects more

transcriptionally influential changes to chromatin architecture at enhancers over promoters. We

believe this indicates that the mechanism for adjusting Pparα target gene transcription levels is

more closely linked to enhancer remodeling than to promoter dynamics.

These expanded insights into Pparα gene regulation provide evidence for discrete

modes by which the liver signals its transcriptional needs in the face of metabolic perturbations

linked to fluctuations in lipid availability. We demonstrate distinct types of Pparα signaling

based on patterns of chromatin state and transcriptional output, show the role for Pparα in

hepatic enhancer maintenance, and further suggest secondary mechanisms which aid and

potentially even modulate Pparα signaling. These data prompt questions as to what role Pparα

may play in regulating the transcriptional landscape associated with lipid sensing in other

tissues, such as the kidney and adipose. This study thus uncovers new avenues of

investigation into lipid sensing at the levels of locus-specificity, tissue-specificity, and broader

metabolic physiology.

MATERIALS & METHODS

Animals

All procedures were performed in accordance with the NIH's Guide for the Care and Use of

Laboratory Animals and under the approval of the Johns Hopkins School of Medicine Animal

Care and Use Committee.

$Cpt2^{fl/fl}$ and Albumin-Cre;$Cpt2^{fl/fl}$ ($Cpt2^{L-/-}$) mice were previously described (Lee et al., 2015,

2016). $Cpt2^{fl/fl}$;$Ppar\alpha^{-/-}$ and $Cpt2^{L-/-}$;$Ppar\alpha^{-/-}$ animals were generated by initially crossing $Cpt2^{fl/fl}$ or

$Cpt2^{L-/-}$ animals with $Ppar\alpha^{-/-}$ mice (Jackson Laboratories; stock no. 008154). Mice for

experiments were bred from either $Cpt2^{fl/fl}$ crossed to $Cpt2^{L-/-}$ animals or $Cpt2^{fl/fl}$;$Ppar\alpha^{-/-}$ crossed

to $Cpt2^{L-/-}$;$Ppar\alpha^{-/-}$ animals. All mice were housed in a facility with ventilated racks on a 14h

light/10h dark cycle with *ad libitum* access to a standard rodent chow (2018SX Teklad Global,

18% protein). For fasting experiments, 9 week old male mice were deprived of food for 24 hours

(3p.m.-3 p.m). Tissues and serum were collected and flash-frozen at time of harvest.

Immunoblotting

Flash-frozen liver tissue was homogenized in RIPA buffer (50 mM Tris-HCl at pH 7.4, 150 mM

NaCl, 1 mM EDTA, 1% Triton X-100, and 0.25% deoxycholate) with PhosSTOP phosphatase

Inhibitor (Roche) and protease inhibitor cocktail (Roche). Homogenates were rotated for 30

minutes at 4°C and then centrifuged at 18,000g for 15 minutes. Total protein concentrations

were quantified by BCA assay (Thermo Fisher Scientific).

Protein lysates (30ug input) were separated by Tris-glycine SDS-PAGE (10-12% polyacrylamide

gels), followed by a transfer to PVDF membranes (Immobilon). Membranes were blocked with

5% nonfat milk in TBST for 1 hour and incubated overnight at 4°C with primary antibodies at

1:1000 in 3% BSA in TBST: Acetylated-Lysine, 9441, Cell Signaling Technologies; Histone H3,

4499, Cell Signaling Technology, Acetyl-Histone H3 Lys9, 9649, Cell Signaling Technologies;

Acetyl-Histone H3 Lys14, 7627, Cell Signaling Technologies; HSC70, 7298, Santa Cruz

Biotechnology. HRP-conjugated anti-rabbit (Cytiva, NA934V) or fluorescence-based (Cy3-

conjugated anti-mouse or Cy5-conjugated anti-rabbit; Invitrogen, Thermo Fisher Scientific)

secondaries were used at 1:1000. Blots were imaged using the Amersham Prime enhanced

chemiluminescent substrate (Cytiva) or epifluorescence on an Alpha Innotech MultiImage III

instrument.

Histones were acid precipitated as previously described using an overnight 0.4N H_2SO_4

incubation and 1 hour TCA precipitation (Shechter et al., 2007). Histones were visualized per

above using 15ug protein input on an 8% polyacrylamide gel.

Mass Spectrometry & Proteomic Analysis

Flash-liver tissue was submitted to the Johns Hopkins Center for Proteomics Core for total

proteome and acetyl-proteome analysis. Samples were prepared in lysis buffer with 8M urea

and 50mM tetraethyl ammoniumbicarbonate. Lysates were treated with LysC and trypsin, then

labeled with 10-plex TMT. Peptide fractionation was done using basic pH reverse phase liquid

chromatography, resulting in 24 fractions. Fractions were analyzed using an Orbitrap Fusion

Lumos (Thermo Scientific) on an Easy nLC 1200 (Thermo Scientific) using MS1 resolution =

120,000 and MS2 resolution = 30,000, HCD fragmentation method, and MS2 collision energy =

35. Each fraction received a two hour run time.

Total proteome analysis was done in Perseus (version 1.6.0.0) (Cox and Mann, 2012). p-value

was calculated by Student's t-test. q-value was calculated by significance analysis of

microarrays (SAM) and permutation based false discovery rate (FDR), with SAM S0 value = 0.1

(Tusher et al., 2001).

Acetyl-proteome was analyzed with Proteome Discoverer 2.1 (Thermo Scientific) using RefSeq

version 78. The following parameters were used: cleavage enzyme was set as trypsin, no more

than five missed cleavages were allowed, fixed modification was carbamidomethyl on cysteine

residue, dynamic modifications were acetyl group on protein N-terminus or oxidation on

methionine and acetyl-lysine, and MS2 level quantification. A 1% FDR was applied for both

peptide and protein levels. Acetyl-proteome data were normalized to median of total proteome

to remove systemic deviation.

The COMPARTMENT dataset was used to assign localization for peptides for all cell

compartments except mitochondria (Binder et al., 2014). Only COMPARTMENTS annotations

with a minimum confidence score of 3 were used. Mitochondrial peptides were assigned using

the MitoCarta 2.0 (Calvo et al., 2016).

Tissue acetyl-CoA

Flash-frozen liver tissue was submitted to the Mouse Metabolic Phenotyping Center at Case

Western Reserve University for acetyl-CoA measurements via liquid chromatography-mass

spectrometry.

Serum metabolites

Blood glucose levels in fasted mice were measured at time of harvest using a Nova Max Plus

glucometer.

Untargeted metabolomics on serum was performed by Metabolon Inc (n=6). For analysis raw

area counts for each biochemical species were rescaled to set the median equal to 1. Heatmap

and PCA were generated by MetaboAnalyst (Chong et al., 2019). Differentially regulated

metabolites for heatmap were determined using a 1-way ANOVA with Fisher's LSD.

RNA-sequencing library preparation

Total RNA was isolated from flash-frozen liver tissue was using TRIzol reagent (Invitrogen,

Thermo Fisher Scientific), followed by addition purification using RNeasy Mini Kit (QIAGEN), per

manufacturer recommendations. RNA quality was assessed by Nanodrop.

RNA was then submitted to Novogene Corporation Inc (China & Davis, CA, USA) for library

construction and sequencing. Four biological replicates were used for each genotype. Reads

were aligned to mouse reference genome (mm10) using STAR software (Dobin et al., 2013).

Investigators performed analyses downstream of read count acquisition.

RNA-seq Analysis

Differential expression was performed on raw read counts in R with DESeq2 (v3.12) using the

Wald test with betaPrior=FALSE and lfcShrink type="apeglm" (Love et al., 2014; Zhu et al.,

2019). Differential expression cutoff was fold change ≥ |2|, padj < 0.05. Normalized read counts

were visualized in GraphPad Prism. PCA plot was generated using pcaExlorer package (Marini

and Binder, 2019). WY14643-responsive genes were determined from GSE140063 using the

same differential expression pipeline (Naiman et al., 2019).

K-means clustering of differentially expressed transcripts was performed in R on rlog

transformed count values from DESeq2 output. Optimal cluster number (n=5) was determined

by gap statistic (factoextra package). K-means clustering was performed and visualized in the

pheatmap package with Z-score row scaling $[(x - \bar{x})/\sigma]$. Tracing diagrams were generated using

the ggplot2 package (Wickham, 2016). Gene ontology and KEGG pathway enrichment was

obtained using the DAVID web application (Huang et al., 2007).

ATAC-seq and ChIP-seq library preparation

ATAC-seq and ChIP-seq were performed using Active Motif sequencing services (Carlsbad,

CA). Flash-frozen liver tissue was submitted to Active Motif, with two biological replicates per

genotype for H3K4me1 and H3K27ac ChIP-seq, and ATAC-seq. One biological replicate was

submitted for H3K9ac ChIP-seq. Library construction was performed according to company

protocol. The following antibodies were used for ChIP-seq: H3K4me1 (Active Motif #39297),

H3K9ac (Active Motif #39917), H3K27ac (Active Motif #39133).

ATAC-Seq & ChIP-seq Analysis

Sequence acquisition, mapping, raw BAM file generation, and peak calling was performed by

Active Motif. For ATAC-seq, paired-end 42nt sequencing reads generated by Illumina NexSeq

500 were mapped to mm10 reference genome using the BWA algorithm with default settings (Li

and Durbin, 2009). For ChIP-seq, single-end 75nt sequencing reads generated by Illumina

NexSeq 500 were mapped to mm10 reference genome using the BWA algorithm with

"bwaln/samse" default settings. For normalization, tag number of all samples within a

comparison group was reduced by random down-sampling to the number of tags present in the

smallest sample. Peak calling was performed using MACS2 with default cutoff values (Zhang et

al., 2008); ATAC-seq paired reads were treated as independent events. ENCODE blacklist

regions were removed (Amemiya et al., 2019). bigWig files for visualization on UCSC Genome

Browser were created using a 32nt bin with a 200bp *in silico* tag extension (Kent et al., 2002).

Investigators performed downstream analysis.

Raw BAM files were processed using samtools and bamtools to remove reads with mapq score

< 25, unmapped reads, mitochondrial reads, and reads with more than 2 mismatches (Barnett

et al., 2011; Li et al., 2009). Aggregation plots and motif analysis were generated

using Hypergeometric Optimization of Motif EnRichment (HOMER) (Heinz et al., 2010). Motif

analysis was calculated using 'findMotifsGenome.pl' with -size 500 on BED file input with that

assumption that a broad scanning window is preferred for histone mark motif analysis.

knownResults motif output are reported in this study. Tag directories for histograms were

generated from BAM files using 'makeTagDirectory.pl' with -fragLength 200 and -tbp 1. Tag

directories combined both biological replicates per genotype. Aggregation plots were produced

using 'annotatePeaks.pl' with -hist 10 and normalization set to the lowest sample tag count

within a comparison group. Plots and corresponding bar charts, reported as tags per bp per

peak, were visualized in GraphPad Prism 9.

Diffbind (v3.13) was used to generate a RangedSummarizedExperiment from BAM and

narrowPeak files with the following parameters: fragmentLength=200,

score=DBA_SCORE_READS, summits=TRUE, bUseSummarizeOverlaps=FALSE,

bRemoveDuplicates=TRUE, and minOverlap=2 (Ross-Innes et al., 2012; Stark and Brown,

2011). mm10 ENCODE blacklist regions were removed by calling dba.blacklist(). Default

normalization with full library size was called with dba.normalize(). ATAC-seq normalization

included background=TRUE due to large changes between genotypes.

RangedSummarizedExperiment was passed to DESeq2 for differential peak analysis using

betaPrior=TRUE. Significance cutoff for differential peak analysis was considered fold change ≥

|2|, padj < 0.05. ATAC-seq MA plots were generated in Diffbind. PCA plots were generated

using pcaExlorer on DESeq2 rlog transformed read counts.

Enhancer Analysis

All genome arithmetic to compare peak coordinates was done using bedtools (Quinlan, 2014;

Quinlan and Hall, 2010). This study only focused on active enhancer elements that were

detected in one or more genotypes; regions enriched for H3K4me1 alone were excluded. Active

enhancers were defined as a region with H3K4me1 and H3K27ac co-occupancy excluding

intervals located within 2kb of a transcription start site or termination site (Calo and Wysocka,

2013; Creyghton et al., 2010). Chromatin accessibility was not used to define active enhancers

due to the severe global depletion in ATAC-seq tag density observed in Pparα-null animals.

Gene coordinates were obtained from the UCSC Table Browser refGene (2020-08-17 update)

(Karolchik et al., 2004). H3K4me1 and H3K27ac overlap was determined with a minimum 50%

H3K27ac interval overlap over an H3K4me1 peak due to account for differences in histone mark

average peak length. Fasting liver Pparα ChIP-seq coordinates were obtained from GSE118788

(Supplementary File Pparα-C57-Fast-peaks) (Sommars et al., 2019).

Differential enhancer groups were defined using the H3K27ac ChIP-seq DESeq2 output from

above with the significance cutoff set at fold change ≥ |2|, padj < 0.05. DESeq2 normalized

values were used for enhancer tag counts. Enhancer heatmaps were generated from bigWig

files using deepTools 2.0 with 'computeMatrix' reference-point mode and 'plotHeatmap'

(Ramírez et al., 2016).

Enhancer ontology was determined with the Genomic Regions Enrichment of Annotations Tool

(GREAT) (McLean et al., 2010). GREAT version 4.0.4 was used in "basal plus extension" mode

with default parameters and a 500kb maximum extension. Curated regulatory domains were

included in calculations.

Statistical Analysis

All statistical comparisons were carried out in GraphPad Prism 9 unless otherwise noted.

Significance was determined using Student's t-test, 1-way ANOVA with Tukey's post hoc

correction, or Kruskal-Wallis test with Dunn's post hoc correction as noted. Shapiro-Wilk test for

normality was used in R to determine whether to use a parametric or non-parametric test for

significance for genomic data.

DATA AND SOFTWARE AVAILABILITY

RNA-seq data were deposited in GSE165701. ChIP-seq and ATAC-seq data were deposited in

GEO SuperSeries GSE179053. Scripts used for RNA-seq, ChIP-seq, and ATAC-seq analysis

can be found at https://github.com/WolfgangLabJHMI/cpt2_ppara_liver_seq. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

the PRIDE partner repository with the dataset identifier PXD027235.

ACKNOWLEDGEMENTS

This work was supported in part by a National Institutes of Health grant R01DK120530 and

R01DK116746 to M.J.W.

We thank Dr. Caitlyn Bowman-Cornelius for initial mouse breeding, Dr. Ebru Selen for advice on

metabolomics analysis, and Dr. Anastasia Kralli for helpful conversations. We also thank Dr.

Chan-Hyun Na and the Johns Hopkins Center for Proteomics Discovery for proteomic data

acquisition.

AUTHOR CONTRIBUTIONS

M.J.W and K.S.C conceptualized the project. K.S.C collected and analyzed data. K.S.C and

M.J.W participated in the writing, review, and editing of the manuscript.

REFERENCES

Amemiya, H.M., Kundaje, A., and Boyle, A.P. (2019). The ENCODE Blacklist: Identification of
Problematic Regions of the Genome. Scientific Reports *9*, 1–5.

Andersson, R., and Sandelin, A. (2020). Determinants of enhancer and promoter activities of
regulatory elements. Nature Reviews Genetics *21*, 71–87.

Aoyama, T., Peters, J.M., Iritani, N., Nakajima, T., Furihata, K., Hashimoto, T., and Gonzalez,
F.J. (1998). Altered constitutive expression of fatty acid-metabolizing enzymes in mice
lacking the peroxisome proliferator-activated receptor α (PPARα). Journal of Biological
Chemistry *273*, 5678–5684.

Asrani, S.K., Devarbhavi, H., Eaton, J., and Kamath, P.S. (2019). Burden of liver diseases in the
world. Journal of Hepatology *70*, 151–171.

Bardot, O., Aldridge, T.C., Latruffe, N., and Green, S. (1993). PPAR-RXR heterodimer activates
a peroxisome proliferator response element upstream of the bifunctional enzyme gene.
Biochemical and Biophysical Research Communications *192*, 37–45.

Barnett, D.W., Garrison, E.K., Quinlan, A.R., Stïmberg, M.P., and Marth, G.T. (2011). Bamtools:
A C++ API and toolkit for analyzing and managing BAM files. Bioinformatics *27*, 1691–
1692.

Binder, J.X., Pletscher-Frankild, S., Tsafou, K., Stolte, C., O'Donoghue, S.I., Schneider, R., and
Jensen, L.J. (2014). COMPARTMENTS: Unification and visualization of protein subcellular
localization evidence. Database *2014*, bau012–bau012.

Boergesen, M., Pedersen, T.A., Gross, B., van Heeringen, S.J., Hagenbeek, D., Bindesboll, C.,
Caron, S., Lalloyer, F., Steffensen, K.R., Nebb, H.I., et al. (2012). Genome-Wide Profiling of
Liver X Receptor, Retinoid X Receptor, and Peroxisome Proliferator-Activated Receptor   in
Mouse Liver Reveals Extensive Sharing of Binding Sites. Molecular and Cellular Biology

32, 852–867.

Bowman, C.E., Selen Alpergin, E.S., Cavagnini, K., Smith, D.M., Scafidi, S., and Wolfgang, M.J.

(2019). Maternal Lipid Metabolism Directs Fetal Liver Programming following Nutrient

Stress. Cell Reports *29*.

Brocker, C.N., Kim, D., Melia, T., Karri, K., Velenosi, T.J., Takahashi, S., Aibara, D., Bonzo,

J.A., Levi, M., Waxman, D.J., et al. (2020). Long non-coding RNA Gm15441 attenuates

hepatic inflammasome activation in response to PPARA agonism and fasting. Nature

Communications *11*.

Calo, E., and Wysocka, J. (2013). Modification of Enhancer Chromatin: What, How, and Why?

Molecular Cell *49*, 825–837.

Calvo, S.E., Clauser, K.R., and Mootha, V.K. (2016). MitoCarta2.0: An updated inventory of

mammalian mitochondrial proteins. Nucleic Acids Research *44*, D1251–D1257.

Carrer, A., Parris, J.L.D., Trefely, S., Henry, R.A., Montgomery, D.C., Torres, A., Viola, J.M.,

Kuo, Y.M., Blair, I.A., Meier, J.L., et al. (2017). Impact of a high-fat diet on tissue Acyl-CoA

and histone acetylation levels. Journal of Biological Chemistry *292*, 3312–3322.

Chong, J., Wishart, D.S., and Xia, J. (2019). Using MetaboAnalyst 4.0 for Comprehensive and

Integrative Metabolomics Data Analysis. Current Protocols in Bioinformatics *68*, e86.

Cox, J., and Mann, M. (2012). 1D and 2D annotation enrichment: a statistical method integrating

quantitative proteomics with complementary high-throughput data. BMC Bioinformatics *13*

Suppl 16, 1–11.

Creyghton, M.P., Cheng, A.W., Welstead, G.G., Kooistra, T., Carey, B.W., Steine, E.J., Hanna,

J., Lodato, M.A., Frampton, G.M., Sharp, P.A., et al. (2010). Histone H3K27ac separates

active from poised enhancers and predicts developmental state. Proceedings of the

National Academy of Sciences of the United States of America *107*, 21931–21936.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., and Gingeras, T.R. (2013). STAR: Ultrafast universal RNA-seq aligner. Bioinformatics

29, 15–21.

Evans, R.M., and Mangelsdorf, D.J. (2014). Nuclear receptors, RXR, and the big bang. Cell

157, 255–266.

Francque, S., Verrijken, A., Caron, S., Prawitt, J., Paumelle, R., Derudas, B., Lefebvre, P.,

Taskinen, M.R., Van Hul, W., Mertens, I., et al. (2015). PPARα gene expression correlates

with severity and histological treatment response in patients with non-alcoholic

steatohepatitis. Journal of Hepatology *63*, 164–173.

Gearing, K.L., Gottlicher, M., Teboul, M., Widmark, E., and Gustafsson, J.A. (1993). Interaction

of the peroxisome-proliferator-activated receptor and retinoid X receptor. Proceedings of

the National Academy of Sciences of the United States of America *90*, 1440–1444.

George, F.C. (2006). Fuel metabolism in starvation. Annual Review of Nutrition *26*, 1–22.

Goldstein, I., and Hager, G.L. (2015). Transcriptional and Chromatin Regulation during Fasting -

The Genomic Era. Trends in Endocrinology and Metabolism *26*, 699–710.

Goldstein, I., Baek, S., Presman, Di.M., Paakinaho, V., Swinstead, E.E., and Hager, G.L.

(2017). Transcription factor assisted loading and enhancer dynamics dictate the hepatic

fasting response. Genome Research *27*, 427–439.

Gong, Z., Tas, E., Yakar, S., and Muzumdar, R. (2017). Hepatic lipid metabolism and non-

alcoholic fatty liver disease in aging. Molecular and Cellular Endocrinology *455*, 115–130.

Greenberg, A.S., Coleman, R.A., Kraemer, F.B., McManaman, J.L., Obin, M.S., Puri, V., Yan,

Q.W., Miyoshi, H., and Mashek, D.G. (2011). The role of lipid droplets in metabolic disease

in rodents and humans. Journal of Clinical Investigation *121*, 2102–2110.

Guan, D., Xiong, Y., Borck, P.C., Jang, C., Doulias, P.T., Papazyan, R., Fang, B., Jiang, C.,

Zhang, Y., Briggs, E.R., et al. (2018). Diet-Induced Circadian Enhancer Remodeling

Synchronizes Opposing Hepatic Lipid Metabolic Processes. Cell *174*, 831-842.e12.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,

Singh, H., and Glass, C.K. (2010). Simple Combinations of Lineage-Determining

Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell

Identities. Molecular Cell *38*, 576–589.

Heinz, S., Romanoski, C.E., Benner, C., and Glass, C.K. (2015). The selection and function of

cell type-specific enhancers. Nature Reviews Molecular Cell Biology *16*, 144–154.

Houten, S.M., Violante, S., Ventura, F. V., and Wanders, R.J.A. (2016). The Biochemistry and

Physiology of Mitochondrial Fatty Acid β-Oxidation and Its Genetic Disorders. Annual

Review of Physiology *78*, 23–44.

Huang, B., Wu, P., Bowker-Kinley, M.M., and Harris, R.A. (2002). Regulation of pyruvate

dehydrogenase kinase expression by peroxisome proliferator-activated receptor-α ligands,

glucocorticoids, and insulin. Diabetes *51*, 276–283.

Huang, D.W., Sherman, B.T., Tan, Q., Kir, J., Liu, D., Bryant, D., Guo, Y., Stephens, R.,

Baseler, M.W., Lane, H.C., et al. (2007). DAVID Bioinformatics Resources: Expanded

annotation database and novel algorithms to better extract biology from large gene lists.

Nucleic Acids Research *35*.

Inagaki, T., Dutchak, P., Zhao, G., Ding, X., Gautron, L., Parameswara, V., Li, Y., Goetz, R.,

Mohammadi, M., Esser, V., et al. (2007). Endocrine Regulation of the Fasting Response by

PPARα-Mediated Induction of Fibroblast Growth Factor 21. Cell Metabolism *5*, 415–425.

Iroz, A., Montagner, A., Benhamed, F., Levavasseur, F., Polizzi, A., Anthony, E., Régnier, M.,

Fouché, E., Lukowicz, C., Cauzac, M., et al. (2017). A Specific ChREBP and PPARα

Cross-Talk Is Required for the Glucose-Mediated FGF21 Response. Cell Reports *21*, 403–

416.

Janssen, A.W.F., Betzel, B., Stoopen, G., Berends, F.J., Janssen, I.M., Peijnenburg, A.A., and

Kersten, S. (2015). The impact of PPARaα activation on whole genome gene expression in

human precision cut liver slices. BMC Genomics *16*.

Jump, D.B., Tripathy, S., and Depner, C.M. (2013). Fatty acid-regulated transcription factors in

the liver. Annual Review of Nutrition *33*, 249–269.

Karagianni, P., and Talianidis, I. (2015). Transcription factor networks regulating hepatic fatty

acid metabolism. Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids

1851, 2–8.

Karolchik, D., Hinricks, A.S., Furey, T.S., Roskin, K.M., Sugnet, C.W., Haussler, D., and Kent,

W.J. (2004). The UCSC table browser data retrieval tool. Nucleic Acids Research *32*.

Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler,

a. D. (2002). The Human Genome Browser at UCSC. Genome Research *12*, 996–1006.

Kersten, S. (2014). Integrated physiology and systems biology of PPARα. Molecular Metabolism

3, 354–371.

Kersten, S., Seydoux, J., Peters, J.M., Gonzalez, F.J., Desvergne, B., and Wahli, W. (1999).

Peroxisome proliferator-activated receptor α mediates the adaptive response to fasting.

Journal of Clinical Investigation *103*, 1489–1498.

Kersten, S., Rakhshandehroo, M., Knoch, B., and Müller, M. (2010). Peroxisome proliferator-

activated receptor alpha target genes. PPAR Research.

Lee, J., Ellis, J.M., and Wolfgang, M.J. (2015). Adipose fatty acid oxidation is required for

thermogenesis and potentiates oxidative stress-induced inflammation. Cell Reports *10*,

266–279.

Lee, J., Choi, J., Scafidi, S., and Wolfgang, M.J. (2016). Hepatic Fatty Acid Oxidation Restrains

Systemic Catabolism during Starvation. Cell Reports *16*, 201–212.

Lee, J.M., Wagner, M., Xiao, R., Kim, K.H., Feng, D., Lazar, M.A., and Moore, D.D. (2014).

Nutrient-sensing nuclear receptors coordinate autophagy. Nature *516*, 112–115.

Lee, S.S., Pineau, T., Drago, J., Lee, E.J., Owens, J.W., Kroetz, D.L., Fernandez-Salguero,

P.M., Westphal, H., and Gonzalez, F.J. (1995). Targeted disruption of the alpha isoform of

the peroxisome proliferator-activated receptor gene in mice results in abolishment of the

pleiotropic effects of peroxisome proliferators. Molecular and Cellular Biology *15*, 3012–

3022.

Leone, T.C., Weinheimer, C.J., and Kelly, D.P. (1999). A critical role for the peroxisome

proliferator-activated receptor α (PPARα) in the cellular fasting response: The PPARα-null

mouse as a model of fatty acid oxidation disorders. Proceedings of the National Academy

of Sciences of the United States of America *96*, 7473–7478.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics *25*, 1754–1760.

Li, G., Brocker, C.N., Xie, C., Yan, T., Noguchi, A., Krausz, K.W., Xiang, R., and Gonzalez, F.J.

(2018). Hepatic peroxisome proliferator-activated receptor alpha mediates the major

metabolic effects of Wy-14643. Journal of Gastroenterology and Hepatology (Australia) *33*,

1138–1145.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics

25, 2078–2079.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biology *15*, 550.

Mandard, S., Müller, M., and Kersten, S. (2004). Peroxisome proliferator-activated receptor α

target genes. Cellular and Molecular Life Sciences *61*, 393–416.

Marini, F., and Binder, H. (2019). PcaExplorer: An R/Bioconductor package for interacting with

RNA-seq principal components. BMC Bioinformatics *20*, 1–8.

Martinez-Jimenez, C.P., Kyrmizi, I., Cardot, P., Gonzalez, F.J., and Talianidis, I. (2010).

Hepatocyte Nuclear Factor 4 Coordinates a Transcription Factor Network Regulating

Hepatic Fatty Acid Metabolism. Molecular and Cellular Biology *30*, 565–577.

McDonnell, E., Crown, S.B., Fox, D.B., Kitir, B., Ilkayeva, O.R., Olsen, C.A., Grimsrud, P.A., and

Hirschey, M.D. (2016). Lipids Reprogram Metabolism to Become a Major Carbon Source

for Histone Acetylation. Cell Reports *17*, 1463–1472.

McLean, C.Y., Bristor, D., Hiller, M., Clarke, S.L., Schaar, B.T., Lowe, C.B., Wenger, A.M., and

Bejerano, G. (2010). GREAT improves functional interpretation of cis-regulatory regions.

Nature Biotechnology *28*, 495–501.

Naiman, S., Huynh, F.K., Gil, R., Glick, Y., Shahar, Y., Touitou, N., Nahum, L., Avivi, M.Y.,

Roichman, A., Kanfi, Y., et al. (2019). SIRT6 Promotes Hepatic Beta-Oxidation via

Activation of PPARα. Cell Reports *29*, 4127-4143.e8.

Patsouris, D., Reddy, J.K., Müller, M., and Kersten, S. (2006). Peroxisome proliferator-activated

894   receptor α mediates the effects of high-fat diet on hepatic gene expression. Endocrinology
895   *147*, 1508–1516.

896 Ponziani, F.R., Pecere, S., Gasbarrini, A., and Ojetti, V. (2015). Physiology and pathophysiology
897   of liver lipid metabolism. Expert Review of Gastroenterology and Hepatology *9*, 1055–1067.

898 Qin, Y., Grimm, S.A., Roberts, J.D., Chrysovergis, K., and Wade, P.A. (2020). Alterations in
899   promoter interaction landscape and transcriptional network underlying metabolic adaptation
900   to diet. Nature Communications *11*, 1–16.

901 Quinlan, A.R. (2014). BEDTools: The Swiss-Army tool for genome feature analysis. Current
902   Protocols in Bioinformatics *2014*, 11.12.1-11.12.34.

903 Quinlan, A.R., and Hall, I.M. (2010). BEDTools: A flexible suite of utilities for comparing genomic
904   features. Bioinformatics *26*, 841–842.

905 Raisner, R., Kharbanda, S., Jin, L., Jeng, E., Chan, E., Merchant, M., Haverty, P.M., Bainer, R.,
906   Cheung, T., Arnott, D., et al. (2018). Enhancer Activity Requires CBP/P300 Bromodomain-
907   Dependent Histone H3K27 Acetylation. Cell Reports *24*, 1722–1729.

908 Rakhshandehroo, M., Sanderson, L.M., Matilainen, M., Stienstra, R., Carlberg, C., De Groot,
909   P.J., Müller, M., and Kersten, S. (2007). Comprehensive analysis of PPARα-dependent
910   regulation of hepatic lipid metabolism by expression profiling. PPAR Research *2007*.

911 Rakhshandehroo, M., Hooiveld, G., Müller, M., and Kersten, S. (2009). Comparative Analysis of
912   Gene Regulation by the Transcription Factor PPARα between Mouse and Human. PLoS
913   ONE *4*, e6796.

914 Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar,
915   F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing
916   data analysis. Nucleic Acids Research *44*, W160–W165.

917 Reddy, J.K. (2001). Nonalcoholic steatosis and steatohepatitis III. Peroxisomal β-oxidation,
918   PPARα, and steatohepatitis. American Journal of Physiology - Gastrointestinal and Liver
919   Physiology *281*.

920 Ross-Innes, C.S., Stark, R., Teschendorff, A.E., Holmes, K.A., Ali, H.R., Dunning, M.J., Brown,
921   G.D., Gojis, O., Ellis, I.O., Green, A.R., et al. (2012). Differential oestrogen receptor binding
922   is associated with clinical outcome in breast cancer. Nature *481*, 389–393.

923 Ruppert, P.M.M., Park, J.G., Xu, X., Hur, K.Y., Lee, A.H., and Kersten, S. (2019).
924   Transcriptional profiling of PPARα-/- and CREB3L3-/- livers reveals disparate regulation of
925   hepatoproliferative and metabolic functions of PPARα. BMC Genomics *20*.

926 Selen, E.S., Choi, J., and Wolfgang, M.J. (2021). Discordant hepatic fatty acid oxidation and
927   triglyceride hydrolysis leads to liver disease. JCI Insight *6*.

Shechter, D., Dormann, H.L., Allis, C.D., and Hake, S.B. (2007). Extraction, purification and
analysis of histones. Nature Protocols *2*, 1445–1457.

Sommars, M.A., Ramachandran, K., Senagolage, M.D., Futtner, C.R., Germain, D.M., Allred,
A.L., Omura, Y., Bederman, I.R., and Barish, G.D. (2019). Dynamic repression by BCL6
controls the genome-wide liver response to fasting and steatosis. ELife *8*.

Song, S., Attia, R.R., Connaughton, S., Niesen, M.I., Ness, G.C., Elam, M.B., Hori, R.T., Cook,
G.A., and Park, E.A. (2010). Peroxisome proliferator activated receptor α (PPARα) and
PPAR gamma coactivator (PGC-1α) induce carnitine palmitoyltransferase IA (CPT-1A) via
independent gene elements. Molecular and Cellular Endocrinology *325*, 54–63.

Stark, R., and Brown, G. (2011). DiffBind: differential binding analysis of ChIP-Seq peak data.
http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf.

Stern, J.H., Rutkowski, J.M., and Scherer, P.E. (2016). Adiponectin, Leptin, and Fatty Acids in
the Maintenance of Metabolic Homeostasis through Adipose Tissue Crosstalk. Cell
Metabolism *23*, 770–784.

Thakur, A., Wong, J.C.H., Wang, E.Y., Lotto, J., Kim, D., Cheng, J.C., Mingay, M., Cullum, R.,
Moudgil, V., Ahmed, N., et al. (2019). Hepatocyte Nuclear Factor 4-Alpha Is Essential for
the Active Epigenetic State at Enhancers in Mouse Liver. Hepatology *70*, 1360–1376.

The, B., and Schulz, H. (1991). Beta oxidation of fatty acids. Biochimica et Biophysica Acta
(BBA) - Lipids and Lipid Metabolism *1081*, 109–120.

Tusher, V.G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays applied to
the ionizing radiation response. Proceedings of the National Academy of Sciences of the
United States of America *98*, 5116–5121.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (New York: Springer-Verlag).

Wu, P., Peters, J.M., and Harris, R.A. (2001). Adaptive increase in pyruvate dehydrogenase
kinase 4 during starvation is mediated by peroxisome proliferator-activated receptor α.
Biochemical and Biophysical Research Communications *287*, 391–396.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nussbaum, C.,
Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biology *9*, R137.

Zhu, A., Ibrahim, J.G., and Love, M.I. (2019). Heavy-Tailed prior distributions for sequence
count data: Removing the noise and preserving large differences. Bioinformatics *35*, 2084–
2092.