**Title Page:**

**Title**: TNF-α levels in respiratory samples are associated with SARS-CoV-2 infection.

**Running title:** Cytokine levels in respiratory samples.

**Authors**:

Matias J. PERESON[a,b] *, Maria Noel BADANO[a] *, Natalia ALOISI[a], Roberto CHUIT[c], MME de

BRACCO[a], Patricia BARE[a]

**Affiliations**:

[a]Instituto de Medicina Experimental (IMEX) CONICET, Instituto de Investigaciones

Hematologicas (IIHEMA), Academia Nacional de Medicina, Buenos Aires, Argentina

[b]Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Instituto de

Investigaciones en Bacteriología y Virología Molecular (IBaViM). Buenos Aires, Argentina.

[c] Instituto de Investigaciones Epidemiológicas, Academia Nacional de Medicina.

*Contributed equally to this work.

**Corresponding Author**:

Patricia Bare, Instituto de Medicina Experimental (IMEX), CONICET, Instituto de

Investigaciones Hematologicas (IIHEMA), Academia Nacional de Medicina, Buenos Aires,

Argentina.

Tel: +541148055759 ext 269; Tel: +5491154975179;

Fax: +541148050712, E-mail: pbare@hematologia.anm.edu.ar; patobare@hotmail.com

**Declarations of interest:** none.

**ABSTRACT**

**Purpose:**

The aim of this study was to measure levels of IL-6 and TNF-α in respiratory samples from

individuals with symptoms compatible with COVID-19 and analyze their association with

SARS-CoV-2 presence.

**Methods:**

SARS-CoV-2 detection was performed using the CDC (USA) real-time RT-PCR primers,

probes and protocols. Cytokine concentrations were measured using commercial reagents

based on enzyme linked immunosorbent assay (ELISA).

**Results**:

TNF-α median levels were greater in COVID19 (+) symptomatic group (5.88 (1.36 - 172.1)

pg/ml) compared to COVID19 (-) symptomatic individuals (2.87 (1.45 – 69.9) pg/ml)

(p=0.0003). No significant differences were shown in IL-6 median values between COVID-19

(+) and (-) symptomatic patients (5.40 (1.7 - 467) pg/ml and 6.07 (1.57 – 466.6) pg/ml

respectively). In addition, increased TNF-α levels (greater than 10 pg/ml), but not IL-6, were

associated with SARS-CoV-2 presence (OR= 5.7; p=0.006; 95% CI= 1,551 to 19,11).

**Conclusions**:

We found a statistically significant association between the production of local TNF-α and

the presence of the virus in early stages of infection. IL-6 showed high levels in swabs from

some symptomatic patients but independent from SARS-CoV-2 presence and viral load,

individual's age and gender. On the contrary, TNF-α evaluation confirmed the presence of

inflammatory response but mostly related to COVID-19. More studies are required in order to

characterize the cytokine profile expressed at the site of infection of SARS-CoV-2 and its

implications in disease outcomes.

**Keywords**: SARS-COV-2; COVID-19; respiratory samples; Interleukin-6; TNF-α

**Introduction**

The coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is one of the current major health concerns. After 15 months since the emergence of the pandemic SARS-CoV-2, up to 194 million cases were confirmed, including more than 4 million deaths worldwide due to severe forms of COVID-19 (WHO, 2021). Severe COVID-19 is widely considered an immunopathology disease driven by an excessive and deleterious immune response mounted against the pathogen rather than by the action of the pathogen *per se* (Mangalmurti & Hunter, 2020; Pedersen, & Ho, 2020; Hu et al. 2021). Many components of the innate immune system have been associated with this phenomenon and have been proposed as biomarker for COVID-19 progression. In this sense, high levels of serum interleukin-6 (IL-6) and tumor necrosis factor alpha (TNF-α) were correlated with increased morbidity and mortality in COVID-19 patients (Costella-Ruiz et al. 2020; Del Valle et al. 2020; Herold et al. 2020; Robinson et al. 2020a; Zhu et al. 2020; Danlos et al. 2021; Sabaka et al. 2021). However, innate immune mediators are necessary for efficient clearance of infectious agents. On the other hand, reports of lower median IL-6 levels in severe COVID-19 compared with other inflammatory conditions, such as acute respiratory distress syndrome and bacterial sepsis were published (Chen et al. 2020; Hambali et al. 2020; Chen & Quach, 2021). Additionally, many studies found no significant differences in TNF-α levels between severe and non-severe COVID-19 cases (Udomsinprasert et al. 2021). Therefore, the role of these cytokines in COVID-19 progression remains contradictory and needs further study.

To our knowledge, IL-6 and TNF-α concentration at the site of infection in COVID-19 patients has not been previously reported. Therefore, the aim of this study was to evaluate IL-6 and TNF-α concentration in swab samples from individuals showing symptoms compatible with COVID-19 who were positive or negative for SARS-CoV-2 genome detection.

## Material and methods

*Patients and study design*

The study was conducted on 127 archived swab samples referred to the Academia Nacional de Medicina for diagnosis of SARS-CoV-2 by real-time polymerase reaction (RT-PCR) between August 2020 and April 2021. Samples were obtained during the initial stages of disease progression (1 to 10 days). Symptomatic individuals with detectable (n=52) or undetectable (n=33) SARS-CoV-2 infection were analyzed. A group consisting of asymptomatic individuals and negative for SARSCoV-2 genome (n=42) were included to obtain cytokines' normal range.

*Collection of samples*

Combined nasopharyngeal and oropharyngeal swabs were collected by trained healthcare staff, placed into a single tube pre-filled with 2 mL of saline solution (0,9% sodium chloride) and were transferred on the same day to the laboratory for SARS-CoV-2 genome detection. Samples were stored at −80 °C until cytokine assays were performed.

*SARS-CoV-2 genome detection and viral load*

Genomic extraction was carried out with the QIAamp Viral RNA minikit (Qiagen) following manufacturer's instructions. A real time-based methodology was performed for viral genome detection using the CDC RT-qPCR protocol for Wuhan virus (nCoV-19), 2019-nCov CDC USA. The procedure incorporates a set of oligonucleotides primers and double-marker hydrolysis probes (Taqman®) (2019 nCov_N1 and N2) that amplify regions of the viral nucleocapsid gene (N). An internal control for RNase P (RP) was also included. Viral load was calculated considering Ct values.

*IL-6 and TNF-α measurement*

Concentrations of IL-6 (standard curve range: 0-300 pg/mL) and TNF-α (standard curve

range: 0 -500 pg/mL) in swabs were determined using commercial reagents based on

enzyme linked immunosorbent assay (ELISA) (BD-Biosciences, San Diego, California,

United States). Recommendations of the supplier were followed for procedure. Duplicates

were performed in selected samples to verify the accuracy of the results.

*Statistical analysis*

Mann-Whitney U test was used for testing differences between two groups. Categorical

variables were compared with Chi-square test or Fisher's exact test. Confidence intervals

were set at 95% (CI95). In all cases, a p value <0.05 was considered significant. Data and

graphs were performed using the GraphPad Prism 9.1.0 software (GraphPad Software, San

Diego, CA, USA).

*Ethics approval*

Experimental protocols and procedures performed in this work have been approved by

the Biosafety Review board of the IMEX-CONICET-Academia Nacional de Medicina and the

Ethical Committee of the Academia Nacional de Medicina.

**Results**

*Study population*

Combined oro- and nasopharyngeal swabs from 127 patients submitted for diagnosis of COVID-19 were examined for SARS-CoV-2 genome presence and IL-6 and TNF-α concentration as described above. Since samples were drawn for SARS-CoV-2 infection diagnostic purposes, an important aspect of this study consisted in the timing of cytokine concentration assessment, performed during the first 10 days of symptoms in most of these patients. The most prevalent symptoms included malaise, myalgia, headache, and fever. Characteristics of the study population are shown in Table 1.

*Cytokine analysis*

Normal ranges of IL-6 and TNF-α for swab samples were set up using a group of asymptomatic COVID-19 (-) individuals. In this group, IL-6 median value was 3.18 pg/ml (0.78-8.39) and TNF-α median value was 4.72 pg/ml (3.19-9.69).

IL-6 was increased in both groups of symptomatic patients compared to asymptomatic patients (Fig. 1a). However, the difference between COVID-19 (+) and (-) symptomatic patients, with median values of 5.40 (1.7-467) pg/ml and 6.07 (1.57-466.6) pg/ml respectively, did not reach statistical significance (Fig. 1a). Despite both groups of symptomatic patients showed higher TNF-α levels compared to the asymptomatic group, no significant differences were observed (Fig. 1b). However, TNF-α levels were greater in COVID (+) symptomatic group [5.88 (1.36-172.1) pg/ml] compared to COVID (-) symptomatic individuals [2.87 (1.45–69.9) pg/ml], displaying a better relationship among this cytokine local concentration and the presence of SARS-CoV-2 infection (p=0.0003) (Fig. 1b). Considering a threshold value of 10 pg/ml for TNF-α, the presence of SARS-CoV-2 was associated with increased levels of TNF-α (OR= 5.7; p=0.006; 95% CI=1,551 to 19,11). On the contrary, IL-6 values did not show any association with the infection.

Disease severity did not correlate with levels of IL-6 or TNF-α since median values in

hospitalized and outpatients were not significantly different. However, COVID-19 (+)

hospitalized individuals showed increased TNF-α values [median 4.89 pg/ml (1.4 – 172.2)]

compared to COVID (-) hospitalized patients [median 3.36 pg/ml (1.4 – 70)] (p=0.0138).

Among the group of COVID-19 (+) individuals, correlation between IL-6 or TNF-α levels and

viral load (Ct) was not found. Furthermore, older age and gender did not seem to be related

with higher values of IL-6 or TNF-α among the COVID-19 (+) group.

**Discussion**

In the present study, we evaluated the concentration of IL6 and TNF at the entry site of SARS-CoV-2. Although levels of IL-6 and TNF-α were both increased in some patients during the first stages of COVID-19, only TNF-α was significantly associated with SARS-CoV-2 presence. This result may indicate a distinct cytokine profile expressed at this site during the symptomatic phase of SARS-CoV-2 infection.

Early in the pandemic, patients with high systemic levels of IL-6 and TNF-α were correlated with an excessive and deleterious immune response driving to COVID-19-related morbidity and mortality (Costella-Ruiz et al. 2020; Del Valle et al. 2020; Herold et al. 2020; Robinson et al. 2020a; Danlos et al. 2021; Sabaka et al. 2021). Focusing on IL-6 as a possible marker to predict the progress of COVID-19, several large studies suggested that systemic IL-6 levels of more than 80 pg/mL were the best laboratory predictor of respiratory failure and death and many IL-6 inhibitors therapies have been approved for the treatment of COVID-19 (Chen et al. 2020; Hambali et al. 2020; Chen & Quach, 2021; NIH, 2021; Udomsinprasert et al. 2021). However, many authors argued that although IL-6 levels may suggest severity of responses, they do not necessarily imply pathogenesis (Zhang et al. 2020; Chen & Quach, 2021; NIH, 2021; Udomsinprasert et al. 2021). Furthermore, some studies reported low median IL-6 levels in severe COVID-19 patients when compared to other inflammatory conditions, such as acute respiratory distress syndrome (ARDS) and bacterial sepsis (Calfee et al. 2014; Sinha et al. 2018). Accordingly, we could not find a correlation between IL-6 or TNF concentration in respiratory samples and severity of disease (analyzed as the requirement of hospitalization). In addition, IL-6 and TNF-α normal values were present even in severe hospitalized COVID-19 patients.

Likewise, reports on anti-TNF-α use are suggestive of a therapeutic benefit in patients with COVID-19 (Robinson et al. 2020a; Robinson et al. 2020b). Observational data from patients already on anti-TNF therapy show a reduced rate of COVID-19 poor outcomes and death compared with other immune-suppressing therapies (Gianfrancesco et al. 2020;

Robinson et al. 2020a; Robinson et al. 2020b). Although randomized controlled trials are

needed, these results may indicate that early anti-TNF therapy could be beneficial in a group

of patients (Robinson et al. 2020b). In this study, we found a statistically significant

association between the production of local TNF-α and the presence of the virus in early

stages of infection. Therefore, measurement of local TNF-α, and probably some other

mediators, could contribute to identify those individuals that may benefit from

immunomodulatory and cytokine inhibiting therapies.

An increase of IL-6 and TNF-α and other cytokines in respiratory samples has been

already described in numerous respiratory viral infections and has been associated with a

worse prognosis of the disease (Seemungal et al. 2000; Patel et al. 2009; Ugonna et al.

2016; Vazquez et al. 2019). Our findings highlight the wide and pleiotropic role of IL-6 in

local inflammation process and show that TNF-α estimation and its association to viral

presence might be a valuable biomarker that could contribute to disease evolution

prognosis. Through our results, we could hypothesize that a proportion of individuals might

take advantage of the early and sequential measurement of cytokine concentration (at local

and systemic levels) acting as supplementary biomarkers for COVID-19.

**Limitations**

Because this study was done in samples obtained for COVID-19 diagnostic purposes,

simultaneous serum samples were not drawn at the time of swab collection. Therefore, no

comparison can be made with the systemic IL-6 or TNF-α values reported in other studies.

Lack of monitoring cytokines at different time points after diagnosis is another limitation of

our study.

241 **Funding:** Some aspects of this investigation could not have been fulfilled without the
242 generous contribution of the Fundación René Baron, Buenos Aires, who provided financial
243 support.

**Declaration of Author Contributions**

246 PB conceived the study; MNB and MJP designed the study protocol and carried out the
247 clinical assessment; PB made the analysis and interpretation of the database. PB, MNB and
248 MJP drafted the manuscript; MMEB and RC critically revised the manuscript for intellectual
249 content. All authors read and approved the final manuscript.

**Acknowledgements**

252 MJP was recipient of a Doctoral fellowship from the Consejo Nacional de Investigaciones
253 Científico y Tecnológicas, CONICET. MNB was recipient of a fellowship from the National
254 Academy of Medicine. PB is member of the National Research Council (CONICET)
255 Research Career Program.

**References**

1. World Healt Organization, 2021. Retrieved from: https://covid19.who.int/ (26 July 2021, date last accessed).

266
267

2. Hu, B., Guo, H., Zhou, P., & Shi, Z. L. (2021). Characteristics of SARS-CoV-2 and COVID-19. Nature reviews. Microbiology, 19(3), 141–154. https://doi.org/10.1038/s41579-020-00459-7

268
269
270

3. Mangalmurti, N., & Hunter, C. A. (2020). Cytokine Storms: Understanding COVID-19. Immunity, 53(1), 19–25. https://doi.org/10.1016/j.immuni.2020.06.017

271
272

4. Pedersen, S. F., & Ho, Y. C. (2020). SARS-CoV-2: a storm is raging. The Journal of clinical investigation, 130(5), 2202–2205. https://doi.org/10.1172/JCI137647

273
274

5. Costela-Ruiz, V. J., Illescas-Montes, R., Puerta-Puerta, J. M., Ruiz, C., & Melguizo-Rodríguez, L. (2020). SARS-CoV-2 infection: The role of cytokines in COVID-19 disease. Cytokine & growth factor reviews, 54, 62–75. https://doi.org/10.1016/j.cytogfr.2020.06.001

275
276
277
278

6. Del Valle, D. M., Kim-Schulze, S., Huang, H. H., Beckmann, N. D., Nirenberg, S., Wang, B., Lavin, Y., Swartz, T. H., Madduri, D., Stock, A., Marron, T. U., Xie, H., Patel, M., Tuballes, K., Van Oekelen, O., Rahman, A., Kovatch, P., Aberg, J. A., Schadt, E., Jagannath, S., … Gnjatic, S. (2020). An inflammatory cytokine signature predicts COVID-19 severity and survival. Nature medicine, 26(10), 1636–1643. https://doi.org/10.1038/s41591-020-1051-9

7. Herold, T., Jurinovic, V., Arnreich, C., Lipworth, B. J., Hellmuth, J. C., von Bergwelt-Baildon, M., Klein, M., & Weinberger, T. (2020). Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in COVID-19. The Journal of allergy and clinical immunology, 146(1), 128–136.e4. https://doi.org/10.1016/j.jaci.2020.05.008

285
286
287
288

8. Robinson, P. C., Richards, D., Tanner, H. L., & Feldmann, M. (2020)a. Accumulating evidence suggests anti-TNF therapy needs to be given trial priority in COVID-19

289
290

291 treatment. The Lancet. Rheumatology, 2(11), e653–e655.

292 https://doi.org/10.1016/S2665-9913(20)30309-X

293 9. Zhu, Z., Cai, T., Fan, L., Lou, K., Hua, X., Huang, Z., & Gao, G. (2020). Clinical value

294 of immune-inflammatory parameters to assess the severity of coronavirus disease

295 2019. International journal of infectious diseases : IJID : official publication of the

296 International Society for Infectious Diseases, 95, 332–339.

297 https://doi.org/10.1016/j.ijid.2020.04.041

298 10. Danlos, F. X., Ackermann, F., Rohmer, J., Roumier, M., Marabelle, A., & Michot, J.

299 M. (2021). High levels of TNFα in patients with COVID-19 refractory to tocilizumab.

300 European journal of cancer (Oxford, England : 1990), 149, 102–104.

301 https://doi.org/10.1016/j.ejca.2021.01.056

302 11. Sabaka, P., Koščálová, A., Straka, I., Hodosy, J., Lipták, R., Kmotorková, B.,

303 Kachlíková, M., & Kušnírová, A. (2021). Role of interleukin 6 as a predictive factor for

304 a severe course of Covid-19: retrospective data analysis of patients from a long-term

305 care facility during Covid-19 outbreak. BMC infectious diseases, 21(1), 308.

306 https://doi.org/10.1186/s12879-021-05945-8

307 12. Chen, L., Hoiland, R. L., Stukas, S., Wellington, C. L., & Sekhon, M. S. (2020).

308 Confronting the controversy: interleukin-6 and the COVID-19 cytokine storm

309 syndrome. *The European respiratory journal*, *56*(4), 2003006.

310 https://doi.org/10.1183/13993003.03006-2020

311 13. Hambali, N. L., Mohd Noh, M., Paramasivam, S., Chua, T. H., Hayati, F., Payus, A.

312 O., Tee, T. Y., Rosli, K. T., Abd Rachman Isnadi, M. F., & Manin, B. O. (2020). A

313 Non-severe Coronavirus Disease 2019 Patient With Persistently High Interleukin-6

314 Level. *Frontiers in public health*, *8*, 584552.

315 https://doi.org/10.3389/fpubh.2020.584552

316 14. Chen, L., & Quach, T. (2021). COVID-19 cytokine storm syndrome: a threshold

317 concept. The Lancet. Microbe, 2(2), e49–e50. https://doi.org/10.1016/S2666-

318 5247(20)30223-8

15. Udomsinprasert, W., Jittikoon, J., Sangroongruangsri, S., & Chaikledkaew, U. (2021).

Circulating Levels of Interleukin-6 and Interleukin-10, But Not Tumor Necrosis Factor-

Alpha, as Potential Biomarkers of Severity and Mortality for COVID-19: Systematic

Review with Meta-analysis. Journal of clinical immunology, 41(1), 11–22.

https://doi.org/10.1007/s10875-020-00899-z

16. National        Institute        of        Health,        2021.        Retrieved        from:

https://www.covid19treatmentguidelines.nih.gov/therapies/immunomodulators/interle

ukin-6-inhibitors/ (14 June 2021, date last accessed).

17. Zhang, Y., Zhong, Y., Pan, L., & Dong, J. (2020). Treat 2019 novel coronavirus

(COVID-19) with IL-6 inhibitor: Are we already that far?. Drug discoveries &

therapeutics, 14(2), 100–102. https://doi.org/10.5582/ddt.2020.03006

18. Calfee, C. S., Delucchi, K., Parsons, P. E., Thompson, B. T., Ware, L. B., Matthay,

M. A., & NHLBI ARDS Network (2014). Subphenotypes in acute respiratory distress

syndrome: latent class analysis of data from two randomised controlled trials. *The

Lancet.   Respiratory   medicine*, *2*(8),   611–620.   https://doi.org/10.1016/S2213-

2600(14)70097-9

19. Sinha, P., Delucchi, K. L., Thompson, B. T., McAuley, D. F., Matthay, M. A., Calfee,

C. S., & NHLBI ARDS Network (2018). Latent class analysis of ARDS

subphenotypes: a secondary analysis of the statins for acutely injured lungs from

sepsis   (SAILS)   study. *Intensive   care   medicine*, *44*(11),   1859–1869.

https://doi.org/10.1007/s00134-018-5378-3

20. Gianfrancesco, M., Hyrich, K. L., Al-Adely, S., Carmona, L., Danila, M. I., Gossec, L.,

Izadi, Z., Jacobsohn, L., Katz, P., Lawson-Tovey, S., Mateus, E. F., Rush, S.,

Schmajuk, G., Simard, J., Strangfeld, A., Trupin, L., Wysham, K. D., Bhana, S.,

Costello, W., Grainger, R., … COVID-19 Global Rheumatology Alliance (2020).

Characteristics associated with hospitalisation for COVID-19 in people with rheumatic

disease: data from the COVID-19 Global Rheumatology Alliance physician-reported

registry. Annals of the rheumatic diseases, 79(7), 859–866.

https://doi.org/10.1136/annrheumdis-2020-217871

21. Robinson, P. C., Liew, D., Liew, J. W., Monaco, C., Richards, D., Shivakumar, S.,

Tanner, H. L., & Feldmann, M. (2020)b. The Potential for Repurposing Anti-TNF as a

Therapy for the Treatment of COVID-19. Med (New York, N.Y.), 1(1), 90–102.

https://doi.org/10.1016/j.medj.2020.11.005

22. Seemungal, T. A., Harper-Owen, R., Bhowmik, A., Jeffries, D. J., & Wedzicha, J. A.

(2000). Detection of rhinovirus in induced sputum at exacerbation of chronic

obstructive pulmonary disease. The European respiratory journal, 16(4), 677–683.

https://doi.org/10.1034/j.1399-3003.2000.16d19.x

23. Patel, J. A., Nair, S., Revai, K., Grady, J., & Chonmaitree, T. (2009). Nasopharyngeal

acute phase cytokines in viral upper respiratory infection: impact on acute otitis

media in children. The Pediatric infectious disease journal, 28(11), 1002–1007.

https://doi.org/10.1097/INF.0b013e3181aa5b13

24. Ugonna, K., Douros, K., Bingle, C. D., & Everard, M. L. (2016). Cytokine responses

in primary and secondary respiratory syncytial virus infections. Pediatric research,

79(6), 946–950. https://doi.org/10.1038/pr.2016.29

25. Vázquez, Y., González, L., Noguera, L., González, P. A., Riedel, C. A., Bertrand, P.,

& Bueno, S. M. (2019). Cytokines in the Respiratory Airway as Biomarkers of

Severity and Prognosis for Respiratory Syncytial Virus Infection: An Update.

Frontiers in immunology, 10, 1154. https://doi.org/10.3389/fimmu.2019.01154

372    **Table 1** Population characteristics

|  | COVID-19 (+) Symptomatic pts | COVID-19 (-) Symptomatic pts | COVID-19 (-) Asymptomatic pts |
|---|---|---|---|
| **Age (years)** | 48,5 (0,16-93) | 56 (0,16-92) | 45 (0,7-93) |
| **Gender (F/M)** | 26/27 | 15/18 | 22/21 |
| **Hospitalized pts (n)** | 26 | 29 | 0 |
| **SARS-CoV-2 viral load (Ct)** | 29 (14-39) | na | na |
| **Days of symptoms** | 4 (1-11) | 2 (0-10) | na |

374    Values are expressed as median (range) unless otherwise noted.
375    *pts* patients, *F* female, *M* male, *Ct* cycle threshold , *na* not applicable
376

392 <u>Figure Captions</u>

393 **Fig.1**: IL-6 (A) and TNF-α (B) concentrations (log pg/ml) in swab samples from symptomatic

394 COVID-19 (+) and COVID-19 (-) symptomatic individuals.

395