Title Page:

Title: TNF-α levels in respiratory samples are associated with SARS-CoV-2 infection.

Running title: Cytokine levels in respiratory samples.

Authors:

Matias J. PERESON[a,b] [*], Maria Noel BADANO[a] [*], Natalia ALOISI[a], Roberto CHUIT[c], MME de

BRACCO[a], Patricia BARE[a]

Affiliations:

[a]Instituto de Medicina Experimental (IMEX) CONICET, Instituto de Investigaciones

Hematologicas (IIHEMA), Academia Nacional de Medicina, Buenos Aires, Argentina

[b]Universidad de Buenos Aires. Facultad de Farmacia y Bioquímica. Instituto de

Investigaciones en Bacteriología y Virología Molecular (IBaViM). Buenos Aires, Argentina.

[c] Instituto de Investigaciones Epidemiológicas, Academia Nacional de Medicina.

*Contributed equally to this work.

Corresponding Author:

Dra. Patricia Bare, Instituto de Medicina Experimental (IMEX), CONICET, Instituto de

Investigaciones Hematologicas (IIHEMA), Academia Nacional de Medicina, Buenos Aires,

Argentina.

Tel: +541148055759 ext 269; Tel: +5491154975179;

Fax: +541148050712, E-mail: pbare@hematologia.anm.edu.ar; patobare@hotmail.com

ABSTRACT

Purpose: Increased serum levels of IL-6 and TNF-α have been proposed as biomarkers for COVID-19 progression. However, the role and the implication of these cytokines in SARS-CoV-2 infection remain controversial. The aim of this study was to measure levels of IL-6 and TNF-α in swab samples from individuals with symptoms compatible with COVID-19 and analyze their association with SARS-CoV-2 presence.

Methods: SARS-CoV-2 detection was performed using the CDC (USA) real-time RT-PCR primers, probes and protocols. Cytokine concentrations were measured using commercial reagents based on enzyme linked immunosorbent assay (ELISA).

Results: TNF-α median levels were greater in COVID19 (+) symptomatic group (5.88 (1.36 - 172.1) pg/ml) compared to COVID19 (-) symptomatic individuals (2.87 (1.45 – 69.9) pg/ml) (p=0.0003). No significant differences were shown in IL-6 median values between COVID-19 (+) and (-) symptomatic patients (5.40 (1.7 - 467) pg/ml and 6.07 (1.57 – 466.6) pg/ml respectively). In addition, increased TNF-α levels (greater than 10 pg/ml), but not IL-6, were associated with SARS-CoV-2 presence (OR= 5.7; p=0.006; 95% CI= 1,551 to 19,11).

Conclusions: IL-6 concentration showed high levels in swabs from some symptomatic patients, suggesting the presence of immune response at viral entry site. However, IL-6 levels were independent from SARS-CoV-2 presence and viral load, individual's age and gender. On the contrary, TNF-α evaluation confirmed the presence of inflammatory response but mostly related to COVID-19. More studies are required in order to characterize the cytokine profile expressed at the site of infection of SARS-CoV-2 and its implications in disease outcomes.

Keywords: SARS-COV-2; COVID-19; respiratory samples; Interleukin-6; TNF-α

Introduction

The coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is one of the current major health concerns. After 15 months since the emergence of the pandemic SARS-CoV-2, up to 178 million cases were confirmed, including more than 3.8 million deaths worldwide due to severe forms of COVID-19 [1]. Severe COVID-19 is widely considered an immunopathology disease driven by an excessive and deleterious immune response mounted against the pathogen rather than by the action of the pathogen *per se* [2-4]. Many components of the innate immune system have been associated with this phenomenon and have been proposed as biomarker for COVID-19 progression. In this sense, high levels of serum interleukin-6 (IL-6) and tumor necrosis factor alpha (TNF-α) were correlated with increased morbidity and mortality in COVID-19 patients [5-11]. However, innate immune mediators are necessary for efficient clearance of infectious agents. On the other hand, reports of lower median IL-6 levels in severe COVID-19 compared with other inflammatory conditions, such as acute respiratory distress syndrome and bacterial sepsis were published [12-14]. Additionally, many studies found no significant differences in TNF-α levels between severe and non-severe COVID-19 cases [15]. Therefore, the role of these cytokines in COVID-19 progression remains contradictory and needs further study.

To our knowledge, IL-6 and TNF-α concentrations at the site of infection in COVID-19 patients has not been previously reported. Therefore, the aim of this study was to evaluate IL-6 and TNF-α concentrations in swab samples from individuals showing symptoms compatible with COVID-19 who were positive or negative for SARS-CoV-2 genome detection.

Material and methods

Patients and study design

The study was conducted on 127 archived swab samples referred to the Academia Nacional de Medicina for diagnosis of SARS-CoV-2 by real-time polymerase reaction (RT-PCR) between August 2020 and April 2021. Samples were obtained during the initial stages of disease progression (1 to 10 days). Symptomatic individuals with detectable (n=52) or undetectable (n=33) SARS-CoV-2 infection were analyzed. A group consisting of asymptomatic individuals and negative for SARSCoV-2 genome (n=42) were included to obtain cytokines' normal range.

Collection of samples

Combined nasopharyngeal and oropharyngeal swabs were collected by trained healthcare staff and placed into a single tube pre-filled with 2 mL of saline solution (0,9% sodium chloride) and were transferred on the same day to the laboratory for SARS-CoV-2 genome detection. Samples were stored at −80 °C until cytokines assays were performed.

SARS-CoV-2 genome detection and viral load

Genomic extraction was carried out with the QIAamp Viral RNA minikit (Qiagen) following manufacturer's instructions. A real time-based methodology was performed for viral genome detection using the CDC RT-qPCR protocol for Wuhan virus (nCoV-19), 2019-nCov CDC USA. The procedure incorporates a set of oligonucleotides primers and double-marker hydrolysis probes (Taqman®) (2019 nCov_N1 and N2) that amplify regions of the viral nucleocapsid gene (N). An internal control for RNase P (RP) was also included. Viral load was calculated considering Ct values.

IL-6 and TNF-α measurement

Concentrations of IL-6 (standard curve range: 0-300 pg/mL) and TNF-α (standard curve

range: 0 -500 pg/mL) in swabs were determined using commercial reagents based on enzyme

linked immunosorbent assay (ELISA) (BD-Biosciences, San Diego, California, United States).

Recommendations of the supplier were followed for procedure. Duplicates were performed in

selected samples to verify the accuracy of the results.

Statistical analyses

Mann-Whitney U test was used for testing differences between two groups. Categorical

variables were compared with Chi-square test or Fisher's exact test. Confidence intervals were

set at 95% (CI95). In all cases, a p value <0.05 was considered significant. Data and graphs

were performed using the GraphPad Prism 9.1.0 software (GraphPad Software, San Diego,

CA, USA).

Ethics approval

Experimental protocols and procedures performed in this work have been approved by the

Biosafety Review board of the IMEX-CONICET-Academia Nacional de Medicina and the

Ethical Committee of the Academia Nacional de Medicina.

Results

Study population

Combined oro- and nasopharyngeal swabs from 127 patients submitted for diagnosis of COVID-19 were examined for SARS-CoV-2 genome presence and IL-6 and TNF-α concentration as described above. Since samples were drawn for SARS-CoV-2 infection diagnostic purposes, an important aspect of this study consisted in the timing of cytokine concentration assessment, performed during the first 10 days of symptoms in most of these patients. The most prevalent symptoms included malaise, myalgia, headache and fever. Characteristics of the study population are shown in Table 1.

Cytokines analysis

Normal ranges of IL-6 and TNF-α for swab samples were set up using a group of asymptomatic COVID-19 (-) individuals. In this group, IL-6 median value was 3.18 pg/ml (0.78-8.39) and TNF-α median value was 4.72 pg/ml (3.19-9.69).

IL-6 was increased in both groups of symptomatic patients compared to asymptomatic patients (Fig. 1a). However, the difference between COVID-19 (+) and (-) symptomatic patients, with median values of 5.40 (1.7-467) pg/ml and 6.07 (1.57-466.6) pg/ml respectively, did not reach statistical significance (Fig. 1a). Despite both groups of symptomatic patients showed higher TNF-α levels compared to the asymptomatic group, no significant differences were observed (Fig. 1b). However, TNF-α levels were greater in COVID (+) symptomatic group [5.88 (1.36-172.1) pg/ml] compared to COVID (-) symptomatic individuals [2.87 (1.45–69.9) pg/ml], displaying a better relationship among this cytokine local concentration and the presence of SARS-CoV-2 infection (p=0.0003) (Fig. 1b). Considering a threshold value of 10 pg/ml for TNF-α, the presence of SARS-CoV-2 was associated with increased levels of TNF-α (OR= 5.7; p=0.006; 95% CI=1,551 to 19,11). On the contrary, IL-6 values did not show any association to the infection.

Disease severity did not correlate with levels of IL-6 or TNF-α since median values in hospitalized and outpatients were not significantly different. However, COVID-19 (+) hospitalized individuals showed increased TNF-α values [median 4.89 pg/ml (1.4 – 172.2)] compared to COVID (-) hospitalized patients [median 3.36 pg/ml (1.4 – 70)] (p=0.0138). Among the group of COVID-19 (+) individuals, correlation between IL-6 or TNF-α levels and viral load (Ct) was not found. Furthermore, older age and gender did not seem to be related with higher values of IL-6 or TNF-α among the COVID-19 (+) group.

Discussion

In the present study, we evaluated the concentration of IL6 and TNF at the entry site of SARS-CoV-2. Although levels of IL-6 and TNF-α were both increased in some patients during the first stages of COVID-19, only TNF-α was significantly associated with SARS-CoV-2 presence. This result may indicate a distinct cytokine profile expressed at this site during the symptomatic phase of SARS-CoV-2 infection.

Early in the pandemic, patients with high systemic levels of IL-6 and TNF-α were correlated with an excessive and deleterious immune response driving to COVID-19-related morbidity and mortality [5-8,10,11]. Focusing on IL-6 as a possible marker to predict the progress of COVID-19, several large studies suggested that systemic IL-6 levels of more than 80 pg/mL were the best laboratory predictor of respiratory failure and death and many IL-6 inhibitors therapies have been approved for the treatment of COVID-19 [12-16]. However, many authors argued that although IL-6 levels may suggest severity of responses, they do not necessarily imply pathogenesis [14,17]. Furthermore, some studies reported low median IL-6 levels in severe COVID-19 patients when compared to other inflammatory conditions, such as acute respiratory distress syndrome (ARDS) and bacterial sepsis [18,19]. Accordingly, we could not find a correlation between IL-6 or TNF concentration in respiratory samples and severity of disease (analyzed as the requirement of hospitalization). In addition, IL-6 and TNF-α normal values were present even in severe hospitalized COVID-19 patients.

Likewise, reports on anti-TNF-α use are suggestive of a therapeutic benefit in patients with COVID-19 [8,21]. Observational data from patients already on anti-TNF therapy show a reduced rate of COVID-19 poor outcomes and death compared with other immune-suppressing therapies [8,20,21]. Although randomized controlled trials are needed, these results may indicate that early anti-TNF therapy could be beneficial in a group of patients [21]. In this study, we found a statistically significant association between the production of local TNF-α and the presence of the virus in early stages of infection. Therefore, measurement of

local TNF-α, and probably some others mediators, could contribute to identify those individuals

that may benefit from immunomodulatory and cytokine inhibiting therapies.

An increase of IL-6 and TNF-α and other cytokines in respiratory samples has been already

described in numerous respiratory viral infections and has been associated with a worse

prognosis of the disease [22,25]. Our findings highlight the wide and pleiotropic role of IL-6 in

local inflammation process and show that TNF-α estimation and its association to viral

presence might be a valuable biomarker that could contribute to disease evolution prognosis.

Through our results, we could hypothesize that a proportion of individuals might take

advantage of the early and sequential measurement of cytokine concentration (at local and

systemic levels) acting as supplementary biomarkers for COVID-19.

Limitations

Because this study was done in samples obtained for COVID-19 diagnostic purposes,

simultaneous serum samples were not drawn at the time of swab collection. Therefore, no

comparison can be made with the systemic IL-6 or TNF-α values reported in other studies.

Lack of monitoring cytokines at different time points after diagnosis is another limitation of our

study.

Funding: Some aspects of this investigation could not have been fulfilled without the generous

contribution of the Fundación René Baron, Buenos Aires, who provided financial support.

Competing interest: All authors have read and approved the final manuscript. The authors

have no conflicts of interest to declare.

Declaration of Author Contributions

PB conceived the study; MNB and MJP designed the study protocol and carried out the clinical

assessment; PB made the analysis and interpretation of the database. PB, MNB and MJP

drafted the manuscript; MMEB and RC critically revised the manuscript for intellectual content.

All authors read and approved the final manuscript.

Acknowledgements

MJP was recipient of a Doctoral fellowship from the Consejo Nacional de Investigaciones

Científico y Tecnológicas, CONICET. MNB was recipient of a fellowship from the National

Academy of Medicine. PB is member of the National Research Council (CONICET) Research

Career Program.

References

1. World Healt Organization, 2021. Retrieved from: https://covid19.who.int/ (22 June 2021, date last accessed).

2. Hu, B., Guo, H., Zhou, P., & Shi, Z. L. (2021). Characteristics of SARS-CoV-2 and COVID-19. Nature reviews. Microbiology, 19(3), 141–154. https://doi.org/10.1038/s41579-020-00459-7

3. Mangalmurti, N., & Hunter, C. A. (2020). Cytokine Storms: Understanding COVID-19. Immunity, 53(1), 19–25. https://doi.org/10.1016/j.immuni.2020.06.017

4. Pedersen, S. F., & Ho, Y. C. (2020). SARS-CoV-2: a storm is raging. The Journal of clinical investigation, 130(5), 2202–2205. https://doi.org/10.1172/JCI137647

5. Costela-Ruiz, V. J., Illescas-Montes, R., Puerta-Puerta, J. M., Ruiz, C., & Melguizo-Rodríguez, L. (2020). SARS-CoV-2 infection: The role of cytokines in COVID-19 disease. Cytokine & growth factor reviews, 54, 62–75. https://doi.org/10.1016/j.cytogfr.2020.06.001

6. Del Valle, D. M., Kim-Schulze, S., Huang, H. H., Beckmann, N. D., Nirenberg, S., Wang, B., Lavin, Y., Swartz, T. H., Madduri, D., Stock, A., Marron, T. U., Xie, H., Patel, M., Tuballes, K., Van Oekelen, O., Rahman, A., Kovatch, P., Aberg, J. A., Schadt, E., Jagannath, S., … Gnjatic, S. (2020). An inflammatory cytokine signature predicts COVID-19 severity and survival. Nature medicine, 26(10), 1636–1643. https://doi.org/10.1038/s41591-020-1051-9

7. Herold, T., Jurinovic, V., Arnreich, C., Lipworth, B. J., Hellmuth, J. C., von Bergwelt-Baildon, M., Klein, M., & Weinberger, T. (2020). Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in COVID-19. The Journal of allergy and clinical immunology, 146(1), 128–136.e4. https://doi.org/10.1016/j.jaci.2020.05.008

8. Robinson, P. C., Richards, D., Tanner, H. L., & Feldmann, M. (2020)a. Accumulating evidence suggests anti-TNF therapy needs to be given trial priority in COVID-19

treatment. The Lancet. Rheumatology, 2(11), e653–e655.
https://doi.org/10.1016/S2665-9913(20)30309-X

9. Zhu, Z., Cai, T., Fan, L., Lou, K., Hua, X., Huang, Z., & Gao, G. (2020). Clinical value
of immune-inflammatory parameters to assess the severity of coronavirus disease
2019. International journal of infectious diseases : IJID : official publication of the
International Society for Infectious Diseases, 95, 332–339.
https://doi.org/10.1016/j.ijid.2020.04.041

10. Danlos, F. X., Ackermann, F., Rohmer, J., Roumier, M., Marabelle, A., & Michot, J. M.
(2021). High levels of TNFα in patients with COVID-19 refractory to tocilizumab.
European journal of cancer (Oxford, England : 1990), 149, 102–104.
https://doi.org/10.1016/j.ejca.2021.01.056

11. Sabaka, P., Koščálová, A., Straka, I., Hodosy, J., Lipták, R., Kmotorková, B.,
Kachlíková, M., & Kušnírová, A. (2021). Role of interleukin 6 as a predictive factor for
a severe course of Covid-19: retrospective data analysis of patients from a long-term
care facility during Covid-19 outbreak. BMC infectious diseases, 21(1), 308.
https://doi.org/10.1186/s12879-021-05945-8

12. Chen, L., Hoiland, R. L., Stukas, S., Wellington, C. L., & Sekhon, M. S. (2020).
Confronting the controversy: interleukin-6 and the COVID-19 cytokine storm
syndrome. *The European respiratory journal*, *56*(4), 2003006.
https://doi.org/10.1183/13993003.03006-2020

13. Hambali, N. L., Mohd Noh, M., Paramasivam, S., Chua, T. H., Hayati, F., Payus, A. O.,
Tee, T. Y., Rosli, K. T., Abd Rachman Isnadi, M. F., & Manin, B. O. (2020). A Non-
severe Coronavirus Disease 2019 Patient With Persistently High Interleukin-6
Level. *Frontiers in public health*, *8*, 584552.
https://doi.org/10.3389/fpubh.2020.584552

14. Chen, L., & Quach, T. (2021). COVID-19 cytokine storm syndrome: a threshold
concept. The Lancet. Microbe, 2(2), e49–e50. https://doi.org/10.1016/S2666-
5247(20)30223-8

15. Udomsinprasert, W., Jittikoon, J., Sangroongruangsri, S., & Chaikledkaew, U. (2021). Circulating Levels of Interleukin-6 and Interleukin-10, But Not Tumor Necrosis Factor-Alpha, as Potential Biomarkers of Severity and Mortality for COVID-19: Systematic Review with Meta-analysis. Journal of clinical immunology, 41(1), 11–22. https://doi.org/10.1007/s10875-020-00899-z

16. National Institute of Health, 2021. Retrieved from: https://www.covid19treatmentguidelines.nih.gov/therapies/immunomodulators/interleukin-6-inhibitors/ (14 June 2021, date last accessed).

17. Zhang, Y., Zhong, Y., Pan, L., & Dong, J. (2020). Treat 2019 novel coronavirus (COVID-19) with IL-6 inhibitor: Are we already that far?. Drug discoveries & therapeutics, 14(2), 100–102. https://doi.org/10.5582/ddt.2020.03006

18. Calfee, C. S., Delucchi, K., Parsons, P. E., Thompson, B. T., Ware, L. B., Matthay, M. A., & NHLBI ARDS Network (2014). Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. *The Lancet. Respiratory medicine*, *2*(8), 611–620. https://doi.org/10.1016/S2213-2600(14)70097-9

19. Sinha, P., Delucchi, K. L., Thompson, B. T., McAuley, D. F., Matthay, M. A., Calfee, C. S., & NHLBI ARDS Network (2018). Latent class analysis of ARDS subphenotypes: a secondary analysis of the statins for acutely injured lungs from sepsis (SAILS) study. *Intensive care medicine*, *44*(11), 1859–1869. https://doi.org/10.1007/s00134-018-5378-3

20. Gianfrancesco, M., Hyrich, K. L., Al-Adely, S., Carmona, L., Danila, M. I., Gossec, L., Izadi, Z., Jacobsohn, L., Katz, P., Lawson-Tovey, S., Mateus, E. F., Rush, S., Schmajuk, G., Simard, J., Strangfeld, A., Trupin, L., Wysham, K. D., Bhana, S., Costello, W., Grainger, R., … COVID-19 Global Rheumatology Alliance (2020). Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: data from the COVID-19 Global Rheumatology Alliance physician-reported

registry. Annals of the rheumatic diseases, 79(7), 859–866.
https://doi.org/10.1136/annrheumdis-2020-217871

21. Robinson, P. C., Liew, D., Liew, J. W., Monaco, C., Richards, D., Shivakumar, S.,
Tanner, H. L., & Feldmann, M. (2020)b. The Potential for Repurposing Anti-TNF as a
Therapy for the Treatment of COVID-19. Med (New York, N.Y.), 1(1), 90–102.
https://doi.org/10.1016/j.medj.2020.11.005

22. Seemungal, T. A., Harper-Owen, R., Bhowmik, A., Jeffries, D. J., & Wedzicha, J. A.
(2000). Detection of rhinovirus in induced sputum at exacerbation of chronic
obstructive pulmonary disease. The European respiratory journal, 16(4), 677–683.
https://doi.org/10.1034/j.1399-3003.2000.16d19.x

23. Patel, J. A., Nair, S., Revai, K., Grady, J., & Chonmaitree, T. (2009). Nasopharyngeal
acute phase cytokines in viral upper respiratory infection: impact on acute otitis media
in children. The Pediatric infectious disease journal, 28(11), 1002–1007.
https://doi.org/10.1097/INF.0b013e3181aa5b13

24. Ugonna, K., Douros, K., Bingle, C. D., & Everard, M. L. (2016). Cytokine responses in
primary and secondary respiratory syncytial virus infections. Pediatric research, 79(6),
946–950. https://doi.org/10.1038/pr.2016.29

25. Vázquez, Y., González, L., Noguera, L., González, P. A., Riedel, C. A., Bertrand, P.,
& Bueno, S. M. (2019). Cytokines in the Respiratory Airway as Biomarkers of Severity
and Prognosis for Respiratory Syncytial Virus Infection: An Update. Frontiers in
immunology, 10, 1154. https://doi.org/10.3389/fimmu.2019.01154

 Tables

371 **Table 1** Population characteristics

	COVID-19 (+) Symptomatic pts	COVID-19 (-) Symptomatic pts	COVID-19 (-) Asymptomatic pts
Age (years)	48,5 (0,16-93)	56 (0,16-92)	45 (0,7-93)
Gender (F/M)	26/27	15/18	22/21
Hospitalized pts (n)	26	29	0
SARS-CoV-2 viral load (Ct)	29 (14-39)	na	na
Days of symptoms	4 (1-11)	2 (0-10)	na

373 Values are expressed as median (range) unless otherwise noted.
374 *pts* patients, *F* female, *M* male, *Ct* cycle threshold , *na* not applicable
375

377 **Figure Captions**

378 **Fig.1**: IL-6 (A) and TNF-α (B) concentrations (log pg/ml) in swab samples from symptomatic

379 COVID-19 (+) and COVID-19 (-) symptomatic individuals.

15