BREAKTHROUGH REPORT

Image-based assessment of plant disease progression identifies new genetic loci for resistance

Valérian Méline[a], Denise L. Caldwell[a], Bong-Suk Kim[a], Sriram Baireddy[b], Changye Yang[b], Erin E. Sparks[c], Edward J. Delp[b], and Anjali S. Iyer-Pascuzzi[a†]

[a] 915 W. State Street, Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana, USA
[b] Video and Image Processing Laboratory (VIPER), School of Electrical and Computer Engineering
[c] Department of Plant and Soil Sciences and the Delaware Biotechnology Institute, University of Delaware, Newark, DE

†Author for correspondence: asi2@purdue.edu

Keywords: Digital phenotyping, QTL, *Ralstonia*, bacterial wilt, tomato

Short title: Disease resistance QTL and image-based phenotyping

Abstract

A major challenge in global crop production is mitigating yield loss due to plant diseases. One of

the best means of disease control is plant resistance, but the identification of genes that promote

resistance has been limited by the subjective quantification of disease, which is typically scored

by the human eye. We hypothesized that image-based, non-destructive quantification of disease

phenotypes would enable the rapid identification of new disease resistance loci. We tested this

using the interaction between tomato and *Ralstonia solanacearum*, a soilborne pathogen that

causes bacterial wilt disease. We acquired over 40,000 time-series images of disease progression

in a tomato recombinant inbred line population, and developed an image analysis pipeline

providing a suite of ten traits to quantify wilt disease based on plant shape and size. Quantitative

trait loci (QTL) analyses using image-based phenotyping identified QTL that were both unique

and shared compared with those identified by human assessment of wilting. When shared loci

were identified, image-based phenotyping could detect some QTL several days earlier than

human assessment. Thus, expanding the phenotypic space of disease with image-based, non-

destructive phenotyping allowed both earlier detection and identified new genetic components of

resistance.

Introduction

Plant diseases are a significant global constraint to crop production. Developing disease resistant crops requires identifying the plant genomic regions and genes that contribute to resistance to the pathogenic microbes that cause disease. This in turn depends on phenotyping large populations of plants for their responses to pathogens. Phenotyping plant diseases is challenging because diseases cause complex, quantitative phenotypes that can occur at different scales – e.g. on parts of leaves, entire leaves, or the whole plant. In addition, disease phenotypes vary over time and depend on environmental conditions, plant age, and pathogen virulence. Disease symptoms such as wilting or necrotic spots have traditionally been scored with the human eye, but these scores are subjective, can vary by individual, and are difficulty to accurately quantify.

The challenging nature of visual disease assessment has led to the use of sensors including RGB, hyperspectral, chlorophyll fluorescence and thermal cameras to assess disease symptoms (Colwell, 1956; Jackson, 1986; Bock et al., 2010; Simko et al., 2017). Compared to assessment by the human eye, image-based phenotyping is faster, more reproducible, and more sensitive to small variations in disease symptoms that can be critical for detecting resistance loci (Bock et al., 2008, 2010; Stewart and McDonald, 2014; Stewart et al., 2016; Simko et al., 2017; Shakoor et al., 2017). Many studies have used or developed tools to assess plant symptoms using different types of sensors (Mahlein, 2016; Mahlein et al., 2017, 2019; Lowe et al., 2017; Shakoor et al., 2017; Mochida et al., 2019; Mir et al., 2019; Pérez-Bueno et al., 2019; Pineda et al., 2021; Simko et al., 2017). However, few studies have used these technologies in QTL or Genome Wide Association (GWA) analyses for responses to plant pathogens, and all have used destructive methods (Yates et al., 2019; Fordyce et al., 2018; Corwin et al., 2016). It has remained challenging to use image-based, non-destructive phenotyping for disease resistance across large populations, both because of technical factors like the expense of phenotyping platforms and the time associated with imaging, and also biological factors such as differences in plant morphology and disease progression within a population.

The soil-borne betaproteobacterium *Ralstonia solanacearum* is the causal agent of bacterial wilt disease and has been ranked as one of the top 10 most destructive plant bacterial pathogens of all time (Mansfield et al., 2012). *Ralstonia* infection causes susceptible plants to wilt, and the

amount of wilting correlates with a plant's level of susceptibility (Genin, 2010; Genin and

Denny, 2011). The bacterium is a major production constraint in Solanaceous crops both

globally and in the United States, where disease loss in tomatoes can exceed 70%. In crops,

resistance to *Ralstonia* is quantitative, but the quantitative trait loci (QTL) underlying resistance

to US strains of *Ralstonia* are largely unknown. QTL for other strains have been mapped

(Danesh et al., 1994; Thoquet et al., 1996a, 1996b; Mangin et al., 1999; Wang et al., 2000;

Carmeille et al., 2006; Jaw-Fen Wang et al., 2013; Shin et al., 2020) but have not been cloned,

and the host determinants necessary for resistance remain mostly unspecified.

The limited identification of QTL for *Ralstonia* resistance can be attributed in part to the

difficulty in accurately scoring plant wilting. Wilting is traditionally measured on a $0 - 4$ scale,

in which 0 indicates a plant with no wilting, $1 = 1 - 25\%$, $2 = 26 - 50\%$, $3 = 51 - 75\%$ and a

score of 4 indicates a plant with $76 - 100\%$ wilted leaves (Schandry, 2017). While it is

straightforward to assess the ends of the spectrum, rating plants with scores of 2 or 3 is

particularly difficult. This is due to the subjective nature of visually determining when a leaf has

lost sufficient turgor to qualify as wilted. Reliable disease phenotyping is critical for identifying

QTL for resistance to *Ralstonia* and the development of resistant varieties.

Here, we used image-based, rapid, non-destructive phenotyping to identify new tomato genetic

resistance loci to *Ralstonia*. We developed a rapid, semi-automated imaging and trait analysis

pipeline to quantify bacterial wilt disease in a recombinant inbred line (RIL) population derived

from *Ralstonia*-resistant and susceptible tomato genotypes. We found both unique and shared

QTL between our image-based traits and plant wilting scored by the human eye. At least one of

the QTL was detected by image-based phenotyping before the onset of visual symptoms,

demonstrating that image-based phenotyping captures the disease phenotype at early stages of

infection. These results demonstrate that imaged-based, non-destructive phenotyping can shed

light on new aspects of disease and improve our ability to identify genetic loci crop resistance.

Results

Development of an aboveground imaging and semi-automated analysis pipeline

We first constructed a simple, low-cost imaging system that allowed us to semi-automate

aboveground disease phenotyping. Each plant was placed on a commercially available turntable,

and plants were imaged with a Canon DSLR (Supplemental Figure 1; details in methods). The

turntable and camera were connected with Photocapture 360 (Ortery technologies), which

allowed us to automatically capture images every 45 degrees (8 images per plant). Using this

system, we were able to non-destructively image each plant in less than 2 minutes, with minimal

manual labor. Each image included a fiducial marker for post-image color correction. Plants in

the F_9 generation from a RIL population derived from a cross between resistant Hawaii 7996

(H7996) and susceptible West Virginia 700 (WV) were imaged the day before inoculation with

R. solanacearum strain K60, and at 3, 4, 5 and 6 days post inoculation (dpi). At 3 dpi, symptoms

were not present in susceptible parent WV, but by 6 dpi these plants were completely wilted. We

imaged five replicate plants of each RIL as well as the parental lines. Using this system, we

captured over 40,000 images for high-resolution disease phenotyping. The same set of plants was

also visually scored by the human eye at 8 dpi. Visual scoring was based on the percentage of

wilted leaves. Tomato phenotypes in the RIL population ranged from highly susceptible to

highly resistant, consistent with the quantitative nature of disease resistance (Supplemental

Figures 2 and 3).

We next developed a set of mathematical descriptors to phenotype wilting over time in our

images. Plant wilting is a composite phenotype, and we used 10 image-based traits

(Supplemental Table 1) to describe different aspects of wilting: convex area, convex width, plant

area, plant height, plant width, X mass, Y mass, center of mass (CM) height, CM width, and

color. Several of these, such as the area and width of the convex hull, are traditional methods of

describing aboveground plant shape. Because the center of mass of a plant leaf changes as a

plant wilts, we developed additional descriptors based on the distance of the leaf center of mass

from the stem (CM width and CM height, X mass and Y mass). We then developed a pipeline

which used the original image as input, performed color correction, and quantified each

descriptor (Figure 1 and Supplemental Figure 4, and methods).

Figure 1: Diagram of the semi-automated analysis pipeline and 10 image descriptors

Image-based traits differentiate resistant and susceptible plants

To validate the efficiency of the image-based descriptors to estimate wilting phenotypes, we tested whether image-based phenotyping descriptors could differentiate resistant from susceptible plants. Supplemental Figure 5 shows the average normalized score for each of the image-based descriptors for each parent and the RIL population from -1 (the day before inoculation) to 6 dpi. Most descriptors, particularly those based on plant width or convex hull, had clearly divergent values in resistant and susceptible plants at a given time point. RIL descriptor values ranged from those of the resistant to susceptible parents and occasionally showed transgressive segregation (Supplemental Figure 5). Because plant shape at 6 dpi depends on shape of the same plant at -1 dpi, we used the evolution of each descriptor from day -1 to 6 dpi in our QTL analysis. This evolution was termed a 'trait'. Trait values were clearly different for resistant and susceptible parents for all traits except color (Figure 2), which was not used further.

Figure 2: Trait evolution during disease and correlation among descriptors. a. Boxplots showing the evolution of the normalized values between -1 and 6 dpi for each of the five biological replicates for the ten image-based traits for resistant Hawaii 7996, susceptible West Virginia parents and the 166 RILs. Color was unable to differentiate resistant from susceptible plants and was not used further. b. Heatmap showing the Pearson correlation values between image-based traits used in our QTL analysis and the human eye based wilting score in 2019. The correlations values were determined using the same plants imaged and visually assessed in 2019.

Image-based phenotyping reveals drivers of the wilting phenotype

We next investigated whether any of our traits were major components of the wilting phenotype.

We visually scored plants and asked how well our image-based traits correlated with human

visual scoring. Wilting is categorized by loss of plant leaf turgor that results in drooping leaves,

and decreased plant width and height. Determining how much a plant has wilted is challenging,

in part because it can be difficult to quantify how much each a leaf has drooped and how much

drooping of one leaf correlates with whole plant wilting.

We aimed to quantify leaf drooping using center of mass traits. Among our image-based traits,

those which were functions of the leaf center of mass were highly inversely correlated with

visual wilting (i.e. as a plant wilts, the center of mass decreases), suggesting that these are major

drivers of the wilting phenotype. These traits included CM width, plant width and X mass ($r > -$

0.75) (Figure 2B).

Several of our traits describe similar aspects of plant shape, such as height or width, through

different methods. These traits tended to be highly correlated with each other. For example, plant

height vs Y mass use different methods to describe plant width (based on the plant mask or the

center of mass of the stem masks; see Methods), and were highly positively correlated with each

other (0.93; Figure 2B).

We trained a random forest, consisting of 1000 decision trees, to use data from 6 dpi to predict

the expert visual score assigned at 8 dpi. We used 969 plants in a 60:40 training-testing data

split, and achieved a classification accuracy of 83%. To identify the traits that provided the most

efficient estimation of wilting symptoms, we randomly scrambled the nine image-based traits

one at a time. Plant width and plant area had the most impact on wilting score prediction

(Supplemental Figure 6).

QTL analysis identifies 20 wilting QTL in 10 clusters across the tomato genome at 6 dpi

Our overall goal was to identify tomato genomic regions that provide resistance or susceptibility

to *Ralstonia*. Prior to this analysis, we first generated a genetic map using Genotyping-by-

Sequencing (GBS). We identified 632 high-quality SNPs for linkage mapping using GBS. We

combined these with 112 SolCap markers, and subsequently generated a linkage map using ICI

mapping software (Meng et al., 2015). Our linkage map consisted of ~1300 cM (Supplemental

Figure 7) with an average per chromosome marker density that varies from 1.8 to 7.48 cM

(Supplemental Table 2).

For QTL analysis, we mapped the evolution of each our nine image-based descriptors from -1 to

6 dpi. In addition to the image-based phenotyping, we used two years (2016 and 2019) of

visually scored phenotyping data at 8 dpi. Visually assessed wilt scores from 2019 were

quantified from the same plants that were used for image-based phenotyping. In 2016, plants

were only visually scored for wilting, and no image-based measurements were taken. Using ICI

mapping software and composite interval mapping (CIM) for all traits, we identified 20 QTL

within the RIL population with a LOD score above 3 (Table 1). To be consistent with previous

studies of tomato-*Ralstonia* QTL mapping, we call these QTL, 'Bacterial wilt resistance (*Bwr*)

QTL'. Each *Bwr* QTL explained approximately 6 to 11% of the variation in response to *R.*

solanacearum strain K60, and together the clusters explained more than 88% of the variation

(calculated using the sum of the QTL with the highest PVE in each cluster).

The parent that donated the favorable allele was determined according to the sign of the QTL

additive effect (Awata et al., 2020), where a positive sign referred to H7996 resistant parent and

a negative sign to WV susceptible parent. Typically, in QTL analysis, the favorable allele has the

Chromosome	Cluster	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI	RightCI
2	Bwr2.1	45	solcap_snp_sl_18519	SL3.0ch02-49265134	plant width	3.4108	6.4576	-0.1669	42.5	47.5
2	Bwr2.2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	X mass	3.4765	9.4924	-0.2178	69.5	73.5
2	Bwr2.2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	CM width	3.2455	8.455	-0.2189	69.5	73.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	convex width	5.3238	10.2506	0.2103	21.5	26.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	4.7309	9.1511	0.1976	21.5	25.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	3.4765	5.8012	0.1705	21.5	26.5
3	Bwr3.1	23	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.6627	6.2716	0.1887	21.5	27.5
3	Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	4.5107	7.6902	0.1896	21.5	31.5
3	Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.3918	9.3283	0.1708	21.5	31.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	Wilting Score 2016	3.5258	7.3736	-0.1726	35.5	39.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	convex area	3.4328	9.0796	0.1953	34.5	39.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	plant area	3.3019	8.6697	0.1902	34.5	39.5
4	Bwr4.1	4	SL3.0ch04-60431371	solcap_snp_sl_11543	convex width	3.1855	6.0199	0.1627	1.5	10.5
5	Bwr5.1	100	solcap_snp_sl_22649	SL3.0ch05-65105663	plant height	3.6115	9.4852	0.2103	95.5	103.5
6	Bwr6.1	79	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2016	4.077	8.7626	-0.1871	76.5	81.5
6	Bwr6.1	81	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2019	3.2673	11.0473	-0.1621	78.5	81.5
10	Bwr10.1	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	7.6897	-0.1811	50.5	54.5
10	Bwr10.1	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	6.8155	-0.1714	50.5	54.5
12	Bwr12.1	37	solcap_snp_sl_1525	solcap_snp_sl_1572	CM height	4.1755	10.7222	0.1837	33.5	40.5
12	Bwr12.2	42	solcap_snp_sl_58807	solcap_snp_sl_9707	Y mass	3.3249	8.5939	0.1651	40.5	43.5

Table 1 Overview of the 20 wilting QTL identified in 10 clusters across the tomato genome at 6 dpi. LOD: maximum value of the Logarithm of the odd. PVE: Percentage of phenotypic variance explained. Add: Additive effect. Left CI and Right CI are the confidence interval calculated by a one-LOD decrease from the estimated QTL position.

higher trait value. Here, a higher trait value is favorable in all cases except the visual wilting

score, in which a higher trait value was associated with susceptibility (e.g. 90% wilting is more

susceptible than 20% wilting). For QTL detected using our visual wilting score, the susceptibility

allele was contributed by the susceptible parent WV (Table 1), consistent with other tomato-*R.*

solanacearum QTL studies based on visual assessment of wilting. For all QTL except those on

chromosome 2 and 10, the favorable allele was contributed by Hawaii 7996. QTL clusters on

chromosome 2 and 10, which contributed to plant height and width, were donated by the

susceptible parent WV. Favorable allelic contribution from both resistant and susceptible parents

is common in QTL studies for resistance (Maschietto et al., 2017; Awata et al., 2020).

Among the 20 individual *Bwr* QTL were 10 QTL clusters (Table 1). We use the term 'QTL

cluster' to describe QTL for different traits that co-localize at the same left and right genetic

marker. There are between one and six *Bwr* QTL within a given cluster, and each QTL within

the cluster has different LOD scores and explains a different percentage of phenotypic variation.

Bwr QTL for traits that are highly correlated with each other (Figure 2B) tended to cluster

together. For example, a cluster of QTL on chromosome 3 (*Bwr3.1*; Table 1) contains six *Bwr*

QTL, including three for area-related traits (convex width, and X mass and CM width), which

are correlated with each other at r = 0.94 - 0.97 (Figure 2B). In another cluster (*Bwr3.2*) on this

same chromosome, *Bwr* QTL for traits that describe plant area were detected together. This co-

localization supports the robust nature of our analyses.

In other clusters, only one trait that described one aspect of the wilting phenotype was present.

For example, despite several metrics that describe width, only convex width was identified as a

QTL on chromosome 4 (*Bwr4.1*; LOD 3.18; PVE = 6%). This suggests that the image-based

phenotyping captured genetic variation that is specific to each trait.

Image-based non-destructive phenotyping identifies three types of QTL clusters

Among our *Bwr* QTL clusters, we identified three types: those found using both image-based

and visual phenotyping (1 cluster, *Bwr3.2*), those found only through visual phenotyping (1

cluster, *Bwr6.1*), and those that were identified only through image-based phenotyping (8

clusters; Table 1 and Figure 3).

For the cluster including both image-based and visual phenotyping, (*Bwr3.2*) on chromosome 3,

we detected QTL for convex area, plant area and visual plant wilting in 2016 (Table 1 and Figure

3a). Previous studies examining tomato responses to other strains of *Ralstonia* have not detected

Bwr QTL in this region, indicating that it may be a novel target for *R. solanacearum* strain K60.

The identification of both image-based and visual QTL within the same cluster suggests that our

phenotyping methods are robust at identifying features of plant wilting. Close to this cluster is

Bwr3.1, which detected QTL for traits based on plant width, including convex width and CM

width. The proximal arm of chromosome 3 may be an important but unexploited region for

defense against *R. solanacearum* strain K60.

Figure 3: Significant QTL clusters on chromosomes 3 (a.), 6 (b.), 12 (c.) and 10 (d.). The image-based and visually assessed traits are represented with different colors for each chromosome. The vertical axis represents the genetic position (cM) and horizontal axis shows the LOD score. For each chromosome, the left panel represents the entire chromosome and the right panel represents the significant QTL cluster regions.

On chromosome 6 we detected *Bwr* (*Bwr6.1*) using only visual assessment for both 2016 and

2019 (Table 1, Figure 3b). This region may be a 'hot spot' for QTL for resistance to multiple

strains of *Ralstonia*. Previous studies using this same tomato population for mapping *Bwr* to

other strains of *Ralstonia* detected QTL approximately 4 Mb from this position (Supplemental

Figure 7).

In all other clusters, QTL were identified using only image-based phenotyping. Two such

clusters, detected on chromosome 12, identified QTL for CM height (*Bwr12.1*) and Y mass

(*Bwr12.2*), and were located near previous QTL detected against other strains of *Ralstonia*

(Supplemental Figure 7 and Figure 3c) . These previous studies (Wang et al., 2000; Jaw-Fen

Wang et al., 2013; Shin et al., 2020) detected QTL using visual wilt assessment of adult plants in

the field.

The remaining QTL detected through image-based phenotyping have not been previously

identified in studies using other strains of *Ralstonia*. These regions may be effective only against

specific strains of *Ralstonia* ('strain-specific' QTL), or may include those that are not easily

detected using visual assessment. In support of the latter, at one of the image-based-only clusters,

Bwr10.1 (Figure 3d), we detected a QTL using visual assessment that did not meet our threshold

for significance after permutation (Supplemental Table 3). Thus, image-based phenotyping

improved our ability to detect QTL.

Image-based phenotyping identified Bwr QTL prior to visual symptom development

In our system, wilting symptoms begin to appear on highly susceptible plants at 4 dpi, and these

plants are nearly 100% wilted by 6 dpi. To test whether we could detect *Bwr* QTL prior to the

onset of visual symptoms, we performed QTL analysis at 3, 4 and 5 dpi. No QTL were identified

based on visual assessment at any of these time points, however several QTL were identified

based on image-based phenotyping. At 3 dpi, three clusters were detected on chromosome 3, two

of which co-localized with those identified at 6 dpi (Supplemental Table 4). *Bwr3.1*, a cluster

detected only by image-based phenotyping, and *Bwr3.2*, a cluster detected by both visual and

image-based phenotyping, were first detected at 3 dpi. *Bwr3.1* was also detected at 5 and 6 dpi,

while *Bwr3.2* was detected 3, 4, 5 and 6 dpi (Supplemental Table 4). A small number of

additional QTL were detected at earlier time points (Supplemental Table 4), but in most cases

these were not stable, since they were not present at 6 dpi. These data suggest that image-based

phenotyping can identify *Bwr* QTL prior to the onset of wilting symptoms.

Plant architecture QTL do not overlap with Bwr QTL

The parents of the RIL population, H7996 and WV, have different aboveground phenotypes

(Supplemental Figure 4) and the shoot architecture of the RILs correspondingly varies. To ensure

that *Bwr* QTL were the result of tomato responses to *Ralstonia*, and not due to differences in

aboveground plant architecture, we used our traits in a QTL analysis at -1 dpi, the day before

plants had been inoculated. We call these 'tomato plant architecture (*Tpa*)' QTL. We identified

16 *Tpa*, within 9 QTL clusters (Table 2). None of the 16 *Tpa* were detected within the same

interval as *Bwr* at 6 dpi. However, two *Tpa* QTL clusters were detected that were near *Bwr* QTL

present at 6 dpi. One of these was detected on chromosome 3 at position 30, between *Bwr3.1*

and *3.2*.

Chromosome	Cluster	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI
2	Tpa2.1	117	solcap_snp_sl_8510	solcap_snp_sl_15574	plant width	3.3851	6.4842	0.129	114.5
2	Tpa2.1	117	solcap_snp_sl_8510	solcap_snp_sl_15574	X mass	4.1424	6.8279	0.139	114.5
3	Tpa3.1	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.6327	8.0934	0.1522	24.5
3	Tpa3.2	58	SL3.0ch03-61078160	solcap_snp_sl_10372	plant width	3.2538	9.4831	0.1555	56.5
3	Tpa3.2	58	SL3.0ch03-61078160	solcap_snp_sl_10372	X mass	3.8291	9.4218	0.1628	56.5
3	Tpa3.2	59	SL3.0ch03-61078160	solcap_snp_sl_10372	CM height	3.794	3.6957	0.1603	56.5
4	Tpa4.1	128	SL3.0ch04-1213025	solcap_snp_sl_63869	CM height	3.8063	12.0286	-0.2891	127.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	convex area	4.741	11.2755	0.1621	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant width	3.5865	6.9265	0.1336	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant area	3.5117	7.3344	0.1457	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	X mass	4.8686	8.016	0.151	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	Y mass	4.902	9.3612	0.1757	8.5
8	Tpa8.1	80	solcap_snp_sl_21461	SL3.0ch08-63373341	X mass	3.2439	5.3622	0.123	74.5
9	Tpa9.1	36	solcap_snp_sl_26683	SL3.0ch09-4516570	CM height	4.8307	6.146	0.2081	32.5
9	Tpa9.2	67	SL3.0ch09-39293163	SL3.0ch09-63338944	convex area	3.1976	7.523	-0.1323	60.5
10	Tpa10.1	71	solcap_snp_sl_8835	Le013158s_161	Y mass	3.2077	5.9734	0.1404	68.5

Table 2 Overview of tomato plant architecture (Tpa) QTL identified at -1 dpi by image-based phenotyping.

While we cannot rule out the possibility that the same genes underlie these *Tpa* and *Bwr* QTL,

their different marker intervals, coupled with the visual wilting QTL that is within the *Bwr3.2*

cluster, suggests that different genes are responsible for these QTL. Together, these data suggest

that our image-based phenotyping detected genomic regions that function in responding to

Ralstonia and *Bwr* QTL are not the result of differential growth patterns within the RILs.

Discussion

Breeding for plant disease resistance is one of the best strategies to combat plant diseases and

prevent major crop loss, but is challenging in part due to the complicated nature of disease

phenotyping. Here we used rapid, non-destructive, image-based phenotyping with RGB images

to identify 20 QTL in 10 clusters for tomato responses to *Ralstonia* at 6 dpi, two of which were

detected as early as 3 dpi. Together, the 20 *Bwr* QTL at 6 dpi explained more than 88% of the

variation in response to *Ralstonia*. Image-based phenotyping for shape-based traits that were

correlated with wilting detected both novel loci and those that overlapped with QTL based on

human visual assessment. These results establish the importance and feasibility of quantitative,

non-destructive, imaged-based phenotyping to identify new genetic targets for crop disease

resistance during disease progression.

Benefits of image-based phenotyping

Imaged-based phenotyping has been used to detect QTL associated with plant root (Topp et al.,

2013) and shoot (Zhang et al., 2017; Knoch et al., 2020; Li et al., 2020) architecture, plant height

(Wang et al., 2019), salt stress (Awlia et al.), and yield (Tanger et al., 2017; Pauli et al., 2016)

among other traits. Although image-based phenotyping has become increasingly common to

quantify plant disease symptoms (Mahlein, 2016; Mahlein et al., 2017, 2019; Lowe et al., 2017;

Shakoor et al., 2017; Mochida et al., 2019; Mir et al., 2019; Pérez-Bueno et al., 2019; Pineda et

al., 2021; Simko et al., 2017), few studies have used this technology to identify new genetic loci

for plant disease resistance (Yates et al., 2019; Fordyce et al., 2018; Corwin et al., 2016). One

reason for this may be that many plant disease symptoms occur at the leaf scale (such as spots

and specks), making it difficult to use some sensors non-destructively and in high-throughput, or

at the proper resolution needed to assess disease. For example, automated digital phenotyping of

Septoria Tritici Blotch on wheat leaves identified novel QTL for resistance, but the destructive

phenotyping required significant manual labor to harvest, mount and scan leaves (Yates et al.,

2019). While wilting does occur at the leaf scale, it is an easier phenotype to assess at the whole

plant scale. Another possibility is that imaging sensors are often expensive, making it more

challenging to phenotype the large number of plants needed in a QTL or GWA study. Our

method is low-cost and rapid (< 2 min/plant), and requires no manual labor other than placing

the plant on the turntable and initiating imaging via a computer.

Our image-based phenotyping identified loci that function in tomato responses to *Ralstonia*, but

did not detect one of the two *Bwr* clusters (*Bwr6.1*) we found using visual assessment of disease.

Visual assessment was performed at 8 dpi, while our image based phenotyping ended at 6 dpi. It

is possible *Bwr6.1* is not apparent until very late stages of disease. Alternatively, these results

could mean that we missed components of wilting in our image-based traits. However, a more

likely explanation is that our image-based traits were averaged across eight two dimensional

images, but wilting is a three dimensional (3D) phenotype. Images in 3D may allow detection of

Bwr6.1. It would also be of interest to test whether using additional types of sensors, such as

hyperspectral, would identify *Bwr 6.1*. Multi and hyperspectral imaging has been used to detect

biochemical changes in disease (Mahlein et al., 2017; Lowe et al., 2017; Zhang et al., 2020).

Given the chemical changes that occur during water stress, this type of imaging would likely

identify additional genetic variation associated with resistance to bacterial wilt disease.

New QTL for responses to Ralstonia in tomato

As in tomato, resistance to *Ralstonia* in other Solanaceous crops is quantitative (Young and

Danesh, 1994; Danesh et al., 1994; Thoquet et al., 1996a, 1996b; Mangin et al., 1999; Wang et

al., 2000; Carmeille et al., 2006; Jaw-Fen Wang et al., 2013). Breeding for resistance QTL is

thus the primary way forward to developing *Ralstonia*-resistant crops in the Solanaceae. This has

not been easy, in part because of the diversity of *Ralstonia*. The *Ralstonia solanacearum* species

complex was recently subdivided into three species (*R. solanacearum*, *R. pseudosolanacearum*,

and *R. syzygii*), each prevalent in a different part of the world (Remenant et al., 2012; Safni et al.;

Prior et al., 2016). Each species has multiple strains, with an overlapping, but distinct set of

virulence proteins that promote disease (Landry et al., 2020). Varieties with effective resistance

will likely have QTL that are effective against local strains (strain-specific QTL) as well as those

effective against multiple strains (broad-spectrum QTL).

Most previously identified QTL have focused on resistance to *R. pseudosolanacearum*, and none

of them used the strain of *R. solanacearum* as in this work (Danesh et al., 1994; Thoquet et al.,

1996a, 1996b; Wang et al., 2000; Carmeille et al., 2006; Jaw-Fen Wang et al., 2013; Shin et al.,

2020; Mangin et al., 1999). Using visual assessment of wilting and the same RIL population of

tomato (H7996 x WV700) used here, one broad-spectrum QTL for resistance to multiple strains

of *R. pseudosolanacearum,* and one strain of *R. solanacearum* (JT-516) had been previously

identified on chromosome 6 (Danesh et al., 1994; Thoquet et al., 1996a, 1996b; Mangin et al.,

1999; Wang et al., 2000; Carmeille et al., 2006; Jaw-Fen Wang et al., 2013; Shin et al., 2020).

We also identified a QTL on chromosome 6, using visual assessment in both 2016 and 2019. In

our study, *Bwr6.1* confers 8 – 11% of the variation, compared to 11.5 – 33% for *Bwr6a - Bwr6d*

(Jaw-Fen Wang et al., 2013; Shin et al., 2020). The genes within the left and right intervals of

Bwr6.1 are particularly interesting because there are two groups of NBS-LRRs located between

the left and right markers for *Bwr6.1* (solcap_snp_sl_14458 and the GBS-identified SNP at

39435402). One of these groups includes the *S. lycopersicum* homolog of Arabidopsis

RESISTANCE to PSEUDOMONAS SYRINGAE 4 (*RPS4*). Fine-mapping of *Bwr6.1* is needed to

determine whether a specific NBS-LRR protein or receptor is responsible for the resistance

conferred by this region.

Previous studies also identified a major QTL on chromosome 12 that is effective for resistance to

R. pseudosolanacearum, and explained between 15.9 – 53.9% of the variation (Shin et al., 2020).

We found two clusters of QTL on chromosome 12, for image-based traits. Within the region

spanning *Bwr12.1* on chromosome 12 is one NBS-LRR disease resistance gene (*Solyc12g10740*)

and one receptor-like kinase (*Solyc12g010660*) while within *Bwr12.2* are genes related to disease

resistance pathways, including a gene encoding a Pathogenesis Response (PR) Protein

(*Solyc12g014310*), and two genes encoding Leucine Rich Repeat – Receptor Like Kinases

(*Solyc12g014350* and *Solyc12g150105*). Our results suggest that this region of chromosome 12

may be important for resistance to multiple strains of *Ralstonia.* The QTL we identified on

chromosomes 2, 3, 4, 5, 8 and 10 were not previously identified, and thus may be specific to *R.*

solanacearum strain K60. With the exception of *Bwr3.2*, these QTL were detected using only

our image-based traits.

Together, our results establish the value of image-based, non-destructive disease phenotyping for

uncovering novel genetic components and new targets for quantitative disease resistance in

crops. By identifying new genetic loci, this type of rapid phenotyping may enable the

identification of broad-spectrum and durable resistance.

Methods

Plant growth

Seeds of 188 Hawaii7996 (H7996) x West Virginia700 (WV) recombinant inbred lines (RILs) in

the F8 generation were obtained from the Asian Vegetable Research and Development Center

(AVRDC) in June 2014. Seeds were propagated to the F_9 generation in field and greenhouse in

West Lafayette, IN in 2014 and 2015 and were used in QTL analysis in 2016 and 2019.

For all plants used in the imaging experiments and 2019 visual assessment of wilting, seeds were

sown into individually labeled 1801 traditional inserts that were placed into 1020 flats (Hummert

International, USA). Seeds in the 2016 experiment were sown into individually labeled 1203

inserts that were placed into 1020 flats. Seeds were sown into ProMix Propagation Mix supplied

by the Lilly Greenhouses and Growth Facility located at Purdue University, West Lafayette,

Indiana, USA. The individual pots were randomized and placed in a growth chamber at 28ºC,

relative humidity of 65% for a lighting cycle of 16 hours light/8 hours dark. The plants began

germination on day 4 and plants were inoculated with *R. solanacearum* strain K60 at 17 days

after planting when three true leaves were present. Trays were rotated in the growth chamber

throughout each experiment. In 2016, four seeds of each RIL were grown in the growth chamber

with parental controls, and each plant was visually assessed for wilting at 8 days post inoculation

(dpi). In 2019, one seed per RIL, along with parental controls, was grown in the growth chamber

for each of five independent replicates. Each plant was imaged at -1, 3, 4, 5, and 6 dpi, and

visually assessed at 8 dpi.

For plants used to genotyping with tomato SolCap markers, F_9 generation RIL seeds and parental

controls were grown in the greenhouse in 2-gallon pots with Metro Mix 510 soil. Plants were

grown for 6 weeks.

Ralstonia solanacearum growth and plant inoculation

Ralstonia solanacearum strain K60 (containing a GFP reporter) was grown on casamino acid-

peptone-glucose (CPG) agar containing tetrazolium chloride (TZC) in the dark for 48 hours at

28ºC as in (Caldwell et al., 2017). Briefly, bacteria were resuspended in sterile water to a

concentration of approximately 2 x 10^8 colony forming units (CFU)/mL for each experiment. For

each experiment, the concentration of inoculum was confirmed through dilution plating. Pots of

three-leaf plants were lightly compressed to induce wounding similar to transplant handling in

field conditions. 60 mL of inoculum was applied to the surrounding soil using a serological

pipet.

Visual assessment of bacterial wilt disease

Wilt scores were visually assessed 8 dpi. Wilt scores were calculated by counting the number of

wilted true leaves divided by the total number of true leaves on the plant. Plants were scored with

a 95% if the plant had all of its leaves wilted, excluding the topmost leaf. Plants were marked as

100% when the topmost portion of the stem was collapsed and wilted (Supplemental Figure 2).

Plant imaging

Plants were imaged the day before inoculation (-1 dpi) and then imaged on 3, 4, 5, and 6 dpi. A

Linco Linstor 2000-watt photo studio (Amazon, USA) was used as the backdrop, and Flora X

fluorescent lighting (Amazon, USA) was used to create a small photo studio. Individual plants

were placed on a Photocapture 360 turntable (Ortery Technologies) that was programmed to

capture eight images around the plant (every 45°). The images were captured using an EOS 6D

Canon camera with an EF 50mm f/1.4 USM lens on a stationary tripod. An imaging carriage

was created with a square petri dish and a fiducial marker attached to each side. A total of five

independent replicates for each RIL and the parental controls were imaged.

Phenotyping Pipeline and Image Analysis

From the plant images, ten traits were acquired for QTL analysis. A detailed description of the

acquisition process can be found in (Yang et al., 2021, 2020). All traits were extracted based on

plant color and shape. To eliminate any color discrepancy between images, images were

automatically color corrected using a fiducial marker consisting of a colored checkerboard with

known physical dimensions and colors.

After color correction, plant pixels were segmented from the rest of the image by thresholding

channels in the L*A*B* color space. The resulting segmentation mask was improved with image

morphological operations to fill holes and remove noise generated by the thresholding. The stem

of the plant was identified using two neural networks, Mask R-CNN and U-Net. By locating the

stem of the plant, metrics were defined that reflect the inner morphology of the plant. To train

the neural networks, stem segmentation ground truth data were generated using Adobe

Photoshop and LabelMe (a Python-based annotation tool) to mark the location of the image

pixels belonging to the stem of the plant. Further details about this procedure are described in

(Yang et al., 2020, 2021).

The plant and stem masks were used in the subsequent color and shape analysis, which acquired

ten traits: total area of the plant mask (plant area), height of the plant mask (plant height),

maximum width of the plant mask (plant width), area of the convex hull (convex area), width of

the convex hull (convex width), color-based weighted average (color average), horizontal

distance between the center of mass of the left and right sides of the plant stem (CM width),

height of the center of mass (CM height), x-axis distribution of the center of mass (X mass), Y-

axis distribution of the center of mass (Y mass).

The plant area, plant height and plant width were calculated from the plant mask. When

calculating plant height, the upper 5% of the plant material was not included, which helped

eliminate the impact of small leaves growing at the top. Using functions available in *OpenCV*

Python Library, a convex hull was fit around the plant mask and the convex area and convex

width were calculated. For color analysis, the weighted average of the pixel values in the A*

channel was calculated, as this channel captures the green-magenta spectrum.

To calculate traits involving the center of mass (CM), the plant mask was split into left and right
halves using the stem mask (Yang et al., 2020), and the CM located for each half. The x-axis
distance between CMs (CM width) and the average height of the CMs (CM height) were used as
traits.
459
The extension of just one leaf can have a major impact on measured plant width, generating a
disproportionately larger value. Capturing the plant material distribution inside the plant mask
can overcome this issue. Using the split plant mask, the horizontal and vertical distribution of
plant material was estimated. The distances at which 90% of the plant material was captured in
the horizontal and vertical directions were used as the traits 'X mass distribution' and 'Y mass
distribution'. The average of eight views around the plant were used for each trait value. Outputs
are in pixels.
467
468   In the random forest classification, visual wilting scores were thresholded to 0 or 1, where $0 = 0$
469   to 50% wilting, $1 = 51 - 100$% wilting. Average trait values from eight views from 6 dpi for 969
470   plants (RILs and parental controls) were used across all five replicates. Some plants did not have
471   a trait value for that day and were not used in the algorithm.
472
DNA extraction, Marker generation and Genotyping by Sequencing (GBS)
Tomato SolCap markers and single nucleotide polymorphisms (SNPs) identified through GBS
were used for map creation. Leaf disc samples were collected from each RIL plant using a
biopsy punch and were sent to LGC, Biosearch Technologies. Genomic DNA was extracted and
genotyped for 128 SNPs from the Tomato SolCAP panel using LGC's KASP assay. One
hundred twelve SNPs were obtained for RILs using this method.
479
480   For GBS, genomic DNA (gDNA) was extracted from 188 F_9 individuals of the RIL population
481   and from parental plants using Trizol. gDNA was RNase-treated and cleaned with phenol
482   chloroform extraction. Sequencing library preparation was as described by (Elshire et al., 2011).
483   Briefly, gDNA was digested with *PstI* and 150 bp paired end sequencing was performed on two
484   lanes of an Illumina Hi-seq 2500 at the Purdue Genomics Facility.
485

Reads were mapped to the *Solanum lycopersicum* 3.0 genome using the methods of (Manching
et al., 2017). In brief, reads were de-multiplexed, and adapter sequences were removed. Reads
were filtered based on the presence of a GBS barcode on the forward read (R1), which accounts
for 95.4% of the PE reads. For these R1, 97.3% had a paired R2 read, and 2.7% did not have a
pair. R1 read pairs and R1 singles reads were combined and assessed independent of pairing and
treated as single end reads for the remaining analyses (924,500,737 reads).
492
Reads were filtered to remove those with the presence of an internal restriction site, the lack of
restriction site hang sequence at the end, and for minimum length, which retained 93.0% of the
reads. There was a minimum of 17,561 reads, a maximum of 16,127,292 reads, and an average
of 4,480,149 reads per sample (Supplemental Figure 8). Reads were mapped using BWA-MEM
for paired end reads, and the GATK haplotype caller was used to generate a genomic Variant
Call Format (gVCF) file for downstream analysis.
There were 74,082 SNPs called between the population and *Solanum lycopersicum* 3.0 reference
genome. Many of these were SNPs between the reference genome and the population and do not
vary in the population used here. Filtering for the presence of two alleles within the RIL
population identified resulted in 2,738 SNPs. Of these, the parental alleles were identified in
both parents for 278 SNPs and in one parent for 698 SNPs. The remaining 1,762 SNPs could not
be assigned a parental origin and were not used for further analysis. The 976 SNPs with parental
origin identified were filtered for a minor allele frequency greater than 0.02 and less than 0.99,
which resulted in 632 high quality SNPs.
507
High quality SNPs identified from GBS were combined with previously defined SolCap markers
from LGC Biosearch Technologies for a total of 748 markers. SNP coverage of the 188 RIL taxa
was high. Of the 748 markers, the minimum coverage was 115 markers, the maximum coverage
was 555 markers, and the average was 366 markers (Supplemental Figure 9). There was some
residual heterozygosity in the taxa (Supplemental Figure 9b), and taxa with 10% or greater
heterozygosity were removed from further analysis (2 RILs).
514
Linkage Map Construction

The software QTL IciMapping (Meng et al., 2015) (version 4.1) was used for the map

construction with all 748 markers. Redundant markers were filtered by taxa coverage using a

missing rate of 85% and a distortion threshold at 0.001 to obtain a total of 408 markers. The 408

markers without anchor information were assigned to 12 groups based on a LOD score threshold

value of 3. After grouping, the markers were ordered using the nearest neighbor algorithm

(nnTwoOpt) using the rippling criterion SARF (Sum of Adjacent Recombination Frequencies)

with a window size of five markers. After ordering, some markers at the end of the chromosomes

were deleted when they were adding an insignificant genetic distance to the chromosome. These

markers were identified after splitting the current chromosome in two sub-chromosomes between

the longest marker interval. If the shortest sub-chromosome contained more than 20% of the

markers before splitting, the two sub-chromosome were re-assembled. Otherwise, the shortest

sub-chromosome was deleted.

QTL detection

QTL were detected using Inclusive Composite Interval Mapping with additive effects (ICIM-

ADD) in IciMapping (Meng et al., 2015) (version 4.1) using a genetic mapping with a 1 cM

scanning step and a probability in stepwise regression of 0.001. The LOD significance threshold

to declare a QTL significant was determined using a Type-I error of 0.0500 after one thousand

permutation. For the QTL analysis, we removed 19 of the 188 RIL (10%) showing the highest

variation according to the visual wilting score and 3 RIL due to missing data across the 5

biological repetitions for a total of 166 RIL into the QTL map (Supplemental Figure 3). The

evolution of the ten image-based traits and two visual wilting scores between -1 dpi and 3, 4, 5

and 6 dpi were determined for the 166 RIL. To assure a normal distribution of the phenotypic

data, the function "orderNorm" was used to perform an ordered quantile normalization (Peterson

and Cavanaugh, 2020) before QTL analysis using the package "bestNormalize" version 1.6.1

with R software version 3.6.1 (R Core Team).

Acknowledgements

We thank Chloe Caldwell for help with imaging, and Kristina Gans for help with GBS.

This work was funded by a Purdue Seed Grant to AIP and ED, a Foundation for Food and

Agriculture Research New Innovator Award to AIP, and the endowment of the Charles William

Harrison Distinguished Professorship at Purdue University to ED.

Author Contributions

A.I.P. and E.D. designed the research; V.M., D.C., and B.K. performed research; V.M, E.E.,

S.B., C.Y, A.I.P, and E.J.D. analyzed data; S.B., C.Y., and E.D. contributed new computational

tools; A.I.P. and V.M. wrote the manuscript; all authors edited and approved the manuscript.

References

Awata, L.A.O., Beyene Y., Gowda M., Suresh L.M., Jumbo M.B. Tongoona P., Danquah E., Ifie
B.E., Marchelo-Dragga P.W., Olsen M., Ogugu V. et al. (2020). Genetic Analysis of
QTL for Resistance to Maize Lethal Necrosis in Multiple Mapping Populations. Genes
11: 32.

Awlia, M., Alshareef, N., Saber, N., Korte, A., Oakey, H., Panzarová, K., Trtílek, M., Negrão,
S., Tester, M., and Julkowska, M.M. (2021).Genetic mapping of the early responses to
salt stress in *Arabidopsis thaliana*. Plant J. doi:10.1111/tpj.15310.

Bock, C., Parker, P., Cook, A.Z., and Gottwald, T. (2008). Visual Rating and the Use of Image
Analysis for Assessing Different Symptoms of Citrus Canker on Grapefruit Leaves. Plant
Dis 92:530-541.

Bock, C., Poole, G., Parker, P., and Gottwald, T.R. (2010). Plant Disease Severity Estimated
Visually, by Digital Photography and Image Analysis, and by Hyperspectral Imaging.
Crit Rev Plant Sci 29: 59-107.

Caldwell, D., Kim, B.-S., and Iyer-Pascuzzi, A.S. (2017). *Ralstonia solanacearum* Differentially
Colonizes Roots of Resistant and Susceptible Tomato Plants. Phytopathology 107: 528–
536.

Carmeille, A., Caranta, C., Dintinger, J., Prior, P., Luisetti, J., and Besse, P. (2006).
Identification of QTLs for *Ralstonia solanacearum* race 3-phylotype II resistance in
tomato. Theor Appl Genet 113: 110–21.

Colwell, R.N. (1956). Determining the prevalence of certain cereal crop diseases by means of
aerial photography. Hilgardia 26: 223–286.

Corwin, J.A., Copeland, D., Feusier, J., Subedy, A., Eshbaugh, R., Palmer, C., Maloof, J., and
Kliebenstein, D.J. (2016). The Quantitative Basis of the Arabidopsis Innate Immune

System to Endemic Pathogens Depends on Pathogen Genetics. PLoS Genet 12:
e1005789.

Danesh, D., Aarons, S., McGill, G.E., and Young, N.D. (1994). Genetic Dissection of
Oligogenic Resistance to Bacterial Wilt in Tomato. Mol Plant Microbe Interact 7: 464–
471.

Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., and Mitchell,
S.E. (2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High
Diversity Species. PLOS ONE 6: e19379.

Fordyce, R.F., Soltis, N.E., Caseys, C., Gwinner, R., Corwin, J.A., Atwell, S., Copeland, D.,
Feusier, J., Subedy, A., Eshbaugh, R., and Kliebenstein, D.J. (2018). Digital Imaging
Combined with Genome-Wide Association Mapping Links Loci to Plant-Pathogen
Interaction Traits. Plant Physiol 178: 1406–1422.

Genin, S. (2010). Molecular traits controlling host range and adaptation to plants in *Ralstonia*
solanacearum. New Phytol 187: 920–928.

Genin, S. and Denny, T.P. (2011). Pathogenomics of the *Ralstonia solanacearum* Species
Complex. Annu Rev Phytopathol 50: 67–89.

Jackson, R.D. (1986). Remote Sensing of Biotic and Abiotic Plant Stress. Annu Rev Phytopathol
24: 265–287.

Jaw-Fen Wang, Fang-I Ho, Hai Thi Hong Truong, Shu-Mei Hugan, Conrado H. Balatero, Vanla
Dittapongpitch, and Nurul Hidayati (2013). Identification of major QTLs associated with
stable resitance of tomato cultivar "Hawaii 7996" to *Ralstonia solanacearum*. Euphytica
190: 241–252.

Knoch, D., Abbadi, A., Grandke, F., Meyer, R.C., Samans, B., Werner, C.R., Snowdon, R.J., and
Altmann, T. (2020). Strong temporal dynamics of QTL action on plant growth
progression revealed through high-throughput phenotyping in canola. Plant Biotechnol J
18: 68–82.

Landry, D., González-Fuente, M., Deslandes, L., and Peeters, N. (2020). The large, diverse, and
robust arsenal of *Ralstonia solanacearum* type III effectors and their in planta functions.
Mol. Plant Pathol. doi:10.1111/mpp.12977

Li, H., Feng, H., Guo, C., Yang, S., Huang, W., Xiong, X., Liu, J., Chen, G., Liu, Q., Xiong, L.,
Liu, K., and Yang, W. (2020). High-throughput phenotyping accelerates the dissection of
the dynamic genetic architecture of plant growth and yield improvement in rapeseed.
Plant Biotechnol J 18: 2345–2353.

Lowe, A., Harrison, N., and French, A.P. (2017). Hyperspectral image analysis techniques for
the detection and classification of the early onset of plant disease and stress. Plant
Methods 13: 80.

Mahlein, A.-K. (2016). Plant Disease Detection by Imaging Sensors - Parallels and Specific
Demands for Precision Agriculture and Plant Phenotyping. Plant Dis. 100: 241-251.

Mahlein, A.-K., Alisaac, E., Al Masri, A., Behmann, J., Dehne, H.-W., and Oerke, E.-C. (2019).
Comparison and Combination of Thermal, Fluorescence, and Hyperspectral Imaging for
Monitoring Fusarium Head Blight of Wheat on Spikelet Scale. Sensors 19: 2281.

Mahlein, A.-K., Kuska, M., Thomas, S., Bohnenkamp, D., Alisaac, E., Behmann, J., Wahabzada,
M., and Kersting, K. (2017). Plant disease detection by hyperspectral imaging: from the
lab to the field. Adv Anim Biosci 8: 238-243.

Manching, H., Sengupta, S., Hopper, K.R., Polson, S.W., Ji, Y., and Wisser, R.J. (2017). Phased
Genotyping-by-Sequencing Enhances Analysis of Genetic Diversity and Reveals
Divergent Copy Number Variants in Maize. G3 Genes Genom Genet 7: 2161–2170.

Mangin, B., Thoquet, P., Olivier, J., and Grimsley, N.H. (1999). Temporal and multiple
quantitative trait loci analyses of resistance to bacterial wilt in tomato permit the
resolution of linked loci. Genetics 151: 1165–1172.

Mansfield, J. et al. (2012). Top 10 plant pathogenic bacteria in molecular plant pathology. Mol
Plant Pathol 13: 614–629.

Maschietto, V., Colombi, C., Pirona, R., Pea, G., Strozzi, F., Marocco, A., Rossini, L., and
Lanubile, A. (2017). QTL mapping and candidate genes for resistance to Fusarium ear rot
and fumonisin contamination in maize. BMC Plant Biology 17: 20.

Meng, L., Li, H., Zhang, L., and Wang, J. (2015). QTL IciMapping: Integrated software for
genetic linkage map construction and quantitative trait locus mapping in biparental
populations. Crop J 3: 269–283.

Mir, R.R., Reynolds, M., Pinto, F., Khan, M.A., and Bhat, M.A. (2019). High-throughput
phenotyping for crop improvement in the genomics era. Plant Sci 282: 60–72.

Mochida, K., Koda, S., Inoue, K., Hirayama, T., Tanaka, S., Nishii, R., and Melgani, F. (2019).
Computer vision-based phenotyping for improvement of plant productivity: a machine
learning perspective. GigaScience 8.

Pauli, D., Andrade-Sanchez, P., Carmo-Silva, A.E., Gazave, E., French, A.N., Heun, J.,
Hunsaker, D.J., Lipka A.E., Setter, T.L., Strand, R.J. et al. (2016). Field-Based High-
Throughput Plant Phenotyping Reveals the Temporal Patterns of Quantitative Trait Loci
Associated with Stress-Responsive Traits in Cotton. G3 Genes Genom Genet 6: 865–879.

Pérez-Bueno, M.L., Pineda, M., and Barón, M. (2019). Phenotyping Plant Responses to Biotic
Stress by Chlorophyll Fluorescence Imaging. Front. Plant Sci. 10.

Peterson, R.A. and Cavanaugh, J.E. (2020). Ordered quantile normalization: a semiparametric
transformation built for the cross-validation era. J Applied Stat 47: 2312–2327.

Pineda, M., Barón, M., and Pérez-Bueno, M.-L. (2021). Thermal Imaging for Plant Stress
651   Detection and Phenotyping. Remote Sens 13: 68.

652   Prior, P., Ailloud, F., Dalsing, B.L., Remenant, B., Sanchez, B., and Allen, C. (2016). Genomic
653   and proteomic evidence supporting the division of the plant pathogen *Ralstonia*
654   *solanacearum* into three species. BMC Genomics 17: 90.

655   R Core Team R: A language and environment for statistical computing. R Foundation for
656   Statistical Computing Vienna: URL https://www.R-project.org/.

657   Remenant, B., Babujee, L., Lajus, A., Médigue, C., Prior, P., and Allen, C. (2012). Sequencing
658   of K60, Type Strain of the Major Plant Pathogen *Ralstonia solanacearum*. J Bacteriol
659   194: 2742–2743.

660   Safni, I., Cleenwerck, I., De Vos, P., Fegan, M., Sly, L., and Kappler, U. 2014 Polyphasic
661   taxonomic revision of the *Ralstonia solanacearum* species complex: proposal to emend
662   the descriptions of *Ralstonia solanacearum* and *Ralstonia syzygii* and reclassify current
663   *R. syzygii* strains as *Ralstonia syzygii* subsp. syzygii subsp. nov., *R. solanacearum*
664   phylotype IV strains as *Ralstonia syzygii* subsp. indonesiensis subsp. nov., banana blood
665   disease bacterium strains as *Ralstonia syzygii* subsp. celebesensis subsp. nov. and *R.*
666   *solanacearum* phylotype I and III strains as *Ralstonia pseudosolanacearum* sp. nov. Int J
667   Syst Evol Microbiol 64: 3087–3103.

668   Schandry, N. (2017). A Practical Guide to Visualization and Statistical Analysis of R.
669   solanacearum Infection Data Using R. Front. Plant Sci. 8.

670   Shakoor, N., Lee, S., and Mockler, T.C. (2017). High throughput phenotyping to accelerate crop
671   breeding and monitoring of diseases in the field. Curr Opin Plant Biol 38: 184–192.

672   Shin, I.S., Hsu, J.-C., Huang, S.-M., Chen, J.-R., Wang, J.-F., Hanson, P., and Schafleitner, R.
673   (2020). Construction of a single nucleotide polymorphism marker based QTL map and
674   validation of resistance loci to bacterial wilt caused by *Ralstonia solanacearum* species
675   complex in tomato. Euphytica 216: 54.

676   Simko, I., Jimenez-Berni, J.A., and Sirault, X.R.R. (2017). Phenomic Approaches and Tools for
677   Phytopathologists. Phytopathology 107: 6–17.

678   Stewart, E. and McDonald, B. (2014). Measuring quantitative virulence in the wheat pathogen
679   *Zymoseptoria tritici* using high-throughput automated image analysis. Phytopathology
680   104: 985-992.

681   Stewart, E.L., Hagerty, C.H., Mikaberidze, A., Mundt, C.C., Zhong, Z., and McDonald, B.A.
682   (2016). An Improved Method for Measuring Quantitative Resistance to the Wheat
683   Pathogen *Zymoseptoria tritici* Using High-Throughput Automated Image Analysis.
684   Phytopathology 106: 782–788.

685   Tanger, P., Klassen S., Mojica J.P., Lovell, J.T., Moyers, B.T., Baraoidan, M., Naredo, M.E.B.,
686   McNally, K.L., Poland, J., Bush, D.R. et al. (2017). Field-based high throughput

phenotyping rapidly identifies genomic regions controlling yield components in rice. Sci
Rep 7: 42839.

Thoquet, P., Olivier, J., Sperisen, C., Rogowsky, P., Laterrot, H., and Grimsley, N. (1996a).
Quantitative trait loci determining resistance to bacterial wilt in tomato cultivar
Hawaii7996. Mol Plant Microbe Interact 9: 826–836.

Thoquet, P., Olivier, J., Sperisen, C., Rogowsky, P., Prior, P., Anais, G., Mangin, B., Bazin, B.,
Nazer, R., and Grimsley, N. (1996b). Polygenic resistance of tomato plants to bacterial
wilt in the French West Indies. Mol Plant-Microbe Interact 9: 837–842.

Topp, C.N., Iyer-Pascuzzi, A.S., Anderson, J.T., Lee, C-R., Zurek, P.R., Symonova, O., Zheng
Y., Bucksch, A., Mileyko, Y., Galkovskyi, T., et al(2013). 3D phenotyping and
quantitative trait locus mapping identify core regions of the rice genome controlling root
architecture. Proc. Natl. Acad. Sci. U.S.A. 110: E1695-1704.

Wang, J.F., Olivier, J., Thoquet, P., Mangin, B., Sauviac, L., and Grimsley, N.H. (2000).
Resistance of tomato line Hawaii7996 to *Ralstonia solanacearum* Pss4 in Taiwan is
controlled mainly by a major strain-specific locus. Mol Plant-Microbe Interact 13: 6–13.

Wang, X. et al. (2019). Dynamic plant height QTL revealed in maize through remote sensing
phenotyping using a high-throughput unmanned aerial vehicle (UAV). Sci Rep 9: 3458.

Yang, C., Baireddy, S., Cai, E., Meline, V., Caldwell, D., Iyer-Pascuzzi, A.S., and Delp, E.J.
(2021). Image-Based Plant Wilting Estimation. arXiv:2105.12926 [cs].

Yang, C., Baireddy, S., Chen, Y., Cai, E., Caldwell, D., Méline, V., Iyer-Pascuzzi, A.S., and
Delp, E.J. (2020). Plant Stem Segmentation Using Fast Ground Truth Generation. In
2020 IEEE Southwest Symposium on Image Analysis and Interpretation (SSIAI), pp. 62–
65.

Yates, S., Mikaberidze, A., Krattinger, S.G., Abrouk, M., Hund, A., Yu, K., Studer, B., Fouche,
S., Meile, L., Pereira, D., Karisto, P., and McDonald, B.A. (2019). Precision Phenotyping
Reveals Novel Loci for Quantitative Resistance to Septoria Tritici Blotch. Plant
Phenomics 2019: 1–11.

Young, N.D. and Danesh, D. (1994). Understanding Bacterial Wilt Resistance in Tomato
through the Use of DNA Genetic-Markers. Bacterial Wilt: 145–155.

Zhang, N., Yang, G., Pan, Y., Yang, X., Chen, L., and Zhao, C. (2020). A Review of Advanced
Technologies and Development for Hyperspectral-Based Plant Disease Detection in the
Past Three Decades. Remote Sensing 12: 3188.

Zhang, X. et al. (2017). High-Throughput Phenotyping and QTL Mapping Reveals the Genetic
Architecture of Maize Plant Growth. Plant Physiol 173: 1554–1564.

 Supplemental Information

 Supplemental Figures

725 Supplemental Figure 1: Design of our low cost phenotyping platform including automatic

726 turntable, backdrop, lightning and RGB camera.

728 Supplemental Figure 2: Raw RGB pictures showing the evolution of wilting symptoms on RIL #
729 646 at -1, 3, 4, 5 and 6 dpi. Visually assessed wilting score are expressed in percentage of wilted
730 leaves.
731
732

733

755 Supplemental Figure 3: Wilting Score RIL: Boxplots showing the normalized wilting score at 6

756 dpi for the 166 RILs, resistant Hawaii 7996 (HA) and susceptible West Virginia (WV)

757 genotypes.

768 Supplemental Figure 4: RGB pictures showing the evolution of the traits at -1, 3, 4, 5 and 6 dpi
769 for A) Hawaii 7996 and B) West Virginia (pw: plant width, ph: plant height, CMh: Center of
770 Mass height, CMw: Center of Mass width, xm: Horizontal mass distribution, ym: Vertical mass
771 distribution). The Hawaii 7996 plant shown here is the same plant used for Figure 1.

Hawaii 7996

West Virginia

Supplemental Figure 5: Boxplots showing the normalized values for image-based traits at -1, 3, 4, 5, 6 dpi for resistant Hawaii 7996, susceptible West Virginia 700, and 166 individuals of the RIL population.

786   Supplemental Figure 6: Histogram showing the image-based descriptors which contribute the

787   most to wilting

808   Supplemental Figure 7: Genetic Linkage map constructed with SolCap and SNP markers.

809   Location of 10 QTL clusters are displayed using blue and green for left and right markers

810   respectively. Location of previously identified QTL on chromosome 6 and 12[27] are also

811   displayed.

818   Supplemental Figure 8: GBS Read Distribution by Sample (includes parental references).

839   Supplemental Figure 9: a) RIL and parental coverage and SNP density from GBS analysis; b)

840   Percent heterozygosity and SNP density

Supplemental Tables

Supplemental Table 1 Descriptive statistics for 10 image-based traits and two years of visually assessed wilting scores (2016 and 2019). Image-based traits were from the same plants as the visually assessed wilting score in 2019. The RIL population was inoculated with Ralstonia and visually scored for wilting in 2016, but images were not taken. Data is shown for resistant parent Hawaii 7996, susceptible parent West Virginia 700, and the RIL population.

year	Variable	Hawaii 7996				West Virginia 700				RILs			
		Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum
2016	Wilting_Score	0	0	0	0	100	0	100	100	44.41	41.4	0	100
	Wilting_Score	0	0	0	0	99.33	5.16	60	100	53.69	42.23	0	100
	convex area	2.7e+06	1.4e+06	-4.1e+04	6.3e+06	-3.6e+05	3.6e+05	-1.1e+06	7.4e+05	1.2e+06	1.4e+06	-2.0e+06	7.9e+06
	convex width	8.7e+02	4.4e+02	-7.0e+01	1.7e+03	-7.4e+02	3.7e+02	-1.4e+03	5.4e+02	-4.6e-01	5.9e+02	-1.9e+03	2.9e+03
	plant area	8.7e+05	5.6e+05	-1.0e+04	2.2e+06	-1.1e+05	1.1e+05	-3.4e+05	2.1e+05	5.0e+05	5.5e+05	-7.4e+05	2.6e+06
	plant height	5.6e+02	1.9e+02	-7.0e+01	8.7e+02	-1.3e+01	1.7e+02	-3.6e+02	3.7e+02	3.3e+02	2.5e+02	-6.5e+02	1.1e+03
2019	plant width	4.0e+02	2.0e+02	-3.6e+01	8.2e+02	-3.5e+02	1.8e+02	-6.6e+02	2.5e+02	-2.1e+01	2.7e+02	-9.7e+02	8.4e+02
	X mass	3.1e+02	1.6e+02	-3.7e+01	5.8e+02	-2.9e+02	1.5e+02	-5.6e+02	2.0e+02	-1.2e+01	2.3e+02	-7.4e+02	6.9e+02
	Y mass	5.4e+02	2.1e+02	-8.4e+01	9.3e+02	-6.7e+01	1.6e+02	-4.1e+02	3.6e+02	2.1e+02	2.2e+02	-7.0e+02	9.6e+02
	CM width	3.1e+02	1.6e+02	-1.0e+02	6.0e+02	-2.9e+02	1.5e+02	-5.2e+02	1.8e+02	2.7e+01	2.3e+02	-7.0e+02	9.8e+02
	CM height	2.6e+02	1.8e+02	-2.1e+02	6.2e+02	-1.3e+02	1.5e+02	-3.9e+02	1.9e+02	-5.2e+01	1.9e+02	-8.8e+02	8.3e+02
	color average	2.88	5.44	-5.96	22.9	7.37	2.84	1.03	14.83	4.06	4.48	-11.35	16.48

882   Supplemental Table 2: Marker density per cM for each chromosome.

Chromosome	Number of marker	Chromosome Length (cM)	Average interval length (cM)	Standard deviation	Standard error
1	36	153.42	4.38	4.15	0.70
2	19	117.42	6.52	3.92	0.92
3	28	87.77	3.25	3.56	0.68
4	44	133.43	3.10	4.25	0.65
5	18	116.49	6.85	5.33	1.29
6	61	116.92	1.80	1.99	0.26
7	21	95.63	4.78	3.94	0.88
8	14	97.27	7.48	4.44	1.23
9	27	143.5	5.52	5.77	1.13
10	18	80.01	4.71	4.80	1.16
11	25	61.22	2.55	3.88	0.79
12	43	120.42	2.87	3.01	0.46

905 Supplemental Table 3: Overview of the 9 genomic regions identified with a LOD score greater

906 than 3 on chromosome 10 after inoculation with R. solanacearum at 6 dpi. According to the ICI

907 Mapping analysis using 1000 permutations, a threshold value of 3.48 represents the minimum

908 LOD to identify a significant QTL. Genomic regions with a lower LOD value did not reach our

909 threshold for significance after permutation.

Chromosome	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add
10	51	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0067	13.3810	0.2077
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0927	10.5192	0.1842
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0832	8.0396	0.1611
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.1634	9.0040	-0.1835
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	7.8538	-0.1714
10	51	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.0398	11.1872	-0.2039
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.5411	10.2424	-0.1951
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	8.8147	-0.1811
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	X mass	3.0584	7.0293	-0.1629

Supplemental Table 4: Overview of QTLs identified across the genome at all time points (-1, 3,

4, 5, 6 dpi) with R. solanacearum. LOD: maximum value of the Logarithm of the odd. PVE:

Percentage of phenotypic variance explained. Add: Additive effect, the positive and negative

values indicated that the alleles are introgressed from the resistant parent (Hawaii 7996) and

susceptible parents (WestVirginia). LeftCI and RightCI are the confidence interval calculated by

a one-LOD decrease from the estimated QTL position. * = Bwr3.1; ** = Bwr3.2

	Timepoint	Chromosome	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI	RightCI
	-1	2	117	solcap_snp_sl_8510	solcap_snp_sl_15574	plant width	3.3851	6.4842	0.1290	114.5	117.0
	-1	2	117	solcap_snp_sl_8510	solcap_snp_sl_15574	X mass	4.1424	6.8279	0.1390	114.5	117.0
	-1	3	58	SL3.0ch03-61078160	solcap_snp_sl_10372	plant width	3.2538	9.4831	0.1555	56.5	64.5
	-1	3	58	SL3.0ch03-61078160	solcap_snp_sl_10372	X mass	3.8291	9.4218	0.1628	56.5	64.5
	-1	3	59	SL3.0ch03-61078160	solcap_snp_sl_10372	CM height	3.7940	3.6957	0.1603	56.5	63.5
	-1	3	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.6327	8.0934	0.1522	24.5	34.5
	-1	4	128	SL3.0ch04-1213025	solcap_snp_sl_63869	CM height	3.8063	12.0286	-0.2891	127.5	130.5
	-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	convex area	4.7410	11.2755	0.1621	8.5	11.5
	-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant width	3.5865	6.9265	0.1336	8.5	11.5
	-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant area	3.5117	7.3344	0.1457	8.5	11.5
	-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	X mass	4.8686	8.0160	0.1510	8.5	11.5
	-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	Y mass	4.9020	9.3612	0.1757	8.5	11.5
	-1	8	80	solcap_snp_sl_21461	SL3.0ch08-63373341	X mass	3.2439	5.3622	0.1230	74.5	83.5
	-1	9	67	SL3.0ch09-39293163	SL3.0ch09-63338944	convex area	3.1976	7.5230	-0.1323	60.5	69.5
	-1	9	36	solcap_snp_sl_26683	SL3.0ch09-4516570	CM height	4.8307	6.1460	0.2081	32.5	40.5
	-1	10	71	solcap_snp_sl_8835	Le013158s_161	Y mass	3.2077	5.9734	0.1404	68.5	73.5
*	3	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	CM height	7.2477	18.2665	0.2584	21.5	30.5
	3	3	27	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.5022	8.8528	0.1536	22.5	32.5
	3	3	26	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	5.5180	13.9255	0.2191	21.5	27.5
**	3	3	26	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	4.5407	11.7470	0.1991	22.5	32.5
	3	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	5.6025	13.8198	0.2120	37.5	42.5
	3	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	plant height	4.5094	10.3345	0.1974	25.5	31.5
	3	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	Y mass	5.7227	15.0239	0.2325	23.5	31.5
**	4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	3.9750	10.2036	0.1829	37.5	43.5
	4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	CM width	4.5294	8.4260	0.1853	37.5	43.5
	4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	plant width	3.8768	9.9017	0.1838	37.5	43.5
	4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	X mass	3.8156	9.7282	0.1837	37.5	43.5
	4	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	convex area	3.1310	7.7809	0.1508	22.5	32.5
	4	3	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.8685	9.9594	0.1672	23.5	33.5
*	5	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	CM height	3.6770	9.3886	0.1414	21.5	27.5
	5	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	3.4953	9.3017	0.1806	21.5	27.5
	5	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	3.6851	8.5381	0.1738	21.5	31.5
	5	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.4687	9.9830	0.1673	21.5	31.5
**	5	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	3.7859	9.8306	0.1891	37.5	43.5
	5	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	CM width	4.0630	7.4955	0.1924	35.5	39.5
	5	5	107	SL3.0ch05-65105663	SL3.0ch05-66111590	plant height	3.4091	8.9327	0.1778	104.5	114.5
	5	12	56	SL3.0ch12-62154155	solcap_snp_sl_19345	CM height	3.2828	8.2102	0.1360	54.5	59.5
	6	2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	X mass	3.4765	9.4924	-0.2178	69.5	73.5
	6	2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	CM width	3.2455	8.4550	-0.2189	69.5	73.5
	6	2	45	solcap_snp_sl_18519	SL3.0ch02-49265134	plant width	3.4108	6.4576	-0.1669	42.5	47.5
*	6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	convex width	5.3238	10.2506	0.2103	21.5	26.5
	6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	4.7309	9.1511	0.1976	21.5	25.5
	6	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	4.5107	7.6902	0.1896	21.5	31.5
	6	3	23	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.6627	6.2716	0.1887	21.5	27.5
	6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	3.4765	5.8012	0.1705	21.5	26.5
	6	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.3918	9.3283	0.1708	21.5	31.5
**	6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	Wilting Score 2016	3.5258	7.3736	-0.1726	35.5	39.5
	6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	convex area	3.4328	9.0796	0.1953	34.5	39.5
	6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	plant area	3.3019	8.6697	0.1902	34.5	39.5
	6	4	4	SL3.0ch04-60431371	solcap_snp_sl_11543	convex width	3.1855	6.0199	0.1627	1.5	10.5
	6	5	100	solcap_snp_sl_22649	SL3.0ch05-65105663	plant height	3.6115	9.4852	0.2103	95.5	103.5
	6	6	79	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2016	4.0770	8.7626	-0.1871	76.5	81.5
	6	6	81	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2019	3.2673	11.0473	-0.1621	78.5	81.5
	6	10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	7.6897	-0.1811	50.5	54.5
	6	10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	6.8155	-0.1714	50.5	54.5
	6	12	37	solcap_snp_sl_1525	solcap_snp_sl_1572	CM height	4.1755	10.7222	0.1837	33.5	40.5
	6	12	42	solcap_snp_sl_58807	solcap_snp_sl_9707	Y mass	3.3249	8.5939	0.1651	40.5	43.5