Image-based assessment of plant disease progression identifies new genetic loci for resistance

Valérian Méline[a], Denise L. Caldwell[a], Bong-Suk Kim[a], Sriram Baireddy[b], Changye Yang[b], Erin E. Sparks[c], Edward J. Delp[b], and Anjali S. Iyer-Pascuzzi[a†]

[a] 915 W. State Street, Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana, USA
[b] Video and Image Processing Laboratory (VIPER), School of Electrical and Computer Engineering
[c] Department of Plant and Soil Sciences and the Delaware Biotechnology Institute, University of Delaware, Newark, DE

†Author for correspondence: asi2@purdue.edu

Keywords: Digital phenotyping, QTL, *Ralstonia*, bacterial wilt, tomato

Running title: Disease resistance QTL from image-based phenotyping

Abstract

A major challenge in global crop production is mitigating yield loss due to plant diseases. One of

the best means of disease control is plant resistance, but the identification of genes that promote

resistance has been limited by the subjective quantification of disease, which is typically scored

by the human eye. We hypothesized that image-based quantification of disease phenotypes

would enable the identification of new disease resistance loci. We tested this using the

interaction between tomato and *Ralstonia solanacearum*, a soilborne pathogen that causes

bacterial wilt disease. We acquired over 40,000 time-series images of disease progression in a

tomato recombinant inbred line population, and developed an image analysis pipeline providing

a suite of ten traits to quantify wilt disease based on plant shape and size. Quantitative trait loci

(QTL) analyses using image-based phenotyping identified QTL that were both unique and shared

compared with those identified by human assessment of wilting. When shared loci were

identified, image-based phenotyping could detect some QTL several days earlier than human

assessment. Thus, expanding the phenotypic space of disease with image-based phenotyping

allowed both earlier detection and identified new genetic components of resistance.

Main

Plant diseases are a significant global constraint to crop production. Developing disease resistant

crops requires identifying the plant genomic regions and genes that contribute to resistance to the

pathogenic microbes that cause disease. This in turn depends on phenotyping large populations

of plants for their responses to pathogens. Phenotyping plant diseases is challenging because

diseases cause complex, quantitative phenotypes that can occur at different scales – e.g. on parts

of leaves, entire leaves, or the whole plant. In addition, disease phenotypes vary over time and

depend on environmental conditions, plant age, and pathogen virulence. Disease symptoms such

as wilting or necrotic spots have traditionally been scored with the human eye, but these scores

are subjective, can vary by individual, and are difficulty to accurately quantify.

The challenging nature of visual disease assessment has led to the use of sensors including RGB,

hyperspectral, chlorophyll fluorescence and thermal cameras to assess disease symptoms [1–7].

Compared to assessment by the human eye, image-based phenotyping is faster, more

reproducible, and more sensitive to small variations in disease symptoms that can be critical for

detecting resistance loci [3–5,8–10]. Although many studies have used or developed tools to assess

plant symptoms using different types of sensors [4–7,11–16], few studies have used these

technologies in QTL or Genome Wide Association (GWA) analyses for responses to plant

pathogens.

The soil-borne betaproteobacterium *Ralstonia solanacearum* is the causal agent of bacterial wilt

disease and has been ranked as one of the top 10 most destructive plant bacterial pathogens of all

time [17]. *Ralstonia* infection causes susceptible plants to wilt, and the amount of wilting correlates

with a plant's level of susceptibility [18,19]. The bacterium is a major production constraint in

Solanaceous crops both globally and in the United States, where disease loss in tomatoes can

exceed 70%. In crops, resistance to *Ralstonia* is quantitative, but the quantitative trait loci (QTL)

underlying resistance to US strains of *Ralstonia* are largely unknown. QTL for other strains have

been mapped [20–27] but have not been cloned, and the host determinants necessary for resistance

remain mostly unspecified.

The limited identification of QTL for *Ralstonia* resistance can be attributed in part to the

difficulty in accurately scoring plant wilting. Wilting is traditionally measured on a $0 - 4$ scale,

in which 0 indicates a plant with no wilting, $1 = 1 - 25\%$, $2 = 26 - 50\%$, $3 = 51 - 75\%$ and a

score of 4 indicates a plant with $76 - 100\%$ wilted leaves [28]. While it is straightforward to assess

the ends of the spectrum, rating plants with scores of 2 or 3 is particularly difficult. This is due to

the subjective nature of visually determining when a leaf has lost sufficient turgor to qualify as

wilted. Reliable disease phenotyping is critical for identifying QTL for resistance to *Ralstonia*

and the development of resistant varieties.

Here, we used image-based phenotyping to identify new tomato genetic resistance loci to

Ralstonia. We developed a semi-automated imaging and trait analysis pipeline to quantify

bacterial wilt disease in a recombinant inbred line (RIL) population derived from *Ralstonia*-

resistant and susceptible tomato genotypes. We found both unique and shared QTL between our

image-based traits and plant wilting scored by the human eye. At least one of the QTL was

detected by image-based phenotyping before the onset of visual symptoms, demonstrating that

image-based phenotyping captures the disease phenotype at early stages of infection. These

results demonstrate that imaged-based phenotyping can shed light on new aspects of disease and

improve our ability to identify genetic loci crop resistance.

Results

Development of an aboveground imaging and semi-automated analysis pipeline

We first constructed a simple, low-cost imaging system that allowed us to semi-automate

aboveground disease phenotyping. Each plant was placed on a commercially available turntable,

and plants were imaged with a Canon DSLR (Supplementary Figure 1; details in methods). The

turntable and camera were connected with Photocapture 360 (Ortery technologies), which

allowed us to automatically capture images every 45 degrees (8 images per plant). Each image

included a fiducial marker for post-image color correction. Plants in the F_9 generation from a

RIL population derived from a cross between resistant Hawaii 7996 (H7996) and susceptible

West Virginia 700 (WV) were imaged the day before inoculation with *R. solanacearum* strain

K60, and at 3, 4, 5 and 6 days post inoculation (dpi). At 3 dpi, symptoms were not present in

susceptible parent WV, but by 6 dpi these plants were completely wilted. We imaged five

replicate plants of each RIL as well as the parental lines. Using this system, we captured over

40,000 images for high-resolution disease phenotyping. The same set of plants was also visually

scored by the human eye at 8 dpi. Visual scoring was based on the percentage of wilted leaves.

Tomato phenotypes in the RIL population ranged from highly susceptible to highly resistant,

consistent with the quantitative nature of disease resistance (Supplementary Figures 2 and 3).

We next developed a set of mathematical descriptors to phenotype wilting over time in our

images. We used 10 image-based traits (Table 1) to describe different aspects of plant wilting:

year	Variable	Hawaii 7996				West Virginia 700				RILs			
		Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum
2016	Wilting_Score	0	0	0	0	100	0	100	100	44.41	41.4	0	100
	Wilting_Score	0	0	0	0	99.33	5.16	60	100	53.69	42.23	0	100
	convex area	2.7e+06	1.4e+06	-4.1e+04	6.3e+06	-3.6e+05	3.6e+05	-1.1e+06	7.4e+05	1.2e+06	1.4e+06	-2.0e+06	7.9e+06
	convex width	8.7e+02	4.4e+02	-7.0e+01	1.7e+03	-7.4e+02	3.7e+02	-1.4e+03	5.4e+02	-4.6e-01	5.9e+02	-1.9e+03	2.9e+03
	plant area	8.7e+05	5.6e+05	-1.0e+04	2.2e+06	-1.1e+05	1.1e+05	-3.4e+05	2.1e+05	5.0e+05	5.5e+05	-7.4e+05	2.6e+06
2019	plant height	5.6e+02	1.9e+02	-7.0e+01	8.7e+02	-1.3e+01	1.7e+02	-3.6e+02	3.7e+02	3.3e+02	2.5e+02	-6.5e+02	1.1e+03
	plant width	4.0e+02	2.0e+02	-3.6e+01	8.2e+02	-3.5e+02	1.8e+02	-6.6e+02	2.5e+02	-2.1e+01	2.7e+02	-9.7e+02	8.4e+02
	X mass	3.1e+02	1.6e+02	-3.7e+01	5.8e+02	-2.9e+02	1.5e+02	-5.6e+02	2.0e+02	-1.2e+01	2.3e+02	-7.4e+02	6.9e+02
	Y mass	5.4e+02	2.1e+02	-8.4e+01	9.3e+02	-6.7e+01	1.6e+02	-4.1e+02	3.6e+02	2.1e+02	2.2e+02	-7.0e+02	9.6e+02
	CM width	3.1e+02	1.6e+02	-1.0e+02	6.0e+02	-2.9e+02	1.5e+02	-5.2e+02	1.8e+02	2.7e+01	2.3e+02	-7.0e+02	9.8e+02
	CM height	2.6e+02	1.8e+02	-2.1e+02	6.2e+02	-1.3e+02	1.5e+02	-3.9e+02	1.9e+02	-5.2e+01	1.9e+02	-8.8e+02	8.3e+02
	color average	2.88	5.44	-5.96	22.9	7.37	2.84	1.03	14.83	4.06	4.48	-11.35	16.48

Table 1 Descriptive statistics for 10 image-based traits and two years of visually assessed wilting scores (2016 and 2019). Image-based traits were from the same plants as the visually assessed wilting score in 2019. The RIL population was inoculated with Ralstonia and visually scored for wilting in 2016, but images were not taken. Data is shown for resistant parent Hawaii 7996, susceptible parent West Virginia 700, and the RIL population.

convex area, convex width, plant area, plant height, plant width, X mass, Y mass, center of mass

(CM) height, CM width, and color. Several of these, such as the area and width of the convex

hull, are traditional methods of describing aboveground plant shape. Because the center of mass

of a plant leaf changes as a plant wilts, we developed additional descriptors based on the distance

of the leaf center of mass from the stem (CM width and CM height, X mass and Y mass). We

then developed a pipeline which used the original image as input, performed color correction,

and quantified each descriptor (Figure 1 and Supplementary Figure 4, and methods).

Image-based traits differentiate resistant and susceptible plants

To validate the efficiency of the image-based descriptors to estimate wilting phenotypes, we

tested whether image-based phenotyping descriptors could differentiate resistant from

susceptible plants. Supplementary Figure 5 shows the average normalized score for each of the

Figure 1: Diagram of the semi-automated analysis pipeline and 10 image descriptors

image-based descriptors for each parent and the RIL population from -1 (the day before

inoculation) to 6 dpi. Most descriptors, particularly those based on plant width or convex hull,

had clearly divergent values in resistant and susceptible plants at a given time point. RIL

descriptor values ranged from those of the resistant to susceptible parents and occasionally

showed transgressive segregation (Supplementary Figure 5). Because plant shape at 6 dpi

depends on shape of the same plant at -1 dpi, we used the evolution of each descriptor from day

-1 to 6 dpi in our QTL analysis. This evolution was termed a 'trait'. Trait values were clearly

different for resistant and susceptible parents for all traits except color (Figure 2), which was not

used further.

Figure 2: Boxplots showing the evolution of the normalized values between -1 and 6 dpi for each of the five biological replicates for the 10 image-based traits for resistant Hawaii 7996, susceptible West Virginia parents and the 166 RILs. Color was unable to differentiate resistant from susceptible plants and was not used further.

Image-based phenotyping reveals drivers of the wilting phenotype

We next investigated whether any of our traits were major components of the wilting phenotype.

We visually scored plants and asked how well our image-based traits correlated with human

visual scoring. Wilting is categorized by loss of plant leaf turgor that results in drooping leaves,

and decreased plant width and height. Determining how much a plant has wilted is challenging,

in part because it can be difficult to quantify how much each a leaf has drooped and how much

drooping of one leaf correlates with whole plant wilting.

We aimed to quantify leaf drooping using center of mass traits. Among our image-based traits,

those which were functions of the leaf center of mass were highly inversely correlated with

visual wilting (i.e. as a plant wilts, the center of mass decreases), suggesting that these are major

drivers of the wilting phenotype. These traits included CM width, plant width and X mass ($r > -$

0.75) (Figure 3).

Figure 3: Heatmap showing the Pearson correlation values between image-based traits used in our QTL analysis and the human eye based wilting score in 2019. The correlations values were determined using the same plants imaged and visually assessed in 2019.

Several of our traits describe similar aspects of plant shape, such as height or width, through

different methods. These traits tended to be highly correlated with each other. For example, plant

height vs Y mass use different methods to describe plant width (based on the plant mask or the

center of mass of the stem masks; see Methods), and were highly positively correlated with each

other (0.93; Figure 3).

We trained a random forest, consisting of 1000 decision trees, to use data from 6 dpi to predict

the expert visual score assigned at 8 dpi. We used 969 plants in a 60:40 training-testing data

split, and achieved a classification accuracy of 83%. To identify the traits that provided the most

efficient estimation of wilting symptoms, we randomly scrambled the nine image-based traits

one at a time. Plant width and plant area had the most impact on wilting score prediction

(Supplementary Figure 6).

QTL analysis identifies 20 wilting QTL in 10 clusters across the tomato genome at 6 dpi

Our overall goal was to identify tomato genomic regions that provide resistance or susceptibility

to *Ralstonia*. Prior to this analysis, we first generated a genetic map using Genotyping-by-

Sequencing (GBS). We identified 632 high-quality SNPs for linkage mapping using GBS. We

combined these with 112 SolCap markers, and subsequently generated a linkage map using ICI

mapping software [29]. Our linkage map consisted of ~1300 cM (Figure 4) with an average per

chromosome marker density that varies from 1.8 to 7.48 cM (Supplementary Table 1).

For QTL analysis, we mapped the evolution of each our nine image-based descriptors from -1 to

6 dpi. In addition to the image-based phenotyping, we used two years (2016 and 2019) of

visually scored phenotyping data at 8 dpi. Visually assessed wilt scores from 2019 were

quantified from the same plants that were used for image-based phenotyping. In 2016, plants

were only visually scored for wilting, and no image-based measurements were taken. Using ICI

mapping software and composite interval mapping (CIM) for all traits, we identified 20 QTL

within the RIL population with a LOD score above 3 (Table 2). To be consistent with previous

studies of tomato-*Ralstonia* QTL mapping, we call these QTL, 'Bacterial wilt resistance (*Bwr*)

QTL'. Each *Bwr* QTL explained approximately 6 to 11% of the variation in response to *R.*

solanacearum strain K60.

Chromosome	Cluster	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI	RightCI
2	Bwr2.1	45	solcap_snp_sl_18519	SL3.0ch02-49265134	plant width	3.4108	6.4576	-0.1669	42.5	47.5
2	Bwr2.2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	X mass	3.4765	9.4924	-0.2178	69.5	73.5
2	Bwr2.2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	CM width	3.2455	8.455	-0.2189	69.5	73.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	convex width	5.3238	10.2506	0.2103	21.5	26.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	4.7309	9.1511	0.1976	21.5	25.5
3	Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	3.4765	5.8012	0.1705	21.5	26.5
3	Bwr3.1	23	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.6627	6.2716	0.1887	21.5	27.5
3	Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	4.5107	7.6902	0.1896	21.5	31.5
3	Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.3918	9.3283	0.1708	21.5	31.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	Wilting Score 2016	3.5258	7.3736	-0.1726	35.5	39.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	convex area	3.4328	9.0796	0.1953	34.5	39.5
3	Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	plant area	3.3019	8.6697	0.1902	34.5	39.5
4	Bwr4.1	4	SL3.0ch04-60431371	solcap_snp_sl_11543	convex width	3.1855	6.0199	0.1627	1.5	10.5
5	Bwr5.1	100	solcap_snp_sl_22649	SL3.0ch05-65105663	plant height	3.6115	9.4852	0.2103	95.5	103.5
6	Bwr6.1	79	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2016	4.077	8.7626	-0.1871	76.5	81.5
6	Bwr6.1	81	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2019	3.2673	11.0473	-0.1621	78.5	81.5
10	Bwr10.1	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	7.6897	-0.1811	50.5	54.5
10	Bwr10.1	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	6.8155	-0.1714	50.5	54.5
12	Bwr12.1	37	solcap_snp_sl_1525	solcap_snp_sl_1572	CM height	4.1755	10.7222	0.1837	33.5	40.5
12	Bwr12.2	42	solcap_snp_sl_58807	solcap_snp_sl_9707	Y mass	3.3249	8.5939	0.1651	40.5	43.5

Table 2 Overview of the 20 wilting QTL identified in 10 clusters across the tomato genome at 6 dpi. LOD: maximum value of the Logarithm of the odd. PVE: Percentage of phenotypic variance explained. Add: Additive effect. Left CI and Right CI are the confidence interval calculated by a one-LOD decrease from the estimated QTL position.

Figure 4: Genetic linkage map constructed with SolCap and SNP markers. Location of the 10 QTL clusters are displayed using blue and green for left and right markers respectively. Location of previously identified QTL on chromosome 6 and 12[27] are also displayed.

The parent that donated the favorable allele was determined according to the sign of the QTL

additive effect [30], where a positive sign referred to H7996 resistant parent and a negative sign to

WV susceptible parent. Typically, in QTL analysis the favorable allele has the higher trait value.

Here, a higher trait value is favorable in all cases except the visual wilting score, in which a

higher trait value was associated with susceptibility (e.g. 90% wilting is more susceptible than

20% wilting). For QTL detected using our visual wilting score, the susceptibility allele was

contributed by the susceptible parent WV (Table 2), consistent with other tomato-*R*.

solanacearum QTL studies based on visual assessment of wilting. For all QTL except those on

chromosome 2 and 10, the favorable allele was contributed by Hawaii 7996. QTL clusters on

chromosome 2 and 10, which contributed to plant height and width, were donated by the

susceptible parent WV. Favorable allelic contribution from both resistant and susceptible parents

is common in QTL studies for resistance [30,31].

Among the 20 individual *Bwr* QTL were 10 QTL clusters (Table 2). We use the term 'QTL

cluster' to describe QTL for different traits that co-localize at the same left and right genetic

marker. There are between one and six *Bwr* QTL within a given cluster, and each QTL within

the cluster has different LOD scores and explains a different percentage of phenotypic variation.

Together the clusters explained more than 88% of the variation (calculated using the sum of the

QTL with the highest PVE in each cluster).

Bwr QTL for traits that are highly correlated with each other (Figure 3) tended to cluster

together. For example, a cluster of QTL on chromosome 3 (*Bwr3.1*; Table 2) contains six *Bwr*

QTL, including three for area-related traits (convex width, and X mass and CM width), which

are correlated with each other at r = 0.94 - 0.97 (Figure 3). In another cluster (*Bwr3.2*) on this

same chromosome, *Bwr* QTL for traits that describe plant area were detected together. This co-

localization supports the robust nature of our analyses.

In other clusters, only one trait that described one aspect of the wilting phenotype was present.

For example, despite several metrics that describe width, only convex width was identified as a

QTL on chromosome 4 (*Bwr4.1*; LOD 3.18; PVE = 6%). This suggests that the image-based

phenotyping captured genetic variation that is specific to each trait.

Image-based phenotyping identifies three types of QTL clusters

Among our *Bwr* QTL clusters, we identified three types: those found using both image-based

and visual phenotyping (1 cluster, *Bwr3.2*), those found only through visual phenotyping (1

cluster, *Bwr6.1*), and those that were identified only through image-based phenotyping (8

clusters; Table 2 and Figure 5).

For the cluster including both image-based and visual phenotyping, (*Bwr3.2*) on chromosome 3,

we detected QTL for convex area, plant area and visual plant wilting in 2016 (Table 2 and Figure

5a). Previous studies examining tomato responses to other strains of *Ralstonia* have not detected

Bwr QTL in this region, indicating that it may be a novel target for *R. solanacearum* strain K60.

The identification of both image-based and visual QTL within the same cluster suggests that our

phenotyping methods are robust at identifying features of plant wilting. Close to this cluster is

Bwr3.1, which detected QTL for traits based on plant width, including convex width and CM

Figure 5: Significant QTL clusters on chromosomes 3 (a.), 6 (b.), 12 (c.) and 10 (d.). The image-based and visually assessed traits are represented with different colors for each chromosome. The vertical axis represents the genetic position (cM) and horizontal axis shows the LOD score. For each chromosome, the left panel represents the entire chromosome and the right panel represents the significant QTL cluster regions.

width. The proximal arm of chromosome 3 may be an important but unexploited region for

defense against *R. solanacearum* strain K60.

On chromosome 6 we detected *Bwr* (*Bwr6.1*) using only visual assessment for both 2016 and

2019 (Table 2, Figure 5b). This region may be a 'hot spot' for QTL for resistance to multiple

strains of *Ralstonia*. Previous studies using this same tomato population for mapping *Bwr* to

other strains of *Ralstonia* detected QTL approximately 4 Mb from this position (Figure 4).

In all other clusters, QTL were identified using only image-based phenotyping. Two such

clusters, detected on chromosome 12, identified QTL for CM height (*Bwr12.1*) and Y mass

(*Bwr12.2*), and were located near previous QTL detected against other strains of *Ralstonia*

(Figure 4 and Figure 5c) . These previous studies [24,26,27] detected QTL using visual wilt

assessment of adult plants in the field.

The remaining QTL detected through image-based phenotyping have not been previously

identified in studies using other strains of *Ralstonia*. These regions may be effective only against

specific strains of *Ralstonia* ('strain-specific' QTL), or may include those that are not easily

detected using visual assessment. In support of the latter, at one of the image-based-only clusters,

Bwr10.1 (Figure 5d), we detected a QTL using visual assessment that did not meet our threshold

for significance after permutation (Supplementary Table 2). Thus, image-based phenotyping

improved our ability to detect QTL.

Image-based phenotyping identified Bwr QTL prior to visual symptom development

In our system, wilting symptoms begin to appear on highly susceptible plants at 4 dpi, and these

plants are nearly 100% wilted by 6 dpi. To test whether we could detect *Bwr* QTL prior to the

onset of visual symptoms, we performed QTL analysis at 3, 4 and 5 dpi. No QTL were identified

based on visual assessment at any of these time points, however several QTL were identified

based on image-based phenotyping. At 3 dpi, three clusters were detected on chromosome 3, two

of which co-localized with those identified at 6 dpi (Table 3). *Bwr3.1*, a cluster detected only by

image-based phenotyping, and *Bwr3.2*, a cluster detected by both visual and image-based

phenotyping, were first detected at 3 dpi. *Bwr3.1* was also detected at 5 and 6 dpi, while *Bwr3.2*

was detected 3, 4, 5 and 6 dpi (Table 3). A small number of additional QTL were detected at

Clusters	Position	LeftMarker	RightMarker	Timepoint	TraitName	LOD	PVE	Add	LeftCI	RightCI
Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	3	CM height	7.2477	18.2665	0.2584	21.5	30.5
Bwr3.1	27	solcap_snp_sl_14355	solcap_snp_sl_9689	3	CM width	3.5022	8.8528	0.1536	22.5	32.5
Bwr3.1	26	solcap_snp_sl_14355	solcap_snp_sl_9689	3	plant width	5.5180	13.9255	0.2191	21.5	27.5
Bwr3.1	26	solcap_snp_sl_14355	solcap_snp_sl_9689	3	X mass	4.5407	11.7470	0.1991	22.5	32.5
Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	5	CM height	3.6770	9.3886	0.1414	21.5	27.5
Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	5	plant width	3.4953	9.3017	0.1806	21.5	27.5
Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	5	plant height	3.6851	8.5381	0.1738	21.5	31.5
Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	5	Y mass	3.4687	9.9830	0.1673	21.5	31.5
Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	6	convex width	5.3238	10.2506	0.2103	21.5	26.5
Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	6	plant width	4.7309	9.1511	0.1976	21.5	25.5
Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	6	plant height	4.5107	7.6902	0.1896	21.5	31.5
Bwr3.1	23	solcap_snp_sl_14355	solcap_snp_sl_9689	6	CM width	3.6627	6.2716	0.1887	21.5	27.5
Bwr3.1	22	solcap_snp_sl_14355	solcap_snp_sl_9689	6	X mass	3.4765	5.8012	0.1705	21.5	26.5
Bwr3.1	25	solcap_snp_sl_14355	solcap_snp_sl_9689	6	Y mass	3.3918	9.3283	0.1708	21.5	31.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	3	convex width	5.6025	13.8198	0.2120	37.5	42.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	4	convex width	3.9750	10.2036	0.1829	37.5	43.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	4	CM width	4.5294	8.4260	0.1853	37.5	43.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	4	plant width	3.8768	9.9017	0.1838	37.5	43.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	4	X mass	3.8156	9.7282	0.1837	37.5	43.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	5	convex width	3.7859	9.8306	0.1891	37.5	43.5
Bwr3.2	39	solcap_snp_sl_21215	solcap_snp_sl_9663	5	CM width	4.0630	7.4955	0.1924	35.5	39.5
Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	6	Wilting Score 2016	3.5258	7.3736	-0.1726	35.5	39.5
Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	6	convex area	3.4328	9.0796	0.1953	34.5	39.5
Bwr3.2	38	solcap_snp_sl_21215	solcap_snp_sl_9663	6	plant area	3.3019	8.6697	0.1902	34.5	39.5

Table 3 Overview of the QTLs Bwr3.1 and Bwr3.2 identified on chromosome 3 at 3, 4, 5 and 6 dpi by image-based and visually assessed traits.

earlier time points (Supplementary Table 3), but in most cases these were not stable, since they

were not present at 6 dpi. These data suggest that image-based phenotyping can identify *Bwr*

QTL prior to the onset of wilting symptoms.

Plant architecture QTL do not overlap with Bwr QTL

The parents of the RIL population, H7996 and WV, have different aboveground phenotypes

(Supplementary Figure 4) and the shoot architecture of the RILs correspondingly varies. To

ensure that *Bwr* QTL were the result of tomato responses to *Ralstonia*, and not due to differences

in aboveground plant architecture, we used our traits in a QTL analysis at -1 dpi, the day before

plants had been inoculated. We call these 'tomato plant architecture (*Tpa*)' QTL. We identified

16 *Tpa*, within 9 QTL clusters (Table 4). None of the 16 *Tpa* were detected within the same

interval as *Bwr* at 6 dpi. However, two *Tpa* QTL clusters were detected that were near *Bwr* QTL

present at 6 dpi. One of these was detected on chromosome 3 at position 30, between *Bwr3.1* and

3.2. While we cannot rule out the possibility that the same genes underlie these *Tpa* and *Bwr*

Chromosome	Cluster	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI
2	Tpa2.1	117	solcap_snp_sl_8510	solcap_snp_sl_15574	plant width	3.3851	6.4842	0.129	114.5
2	Tpa2.1	117	solcap_snp_sl_8510	solcap_snp_sl_15574	X mass	4.1424	6.8279	0.139	114.5
3	Tpa3.1	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.6327	8.0934	0.1522	24.5
3	Tpa3.2	58	SL3.0ch03-61078160	solcap_snp_sl_10372	plant width	3.2538	9.4831	0.1555	56.5
3	Tpa3.2	58	SL3.0ch03-61078160	solcap_snp_sl_10372	X mass	3.8291	9.4218	0.1628	56.5
3	Tpa3.2	59	SL3.0ch03-61078160	solcap_snp_sl_10372	CM height	3.794	3.6957	0.1603	56.5
4	Tpa4.1	128	SL3.0ch04-1213025	solcap_snp_sl_63869	CM height	3.8063	12.0286	-0.2891	127.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	convex area	4.741	11.2755	0.1621	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant width	3.5865	6.9265	0.1336	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant area	3.5117	7.3344	0.1457	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	X mass	4.8686	8.016	0.151	8.5
6	Tpa6.1	10	SL3.0ch06-13533085	SL3.0ch06-3310762	Y mass	4.902	9.3612	0.1757	8.5
8	Tpa8.1	80	solcap_snp_sl_21461	SL3.0ch08-63373341	X mass	3.2439	5.3622	0.123	74.5
9	Tpa9.1	36	solcap_snp_sl_26683	SL3.0ch09-4516570	CM height	4.8307	6.146	0.2081	32.5
9	Tpa9.2	67	SL3.0ch09-39293163	SL3.0ch09-63338944	convex area	3.1976	7.523	-0.1323	60.5
10	Tpa10.1	71	solcap_snp_sl_8835	Le013158s_161	Y mass	3.2077	5.9734	0.1404	68.5

Table 4 Overview of tomato plant architecture (Tpa) QTL identified at -1 dpi by image-based phenotyping.

QTL, their different marker intervals, coupled with the visual wilting QTL that is within the

Bwr3.2 cluster, suggests that different genes are responsible for these QTL. Together, these data

suggest that our image-based phenotyping detected genomic regions that function in responding

to *Ralstonia* and *Bwr* QTL are not the result of differential growth patterns within the RILs.

Discussion

Breeding for plant disease resistance is one of the best strategies to combat plant diseases and

prevent major crop loss, but is challenging in part due to the complicated nature of disease

phenotyping. Here we used image-based phenotyping with RGB images to identify 20 QTL in

10 clusters for tomato responses to *Ralstonia* at 6 dpi, two of which were detected as early as 3

dpi. Together, the 20 *Bwr* QTL at 6 dpi explained more than 88% of the variation in response to

Ralstonia. Image-based phenotyping for shape-based traits that were correlated with wilting

detected both novel loci and those that overlapped with QTL based on human visual assessment.

These results establish the importance and feasibility of quantitative imaged-based phenotyping

to identify new genetic targets for crop disease resistance.

Benefits of image-based phenotyping

Imaged-based phenotyping has been used to detect QTL associated with plant root [32] and shoot

[33–35] architecture, plant height [36], salt stress [37], and yield [38,39] among other traits. Although image-

based phenotyping has become increasingly common to quantify plant disease symptoms [4–7,11–

16], to our knowledge, it has yet to be used in the identification of new genetic loci for disease

resistance. One reason may be that many plant disease symptoms often occur at the leaf scale

(such as spots and specks), making it difficult to use some sensors in high-throughput or at the

proper resolution needed to assess disease. While wilting does occur at the leaf scale, it is an

easier phenotype to assess at the whole plant scale. Another possibility is that imaging sensors

are often expensive, making it more challenging to phenotype the large number of plants needed

in a QTL or GWA study.

Our image-based phenotyping identified loci that function in tomato responses to *Ralstonia*, but

did not detect one of the two *Bwr* clusters (*Bwr6.1*) we found using visual assessment of disease.

Visual assessment was performed at 8 dpi, while our image based phenotyping ended at 6 dpi. It

is possible *Bwr6.1* is not apparent until very late stages of disease. Alternatively, these results

could mean that the visually assessed wilting phenotype relies on human complexity that cannot

be broken into individual traits, or that we missed components of wilting in our image-based

traits. However, a more likely explanation is that our image-based traits were averaged across

eight two dimensional images, but wilting is a three dimensional (3D) phenotype. Images in 3D

may allow detection of *Bwr6.1*. It would also be of interest to test whether using additional types

of sensors – hyperspectral for example - would identify *Bwr 6.1*. Multi and hyperspectral

imaging has been used to detect the chemical changes in disease [12,14,40] and would likely be very

useful for quantifying water-stress changes associated with diseases like wilting.

New QTL for responses to Ralstonia in tomato

As in tomato, resistance to *Ralstonia* in other Solanaceous crops is quantitative [20–26,41]. Breeding

for resistance QTL is thus the primary way forward to developing *Ralstonia*-resistant crops in the

Solanaceae. This has not been easy, in part because of the diversity of *Ralstonia*. The *Ralstonia*

solanacearum species complex was recently subdivided into three species (*R. solanacearum*, *R.*

pseudosolanacearum, and *R. syzygii*), each prevalent in a different part of the world [42–44]. Each

species has multiple strains, with an overlapping, but distinct set of virulence proteins that

promote disease [45]. Varieties with effective resistance will likely have QTL that are effective

against local strains (strain-specific QTL) as well as those effective against multiple strains

(broad-spectrum QTL).

Most previously identified QTL have focused on resistance to *R. pseudosolanacearum*, and none

of them used the strain of *R. solanacearum* as in this work [20–27]. Using visual assessment of

wilting and the same RIL population of tomato (H7996 x WV700) used here, one broad-

spectrum QTL for resistance to multiple strains of *R. pseudosolanacearum,* and one strain of *R.*

solanacearum (JT-516) had been previously identified on chromosome 6 [20–27]. We also identified

a QTL on chromosome 6, using visual assessment in both 2016 and 2019. In our study, *Bwr6.1*

confers 8 − 11% of the variation, compared to 11.5 – 33% for *Bwr6a - Bwr6d* [26,27]. The genes

within the left and right intervals of *Bwr6.1* are particularly interesting because there are two

groups of NBS-LRRs located between the left and right markers for *Bwr6.1*

(solcap_snp_sl_14458 and the GBS-identified SNP at 39435402). One of these groups includes

the *S. lycopersicum* homolog of Arabidopsis *RESISTANCE to PSEUDOMONAS SYRINGAE 4*

(*RPS4*). Fine-mapping of *Bwr6.1* is needed to determine whether a specific NBS-LRR protein or

receptor is responsible for the resistance conferred by this region.

Previous studies also identified a major QTL on chromosome 12 that is effective for resistance to

R. pseudosolanacearum, and explained between 15.9 – 53.9% of the variation [27]. We found two

clusters of QTL on chromosome 12, for image-based traits. Within the region spanning *Bwr12.1*

on chromosome 12 is one NBS-LRR disease resistance gene (*Solyc12g10740*) and one receptor-

like kinase (*Solyc12g010660*) while within *Bwr12.2* are genes related to disease resistance

pathways, including a gene encoding a Pathogenesis Response (PR) Protein (*Solyc12g014310*),

and two genes encoding Leucine Rich Repeat – Receptor Like Kinases (*Solyc12g014350* and

Solyc12g150105). Our results suggest that this region of chromosome 12 may be important for

resistance to multiple strains of *Ralstonia*. The QTL we identified on chromosomes 2, 3, 4, 5, 8

and 10 were not previously identified, and thus may be specific to *R. solanacearum* strain K60.

With the exception of *Bwr3.2*, these QTL were detected using only our image-based traits.

Together, our results establish the value of image-based disease phenotyping for uncovering

novel genetic components and new targets for quantitative disease resistance in crops. By

identifying new genetic loci, this type of phenotyping may enable more broad-spectrum and

durable resistance.

Methods

Plant growth

Seeds of 188 Hawaii7996 (H7996) x West Virginia700 (WV) recombinant inbred lines (RIL) in

the F8 generation were obtained from the Asian Vegetable Research and Development Center

(AVRDC) in June 2014. Seeds were propagated to the F_9 generation in field and greenhouse in

West Lafayette, IN in 2014 and 2015 and were used in QTL analysis in 2016 and 2019.

For all plants used in the imaging experiments and 2019 visual assessment of wilting, seeds were

sown into individually labeled 1801 traditional inserts that were placed into 1020 flats (Hummert

International, USA). Seeds in the 2016 experiment were sown into individually labeled 1203

inserts that were placed into 1020 flats. Seeds were sown into ProMix Propagation Mix supplied

by the Lilly Greenhouses and Growth Facility located at Purdue University, West Lafayette,

Indiana, USA. The individual pots were randomized and placed in a growth chamber at 28ºC,

relative humidity of 65% for a lighting cycle of 16 hours light/8 hours dark. The plants began

germination on day 4 and plants were inoculated with *R. solanacearum* strain K60 at 17 days

after planting when three true leaves were present. Trays were rotated in the growth chamber

throughout each experiment. In 2016, four seeds of each RIL were grown in the growth chamber

with parental controls, and each plant was visually assessed for wilting at 8 days post inoculation

(dpi). In 2019, one seed per RIL, along with parental controls, was grown in the growth chamber

for each of five independent replicates. Each plant was imaged at -1, 3, 4, 5, and 6 dpi, and

visually assessed at 8 dpi.

For plants used to genotyping with tomato SolCap markers, F_9 generation RIL seeds and parental

controls were grown in the greenhouse in 2-gallon pots with Metro Mix 510 soil. Plants were

grown for 6 weeks.

Ralstonia solanacearum growth and plant inoculation

Ralstonia solanacearum strain K60 (containing a GFP reporter) was grown on casamino acid-

peptone-glucose (CPG) agar containing tetrazolium chloride (TZC) in the dark for 48 hours at

28ºC as in [46]. Briefly, bacteria were resuspended in sterile water to a concentration of

approximately 2×10^8 colony forming units (CFU)/mL for each experiment. For each

experiment, the concentration of inoculum was confirmed through dilution plating. Pots of three-

leaf plants were lightly compressed to induce wounding similar to transplant handling in field

conditions. 60 mL of inoculum was applied to the surrounding soil using a serological pipet.

Visual assessment of bacterial wilt disease

Wilt scores were visually assessed 8 dpi. Wilt scores were calculated by counting the number of

wilted true leaves divided by the total number of true leaves on the plant. Plants were scored with

a 95% if the plant had all of its leaves wilted, excluding the topmost leaf. Plants were marked as

100% when the topmost portion of the stem was collapsed and wilted (Supplementary Figure 2).

Plant imaging

Plants were imaged the day before inoculation (-1 dpi) and then imaged on 3, 4, 5, and 6 dpi. A

Linco Linstor 2000-watt photo studio (Amazon, USA) was used as the backdrop, and Flora X

fluorescent lighting (Amazon, USA) was used to create a small photo studio. Individual plants

were placed on a Photocapture 360 turntable (Ortery Technologies) that was programmed to

capture eight images around the plant (every 45°). The images were captured using an EOS 6D

Canon camera with an EF 50mm f/1.4 USM lens on a stationary tripod. An imaging carriage

was created with a square petri dish and a fiducial marker attached to each side. A total of five

independent replicates for each RIL and the parental controls were imaged.

Phenotyping Pipeline and Image Analysis

From the plant images, ten traits were acquired for QTL analysis. A detailed description of the

acquisition process can be found in [47,48]. All traits were extracted based on plant color and shape.

To eliminate any color discrepancy between images, images were automatically color corrected

using a fiducial marker consisting of a colored checkerboard with known physical dimensions

and colors.

After color correction, plant pixels were segmented from the rest of the image by thresholding

channels in the L*A*B* color space. The resulting segmentation mask was improved with image

morphological operations to fill holes and remove noise generated by the thresholding. The stem

of the plant was identified using two neural networks, Mask R-CNN and U-Net. By locating the

stem of the plant, metrics were defined that reflect the inner morphology of the plant. To train

the neural networks, stem segmentation ground truth data were generated using Adobe

Photoshop and LabelMe (a Python-based annotation tool) to mark the location of the image

pixels belonging to the stem of the plant. Further details about this procedure are described in

[47,48].

The plant and stem masks were used in the subsequent color and shape analysis, which acquired

ten traits: total area of the plant mask (plant area), height of the plant mask (plant height),

maximum width of the plant mask (plant width), area of the convex hull (convex area), width of

the convex hull (convex width), color-based weighted average (color average), horizontal

distance between the center of mass of the left and right sides of the plant stem (CM width),

height of the center of mass (CM height), x-axis distribution of the center of mass (X mass), Y-

axis distribution of the center of mass (Y mass).

The plant area, plant height and plant width were calculated from the plant mask. When

calculating plant height, the upper 5% of the plant material was not included, which helped

eliminate the impact of small leaves growing at the top. Using functions available in *OpenCV*

Python Library, a convex hull was fit around the plant mask and the convex area and convex

width were calculated. For color analysis, the weighted average of the pixel values in the A*

channel was calculated, as this channel captures the green-magenta spectrum.

To calculate traits involving the center of mass (CM), the plant mask was split into left and right

halves using the stem mask [48], and the CM located for each half. The x-axis distance between

CMs (CM width) and the average height of the CMs (CM height) were used as traits.

The extension of just one leaf can have a major impact on measured plant width, generating a

disproportionately larger value. Capturing the plant material distribution inside the plant mask

can overcome this issue. Using the split plant mask, the horizontal and vertical distribution of

plant material was estimated. The distances at which 90% of the plant material was captured in

the horizontal and vertical directions were used as the traits 'X mass distribution' and 'Y mass

distribution'. The average of eight views around the plant were used for each trait value.

In the random forest classification, visual wilting scores were thresholded to 0 or 1, where $0 = 0$

to 50% wilting, $1 = 51 - 100\%$ wilting. Average trait values from eight views from 6 dpi for 969

plants (RILs and parental controls) were used across all five replicates. Some plants did not have

a trait value for that day and were not used in the algorithm.

DNA extraction, Marker generation and Genotyping by Sequencing (GBS)

Tomato SolCap markers and single nucleotide polymorphisms (SNPs) identified through GBS

were used for map creation. Leaf disc samples were collected from each RIL plant using a

biopsy punch and were sent to LGC, Biosearch Technologies. Genomic DNA was extracted and

genotyped for 128 SNPs from the Tomato SolCAP panel using LGC's KASP assay. One

hundred twelve SNPs were obtained for RILs using this method.

For GBS, genomic DNA (gDNA) was extracted from 188 F_9 individuals of the RIL population

and from parental plants using Trizol. gDNA was RNase-treated and cleaned with phenol

chloroform extraction. Sequencing library preparation was as described by [49]. Briefly, gDNA

was digested with *PstI* and 150 bp paired end sequencing was performed on two lanes of an

Illumina Hi-seq 2500 at the Purdue Genomics Facility.

Reads were mapped to the *Solanum lycopersicum* 3.0 genome using the methods of [50]. In brief,

reads were de-multiplexed, and adapter sequences were removed. Reads were filtered based on

the presence of a GBS barcode on the forward read (R1), which accounts for 95.4% of the PE

reads. For these R1, 97.3% had a paired R2 read, and 2.7% did not have a pair. R1 read pairs

and R1 singles reads were combined and assessed independent of pairing and treated as single

end reads for the remaining analyses (924,500,737 reads).

Reads were filtered to remove those with the presence of an internal restriction site, the lack of

restriction site hang sequence at the end, and for minimum length, which retained 93.0% of the

reads. There was a minimum of 17,561 reads, a maximum of 16,127,292 reads, and an average

of 4,480,149 reads per sample (Supplementary Figure 7). Reads were mapped using BWA-

MEM for paired end reads, and the GATK haplotype caller was used to generate a genomic

Variant Call Format (gVCF) file for downstream analysis.

There were 74,082 SNPs called between the population and *Solanum lycopersicum* 3.0 reference

genome. Many of these were SNPs between the reference genome and the population and do not

vary in the population used here. Filtering for the presence of two alleles within the RIL

population identified resulted in 2,738 SNPs. Of these, the parental alleles were identified in

both parents for 278 SNPs and in one parent for 698 SNPs. The remaining 1,762 SNPs could not

be assigned a parental origin and were not used for further analysis. The 976 SNPs with parental

origin identified were filtered for a minor allele frequency greater than 0.02 and less than 0.99,

which resulted in 632 high quality SNPs.

High quality SNPs identified from GBS were combined with previously defined SolCap markers

from LGC Biosearch Technologies for a total of 748 markers. SNP coverage of the 188 RIL taxa

was high. Of the 748 markers, the minimum coverage was 115 markers, the maximum coverage

was 555 markers, and the average was 366 markers (Supplementary Figure 8a). There was some

residual heterozygosity in the taxa (Supplementary Figure 8b), and taxa with 10% or greater

heterozygosity were removed from further analysis (2 RILs).

Linkage Map Construction

The software QTL IciMapping [29] (version 4.1) was used for the map construction with all 748

markers. Redundant markers were filtered by taxa coverage using a missing rate of 85% and a

distortion threshold at 0.001 to obtain a total of 408 markers. The 408 markers without anchor

information were assigned to 12 groups based on a LOD score threshold value of 3. After

grouping, the markers were ordered using the nearest neighbor algorithm (nnTwoOpt) using the

rippling criterion SARF (Sum of Adjacent Recombination Frequencies) with a window size of

five markers. After ordering, some markers at the end of the chromosomes were deleted when

they were adding an insignificant genetic distance to the chromosome. These markers were

identified after splitting the current chromosome in two sub-chromosomes between the longest

marker interval. If the shortest sub-chromosome contained more than 20% of the markers before

splitting, the two sub-chromosome were re-assembled. Otherwise, the shortest sub-chromosome

was deleted.

QTL detection

QTL were detected using Inclusive Composite Interval Mapping with additive effects (ICIM-

ADD) in IciMapping [29] (version 4.1) using a genetic mapping with a 1 cM scanning step and a

probability in stepwise regression of 0.001. The LOD significance threshold to declare a QTL

significant was determined using a Type-I error of 0.0500 after one thousand permutation. For

the QTL analysis, we removed 19 of the 188 RIL (10%) showing the highest variation according

to the visual wilting score and 3 RIL due to missing data across the 5 biological repetitions for a

total of 166 RIL into the QTL map (Supplementary Figure 3). The evolution of the ten image-

based traits and two visual wilting scores between -1 dpi and 3, 4, 5 and 6 dpi were determined

for the 166 RIL. To assure a normal distribution of the phenotypic data, the function

"orderNorm" was used to perform an ordered quantile normalization [51] before QTL analysis

using the package "bestNormalize" version 1.6.1 with R software version 3.6.1 [52].

References
508

1. Colwell, R. N. Determining the prevalence of certain cereal crop diseases by means of aerial

photography. *Hilgardia* **26**, 223–286 (1956).

2. Jackson, R. D. Remote Sensing of Biotic and Abiotic Plant Stress. *Annu. Rev. Phytopathol.*

24, 265–287 (1986).

3. Bock, C., Poole, G., Parker, P. & Gottwald, T. R. Plant Disease Severity Estimated Visually,

by Digital Photography and Image Analysis, and by Hyperspectral Imaging. *Crit. Rev. Plant*

Sci. **29**, 59-107. (2010).

4. Simko, I., Jimenez-Berni, J. A. & Sirault, X. R. R. Phenomic Approaches and Tools for

Phytopathologists. *Phytopathology* **107**, 6–17 (2017).

5. Shakoor, N., Lee, S. & Mockler, T. C. High throughput phenotyping to accelerate crop

breeding and monitoring of diseases in the field. *Curr. Opin. Plant Biol.* **38**, 184–192 (2017).

6. Mochida, K. *et al.* Computer vision-based phenotyping for improvement of plant

productivity: a machine learning perspective. *GigaScience* **8**, (2019).

7. Mir, R. R., Reynolds, M., Pinto, F., Khan, M. A. & Bhat, M. A. High-throughput

phenotyping for crop improvement in the genomics era. *Plant Sci.* **282**, 60–72 (2019).

8. Bock, C., Parker, P., Cook, A. Z. & Gottwald, T. Visual Rating and the Use of Image

Analysis for Assessing Different Symptoms of Citrus Canker on Grapefruit Leaves. *Plant*

Dis. **92**, 530 - 541 (2008).

9. Stewart, E. & McDonald, B. Measuring quantitative virulence in the wheat pathogen

Zymoseptoria tritici using high-throughput automated image analysis. *Phytopathology* **104**,

985 -992 (2014).

10. Stewart, E. L. *et al.* An Improved Method for Measuring Quantitative Resistance to the

Wheat Pathogen *Zymoseptoria tritici* Using High-Throughput Automated Image Analysis.

Phytopathology **106**, 782–788 (2016).

11. Mahlein, A.-K. Plant Disease Detection by Imaging Sensors - Parallels and Specific

Demands for Precision Agriculture and Plant Phenotyping. *Plant Dis.* **100**, 241 - 251 (2016).

12. Mahlein, A.-K. *et al.* Plant disease detection by hyperspectral imaging: from the lab to the

field. *Adv. Anim. Biosci.* **8,** 238-243 (2017).

13. Mahlein, A.-K. *et al.* Comparison and Combination of Thermal, Fluorescence, and

Hyperspectral Imaging for Monitoring Fusarium Head Blight of Wheat on Spikelet Scale.

Sensors **19**, 2281 (2019).

14. Lowe, A., Harrison, N. & French, A. P. Hyperspectral image analysis techniques for the

detection and classification of the early onset of plant disease and stress. *Plant Methods* **13**,

80 (2017).

15. Pérez-Bueno, M. L., Pineda, M. & Barón, M. Phenotyping Plant Responses to Biotic Stress

by Chlorophyll Fluorescence Imaging. *Front. Plant Sci.* **10**, (2019).

16. Pineda, M., Barón, M. & Pérez-Bueno, M.-L. Thermal Imaging for Plant Stress Detection

and Phenotyping. *Remote Sensing* **13**, 68 (2021).

17. Mansfield, J. *et al.* Top 10 plant pathogenic bacteria in molecular plant pathology. *Mol. Plant*

Pathol. **13**, 614–629 (2012).

18. Genin, S. Molecular traits controlling host range and adaptation to plants in Ralstonia

solanacearum. *New Phytol.* **187**, 920–928 (2010).

19. Genin, S. & Denny, T. P. Pathogenomics of the *Ralstonia solanacearum* Species Complex.

Annu. Rev. Phytopathol. **50**, 67–89 (2011).

20. Danesh, D., Aarons, S., McGill, G. E. & Young, N. D. Genetic Dissection of Oligogenic

Resistance to Bacterial Wilt in Tomato. *Mol. Plant Microbe Interact.* **7**, 464–471 (1994).

21. Thoquet, P. *et al.* Quantitative trait loci determining resistance to bacterial wilt in tomato

cultivar Hawaii7996. *Mol. Plant Microbe Interact.* **9**, 826–836 (1996).

22. Thoquet, P. *et al.* Polygenic resistance of tomato plants to bacterial wilt in the French West

Indies. *Mol. Plant Microbe Interact.* **9**, 837–842 (1996).

23. Mangin, B., Thoquet, P., Olivier, J. & Grimsley, N. H. Temporal and multiple quantitative

trait loci analyses of resistance to bacterial wilt in tomato permit the resolution of linked loci.

Genetics **151**, 1165–1172 (1999).

24. Wang, J. F. *et al.* Resistance of tomato line Hawaii7996 to *Ralstonia solanacearum* Pss4 in

Taiwan is controlled mainly by a major strain-specific locus. *Mol. Plant Microbe Interact.*

13, 6–13 (2000).

25. Carmeille, A. *et al.* Identification of QTLs for *Ralstonia solanacearum* race 3-phylotype II

resistance in tomato. *Theor. Appl. Genet.* **113**, 110–21 (2006).

26. Wang J-W. *et al.* Identification of major QTLs associated with stable resistance of tomato

cultivar 'Hawaii 7996' to *Ralstonia solanacearum*. *Euphytica* **190**, 241–252 (2013).

27. Shin, I. S. *et al.* Construction of a single nucleotide polymorphism marker based QTL map

and validation of resistance loci to bacterial wilt caused by *Ralstonia solanacearum* species

complex in tomato. *Euphytica* **216**, 54 (2020).

28. Schandry, N. A Practical Guide to Visualization and Statistical Analysis of *R. solanacearum*

Infection Data Using R. *Front. Plant Sci.* **8**, (2017).

29. Meng, L., Li, H., Zhang, L. & Wang, J. QTL IciMapping: Integrated software for genetic

linkage map construction and quantitative trait locus mapping in biparental populations.

Crop J. **3**, 269–283 (2015).

30. Awata, L. A. O. *et al.* Genetic Analysis of QTL for Resistance to Maize Lethal Necrosis in

Multiple Mapping Populations. *Genes* **11**, 32 (2020).

31. Maschietto, V. *et al.* QTL mapping and candidate genes for resistance to Fusarium ear rot

and fumonisin contamination in maize. *BMC Plant Biol.* **17**, 20 (2017).

32. Topp, C. N. *et al.* 3D phenotyping and quantitative trait locus mapping identify core regions

of the rice genome controlling root architecture. *Proc. Natl. Acad. Sci. U.S.A.* **110**, E1695-

1704 (2013).

33. Zhang, X. *et al.* High-Throughput Phenotyping and QTL Mapping Reveals the Genetic

Architecture of Maize Plant Growth. *Plant Physiol.* **173**, 1554–1564 (2017).

34. Knoch, D. *et al.* Strong temporal dynamics of QTL action on plant growth progression

revealed through high-throughput phenotyping in canola. *Plant Biotechnology Journal* **18**,

68–82 (2020).

35. Li, H. *et al.* High-throughput phenotyping accelerates the dissection of the dynamic genetic

architecture of plant growth and yield improvement in rapeseed. *Plant Biotechnol. J.* **18**,

2345–2353 (2020).

36. Wang, X. *et al.* Dynamic plant height QTL revealed in maize through remote sensing

phenotyping using a high-throughput unmanned aerial vehicle (UAV). *Sci. Rep.* **9**, 3458

(2019).

37. Awlia, M. *et al.* Genetic mapping of the early responses to salt stress in *Arabidopsis*

thaliana. *Plant J.* doi:10.1111/tpj.15310 (2021).

38. Tanger, P. *et al.* Field-based high throughput phenotyping rapidly identifies genomic regions

controlling yield components in rice. *Sci. Rep.* **7**, 42839 (2017).

39. Pauli, D. *et al.* Field-Based High-Throughput Plant Phenotyping Reveals the Temporal

Patterns of Quantitative Trait Loci Associated with Stress-Responsive Traits in Cotton. *G3*

Genes Genom. Genet. **6**, 865–879 (2016).

40. Zhang, N. *et al.* A Review of Advanced Technologies and Development for Hyperspectral-

Based Plant Disease Detection in the Past Three Decades. *Remote Sens.* **12**, 3188 (2020).

41. Young, N. D. & Danesh, D. Understanding Bacterial Wilt Resistance in Tomato through the

Use of DNA Genetic-Markers. In: *Bacterial Wilt*: *the disease and its causative agent,*

Pseudomonas solanacearum, CAB International, Wallingford, 145–155 (1994).

42. Remenant, B. *et al.* Sequencing of K60, Type Strain of the Major Plant Pathogen *Ralstonia*

solanacearum. *J Bacteriol* **194**, 2742–2743 (2012).

43. Safni, I. *et al.* Polyphasic taxonomic revision of the *Ralstonia solanacearum* species

complex: proposal to emend the descriptions of *Ralstonia solanacearum* and *Ralstonia*

syzygii and reclassify current *R. syzygii* strains as *Ralstonia syzygii* subsp. *syzygii* subsp.

nov., *R. solanacearum* phylotype IV strains as *Ralstonia syzygii* subsp. *indonesiensis* subsp.

nov., banana blood disease bacterium strains as *Ralstonia syzygii* subsp. *celebesensis* subsp.

nov. and *R. solanacearum* phylotype I and III strains as *Ralstonia pseudosolanacearum* sp.

nov. *Int. J. Syst. Evol. Micr.* **64**, 3087–3103.

44. Prior, P. *et al.* Genomic and proteomic evidence supporting the division of the plant

pathogen *Ralstonia solanacearum* into three species. *BMC Genomics* **17**, 90 (2016).

45. Landry, D., González-Fuente, M., Deslandes, L. & Peeters, N. The large, diverse, and robust

arsenal of *Ralstonia solanacearum* type III effectors and their *in planta* functions. *Mol. Plant*

Pathol. (2020) doi:10.1111/mpp.12977.

46. Caldwell, D., Kim, B.-S. & Iyer-Pascuzzi, A. S. *Ralstonia solanacearum* Differentially

Colonizes Roots of Resistant and Susceptible Tomato Plants. *Phytopathology* **107**, 528–536

(2017).

47. Yang, C. *et al.* Image-Based Plant Wilting Estimation. *arXiv:2105.12926 [cs]* (2021).

48. Yang, C. *et al.* Plant Stem Segmentation Using Fast Ground Truth Generation. *2020 IEEE*

Southwest Symposium on Image Analysis and Interpretation (SSIAI) 62–65 (2020).

49. Elshire, R. J. *et al.* A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High

Diversity Species. *PLOS ONE* **6**, e19379 (2011).

50. Manching, H. *et al.* Phased Genotyping-by-Sequencing Enhances Analysis of Genetic

Diversity and Reveals Divergent Copy Number Variants in Maize. *G3 Genes Genom. Genet*

7, 2161–2170 (2017).

51. Peterson, R. A. & Cavanaugh, J. E. Ordered quantile normalization: a semiparametric

transformation built for the cross-validation era. *J. Appl. Stat.* **47**, 2312–2327 (2020).

52. R Core Team. R: A language and environment for statistical computing. *R Foundation for*

Statistical Computing **Vienna**, URL https://www.R-project.org/.

Acknowledgements

We thank Chloe Caldwell for help with imaging, and Kristina Gans for help with GBS.

This work was funded by a Purdue Seed Grant to AIP and ED, a Foundation for Food and

Agriculture Research New Innovator Award to AIP, and the endowment of the Charles William

Harrison Distinguished Professorship at Purdue University to ED.

Author Contributions

A.I.P. and E.D. designed the study and coordinated the project; V.M., D.C., and B.K. performed

experiments; E.E. carried out the GBS analysis; V.M. conducted the QTL analysis; V.M, D.C,

A.I.P., S.B., C.Y., and E.D. developed image descriptors; S.B. and C.Y. conducted the image

analysis; A.I.P. and V.M. wrote the manuscript; all authors edited the manuscript.

Supplementary Information

Supplementary Figures

Supplementary Figure 1: Design of our low cost phenotyping platform including automatic

turntable, backdrop, lightning and RGB camera.

Supplementary Figure 2: Raw RGB pictures showing the evolution of wilting symptoms on RIL

646 at -1, 3, 4, 5 and 6 dpi. Visually assessed wilting score are expressed in percentage of

wilted leaves.

689 Supplementary Figure 3: Wilting Score RIL: Boxplots showing the normalized wilting score at 6

690 dpi for the 166 RILs, resistant Hawaii 7996 (HA) and susceptible West Virginia (WV)

691 genotypes.

699   Supplementary Figure 4: RGB pictures showing the evolution of the traits at -1, 3, 4, 5 and 6 dpi

700   for A) Hawaii 7996 and B) West Virginia (pw: plant width, ph: plant height, CMh: Center of

701   Mass height, CMw: Center of Mass width, xm: Horizontal mass distribution, ym: Vertical mass

702   distribution).

Hawaii 7996

Supplementary Figure 5: Boxplots showing the normalized values for image-based traits at -1, 3, 4, 5, 6 dpi for resistant Hawaii 7996, susceptible West Virginia 700, and 166 individuals of the RIL population.

715   Supplementary Figure 6: Histogram showing the image-based descriptors which contribute the

716   most to wilting

733   Supplementary Figure 7: GBS Read Distribution by Sample (includes parental references).

748   Supplementary Figure 8: a) RIL and parental coverage and SNP density from GBS analysis; b)

749   Percent heterozygosity and SNP density

750
751

766   **Supplementary Tables**

767   Supplementary Table 1: Marker density per cM for each chromosome.

Chromosome	Number of marker	Chromosome Length (cM)	Average interval length (cM)	Standard deviation	Standard error
1	36	153.42	4.38	4.15	0.70
2	19	117.42	6.52	3.92	0.92
3	28	87.77	3.25	3.56	0.68
4	44	133.43	3.10	4.25	0.65
5	18	116.49	6.85	5.33	1.29
6	61	116.92	1.80	1.99	0.26
7	21	95.63	4.78	3.94	0.88
8	14	97.27	7.48	4.44	1.23
9	27	143.5	5.52	5.77	1.13
10	18	80.01	4.71	4.80	1.16
11	25	61.22	2.55	3.88	0.79
12	43	120.42	2.87	3.01	0.46

772   Supplementary Table 2: Overview of the 9 genomic regions identified with a LOD score superior

773   at 3 on chromosome 10 after inoculation with *R. solanacearum* at 6 dpi. According to the ICI

774   Mapping analysis using 1000 permutations, a threshold value of 3.48 represents the minimum

775   LOD to identify a significant QTL. Genomic regions with a lower LOD value did not reach our

776   threshold for significance after permutation.

Chromosome	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add
10	51	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0067	13.3810	0.2077
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0927	10.5192	0.1842
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	Wilting Score 2019	3.0832	8.0396	0.1611
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.1634	9.0040	-0.1835
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	7.8538	-0.1714
10	51	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.0398	11.1872	-0.2039
10	52	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.5411	10.2424	-0.1951
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	8.8147	-0.1811
10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	X mass	3.0584	7.0293	-0.1629

Supplementary Table 3: Overview of QTLs identified across the genome at -1, 3, 4, 5, 6 dpi with

R. solanacearum. LOD: maximum value of the Logarithm of the odd. PVE: Percentage of

phenotypic variance explained. Add: Additive effect, the positive and negative values indicated

that the alleles are introgressed from the resistant parent (Hawaii 7996) and susceptible parents

(WestVirginia). LeftCI and RightCI are the confidence interval calculated by a one-LOD

decrease from the estimated QTL position.

Timepoint	Chromosome	Position	LeftMarker	RightMarker	TraitName	LOD	PVE	Add	LeftCI	RightCI
-1	2	117	solcap_snp_sl_8510	solcap_snp_sl_15574	plant width	3.3851	6.4842	0.1290	114.5	117.0
-1	2	117	solcap_snp_sl_8510	solcap_snp_sl_15574	X mass	4.1424	6.8279	0.1390	114.5	117.0
-1	3	58	SL3.0ch03-61078160	solcap_snp_sl_10372	plant width	3.2538	9.4831	0.1555	56.5	64.5
-1	3	58	SL3.0ch03-61078160	solcap_snp_sl_10372	X mass	3.8291	9.4218	0.1628	56.5	64.5
-1	3	59	SL3.0ch03-61078160	solcap_snp_sl_10372	CM height	3.7940	3.6957	0.1603	56.5	63.5
-1	3	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.6327	8.0934	0.1522	24.5	34.5
-1	4	128	SL3.0ch04-1213025	solcap_snp_sl_63869	CM height	3.8063	12.0286	-0.2891	127.5	130.5
-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	convex area	4.7410	11.2755	0.1621	8.5	11.5
-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant width	3.5865	6.9265	0.1336	8.5	11.5
-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	plant area	3.5117	7.3344	0.1457	8.5	11.5
-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	X mass	4.8686	8.0160	0.1510	8.5	11.5
-1	6	10	SL3.0ch06-13533085	SL3.0ch06-3310762	Y mass	4.9020	9.3612	0.1757	8.5	11.5
-1	8	80	solcap_snp_sl_21461	SL3.0ch08-63373341	X mass	3.2439	5.3622	0.1230	74.5	83.5
-1	9	67	SL3.0ch09-39293163	SL3.0ch09-63338944	convex area	3.1976	7.5230	-0.1323	60.5	69.5
-1	9	36	solcap_snp_sl_26683	SL3.0ch09-4516570	CM height	4.8307	6.1460	0.2081	32.5	40.5
-1	10	71	solcap_snp_sl_8835	Le013158s_161	Y mass	3.2077	5.9734	0.1404	68.5	73.5
3	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	CM height	7.2477	18.2665	0.2584	21.5	30.5
3	3	27	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.5022	8.8528	0.1536	22.5	32.5
3	3	26	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	5.5180	13.9255	0.2191	21.5	27.5
3	3	26	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	4.5407	11.7470	0.1991	22.5	32.5
3	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	5.6025	13.8198	0.2120	37.5	42.5
3	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	plant height	4.5094	10.3345	0.1974	25.5	31.5
3	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	Y mass	5.7227	15.0239	0.2325	23.5	31.5
4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	3.9750	10.2036	0.1829	37.5	43.5
4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	CM width	4.5294	8.4260	0.1853	37.5	43.5
4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	plant width	3.8768	9.9017	0.1838	37.5	43.5
4	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	X mass	3.8156	9.7282	0.1837	37.5	43.5
4	3	28	solcap_snp_sl_9689	solcap_snp_sl_9681	convex area	3.1310	7.7809	0.1508	22.5	32.5
4	3	30	solcap_snp_sl_9689	solcap_snp_sl_9681	plant area	3.8685	9.9594	0.1672	23.5	33.5
5	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	CM height	3.6770	9.3886	0.1414	21.5	27.5
5	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	3.4953	9.3017	0.1806	21.5	27.5
5	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	3.6851	8.5381	0.1738	21.5	31.5
5	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.4687	9.9830	0.1673	21.5	31.5
5	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	convex width	3.7859	9.8306	0.1891	37.5	43.5
5	3	39	solcap_snp_sl_21215	solcap_snp_sl_9663	CM width	4.0630	7.4955	0.1924	35.5	39.5
5	5	107	SL3.0ch05-65105663	SL3.0ch05-66111590	plant height	3.4091	8.9327	0.1778	104.5	114.5
5	12	56	SL3.0ch12-62154155	solcap_snp_sl_19345	CM height	3.2828	8.2102	0.1360	54.5	59.5
6	2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	X mass	3.4765	9.4924	-0.2178	69.5	73.5
6	2	72	SL3.0ch02-45885016	SL3.0ch02-44375028	CM width	3.2455	8.4550	-0.2189	69.5	73.5
6	2	45	solcap_snp_sl_18519	SL3.0ch02-49265134	plant width	3.4108	6.4576	-0.1669	42.5	47.5
6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	convex width	5.3238	10.2506	0.2103	21.5	26.5
6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	plant width	4.7309	9.1511	0.1976	21.5	25.5
6	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	plant height	4.5107	7.6902	0.1896	21.5	31.5
6	3	23	solcap_snp_sl_14355	solcap_snp_sl_9689	CM width	3.6627	6.2716	0.1887	21.5	27.5
6	3	22	solcap_snp_sl_14355	solcap_snp_sl_9689	X mass	3.4765	5.8012	0.1705	21.5	26.5
6	3	25	solcap_snp_sl_14355	solcap_snp_sl_9689	Y mass	3.3918	9.3283	0.1708	21.5	31.5
6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	Wilting Score 2016	3.5258	7.3736	-0.1726	35.5	39.5
6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	convex area	3.4328	9.0796	0.1953	34.5	39.5
6	3	38	solcap_snp_sl_21215	solcap_snp_sl_9663	plant area	3.3019	8.6697	0.1902	34.5	39.5
6	4	4	SL3.0ch04-60431371	solcap_snp_sl_11543	convex width	3.1855	6.0199	0.1627	1.5	10.5
6	5	100	solcap_snp_sl_22649	SL3.0ch05-65105663	plant height	3.6115	9.4852	0.2103	95.5	103.5
6	6	79	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2016	4.0770	8.7626	-0.1871	76.5	81.5
6	6	81	solcap_snp_sl_14458	SL3.0ch06-39435402	Wilting Score 2019	3.2673	11.0473	-0.1621	78.5	81.5
6	10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	plant width	3.8661	7.6897	-0.1811	50.5	54.5
6	10	53	solcap_snp_sl_33168	SL3.0ch10-63377492	convex width	3.4883	6.8155	-0.1714	50.5	54.5
6	12	37	solcap_snp_sl_1525	solcap_snp_sl_1572	CM height	4.1755	10.7222	0.1837	33.5	40.5
6	12	42	solcap_snp_sl_58807	solcap_snp_sl_9707	Y mass	3.3249	8.5939	0.1651	40.5	43.5