Active enhancers strengthen insulation by RNA-mediated CTCF binding at TAD boundaries

Zubairul Islam[1,2*], Bharath Saravanan[1,2*], Kaivalya Walavalkar[1], Jitendra Thakur [3], Umer Farooq[1,4], Anurag Kumar Singh[1], Radhakrishnan Sabarinathan[1], Awadhesh Pandit[1], Steven Henikoff [5], Dimple Notani[1#]

Affiliations:

1. Genetics and Development, National Centre for Biological Sciences, Tata Institute for Fundamental Research, Bangalore, Karnataka, 560065. India

2. Sastra Deemed University, Thanjavur, Tamil Nadu, 613401. India

3. Department of Biology, Emory University, Atlanta, Georgia, 30322. USA

4. The University of Trans-Disciplinary Health Sciences and Technology, IVRI road, Bangalore, Karnataka, 560064. India

5. Basic Sciences Division, Fred Hutchinson Cancer Research Centre, 1100 N. Fairview Ave, Seattle, WA, 98109, USA

* Equal contributions

Corresponding author: dnotani@ncbs.res.in

Abstract

Vertebrate genomes are partitioned into Topologically Associating Domains (TADs), which are typically bound by head-to-head pairs of CTCF binding sites. Transcription at TAD boundaries correlates with better insulation, however, it is not known whether the boundary transcripts themselves contribute to boundary function. Here we characterize boundary-associated RNAs genome-wide, focusing on the disease-relevant *INK4a/ARF* TAD. Using CTCF site deletions and boundary-associated RNA knockdowns, we observe that boundary-associated RNAs facilitate recruitment and clustering of CTCF at TAD borders. The resulting CTCF enrichment enhances TAD insulation, enhancer:promoter interactions and TAD gene expression. Importantly, knockdown of boundary-associated RNAs results in loss of boundary insulation function. Using enhancer deletions and CRISPRi of promoters we show that active TAD enhancers but not promoters induce boundary-associated RNA transcription, thus defining a novel class of regulatory enhancer RNAs.

Introduction

The genomes of multicellular eukaryotes are organized into self-interacting units known as Topologically Associating Domains (TADs) (1). TADs are identified in Hi-C interaction matrices as contact domains, and contact domains that exhibit corner dots (at the apex of a loop) are known as loop domains (2). Loop and contact domains are both proposed to be formed by loop extrusion (1,3-7) and are dependent on CTCF and cohesin (6-10). The length of an extruded loop is determined by the presence and orientation of CTCF bound regions that interact within the loop (11). By preventing enhancers and promoters within the loop from interacting outside of the loops, boundary CTCF pairs insulate genes within a TAD (7-8,12-13). However, the presence of CTCF consensus motifs or their orientation at contact domains do not accurately predict TAD boundaries, suggesting that additional factors must influence CTCF binding and, therefore, TAD formation and insulation.

Mutation of CTCF sites at TAD boundaries affects gene expression, ranging from no effect to severe effect in the same and neighboring TADs (14-21). Mutation of all CTCF sites within a boundary region or depletion of CTCF may be required to display effects on transcription (20,22), suggesting redundancy among multiple CTCF sites at a TAD boundary. However, it is unclear how multiple CTCF sites within the boundary coordinate to achieve better insulation. Active transcription at TAD boundaries correlates with better insulation and CTCF occupancy (22-24) and has been speculated to increase TAD strength (25-27), consistent with observations that inhibition of transcription or Ribonuclease A (RNaseA) treatment disrupts 3D chromatin architecture (28-30). Furthermore, CTCF dimerization, important for loop extrusion, is RNA-dependent, and RNA binding of CTCF is required for TAD chromatin organization (31-32), but how CTCF:RNA interactions maintain TADs is not clear.

In some instances, TAD boundaries are found near sites of enhancers (33). Also, TADs containing
super-enhancers consist of multiple enhancers exhibit strong boundaries (25,34). However, it is
not known whether these enhancers play a role in boundary insulation. Moreover, boundaries
themselves display enhancer-like features (30,35), although the levels of H3K27ac, RNA
Polymerase II (PolII) and H3K4me3 on TAD boundaries are far less than are found on active
enhancers in general (36). These observations suggest a relationship between certain TAD
boundaries and enhancers, but the mechanistic basis for their putative interactions remains
unknown.

Here we show that boundaries of about half of human TADs are transcribed, either from a gene-
body that overlaps a boundary region or from a non-genic region that is transcribed within the
boundary. Compared to non-transcribed TAD boundaries, transcribed genic and non-genic
boundaries are more enriched for CTCF binding and are better insulators of genes within TADs,
which are more highly expressed. RNAs transcribed from non-genic (i.e. enhancer-like regions)
boundaries are more stable than RNAs expressed from enhancers at non-boundary regions.
Similarly, genic transcripts from boundaries are also more stable than other mRNAs. To test for
functional relevance of boundary-associated RNAs, we focused on the well-characterized
INK4a/ARF TAD, which is bound by CTCF sites and harbors a super-enhancer. Both deletions of
CTCF sites and knockdown of boundary-associated RNAs resulted in loss of enhancer:promoter
interactions within TADs while strengthening interactions with enhancers in regions outside the
TAD. These observations suggest that active enhancers within TADs promote TAD insulation by
triggering transcription of the boundaries. We propose that the RNAs produced at boundaries
recruit CTCF, resulting in robust intra-TAD transcription.
102
Results

Transcribed TAD boundaries insulate better than non-transcribed boundaries

To determine the frequency of transcribed TAD boundaries, we measured run-on transcription
with GRO-seq (37) on TAD boundaries identified by HOMER's findTADsAndLoops.pl. This
program identifies both contact and loop domain TADs based on intra-domain interaction
frequency. Both TADs and loops are thought to be formed by loop extrusion and to be dependent
on CTCF and cohesin (6, 8-10). HOMER identified 15,397 TAD boundaries at 5 kb resolution
using HiC data from HeLa cells (11). GRO-seq detected at least ten normalised counts within
about half of the TADs identified by HOMER (8134, 52.8%, Fig. 1A). Almost 80% of transcribed
boundaries are genic (Fig. 1B-C).

TAD boundaries are enriched for CTCF binding. We determined if strong CTCF binding at TAD
boundaries is associated with increased transcription. Using high-resolution CTCF CUT&RUN
data (38), we observed that both genic and non-genic transcribed TAD boundaries had
significantly higher CTCF enrichment than non-transcribed boundaries (Fig. 1D). This
relationship between transcription and CTCF enrichment at TAD boundaries was also observed
for different RNA levels (Fig. 1E). Within non-genic transcribed boundaries, CTCF enrichment

was also correlated with GRO-seq signal (Fig. S1A-B). Similar results were seen using CTCF
ChIP-seq (39) (Fig. S1C-E). We conclude that transcribed TAD boundaries are more enriched for
CTCF binding than non-transcribed boundaries.

We next asked if transcribed TAD boundaries that show high CTCF binding are stronger
insulators. We noted a positive correlation between the levels of RNA at the transcribed boundaries
and the TAD insulation (Fig. 1F). Even TADs with the least transcribed boundaries were better
insulated as compared to non-transcribed boundaries (Fig. 1F). We observed a similar positive
correlation between boundary RNA levels and TAD insulation using two different bin sizes (5 kb
and 20 kb) in our analyses, thereby ruling out resolution-related biases (Fig. S2). A total of 69.2%
(10658 out of 15397) of TADs boundaries were found to be transcribed (Fig. S2A). The CTCF
enrichment was stronger on transcribed genic and non-genic boundaries at both 5 kb and 20 kb
bin sizes (Fig. S2B-E). Insulation scores on transcribed genic and non-genic boundaries were
correlated with levels of RNA at the boundaries (Fig. S2F-H)

We also examined the relationship between transcription of TAD boundaries and CTCF binding
using HICCUPS, which identifies loop anchors. In total, 3525 loop anchors were identified with 5
kb bin size, of which 43% (1519) were genic and 19% (675) were non-genic (Fig. S3A). As was
the case for TAD boundaries identified by HOMER, we found a positive correlation between
transcription of loop anchors and CTCF enrichment using both CUT&RUN (Fig. S3B-C) and
ChIP-seq datasets (Fig. S3D-E). For subsequent analyses, we focused on TADs called by HOMER
with 5 kb boundary regions.

We also observed positive associations between CTCF enrichment, insulation score and
transcription at TAD boundaries in IMR90 fibroblasts (11) (Fig. S4A-D) and lymphoblastoid cells
K562 (40) (Fig. S4E-H). Together, our results suggest that CTCF enrichment and enhanced
insulation are general features of transcribed TAD boundaries.

RNA increases CTCF occupancy at TAD boundaries

We wondered whether elevated levels of transcription and CTCF at TAD boundaries are causally
related. To test this possibility, we treated HeLa cells with RNaseA and performed Western blot
analysis on the nuclear and chromatin fractions. We observed a significant loss of CTCF in the
chromatin fraction (Fig. 2A, S6A), consistent with the reported dependence of CTCF on RNA for
chromatin binding (29,31-32). To map the genomic sites of RNA-dependent CTCF binding, we
performed CUT&RUN for CTCF with RNaseA treatment (CUT&RUN.RNaseA). Upon RNA
depletion, we observed an overall decrease in CTCF occupancy, especially at the TAD boundaries
where CTCF is most enriched (Fig. 2B). Genome-wide, transcribed boundaries exhibited greater
loss of CTCF with RNaseA treatment as compared to non-transcribed and randomly selected
boundaries (Fig. 2C).

We next asked if the loss of CTCF at TAD boundaries is due to the digestion of RNA at direct
binding sites as opposed to a general effect of RNA degradation. We chose the ~500-kb

INK4a/ARF TAD, which spans the *CDKN2A*, *CDKN2B* and *MTAP* cell cycle regulatory genes and
a super-enhancer (Fig. 2D, highlighted region) (41-42). The super-enhancer within this TAD
contains several active enhancers that are enriched for H3K27ac modification and enhancer RNAs
(eRNAs) (Fig. 2D, mNET-seq panel) (41). We confirmed 5' TAD boundary locations on the
transcribed gene *MTAP* and 3' boundary on the non-genic region containing three CTCF ChIP-
seq peaks (B1, B2 and B3). The position of these boundary regions was persistent in HiC data
from other cell types (Data not shown). CTCF peaks B1 and B2 but not B3 exhibit strong nascent
transcription based on publicly available mNET-seq, GRO-seq and PRO-seq data (43-45) (Fig.
S5) and so are classified as eRNAs. In our CUT&RUN.RNaseA data, we observed a loss of CTCF
at the boundary regions of the *INK4a/ARF* TAD (Fig. S6B), which suggests that eRNAs help
maintain CTCF at B1 and B2 CTCF sites. To test this interpretation, we depleted sense and anti-
sense eRNAs at B1 and B2 CTCF sites using shRNAs (Fig. 2E-F) and confirmed the drop of these
eRNAs from chromatin fraction by qRT-PCR (Fig. S6C). Knockdowns using shRNAs to B1 and
B2 but not B3 and nearby TAD boundaries (T1 and T2) resulted in the loss of CTCF binding at
these sites (Fig. 2G-H and Fig. S6D). Notably, the loss of RNAs at B1 reduced CTCF occupancy
not only at B1 but also at B2 and vice versa (Fig. 2G-H). This suggests that CTCF on transcribed
boundaries is maintained in part by eRNAs, and different transcribed CTCF sites within a
boundary are potentially interdependent. Notably, shRNA-mediated knockdown or over-
expression of CTCF had at most minor effects on eRNA expression (Fig. 2I and Fig. S6E), which
suggests that CTCF enrichment at TAD boundaries depends on RNA, but not vice-versa.

Non-coding RNAs and CTCF interact at the *INKa/ARF* TAD boundary

We next asked whether RNA and CTCF interact at functional boundaries. We first performed
RNA pulldowns of sense and anti-sense RNAs transcribed from B1 and B2 CTCF sites in the
INKa/ARF TAD followed by immunoblotting with a CTCF antibody (Fig. 3A). We observed
CTCF interactions with B1 and B2 sense and anti-sense RNA and also with an eRNA from the
CTCF-bound E8 enhancer within the super-enhancer, but not with the Box B negative control.
CTCF binding to B1 and B2 sense and anti-sense RNAs was confirmed by UV-RIP, an RNA
immunoprecipitation method (Fig. 3B). Surprisingly, the CTCF motifs present at transcribed
boundaries are much sparser than at non-transcribed or randomly selected boundaries (Fig. 3C-D).
However, we found that transcribed boundaries are more densely packed with CTCF peaks than
are non-transcribed or randomly selected boundaries (Fig. 3E). These data suggest that
transcription or the presence of eRNA can reinforce CTCF binding to multiple weak CTCF motifs.

Next, we compared the stability, polyadenylation and splicing features of genic RNAs and eRNAs
at TAD boundaries with those that are not within boundaries. In general, eRNAs exhibit fast
turnover as seen by a higher ratio of nascent to mature eRNAs as compared to genic RNAs (37).
By this measure eRNAs on non-genic boundaries were significantly more stable than other
eRNAs, and RNAs on genic boundaries were slightly more stable than other genic RNAs (Fig.
3G).

The AATAAA polyadenylation motif is rare around Transcriptional Start sites (TSSs) of genic
transcripts, increasing in abundance downstream as expected (Fig. 3H). Non-genic eRNAs on
boundaries showed less enrichment of pre-mature AATAAA than other eRNAs (Fig. 3H),
consistent with their higher stability. As expected, 5' splice consensus sequences were strongly
enriched in mRNAs relative to eRNAs (Fig. 3I), consistent with their greater stability. Together,
enhanced stability, higher CTCF enrichment and insulation promoting activity of boundary eRNAs
suggest that they have evolved to facilitate CTCF binding by direct physical interaction.

Boundary RNAs enhance TAD transcription

The enhanced insulation we observed at sites of boundary eRNAs suggested that genes within the
TADs of transcribed boundaries might be expressed at higher levels than those in TADs of non-
transcribed boundaries. Accordingly, we compared the transcriptional levels of TADs with high,
low, and randomly selected transcribed or non-transcribed boundaries. We found that the
transcriptional output of entire TADs (Fig. 4A) or of individual genes within TADs (Fig. S7A) is
correlated with the levels of RNAs at boundaries, and we obtained similar results for non-genic
boundaries (Fig. S7B). To determine whether this positive association resulted from direct
interactions between promoters and boundaries we analysed available promoter capture-seq data
in HeLa cells (46). Boundaries were captured by promoters far more than by random regions (Fig.
4B), and genes that interacted with boundaries were relatively more active than genes that
interacted with other genomic regions (Fig. 4C). Together, these data suggest that transcribed
boundaries tend to interact with promoters and are associated with higher rates of TAD
transcription. We confirmed the interaction of the *CDKN2A* promoter with the nearby transcribed
boundary within the *INK4a/ARF* TAD using 4C-seq assays (Fig. 4D).

Next, we asked if boundary RNAs play a direct role in transcriptional activity of genes within this
TAD. Using shRNAs to sense and anti-sense B1 and B2 CTCF sites, we observed downregulation
of *INK4a/ARF* (transcribed from *CDKN2A*), *INK4b* (transcribed from *CDKN2B*) and *MTAP*,
whereas the transcription of *DDX5*, a non-*INK4a/ARF* TAD gene, remained unchanged (Fig. 4E).
These data suggest that boundary eRNAs positively regulate the transcription of the genes within
the *INK4a/ARF* TAD by direct physical interaction between promoters and boundaries.

Since the loss of specific boundary RNAs also caused loss of CTCF binding at the TAD boundary,
we asked if the role of boundary RNA in TAD transcription is CTCF-dependent. Depletion of
global CTCF using a pool of CTCF-specific siRNAs resulted in downregulation of genes within
and outside of the *INK4a/ARF* TAD as expected (Fig. 4F-G). To test the effects of the loss of
CTCF binding only at the *INK4a/ARF* TAD boundary, we deleted the B1 and B2 CTCF sites
individually in HeLa cells (Fig. S8A-B) and observed downregulation of *INK4a/ARF* genes within
the TAD (Fig. 4H) and the loss of PolII at promoters (Fig. 4I). We also observed that boundary
eRNA knockdown and boundary deletion reduced the levels of the p16 protein encoded by
INK4a/ARF (Fig. S8C). These data suggest that binding of CTCF mediated by boundary RNAs
regulates the expression of genes within the *INK4a/ARF* TAD.

Active enhancers within the TAD directly regulate boundary eRNA transcription

What is responsible for *INK4a/ARF* boundary transcription? Because HeLa cells are addicted to high levels of proteins encoded within the TAD (47-48), instead of deleting gene promoters, we used CRISPR inhibition (CRISPRi) using specific guide RNAs. We applied dCas9-KRAB on promoters of *CDKN2A*, *CDKN2B* and *MTAP* and assayed transcript levels at B1 and B2 CTCF sites (Fig. 5A). CRISPRi of these promoters resulted in transcriptional inhibition of these genes (Fig. 5B, red bars). However, boundary RNA levels were unaffected (Fig. 5B, blue bars). Consistent with unperturbed boundary RNA expression, there was no change in CTCF occupancy at B1 and B2 CTCF sites. However, we did observe loss of CTCF on the 5' boundary that overlaps the MTAP gene, likely due to down-regulation of *MTAP* (Fig. 5C). The boundary of flanking TADs also did not exhibit any perturbations in CTCF occupancy (Fig. 5C). These data suggest that initiation of transcription from *CDKN2A* and *CDKN2B* promoters is not responsible for eRNAs at the boundary.

We next asked whether the *INK4a/ARF* TAD enhancers regulate genes within the TAD and boundary eRNAs. Genome-wide, transcribed boundary TADs are enriched in H3K27ac (Fig. 6A-B) and PolII (Fig. 6C) relative to non-transcribed boundaries and randomly selected genic or non-genic boundaries (Fig. 6C, Fig. S9A-B). We first tested the super-enhancer using E8, E12 and E17 enhancer deletion lines (Fig. 6D) in which *INK4a/ARF* genes were seen to be downregulated based on RNA-seq data (Fig. 6E) (41), confirming that these enhancers are functional enhancers. We then asked if these enhancers indeed regulate the boundaries as well. Interestingly, we observed down-regulation of sense and anti-sense boundary RNAs at the B1 and B2 CTCF sites upon E8, E12 or E17 deletions (Fig. 6F), which suggests that enhancers activate boundary transcription. To determine whether down-regulation might be caused by direct contact, we also performed 4C-seq on the E8 enhancer. We observed strong interactions with other enhancers in the super-enhancer cluster, with *CDKN2A* and *CDKN2B* promoters and with the boundary region (Fig. 6G, right highlighted pink). Importantly, deletion of the E8 enhancer conspicuously reduced 4C-seq signal, confirming that the interaction between the super-enhancer and distant B1 and B2 CTCF sites regulates boundary transcription (Fig. 6G). Although interactions between boundaries and enhancers have been previously reported (35) no regulatory relationship had been established, whereas our data demonstrate that apart from activating their target promoters, functional enhancers positively regulate the transcription at the boundaries by direct interactions.

To ascertain whether high levels of boundary RNAs enhance insulation of the boundary *in vivo*, we interrogated the *INK4a/ARF* TAD upon senescence, where TAD transcription increases due to its cell cycle regulatory role. Upon senescence, bidirectional eRNA expression increases several-fold at the super-enhancer cluster (Fig. S10A, blue highlighted region), and *INK4a/ARF* gene expression and boundary RNA expression also increases (Fig. S10A, pink highlighted region). We validated these increases in gene transcription, boundary RNA expression and CTCF enrichment upon X-ray induced senescence (non-replicative senescence) (Fig. S10B-D), and also observed elevated p16 and p21 protein levels (Fig. S10E). Using HiC data, we observed increase in intra-

TAD interactions at this TAD in senescent cells as compared to proliferative cells (Fig. S10F).
Together these data suggest that increases in TAD enhancer activity accompany increases in their
interactions with boundaries to induce boundary RNA expression, which in turn recruits more
CTCF to enhance insulation.

Boundary eRNAs strengthen TAD insulation

We noticed that TADs with transcribed boundaries exhibited higher strength of enhancer/promoter
interactions than TADs with non-transcribed boundaries genome-wide (Fig. 7A). To test whether
boundary function of the *INK4a/ARF* TAD affects super-enhancer/promoter interactions in a
boundary RNA dependent manner, we performed 4C-seq assays at the E8 enhancer upon
knockdown of sense and anti-sense boundary RNAs at the B1 CTCF site. As expected, we
observed loss of boundary/enhancer interactions, but in addition, we observed weakening of
INK4a/ARF enhancer/promoter interactions (Fig. 7B). Knockdown of B1 RNAs also resulted in
increased interaction with the +T1 boundary of the neighboring TAD (Fig. 7B), which also showed
a gain in CTCF binding (Fig. 2G). To confirm that boundary RNA loss was accompanied by
reduced CTCF binding, we performed 4C-seq at the E8 enhancer using cells with a B1 CTCF site
deletion. We again observed a loss of enhancer/promoter interaction and a concomitant increase
in the interaction with the +1 T1 boundary of the neighboring TAD (Fig. 7B). These observations
demonstrate that loss of a boundary RNA reduces CTCF binding and weakens insulation, resulting
in aberrant enhancer interaction with an element outside of the TAD. We conclude that boundary
RNAs are functional components of TAD boundaries (Fig. 7C).

Discussion

Interactions between CTCF and RNA have been implicated in genome organization (31-32), but
the exact role of these interactions remains unknown. Here we show that eRNAs transcribed from
TAD boundaries function in CTCF binding, TAD insulation and gene regulation. Our data provide
direct evidence that boundary eRNAs create an insulated neighbourhood by associating with
CTCF-bound sites and facilitating enhancer:promoter interactions for robust gene activation
within a given TAD. We found that the transcriptional activity of promoters present within the
TAD does not regulate non-genic boundary transcription. Using a series of enhancer and CTCF
site deletions at the *INK4a/ARF* TAD, we have shown that enhancers interact with and positively
regulate the expression of both genic and non-genic TAD boundary RNAs. Boundary RNAs, in
turn, increase CTCF binding at TAD boundaries, strengthening weak CTCF motifs and increasing
insulation. Transcription of boundary RNAs at the *INK4a/ARF* TAD is regulated by the super-
enhancer within the TAD to facilitate CTCF binding and increased insulation at the boundary. This
may explain why super-enhancers containing TADs exhibit stronger boundaries (34-36).

Boundary RNAs resemble eRNAs at active enhancers in being bidirectional (49-50), and as our
study demonstrates, making direct contact with distant regulatory elements. However, unlike
eRNAs, boundary RNAs contact enhancers that also contact promoters to enhance TAD gene

activation. Popular chromosome conformation capture assays are strictly pairwise and cannot determine if enhancer elements activate promoters and boundary elements independently or together. However, more recently introduced 3D contact methods, including SPRITE (51) and GAM (52) are able to capture multiple simultaneous contacts, and in the future, may be applied to understand how enhancers regulate both gene promoter activity and transcription at boundaries to strengthen TADs.

Furthermore, in the absence of the RNA binding domain of CTCF, cells exhibit a decrease in the total number of TADs and an increase in the size of individual TADs due to reduced insulation between neighboring TADs (31). A clear mechanistic basis for this observation comes from our findings in which TAD enhancer-mediated CTCF-RNA interactions maintain strong TAD boundaries by increasing insulation between neighboring TADs. Thus, the inability of CTCF mutants to interact with RNA will compromise CTCF enrichment and its insulating function at TAD boundaries, leading to the merging of adjacent TADs into larger TADs. In such cases, less CTCF at the boundary might allow the cohesin ring to slip to the next TAD thereby compromising the insulation (53-54). Thus a positive feedback loop is established whereby super-enhancers maintain high specificity for promoters within the TAD by also activating boundary transcription and preventing interactions with neighboring promoters.

TAD boundaries are enriched for H3K27ac and PolII, suggesting the presence of active regulatory elements and transcription (24,34-36). However, activation of transcription at the boundary alone is not sufficient to maintain a TAD boundary (23,55). Our study shows that the products of the TAD boundary transcription *i.e.*, boundary RNAs, are required to strengthen TAD insulation by increasing CTCF occupancy. Transcription-assisted TAD boundary strengthening may also be a factor in zygotic activation-dependent boundary formation where early transcribing genes serve as the nucleation sites for TAD boundaries (55). Similarly, transcription and the resulting RNA may be involved in forming distinct TADs boundaries around transcribed escape genes on the inactive chromosome during X chromosome inactivation (26).

Materials and Methods

Cell Culture:

HeLa cells were obtained from ATCC and were cultured in (DMEM) Dulbecco's Modified Eagle's medium supplemented with 10% Fetal Bovine Serum and 1% Penicillin/Streptomycin. The cells were maintained at 37°C in the presence of 5% CO_2 in a humidified incubator.

Chromatin Immunoprecipitation:

Cells were crosslinked with 1% formaldehyde at room temperature for 10 min with constant shaking. Glycine was added to a final concentration of 125 mM to quench the formaldehyde. Cells were washed three times with 1X ice-cold PBS then were scraped and pelleted down 1X PBS at 4 °C. Cells were gently resuspended in nuclear lysis buffer (50 mM Tris-HCl pH 7.4, 1% SDS, 10 mM EDTA pH 8.0) supplemented with 1X PIC and incubated on ice for 10 min. The lysate was

subjected to fragmentation using a Diagenode Bioruptor Pico for 20 cycles (30 s ON and 30 s OFF)
to generate fragments of around ~500 bp. The cell lysate was cleared at 12K rpm for 12 min. For
each IP, 100 μg of sheared chromatin was used. The lysates were diluted by adding Dilution Buffer
(DB) (20 mM Tris-HCl pH 7.4, 100 mM NaCl, 2 mM EDTA pH 8.0, 0.5% Triton X-100)
supplemented with 1X PIC in 1:1.5 ratio (1 volume of sheared chromatin and 1.5 volumes of
Dilution Buffer). One μg of antibody was added to immunoprecipitate the DNA and incubated on
a rocking platform overnight at 4°C. Fifteen μl of Pre-blocked (with 1% BSA) Protein G
Dynabeads (10004D, Invitrogen) was added to the tubes and rotated for 4h at 4 °C. Beads were
collected, flow-through was discarded and the samples were washed with Wash Buffer I (20 mM
Tris-HCl pH 7.4, 150 mM NaCl, 0.1% SDS, 2 mM EDTA pH 8.0, 1% Triton X-100) at 4 °C on a
rocking platform. Washes were sequentially repeated with Wash Buffer II (20 mM Tris-HCl pH
7.4, 500 mM NaCl, 2 mM EDTA pH 8.0, 1% Triton X-100), Wash Buffer III (10 mM Tris-HCl
pH 7.4, 250 mM LiCl, 1% NP-40, 1% Sodium Deoxycholate, 1 mM EDTA pH 8.0 and 1X TE (10
mM Tris pH 8.0, 1mM EDTA pH 8.0). Chromatin was eluted with 200 ul of Elution Buffer (100
mM NaHCO3, 1% SDS) for 30 min at 37°C in a thermomixer. The supernatant was collected in
separate tubes and 14 μl NaCl (5 M) was added to eluted samples and kept at 65 °C overnight for
de-crosslinking. Immunoprecipitated DNA was purified by phenol:chloroform:isoamyl alcohol,
followed by ethanol precipitation. The final air-dried DNA pellet was dissolved in 100 μl of 1X
TE. These samples were then used for qRT-PCRs using SYBR Green. Mean values for all the
regions analyzed in different conditions were expressed as fold-enrichment compared over beads.
Statistically significant differences between different conditions were computed using a T-test (p-
value < 0.05), ChIP primers are listed in Supplementary Table. 1.

RNA Isolation and cDNA synthesis:

Cells were lysed in 1 ml of Trizol (Thermo Fisher Inc.). To each sample, 200 μl chloroform was
added, briefly vortexed and centrifuged at 12K rpm for 12 min. The aqueous phase was carefully
collected and transferred to the fresh tube. One volume of isopropanol was added to the sample
and incubated at room temperature for 10 min to precipitate the RNA. The samples were
centrifuged at 12K rpm for 12 min, and supernatants were discarded without disturbing the pellet.
The pellet was washed with 75% ethanol. The pellet was air-dried and dissolved in RNase-free
water. The RNA was treated with ezDNase (Invitrogen) to remove the traces of contaminating
DNA. One μg of RNA was used for each cDNA synthesis reaction by Superscript IV (Invitrogen)
and random hexamers as per the manufacturer's recommendation. The CFX96 Touch (Biorad)
real-time PCR instrument was used for qRT-PCRs. qRT-PCRs were performed using three
biological replicates for each sample. Fold change was calculated by the $\Delta\Delta$Ct method and
individual expression data were normalized to GAPDH mRNA. The p-values were calculated by
Student's unpaired two-tailed t-test. qRT-PCR primers are listed in Supplementary Table 1.

CRISPR Cas9 mediated deletion:

Guide RNAs (gRNAs) were designed with the crispr.mit.edu tool. gRNAs were selected based on
the highest score and the least number of off-targets. gRNAs were cloned in a pgRNA humanized

vector (Addgene #44248, a gift from the Stanley Qi Lab) between BstX1 and Xho1 restriction
sites. gRNAs were co-transduced with a lenti-Cas9 vector. Cells were placed under selection using
puromycin (3µg/ml) for 48 hours. Single cells were seeded in a 96 well plate. Wells with single
cells were marked and allowed to grow until they formed colonies. The cells were trypsinized and
shifted to 48 well plates and allowed to grow to confluence, and then half of the cells were taken
for the Surveyor assay and the other half was plated again. Surveyor assays were performed using
a PCR-based method using primers listed in the Supplementary Table 1.

CRISPRi gRNA design and cloning:

We used GPP sgRNA Designer (https://portals.broadinstitute.org/gpp/public/analysis-tools/sgrna-
design) to design the gRNAs against TSSs of *MTAP*, *CDKN2A* and *CDKN2B* in the *INK4a/ARF*
TAD. For each gene, two gRNAs were designed. The gRNAs were cloned in a customized PX459
vector (pSpCas9(BB)-2A-Puro V2.0, Addgene #62988, a gift from the Feng Zhang lab). The Cas9
enzyme cassette was removed from PX459 by digestion with Xba1 and Not1 followed by blunt-
end ligation. The cloning of the gRNAs in the customized PX459 plasmid was performed as per
the Zhang Lab's general cloning protocol. The dCas9 plasmid, together with gRNA specific to
targeted genes, were transfected in 70% confluent cells. Transfections were done using
Lipofectamine 2000 (Invitrogen, cat no #11668027). CRISPRi cells were harvested for ChIP and
gene expression analysis.

shRNA design and transfection:

We used an shRNA design tool (https://portals.broadinstitute.org/gpp/public/seq/search) to design
the shRNAs against RNA at B1 and B2 CTCF sites within the *INK4a/ARF* TAD boundary. The
shRNAs were cloned in a customized pLKO.1 puro, (Addgene #8453, a gift from Bob Weinberg.
The cloning of the shRNAs in the pLKO.1 plasmid was performed as per the Addgene's pLKO.1
cloning protocol. shRNA-specific plasmid together with lentiviral packaging plasmids like VSVG
(a gift from Bob Weinberg, Addgene #8454), and PAX2, a gift from Didier Trono (addgene
#12260) were co-transfected using Lipofectamine 2000 Invitrogen (cat #11668027). Cells were
selected by puromycin (3µg/ml) (GIBCO cat # A11138-03) and knockdowns were confirmed
using qPCR oligos listed in Supplementary Table 1.

siRNA transfection:

SMARTpools was used to design scrambled (cat no D-001810-10-05) and CTCF siRNAs (L-
020165-00-0005) were purchased from GE Dharmacon. Transfections were done using
Lipofectamine 2000 (Invitrogen cat no 11668027).

Lentiviral transduction:

HEK293FT cells were grown in poly-D lysine coated culture dishes. These cells were co-
transfected with lentiviral packaging plasmids like VSVG (a gift from Bob Weinberg, Addgene
#8454) and PAX2 (a gift from Didier Trono, Addgene #12260) along with the plasmid of interest

using Lipofectamine 2000 Invitrogen (cat# 11668027). The medium was changed after 6 hours.
The viral soup was collected after 48 h and 72 hours, pooled together, filtered with a 0.44 μm
syringe filter and then finally added to cells along with 8 μg/ml of polybrene. Transduction was
stopped after 16 hours.

RNA Immunoprecipitation:

For RNA, primers were designed to amplify desired genomic regions that correspond to peaks of
boundary RNA (GRO-seq) at B1 and B2 CTCF sites of the *INK4a/ARF* TAD boundary. The
amplified products were cloned into the pcDNA3 BoxB plasmid (a gift from Howard Chang,
Addgene #29729). All clones were confirmed by Sanger sequencing and subsequently used for
RNA synthesis using T7 RNA polymerase (Promega Inc. P207e) and biotin RNA labeling mix
(Roche 11685597910). Primer sequences are listed in Supplementary Table 1.

RNA immunoprecipitation was adopted from (56), with some modifications. HeLa cells were
grown to 90–95% confluency and were harvested by scraping and washed twice with ice-cold 1X
PBS to gently resuspend in 2 ml of PBS. A volume of 2 ml of nuclear isolation buffer [NIB: 40
mM Tris–HCl pH 7.5, 20 mM $MgCl_2$, 1.28 M sucrose, 4 % Triton X-100, 1 mM PMSF, protease
inhibitors, and 20 U/ml SUPERase inhibitor (Thermo Fisher Scientific, catalog # AM2694)] was
added and the pellet was gently resuspended. Then, 6 ml of distilled water was added and kept on
ice for 20 min with intermittent gentle shaking. Nuclei were pelleted by centrifugation at $2500 \times g$
for 5 min at 4°C. The pellet was resuspended in 1 ml of RIP buffer (25 mM Tri–HCl pH 7.4, 150
mM KCl, 0.5 mM DTT, 0.5 % NP40, 1 mM PMSF, protease inhibitors, and 20 U/ml SUPERase
inhibitor) and incubated on ice for 5 min. Nuclei were sheared by 10 cycles of sonication (30s on
and 30 s off) in a Bioruptor (Diagenode) followed by centrifugation at 12K rpm for 10 min at 4°C.
One μg of biotinylated RNA was incubated with 20 μl of RNA structure buffer (RSB) (10 mM
Tris–HCl pH 7.0, 100 mM KCl, 10 mM $MgCl_2$, 1 mM PMSF, protease inhibitors, and 20 U/ml
SUPERase inhibitor) at room temperature for 5 min. Folded RNA was mixed with 1 mg of nuclear
extract in 500 μl of RIP buffer and rotated at 4°C for 1h. Fifteen μl of Dynabeads MyOne
Streptavidin beads was added and rotation continued for one more hour. Samples were washed
three times with RIP buffer and beads with proteins were boiled in 2XSDS for 10 min.
Immunoblotting for CTCF was performed using a CTCF antibody (Cell Signaling Technology,
#3418).

UV RIP:

Ultraviolet-RNA Immunoprecipitation was performed as in (57) with some modifications. HeLa
cells at a confluency of 80–90% were cross-linked by UV irradiation in Stratalinker UV cross-
linker. Cells were washed three times with cold 1XPBS and scraped in PBS before subjecting to
centrifugation at 2500 rpm. Pellets were lysed in RIP lysis buffer [25 mM HEPES-KOH at pH 7.5,
150 mM KCl, 0.5% NP40, 1.5 mM $MgCl_2$, 10% glycerol, 1 mM EDTA, 0.4 U RNase inhibitor
(Invitrogen, 18091050)], and 1X protease inhibitor cocktail (PIC) on ice for 30 min. Cleared cell
lysates were subjected to IP with CTCF antibody-bound Protein G Dynabeads (Invitrogen,
10004D) overnight. Samples were subsequently washed three times with RIP lysis buffer and RNA

samples were eluted using TRIzol reagent. Prior to cDNA synthesis, RNA was treated with RNase-
Free DNase Qiagen (cat no.79256) to remove the traces of contaminating DNA. cDNA was
prepared using random hexamers by Superscript IV (Invitrogen 18091050) as per the
manufacturer's recommendation and analyzed by PCR primers listed in Supplementary Table 1
and were run on 1% Agarose gel.

Cell Fractionation:

Cell fractionation and RNAseA treatment of nuclei was adapted from (58). Briefly, cells were
washed three times with cold 1XPBS and scraped in 1ml of PBS, pelleted down at 2500 rpm for 5
min at 4^0C. Pellets were then resuspended in Hypotonic Lysis Buffer (HLB) (10 mM Tris HCl pH
7.5, 10 mM NaCl, 3 mM $MgCl_2$, 0.3% NP-40, 10% glycerol, 1XPIC), the mix was incubated on
ice for 10 min and then centrifuged at 800g for 8 min at 4^0C. The supernatant was transferred to a
new tube and marked as Cytoplasmic fraction. The nuclei pellet after centrifugation was washed
twice with HLB and centrifuged at 200g for 2 min at 4^0C. The pellet was then resuspended in 700
ul of ice cold Modified Wuarin-Schiebler buffer (MWS: 10 mM Tris-HCl, pH 7.5, 300 mM NaCl,
4 mM EDTA, 1 M urea, 1% NP-40, 1% glycerol, 1XPIC) with or without 0.1l of 100mg/ml
RNaseA (Qiagen cat. no. 19101) for 15 min on ice briefly tapped few times during incubation time
and then centrifuged at 1000g for 5 min at 4^0C. The supernatant was collected as a nucleoplasmic
fraction. The chromatin pellet was washed twice with MWS buffer and centrifuged at 500g for
3min at 4^0C. The pellet was then dissolved in 500 µl of Nuclear Lysis Buffer (NLB: 20 mM Tris
HCl pH 7.5, 150 mM KCl, 3 mM $MgCl_2$, 0.3% NP-40, 10% glycerol, 1XPIC) and sonicated for
10 cycles (30 s on, 30 s off) in a Bioruptor (Diagenode). SDS loading dye was added and the
samples were boiled before immunoblotting.

Chromatin-associated RNA isolation:

Chromatin-associated RNA isolation was adapted from (59). Trypsinized cells were collected in
1.5 ml tubes and washed twice with 1xPBS and pellet down at 200xg for 2 min. To each pellet
was added 400 µl of Cell lysis buffer (10 mM Tris HCl ph7.4, 150 mM NaCl, 0.15% Igepal),
gently pipetted 3-5 times and then incubated for 5 min on ice. To the Lo-bind 1.5ml tube was
added 1 ml (2.5 volume) of cold sucrose buffer (10mM Tris HCl ph7.4, 150mM NaCl, 24%
Sucrose) and gently overlaid with the cell lysate followed by centrifugation at 3500xg for 5 min
to collect the nuclear pellet. Isolated nuclei were rinsed with 1 ml of ice-cold 1xPBS-EDTA
solution followed by a short spin at 3500xg. The nuclei were resuspended in 250 µl of glycerol
buffer (20 mM Tris HCl ph7.4, 75 mM NaCl, 0.5 mM EDTA pH 8.0, 50% Glycerol) and then
immediately 250 µl of Urea buffer (10 mM Tris HCl ph7.4, 1 M Urea, 0.3 M NaCl, 7.5 mM $MgCl_2$,
0.2 mM EDTA, 1% Igepal) was added, mixed by vortexing for 4 s and then incubated on ice for 2
min. Chromatin was centrifuged at 13000xg for 2 min to collect the Chromatin-RNA complex.
The chromatin-RNA pellet was washed with 1xPBS-EDTA once and RNA was isolated using
Trizol Reagent. RNA obtained was treated with ezDNase (Ambion) to remove the traces of
contaminating DNA. One µg of RNA was used for each cDNA synthesis reaction by Superscript
IV (Invitrogen) and random hexamers as per the manufacturer's recommendation. The CFX96

Touch (Biorad) real-time PCR was used for qRT-PCRs. qRT-PCRs were performed using three biological replicates for each sample. Fold changes were calculated by the ΔΔCt method and individual expression data was normalized to pre-c-myc. The *p*-values were calculated by Student's unpaired two-tailed *t*-test. qRT-PCR primers are listed in Supplementary Table 1.

X-ray induced Senescence:

BJ fibroblasts were cultured in (DMEM) Dulbecco's Modified Eagle's medium supplemented with 10% Fetal Bovine Serum and 1% Penicillin/Streptomycin at 37 °C in a humidified incubator containing 5% CO_2. Cells were first immortalized by expressing hTERT. Cells were then seeded at around 40% confluency. The day after seeding, a single dose of 10 Gy was delivered using a CIX3 cabinet x-ray irradiator (xstrahl Life Sciences) (195 kV, 10 mA). Irradiated cells were not passaged, but the medium was changed on Days 4, 8,12 and 16 and cells were either harvested on day 10 or day 20 and used for RNA isolation and Chip experiments respectively. Senescence was confirmed by SA-β-gal staining (Cell Signaling technology #9860) and p21 expression (p21 Santacruz Biotechnology sc-397).

Antibodies:

Antibodies used were: CTCF (Cell Signaling Technologies, #3418 for immunoblotting and Millipore Sigma #07-729 for CUT&RUN.RNase), DDX5 (Abcam, ab-21696), Cdkn2A, p16 (Abcam ab-108349), ARF, p14 (Santacruz Biotechnology, sc-53639), PolII (Santacruz Biotechnology, sc-55492), GAPDH (Santacruz Biotechnology, sc-32233), Histone H3 (Sigma-Aldrich H0164).

ChIP-seq and CUT&RUN analysis:

Sequenced reads were aligned to the hg19 assembly using default Bowtie2 options. Tag directories were made from the aligned reads using the HOMER (http://homer.ucsd.edu/homer/) makeTagDirectory command. A 200 bp sliding window was used to identify narrow peaks of transcription factors' ChIP-seq and CTCF CUT&RUN. CUT&RUN.RNAse was performed as described in (26). Artifacts from clonal amplification were neglected as only one tag from each unique genomic position was considered (-tbp 1). The threshold was set at a false discovery rate (FDR) of 0.001 determined by peak-finding using randomized tag positions in a genome with an effective size of 2×10^9 bp. For histone marks, seed regions were initially found using a peak size of 500 bp (FDR<0.001) to identify enriched loci. Enriched regions separated by <1kb were merged and considered as blocks of variable lengths. The HOMER makeUCSCfile command was used to generate bedgraph files of the read densities across the genome and this track was uploaded to the UCSC genome browser. The tag directories generated were used to quantify signals at various regions of interest using annotatePeaks.pl. The datasets used in the study are listed in Supplementary Table 3.

GRO-seq analysis:

The sequenced reads were aligned to the hg19 assembly using default Bowtie2 options. Tag
directories were made from the aligned reads using the HOMER makeTagDirectory command.
Tag directories were used to quantify signal and gene expression. The HOMER makeUCSCfile
command was used to generate strand-specific bedgraph files of the read densities across the
genome and this track was uploaded to the UCSC genome browser.

TADs and loop domain calling:

Raw paired-end reads were trimmed and individually mapped to the hg19 assembly using bowtie2.
The HOMER makeTagDirectory was first used to create tag directories with tbp 1 option. Datasets
were further processed by HOMER in order to remove small fragment and self-ligations using
makeTagDirectory with the following options: -removePEbg -removeSpikes 10000 5. Next,
findTADsAndLoops.pl was used to obtain overlapping TADs (TADs and sub-TADs), produced
at 5 kb resolution with 10 kb windows. This program was also used to generate two bedGraph files
that describe the directionality index and the insulation score. tagDir2hicFile.pl was used to
generate .hic files. The WashU epigenome browser was used to visualize the HiC data. Boundaries
of the TADs were defined as 5 kb or 20 kb regions centered at the end of the TADs called.
HICCUPS was used to obtain loop anchors from the .hic file at a 5kb resolution. HOMER
annotatePeaks.pl was used to quantify signal from tagDirectories, count peaks, quantify insulation
and annotate features at the TAD boundaries. Non-genic TAD boundaries were identified as those
that did NOT intersect with an annotated gene body. GRO-seq signal was used to rank the
boundaries and to then select 10% of unique boundaries in each category (high transcription, low
transcription, no transcription and random). The .hic file was converted to .cool. The file was then
used to plot the averaged insulation score valleys at 5 kb resolution and 100 kb around the
boundary using coolpup.py.

NET-CAGE and CAGE Analysis:

NET-CAGE and CAGE sequenced reads were aligned to hg19 using bowtie2. The HOMER
program makeTagDirectory was first used to create tag directories. findPeaks with "–style tss"
was used to obtain the NET-CAGE TSSs. TSSs identified within 1 kb of each other on the same
strand were merged. NET-CAGE identified TSSs intersecting the refseq annotated TSSs were
called mRNA TSSs and the rest were defined as eRNA TSSs. Those TSSs which fell in boundaries
were identified as bmRNAs or beRNAs. A span of 2kb around each identified TSSs was used to
quantify NET-CAGE and CAGE signal and to calculate the $\log_2$ fold difference between the two.

5' Splice site and Transcription Termination Site Sequence Analysis:

To identify the 5' SS motif, HOMER denovo motif analysis was performed on the 500bp region
after the Refseq annotated TSSs with other H3K27ac 500 bp regions as background. The top hit
was observed to be the splice acceptor site. The "AATAAA" motif was used as the 3' polyA
sequence. HOMER scanMotifGenomeWide was used to identify "predicted" splice sites and
polyA terminator sites using previously identified motifs. The occurrence of predicted sites was
plotted at the previously identified TSSs.

Circular Chromatin Conformation Capture-seq:

4C was performed as per the protocol described in van de Werken et al., 2012 with minor modifications. HeLa cells were fixed with fresh formaldehyde (1.5%) and quenched with glycine (125mM) followed by washes with ice-cold PBS (2x) and scraped, pelleted and stored at -80^{0}C. Lysis buffer [Tris-Cl pH 8.0 (10mM), NaCl (10mM), NP-40 (0.2%), PIC (1x)] was added to the pellets and homogenized with a Dounce homogenizer (20 stroked with pestle A followed by pestle B). The 3C digestion was performed with HindIII (400units, NEB) and ligation was performed using the T4 DNA ligase and ligation mix (1% Triton X-100, 0.105 mg/ml BSA, 1.05 mM ATP, 1x Ligation buffer, where 10x Ligation buffer is 500 mM Tris-Cl pH 7.5, 100 mM MgCl$_2$, 100 mM DTT). The ligated samples were then purified by PCI (Phenol/Chloroform/Isoamyl alcohol) extraction, subjected to ethanol precipitation, and the pellet was eluted in TE (pH 8.0) to obtain the 3C library. The 4C digestion was performed using 50 units DpnII (NEB) and the samples were ligated, purified and precipitated similar to the 3C library to obtain the 4C library. The 4C library was subjected to RNaseA treatment and purified by the QIAquick PCR purification kit. The concentration of the library was then measured by Nanodrop and subjected to PCR using the oligos for the enhancer viewpoint. The samples were next PCR purified using the same kit and subjected to Illumina HiSeq2500 sequencing using 50 bp single-end reads.

4C method and analysis:

The sequenced reads were aligned to the hg19 assembly as described in R.4Cker (https://github.com/rr1859/R.4Cker) with Bowtie2. Data analysis was further performed using 4C-ker with default parameters. 4Cseqpipe (http://compgenomics.weizmann.ac.il/tanay/?page_id=367) was also used to process the sequenced data. 4C-seq images were generated using truncated mean at a 10kb resolution.

Motif analysis:

annotatePeaks.pl was used with –hist option with the CTCF motif probability matrix to plot the motif occurrence in the peaks.

Promoter capture analysis:

Processed CHiCAGO data was obtained from Thiecke et al. (https://osf.io/brzuc/). The BEDTools suite was used to obtain interactions of interest and perform further analysis. The promoter interacting region of the Bedpe files of pcHiC interactions were then intersected with boundaries to obtain promoters that interact with boundaries. The strength of the interactions obtained from the processed CHiCAGO data was plotted for interactions from boundary interacting promoters to enhancers (H3K27ac regions).

RNA-seq analysis:

The raw reads were mapped to the hg19 assembly using hisat2 in a strand-specific manner. Tag directories were made from the aligned reads using the HOMER makeTagDirectory command.

616   The HOMER makeUCSCfile command was used to generate strand-specific bedgraph files of the
617   read densities across the genome and this track was uploaded to the UCSC genome browser.
618

619   **Acknowledgments:**
620
621   We are grateful to Henrietta Lacks and to her family members for their contributions to
622   biomedical research. We thank DN lab members, Ranveer Jayani, Amanjot Singh for the
623   discussions.
624
625   **Funding:**
626
The Department of Atomic Energy, Government of India, under project no.12-R&D-TFR-5.04-
0800. Intramural funds from NCBS-TIFR (to DN)
DN is a EMBO Global Investigator.
Welcome-IA: IA/1/14/2/501539 (to DN).
UF is supported by CSIR (Govt. of India) senior research fellowship.
KW is supported by the TIFR-NCBS graduate program and CSIR-SPM (Govt. of India)
fellowship.

634
635   **Author contributions:**
636
Conceptualization and design: ZI, BS and DN
NGS analysis: BS
Methodology: ZI, BS, UF and KW
CTCF CUT&RUN: JT and SH
NGS library preparation and sequencing: AP
Writing—original draft: DN, SH, ZI and BS
Reviewing and editing: DN, ZI, BS, KW, UF, JT, AKS, AP, RS
644
645   **Competing interests:** All other authors declare they have no competing interests.
646
647   **Data and materials availability:** The accession codes for data used in the study are mentioned
648   in Table. 3. The CUT&RUN raw and processed data for control and RNaseA- treated HeLa cells
649   have been deposited in the GEO repository with accession code GSE180659. The data can be
650   reviewed using token ehwnowswbbcdxyl into the box.
651   Goto https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE180659
652

References

1. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B. 2012. Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature*. doi:10.1038/nature11082

2. Beagan JA, Phillips-Cremin JE. 2020. On the existence and functionality of topologically associating domains. *Nat Genet.* doi: 10.1038/s41588-019-0561-1.

3. Dekker J, Mirny L. 2016. The 3D Genome as Moderator of Chromosomal Communication. *Cell*. doi:10.1016/j.cell.2016.02.007

4. Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA. 2016. Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep*. doi:10.1016/j.celrep.2016.04.085

5. Nuebler J, Fudenberg G, Imakaev M, Abdennur N, Mirny LA. 2018. Chromatin organization by an interplay of loop extrusion and compartmental segregation. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1717730115

6. Rao SSP, Huang SC, Glenn St Hilaire B, et al. 2017. Cohesin loss eliminates all loop domains. *Cell*.;171(2):305–320.e24.

7. Szabo Q, Donjon A, Jerković I, Papadopoulos GL, Cheutin T, Bonev B, Nora EP, Bruneau BG, Bantignies F, Cavalli G. 2020. Regulation of single-cell genome organization into TADs and chromatin nanodomains. *Nat Genet*. doi:10.1038/s41588-020-00716-8

8. Nora EP, Goloborodko A, Valton AL, Gibcus JH, Uebersohn A, Abdennur N, Dekker J, Mirny LA, Bruneau BG. 2017. Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. *Cell*. doi:10.1016/j.cell.2017.05.004

9. Schwarzer W, Abdennur N, Goloborodko A, et al. 2017. Two independent modes of chromatin organization revealed by cohesin removal. *Nature*. 2017;551(7678):51–56

10. Wutz G, Várnai C, Nagasaka K, et al. 2017. Topologically associating domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. *Embo J*. 2017;36(24):3573–3599

11. Rao SSP, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES, Aiden EL. 2014. A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell*. doi:10.1016/j.cell.2014.11.021

12. Krivega I, Dean A. 2017. CTCF fences make good neighbours. *Nat Cell Biol*. doi:10.1038/ncb3584

13. Rowley MJ, Nichols MH, Lyu X, Ando-Kuri M, Rivera ISM, Hermetz K, Wang P, Ruan Y, Corces VG. 2017. Evolutionarily Conserved Principles Predict 3D Chromatin Organization. *Mol Cell*. doi:10.1016/j.molcel.2017.07.022

14. Despang A, Schöpflin R, Franke M, Ali S, Jerković I, Paliou C, Chan WL, Timmermann B, Wittler L, Vingron M, Mundlos S, Ibrahim DM. 2019. Functional dissection of the Sox9–Kcnj2 locus identifies nonessential and instructive roles of TAD architecture. *Nat Genet*. doi:10.1038/s41588-019-0466-z

15. Flavahan WA, Drier Y, Liau BB, Gillespie SM, Venteicher AS, Stemmer-Rachamimov AO, Suvà ML, Bernstein BE. 2016. Insulator dysfunction and oncogene activation in IDH mutant gliomas. *Nature*. doi:10.1038/nature16490

16. Hanssen LLP, Kassouf MT, Oudelaar AM, Biggs D, Preece C, Downes DJ, Gosden M, Sharpe JA, Sloane-Stanley JA, Hughes JR, Davies B, Higgs DR. 2017. Tissue-specific CTCF-cohesin-mediated chromatin architecture delimits enhancer interactions and function in vivo. *Nat Cell Biol*. doi:10.1038/ncb3573

17. Kloetgen A, Thandapani P, Ntziachristos P, Ghebrechristos Y, Nomikou S, Lazaris C, Chen X, Hu H, Bakogianni S, Wang J, Fu Y, Boccalatte F, Zhong H, Paietta E, Trimarchi T, Zhu Y, Van Vlierberghe P, Inghirami GG, Lionnet T, Aifantis I, Tsirigos A. 2020. Three-dimensional chromatin landscapes in T cell acute lymphoblastic leukemia. *Nat Genet*. doi:10.1038/s41588-020-0602-9

18. Narendra V, Rocha PP, An D, Raviram R, Skok JA, Mazzoni EO, Reinberg D. 2015. CTCF establishes discrete functional chromatin domains at the Hox clusters during differentiation. *Science (80-)*. doi:10.1126/science.1262088

19. Sima J, Chakraborty A, Dileep V, Michalski M, Klein KN, Holcomb NP, Turner JL, Paulsen MT, Rivera-Mulia JC, Trevilla-Garcia C, Bartlett DA, Zhao PA, Washburn BK, Nora EP, Kraft K, Mundlos S, Bruneau BG, Ljungman M, Fraser P, Ay F, Gilbert DM. 2019. Identifying cis Elements for Spatiotemporal Control of Mammalian DNA Replication. *Cell*. doi:10.1016/j.cell.2018.11.036

20. Williamson I, Kane L, Devenney PS, Flyamer IM, Anderson E, Kilanowski F, Hill RE, Bickmore WA, Lettice LA. 2019. Developmentally regulated Shh expression is robust to TAD perturbations. *Dev*. doi:10.1242/dev.179523

21. Yang M, Vesterlund M, Siavelis I, Moura-Castro LH, Castor A, Fioretos T, Jafari R, Lilljebjörn H, Odom DT, Olsson L, Ravi N, Woodward EL, Harewood L, Lehtiö J, Paulsson K. 2019. Proteogenomics and Hi-C reveal transcriptional dysregulation in high hyperdiploid childhood acute lymphoblastic leukemia. *Nat Commun*. doi:10.1038/s41467-019-09469-3

22. Khoury A, Achinger-Kawecka J, Bert SA, Smith GC, French HJ, Luu PL, Peters TJ, Du Q, Parry AJ, Valdes-Mora F, Taberlay PC, Stirzaker C, Statham AL, Clark SJ. 2020. Constitutively bound CTCF sites maintain 3D chromatin architecture and long-range epigenetically regulated domains. *Nat Commun*. doi:10.1038/s41467-019-13753-7

23. Bonev B, Mendelson Cohen N, Szabo Q, Fritsch L, Papadopoulos GL, Lubling Y, Xu X, Lv X, Hugnot JP, Tanay A, Cavalli G. 2017. Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell*. doi:10.1016/j.cell.2017.09.043

24. Hsieh THS, Cattoglio C, Slobodyanyuk E, Hansen AS, Rando OJ, Tjian R, Darzacq X. 2020. Resolving the 3D Landscape of Transcription-Linked Mammalian Chromatin Folding. *Mol Cell*. doi:10.1016/j.molcel.2020.03.002

25. Zhang D, Huang P, Sharma M, Keller CA, Giardine B, Zhang H, Gilgenast TG, Phillips-Cremin JE, Hardison RC, Blobel GA. 2020. Alteration of genome folding via contact domain boundary insertion. *Nat Genet*. doi:10.1038/s41588-020-0680-8

26. Giorgetti L, Lajoie BR, Carter AC, Attia M, Zhan Y, Xu J, Chen CJ, Kaplan N, Chang HY, Heard E, Dekker J. 2016. Structural organization of the inactive X chromosome in the mouse. *Nature*. doi:10.1038/nature18589

27. Zhang Y, Li T, Preissl S, Amaral ML, Grinstein JD, Farah EN, Destici E, Qiu Y, Hu R, Lee AY, Chee S, Ma K, Ye Z, Zhu Q, Huang H, Fang R, Yu L, Izpisua Belmonte JC, Wu J, Evans SM, Chi NC, Ren B. 2019. Transcriptionally active HERV-H retrotransposons demarcate topologically associating domains in human pluripotent stem cells. *Nat Genet*. doi:10.1038/s41588-019-0479-7

28. Barutcu AR, Blencowe BJ, Rinn JL. 2019. Differential contribution of steady-state RNA and active transcription in chromatin organization . *EMBO Rep*. doi:10.15252/embr.201948068

29. Thakur J, Fang H, Llagas T, Disteche CM, Henikoff S. 2019. Architectural RNA is required for heterochromatin organization. *bioRxiv*. doi: https://doi.org/10.1101/784835

30. Swygert SG, Lin D, Portillo-Ledesma S, Lin PY, Hunt DR, Kao CF, Schlick T, Noble WS, Tsukiyama T. 2020. Chromatin Fiber Folding Represses Transcription and Loop Extrusion in Quiescent Cells. *bioRxiv*. doi: https://doi.org/10.1101/2020.11.24.396713

31. Hansen AS, Hsieh THS, Cattoglio C, Pustova I, Saldaña-Meyer R, Reinberg D, Darzacq X, Tjian R. 2019. Distinct Classes of Chromatin Loops Revealed by Deletion of an RNA-Binding Region in CTCF. *Mol Cell*. doi:10.1016/j.molcel.2019.07.039

32. Saldaña-Meyer R, Rodriguez-Hernaez J, Escobar T, Nishana M, Jácome-López K, Nora EP, Bruneau BG, Tsirigos A, Furlan-Magaril M, Skok J, Reinberg D. 2019. RNA Interactions Are Essential for CTCF-Mediated Genome Organization. *Mol Cell*. doi:10.1016/j.molcel.2019.08.015

33. Nandy Mazumdar M, Yin S, Paranjapye A, Kerschner JL, Swahn H, Ge A, Leir SH, Harris A. 2020. Looping of upstream cis-regulatory elements is required for CFTR expression in human airway epithelial cells. *Nucleic Acids Res*. doi:10.1093/nar/gkaa089

34. Gong Y, Lazaris C, Sakellaropoulos T, Lozano A, Kambadur P, Ntziachristos P, Aifantis I, Tsirigos A. 2018. Stratification of TAD boundaries reveals preferential insulation of super-enhancers by strong boundaries. *Nat Commun*. doi:10.1038/s41467-018-03017-1

35. Sun F, Chronis C, Kronenberg M, Chen XF, Su T, Lay FD, Plath K, Kurdistani SK, Carey MF. 2019. Promoter-Enhancer Communication Occurs Primarily within Insulated Neighborhoods. *Mol Cell*. doi:10.1016/j.molcel.2018.10.039

36. Matthews BJ, Waxman DJ. 2018. Computational prediction of CTCF/cohesin-based intra-TAD loops that insulate chromatin contacts and gene expression in mouse liver. *Elife*. 2018 May 14;7:e34077. doi: 10.7554/eLife.34077

37. Andersson R, Gebhard C, Miguel-Escalada I. et al. 2014. An atlas of active enhancers across human cell types and tissues. *Nature* 507, 455–461 (2014). https://doi.org/10.1038/nature12787

38. Skene PJ, Henikoff S. 2017. An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. *Elife*. 2017 Jan 16;6:e21856. doi: 10.7554/eLife.21856

39. Zhang J, Lee D, Dhiman V, Jiang P, Xu Jie, McGillivray P, Yang H, Liu J, Meyerson W, Clarke D, Gu M, Li S, Lou S, Xu Jinrui, Lochovsky L, Ung M, Ma L, Yu S, Cao Q,

Harmanci A, Yan KK, Sethi A, Gürsoy G, Schoenberg MR, Rozowsky J, Warrell J, Emani P, Yang YT, Galeev T, Kong X, Liu S, Li X, Krishnan J, Feng Y, Rivera-Mulia JC, Adrian J, Broach JR, Bolt M, Moran J, Fitzgerald D, Dileep V, Liu T, Mei S, Sasaki T, Trevilla-Garcia C, Wang S, Wang Y, Zang C, Wang D, Klein RJ, Snyder M, Gilbert DM, Yip K, Cheng C, Yue F, Liu XS, White KP, Gerstein M. 2020. An integrative ENCODE resource for cancer genomics. *Nat Commun*. doi:10.1038/s41467-020-14743-w

40. Zirkel A, Nikolic M, Sofiadis K, Mallm JP, Brackley CA, Gothe H, Drechsel O, Becker C, Altmüller J, Josipovic N, Georgomanolis T, Brant L, Franzen J, Koker M, Gusmao EG, Costa IG, Ullrich RT, Wagner W, Roukos V, Nürnberg P, Marenduzzo D, Rippe K, Papantonis A. 2018. HMGB2 Loss upon Senescence Entry Disrupts Genomic Organization and Induces CTCF Clustering across Cell Types. *Mol Cell*. doi:10.1016/j.molcel.2018.03.030

41. Farooq U, Saravanan B, Islam Z, Walavalkar K, Singh AK, Jayani RS, Meel S, Swaminathan S, Notani D. 2021. An interdependent network of functional enhancers regulates transcription and EZH2 loading at the INK4a/ARF locus. *Cell Rep*. doi: 10.1016/j.celrep.2021.108898.

42. Harismendy O, Notani D, Song X, Rahim NG, Tanasa B, Heintzman N, Ren B, Fu XD, Topol EJ, Rosenfeld MG, Frazer KA. 2011. 9p21 DNA variants associated with coronary artery disease impair interferon-γ 3 signalling response. *Nature*. doi:10.1038/nature09753

43. Bouvy-Liivrand M, De Sande AH, Pölönen P, Mehtonen J, Vuorenmaa T, Niskanen H, Sinkkonen L, Kaikkonen MU, Heinäniemi M. 2017. Analysis of primary microRNA loci from nascent transcriptomes reveals regulatory domains governed by chromatin architecture. *Nucleic Acids Res*. doi:10.1093/nar/gkx680

44. Nilson KA, Lawson CK, Mullen NJ, Ball CB, Spector BM, Meier JL, Price DH. 2017. Oxidative stress rapidly stabilizes promoter-proximal paused Pol II across the human genome. *Nucleic Acids Res*. doi:10.1093/nar/gkx724

45. Schlackow M, Nojima T, Gomes T, Dhir A, Carmo-Fonseca M, Proudfoot NJ. 2017. Distinctive Patterns of Transcription and RNA Processing for Human lincRNAs. *Mol Cell*. doi:10.1016/j.molcel.2016.11.029

46. Thiecke MJ, Wutz G, Muhar M, Tang W, Bevan S, Malysheva V, Stocsits R, Neumann T, Zuber J, Fraser P, Schoenfelder S, Peters JM, Spivakov M. 2020. Cohesin-Dependent and -Independent Mechanisms Mediate Chromosomal Contacts between Promoters and Enhancers. *Cell Rep*. doi:10.1016/j.celrep.2020.107929

47. McLaughlin-Drubin ME, Park D, Munger K. 2013. Tumor suppressor p16INK4A is necessary for survival of cervical carcinoma cell lines. *Proc Natl Acad Sci U S A*. 2013;110(40):16175-16180. doi:10.1073/pnas.1310432110

48. Pauck A, Lener B, Hoell M, Kaiser A, Kaufmann AM, Zwerschke W, Jansen-Dürr P. 2014. Depletion of the cdk inhibitor p16INK4a differentially affects proliferation of established cervical carcinoma cells. *J Virol*. 2014 May;88(10):5256-62. doi: 10.1128/JVI.03817-13.

49. Li W, Notani D, Ma Q, Tanasa B, Nunez E, Chen AY, Merkurjev D, Zhang J, Ohgi K, Song X, Oh S, Kim HS, Glass CK, Rosenfeld MG. 2013. Functional roles of enhancer RNAs for oestrogen-dependent transcriptional activation. *Nature*. doi:10.1038/nature12210

50. Li W, Notani D, Rosenfeld MG. 2016. Enhancers as non-coding RNA transcription units: Recent insights and future perspectives. *Nat Rev Genet*. doi:10.1038/nrg.2016.4

51. Quinodoz SA, Ollikainen N, Tabak B, Palla A, Schmidt JM, Detmar E, Lai MM, Shishkin AA, Bhat P, Takei Y, Trinh V, Aznauryan E, Russell P, Cheng C, Jovanovic M, Chow A, Cai L, McDonel P, Garber M, Guttman M. 2018. Higher-Order Inter-chromosomal Hubs Shape 3D Genome Organization in the Nucleus. *Cell*. 2018 Jul 26;174(3):744-757.e24. doi: 10.1016/j.cell.2018.05.024.

52. Beagrie RA, Scialdone A, Schueler M, Kraemer DC, Chotalia M, Xie SQ, Barbieri M, de Santiago I, Lavitas LM, Branco MR, Fraser J, Dostie J, Game L, Dillon N, Edwards PA, Nicodemi M, Pombo A. 2017. Complex multi-enhancer contacts captured by genome architecture mapping. *Nature*. 2017 Mar 23;543(7646):519-524. doi: 10.1038/nature21411.

53. Hansen AS. 2020. CTCF as a boundary factor for cohesin-mediated loop extrusion: evidence for a multi-step mechanism. *Nucleus*. doi:10.1080/19491034.2020.1782024

54. Kentepozidou E, Aitken SJ, Feig C, Stefflova K, Ibarra-Soria X, Odom DT, Roller M, Flicek P. 2020. Clustered CTCF binding is an evolutionary mechanism to maintain topologically associating domains. *Genome Biol*. doi:10.1186/s13059-019-1894-x

55. Hug CB, Grimaldi AG, Kruse K, Vaquerizas JM. 2017. Chromatin Architecture Emerges during Zygotic Genome Activation Independent of Transcription. *Cell*. doi:10.1016/j.cell.2017.03.024

56. Jayani RS, Singh A, Notani D. 2017. Isolation of nuclear RNA-associated protein complexes. *Methods in Molecular Biology*. doi:10.1007/978-1-4939-6716-2_9

57. Rahnamoun H, Lee J, Sun Z, Lu H, Ramsey KM, Komives EA, Lauberth SM. 2018. RNAs interact with BRD4 to promote enhanced chromatin engagement and transcription activation. *Nat Struct Mol Biol*. doi:10.1038/s41594-018-0102-0

Islam and Saravanan et al. Figure 1

Figure 1. Transcribed TAD boundaries insulate better than non-transcribed boundaries: A. Pie chart displaying the percentage of boundaries that show genic transcription, non-genic de novo transcription and no transcription. **B.** Schematic depicting genic and non-genic transcribed boundaries. **C.** Browser shot displaying a TAD structure from HiC data. The zoomed in box shows a *de novo* non-genic transcribed boundary region that is overlaid with CTCF and mNET-seq signal. **D.** Box plots showing CTCF enrichment on genic, non-genic and non-transcribed boundaries. **E.** Box plots showing CTCF enrichment on all boundaries exhibiting varying levels of RNAs versus non-transcribed and random boundaries. **F.** Pile up (Aggregated normalized HiC interactions) plot centered at high transcribed, low transcribed, non-transcribed and random boundaries at a 5 kb resolution. The 100 kb distances are taken from the boundary region. The p-values in boxplots were calculated using the Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al Figure 2

Figure 2. RNA increases CTCF occupancy at TAD boundaries: A. Immunoblotting with CTCF and Histone H3 on soluble nucleoplasm and chromatin bound fractions of the nucleus treated with or without RNaseA (Left panel). The right panel shows the quantified loss of CTCF from chromatin fractions from two replicates. **B.** Normalised CTCF CUT&RUN signal in control and RNAseA-treated conditions plotted at equally scaled TADs of 50 kb length each and 25 kb flanks**. C.** Box plot showing the fold loss in CTCF upon RNAseA treatment at boundaries with high RNA levels, non-transcribed and random boundaries. **D.** TAD structure at *INK4a/ARF* locus on 9p21 region (Rao et al., 2014). The loop domain (Blue bar) is overlaid by the position of CTCF peaks, CTCF motif orientation, H3K4me3, H3K27ac (enhancers), mNET-seq tracks and gene annotations. The highlighted region shows the B1, B2 and B3 CTCF sites on the 3' boundary and super-enhancer with the position of enhancer E8 marked by an arrow. **E.** qRT-PCRs showing the levels of sense and anti-sense non-coding RNA at B1 CTCF site upon B1 RNA knockdown. **F.** qRT-PCRs showing the levels of sense and anti-sense RNA at the B2 CTCF site upon B2 RNA knockdown. **G.** CTCF ChIP enrichment before and after shRNA mediated knockdown of sense and anti-sense RNA at B1 CTCF sites. The bars show CTCF fold enrichment on three CTCF sites (B1, B2 and B3) at the 3' boundary, on -T1 (5' boundary) of the *INK4a/ARF* TAD, on -T2 on the adjacent TAD boundary upstream, +T1 (on the adjacent TAD boundary downstream) and +T2, the boundary of the following TAD downstream. **H.** CTCF ChIP enrichment before and after shRNA mediated knockdown of sense and anti-sense RNA at B2 CTCF sites. The bars show CTCF on three CTCF sites (B1, B2 and B3) at the 3' boundary, on -T1 (5' boundary) of the *INK4a/ARF* TAD, on -T2 of the adjacent TAD boundary upstream, on +T1 of the adjacent TAD boundary downstream, and on +T2 of the boundary of the following TAD downstream. **I.** qRT-PCRs of sense and anti-sense RNAs from B1 and B2 CTCF sites upon CTCF knockdown. The drops in CTCF levels are shown in the last two bars. Error bars denote SEM from three biological replicates. p-values were calculated by Student's two-tailed unpaired t-test in **E, F, G and H .** $**p < 0.01, *p < 0.05, {}^{ns}p > 0.05$.

Islam and Saravanan et al. Figure 3

Figure 3. Non-coding RNAs and CTCF interact at the *INKa/ARF* TAD boundary:
A. Immunoblot for CTCF on RNA pulldowns performed on various RNAs (n=2). **B.** Agarose gel image showing the UV-RIP RT-PCRs for sense and anti-sense RNA arising from B1 and B2 CTCF sites (n=2). **C.** Line plots show the CTCF motif per 10 base pair per peak on various classes of boundaries. **D.** Fraction of boundaries with differential occupancy of CTCF peaks over the presence of the CTCF motif. **E.** CTCF peaks per kb on all transcribed boundaries with high RNA, no RNAs and random boundaries (50 kb flanks on both side). **F.** CTCF peaks per kb on non-genic transcribed boundaries with high RNAs, no RNAs and random boundaries. **G.** Log_2FC of NET-CAGE+1/CAGE+1, indicating stability of RNA at TSSs identified from NET-CAGE at boundaries and TSSs not at boundaries. eRNA denotes the eRNA from non-genic boundaries, mRNA denotes the mRNA from genic boundaries **H.** Average predicted TTS sites from the TSSs identified in different classes of RNA by NET-CAGE. **I.** Average predicted 5'SS sites from the TSSs identified by NET-CAGE. The p-values in boxplots were calculated by the Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al. Figure 4

Figure 4. Boundary RNAs enhance TAD transcription: A. Box plots showing GRO-seq tags from the entire TAD with high, low, non-transcribed and random boundaries. **B.** Graph showing the number of interactions between genes and boundaries, and 100 randomizations of interactions between boundaries and random regions. **C.** Box plots showing the GRO-seq tag counts from genes when they interact with boundaries or any region vs. random interactions. **D.** 4C plot at the *CDKN2A* promoter viewpoint, where the interaction with the boundary is highlighted in blue. Below are H3K27ac, CTCF and gene annotations for HeLa cells. **E.** qRT-PCR plot shows the change in gene expression upon shRNA mediated knockdowns of boundary RNAs arising from B1 and B2 CTCF sites at the *INK4a/ARF* TAD boundary. **F.** Immunoblot showing the drop in CTCF protein levels upon its siRNA mediated knockdown. **G**. qRT-PCRs showing down-regulation of *INK4a/ARF* genes and levels of DDX5 and CTCF upon CTCF knockdown. **H.** qRT-PCR showing the changes in expression of different genes in the *INK4a/ARF* TAD and DDX5 upon Cas9-mediated independent deletion of B1 and B2 CTCF sites. **I.** PolII ChIP signal at *INK4a*, *ARF* and *INK4b* promoters in WT and in cells with B1 and B2 CTCF sites deletions. Error bars denote SEM from three biological replicates. *p*-values were calculated by Student's two-tailed unpaired *t*-test in **E, G, H** and **I**. ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, $^{ns}p > 0.05$. p-values in boxplots were calculated by the Wilcoxon two-sided rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

A

chr9:21,790,079-22,269,127

B1 B2 B3 CBS

CTCF

H3K4me3

H3K27ac

NET-seq
(+strand)

[2.4, 30]

[-2.4, -30]

NET-seq
(-strand)

MTAP

CDKN2A
CDKN2A
CDKN2B-AS1
CDKN2B

B

C

Islam and Saravanan et al. Figure 5

Figure 5. Transcriptional activity of promoters does not regulate non-genic transcription at boundaries: A. Browser shots displaying the promoters targeted by CRISPRi (Black arrows), the track is overlaid with CTCF, H3K4me3, H3K27ac and mNET-seq tracks. **B.** qRT-PCR assessing the changes in *INK4a/ARF* genes, sense and anti-sense boundary RNA arising from B1 and B2 CTCF sites upon CRISPRi on promoters. **C.** CTCF enrichment on boundaries of *INK4a/ARF* and its flanking TADs upon CRISPRi on *INK4a/ARF* promoters. Error bars denote SEM from three biological replicates. *p*-values were calculated by Student's two-tailed unpaired *t*-test in **B** and **C.** **$p < 0.01$, *$p < 0.05$, ns$p > 0.05$.

Islam and Saravanan et al., Figure 6

Figure 6. Active enhancers within a TAD directly regulate boundary eRNA transcription : A. Boxplots displaying enrichment of H3K27ac in TADs with high, low, non-transcribed and random boundaries. **B.** Boxplot showing number of H3K27ac peaks in TADs with various boundaries. **C.** Levels of PolII in TADs with different boundaries. **D.** UCSC browser shot showing H3K27ac, p300, PolII ChIP-seq and mNET-seq on the *INK4a/ARF* TAD. The tracks are overlaid by the gene annotations. The highlighted regions represent the enhancers (E8, E12 and E17) that were deleted. **E.** Browser shot shows RNA-seq at *INK4a/ARF* TAD in WT and E8 enhancer delete lines. **F.** qRT-PCR shows levels of sense and anti-sense boundary RNA arising from B1 and B2 CTCF sites upon E8, E12 and E17 enhancer deletions. **G.** 4C-seq plot in WT and in E8 deletion lines on an enhancer with the E8 enhancer as viewpoint (Yellow highlight) exhibiting interactions with promoters and 3' boundary region (Pink highlight). Below are PolII, H3K27ac ChIP-seq tracks and gene annotations. Error bars denote SEM from three biological replicates. p-values were calculated by Student's two-tailed unpaired t-test in **F.** ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$. p-values in boxplots were calculated by the Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

A

Prom:enh interaction strength

3.054e-05

bRNA: High None

B

-T1 Boundary   CDKN2B / CDKN2A   SE   3' Boundary   +T1Boundary

Wild Type

ΔB1 CTCF

ShB1 sense
+ anti-sense
bRNA

Chr 9: 21.6   21.7   21.8   21.9   22   22.1   22.2   22.3   22.4   22.5   22.6 Mb

PolII   [0-2]
H3K27ac   [0-100]
CTCF   [0-1]

MTAP   CDKN2B   DMRTA1
CDKN2A
CDKN2A
CDKN2B-AS1

C

4. Robust E:P interactions   1. Enhancers induce bRNA

CTCF

5'B   P   E   CTCF

                                   2. bRNAs stabilize CTCF

                                   3. Better insulation

TAD

Enhancer

POLII

CTCF   bRNA   bRNA Knockdown ───▶   CTCF   CTCF

Enhancer Deletion

Islam and Saravanan et al. Figure 7

Figure 7. Boundary eRNAs strengthen TAD insulation: A. Enhancer:promoter interaction strength within TADs with high and non-transcribed boundaries. **B.** 4C-seq plot with the E8 enhancer as viewpoint (Yellow highlight) in WT and upon B1 CTCF site deletion (ΔB1) and knockdown of B1 sense and anti-sense boundary RNAs. The anchored region exhibit interactions with promoters and the 3' and +T1 boundaries (Pink highlights). Below are PolII, H3K27ac and CTCF ChIP-seq tracks and gene annotations. **C.** Model: (1) An active enhancer physically interacts with a boundary and activates boundary RNA transcription. (2) The boundary RNA in return stabilizes the CTCF at these boundaries, thereby (3) strengthening insulation of these TADs. (4) This favors intra-TAD enhancer: promoter interactions to facilitate robust gene transcription. The loss of boundary RNA/enhancer deletion reduces the boundary RNA levels which triggers the loss of CTCF and insulation of TADs. Enhancer:promoter interactions weaken in these scenarios which causes concomitant loss of gene transcription. p-values in boxplots were calculated by the Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al Sup Fig. 1

Supp Figure 1. Transcribed boundaries exhibit better insulation: A. CTCF enrichment (CTCF CUT&RUN) at non-genic transcribed boundaries with high, low RNA versus non-transcribed and random boundaries **B.** levels of RNA and CTCF occupancy at TAD boundaries are positively correlated. **C.** CTCF enrichment (ChIP-seq) at transcribed genic, non-genic and non-transcribed boundaries. **D.** CTCF enrichment at all transcribed boundaries with high, low RNA versus non-transcribed and random boundaries. **E.** CTCF enrichment at non-genic transcribed boundaries with high, low non-coding RNA versus non-transcribed and random boundaries. p-values in boxplots were calculated by the Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al Sup Fig. 2

Supp Figure 2. boundaries defined at 20 kb resolution show the similar features as at 5 kb resolution:

A. Pie chart displaying the percentage of boundaries that show genic and non-genic transcribed boundaries and not transcribed boundaries. **B.** Levels of RNA and CTCF occupancy at TAD boundaries are positively correlated. **C.** CTCF enrichment on transcribed genic, non-genic and non-transcribed boundaries. **D.** CTCF enrichment on all transcribed boundaries with high and low RNA versus non-transcribed and random boundaries. **E.** CTCF enrichment on non-genic transcribed boundaries with high and low non-coding RNA versus non-transcribed and random boundaries. **F.** Density plot showing insulation scores for transcribed genic, non-genic and non-transcribed boundaries. **G.** Density plot showing insulation scores for all transcribed boundaries exhibiting varying levels of RNA vs. non-transcribed and random boundaries. **H.** Density plot showing insulation scores of non-genic transcribed boundaries exhibiting varying levels of RNA versus non-transcribed and random boundaries. p-values in boxplots were calculated by Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al Sup Fig. 3

Supp Figure 3. Transcribed Loop Anchors exhibit more CTCF enrichment: A. Pie chart showing the percentage of loop anchors that overlap with regions of genic transcription and non-genic de novo transcription and regions without transcription. **B.** CTCF enrichment (CTCF CUT&RUN) at transcribed genic, non-genic and non-transcribed loop anchors. **C.** CTCF enrichment on all transcribed loop anchors with high and low RNA versus non-transcribed and random loop anchors. **D.** CTCF enrichment (CTCF ChIP-seq) at transcribed genic, non-genic and non-transcribed loop anchors. **E.** CTCF enrichment at all transcribed loop anchors with high and low RNA versus non-transcribed and random loop anchors. p-values in boxplots were calculated by Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al Sup Fig. 4

Supp Figure 4. Transcribed boundaries in IMR90 and K562 exhibit better insulation: A. CTCF enrichment at transcribed genic, non-genic and non-transcribed boundaries in IMR90 cells. **B.** CTCF enrichment at non-genic transcribed boundaries with high and low non-coding RNA versus non-transcribed and random boundaries in IMR90 cells. **C.** Density plots showing insulation scores for transcribed genic, non-genic and non-transcribed boundaries in IMR90 cells**. D.** Density plots showing insulation scores for non-genic transcribed boundaries exhibiting varying levels of non-coding RNAs versus non-transcribed and random boundaries in IMR90. **E-F**. Same as A-B in K562 cells. **G-H.** Same as C-D in K562 cells. p-values in boxplots were calculated by Wilcoxon rank-sum test. The boxplots depict the minimum (Q1-1.5*IQR), first quartile, median, third quartile and maximum (Q3+1.5*IQR) without outliers.

Islam and Saravanan et al. Supp Figure 5

Supp Figure 5. Browser shots of the *INK4a/ARF* loop domain: A. A browser shot showing the *INK4a/ARF* locus in the 9p21 region. Below are H3K4me3, H3K27ac (enhancers), PolII, CTCF, mNET-seq tracks, GRO-seq tracks, PRO-seq tracks and gene annotations. The highlighted region shows the B1, B2 and B3 CTCF sites at the 3' boundary and super-enhancer. **B**. Browser shot showing the zoomed-in region of the 3' boundary (B1 and B2 CTCF sites). Below are H3K27ac, H3K4me1, H3K4me2, PolII, CAGE seq tracks, mNET-seq tracks and CTCF peaks. Arrowheads at the bottom show CTCF motif orientation at the B1 and B2 CTCF sites.

Islam and Saravanan et al. Supp Figure 6

Supp Figure 6. RNA increases CTCF occupancy at TAD boundary: A. Immunoblotting with CTCF, Histone H3 and GAPDH on soluble nucleoplasm and chromatin‑bound fractions of nuclei treated with or without RNaseA. **B.** Browser shot showing CTCF CUT&RUN peaks at the *INK4a/ARF* TAD with and without RNaseA treatment. **C.** qRT-PCRs showing the levels of sense and anti-sense non-coding RNA at B1and B2 CTCF sites and of snoRNA upon B1 and B2 RNA knockdown in the chromatin associated RNA fraction. **D.** TADs around the *INK4a/ARF* TAD. Positions of arrows show the CTCF sites at upstream and downstream TAD boundaries where CTCF occupancy was interrogated. **E.** qRT-PCRs showing the fold-change in sense and anti-sense RNAs from B1 and B2 CTCF sites upon Flag-CTCF overexpression. The last two bars show the increase in CTCF levels. Error bars denote SEM from three biological replicates. p-values were calculated by Student's two-tailed unpaired t test in **C** and **E**. ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, [ns]$p > 0.05$.

A

Genic Boundaries

All Boundary
RNA

B

Non-genic Boundaries

Non-genic
Boundaries

Islam and Saravanan et al. Supp Figure 7

Supp Figure 7. TAD transcription is correlated with transcribed boundaries: A. Box plots showing the normalized GRO-seq reads per gene within TADs of high, low, non-transcribed and random boundaries. **B.** Box plots showing the normalized GRO-seq tags within the TADs of transcribed non-genic boundaries with high and low boundary RNAs, non-transcribed and random boundaries.

A

B1 CTCF   chr9:22,214,590-22,240,888   B2 CTCF

10 kb ⊢————————————————————————⊣ hg19

H3K27ac
H3K4me1
H3K4me2
PolII
CAGE-seq
mNET-seq ⌈ +strand
         ⌊ -strand
CTCF   1189bp   1051bp
CTCF Motif Orientation

WT GGCCCCACCGATCA TCCCCCAACCTACCTCCGCA - 1167nt- GTAATATTCACTACAAGCTG ACTGAAGATACTTTAAAC
B1 Del GGCCCCACCGATCA TCCCCCAACCTACCTCCGCA -C-GTAATATTCACTACAAGCTG ACTGAAGATACTTTAAAC
#c62

WT AGCATTTG AAAGAGGGCAGGTCTTTTT -1059nt- TGGGCCTCTGAGCCTGTAGA TTTCCTCCTCGCC
B2 Del AGCATTTG AAAGAGGGCAGGTCTTTTT-TGGGCCTCTGAGCCTGTAGA TTTCCTCCTCGCC
#c110

B

L   WT   B1 Del #C62
2 Kb
1.65Kb
                                              WT Amplicon
                                              1553bp
1 Kb

500bp
                                              Deleted Amplicon
300bp   364bp

L   WT   B2 Del #C110
2 Kb
1.65Kb
                                              WT Amplicon
                                              1668bp
1 Kb

650bp   Deleted Amplicon
500bp   617bp
300bp

C

              shscr shB1S+AS shB2S+AS
P16   16kDa
GAPDH   37kDa

              B1 DEL   B2 DEL
        WT C50 C62 C95 C110
P14   14kDa
P16   16kDa
DDX5   70kDa
GAPDH   37kDa

Islam and Saravanan et al Supp Fig. 8

Supp Figure 8. Deletion of B1 and B2 CTCF sites in the *INK4a/ARF* TAD boundary: A. UCSC genome browser track depicting the deleted B1 and B2 CTCF sites (boxed region at the bottom). Below are CAGE, NET-seq signal, CTCF peaks and CTCF orientation (Green arrows). The tracks zoom in to the chromatograms from Sanger sequencing on deleted amplicons to verify the deletion. **B.** Agarose gel images showing the amplicons from a Surveyor assay of B1 and B2 deletions. **C.** Immunoblot showing the protein levels of p16 after Scr shRNA knockdown against sense and anti-sense RNA at B1 and B2 CTCF sites (Upper panel). Immunoblot showing the protein levels of p14, p16, DDX5 and GAPDH in WT and in cells with B1 and B2 CTCF site deletions (Lower panel).

A

Non-genic Boundaries

B

Non-genic Boundaries

Islam and Saravanan Supp Fig. 9

Supp Figure 9. TADs with transcribed boundaries harbor more active enhancers: **A.** Boxplots displaying enrichment of H3K27ac in TADs with high and low transcribed non-genic HeLa boundaries versus non-transcribed and random boundaries. **B.** Levels of PolII in TADs with different non-genic transcribed boundaries vs. non-transcribed and random boundaries.

Islam and Saravanan et al. Supp Figure 10

Supp Figure 10. Transcriptional state of *INK4a/ARF* TAD in senescence : A. Browser images showing H3K27ac, PRO-seq reads on *INK4a/ARF* TAD from young and replicative senescent IMR90 fibroblasts. Highlighted regions depict the super-enhancer and the boundary. **B**. qRT-PCR plots show the fold change in expression of *INK4a* and *INK4b* in young and DNA damage-induced senescent BJ fibroblasts. **C.** qRT-PCR plots show the fold change in expression of boundary RNA from B1 and B2 CTCF sites in young and DNA damage-induced senescent BJ fibroblasts. **D.** ChIP qRT-PCRs showing the changes in fold enrichment of CTCF at B1 and B2 CTCF sites in young and DNA damage-induced senescent BJ fibroblasts. Error bars denote SEM from three biological replicates. *p*-values were calculated by Student's two-tailed unpaired *t*-test in **B**, **C**, and **D** . ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, $^{ns}p > 0.05$. **E.** Immunoblot showing the changes in protein levels of CTCF, p21, p16 and GAPDH in young and DNA damage-induced senescent BJ fibroblasts. **F.** *INK4a/ARF* TAD structure in proliferating and senescent cells. Lower boxes represent the transcriptional levels.

Table 1: Oligonucleotide sequences used in the study

Primers used for ChIP-qPCRs

CTCF sites	Forward Primer (5'-3')	Reverse Primer (5'-3')
-T2	TAAATCTCCTGCCAGGAGG	GGCAGGTTCCATTTTACTTTGAG
-T1	GGCAGTACTGAGTTGACAC	CCCAGGAAACTGGACCTCTAA
B1	AGCTCTCTGCACCACATTGCCACTGCC	CCAACCTACCTCCGCAAGGACACC
B2	TTGAGGACATCTAGCGGCG	GCTGCAGATTCTGGAATGAGC
B3	GTCAGGACTAGGCCCATCTGAC	TAGGTGTCCAAGTGAAGATTAAAGAG
+T1	CAACTCCTTCCGAGTGGAAA	ACACTGTGACCGCTCCATTC
+T2	GCATAGATGTTTGGAGGGCAAC	TTACAGAATCTGGCTACTGCCC
INK4a	TCGCCAGGAGGAGGTCTGTGATTAC	CAGGTGGGTAGAGGGTCTGCAGC
ARF	ACCCAGGATATTCGGGACTCACTGAC	CGTCTCTAGCCCAGGCTAGGAGG
INK4b	CCGTCGTCCTTCTGCGGCTTG	AGTGAGGACTCCGCGACGCGT

Primers used for qRT-PCRs

mRNA/bRNA	Forward Primer (5'-3')	Reverse Primer (5'-3')
MTAP	TTGACAGGACCACTATGAGACCTC	AGGTCCCTCGATTGTGACCATT
INK4a	ATGGAGCCTTCGGCTGACT	CACCAGCGTGTCCAGGAAG
ARF	AACATGGTGCGCAGGTTCT	CACCAGCGTGTCCAGGAAG
INK4b	CGTTAAGTTTACGGCCAACG	CCATCATCATGACCTGGATCG
DDX5	CCTACCTTGTCCTTGATGAAGC	CACTCAGTTCAAGTGCACCA
CTCF	TACAAACACACCCACGAGAAG	GGGTAAACCGAGCATGACAA
b1S	CCACAGCAGAATAGAGCACC	AAGTCCAGCCAGTCTCAGCA
b1AS	TGAGGCTTGTAGGAACTCAAGTTC	AGTGGCAATGTGGTGCAGAG
b2S	ACAGAGGTCGGATGTTCGTG	TAAACCGCCGCTAGATGTCC
b2AS	CTCCTGAGCTCTAAGTGAGATAGAGG	GGTTATTCACTCACAGTGCTCTTCAG
sno RNA	CGTGTAGAGCACCGAAAACC	CACTCAGACCGCGTTCTCTC
pre c-MYC	CTTCTCTCCGTCCTCGGATTC	CCTTCCTAATAAGAGTGGCCCG
GAPDH	CGCTCTCTGCTCCTCCTGTT	CCATGGTGTCTGAGCGATGT

shRNAs used for knockdown

shRNA	Primer sequence (5'-3')
shRNA1 Boundary 1 S F	CCGGGAGTTGTGAGACCCTCATTTCCTCGAGGAAATGAGGGTCTCACAACTCTTTTTG
shRNA1 Boundary 1 S R	AATTCAAAAAGAGTTGTGAGACCCTCATTTCCTCGAGGAAATGAGGGTCTCACAACTC
shRNA2 Boundary 1 F	CCGGGGACCCTGAGCTGTGAATAACCTCGAGGTTATTCACAGCTCAGGGTCCTTTTTG
shRNA2 Boundary 1 R	AATTCAAAAAGGACCCTGAGCTGTGAATAACCTCGAGGTTATTCACAGCTCAGGGTCC
shRNA3 Boundary 1 S F	CCGGGAGTCCTAGCCAGGTAATATTCTCGAGAATATTACCTGGCTAGGACTCTTTTTG
shRNA3 Boundary 1 S R	AATTCAAAAAGAGTCCTAGCCAGGTAATATTCTCGAGAATATTACCTGGCTAGGACTC
shRNA1 Boundary 1 AS	CCGGCTGGCTCCGGCTGGTTTAAATCTCGAGATTTAAACCAGCCGGAGCCAGTTTTTG

F	
shRNA1 Boundary 1 AS R	AATTCAAAAACTGGCTCCGGCTGGTTTAAATCTCGAGATTTAAACCAGCCGGAGCCAG
shRNA2 Boundary 1 AS F	CCGGCGGCTCTGCTTTCAGCTATAACTCGAGTTATAGCTGAAAGCAGAGCCGTTTTTG
shRNA2 Boundary 1 AS R	AATTCAAAAACGGCTCTGCTTTCAGCTATAACTCGAGTTATAGCTGAAAGCAGAGCCG
shRNA3 Boundary 1 AS F	CCGGAGGTAAGATGTAAGCATTAATCTCGAGATTAATGCTTACATCTTACCTTTTTTG
shRNA3 Boundary 1 AS R	AATTCAAAAAAGGTAAGATGTAAGCATTAATCTCGAGATTAATGCTTACATCTTACCT
shRNA1 Boundary 2 S F	CCGGCGCAATCCAACTATTCAATTTCTCGAGAAATTGAATAGTTGGATTGCGTTTTTG
shRNA1 Boundary 2 S R	AATTCAAAAACGCAATCCAACTATTCAATTTCTCGAGAAATTGAATAGTTGGATTGCG
shRNA2 Boundary 2 S F	CCGGGCTTTCCTGAAGGAGAGTAATCTCGAGATTACTCTCCTTCAGGAAAGCTTTTTG
shRNA2 Boundary 2 S R	AATTCAAAAAGCTTTCCTGAAGGAGAGTAATCTCGAGATTACTCTCCTTCAGGAAAGC
shRNA3 Boundary 2 S F	CCGGTAGCAATGGCAGAACCTATTTCTCGAGAAATAGGTTCTGCCATTGCTATTTTTG
shRNA3 Boundary 2 S R	AATTCAAAAATAGCAATGGCAGAACCTATTTCTCGAGAAATAGGTTCTGCCATTGCTA
shRNA1 Boundary 2 AS F	CCGGAGGAACTACATTGACTTATTTCTCGAGAAATAAGTCAATGTAGTTCCTTTTTTG
shRNA1 Boundary 2 AS R	AATTCAAAAAAGGAACTACATTGACTTATTTCTCGAGAAATAAGTCAATGTAGTTCCT
shRNA2 Boundary 2 AS F	CCGGCGACCTCTGTAAATGAAATAGCTCGAGCTATTTCATTTACAGAGGTCGTTTTTG
shRNA2 Boundary 2 AS R	AATTCAAAAACGACCTCTGTAAATGAAATAGCTCGAGCTATTTCATTTACAGAGGTCG
shRNA3 Boundary 2 AS F	CCGGTGGATTGCGATTTCCTGAAATCTCGAGATTTCAGGAAATCGCAATCCATTTTTG
shRNA3 Boundary 2 AS R	AATTCAAAAATGGATTGCGATTTCCTGAAATCTCGAGATTTCAGGAAATCGCAATCCA

Primers used for RNA IP

Target		Primer sequence (5'-3')
B1 S in pCDNA 3 Box B	F	AGACTGGCGGCCGCTAGGCCTCAAGGACTGCAA
B1 S in pCDNA 3 Box B	R	AGACTGTCTAGACTCCTGGGAGCCAGAGAA
B1 AS in pCDNA 3 Box B	F	AGACTGGAATTCAGTAGGTCGCATGCAACCACA
B1 AS in pCDNA 3 Box B	R	AGACTGGCGGCCGCTCCTACATCAGCTGTGGCACA
B2 S in pCDNA 3 Box B	F	AGACTGGAATTCGCCTCTGAGCCTGTAGAGACCAA
B2 S in pCDNA 3 Box B	R	AGACTGGCGGCCGCCTCTGGGTGAGGCAGTGAACAC
B2 AS in pCDNA 3 Box B	F	AGACTGGAATTCATCCTCCAGGACCACAGCAGT
B2 AS in pCDNA 3 Box B	R	AGACTGGCGGCCGCGCATACCCTGGAGTGAGAGGTTG
E8 in pCDNA 3 Box B	F	AGACTGGAATTCAATGCCATATGGTGGCAATG
E8 in pCDNA 3 Box B	R	AGACTGGCGGCCGCATCTGTTCCTTGCCTCTTCCAC

gRNAs used for deletions and CRISPRi

gRNA	Sequence
B1 DEL gRNA 1	TTAAATTGGTGTCCTTGCGG
B1 DEL gRNA 2	GTAGTGAATATTACCTGGCT
B2 DEL gRNA 1	CAGACTTCCGTAGTCCCCAT
B2 DEL gRNA 2	TTGGGTAGCATCTTCATCGG
MTAP gRNA 1	AGCAGTGCGGAGACCAGGGG
MTAP gRNA 2	CTCACTTGCCGCGCGAGGAG
INK4a gRNA 1	AGCCGTTTTACACGCAGGAG
INK4a gRNA 2	GACAGGACAGTATTTGAAGC
ARF gRNA 1	ATGCGCCCCGGACTTTTCGA
ARF gRNA 2	CTGCCCCCTTAACTGCAGAC
INK4b gRNA 1	AAAGAATTCCGTTTTCAGCT
INK4b gRNA 2	CTAGGAAGGAGAGAGTGCGC

Surveyor Oligos

Target	Forward Primer (5'-3')	Reverse Primer (5'-3')
B1 DEL	TGGAGTGGTAGTAGTGTGGGAAG	GTACGAGGGTTCTCTTTTGCTTG
B2 DEL	CACAACCCCCATCTCTCTGTAAC	TGAAGTGAGGGGAGTAGACAAGG

4C oligos for enhancer and Promoter viewpoints

View Point	Primer sequence (5'-3')
Enhancer F	AATGATACGGCGACCACCGAACACTCTTTCCCTACACGACGCTCTTCCGATCTGGTATTCTGACTAGAAGCTT
Enhancer R	CAAGCAGAAGACGGCATACGATAGTGGTGCTACAGTACCAC
Promoter F	AATGATACGGCGACCACCGAACACTCTTTCCCTACACGACGCTCTTCCGATCTGGAGGAGCTAGGGCAAGCTT
Promoter R	CAAGCAGAAGACGGCATACGACAGATGGATAAGTAGTAGCAAG

Table 3: Datasets used in the study

Experiment	Annotation	Accession Number	References
HeLa GRO-seq	SRR1596500	GSM1518913	Andersson et al., 2014. Nat Commun.
HeLa GRO-seq	SRR1596501	GSM1518914	Andersson et al., 2014. Nat Commun.
HeLa HiC	SRR1050968	-	Burton et al., 2013. *Nat Biotechnol.*
HeLa HiC	SRR1050969	-	Burton et al., 2013. *Nat Biotechnol.*
HeLa HiC	SRR3203899-SRR3203906	-	Rao et al., 2014. Cell.
HeLa PolII	wgEncodeSydhTfbsHelas3Pol2StdSig	GSM935395	ENCODE
HeLa CTCF	wgEncodeUwTfbsHelas3CtcfStdRawRep1	GSM749729	ENCODE
HeLa CTCF	wgEncodeUwTfbsHelas3CtcfStdRawRep2	GSM749739	ENCODE
HeLa H3K27ac	wgEncodeBroadHistoneHelas3H3k27acStdSig	GSM733684	ENCODE
IMR90 HiC	SRR5494761	GSM2595584	Zirkel etal., 2018. Mol Cell
IMR90 PRO seq	SRR6205688	GSM2828779	Sen et al., 2019. Mol Cell
IMR90 CTCF	wgEncodeSydhTfbsImr90CtcfbIggrabSig	GSM935404	ENCODE
K562 HiC	SRR1658693-SRR1658702	GSM1551618-GSM1551623	Rao et al., 2014. Cell.
K562 CTCF	wgEncodeUwTfbsGm12878CtcfStdAlnRep1	GSM749690	ENCODE
K562 GROseq	SRR1552484	GSM1480325	Core et al., 2014. Nat Genet.
HeLa_GRO-Seq	SRR5109939	GSM2428725	Bouvy-Liivrand et al., 2017. Nucleic Acids Res
PRO-Seq-DMSO	SRR5799635	GSM2692352	Nilson et al., 2017.Nucleic Acids Res
Total mNET-Seq	SRR4436479	GSM2357382	Schlackow et al., 2017. Mol Cell