Title: **Citrus photosynthesis and morphology acclimate to phloem-affecting huanglongbing**

**disease at the leaf and shoot levels**

Authors: Mark Keeley[1], Diane Rowland[1], and Christopher Vincent[2]*

[1] Agronomy Department, University of Florida, Gainesville, FL

[2] Horticultural Sciences Department, Citrus Research and Education Center, University of

Florida

*To whom correspondence should be directed:

civince@ufl.edu

Citrus Research and Education Center

University of Florida

700 Experiment Station Rd.

Lake Alfred FL 33850

Keywords:

Photosynthesis, phloem function, Citrus, huanglongbing, gas exchange

Running title:

Citrus photosynthesis acclimates to huanglongbing

**Abstract**

Huanglongbing (HLB) is a phloem-affecting disease of citrus that reduces growth and has impacted citrus production in most global production regions.  HLB is caused by a phloem-limited bacterium (*Candidatus* Liberibacter asiaticus; *C*Las). By inhibiting phloem function, HLB stunts sink growth, including reducing production of new shoots and leaves, and induces hyperaccumulation of foliar starch. Despite evidence that HLB induces feedback inhibition of photosynthesis by reducing foliar carbohydrate export, its effects on net $CO_2$ assimilation ($A_{net}$) have not been reported. In this work we assessed the relationship of bacterial distribution within the foliage, foliar starch accumulation, and $A_{net}$. Because HLB impacts canopy morphology, we developed a chamber to measure whole shoot $A_{net}$, and tested the effects of HLB at both leaf and shoot levels. Starch accumulation was correlated with bacterial population, and starch was negatively correlated with $A_{net}$ at the leaf but not at the shoot level. Starch increased between the uninfected group and the shortest duration of infection, then decreased progressively with increasing length of infection. HLB infection reduced $A_{net}$ at the leaf level, but increased it at the whole shoot level. We attribute this enhancement of whole shoot $A_{net}$ to the increased contribution of stem photosynthesis due to the altered shoot morphology induced by the disease. Despite the increased photosynthetic efficiency, total carbon fixation per shoot decreased because shoot size and leaf area were reduced. Overall, our results indicate a localized relationship of *C*Las distribution with negative impacts on foliar carbohydrate export and photosynthesis. Additionally, starch accumulation and $A_{net}$ acclimate over time at the shoot level to the localized impacts of the disease. Stems contribute important proportions of whole shoot $A_{net}$, and these contributions are likely increased by the morphological acclimation induced by

HLB. This study highlights the importance of temporal and spatial scale in assessing

photosynthesis.

**Introduction**

Huanglongbing (HLB) is a disease of citrus causing economic loss to the citrus industry

around the world (Bové, 2006; Singerman and Useche, 2016). Its causal agent, *Candidatus*

Liberibacter asiaticus (*C*Las), vectored by the Asian citrus psyllid (ACP; *Diaphorina citri*

Kuwayama), is a phloem-limited bacterium which affects phloem dysfunction leading to starch

accumulation in leaves and loss of sink growth (Achor et al., 2020; Bové, 2006), as well as

eventual development of thin canopies (Tang et al., 2019).

Because of the reduction in phloem function, HLB is hypothesized to lead to feedback

inhibition of photosynthesis (Cimò et al. 2013). Assessing chlorophyll fluorescence, a reduction

in maximal PSII quantum yield was found in *C*Las-infected citrus leaves compared to uninfected

leaves (Cen et al., 2017; Sagaram & Burns, 2009). On the one hand, starch accumulation in the

leaves has been noted as a symptom of HLB, which indicates a surplus of fixation relative to

export (Etxeberria et al., 2009; Folimonova & Achor, 2010; Schneider, 1968). On the other hand,

starch accumulation can disintegrate the chloroplast membrane (Nebauer et al., 2011; Snyder-

Leiby & Wang, 2008), which may contribute to the reduction in PSII yield noted above.

However, no measures of the impacts of HLB on net $CO_2$ assimilation ($A_{net}$) have been reported.

Measures of $A_{net}$ in citrus have previously been taken at the leaf level (Kriedemann, 1968;

Mesquita et al., 2016; Tsagarakis et al., 2012; Vu and Yelenosky, 1988; Vu et al., 1986;

Zambrosi et al., 2011) and at the field level (Contreras et al. 2017; Er-Raki et al., 2009; Maestre-

Valero et al., 2017; Penddinti and Kambhammettu, 2019). Due to the specificity of the leaf level

measurements (light levels, angle of radiation incidence, leaf position within the canopy), leaf-

level data is difficult to extrapolate to the whole canopy. Because, HLB induces reductions in

canopy density (Tang et al. 2019), larger scale measures may be more indicative of changes in

carbon assimilation than leaf-level measures.

In this study, our objective was to compare relationships of $A_{net}$, $C$Las infection, and

foliar symptoms at the leaf and shoot level. To achieve our objective, we first developed and

validated the use of a gas exchange chamber capable of measuring whole shoots. We assessed

the value of such measures by comparing leaf-level and shoot-level light response curves. In our

subsequent HLB experiment, we hypothesized that: 1) greater time of $C$Las inoculation exposure

would increase starch content; 2) higher starch content, as an indicator of reduced leaf

carbohydrate export, would correlate with downregulation of $A_{net}$; and 3) because HLB causes

morphological changes at the canopy level, mean leaf area, leaf number, and stem surface area

and length were expected to affect $A_{net}$ at the shoot level.

**Materials and Methods**

*Study 1: Whole shoot gas exchange chamber development and testing*

*Plant Material*

The trees utilized were greenhouse-raised, four-year-old sweet orange 'Valencia' (*Citrus*

*x sinensis*) on 'Kuharske' (*C. sinensis* x *Poncirus trifoliata* (L.) Raf.) rootstock in 20-L pots.

Plants were maintained with recommended rates of slow-release fertilizer and 3x weekly

irrigation with sufficient volume to exceed field capacity of the pots.

*Chamber construction*

The chamber was built of acrylic tubing, acrylic glue, and a 3-D printed adapter designed

and provided by Jason Hupp, LI-COR Bioscience, Inc. Lincoln, NE, USA. The chamber was fit

to an infrared gas analyzer (LI-6800, LI-COR Bioscience, Inc. Lincoln, NE, USA) using a

custom chamber adapter (LI-6800-19, LI-COR Bioscience, Inc. Lincoln, NE, USA). To seal the

opening, a large conical stopper was step-drilled and cut to provide an easy-to-fit cap around the

stem of the branch (Appendix S1). The chamber could then be made airtight using plumbers'

putty.

Leaves and shoots were measured on trees recently acclimated to the relative darkness of

the laboratory environment (2-4 µmol photons $m^{-2}$ $s^{-1}$). Leaf-level measurements were recorded

for 15 minutes on three leaves at photosynthetic photon flux density (PPFD) of 1000 µmols $m^{-2}$

$s^{-1}$. Environmental settings for the leaf level were set at flow 300 µmol $s^{-1}$, 0.2 kPa pressure, 60%

relative humidity, reference $CO_2$ at 400 µmol $CO_2$ $mol^{-1}$, and a fan speed of 10,000 RPM.

Chamber measurements were logged every 15 seconds for five minutes after a 15-minute

stabilization period under controlled conditions at PPFD of 1000 µmol $m^{-2}$ $s^{-1}$.  Environmental

settings for the whole-shoot chamber were set based on volume, 1000 cubic centimeters, at flow

1000 µmol $s^{-1}$, 0.2 kPa pressure, 60% relative humidity, reference $CO_2$ of 400 µmol $CO_2$ $mol^{-1}$,

and a fan speed of 14,000 RPM to maximize air mixture. Increases in air flow rate from 300

µmol $s^{-1}$ in leaf measurements to 1000 µmol $s^{-1}$ in whole -shoot chamber compensated for the

increase in photosynthetic material and increased volume present in the custom chamber. The fan

speed was increased to 14,000 RPM to help mix air in the chamber as recommended by the

IRGA manufacturer. Using the red-light capability of the red, far red, and blue light adjustable

fluorometer head unit of the LI-6800F, light intensity was controlled for the leaf-level

measurements. An LED Grow Light of similar spectral composition (ZXMean 1500W Full

Spectrum LED, BliGli, Wuhan HuBei, China) was manipulated by distance and a dimming

switch (Credenza Plug-In Lamp CFL-LED Dimmer, Lutron Electronics Co., Coopersburg, PA,

USA) to achieve desired PPFD for the whole-shoot chamber. PPFD for whole-shoot

measurements was measured using the quantum sensor of the LI-6800 at the mid-point of the

shoot adjacent to the whole-shoot chamber.

*Stem assimilation*

To determine stem contribution to $A_{net}$ of the shoot (the leaf and stem material together),

the stems were covered with fitted black drinking straws following the method of Nilsen and Bao

(1990) testing stem contributions to photosynthesis in soybean (*Glycine max* L.). Shoots were

exposed to PPFD 1400 µmol m$^{-2}$ s$^{-1}$ as described above. The shoot was then removed from the

chamber and black straws were measured to fit the space between petioles on the stem, sliced

lengthwise and wrapped around the stem (Appendix S2). Gas exchange measurements then

rerecorded with the darkened stems. The LED Grow Light was then turned off and the custom

chamber was covered by a black cloth to achieve 0 µmol m$^{-2}$ s$^{-1}$ PPFD to determine respiration of

the shoot, again following the same stabilization and logging duration. Stem (shoot $A_{net}$), without

stem ($A_{net}$ with darkened stem), and at 0 µmol m$^{-2}$ s$^{-1}$ PPFD (shoot respiration) were used for

analysis. Measurements of $A$ under the different measurement conditions were used to calculate

gross assimilation ($A_{gross}$) contributions of the stem and leaves as follows:

• Shoot $A_{net}$: shoot net photosynthesis, measured as $A$ of shoot at 1400 µmol m$^{-2}$ s$^{-1}$ PPFD;

inherently the sum of stem and leaf $A_{gross}$ minus respiration (shoot $R$).

•   Stem $A_{gross}$: stem gross $A$; measured as the difference between shoot $A_{gross}$ under full

illumination and $A_{gross}$ with darkened stem.

•   Shoot $R$: whole shoot respiration, measured as $|A|$ at 0 μmol m$^{-2}$ s$^{-1}$ PPFD.

•   Shoot $A_{gross}$: shoot gross assimilation, estimated as shoot $A_{net}$ – shoot $R$.

•   Leaf $A_{gross}$: leaf gross assimilation of all leaves on the shoot, estimated as Shoot $A_{gross}$ –

stem $A_{gross}$

Immediately following gas exchange measurements, shoots were excised and defoliated

for leaf area analysis using the LI-3100 (LI-COR Biosciences, Inc. Lincoln, NE, USA). Stems

were measured for length and diameter (Appendix S3) to calculate surface area using an adjusted

formula for the lateral area of a truncated cone (Equation 1). Due to the irregular shape of citrus

stems, two diameter measurements were taken at the widest and narrowest widths at the base,

middle, and tip of the stem. The circumference of an ellipse is approximately equal to that of a

circle with a radius of the average of the widest and narrowest point of the ellipse.

$Stem\ Surface\ Area\ =\ \pi(r_1 + r_2)\sqrt{(r_1 - r_2)^2 + h_1{}^2} + \pi(r_2 + r_3)\sqrt{(r_2 - r_3)^2 + h_2{}^2}$   (Eq. 1)

Where:

$r_1$= radius of stem base

$r_2$= radius of stem middle

$r_3$= radius of stem tip

$h_1$ =length between measurement point of $r_1$ and $r_2$

$h_2$ =length between measurement point of $r_2$ and $r_3$

Leaves and stems were dried at 60 °C for 72 hours and weighed separately. Leaf and

stem surface area and dry weight were used to calculate $A_{\text{gross}}$ rates per meter squared and per

gram dry weight.

*Leaf and shoot light response curves*

Nine fully mature young shoots, three from each of the same trees were measured in the

laboratory. Shoots were placed in the shoot gas exchange chamber and carbon assimilation was

measured at PPFD of 0, 25, 50, 100, 150, 200, 400, 600, 800, 1000, 1400, and 1800 μmols m$^{-2}$ s$^{-}$

$^{1}$. Environmental settings, stability criteria, and logging were the same as previous whole-shoot

measurements (Appendix S4). The largest, healthiest leaf (assessed visually) on each of the nine

measured shoots was selected to measure a leaf-level comparison (Appendix S5) measurement at

PPFD of 0, 25, 50, 100, 150, 200, 400, 600, 800, 1000, 1400, and 1800 μmols m$^{-2}$ s$^{-1}$.

Environmental settings and stability criteria were set as in the previously described

measurements.

*Study 2: HLB impacts on leaf and shoot net $CO_2$ assimilation*

*Plant material selection*

Trees were selected based on length of exposure to citrus greening infection. The trees

utilized were greenhouse-raised, four to five-year-old sweet orange 'Valencia' on 'Kuharske'

rootstock in 20-L pots. These trees had been exposed to ACP inoculation of *C*Las by growing

plants in infested screenhouses for varying lengths of time. Exposure categories were defined as

follows: the High category of exposure had been exposed to infected ACP feeding and breeding

for 2.5 years; the Mid category of exposure had been exposed to infected ACP feeding and

breeding for 1.5 years; and the Low category of exposure had been exposed to infected ACP

feeding and breeding for 1 year. The uninfected trees had not been exposed to infected ACP.

Three trees of each category were arranged in a greenhouse in a randomized complete block

design, totaling 12 trees for the study.

Shoots and leaves were selected on each tree for measurements. Three shoots per tree

were selected from the exterior of the canopy. Shoots were chosen based on accessibility of the

instrument using the custom chamber, and only those with at least three leaves suitable for leaf

measurement using the standard leaf chamber were selected. Thirty-six shoots were measured in

this study (3 shoots x 3 trees x 4 exposure groups).Three leaves were selected per shoot. These

were distributed along the length of the shoot, always using one apical leaf. To be selected,

leaves had to be large enough to completely cover the leaf chamber (6-cm$^2$ circle). A total of 108

leaves were measured for this study (36 shoots x 3 leaves).

*Gas exchange measurements*

$A_{net}$ was measured in using the LI-6800 (LI-COR Bioscience, Inc. Lincoln, NE, USA).

Leaf chamber environmental settings were set at a PPFD of 1000 µmol m$^{-2}$ s$^{-1}$ and shoot

chamber was placed at PPFD of 1400 µmol m$^{-2}$ s$^{-1}$ with all other settings according to those

described above for Study 1 for leaf and shoot gas exchange measurements.

*CLas quantification*

Midrib and petiole tissues excised from selected leaves were chopped using a razor blade

and bead-beaten using Vortex Genie 2 (Scientific Industries Bohemia, NY, USA) to a fine

homogenized powder in liquid nitrogen. Total genomic DNA was extracted and purified from

100 mg of homogenized material using GeneJET Plant Genomic DNA Purification Mini Kits

(Thermo Fisher Scientific Waltham, MA, USA). Samples were processed following

manufacturer instructions and stored at -20° C. Primers and TaqMan probes were obtained for

CLas based on method developed by Coy et al. (2014). The PCR master mix was used with 10

μL TaqMan Buffer, 8.6 μL sterile deionized water, 0.4 μL CQUL primer, and 1 μL of extracted

purified DNA. PCR reactions performed with standards using QuantStudio 3 Real-time PCR

(Applied Biosystems-Thermo Fisher Scientific Waltham, MA, USA) yielded cycle threshold

values (Ct). Bacterial copy-count estimates were generated by regression using processed Ct

results from standards of known copy quantity, processed in each plate batch. At the shoot level

Ct and bacterial copy number were considered to be the mean of the measurements in the three

selected leaves on each shoot.

*Starch content*

Starch content was determined by extracting starch from a single 28.27 mm$^2$ hole punch

on the selected leaf blade (Whitaker et al., 2014). The leaf punch was homogenized for 2.5

minutes at 6500 RPM in 500 μL of distilled water with five beads. Samples were boiled for 10

minutes at 200° C and agitated prior to a two-minute centrifuge at 2500 rpm, after which 300 μL

of the supernatant was added to 900 μL pure ethanol and agitated for three seconds to precipitate

starch. Prepared samples were centrifuged at 14000 RPM for 10 minutes and the supernatant was

discarded and replaced by one mL of deionized water. Samples were then shaken for four-

minutes to dissolve the starch pellet and two 250 μL of prepared sample were each placed in

individual wells of a 96 well microplate, and 50 μL of 2% iodine was added to each well.

Absorbance at 595nm was read using SpectraMax 250 microplate reader (Molecular Devices San

Jose, CA, USA). Linear functions based on standard curve concentrations included in each plate

were generated with the absorbance of standards to estimate starch content. As for $C$Las

variables, at the shoot level starch values were considered to be the mean of the measurements of

the three selected leaves on each shoot.

*Visual rating*

A rating of the visual symptoms of $C$Las infection was assessed on a scale of 0-10, where

0 = no symptom, 10= full symptom expression, on each leaf. Prior to assessment a scale of leaf

symptoms was generated using photos of sampled $C$Las infected leaves to ensure consistency in

ratings. All disease ratings were performed by the same individual. Visual symptom ratings used

in the leaf level measurements were recorded for every leaf on the measured shoot; an average of

the assessment of all leaves was used for the shoot value.

*Shoot morphology*

Total leaf area per shoot was measured using a LI-3100 (LI-COR Bioscience, Inc.

Lincoln, NE, USA) immediately following the assimilation measurements and visual ratings,

prior to sampling for qPCR and starch content. Stem length and diameter in two directions was

measured using a micrometer to generate a stem surface area estimates using Eq. 3. Average

internode length was calculated by dividing the stem length by the total number of leaves per

stem.

*Analysis*

Analysis of variance was performed using the anova() command in base R, after

assessing appropriateness of parametric tests by testing for normal distribution, to assess the

effect of treatment groups (R Core Team, 2019, Version 3.6.1). Where exposure group was

significant, Fisher's LSD was used as a post hoc test for least significant difference between the

four tree categories used for all variables using the {agricolae} package (de Mendiburu, 2020).

Correlation analysis was generated among the variables at the shoot and leaf-level and

correlation matrices were tested for significance of Pearson's correlations using the

{Hmisc}package (Harrell, 2020). For correlation analyses, because some variables might co-vary

in the case of disease, tests were performed with and without the HLB- plants. For variables with

significant correlations, linear regression models were generated at both shoot and leaf-level

using the lm() command, similarly comparing relationships with and without HLB- plants. The

models were tested for significant relationships using anova(). All linear models nested shoot

within tree, and leaf-level data was nested leaf within shoot. A natural log transformation of

bacterial copy count and starch content facilitated a linear relationship between the transformed

data. Light response curves for leaf and shoot were fit using an nonlinear least squares regression

using the self-starting asymptotic function in R (Pinheiro et al., 2014) with a function supplied

by Tomeo (2017). This function fits an asymptotic function and calculates the parameters of

fully saturated photosynthesis ($A_{sat}$), quantum yield of photosynthesis ($\Phi$), dark respiration ($R_d$),

light compensation point (LCP), and PPFD of 75% saturation ($Q_{sat75}$). These parameters were

then subjected to analysis of variance to test the effects of shoot vs. leaf light responses.

**Results**

*Stem contributes significantly to whole-shoot photosynthesis*

Vegetative stem on young shoots contributed 5.7% to total shoot surface area, 17.0% to

total shoot mass, and 10.7% to shoot $A_{gross}$ ($\mu$mol $CO_2$ $s^{-1}$; Table 1). Mean stem $A_{gross}$ $m^{-2}$ was

nearly double leaf $A_{gross}$ $m^{-2}$ (9.62 $\mu$mol $CO_2$ $m^{-2}$ $s^{-1}$ and 5.13 $\mu$mol $CO_2$ $m^{-2}$ $s^{-1}$, respectively;

Figure 1).

*Light response curves differ between shoots and leaves*

Leaves and shoots had distinct light response curves with differences in all parameters

tested, except for quantum yield (Table 2). Leaf level $R_d$ was more than 3x that of the shoot level

$R_d$, resulting in a correspondingly higher LCP, despite the less acute rise in $A_{net}$ response to

PPFD (Figure 2).  Leaf level 75% saturation was achieved at approximately 528 $\mu$mol photons

$m^{-2}$ $s^{-1}$, while that of shoots was estimated at 1532  $\mu$mol $m^{-2}$ $s^{-1}$. The corresponding $A_{sat}$ was also

approximately 20% higher in shoots than in leaves.

*Photosynthesis and starch are affected by CLas exposure*

Uninfected leaves had a higher $A_{net}$ (mean of 3.97 $\mu$mol $CO_2$ $m^{-2}$ $s^{-1}$) than infected leaves;

however, as $C$Las exposure increased from low, to medium, to high so did $A_{net}$ (means of 1.91,

2.61, 2.70 $\mu$mol $CO_2$ $m^{-2}$ $s^{-1}$, respectively; Table 3). Uninfected leaves had a lower starch content

(mean of 1.08 $\mu$g $mm^{-2}$) than infected leaves; however, as $C$Las exposure increased from low to

high starch content decreased (means of 11.31, 5.69, 3.55 $\mu$g $mm^{-2}$, respectively). Uninfected

leaves had the lowest visual rating score, which increased as $C$Las exposure increased from a

mean of 0.22 in the uninfected to a mean of 2.22 in the highest exposure group.

$A_{net}$ had a positive relationship with $C$Las exposure at the shoot-level. High exposure

shoots had the highest $A_{net}$ at a mean of 3.37 $\mu$mol $m^{-2}$ $s^{-1}$, which decreased with a shortened

exposure down to a mean of 1.91 μmol m$^{-2}$ s$^{-1}$ in the uninfected shoots (Table 4). Reductions in

mean leaf area (17.59 cm$^2$ in the uninfected to 11.38 cm$^2$ in the highest exposed group) and total

shoot surface area (133.5 cm$^2$ in the uninfected to 68.3 cm$^2$ in the highest exposed group) were

found as $C$Las exposure increased.  Exposure did not affect internode length.

*Infection, starch, photosynthesis, and morphological variables were correlated*

At the leaf level, when including uninfected trees, positive correlations were found

between bacterial copy count and starch content ($r$=0.609, $P$<0.00001), $C$Las copy number and

visual rating (r=0.459, $P$<0.00001), and starch content and visual rating ($r$ =0.276, $P$=0.005;

Appendix S6). Negative correlations were found between $A_{net}$ and $C$Las copy number ($r$ =-0.322,

$P$=0.0009) as well as starch content ($r$= -0.243, $P$=0.013). No correlations were found with leaf

position or between $A_{net}$ and visual rating. No additional correlations were found when excluding

uninfected trees.

At the shoot level, when including uninfected trees, positive correlations were found

between $C$Las copy number and starch content ($r$=0.716, $P$<0.00001), visual rating ($r$ =0.539,

$P$=0.001), and $A_{net}$ ($r$=0.413, $P$=0.015); between shoot surface area and mean leaf area ($r$=0.812,

$P$<0.00001) and shoot surface area and shoot length (r=0.58, $P$=0.0003); and between internode

length and shoot length (r=0.344, $P$=0.046; Appendix S8). When excluding uninfected trees at

the shoot level, additional positive correlations were found between starch content and mean leaf

area (r=0.421, $P$=0.032) and shoot surface area (r=0.455, $P$=0.019); between shoot surface area

and mean leaf area (r=0.809, $P$<0.00001) and shoot length (r=0.644, $P$=0.0004); and between

internode length and $A_{net}$ (r=0.558, $P$=0.003; Appendix S9).

*Linear regressions reveal additional relationships between infection and physiology*

Based on the correlations we tested linear regressions of the following variables: $C$Las

copy number by starch content, internode length by $A_{net}$, and foliar starch content by shoot

surface area. After a natural log transformation of both starch content and bacterial copy count, a

positive linear relationship was found at both the leaf and shoot-level (Table 5). Further

observation indicated a change in the slope as $C$Las gene copy # increased and assessment using

breakpoint analysis indicated that the increased starch content at the leaf level increased at a

higher rate with a breakpoint at approximately $10^{11.2}$ $C$Las gene copies (Figure 3A), though the

shoot level data did not indicate such a change in slope (Figure 3B). A positive linear

relationship ($P$=0.048) was found between foliar starch with shoot surface area when excluding

uninfected shoots, though this relationship also had a low $R^2$ (0.18) (Figure 4 and Table 5). This

relationship is inverse to the $C$Las exposure duration, with the highly exposed shoots having

lower starch content and total shoot surface area. When comparing morphological and $A$

measurements, a positive linear relationship ($P$=0.003) was found between internode length and

$A_{net}$ when excluding uninfected shoots (Figure 5 and Table 5).

**Discussion**

*Leaf and shoot light response dynamics differ*

The distinct shoot light response dynamic helps to explain the success of *Citrus* in high

light regions, where maximum $A_{sat}$ is attained near the maximum ambient PPFD occurring in

subtropical regions. The leaf-level light response curves indicating $Q_{sat}$ occurs between 600-700

$\mu$mol m$^{-2}$ s$^{-1}$ PPFD was similar to earlier studies (Habermann et al., 2003). The increased light

saturation point of citrus shoots compared to individual leaves could be explained by varied

angle of incident radiation, as well as by partial shading of basal leaves by more distal leaves.

However, $R_d$ was much lower at the shoot level than at the leaf level. Lower $R_d$ is an important

factor in reducing the LCP. Thus, at the shoot level citrus exhibits higher $A_{net}$ at both the lower

and upper ends of irradiance than is detected at the leaf level, while at PPFD ranges between 50-

1000 µmol m$^{-2}$ s$^{-1}$ shoots exhibit lower $A_{net}$ than leaves. The lack of a difference in Φ is further

confirmation that differences between shoots and leaves are primarily determined by factors such

as morphology, not by enzymatic or photochemical factors. This might be assumed given that

some of the same leaves were involved in both measurements, though the lower rate of $R_d$ in

shoots merits further investigation. The fact that much of the difference between shoot and leaf

irradiance responses may be attributed to the irradiance variance within the shoot indicates that

morphology is likely to have a strong impact on photosynthesis at the whole shoot level.

*Stem assimilation contributes to whole shoot photosynthesis*

Citrus vegetative stems significantly contributed to shoot $A_{gross}$. The stem was also more

efficient in $A_{gross}$ on a surface area basis than leaves. The high assimilation rate of citrus stems

could be explained by area-related chlorophyll content where young stems contain 50–70% more

chlorophyll than adjacent leaves (Kharouk et al. 1995; Pfanz et al., 2002; Pilarski 1984; Schmidt

et al. 2000; Solhaug et al. 1995). Although many factors contribute to whole-shoot assimilation,

the previously unquantified contribution of the photosynthetic stem to canopy wide assimilation

may be an additional factor in the lack of relationship between leaf-level gas exchange

measurements to canopy-wide measurements. Stem internal re-fixation of $CO_2$ in young

branches of perennial woody species have been shown to compensate for 60–90% of the

potential respiratory carbon loss (Pfanz et al., 2002). Although, its contributions compared to the

leaves on whole shoot assimilation are relatively small, stem contribution may vary depending

on shoot morphological characteristics such as internode length, which may affect light

interception in the stem.

*HLB impacts shoot morphology and photosynthesis*

As with light responses, shoots and individual leaves differed in their responses to HLB.

While $C$Las infection reduced $A_{net}$ at the leaf level, it increased $A_{net}$ at the whole shoot level,

while inducing whole-shoot morphological adjustments over time. These morphological impacts

explain the HLB effects on whole-shoot photosynthetic rates. The decreased leaf and shoot

surface area effected by $C$Las exposure reflect the symptom of canopy thinning observed in

infected citrus trees in the field. This process appears to occur on a scale of years. Given that

most citrus leaves senesce by 17 months (Wallace et al. 1954), the changes in shoot morphology

appear to occur after a complete turnover of old to new tissues.

We have not found any report in which a plant produced both longer internodes and

smaller leaves in response to a disease or environmental condition. Typically, an increase in leaf

size has coincided with increased internode length to reduce self-shading, such as in acclimation

to low light conditions (Kavanagh, 2014; Pieters et al., 1998; Poorter and Rozendaal, 2008;

Robertson, 1994), thus the particular morphological changes induced by HLB may be responses

to the specific challenges imposed by the disease.

Shoot morphological changes were associated with the shoot-level gas exchange response

to HLB. Light penetration and absorption are the primary drivers of photosynthesis, and these are

heavily influenced by canopy structure (Niinemets, 2007). Canopy thinning by $C$Las-induced

defoliation, as described by de Graça (1991), within the high $C$Las-exposed category (fewest and

smallest leaves and longest internode length) further facilitate light penetration into the canopy.

With increased light penetration and internode length, typically shaded leaves, further from the

shoot apex, and the photosynthetic stem contribute more to shoot-level assimilation, which help

explain the increased assimilation efficiency found in shoots of the high $C$Las exposure group.

Considering the contribution of stems to shoot $A_{gross}$, the relative increase in stem surface area,

could also contribute to the enhanced shoot $A_{net}$. The increased light exposure of all parts of the

shoot may also lead to compensation for the localized disease effects discussed below in

reference to leaves. Although there are likely other factors that contribute to the increased

assimilation efficiency of HLB-affected shoots, the correlation of these morphological variables

with $A_{net}$ indicates that whole-shoot morphology plays a sufficiently important role in citrus

shoot photosynthesis as to be a defining feature of shoot acclimation that leads to an increase in

$A_{net}$.

The increased efficiency of $A_{net}$ should not be confused with increased total

photosynthesis. While the highly $C$Las-exposed shoots had higher $A_{net}$ than the shoots from the

low exposure shoots, these measures are surface-area based and do not offset carbon fixation lost

in the reduction in shoot surface area with increased $C$Las exposure. Total net shoot assimilation

was 40% higher for the low $C$Las-exposed shoots (0.032 μmol $CO_2$ s$^{-1}$) than high $C$Las-exposed

shoots (0.023 μmol $CO_2$ s$^{-1}$). Although, growth is necessarily dependent on photosynthesis,

numerous complexities have led to a low degree of correlation between net assimilation and

growth.  For example, Poorter and Remkes (1990) did not find net assimilation rate to be

correlated with relative growth rate in a comparison of numerous species, though other multi-

species comparisons have found effects of net assimilation rate on relative growth rate (Li et al.

2016).  Measurements of $A_{net}$ using gas exchange are further complicated in that they are limited

in both spatial (canopy dimensions) and time scales, and neither growth-analysis-based net

assimilation rate nor gas-exchange-based $A_{net}$ include the complexities of allocation and their

subsequent impacts on growth (Pereira 1995). In most studies, however, total photosynthetic area

has been found to affect growth more than photosynthetic activity on a leaf area basis (Pereira

1995, Sinclair et al. 2019). The present study provides an example of how morphological

changes could overwhelm changes in $A_{net}$. Despite increases in photosynthetic efficiency, $C$Las

infected shoots had lower total assimilation rates because the reduction in photosynthetic surface

area was greater than the relative increase in $A_{net}$, leading to a reduction in assimilation per shoot.

Thus citrus plants mitigate losses by increasing photosynthesis per leaf area but do not increase

total assimilation.

*CLas infection induces local starch and photosynthetic responses*

420            Because $C$Las is a phloem-limited bacterium, the effects of infections on leaf

carbohydrate dynamics have been the focus of much HLB-related research. Achor et al. (2010)

observed starch disruption of chloroplast inner grana in association with local phloem blockage

and collapse. Further, Fan et al. (2010) found increased foliar sucrose concentrations in

association with starch accumulation in infected trees. The current study suggests that these

effects are localized, where $C$Las population increases induced starch accumulation at a much

higher rate after reaching a population threshold at the leaf level.

427            Given these earlier findings regarding the impact of HLB on citrus carbohydrate

relations, Cimò et al. (2013) simulated the foliar response of HLB by girdling citrus trunks,

resulting in starch accumulation and a reduction in leaf level $A_{net}$. This reduction had been

hypothesized because of the source-sink hypothesis (Ainsworth and Bush 2011), which observes

that plants downregulate $A_{net}$ in response to reduced leaf carbohydrate export, a condition in

which various carbohydrates may accumulate. In the case of citrus, this hypothesis was

demonstrated by Syvertsen (1994), and later by Iglesias (2002) and Nebauer et al. (2011), who

found $A_{net}$ responded to changes in soluble sugars, which in turn also resulted in starch

accumulation or depletion. Although none of the above studies addressed the condition of HLB,

their interpretation of reduced photosynthesis as a response to leaf export limitation has been

applied to studies of HLB, in which gene expression of photosynthetic pathways has been found

to be down regulated (Balan et al. 2018). The present study found the expected reduction of $A_{net}$

at the leaf level, but not at the shoot level.

The comparison of local versus systemic effects has not been addressed explicitly by

studies of HLB. However, the authors of some studies have implicitly interpreted systemic

effects on leaves (all studies used in Balan et al. 2018) while others have hypothesized that HLB

primarily affects roots (Johnson et al. 2014). The indication in the present study of a localized

impact of CLas on leaf carbohydrate export and photosynthesis supports the interpretation of

Koh et al. (2012), who observed callose blockage of leaf phloem as well as reduced export of

[14]C-labeled photosynthates in infected trees, while also agreeing with those of Fan et al. (2010),

Rao et al. (2018), and Balan et al. (2018), all of which found alterations in carbohydrate

metabolism and photosynthesis in leaves. A unique contribution of the present study is to

indicate that these effects are at least partly local effects of presence of the bacterial population in

the leaf, rather than strictly systemic responses to infection.

Starch accumulation has been associated with CLas infection and has been attributed as

the cause of the blotchy mottle and chlorosis symptoms seen in infected citrus leaf tissue

(Etxeberria et al., 2009; Gonzalez et al., 2012; and Schaffer et al., 1986). Interestingly, we found

high starch accumulation in the most recently exposed group and a relative reduction in starch

content with longer exposure durations. Meanwhile, visual estimates of symptom severity

increased across exposure groups, indicating that more severe symptoms were not the result of

greater starch accumulation.

*Conclusion*

This study provides the first direct comparison of $CO_2$ assimilation between leaf and

shoot levels, as well as the first comparison of assimilation in HLB-affected and healthy citrus.

Shoot- and leaf-level responses were distinct. Shoot light response curves were much more

gradual than those of leaves with $A_{max}$ at approximately 2x the irradiance of the $A_{max}$ of leaves.

This was explained by the combination of leaf angles and photosynthetic stems, which

contributed a mean of 10% of the total shoot $A_{gross}$. Likewise shoot and leaf responses to $C$Las

infection differed. Where $A_{net}$ was reduced at the leaf level, shoot morphological acclimation

partially compensated for this downregulation in leaf-level photosynthetic activity. These

morphological changes further confirm the importance of canopy density as an important

measure of disease severity. This study is an example of the value of assessing physiological

processes along temporal and spatial scales, with whole-shoot photosynthesis providing one tool

for such efforts.

**Supplemental materials**

Appendix S1.  Cap design of whole shoot assimilation chamber. A) bottom view – stepped

drilled hole to limit low leaf interference, B) top view – straight cut to allow for stem placement,

C) fit of cap into chamber, D) showing fit into chamber to prevent leakage.

478 Appendix S2. Straw pieces fit to cover stem of 'Valencia' sweet orange (*Citrus x sinensis* [L.]

479 Osbeck) shoot to measure whole shoot gas exchange excluding stem photosynthesis.

481 Appendix S3.  Diagram describing the measurements taken to estimate stem surface area.

483 Appendix S4. Standard deviation of $A_{net}$ at photosynthetic photon flux density of 1000 μmols m$^{-2}$

484 s$^{-1}$ demonstrating measurement stabilization for 'Valencia' sweet orange (*Citrus x sinensis* [L.]

485 Osbeck) in leaf chamber of whole shoot chamber. Dashed vertical line indicates stabilization

486 point for leaf; dotted line indicates stabilization point for shoots.

488 Appendix S5. Illustration of 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) shoots and

489 leaves selected for to compare assimilation in leaves vs. shoots.

491 Appendix S6. Correlation of leaf disease and photosynthetic variables of healthy and *Candidatus*

492 Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) exposed

493 to 0, 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri* Kuwayama)

494 feeding and reproduction. Copy # - log transformed bacterial copy count, Starch – log

495 transformed starch concentration (μg mm$^{-2}$), Visual – visual  rating  of citrus greening leaf

496 chlorosis and mottling symptoms, $A$ – net $CO_2$ assimilation (μmol m$^{-2}$ s$^{-1}$), and Position – leaf

497 position on stem (1-n starting at apical leaf ). Darkened correlation boxes denote correlations

498 where P≤0.05

Appendix S7.  Correlation of leaf disease and photosynthetic variables of *Candidatus*

Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) exposed

to 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri* Kuwayama)

feeding and reproduction. Copy # - log transformed bacterial copy count, Starch – log

transformed starch concentration ($\mu$g mm$^{-2}$), Visual – visual  rating  of citrus greening leaf

chlorosis and mottling symptoms, *A* – net $CO_2$ assimilation ($\mu$mol m$^{-2}$ s$^{-1}$), and Position – leaf

position on stem (1-n starting at apical leaf ). Darkened correlation boxes denote correlations

where P≤0.05.assimilation ($\mu$mol $CO_2$ m$^{-2}$ s$^{-1}$), and Position – leaf position on stem (1-n starting

at apical leaf ). Darkened correlation boxes denote correlations where P≤0.05.

Appendix S8.  Correlation of shoot disease and photosynthetic variables of healthy and

*Candidatus* Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) exposed to 0, 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina*

*citri* Kuwayama) feeding and reproduction. Copy # - log transformed bacterial copy count,

Starch – log transformed starch concentration ($\mu$g mm$^{-2}$), Visual – visual  rating  of citrus

greening leaf chlorosis and mottling symptoms, *A* – net $CO_2$ assimilation ($\mu$mol m$^{-2}$ s$^{-1}$), LA –

mean leaf surface area (cm$^2$), SA – total shoot surface area (cm$^2$), Length – stem length (mm),

and Inter – average internode length (mm). Darkened correlation boxes denote correlations

where P≤0.05.

Appendix S9.  Correlation of shoot disease and photosynthetic variables of healthy and

*Candidatus* Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) exposed to 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri*

Kuwayama) feeding and reproduction. Copy # - log transformed bacterial copy count, Starch –

log transformed starch concentration ($\mu g\ mm^{-2}$), Visual – visual  rating  of citrus greening leaf

chlorosis and mottling symptoms, $A$ – net $CO_2$ assimilation ($\mu mol\ m^{-2}\ s^{-1}$), LA – mean leaf

surface area ($cm^2$), SA – total shoot surface area ($cm^2$), Length – stem length (mm), and Inter –

average internode length (mm). Darkened correlation boxes denote correlations where $P \leq 0.05$.

*Conflict of Interest*

The authors declare they have not conflict of interest with this study, including no stake in any

product described in the manuscript.

*Funding*

Funding for this project comes from the United States Department of Agriculture Multi-Agency

Coordination System (AP19PPQS&T00C165).

*Acknowledgements*

The authors wish to thank Myrtho Pierre for assistance in gas exchange measurements, Rebecca

Ebert for assistance with starch measurements, and Jason Hupp for providing the design for the

adapter with which the whole-shoot chamber was developed.

*Authors' contributions*

MK, CV, and DR conceived experiments.  MK developed whole-shoot chamber and conducted

experiments.  MK analyzed data and wrote manuscript.  CV supervised experiments. CV and DR

supervised analyses and manuscript writing, and revised manuscript.

**References**

Achor D, Welker S, Ben-Mahmoud S, Wand C, Folimonova SY, Dutt M, Gowda S, Levy A

(2020) Dynamics of Candidatus Liberibacter asiaticus Movement and Seive-Pore Plugging in

Citrus Sink Cells. Plant Physiology.182(20):882-891.

Ainsworth EA, Bush DR (2011) Carbohydrate Export from the Leaf: A Highly Regulated

Process and Target to Enhance Photosynthesis and Productivity. Plant Physiology 155:64–

69.

Balan B, Ibáñez AM, Dandekar AM, Caruso T, Martinelli F (2018) Identifying host molecular

features strongly linked with responses to huanglongbing disease in citrus leaves. Frontiers in

plant science 9:277.

Bové JM (2006) Huanglongbing: A Destructive, Newly-Emerging, Century-Old Disease of

Citrus. Journal of Plant Pathology 88(1):7-37.

Cen H, Weng H, Yao J, He M, Lv J, Hua S, Li H, He Y (2017) Chlorophyll Fluorescence

Imaging Uncovers Photosynthetic Fingerprint of Citrus Huanglongbing. Frontiers in Plant

Science 8:1509.

Cimò G, Bianco RL, Gonzalez P, Bandaranayake W, Etxeberria E, Syvertsen JP (2013)

Carbohydrate and Nutritional Responses to Stem Girdling and Drought Stress with Respect

to Understanding Symptoms of Huanglongbing in Citrus. HortScience 48:920–928.

Contreras S, Hunink JE, Baille A (2017) Water and carbon fluxes in irrigated citrus orchards

assessed from satellite data. FutureWater Report 174.

Coy MR, Hoffman M, Gibbard KHN, Kuhns EH, Pelz-Stelinski KS, Stelinski LL (2014) Nested-

quantitative PCR approach with improved sensitivity for the detection of low titer levels of

Candidatus Liberibacter asiaticus in the Asian citrus psyllid , Diaphorina citri Kuwayama.

Journal of Microbiological Methods 102:15-22.

da Graça JV (1991) Citrus Greening Disease. Annual Review Phytopathology 29:109-136.

Das AK, Nerkar S, Bawage S, Kumar A (2013) Current Distribution of Huanglongbing (citrus

greening disease) in India as Diagnosed by Real-Time PCR. Journal of Phytopathology

162(6):402-406.

de Mendiburu F (2020) agricolae: Statistical Procedures for Agricultural Research. R package

version 1.3-2. https://CRAN.R-project.org/package=agricolae

Er-Raki S, Chehbouni A, Guemouria N, Ezzahar J, Khabba S, Boulet G, Hanich L (2009) Citrus

orchard evapotranspiration: Comparison between eddy covariance measurements and the

FAO 56 approach estimates. Plant Biosystems 143(1):201-208.

Etxeberria E, Gonzalez P, Achor D, Albrigo G (2009) Anatomical distribution of abnormally

high levels of starch in HLB-affected Valencia orange trees. Physiological and Molecular

Plant Pathology 74(1):76-83.

Etxeberria E, Gonzalez P, Dawson W, Spann T (2007) An iodine-based starch test to assist in

selecting leaves for HLB testing. University of Florida, IFAS Extension, Gainesville,

HS1122.

Eyles A, Pinkard EA, O'Grady AP, Worledge D, Warren CR (2009) Role of corticular

photosynthesis following defoliation in Eucalyptus globulus. Plant, Cell and Environment.

32(8):1004-1014.

Falster D, Westoby M (2003) Leaf size and angle vary widely across species: What

consequences for light interception? New Phytologist 158(3):509-525.

Fan J, Chen C, Brlansky RH, Gmitter Jr. FG, Li Z-G (2010) Changes in carbohydrate

metabolism in Citrus sinensis infected with 'Candidatus Liberibacter asiaticus'. Plant

Pathology 59:1037–1043.

Folimonova SY, Achor DS (2010) Early events of citrus greening (Huanglongbing) disease

development at the ultrastructural level. Phytopathlogy 100(9):949-958.

Gonzalez P, Reyes J, Etxeberria E (2012) Characterization of leaf starch from HLB-affected and

unaffected-girdled citrus trees. Physiological and Molecular Plant Pathology 79:71-78.

Habermann G, Machado EC, Rodrigues JD, Medina CL (2003) CO2 assimilation, photosynthetic

light response curves, and water relations of 'Pera' sweet orange plants infected with Xylella

fastidiosa. Brazilian Journal of Plant Physiology 15(2):79-87.

Harrell FE, Jr (2020) Hmisc: Harrell Miscellaneous. R package version 4.4-0. http://CRAN.R-

project.org/package=Hmisc

Iglesias DJ, Lliso I, Tadeo FR, Talon M (2002) Regulation of photosynthesis through source:

sink imbalance in citrus is mediated by carbohydrate content in leaves. Physiologia

Plantarum 116(4):563-572.

Jifon JL, Syvertson JP (2003) Moderate shade can increase net gas exchange and reduce

photoinhibition in citrus leaves. Tree Physiology 23(2):119-127.

Johnson EG, Wu J, Bright DB, Graham JH (2014) Association of 'Candidatus liberibacter

asiaticus' Root Infection, but not Phloem Plugging with Root Loss on

Huanglongbing-affected Trees Prior to Appearance of Foliar Symptoms. Plant Pathology

63(2):290-298.

Kavanagh PH (2014) Herbivory and the evolution of divaricate plants: Structural defences lost

on an offshore island. Austral Ecology 40(2):206-211.

Kharouk VI, Middleton EM, Spencer SL, Rock BN, Williams DL (1995) Aspen bark

photosynthesis and its significance to remote sensing and carbon budget estimate in the

boreal ecosystem.  Water, Air, and Soil Pollution 82:483-497.

Koh E-J, Zhou L, Williams DS, Park J, Ding N, Duan Y-P, Kang B-H (2012) Callose deposition

in the phloem plasmodesmata and inhibition of phloem transport in citrus leaves infected

with "Candidatus Liberibacter asiaticus". Protoplasma 249:687–697.

Kriedemann PE (1968) Some Photosynthetic Characteristics of Citrus Leaves. Australian Journal

of Biological Science 21:895-905.

Maestre-Valero JF, Testi L, Jiménez-Bello MA, Castel JR, Intrigliolo DS (2017)

Evaportranspiration and carbon exchange in a citrus orchard using eddy covariance.

Irrigation Science 35:397-408.

Mesquita GL, Zambrosi FCB, Tanaka FAO, Boaretto RM, Quaggio JA, Ribeiro RV, Mattos D

(2016) Anatomical and Physiological Response of Citrus Trees to Varying Boron

Availability Are Dependent on Rootstock. Frontiers of Plant Science. 7:224.

Nebauer SG, Renau-Morata B, Guardiola JL, Molina R (2011) Photosynthesis down-regulation

precedes carbohydrate accumulation under sink limitation in Citrus. Tree Physiology 31:169-

177.

Niinemets U (2007) Photosynthesis and resource distribution through plant canopies. Plant, Cell

and Environment 30(9):1052-1071.

Nilsen ET, Bao Y (1990) The Influence of Water Stress on Stem and Leaf Photosynthesis in

Glycine Max and Sparteum Junceum (Leguminosae). American Journal of Botany

77(8):1007-1015.

Pfanz H, Aschan G, Langenfeld-Heyser R, Wittman C, Loose M (2002) Ecology and

ecophysiology of tree stems: corticular and wood photosynthesis. Naturwissenschaften

89:147-162.

Pieters GA, Van Den Noort ME, Nijkerken V (1998) Growth adaptation of leaves and internodes

of popular to irradiance, day length and temperature. Tree Physiology 19:933-942.

Pinheiro JC, Bates D, DebRoy S, Sarkar D, R Core Team (2014) nlme: Linear and Nonlinear

Mised Effects Models. R package version 3,1-122, http://CRAN,R-

project.org/package=nlme.

Poorter L, Rozendall DMA (2008) Leaf size and leaf display of thirty-eight tropical tree species.

Physiological Ecology 158:35-46.

R Core Team (2019). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

URL https://www.R-project.org/.

Rao, M. J., Ding, F., Wang, N., Deng, X., & Xu, Q. (2018). Metabolic mechanisms of host

species against citrus Huanglongbing (Greening Disease). *Critical Reviews in Plant*

*Sciences*, *37*(6), 496-511.

Rasowo BA, Khamis FM, Mohamed SA, Ajene IJ, Aidoo OF, Ombura L, Setamou M, Ekesi S,

Borgemeister C (2019) African Citrus Greening Disease in East Africa: Incidence, Severity,

and Distribution Patterns. Journal of Economic Enotomology 112(5):2389-2397.

Robertson MJ (1994) Relationships between internode elongation, plant height and leaf

appearance in maize. Field Crop Research 38(3):135-145.

Sagaram M, Burns JK (2009) Leaf chlorophyll fluorescence parameters and Huanglongbing. J

Am Soc Hortic Sci 134:194–201.

Schaffer AA, Liu K, Goldschmidt EE, Boyer CD, Goren R (1986) Citrus Leaf Chlorosis Induced

by Sink Removal: Starch, Nitrogen, and Chloroplast Ultrastructure. Journal of Plant

Physiology 124 (1-2):111-121.

Schmidt J, Batic F, Pfanz H (2000) Photosynthetic performance of leaves and twigs of evergreen

holly (Ilex aquifolium L.) Phyton 40:179-190.

Schneider H (1968) Anatomy of greening diseased sweet orange shoots. Phytopathology

58:1155-1160.

Singerman A, Usech P (2016) Impact of Citrus Greening on Citrus Operations in Florida.

University of Florida IFAS Extension #FE-983.

Slinski S (2016) Evaluation of HLB treatments. Citrus Industry August 2016:10-12.

Snyder-Leiby TE, Wang S (2008) Role of crop load in chloroplast ultrastructure and zonal

chlorosis, a physiological disorder in 'Honeycrisp' apple trees. Horticultural Science

43(6):1819-1822.

Solhaug KA, Gauslaa Y, Haugen J (1995) Adverse effect of epiphytic crustose lichens upon stem

photosynthesis and chlorophyll of Populus tremula L. Botanica Acta 108:233-239.

Syvertsen JP (1994) Partial shoot removal increases net CO2 assimilation and alters water

relations of Citrus seedlings. Tree Physiology 14:497–508.

Tang L, Chhajed S, Vashisth T (2019) Preharvest Fruit Drop in Huanglongbing-affected

'Valencia' Sweet Orange. J. Amer. Soc. Hort. Sci. 144(2):107-117.

Tinoco-Ojanguren C, Pearcy RW (1992) Dynamic stomatal behavior and its role in carbon gain

during lightflecks of a gap phase and an understory Piper species acclimated to high and low

light. Oecologia 92:222–228.

Tinoco-Ojanguren C, Pearcy RW (1993) Stomatal dynamics and its importance to carbon gain in

two rainforest Piper species I. VPD effects on the transient stomatal response to lightflecks.

Oecologia 94:388–394.

Tomeo, N. (2017). AQ curves example. GitHub Repository.

https://github.com/Tomeopaste/AQ_curves/blob/master/Introduction_to_AQ_curve_fitting.R

md

Tsagarakis AE, Rogers ME, Spann TM (2012) Applications of Plant Growth Regulators to

Container-grown Citrus Trees Affect the Biology and Behavior of the Asian Citrus Psyllid.

Journal of the American Society of Horticultural Science 137(1):3-10.

Vu JCV, Yelenosky G (1988) Biomass and Photosynthesis Response of Rough Lemon (Citrus

jambhiri Lush.) to Growth Regulators. Canadian Journal of Plant Sciences 68:261-266.

Vu JCV, Yelenosky G, Bausher MG (1986) $CO_2$ Exchange Rate, Stomatal Conductance, and

Transpiration in Attached Leaves of Valencia Orange. HortScience 21:143-144.

Wallace A, Zidan ZI, Mueller RT, North CN (1954) Translocation of nitrogen in citrus trees.

Proceedings of American Society of Horticultural Science 64:87-104.

Whitaker DC, Giurcanu MC, Young LJ, Gonzalez P, Etxeberria E, Roberts P, Hendricks K,

Roman F (2014) Starch Content of Citrus Leaves Permits Diagnosis of Huanglongbing in the

Warm Season but Not Cool Season. HortScience 49(6):757-762.

Yanhong T, Hiroshi K, Mitsumasa S, Izumi W (1994) Characteristics of transient photosynthesis

in Quercus serrata seedlings grown under lightfleck and constant light regimes. Oecologia

100:463–469.

Zambrosi FCB, Mattos D, Syvertsen JP (2011) Plant Growth, leaf photosynthesis, and nutrient-

use efficiency of citrus rootstocks decreased with phosphite supply. Journal of Plant

Nutrition and Soil Science 174(3):487-495.

Zheng Z, Chen J, Deng X (2018) Historical perspectives, management and current research of

citrus HLB in Guangdong of China, where the disease has been endemic for over a hundred

years. Phytopathology 108(11):1224-1236.

**List of figures**

Figure 1.  Gross assimilation for 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) leaf,

stem and shoot. Letters within each plot denote means that are significantly different at $P \leq 0.05$

according to Fisher's least significant differences.

Figure 2.  Distinct carbon assimilation light response curves of 'Valencia' sweet orange (*Citrus x*

*sinensis* [L.] Osbeck) leaf (dashed line) and shoot (dotted line).

Figure 3.  Relationship of log transformed bacterial copy count and log transformed starch

content of *Candidatus* Liberibacter asiaticus-infected and uninfected 'Valencia' sweet orange

(*Citrus x sinensis* [L.] Osbeck) A) leaves and B) shoots.

Figure 4.  Relationship of starch and shoot surface area of *Candidatus* Liberibacter asiaticus-

infected and uninfected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck). Data point

shape and color denote *C*Las-infection exposure category: Low – 1 year *C*Las-infected Asian

citrus psyllid feeding and breeding, Mid – 1.5 years *C*Las-infected Asian citrus psyllid feeding

and breeding, High – 2.5 years *C*Las-infected Asian citrus psyllid feeding and breeding.

Figure 5.  Relationship of average internode length and net $CO_2$ assimilation (*A*) of *Candidatus*

Liberibacter asiaticus-infected and uninfected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck).

**Tables**

Table 1. Percent contribution of leaves and stems to the shoot surface area and net $CO_2$

assimilation ($A_{net}$) in 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck).

| Measurement basis | Leaf | Stem |
|---|---|---|
| Surface area | $94.3 \pm 1.6$ % | $5.6 \pm 1.6$ % |
| Mass | $83.0 \pm 4.8$% | $17.0 \pm 4.8$% |
| Per shoot $A_{net}$ | $89.2 \pm 8.5$ % | $10.7 \pm 8.5$% |

Table 2. Analysis of variance and means comparison of parameters of a light response curve

function fit to leaf and whole shoot light response curves of 'Valencia' sweet orange (*Citrus x*

*sinensis* [L.] Osbeck). Values are means ± standard error. Values followed by different letters are

significantly different at P<0.05. $A_{sat}$: light-saturated net $CO_2$ assimilation (µmol $CO_2$ m$^{-2}$ s$^{-1}$);

Φ:quantum yield (mols $CO_2$ fixed:mol photons absorbed) ; $R_d$: dark respiration (µmol $CO_2$ m$^{-2}$ s$^{-}$

$^{1}$); LCP: light compensation point (µmol photosynthetically active radiation m$^{-2}$ s$^{-1}$); Qsat75:

photosynthetic flux density at 75% saturation (µmol photosynthetically active radiation m$^{-2}$ s$^{-1}$).

| Function parameter | Leaf | Shoot | F-value | *P*-value |
|---|---|---|---|---|
| $A_{sat}$ | 7.7 ± 0.46 b | 10.4 ± 0.74 a | 8.7 | 0.01 |
| Φ | 0.044 ± 0.002 | 0.053 ± 0.02 | 0.2 | 0.65 |
| $R_d$ | 1.2 ± 0.12 a | 0.35 ± 0.04 b | 42.4 | <0.0001 |
| LCP | 28.0 ± 8.4 a | 14.7 ± 10.8 b | 7.9 | 0.01 |
| $Q_{sat75}$ | 528 ± 80 b | 1532 ± 226 a | 17.3 | 0.0016 |

Table 3.  Effect of huanglongbing symptom category on  leaf-level net $CO_2$ assimilation ($A_{net}$),

cycle threshold (Ct)  and Copy # count of *Candidatus* Liberibacter asiaticus, foliar starch

content, and visual disease rating in 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck). *P*-

value of analysis of variance with infection categories as the fixed effect. Means (± standard

error) followed by different letters are significantly different at *P*<0.05 according to Fisher's

LSD.

|  | Uninfected | Low | Medium | High | *P*-value |
|---|---|---|---|---|---|
| $A_{net}$ ($\mu$mol m$^{-2}$ s$^{-1}$) | 3.97 ± 1.73 a | 1.91 ± 1.24 b | 2.61 ± 1.62 b | 2.70 ± 1.75 b | 0.00027 |
| qPCR Ct | 37.87 ± 2.35 a | 24.69 ± 4.14 b | 24.16 ± 3.13 b | 24.10 ± 2.52 b | <0.00001 |
| qPCR Copy # (log) | 2.60 ± 2.65 b | 11.78 ± 2.90 a | 12.27 ± 1.93 a | 12.30 ± 1.66 a | <0.00001 |
| Starch ($\mu$g mm$^{-2}$) | 1.08 ± 1.27 d | 11.31 ± 4.30 a | 5.69 ± 4.39 b | 3.55 ± 3.17 c | <0.00001 |
| Visual Rating (0-10) | 0.22 ± 0.60 c | 1.26 ± 0.98 b | 2.11 ± 1.53 a | 2.22 ± 1.42 a | <0.00001 |

758    Table 4. Effect of huanglongbing symptom category on shoot-level net $CO_2$ assimilation ($A_{net}$)

759    and shoot morphological variables. *P*-value of analysis of variance with infection categories as

760    the fixed effect 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck). Means ($\pm$ standard

761    error) followed by different letters are significantly different at *P<0.05* according to Fisher's

762    LSD.

| | Uninfected | Low | Medium | High | P-value |
|---|---|---|---|---|---|
| $A_{net}$ ($\mu$mol m$^{-2}$ s$^{-1}$) | $1.91 \pm 1.07$ b | $2.42 \pm 0.88$ b | $2.44 \pm 0.52$ ab | $3.37 \pm 1.34$ a | 0.03167 |
| Stem length (mm) | $112.50 \pm 22.5$ | $112.30 \pm 19.7$ | $91.20 \pm 26.1$ | $90.30 \pm 30.6$ | 0.12900 |
| Mean leaf area (cm$^2$) | $15.79 \pm 3.39$ ab | $17.59 \pm 5.86$ a | $15.40 \pm 6.33$ ab | $11.38 \pm 2.08$ b | 0.05469 |
| Shoot surface area (cm$^2$) | $117.40 \pm 42.0$ a | $133.50 \pm 42.8$ a | $100.40 \pm 48.8$ ab | $68.30 \pm 27.1$ b | 0.01398 |
| Leaves per shoot | $6.75 \pm 1.49$ | $7.22 \pm 1.48$ | $6.25 \pm 2.25$ | $5.33 \pm 1.58$ | 0.14980 |
| Internode length (mm) | $17.41 \pm 5.49$ | $15.99 \pm 3.92$ | $15.33 \pm 3.99$ | $16.99 \pm 2.67$ | 0.73790 |

766     Table 5.  Linear regressions of effects of gene copy number of *Candidatus* Liberibacter asiaticus

767     on foliar starch content, of internode length on net $CO_2$ assimilation, and of foliar starch content

768     on shoot surface area each assessed across three huanglongbing symptom categories including or

769     not including a fourth category of uninfected trees, and measured at leaf or whole shoot levels.

770     *P*-values below 0.05 are labeled with asterisks.  "/" denotes nested variables.

| Dependent variable | Including uninfected | Tissue | Effect | F-value | *P*-value |
|---|---|---|---|---|---|
| Starch content | yes | leaf | Bacterial copy number | 52.2130 | <0.00001* |
| | | | Shoot/tree | 8.1480 | 0.0053* |
| | | | Leaf/shoot/tree | 0.4403 | 0.5085 |
| | | shoot | Bacterial copy number | 33.1526 | <0.00001* |
| | | | Shoot/tree | 3.7404 | 0.0623 |
| Net $CO_2$ assimilation | yes | shoot | Internode length | 1.3619 | 0.2521 |
| | | | Shoot/tree | 3.9077 | 0.0570 |
| | no | shoot | Internode length | 10.7940 | 0.0032* |
| | | | Shoot/tree | 0.8399 | 0.3689 |
| Shoot surface area | yes | shoot | Starch content | 1.7849 | 0.1913 |
| | | | Shoot/tree | 0.3502 | 0.5583 |
| | no | shoot | Starch content | 4.3675 | 0.0479* |
| | | | Shoot/tree | 0.7133 | 0.4071 |

**A** $r^2=0.42$

Log Starch (µg mm$^{-2}$)

Log *C*Las Gene Copy #

**B** $y=0.15x+0.28$
$r^2=0.54$

Log *C*Las Gene Copy #

Infection status: ■ High  ▲ Low  ■ Mid

$y = 4.9x + 54.2$
$r^2 = 0.18$

$$y = 0.16x - 0.06$$
$$r^2 = 0.34$$

$A_{tot}$ ($\mu$mol m$^{-2}$ s$^{-1}$)

Internode length (mm)