Title: **Citrus photosynthesis and morphology acclimate to phloem-affecting huanglongbing**

disease at the leaf and shoot levels

Authors: Mark Keeley[1], Diane Rowland[1], and Christopher Vincent[2]*

[1] Agronomy Department, University of Florida, Gainesville, FL, USA

[2] Horticultural Sciences Department, Citrus Research and Education Center, University of

Florida, USA

*To whom correspondence should be directed:

civince@ufl.edu

Citrus Research and Education Center

University of Florida

700 Experiment Station Rd.

Lake Alfred FL 33850

USA

Keywords:

Photosynthesis, Citrus, huanglongbing, gas exchange

Running title:

Citrus photosynthesis acclimates to huanglongbing

Abstract

Huanglongbing (HLB) is a phloem-affecting disease of citrus that reduces growth and has

impacted global citrus production. HLB is caused by a phloem-limited bacterium (*Candidatus*

Liberibacter asiaticus; *C*Las). By inhibiting phloem function, HLB stunts sink growth, including

reducing production of new shoots and leaves, and induces hyperaccumulation of foliar starch.

HLB induces feedback inhibition of photosynthesis by reducing foliar carbohydrate export. In

this work we assessed the relationship of bacterial distribution within the foliage, foliar starch

accumulation, and net CO_2 assimilation (A_{net}). Because HLB impacts canopy morphology, we

developed a chamber to measure whole-shoot A_{net} to test the effects of HLB at both leaf and

shoot levels. Whole-shoot-level A_{net} saturated at high irradiance, and green stems had high

photosynthetic rates compared to leaves. Starch accumulation was correlated with bacterial

population, and starch was negatively correlated with A_{net} at the leaf level but not at the shoot

level. Starch increased initially after infection, then decreased progressively with increasing

length of infection. HLB infection reduced A_{net} at the leaf level, but increased it at the whole

shoot level, in association with reduced leaf size and greater relative contribution of stems to

photosynthetic surface area. Although HLB-increased photosynthetic efficiency, total carbon

fixed per shoot decreased because photosynthetic surface area was reduced. We conclude that the

localized effects of infection on photosynthesis are mitigated by whole shoot morphological

acclimation over time. Stems contribute important proportions of whole shoot A_{net}, and these

contributions are likely increased by the morphological acclimation induced by HLB.

Introduction

Huanglongbing (HLB) is a disease of citrus causing economic loss to the citrus industry

around the world (Bové, 2006; Singerman and Useche, 2016). Its causal agent, *Candidatus*

Liberibacter asiaticus (*C*Las), vectored by the Asian citrus psyllid (ACP; *Diaphorina citri*

Kuwayama), is a phloem-limited bacterium which effects phloem dysfunction leading to starch

accumulation in leaves and loss of sink growth (Achor et al., 2020; Bové, 2006), as well as

eventual development of thin canopies (Tang et al., 2019).

Because of the reduction in phloem function, HLB is hypothesized to lead to feedback

inhibition of photosynthesis (Cimò et al. 2013). Assessing chlorophyll fluorescence, a reduction

in maximal PSII quantum yield was found in *C*Las-infected citrus leaves compared to uninfected

leaves (Cen et al., 2017; Sagaram & Burns, 2009). On the one hand, starch accumulation in the

leaves has been noted as a symptom of HLB, which indicates a surplus of fixation relative to

export (Etxeberria et al., 2009; Folimonova & Achor, 2010; Schneider, 1968). Starch

accumulation can disintegrate the chloroplast membrane (Nebauer et al., 2011; Snyder-Leiby &

Wang, 2008), which may contribute to the reduction in PSII yield noted above. However, no

measures of the impacts of HLB on net CO_2 assimilation (A_{net}) have been reported.

Measures of A_{net} in citrus have previously been taken at the leaf level (Kriedemann, 1968;

Mesquita et al., 2016; Tsagarakis et al., 2012; Vu and Yelenosky, 1988; Vu et al., 1986;

Zambrosi et al., 2011) and at the field level (Contreras et al. 2017; Er-Raki et al., 2009; Maestre-

Valero et al., 2017; Penddinti and Kambhammettu, 2019). Due to the specificity of the leaf level

measurements (light levels, angle of radiation incidence, leaf position within the canopy), leaf-

level data is difficult to extrapolate to the whole canopy. Because HLB induces reductions in

canopy density (Tang et al. 2019), larger scale measures may be more indicative of changes in

carbon assimilation than leaf-level measures.

In this study, our objective was to compare relationships of A_{net}, CLas infection, and

foliar symptoms at the leaf and shoot level. To achieve our objective, we first developed and

validated the use of a gas exchange chamber capable of measuring whole shoots. We assessed

the value of such measures by comparing leaf-level and shoot-level light response curves. In our

subsequent HLB experiment, we hypothesized that: 1) greater time of CLas inoculation exposure

would increase starch content; 2) higher starch content, as an indicator of reduced leaf

carbohydrate export, would correlate with downregulation of A_{net}; and 3) because HLB causes

morphological changes at the canopy level, mean leaf area, leaf number, and stem surface area

and length were expected to affect A_{net} at the shoot level.

Materials and Methods

Study 1: Whole shoot gas exchange chamber development and testing

Plant Material

The trees utilized were greenhouse-raised, four-year-old 'Valencia' sweet orange (*Citrus*

x sinensis) on 'Kuharske' (*C. sinensis* x *Poncirus trifoliata* (L.) Raf.) rootstock in 20-L pots.

Plants were maintained with recommended rates of slow-release fertilizer and 3x weekly

irrigation with sufficient volume to exceed field capacity of the pots.

Chamber construction

The chamber was built of acrylic tubing, acrylic glue, and a 3-D printed adapter designed

and provided by Jason Hupp, LI-COR Bioscience, Inc. Lincoln, NE, USA. The chamber was fit

to an infrared gas analyzer (LI-6800, LI-COR Bioscience, Inc. Lincoln, NE, USA) using a

custom chamber adapter (LI-6800-19, LI-COR Bioscience, Inc. Lincoln, NE, USA). To seal the

opening, a large conical stopper was step-drilled and cut to provide an easy-to-fit cap around the

stem of the branch (Appendix S1). The chamber could then be made airtight using plumbers'

putty.

Leaves and shoots were measured on trees recently acclimated to the relative darkness of

the laboratory environment (2-4 μmol photons m^{-2} s^{-1}). Leaf-level measurements were recorded

for 15 minutes on three leaves at photosynthetic photon flux density (PPFD) of 1000 μmols m^{-2}

s^{-1}. Environmental settings for the leaf level were set at flow 300 μmol s^{-1}, 0.2 kPa pressure, 60%

relative humidity, reference CO_2 at 400 μmol CO_2 mol^{-1}, and a fan speed of 10,000 RPM.

Chamber measurements were logged every 15 seconds for five minutes after a 15-minute

stabilization period under controlled conditions at PPFD of 1000 μmol m^{-2} s^{-1}. Environmental

settings for the whole-shoot chamber were set based on volume, 1000 cubic centimeters, at flow

1000 μmol s^{-1}, 0.2 kPa pressure, 60% relative humidity, reference CO_2 of 400 μmol CO_2 mol^{-1},

and a fan speed of 14,000 RPM to maximize air mixture. Increases in air flow rate from 300

μmol s^{-1} in leaf measurements to 1000 μmol s^{-1} in whole -shoot chamber compensated for the

increase in photosynthetic material and increased volume present in the custom chamber. The fan

speed was increased to 14,000 RPM to help mix air in the chamber as recommended by the

IRGA manufacturer. Using the red-light capability of the red, far red, and blue light adjustable

fluorometer head unit of the LI-6800F, light intensity was controlled for the leaf-level

measurements. An LED Grow Light of similar spectral composition (ZXMean 1500W Full

Spectrum LED, BliGli, Wuhan HuBei, China) was manipulated by distance and a dimming

switch (Credenza Plug-In Lamp CFL-LED Dimmer, Lutron Electronics Co., Coopersburg, PA,

USA) to achieve desired PPFD for the whole-shoot chamber. PPFD for whole-shoot

measurements was measured using the quantum sensor of the LI-6800 at the mid-point of the

shoot adjacent to the whole-shoot chamber.

Stem assimilation

To determine stem contribution to A_{net} of the shoot (the leaf and stem material together),

the stems were covered with fitted black drinking straws following the method of Nilsen and Bao

(1990) testing stem contributions to photosynthesis in soybean (*Glycine max* L.). Shoots were

exposed to PPFD 1400 µmol m^{-2} s^{-1} as described above. The shoot was then removed from the

chamber and black straws were measured to fit the space between petioles on the stem, sliced

lengthwise and wrapped around the stem (Appendix S2). Gas exchange measurements then

rerecorded with the darkened stems. The LED Grow Light was then turned off and the custom

chamber was covered by a black cloth to achieve 0 µmol m^{-2} s^{-1} PPFD to determine respiration of

the shoot, again following the same stabilization and logging duration. Measurements used were:

whole shoot in the light (Shoot A_{net}), lit shoot with only the stem darkened stem, and shoot dark

respiration R_d, measured as net CO_2 exchange at 0 µmol m^{-2} s^{-1} PPFD (shoot respiration) were

used for analysis. Past measurements of whole-plant photosynthesis have considered gross

photosynthesis to be $A_{net} - R_d$ (Hwang and Morris 1994), other authors have pointed out that

gross photosynthesis is more accurately interpreted as $A_{net} - R_d -$ photorespiration, and that $A_{net} -$

R_d is better described as apparent photosynthesis (A_{app}) (Wohlfahrt and Gu 2015). Measurements

of A under the different measurement conditions were used to calculate A_{app} contributions of the

stem and leaves as follows:

• Shoot A_{net}: shoot net photosynthesis, measured as A of shoot at 1400 µmol m^{-2} s^{-1} PPFD;

inherently the sum of stem and leaf A_{app} minus respiration (shoot R_d).

• Stem A_{app}: stem apparent A; measured as the difference between shoot A_{app} under full

illumination and A_{app} with darkened stem.

• Shoot R: whole shoot respiration, measured as $|A|$ at 0 µmol m^{-2} s^{-1} PPFD.

• Shoot A_{app}: shoot apparent assimilation, estimated as shoot A_{net} – shoot R_d .

• Leaf A_{app}: leaf apparent assimilation of all leaves on the shoot, estimated as shoot A_{app} –

stem A_{app}

Immediately following gas exchange measurements, shoots were excised and defoliated

for leaf area analysis using the LI-3100 (LI-COR Biosciences, Inc. Lincoln, NE, USA). Stems

were measured for length and diameter (Appendix S3) to calculate surface area using an adjusted

formula for the lateral area of a truncated cone (Equation 1). Due to the irregular shape of citrus

stems, two diameter measurements were taken at the widest and narrowest widths at the base,

middle, and tip of the stem. The circumference of an ellipse is approximately equal to that of a

circle with a radius of the average of the widest and narrowest point of the ellipse.

$Stem\ Surface\ Area\ =\ \pi(r_1 + r_2)\sqrt{(r_1 - r_2)^2 + h_1{}^2} + \pi(r_2 + r_3)\sqrt{(r_2 - r_3)^2 + h_2{}^2}$ (Eq. 1)

Where:

r_1= radius of stem base

r_2= radius of stem middle

r_3= radius of stem tip

h_1 =length between measurement point of r_1 and r_2

h_2 =length between measurement point of r_2 and r_3

Leaves and stems were dried at 60 °C for 72 hours and weighed separately. Leaf and

stem surface area and dry weight were used to calculate A_{app} rates per meter squared and per

gram dry weight.

Leaf and shoot light response curves

Nine fully mature young shoots, three from each of the same trees were measured in the

laboratory, because ambient light in the greenhouse prevented low-light manipulation of shoot

level environment. Shoots were placed in the shoot gas exchange chamber and carbon

assimilation was measured at PPFD of 0, 25, 50, 100, 150, 200, 400, 600, 800, 1000, 1400, and

1800 µmols m^{-2} s^{-1}. Environmental settings, stability criteria, and logging were the same as

previous whole-shoot measurements (Appendix S4). The largest, healthiest leaf (assessed

visually) on each of the nine measured shoots was selected to measure a leaf-level comparison

(Appendix S5) measurement at PPFD of 0, 25, 50, 100, 150, 200, 400, 600, 800, 1000, 1400, and

1800 µmols m^{-2} s^{-1}. Environmental settings and stability criteria were set as in the previously

described measurements.

Study 2: HLB impacts on leaf and shoot net CO_2 assimilation

Plant material selection

Trees were selected based on length of exposure to citrus greening infection. The trees

utilized were greenhouse-raised, four to five-year-old sweet orange 'Valencia' on 'Kuharske'

rootstock in 20-L pots. These trees had been exposed to ACP inoculation of *C*Las by growing

plants in infested screenhouses for varying lengths of time. Exposure categories were defined as

follows: the High category of exposure had been exposed to infected ACP feeding and breeding

for 2.5 years; the Mid category of exposure had been exposed to infected ACP feeding and

breeding for 1.5 years; and the Low category of exposure had been exposed to infected ACP

feeding and breeding for 1 year. The uninfected trees had not been exposed to infected ACP.

Three trees of each category were arranged in a greenhouse in a randomized complete block

design, totaling 12 trees for the study.

Shoots and leaves were selected on each tree for measurements. Three shoots per tree

were selected from the exterior of the canopy. Shoots were chosen based on accessibility of the

instrument using the custom chamber, and only those with at least three leaves suitable for leaf

measurement using the standard leaf chamber were selected. Thirty-six shoots were measured in

this study (3 shoots x 3 trees x 4 exposure groups).Three leaves were selected per shoot. These

were distributed along the length of the shoot, always using one apical leaf. To be selected,

leaves had to be large enough to completely cover the leaf chamber (6-cm^2 circle). A total of 108

leaves were measured for this study (36 shoots x 3 leaves).

Gas exchange measurements

A_{net}, transpiration (E), and conductance were measured in using the LI-6800 (LI-COR

Bioscience, Inc. Lincoln, NE, USA). In the case of leaf-level measurements, conductance

represented stomatal conductance to water (g_s) while the shoot level measurements estimate

shoot conductance to water (shoot g), because non-stomatal pores, such as lenticels may

contribute significantly conductance at the shoot level. Leaf chamber environmental settings

were set at a PPFD of 1000 μmol m^{-2} s^{-1} and shoot chamber was placed at PPFD of 1400 μmol

m^{-2} s^{-1} with all other settings according to those described above for Study 1 for leaf and shoot

gas exchange measurements.

CLas quantification

Midrib and petiole tissues excised from selected leaves were chopped using a razor blade

and bead-beaten using Vortex Genie 2 (Scientific Industries Bohemia, NY, USA) to a fine

homogenized powder in liquid nitrogen. Total genomic DNA was extracted and purified from

100 mg of homogenized material using GeneJET Plant Genomic DNA Purification Mini Kits

(Thermo Fisher Scientific Waltham, MA, USA). Samples were processed following

manufacturer instructions and stored at -20° C. Primers and TaqMan probes were obtained for

*C*Las based on method developed by Coy et al. (2014). The PCR master mix was used with 10

μL TaqMan Buffer, 8.6 μL sterile deionized water, 0.4 μL CQUL primer, and 1 μL of extracted

purified DNA. PCR reactions performed with standards using QuantStudio 3 Real-time PCR

(Applied Biosystems-Thermo Fisher Scientific Waltham, MA, USA) yielded cycle threshold

values (Ct). Bacterial copy-count estimates were generated by regression using processed Ct

results from standards of known copy quantity, processed in each plate batch. At the shoot level

Ct and bacterial copy number were considered to be the mean of the measurements in the three

selected leaves on each shoot.

Starch content

Starch content was determined by extracting starch from a single 28.27 mm^2 hole punch

on the selected leaf blade (Whitaker et al., 2014). The leaf punch was homogenized for 2.5

minutes at 6500 RPM in 500 μL of distilled water with five beads. Samples were boiled for 10

minutes at 200° C and agitated prior to a two-minute centrifuge at 2500 rpm, after which 300 μL

of the supernatant was added to 900 μL pure ethanol and agitated for three seconds to precipitate

starch. Prepared samples were centrifuged at 14000 RPM for 10 minutes and the supernatant was

discarded and replaced by one mL of deionized water. Samples were then shaken for four-

minutes to dissolve the starch pellet and two 250 µL of prepared sample were each placed in

individual wells of a 96 well microplate, and 50 µL of 2% iodine was added to each well.

Absorbance at 595nm was read using SpectraMax 250 microplate reader (Molecular Devices San

Jose, CA, USA). Linear functions based on standard curve concentrations included in each plate

were generated with the absorbance of standards to estimate starch content. As for CLas

variables, at the shoot level starch values were considered to be the mean of the measurements of

the three selected leaves on each shoot.

Visual rating

A rating of the visual symptoms of CLas infection was assessed on a scale of 0-10, where

0 = no symptom, 10= full symptom expression, on each leaf. Prior to assessment a scale of leaf

symptoms was generated using photos of sampled CLas infected leaves to ensure consistency in

ratings. All disease ratings were performed by the same individual. Visual symptom ratings used

in the leaf level measurements were recorded for every leaf on the measured shoot; an average of

the assessment of all leaves was used for the shoot value.

Shoot morphology

Total leaf area per shoot was measured using a LI-3100 (LI-COR Bioscience, Inc.

Lincoln, NE, USA) immediately following the assimilation measurements and visual ratings,

prior to sampling for qPCR and starch content. Stem length and diameter in two directions was

measured using a micrometer to generate a stem surface area estimates using Eq. 3. Average

internode length was calculated by dividing the stem length by the total number of leaves per

stem.

Analysis

Analysis of variance was performed using the anova() command in base R, after

assessing appropriateness of parametric tests by testing for normal distribution, to assess the

effect of treatment groups (R Core Team, 2019, Version 3.6.1). Where exposure group was

significant, Fisher's LSD was used as a post hoc test for least significant difference between the

four tree categories used for all variables using the {agricolae} package (de Mendiburu, 2020).

Correlation analysis was generated among the variables at the shoot and leaf-level and

correlation matrices were tested for significance of Pearson's correlations using the

{Hmisc}package (Harrell, 2020). For correlation analyses, because some variables might co-vary

in the case of disease, tests were performed with and without the HLB- plants. For variables with

significant correlations, linear regression models were generated at both shoot and leaf-level

using the lm() command, similarly comparing relationships with and without HLB- plants. The

models were tested for significant relationships using anova(). All linear models nested shoot

within tree, and leaf-level data was nested leaf within shoot. A natural log transformation of

bacterial copy count and starch content facilitated a linear relationship between the transformed

data. Light response curves for leaf and shoot were fit using an nonlinear least squares regression

using the self-starting asymptotic function in R (Pinheiro et al., 2014) with a function supplied

by Tomeo (2017). This function fits an asymptotic function and calculates the parameters of

fully saturated photosynthesis (A_{sat}), quantum yield of photosynthesis (Φ), dark respiration (R_d),

light compensation point (LCP), and PPFD of 75% saturation (Q_{sat75}). These parameters were

then subjected to analysis of variance to test the effects of shoot vs. leaf light responses.

Results

Stem contributes significantly to whole-shoot photosynthesis

Vegetative stem on young shoots contributed 5.7% to total shoot surface area, 17.0% to

total shoot mass, and 10.7% to total shoot photosynthesis (μmol CO_2 s^{-1}; Table 1). Mean stem

A_{app} was nearly double leaf A_{app} (9.62 μmol CO_2 m^{-2} s^{-1} and 5.13 μmol CO_2 m^{-2} s^{-1}, respectively;

Figure 2).

Light response curves differ between shoots and leaves

Leaves and shoots had distinct light response curves with differences in all parameters

tested, except for quantum yield (Table 2). Leaf level R_d was more than 3x that of the shoot level

R_d, resulting in a correspondingly higher LCP, despite the less acute rise in A_{net} response to

PPFD (Figure 3). Leaf level 75% saturation was achieved at approximately 528 μmol photons

m^{-2} s^{-1}, while that of shoots was estimated at 1532 μmol m^{-2} s^{-1}. The corresponding A_{sat} was also

approximately 20% higher in shoots than in leaves.

Photosynthesis and starch are affected by CLas exposure

Uninfected leaves had a higher A_{net} (mean of 3.97 μmol CO_2 m^{-2} s^{-1}) than infected leaves;

however, as CLas exposure increased from low, to medium, to high so did A_{net} (means of 1.91,

2.61, 2.70 μmol CO_2 m^{-2} s^{-1}, respectively; Table 3). Leaf-level g_{sw} intermediate in the uninfected

category, while it was lower in the low and medium exposure groups, while g_{sw} was highest in

the high exposure group. A similar pattern was observed for leaf-level E, though the P-value for

the effect of exposure category was inconclusive at $P=0.058$ (Table 3). Uninfected leaves had a

lower starch content (mean of 1.08 µg mm^{-2}) than infected leaves; however, as CLas exposure

increased from low to high starch content decreased (means of 11.31, 5.69, 3.55 µg mm^{-2},

respectively). Uninfected leaves had the lowest visual rating score, which increased as CLas

exposure increased from a mean of 0.22 in the uninfected to a mean of 2.22 in the highest

exposure group.

A_{net} had a positive relationship with CLas exposure at the shoot-level. High exposure

shoots had the highest A_{net} at a mean of 3.37 µmol m^{-2} s^{-1}, which decreased with a shortened

exposure down to a mean of 1.91 µmol m^{-2} s^{-1} in the uninfected shoots (Table 4). This pattern

was similar for both g and E, with those of two or more infected exposure groups, including the

high exposure group for both variables, showing higher values than the uninfected plants.

However, as with leaf-level E, the P-value for exposure category differences was inconclusive at

$P=0.061$ (Table 4). Reductions in mean leaf area (17.59 cm^2 in the uninfected to 11.38 cm^2 in

the highest exposed group) and total shoot surface area (133.5 cm^2 in the uninfected to 68.3 cm^2

in the highest exposed group) were found as CLas exposure increased. Exposure did not affect

internode length. Despite the different pattern of impact of CLas exposure, the overall range of

values for shoot- and leaf-level A, E, and g were similar, except in the unexposed category, in

which shoot-level values were markedly lower than at the leaf level.

Infection, starch, photosynthesis, and morphological variables were correlated

At the leaf level, when including uninfected trees, positive correlations were found

between bacterial copy count and starch content ($r=0.609$, $P<0.00001$), CLas copy number and

visual rating (r=0.459, P<0.00001), and starch content and visual rating (r =0.276, P=0.005;

Appendix S6). Negative correlations were found between A_{net} and CLas copy number (r = -

0.322, P=0.0009) as well as starch content (r= -0.243, P=0.013). No correlations were found

with leaf position or between A_{net} and visual rating. No additional correlations were found when

excluding uninfected trees.

At the shoot level, when including uninfected trees, positive correlations were found

between CLas copy number and starch content (r=0.716, P<0.00001), visual rating (r =0.539,

P=0.001), and A_{net} (r=0.413, P=0.015); between shoot surface area and mean leaf area (r=0.812,

P<0.00001) and shoot surface area and shoot length (r=0.58, P=0.0003); and between internode

length and shoot length (r=0.344, P=0.046; Appendix S8). When excluding uninfected trees at

the shoot level, additional positive correlations were found between starch content and mean leaf

area (r=0.421, P=0.032) and shoot surface area (r=0.455, P=0.019); between shoot surface area

and mean leaf area (r=0.809, P<0.00001) and shoot length (r=0.644, P=0.0004); and between

internode length and A_{net} (r=0.558, P=0.003; Appendix S9).

Linear regressions reveal additional relationships between infection and physiology

Based on the correlations we tested linear regressions of the following variables: CLas

copy number by starch content, internode length by A_{net}, and foliar starch content by shoot

surface area. After a natural log transformation of both starch content and bacterial copy count, a

positive linear relationship was found at both the leaf and shoot-level (Table 5). Further

observation indicated a change in the slope as CLas gene copy # increased and assessment using

breakpoint analysis indicated that the increased starch content at the leaf level increased at a

higher rate with a breakpoint at approximately $10^{11.2}$ CLas gene copies (Figure 4A), though the

shoot level data did not indicate such a change in slope (Figure 4B). A positive linear

relationship (P=0.048) was found between foliar starch with shoot surface area when excluding

uninfected shoots, though this relationship also had a low R^2 (0.18) (Figure 5 and Table 5). This

relationship is inverse to the CLas exposure duration, with the highly exposed shoots having

lower starch content and lower total shoot surface area. When comparing morphological and A_{net}

measurements, a positive linear relationship (P=0.003) was found between internode length and

A_{net} when excluding uninfected shoots (Figure 6 and Table 5). These morphological adjustments

are illustrated in Figure 7.

Conductance also influenced A_{net} at both leaf and shoot levels, where exposure category

also influenced the slope of the relationship. In leaf-level measurements medium and low

exposure categories of CLas exposure increased the slope of the A_{net} response to g_{sw}, while the

high exposure category decreased the slope below that of the uninfected leaves (Figure 8).

However, the high exposure category had more disperse data, and slope calculations may be

more subject to influence of single values. At the shoot level, CLas infection reduced the slope of

the A_{net} response to shoot g, while increasing exposure led to increasing slope.

Discussion

Leaf and shoot light response dynamics differ

The distinct shoot light response dynamic helps to explain the success of *Citrus* in high

light regions, where maximum A_{sat} is attained near the maximum ambient PPFD occurring in

subtropical regions. The leaf-level light response curves indicating Q_{sat} occurs between 600-700

$\mu mol\ m^{-2}\ s^{-1}$ PPFD agreed with earlier studies (Habermann et al., 2003). The increased light

saturation point of citrus shoots compared to individual leaves could be explained by varied

angle of incident radiation, as well as by partial shading of basal leaves by more distal leaves.

However, R_d was much lower at the shoot level than at the leaf level. The light response curve R_d

estimates are supported by the measurements at 0 µmol PAR m^{-2} s^{-1}, where the estimates were

similar: 1.3 ± 0.13 µmol CO_2 m^{-2} s^{-1} for leaf and 0.47 ± 0.05 µmol CO_2 m^{-2} s^{-1} for shoots to the

estimates produced by fitting the asymptotic function. Lower R_d is an important factor in

reducing the LCP. Thus, at the shoot level, citrus exhibits higher A_{net} at both the lower and upper

ends of irradiance than is detected at the leaf level, while at PPFD ranges between 50-1000 µmol

PAR m^{-2} s^{-1}, shoots exhibit lower A_{net} than leaves. The lack of a difference in Φ is further

confirmation that differences between shoots and leaves are primarily determined by factors such

as morphology, not by enzymatic or photochemical factors. This might be assumed, given that

some of the same leaves were involved in both measurements, though the lower rate of R_d in

shoots merits further investigation. Because variations in irradiance within shoots may explain

much of the difference between shoot and leaf irradiance responses, morphology is likely to have

a strong impact on photosynthesis at the whole-shoot level. Thus, morphological impacts of HLB

may help explain the impact of the disease on photosynthesis and growth.

Stem assimilation contributes to whole shoot photosynthesis

Citrus vegetative stems significantly contributed to shoot A_{app}. The stem was also more

efficient in A_{app} on a surface area basis than leaves. The high assimilation rate of citrus stems

could be explained by area-related chlorophyll content where young stems contain 50–70% more

chlorophyll than adjacent leaves (Kharouk et al. 1995; Pfanz et al., 2002; Pilarski 1984; Schmidt

et al. 2000; Solhaug et al. 1995). Although many factors contribute to whole-shoot assimilation,

the previously unquantified contribution of the photosynthetic stem to canopy wide assimilation

may be an additional factor limiting extrapolability of leaf-level gas exchange measurements to

canopy-wide measurements. Stem internal re-fixation of CO_2 in young branches of perennial

woody species have been shown to compensate for 60–90% of the potential respiratory carbon

loss (Pfanz et al., 2002). Although, its contributions compared to the leaves on whole shoot

assimilation are relatively small, the proportion of A_{net} coming from the stem may depend on

shoot morphological characteristics such as internode length that affect light interception in the

stem. This observation highlights the importance of morphological impacts of HLB on the

whole-shoot photosynthetic capacity.

Shoot morphology and the effects of HLB of leaf and shoot photosynthesis

As with light responses, shoots and individual leaves differed in their responses to HLB.

While *C*Las infection reduced A_{net} at the leaf level, it increased A_{net} at the shoot level, while

inducing whole-shoot morphological adjustments over time. Although the measured values of

A_{net} were low (~2-4 µmol m^{-2} s^{-1}), these values are typical of leaves of greenhouse-grown citrus

trees (Syvertsen 1994). The morphological impacts explain the HLB effects on whole-shoot

photosynthetic rates. The decreased leaf and shoot surface area effected by *C*Las exposure reflect

the symptom of canopy thinning observed in infected citrus trees in the field. This process

appears to occur on a scale of years. Given that most citrus leaves senesce by 17 months

(Wallace et al. 1954), the changes in shoot morphology appear to occur over a complete turnover

of old to new tissues.

We have not found any report in which a plant produced both longer internodes and

smaller leaves in response to a disease or environmental condition. Typically, an increase in leaf

size has coincided with increased internode length to reduce self-shading, such as in acclimation

to low light conditions (Kavanagh, 2014; Pieters et al., 1998; Poorter and Rozendaal, 2008;

Robertson, 1994), thus the peculiar morphological changes induced by HLB may be responses to

the specific challenges imposed by the disease.

These specific morphological changes were associated with the shoot-level gas exchange

response to HLB. Light penetration and absorption are the primary drivers of photosynthesis, and

these are heavily influenced by canopy structure (Niinemets, 2007). Canopy thinning by CLas-

induced defoliation, as described by de Graça (1991), within the high CLas-exposed category

(fewest and smallest leaves and longest internode length) further facilitate light penetration into

the canopy. With increased light penetration due to smaller leaves and greater internode length,

typically shaded leaves, further from the shoot apex, and the photosynthetic stem contribute

more to shoot-level assimilation, which help explain the increased assimilation efficiency found

in shoots of the high CLas exposure group. Considering the contribution of stems to shoot A_{app},

the relative increase in stem surface area could also contribute to the enhanced shoot A_{net}. The

increased light exposure of all parts of the shoot may also lead to compensation for the localized

disease effects discussed below in reference to leaves. Although there are likely other factors that

contribute to the increased assimilation efficiency of HLB-affected shoots, the correlation of

these morphological variables with A_{net} indicates that whole-shoot morphology is a defining

feature of shoot acclimation that leads to an increase in A_{net}.

Another defining feature of the acclimation of citrus trees to CLas infection appears to be

impacts on conductance and E and their interactions with A_{net}. Citrus are low conductance

species, and other studies have found strong dependence of leaf-level A_{net} on g_s under a variety of

abiotic conditions, with linear responses up to g_s of 0.3 mol m^{-2} s^{-1}, and changes in this

relationship have been interpreted as relating to impacts on nonstomatal photosynthetic processes

directly (Khalid et al. 2021, Shafqat et al. 2021). Depending on whether shoot or leaf levels and

whether conductance or transpiration are considered, infection either initially reduced and then

increased or simply increased gaseous water exchange of leaves and shoots. This may seem to

contrast with the observations of reduced sap flow in HLB-affected trees (Kadyampakeni et al.

2014), but the progressively reduced shoot surface area led to a total per shoot E that was similar

in uninfected (52.7 ± 3.78 mmol s^{-1}) and high exposure infected (57.8 ± 2.78 mmol s^{-1}) groups,

and these instantaneous measurements do not include possible limitations to transpiration of

infected plants under high diurnal evaporative demand. CLas infection may increase conductance

by morphological (increased stomatal density) or behavioral (preventing stomatal closure)

mechanisms, but the present study does not reflect on which mechanisms may be at work.

The impact of CLas exposure categories on the relationship between conductance and

A_{net}, however, helps clarify the impact of HLB on photosynthesis. At the leaf level, mid and low

exposure categories reduced g_s but showed an increased relative response of A_{net} to g_s. This

indicates that reduced g_s drove the photosynthetic reduction in the lower exposure group.

However, at the shoot level the impact of exposure was nearly opposite, with the low exposure

group showing a marked decrease in the photosynthetic response to shoot g. This indicates that at

the shoot level, non-conductance factors are more important, implicating carbon fixation

processes directly, perhaps driven by source-sink attenuation, discussed in detail below.

The increased efficiency of A_{net} should not be confused with increased total

photosynthesis. While the highly CLas-exposed shoots had higher A_{net} than the shoots from the

low exposure group, these measures are surface-area based and do not offset carbon fixation lost

in the reduction in shoot surface area with increased CLas exposure. Total net shoot assimilation

was 40% higher for the uninfected shoots (0.032 µmol CO_2 s^{-1}) than high CLas-exposed shoots

(0.023 μmol CO_2 s^{-1}). The shoot-level mitigation effect is evident here in that, if leaf level A_{net}

are extrapolated to the whole shoot surface area, the estimates values would be 0.048 μmol CO_2

s^{-1} for the uninfected shoots and 0.018 μmol CO_2 s^{-1} for the high-exposure shoots, over- and

underestimations for uninfected and infected shoots, respectively.

Although, growth is necessarily dependent on photosynthesis, numerous complexities

have led to a low degree of correlation between net assimilation and growth. For example,

Poorter and Remkes (1990) did not find leaf-level A_{net} to be correlated with relative growth rate

in a comparison of numerous species, though other multi-species comparisons have found effects

of net assimilation rate on relative growth rate (Li et al. 2016). In the context of growth-limiting

abiotic conditions, Zait et al. (2019) found that impacts of abiotic challenges on A_{net} versus

effects on growth rate were different among water deficit, salinity, and high temperature,

identifying contributions of several components of gas exchange. Measurements of A_{net} using gas

exchange are further complicated in that they are limited in both spatial (canopy dimensions) and

time scales, and neither growth-analysis-based net assimilation rate nor gas-exchange-based A_{net}

include the complexities of allocation and their subsequent impacts on growth (Pereira 1995). In

most studies, however, total whole canopy leaf area has been found to affect growth more than

photosynthetic activity on a leaf area basis (Pereira 1995, Sinclair et al. 2019). The present study

provides an example of how morphological changes could overwhelm changes in A_{net}. Despite

increases in A_{net}, CLas infected shoots had lower total assimilation rates because the reduction in

photosynthetic surface area was greater than the relative increase in A_{net}, leading to a reduction in

assimilation per shoot. Thus, citrus plants mitigate HLB-driven losses by increasing A_{net} but do

not increase total assimilation.

CLas infection induces local starch and photosynthetic responses

Because *C*Las is a phloem-limited bacterium, the effects of infections on leaf

carbohydrate dynamics have been the focus of much HLB-related research. Achor et al. (2010)

observed starch disruption of chloroplast inner grana in association with local phloem blockage

and collapse. Further, Fan et al. (2010) found increased foliar sucrose concentrations in

association with starch accumulation in infected trees, where Welker et al. (*en presse*) confirmed

the sucrose results and observed reduced leaf export rates. The current study suggests that these

effects are localized, where *C*Las population increases induced starch accumulation at a much

higher rate after reaching a population threshold at the leaf level.

Given these earlier findings regarding the impact of HLB on citrus carbohydrate

relations, Cimò et al. (2013) simulated the foliar response of HLB by girdling citrus trunks,

resulting in starch accumulation and a reduction in leaf level A_{net}. This reduction had been

hypothesized because of the source-sink hypothesis (Ainsworth and Bush 2011), which observes

that plants downregulate A_{net} in response to reduced leaf carbohydrate export, a condition in

which various carbohydrates may accumulate. In the case of citrus, this process was

demonstrated by Syvertsen (1994), and later by Iglesias (2002) and Nebauer et al. (2011), who

found A_{net} responded to changes in soluble sugars, which in turn also resulted in starch

accumulation or depletion. Although none of the above studies addressed the condition of HLB,

their interpretation of reduced photosynthesis as a response to leaf export limitation has been

applied to studies of HLB, in which gene expression of photosynthetic pathways has been found

to be downregulated (Balan et al. 2018). The present study found the expected reduction of A_{net}

at the leaf level, but not at the shoot level.

The comparison of local versus systemic effects has not been addressed explicitly by

studies of HLB. However, the authors of some studies have implicitly interpreted systemic

effects on leaves (all studies reviewed by Balan et al. 2018) while others have hypothesized that

HLB primarily affects roots (Johnson et al. 2014). The indication in the present study of a

localized impact of CLas on leaf carbohydrate export and photosynthesis supports the

interpretation of Koh et al. (2012) and Welker et al. (*en presse*), who observed callose blockage

of leaf phloem as well as reduced export of ^{14}C-labeled photosynthates in infected trees, while

also agreeing with those of Fan et al. (2010), Rao et al. (2018), and Balan et al. (2018), all of

which found alterations in carbohydrate metabolism and photosynthesis in leaves. A unique

contribution of the present study is to indicate that these effects are at least partly local effects of

presence of the bacterial population in the leaf, rather than strictly systemic responses to

infection. One recent study found that impacts of HLB on stem transport speeds also correlated

with CLas populations in the same stem segments, indicating an additional localized impact on

transport (Welker et al. *en presse*).

Starch accumulation has been associated with CLas infection and has been attributed as

the cause of the blotchy mottle and chlorosis symptoms seen in infected citrus leaf tissue

(Etxeberria et al., 2009; Gonzalez et al., 2012; and Schaffer et al., 1986). Interestingly, we found

high starch accumulation in the most recently exposed group and a relative reduction in starch

content with longer exposure durations. This observation, including the recovery of the A_{net}

response to g in the high exposure category indicates that the morphological acclimation leads

toward an equilibrium between source area and export and transport capacity under the

constraints of the disease. However, the starch levels in the high exposure group remained

elevated above those of healthy trees, suggesting that a complete source-sink balance is not

possible with CLas infection. Why photosynthesis does not further acclimate to avoid starch

accumulation is an important consideration for future studies. Meanwhile, visual estimates of

symptom severity increased across exposure groups, indicating that more severe symptoms were

not the result of greater starch accumulation.

The CLas exposure categories themselves did not correspond to differences in leaf CLas

population. Given the systemic nature of CLas infection, this indicates that even the mild

exposure category was sufficient to attain systemic distribution of the pathogen. The lack of

correlation between CLas titers and symptoms or disease severity has been born out in other

studies, and these studies suggest wide sources of variation in titers that are not directly related to

disease severity (Stover et al. 2014, 2016, Ibanez and Stelinski 2020, Vasconcelos et al. 2020).

The increasing disease severity across increasing exposure groups without increases in titer

indicates the cumulative chronic impact of infection, which does not appear to include increasing

bacterial populations.

Conclusion

This study provides the first direct comparison of CO_2 assimilation between leaf and

shoot levels, as well as the first comparison of assimilation in HLB-affected and healthy citrus.

Shoot- and leaf-level responses were distinct. Shoot light response curves were much more

gradual than those of leaves with A_{max} at approximately 2x the irradiance of the A_{max} of leaves.

This was explained by the combination of leaf angles and photosynthetic stems, which

contributed a mean of 10% of the total shoot A_{app}. Likewise shoot and leaf responses to CLas

infection differed. Where A_{net} was reduced at the leaf level, shoot morphological acclimation

partially compensated for this downregulation in leaf-level photosynthetic activity. These

morphological changes further confirm the importance of canopy density as an important

measure of disease severity. This study is an example of the value of assessing physiological

processes along temporal and spatial scales, with whole-shoot photosynthesis providing one tool

for such efforts. Further studies should consider the impacts of HLB on photosynthetic

acclimation to understand the limitations of source-sink acclimation. Understanding the nature of

citrus stem gas exchange would further

Supplemental materials

Appendix S1. Cap design of whole shoot assimilation chamber. A) bottom view – stepped

drilled hole to limit low leaf interference, B) top view – straight cut to allow for stem placement,

C) fit of cap into chamber, D) showing fit into chamber to prevent leakage.

Appendix S2. Straw pieces fit to cover stem of 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) shoot to measure whole shoot gas exchange excluding stem photosynthesis.

Appendix S3. Diagram describing the measurements taken to estimate stem surface area.

Appendix S4. Standard deviation of A_{net} at photosynthetic photon flux density of 1000 µmols m^{-2}

s^{-1} demonstrating measurement stabilization for 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) in leaf chamber of whole shoot chamber. Dashed vertical line indicates stabilization

point for leaf; dotted line indicates stabilization point for shoots.

Appendix S5. Illustration of 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) shoots and

leaves selected for to compare assimilation in leaves vs. shoots.

Appendix S6. Correlation of leaf disease and photosynthetic variables of healthy and *Candidatus*

Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) exposed

to 0, 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri* Kuwayama)

feeding and reproduction. Copy # - log transformed bacterial copy count, Starch – log

transformed starch concentration (μg mm^{-2}), Visual – visual rating of citrus greening leaf

chlorosis and mottling symptoms, A – net CO_2 assimilation (μmol m^{-2} s^{-1}), and Position – leaf

position on stem (1-n starting at apical leaf). Darkened correlation boxes denote correlations

where $P \leq 0.05$

Appendix S7.   Correlation of leaf disease and photosynthetic variables of *Candidatus*

Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck) exposed

to 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri* Kuwayama)

feeding and reproduction. Copy # - log transformed bacterial copy count, Starch – log

transformed starch concentration (μg mm^{-2}), Visual – visual rating of citrus greening leaf

chlorosis and mottling symptoms, A – net CO_2 assimilation (μmol m^{-2} s^{-1}), and Position – leaf

position on stem (1-n starting at apical leaf). Darkened correlation boxes denote correlations

where $P \leq 0.05$.assimilation (μmol CO_2 m^{-2} s^{-1}), and Position – leaf position on stem (1-n starting

at apical leaf). Darkened correlation boxes denote correlations where $P \leq 0.05$.

Appendix S8. Correlation of shoot disease and photosynthetic variables of healthy and

Candidatus Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) exposed to 0, 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina*

citri Kuwayama) feeding and reproduction. Copy # - log transformed bacterial copy count,

Starch – log transformed starch concentration (μg mm^{-2}), Visual – visual rating of citrus

greening leaf chlorosis and mottling symptoms, A – net CO_2 assimilation (μmol m^{-2} s^{-1}), LA –

mean leaf surface area (cm^2), SA – total shoot surface area (cm^2), Length – stem length (mm),

and Inter – average internode length (mm). Darkened correlation boxes denote correlations

where P$\leq$0.05.

Appendix S9. Correlation of shoot disease and photosynthetic variables of healthy and

Candidatus Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck) exposed to 1, 1.5, and 2.5 years of inoculation via Asian citrus psyllid (*Diaphorina citri*

Kuwayama) feeding and reproduction. Copy # - log transformed bacterial copy count, Starch –

log transformed starch concentration (μg mm^{-2}), Visual – visual rating of citrus greening leaf

chlorosis and mottling symptoms, A – net CO_2 assimilation (μmol m^{-2} s^{-1}), LA – mean leaf

surface area (cm^2), SA – total shoot surface area (cm^2), Length – stem length (mm), and Inter –

average internode length (mm). Darkened correlation boxes denote correlations where P$\leq$0.05.

Conflict of Interest

The authors declare they have not conflict of interest with this study, including no stake in any

product described in the manuscript.

Funding

Funding for this project comes from the United States Department of Agriculture Multi-Agency

Coordination System (AP19PPQS&T00C165).

Acknowledgements

The authors wish to thank Myrtho Pierre for assistance in gas exchange measurements, Rebecca

Ebert for assistance with starch measurements, and Jason Hupp for providing the design for the

adapter with which the whole-shoot chamber was developed.

Authors' contributions

MK, CV, and DR conceived experiments. MK developed whole-shoot chamber and conducted

experiments. MK analyzed data and wrote manuscript. CV supervised experiments. CV and DR

supervised analyses and manuscript writing, and revised manuscript.

Data availability

All data available upon reasonable request to corresponding author.

References

Achor DS, Etxeberria E, Wang N, Folimonova SY, Chung KR, Albrigo LG (2010) Sequence of

    anatomical symptom observations in citrus affected with huanglongbing disease. Plant Pathol

    J 9:56–64.

Achor D, Welker S, Ben-Mahmoud S, Wand C, Folimonova SY, Dutt M, Gowda S, Levy A

(2020) Dynamics of *Candidatus* Liberibacter asiaticus movement and seive-pore plugging in

citrus sink cells. Plant Physiology.182(20):882-891.

Ainsworth EA, Bush DR (2011) Carbohydrate export from the leaf: a highly regulated process

and target to enhance photosynthesis and productivity. Plant Physiology 155:64–69.

Balan B, Ibáñez AM, Dandekar AM, Caruso T, Martinelli F (2018) Identifying host molecular

features strongly linked with responses to huanglongbing disease in citrus leaves. Frontiers in

plant science 9:277.

Bové JM (2006) Huanglongbing: A destructive, newly-emerging, century-old disease of citrus.

Journal of Plant Pathology 88(1):7-37.

Cen H, Weng H, Yao J, He M, Lv J, Hua S, Li H, He Y (2017) Chlorophyll fluorescence

imaging uncovers photosynthetic fingerprint of citrus huanglongbing. Frontiers in Plant

Science 8:1509.

Cimò G, Bianco RL, Gonzalez P, Bandaranayake W, Etxeberria E, Syvertsen JP (2013)

Carbohydrate and nutritional responses to stem girdling and drought stress with respect to

understanding symptoms of huanglongbing in citrus. HortScience 48:920–928.

Contreras S, Hunink JE, Baille A (2017) Water and carbon fluxes in irrigated citrus orchards

assessed from satellite data. FutureWater Report 174.

Coy MR, Hoffman M, Gibbard KHN, Kuhns EH, Pelz-Stelinski KS, Stelinski LL (2014) Nested-

quantitative PCR approach with improved sensitivity for the detection of low titer levels of

Candidatus Liberibacter asiaticus in the Asian citrus psyllid, *Diaphorina citri* Kuwayama.

Journal of Microbiological Methods 102:15-22.

da Graça JV (1991) Citrus greening disease. Annual Review Phytopathology 29:109-136.

Das AK, Nerkar S, Bawage S, Kumar A (2013) Current distribution of huanglongbing (citrus

greening disease) in india as diagnosed by real-time PCR. Journal of Phytopathology

162(6):402-406.

de Mendiburu F (2020) agricolae: Statistical procedures for agricultural research. R package

version 1.3-2. https://CRAN.R-project.org/package=agricolae

Er-Raki S, Chehbouni A, Guemouria N, Ezzahar J, Khabba S, Boulet G, Hanich L (2009) Citrus

orchard evapotranspiration: Comparison between eddy covariance measurements and the

FAO 56 approach estimates. Plant Biosystems 143(1):201-208.

Etxeberria E, Gonzalez P, Achor D, Albrigo G (2009) Anatomical distribution of abnormally high

levels of starch in HLB-affected Valencia orange trees. Physiological and Molecular Plant

Pathology 74(1):76-83.

Etxeberria E, Gonzalez P, Dawson W, Spann T (2007) An iodine-based starch test to assist in

selecting leaves for HLB testing. University of Florida, IFAS Extension, Gainesville,

HS1122.

Eyles A, Pinkard EA, O'Grady AP, Worledge D, Warren CR (2009) Role of corticular

photosynthesis following defoliation in *Eucalyptus globulus*. Plant, Cell and Environment.

32(8):1004-1014.

Falster D, Westoby M (2003) Leaf size and angle vary widely across species: What

consequences for light interception? New Phytologist 158(3):509-525.

Fan J, Chen C, Brlansky RH, Gmitter Jr. FG, Li Z-G (2010) Changes in carbohydrate

metabolism in Citrus sinensis infected with '*Candidatus* Liberibacter asiaticus'. Plant

Pathology 59:1037–1043.

Folimonova SY, Achor DS (2010) Early events of citrus greening (huanglongbing) disease

development at the ultrastructural level. Phytopathlogy 100(9):949-958.

Gonzalez P, Reyes J, Etxeberria E (2012) Characterization of leaf starch from HLB-affected and

unaffected-girdled citrus trees. Physiological and Molecular Plant Pathology 79:71-78.

Habermann G, Machado EC, Rodrigues JD, Medina CL (2003) CO_2 assimilation, photosynthetic

light response curves, and water relations of 'Pera' sweet orange plants infected with *Xylella*

fastidiosa. Brazilian Journal of Plant Physiology 15(2):79-87.

Harrell FE, Jr (2020) Hmisc: Harrell Miscellaneous. R package version 4.4-0. http://CRAN.R-

project.org/package=Hmisc

Hwang Y-H, Morris JT (1994) Whole-plant gas exchange responses of *Spartina alterniflora*

(Poaceae) to a range of constant and transient salinities. American Journal of Botany 81:659–

665.

Iglesias DJ, Lliso I, Tadeo FR, Talon M (2002) Regulation of photosynthesis through source:

sink imbalance in citrus is mediated by carbohydrate content in leaves. Physiologia

Plantarum 116(4):563-572.

Jifon JL, Syvertson JP (2003) Moderate shade can increase net gas exchange and reduce

photoinhibition in citrus leaves. Tree Physiology 23(2):119-127.

Johnson EG, Wu J, Bright DB, Graham JH (2014) Association of '*Candidatus* liberibacter

asiaticus' root infection, but not phloem plugging with root loss on huanglongbing-affected

trees prior to appearance of foliar symptoms. Plant Pathology 63(2):290-298.

Kadyampakeni DM, Morgan KT, Schumann AW, Nkedi-Kizza P (2014) Effect of irrigation

pattern and timing on root density of young citrus trees infected with huanglongbing disease.

HortTechnology 24:209–221.

Kavanagh PH (2014) Herbivory and the evolution of divaricate plants: Structural defences lost

on an offshore island. Austral Ecology 40(2):206-211.

Khalid MF, Vincent C, Morillon R, Anjum MA, Ahmad S, Hussain S (2021) Different strategies

lead to a common outcome: Different water-deficit scenarios highlight physiological and

biochemical strategies of water deficit tolerance in diploid vs. tetraploid Volkamer lemon.

Tree Physiology: tpab074 https://doi.org/10.1093/treephys/tpab074

709
Kharouk VI, Middleton EM, Spencer SL, Rock BN, Williams DL (1995) Aspen bark

photosynthesis and its significance to remote sensing and carbon budget estimate in the

boreal ecosystem. Water, Air, and Soil Pollution 82:483-497.

Koh E-J, Zhou L, Williams DS, Park J, Ding N, Duan Y-P, Kang B-H (2012) Callose deposition

in the phloem plasmodesmata and inhibition of phloem transport in citrus leaves infected

with "*Candidatus* Liberibacter asiaticus". Protoplasma 249:687–697.

Kriedemann PE (1968) Some photosynthetic characteristics of citrus leaves. Australian Journal

of Biological Science 21:895-905.

Maestre-Valero JF, Testi L, Jiménez-Bello MA, Castel JR, Intrigliolo DS (2017)

Evaportranspiration and carbon exchange in a citrus orchard using eddy covariance.

Irrigation Science 35:397-408.

Mesquita GL, Zambrosi FCB, Tanaka FAO, Boaretto RM, Quaggio JA, Ribeiro RV, Mattos D

(2016) Anatomical and physiological response of citrus trees to varying boron availability are

dependent on rootstock. Frontiers of Plant Science. 7:224.

Nebauer SG, Renau-Morata B, Guardiola JL, Molina R (2011) Photosynthesis down-regulation

precedes carbohydrate accumulation under sink limitation in Citrus. Tree Physiology 31:169-

177.

Niinemets U (2007) Photosynthesis and resource distribution through plant canopies. Plant, Cell

and Environment 30(9):1052-1071.

Nilsen ET, Bao Y (1990) The influence of water stress on stem and leaf photosynthesis in

Glycine max and *Sparteum junceum* (Leguminosae). American Journal of Botany

77(8):1007-1015.

Pereira JS (1995) Gas exchange and growth. In: Ecophysiology of photosynthesis. Springer, pp

147–181.

Pfanz H, Aschan G, Langenfeld-Heyser R, Wittman C, Loose M (2002) Ecology and

ecophysiology of tree stems: corticular and wood photosynthesis. Naturwissenschaften

89:147-162.

Pieters GA, Van Den Noort ME, Nijkerken V (1998) Growth adaptation of leaves and internodes

of popular to irradiance, day length and temperature. Tree Physiology 19:933-942.

Pinheiro JC, Bates D, DebRoy S, Sarkar D, R Core Team (2014) nlme: Linear and nonlinear

mixed effects models. R package version 3,1-122, http://CRAN,R-project.org/package=nlme.

Poorter L, Rozendall DMA (2008) Leaf size and leaf display of thirty-eight tropical tree species.

Physiological Ecology 158:35-46.

R Core Team (2019). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

URL https://www.R-project.org/.

Rao, M. J., Ding, F., Wang, N., Deng, X., & Xu, Q. (2018). Metabolic mechanisms of host

species against citrus huanglongbing (greening disease). *Critical Reviews in Plant*

Sciences, *37*(6), 496-511.

Rasowo BA, Khamis FM, Mohamed SA, Ajene IJ, Aidoo OF, Ombura L, Setamou M, Ekesi S,

Borgemeister C (2019) African citrus greening disease in east africa: incidence, severity, and

distribution patterns. Journal of Economic Enotomology 112(5):2389-2397.

Robertson MJ (1994) Relationships between internode elongation, plant height and leaf

appearance in maize. Field Crop Research 38(3):135-145.

Sagaram M, Burns JK (2009) Leaf chlorophyll fluorescence parameters and huanglongbing. J

Am Soc Hortic Sci 134:194–201.

Schaffer AA, Liu K, Goldschmidt EE, Boyer CD, Goren R (1986) Citrus leaf chlorosis induced

by sink removal: starch, nitrogen, and chloroplast ultrastructure. Journal of Plant Physiology

124 (1-2):111-121.

Schmidt J, Batic F, Pfanz H (2000) Photosynthetic performance of leaves and twigs of evergreen

holly (*Ilex aquifolium* L.) Phyton 40:179-190.

Schneider H (1968) Anatomy of greening diseased sweet orange shoots. Phytopathology

58:1155-1160.

Shafqat W, Jaskani MJ, Maqbool R, Chattha WS, Ali Z, Naqvi SA, Haider MS, Khan IA,

Vincent CI (2021) Heat shock protein and aquaporin expression enhance water conserving

behavior of citrus under water deficits and high temperature conditions. Environmental and

Experimental Botany 181:104270.

Sinclair TR, Rufty TW, Lewis RS (2019) Increasing photosynthesis: unlikely solution for world

food problem. Trends in plant science 24:1032–1039.

Singerman A, Useche P (2016) Impact of citrus greening on citrus operations in Florida.

University of Florida IFAS Extension #FE-983.

Slinski S (2016) Evaluation of HLB treatments. Citrus Industry August 2016:10-12.

Snyder-Leiby TE, Wang S (2008) Role of crop load in chloroplast ultrastructure and zonal

chlorosis, a physiological disorder in 'Honeycrisp' apple trees. Horticultural Science

43(6):1819-1822.

Solhaug KA, Gauslaa Y, Haugen J (1995) Adverse effect of epiphytic crustose lichens upon stem

photosynthesis and chlorophyll of *Populus tremula* L. Botanica Acta 108:233-239.

Stover E, Inch S, Richardson ML, Hall DG (2016) Conventional citrus of some scion/rootstock

combinations show field tolerance under high huanglongbing disease pressure. HortScience

51:127–132.

Stover E, McCollum G, Ramos J, Shatters Jr RG (2014) Growth, health and Liberibacter

asiaticus titer in diverse citrus scions on mandarin versus trifoliate hybrid rootstocks in a field

planting with severe huanglongbing. In: Proceedings of the Florida State Horticultural

Society.pp 53–59.

Syvertsen JP (1994) Partial shoot removal increases net CO_2 assimilation and alters water

relations of Citrus seedlings. Tree Physiology 14:497–508.

Tang L, Chhajed S, Vashisth T (2019) Preharvest fruit drop in huanglongbing-affected

'Valencia' sweet orange. J. Amer. Soc. Hort. Sci. 144(2):107-117.

Tinoco-Ojanguren C, Pearcy RW (1992) Dynamic stomatal behavior and its role in carbon gain

during lightflecks of a gap phase and an understory Piper species acclimated to high and low

light. Oecologia 92:222–228.

Tinoco-Ojanguren C, Pearcy RW (1993) Stomatal dynamics and its importance to carbon gain in

two rainforest Piper species I. VPD effects on the transient stomatal response to lightflecks.

Oecologia 94:388–394.

Tomeo, N. (2017). AQ curves example. GitHub Repository.

https://github.com/Tomeopaste/AQ_curves/blob/master/Introduction_to_AQ_curve_fitting.R

md

Tsagarakis AE, Rogers ME, Spann TM (2012) Applications of plant growth regulators to

container-grown citrus trees affect the biology and behavior of the Asian citrus psyllid.

Journal of the American Society of Horticultural Science 137(1):3-10.

Vasconcelos FN, Li J, Pang Z, Vincent C, Wang N (2021) The total population size of

'Candidatus Liberibacter asiaticus' inside the phloem of citrus trees and the corresponding

metabolic burden related to huanglongbing disease development. Phytopathology: Sep

7:PHYTO-09.

Vu JCV, Yelenosky G (1988) Biomass and photosynthesis response of rough lemon (Citrus

jambhiri Lush.) to Growth Regulators. Canadian Journal of Plant Sciences 68:261-266.

Vu JCV, Yelenosky G, Bausher MG (1986) CO_2 exchange rate, stomatal conductance, and

transpiration in attached leaves of valencia orange. HortScience 21:143-144.

Wallace A, Zidan ZI, Mueller RT, North CN (1954) Translocation of nitrogen in citrus trees.

Proceedings of American Society of Horticultural Science 64:87-104.

Welker S, Pierre M, Santiago JP, Dutt M, Vincent C, Levy A (2021) Phloem transport limitation

in huanglongbing affected sweet orange is dependent on phloem-limited bacteria and callose.

Tree Physiology (en presse).

Whitaker DC, Giurcanu MC, Young LJ, Gonzalez P, Etxeberria E, Roberts P, Hendricks K,

Roman F (2014) Starch content of citrus leaves permits diagnosis of huanglongbing in the

warm season but not cool season. HortScience 49(6):757-762.

Wohlfahrt G, Gu L (2015) The many meanings of gross photosynthesis and their implication for

photosynthesis research from leaf to globe. Plant, cell & environment 38:2500.

Yanhong T, Hiroshi K, Mitsumasa S, Izumi W (1994) Characteristics of transient photosynthesis

in Quercus serrata seedlings grown under lightfleck and constant light regimes. Oecologia

100:463–469.

Zambrosi FCB, Mattos D, Syvertsen JP (2011) Plant Growth, leaf photosynthesis, and nutrient-

use efficiency of citrus rootstocks decreased with phosphite supply. Journal of Plant

Nutrition and Soil Science 174(3):487-495.

Zait Y, Shtein I, Schwartz A (2019) Long-term acclimation to drought, salinity and temperature

in the thermophilic tree *Ziziphus spina-christi*: revealing different tradeoffs between

mesophyll and stomatal conductance. Tree Physiology 39:701–716.

Zheng Z, Chen J, Deng X (2018) Historical perspectives, management and current research of

citrus HLB in Guangdong of China, where the disease has been endemic for over a hundred

years. Phytopathology 108(11):1224-1236.

List of figures

Figure 1. Illustration of the use of a whole-shoot chamber to estimate apparent photosynthesis

(A_{app}) of whole shoots, leaves, and stem.

Figure 2. Apparent assimilation (net assimilation – dark respiration) for 'Valencia' sweet orange

(*Citrus x sinensis* [L.] Osbeck) leaf, stem and shoot. Letters within each plot denote means that

are significantly different at P≤0.05 according to Fisher's least significant differences.

Figure 3. Distinct carbon assimilation light response curves of 'Valencia' sweet orange (*Citrus x*

sinensis [L.] Osbeck) leaf (dashed line) and shoot (dotted line). All points are biological

replicates.

Figure 4. Relationship of log transformed bacterial copy count and log transformed starch

content of *Candidatus* Liberibacter asiaticus-infected and uninfected 'Valencia' sweet orange

(*Citrus x sinensis* [L.] Osbeck) A) leaves and B) shoots.

Figure 5. Relationship of starch and shoot surface area of *Candidatus* Liberibacter asiaticus-

infected and uninfected 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck). Data point

shape and color denote *C*Las-infection exposure category: Low – 1 year *C*Las-infected Asian

citrus psyllid feeding and breeding, Mid – 1.5 years *C*Las-infected Asian citrus psyllid feeding

and breeding, High – 2.5 years *C*Las-infected Asian citrus psyllid feeding and breeding.

Figure 6. Relationship of average internode length and net CO_2 assimilation (*A*) of *Candidatus*

Liberibacter asiaticus-infected and uninfected 'Valencia' sweet orange (*Citrus x sinensis* [L.]

Osbeck).

Figure 7. Illustration of morphological changes in citrus (*Citrus x sinensis*) induced by

huanglongbing (HLB) disease caused by infection with *Candidatus* Liberibacter asiaticus.

Figure 8. Leaf- (A) and shoot- (B) level relationships of conductance with net CO_2 assimilation

(A_{net}) of healthy and *Candidatus* Liberibacter asiaticus-infected 'Valencia' sweet orange (*Citrus*

x sinensis [L.] Osbeck) exposed to 1 (Low), 1.5 (Mid), and 2.5 (High) years of inoculation via

Asian citrus psyllid (*Diaphorina citri* Kuwayama) feeding and reproduction. g_s is leaf-level

stomatal conductance to water. g is shoot conductance to water. For both models $P < 0.0001$. For

leaf-level data $R^2=0.57$. For shoot level data $R^2=0.80$. The interaction term of g_s by infection

category $P=0.002$; the interaction term of shoot g by infection category $P=0.039$. Slope values

labeled with different letters are different at $P<0.05$ within leaf- or shoot-level values.

List of Tables

Table 1. Percent contribution of leaves and stems to the shoot surface area and apparent CO_2

assimilation (A_{app}) in 'Valencia' sweet orange (*Citrus x sinensis* [L.] Osbeck).

Measurement basis	Leaf	Stem
Surface area	94.3 ± 1.6 %	5.6 ± 1.6 %
Mass	83.0 ± 4.8%	17.0 ± 4.8%
Per shoot A_{app}	89.2 ± 8.5 %	10.7 ± 8.5%

Table 2. Analysis of variance and means comparison of parameters of a light response curve

function fit to leaf and whole shoot light response curves of 'Valencia' sweet orange (*Citrus x*

sinensis [L.] Osbeck). Values are means ± standard error. Values followed by different letters are

significantly different at P<0.05. A_{sat}: light-saturated net CO_2 assimilation (μmol CO_2 m^{-2} s^{-1});

Φ:quantum yield (mols CO_2 fixed:mol photons absorbed) ; R_d: dark respiration (μmol CO_2 m^{-2} s^{-}

^{1}); LCP: light compensation point (μmol photosynthetically active radiation m^{-2} s^{-1}); Qsat75:

photosynthetic flux density at 75% saturation (μmol photosynthetically active radiation m^{-2} s^{-1}).

Function parameter	Leaf	Shoot	F-value	*P*-value
A_{sat}	7.7 ± 0.46 b	10.4 ± 0.74 a	8.7	0.01
Φ	0.044 ± 0.002	0.053 ± 0.02	0.2	0.65
R_d	1.2 ± 0.12 a	0.35 ± 0.04 b	42.4	<0.0001
LCP	28.0 ± 8.4 a	14.7 ± 10.8 b	7.9	0.01
Q_{sat75}	528 ± 80 b	1532 ± 226 a	17.3	0.0016

Table 3. Effect of huanglongbing symptom category on leaf-level net CO_2 assimilation (A_{net}),

stomatal conductance (g_s), transpiration (E), cycle threshold (Ct) and Copy # count of

Candidatus Liberibacter asiaticus, foliar starch content, and visual disease rating in 'Valencia'

sweet orange (*Citrus x sinensis* [L.] Osbeck). *P*-value of analysis of variance with infection

categories as the fixed effect. Means (± standard error) followed by different letters are

significantly different at *P*<0.05 according to Fisher's LSD.

	Uninfected	Low	Medium	High	*P*-value
A_{net} (μmol m^{-2} s^{-1})	3.97 ± 1.73 a	1.91 ± 1.24 b	2.61 ± 1.62 b	2.70 ± 1.75 b	0.0003
g_s (mol m^{-2} s^{-1})	0.045 ± 0.007 ab	0.026 ± 0.0.004 b	0.033 ± 0.005 b	0.057 ± 0.011a	0.013
E (mmol m^{-2} s^{-1})	0.94 ± 0.16 a	0.59 ± 0.07 b	0.69 ± 0.11ab	0.89 ± 0.15 a	0.058
qPCR Ct	37.87 ± 2.35 a	24.69 ± 4.14 b	24.16 ± 3.13 b	24.10 ± 2.52 b	<0.00001
qPCR Copy # (log)	2.60 ± 2.65 b	11.78 ± 2.90 a	12.27 ± 1.93 a	12.30 ± 1.66 a	<0.00001
Starch (μg mm^{-2})	1.08 ± 1.27 d	11.31 ± 4.30 a	5.69 ± 4.39 b	3.55 ± 3.17 c	<0.00001
Visual Rating (0-10)	0.22 ± 0.60 c	1.26 ± 0.98 b	2.11 ± 1.53 a	2.22 ± 1.42 a	<0.00001

903 Table 4. Effect of huanglongbing symptom category on shoot-level net CO_2 assimilation (A_{net}),

904 shoot conductance (g), transpiration (E), and shoot morphological variables in 'Valencia' sweet

905 orange (*Citrus x sinensis* [L.] Osbeck). *P*-value of analysis of variance with infection categories

906 as the fixed effect. Means (± standard error) followed by different letters are significantly

907 different at *P*<0.05 according to Fisher's LSD.

	Uninfected	Low	Medium	High	P-value
A_{net} (μmol m^{-2} s^{-1})	1.91 ± 1.07 b	2.42 ± 0.88 b	2.44 ± 0.52 ab	3.37 ± 1.34 a	0.032
g (mol m^{-2} s^{-1})	0.03 ± 0.005 b	0.05 ± 0.007 a	0.044 ± 0.005 ab	0.059 ± 0.008 a	0.039
E (mmol m^{-2} s^{-1})	0.45 ± 0.09 b	0.72 ± 0.1 ab	0.65 ± 0.08 ab	0.85 ± 0.11 a	0.061
Stem length (mm)	112.50 ± 22.5	112.30 ± 19.7	91.20 ± 26.1	90.30 ± 30.6	0.13
Mean leaf area (cm^2)	15.79 ± 3.39 ab	17.59 ± 5.86 a	15.40 ± 6.33 ab	11.38 ± 2.08 b	0.054
Shoot surface area (cm^2)	117.40 ± 42.0 a	133.50 ± 42.8 a	100.40 ± 48.8 ab	68.30 ± 27.1 b	0.014
Leaves per shoot	6.75 ± 1.49	7.22 ± 1.48	6.25 ± 2.25	5.33 ± 1.58	0.15

| Internode length (mm) | 17.41 ± 5.49 | 15.99 ± 3.92 | 15.33 ± 3.99 | 16.99 ± 2.67 | 0.74 |

Table 5. Linear regressions of effects of gene copy number of *Candidatus* Liberibacter asiaticus

on foliar starch content, of internode length on net CO_2 assimilation, and of foliar starch content

on shoot surface area each assessed across three huanglongbing symptom categories including or

not including a fourth category of uninfected trees, and measured at leaf or whole shoot levels.

P-values below 0.05 are labeled with asterisks. All models included shoot nested within tree, and

leaf models included leaf nested within shoot, as random variables.

Dependent variable	Including uninfected	Tissue	Effect	F-value	*P*-value
Starch content	yes	leaf	Bacterial copy number	52.2130	<0.00001*
		shoot	Bacterial copy number	33.1526	<0.00001*
Net CO_2 assimilation	yes	shoot	Internode length	1.3619	0.2521
	no	shoot	Internode length	10.7940	0.0032*
Shoot surface area	yes	shoot	Starch content	1.7849	0.1913
	no	shoot	Starch content	4.3675	0.0479*

918