Title: Gene-Specific Nonsense-Mediated mRNA Decay Targeting for Cystic Fibrosis Therapy

Authors: Young Jin Kim[1, 2, 3], Tomoki Nomakuchi [1, 3, 4], Foteini Papaleonidopoulou [1,5], Adrian R.

Krainer[1, *].

[1] Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, USA.

[2] Department of Genetics, Stony Brook University, Stony Brook, New York, USA.

[3] Medical Scientist Training Program, Stony Brook University School of Medicine, Stony Brook,

New York, USA.

[4] Present address: Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA.

[5] Present address: Francis Crick Institute, London, UK.

Correspondence to: A.R.K. (krainer@cshl.edu).

<u>Abstract</u>

Low *CFTR* mRNA expression due to nonsense-mediated mRNA decay (NMD) is a major hurdle

in developing a therapy for cystic fibrosis (CF) caused by the *W1282X* mutation in the *CFTR* gene.

CFTR-W1282X truncated protein retains partial function, so increasing its levels by inhibiting NMD

of its mRNA will likely be beneficial. Because NMD regulates the normal expression of many

genes, gene-specific stabilization of *CFTR-W1282X* mRNA expression is more desirable than

general NMD inhibition. Synthetic antisense oligonucleotides (ASOs) designed to prevent binding

of exon junction complexes (EJC) downstream of premature termination codons (PTCs) attenuate

NMD in a gene-specific manner. We developed a cocktail of three ASOs that specifically increases

the expression of *CFTR* W1282X mRNA and CFTR protein in ASO-transfected human bronchial

epithelial cells. This treatment increased the CFTR-mediated chloride current. These results set

the stage for clinical development of an allele-specific therapy for CF caused by the W1282X

mutation.

Introduction

CFTR-W1282X, the 6[th] most common CF-causing mutation, causes a severe form of CF and is present in 1.2% of CF patients worldwide [1], 2.2% of U.S. CF patients [2], and up to 40% of Israeli CF patients [3]. The CFTR-W1282X truncated protein retains partial function [4–6], but is expressed at a very low level, due to nonsense-mediated mRNA decay (NMD). In general, NMD prevents the accumulation of potentially harmful truncated proteins translated from premature termination codon (PTC)-containing mRNAs. However, when NMD reduces the expression of a mutant CFTR protein that has partial activity, it exacerbates the phenotype, so that patients homozygous for the CFTR-W1282X mutation or compound heterozygous for CFTR-W1282X and another CF-causing mutation have poor clinical outcomes. As low as 10% of normal CFTR function provides a significant therapeutic benefit for CF patients [7,8]. Thus, increasing the expression of mutant CFTR protein with residual activity is expected to be beneficial.

NMD is a major hurdle for developing a targeted therapy for CF caused by the CFTR-W1282X mutation. The approval of CFTR correctors that enhance post-translational CFTR processing, and potentiators that improve CFTR channel opening, brought benefit to the majority of CF patients [9]. However, these therapeutic options are not effective against CF caused by CFTR-W1282X, due to the low expression of CFTR-W1282X mRNA. One approach to treat CF caused by this mutation involves read-through compounds (RTCs) that increase the level of full-length protein by reducing the fidelity of the ribosome at the PTC [10]. Gentamicin is a type of RTC that can increase full-length CFTR protein *in vitro,* but its clinical efficacy for various CF nonsense mutations is limited by NMD [11,12]. Likewise, ataluren is another non-aminoglycoside RTC with a very good safety profile, but it did not improve forced expiratory volume (FEV) in CF patients with various nonsense mutations, including W1282X, in clinical trials [13,14].

Preclinical models of CF caused by the W1282X mutation showed that the efficacy of

RTCs can be increased *in vivo* by knocking down key components of the NMD pathway [15].

However, a clinically viable NMD-suppression approach does not exist yet. CFTR potentiators

and correctors such as ivacaftor (VX-770) and lumacaftor (VX-809) enhance CFTR-W1282X

activity *in vitro*, and may potentially benefit patients with the W1282X mutation, but they are not

effective if the truncated protein expression is too low [4–6,16]. Thus, there is a pressing need for

strategies to overcome NMD of the *CFTR-W1282X* mRNA.

Several NMD-suppression strategies have been developed for potential application to

diseases caused by NMD-sensitive nonsense mutations. These include inhibition of NMD by

small-molecule inhibitors [17–19] or knockdown of key NMD factors [20, 19]. However, global inhibition

of NMD may be detrimental, because the NMD machinery targets a subset of normal and

physiologically functional mRNA isoforms, thereby post-transcriptionally regulating gene

expression [21]. Therefore, global inhibition of NMD could disrupt mRNA homeostasis in a broad

range of tissues [15].

NMD is strongly dependent on a complex of RNA-binding proteins called the exon junction

complex (EJC). In contrast to many RNA-binding proteins, EJCs bind mRNA in a position-

dependent, sequence-independent manner [22–25]. More than 80% of EJCs are positioned 20~24

nucleotides (nt) upstream of an exon-exon junction [22–26]. Normal stop codons are typically in the

last exon, whereas many PTCs are upstream of one or more exon-exon junctions, and thus are

upstream of at least one EJC [27]. The '55-nt rule' predicts that mRNAs with a PTC > 55 nt upstream

of the last exon-exon junction are degraded by NMD, reflecting the footprint of the stalled

ribosome [23]. A downstream EJC interacts with the ribosome stalled at the PTC, and recruits NMD

factors to form a degradation complex that promotes decapping, deadenylation, and

endocleavage of the target mRNA, which is subsequently degraded [28]. Thus, the EJC is a major

enhancer of NMD.

Disrupting the downstream EJC association with PTC-containing mRNA can be used to

inhibit NMD [29]. Uniformly 2'-*O*-(2-methoxyethyl) (MOE)-modified antisense oligonucleotides

(ASOs) can stably hybridize to complementary RNAs without triggering RNase H-mediated

degradation [30]. Such ASOs are effective tools for disrupting the interaction between an RNA and

its binding proteins, and can be used to alter mRNA processing and translation *in vitro* and *in vivo*

[30–34]. We previously developed ASOs that target presumptive downstream EJC sites of PTC-

containing mRNAs. These ASOs efficiently attenuate NMD of their target genes, restoring mRNA

and proteins levels [29]. In the present study, we demonstrate that a cocktail of three ASOs targeting

presumptive downstream EJC binding sites specifically increases the expression of endogenous

CFTR-W1282X mRNA in human bronchial epithelial (HBE) cells. Furthermore, the ASO cocktail

increases partially active CFTR protein and CFTR-mediated chloride current in HBE cells. These

results set the stage for the clinical development of an allele-specific therapy for CF caused by

the *W1282X* mutation.

<u>Results</u>

CFTR-W1282X mRNA has four downstream exon-exon junctions, on exons 23-24, 24-25,

25-26, and 26-27. However, the predicted EJC binding site on exon 23 is approximately 5 nt

downstream of the PTC—within the ribosome footprint—and thus only three exons (24, 25, and

26) are predicted to harbor EJCs that can induce NMD. To investigate the impact of each EJC on

NMD of *CFTR-W1282X* mRNA, we generated U2OS cells stably expressing doxycycline-

inducible *CFTR*-minigene NMD reporters, each with only one presumptive downstream EJC site

(Figure 1A, Supplementary Fig.1A). These NMD reporters are three-exon minigenes downstream

of GFP, and comprise *CFTR* cDNA sequence for exons 22-27 and intervening sequences (IVSs)

that are shortened natural *CFTR* introns.

The reporters showed some intron retention, and *pW1282X-IVS23* generated an

additional isoform by use of an alternative 3' splice site (3'ss; based on size and motif predictions)

(Supplementary Fig.1B-F). The '55-nt rule' predicts that an exon-exon junction <55 nt downstream

of the PTC does not induce NMD [23]. As mentioned above, only the EJCs on exon 24, 25, and 26

are predicted to induce NMD. Inhibiting NMD with cycloheximide increased the mRNA levels of

the reporters harboring the PTC and intron 24, 25, or 26 (*pW1282X-IVS24, pW1282X-IVS25,* or

pW1282X-IVS26) (Supplementary Fig.1D-F); conversely, the reporters harboring the PTC and

intron 23 (*pW1282X-IVS23*) or harboring intron 24 but not the PTC (*pWT-IVS24*) were not

sensitive to cycloheximide (Supplementary Fig.1B-C).

Uniformly 2'-*O*-(2-methoxyethyl) (MOE)-modified ASOs can stably hybridize to complementary

mRNAs without inducing RNAse-H-mediated degradation, and modulate their posttranscriptional

processing, including NMD [31]. Using our previously described ASO screening strategy for gene-

specific antisense inhibition of NMD--dubbed "GAIN" [29]—we designed sets of 19 overlapping

15mer ASOs to target each of the presumptive EJC binding sites on the NMD reporters *pW1282X-*

IVS24, pW1282X-IVS25, and *pW1282X-IVS26*, respectively (Fig. 1B). We screened a total of 57

ASOs, uniformly modified with MOE ribose and a phosphorothioate (P=S) backbone. ASOs H24,

H26, and M33 are uniformly MOE and P=S modified negative-control ASOs that are not

complementary to any gene expressed in the reporter-expressing cells [29]. Based on the screen,

we chose C478 and C515 as the initial lead ASOs targeting exons 24 and 26, respectively (Fig.

1C-D). Because the retention of IVS25 in *pW1282X-IVS25* caused by some ASOs prevented a

clear assessment of NMD inhibition by the screened ASOs (Supplementary Fig. 2A), we

generated a new *pW1282X-IVS25* NMD reporter with a stronger 5' splice site (5'ss) but the same

amino acid sequence (Supplementary Fig. 2B). Among several ASOs that increased the new

NMD reporter levels, we chose C495 as the lead ASO (Fig. 1E). The candidate ASOs inhibited

NMD of the reporters in a dose-dependent manner (Supplementary Fig. 2C-H).

Endogenous *CFTR* mRNA is targeted for NMD in human bronchial epithelial cells and

colon cancer cells harboring the homozygous *CFTR-W1282X* mutation (16HBE-W1282X and

DLD1-W1282X cells) (Fig. 1F). We used two negative-control treatments: i) a scrambled-

sequence ASO based on C494; and ii) a cocktail composed of C488+C507+C526 ASOs, which

did not stabilize the NMD reporters. We first tested a lead GAIN ASO cocktail composed of

C478+C495+C515, based on the above NMD-reporter screening results. Compared to the

negative-control ASOs, the C478+C495+C515 ASO cocktail significantly increased *CFTR* mRNA

levels in both 16HBE-W1282X and DLD1-W1282X cells (Fig. 1G, Supplementary Fig. 3).

Length is an important parameter in ASO design that can affect the efficacy and specificity

of uniformly modified ASOs [35–37]. Based on the results of the above 15mer ASO screens, we

designed a new 18mer-ASO cocktail and an 18mer scramble-ASO control. Transfection of the

scramble ASO did not increase *CFTR-W1282X* mRNA levels in 16HBE-W1282X cells, whereas

the 18mer ASO cocktail C24+C25+C26 increased *CFTR-W1282X* mRNA levels in a dose-

dependent manner (Fig. 1H). The 18mer ASO cocktail did not affect two other endogenous NMD-

sensitive mRNAs, *eIF4A2* and *SRSF2* (Supplementary Fig. 4). Thus, the lead GAIN ASO 18mer

cocktail inhibited NMD of *CFTR-W1282X* mRNA in a gene-specific manner.

Based on the presumptive mechanism of action of the ASO cocktails, they should not

affect the total mRNA levels of wild-type (WT) or missense-mutant *CFTR* mRNAs that are not

sensitive to NMD (Fig. 1I). Indeed, *CFTR* mRNA levels were insensitive to transfection of the

control ASO or C478+C495+C515 ASO cocktail in DLD1-WT, 16HBE-F508del, and 16HBE-

G551D cells (Fig. 1J). Thus, the C478+C495+C515 ASO cocktail allele-specifically increased the

mRNA levels of nonsense-mutant *CFTR-W1282X* mRNA. To test whether C478+C495+C515

inhibits NMD of *CFTR-W1282X* mRNA specifically, as opposed to somehow affecting global NMD,

we used RT-qPCR to survey seven other endogenous NMD-sensitive transcripts that are

upregulated upon NMD inhibition by cycloheximide treatment [38,39]. As expected,

C478+C495+C515 increased *CFTR-W1282X* mRNA, without significantly changing any of the

other NMD-sensitive mRNAs (Fig. 1K).

To identify the optimal GAIN ASO cocktail, we performed a more comprehensive screening

in 16HBE-W1282X cells (Fig. 2A-C). To systematically screen ASOs targeting each exon, we

tested cocktails composed of two constant ASOs and one varying ASO. For example, to screen

exon-24-targeting ASOs, we tested 19 cocktails composed of varying exon-24-targeting ASOs

and the same two ASOs targeting exons 25 and 26. After testing 57 such ASO cocktails, we

identified C478+C494+C514 as the new lead GAIN ASO cocktail, which was more potent than

C478+C495+C515 (Fig. 2D). At the highest concentration tested, both ASO cocktails increased

CFTR-W1282X mRNA similarly, but at a lower concentration, the new lead cocktail had

significantly higher potency.

G542X and *R1162X CFTR* mutations are on exons 12 and 22, respectively. As these

mutant mRNAs harbor more than three EJCs downstream of the premature termination codon,

we tested whether their levels would be insensitive to the lead ASO cocktail C478+C494+C514.

We transfected 16HBE-G542X and 16HBE-R1162X cells harboring homozygous *G542X* and

R1162X mutations, respectively, with control ASOs or the lead ASO cocktail C478+C494+C514

(Fig. 2E and F). Compared to the no-treatment control, transfection of 120 nM scramble ASO,

control ASO cocktail, or C478+C494+C514 did not affect the levels of *CFTR-G542X, CFTR-*

R1162X, and NMD-sensitive *eIF4A2* mRNAs. Only cycloheximide treatment caused significant

increases in these NMD-sensitive mRNA levels. These results are consistent with the EJC-centric

model of NMD, according to which at least one EJC >55nt downstream of a PTC is sufficient to

induce strong NMD [23].

Using various combinations of the NMD-inhibiting ASOs, we next tested whether all three

presumptive downstream EJC binding sites on *CFTR-W1282X* mRNA must be targeted with the

corresponding ASOs for effective mRNA stabilization. Targeting only one or two EJC binding sites

with the respective lead ASOs partially stabilized the *CFTR-W1282X* mRNA, but the most

significant and efficient increase in *CFTR-W1282X* mRNA was obtained by simultaneous

transfection of all three lead ASOs, in both 16HBE-W1282X and DLD1-W1282X cells (Fig. 2G,

Supplementary Fig. 5). Because ASO cocktails composed of one or two ASOs elicited a small

increase in *CFTR-W1282X* mRNA levels, we next asked whether certain presumptive EJC

binding sites may be more important than others. To test this possibility, we transfected 16HBE-

W1282X cells with ASO cocktails with varying ratios of the individual ASOs (Fig. 2H). The

C478+C495+C515 cocktail with the highest total ASO concentration and an equimolar ratio of

40:40:40 nM increased *CFTR-W1282X* mRNA the most, and the increase was dependent on the

total ASO concentration. In general, limiting the concentration of C495 in the cocktail reduced the

CFTR-W1282X mRNA levels to the greatest extent (Supplementary Fig. 6A-C). Interestingly, ASO

cocktails with equal total concentrations, but different ASO ratios, did not have equivalent effects

on *CFTR-W1282X* mRNA levels. Also, some ASO cocktails increased *CFTR-W1282X* mRNA

similarly or more than others, despite their lower total ASO concentration. The differences among

CFTR-W1282X mRNA expression changes caused by the various ASO cocktails may be

attributable to various factors, including partial EJC occupancy on different exons, differences in

ASO uptake and target accessibility or affinity, involvement of EJC-independent NMD pathways,

and RNA secondary structure [40,41].

Some EJC-targeting ASOs may affect splicing, if their binding site overlaps with cis-

elements that regulate splicing. We monitored exon 24-26 splicing by RT-PCR in DLD1-WT cells

transfected with ASOs (Supplementary Fig. 7A). Exon-26-targeting ASOs did not detectably

disrupt *CFTR* mRNA splicing. On the other hand, all exon-25-targeting ASOs caused slight exon-

25 skipping, and some exon-24-targeting ASOs caused substantial exon-24 skipping. As

disrupted binding of serine-rich (SR) proteins to exonic splicing enhancers (ESEs) by uniformly

modified MOEPS ASOs can cause exon skipping [42–45], we used ESEfinder [46] to identify putative

ESEs on *CFTR* exons 24-26 that might be blocked by the lead ASOs (Supplementary Fig. 7B).

mRNA sequences complementary to the lead ASOs C478 and C494 overlap with SR protein

motifs; however, overlap with an SR protein motif is insufficient to predict an ASO's interference

with splicing.

Transfection of the 15-mer or 18-mer lead ASOs caused dose-dependent, multiple exon

skipping in human bronchial cells (Fig. 2I, Supplementary Fig. 8A-F): single (exon 24 and 25

skipping: d24 and d25), double (exon 24-25 skipping: d24-25), and triple exon skipping (exon 24-

25-26 skipping: d24-25-26) of *CFTR* mRNA. Similar splicing changes occurred with the 15mer or

18mer lead ASO cocktail treatment by free uptake (Supplementary Fig. 9A-B). The splicing

changes were not cell-line-specific, as the lead ASO cocktail promoted similar splicing changes

in 16HBE-W1282X and DLD-W1282X cells (Supplementary Fig. 10A-B).

Control ASO cocktails caused a smaller degree of exon 24 and 25 skipping, consistent

with the results in DLD1-WT cells (Fig. 2I). The 15mer and 18mer lead ASO cocktails generated

the same *CFTR* isoforms, but with varying degrees of percent-spliced-in (PSI); for example, the

PSI of the exon 24-25 double-skipping event was higher in 16HBE-W1282X cells treated with the

18mer lead cocktail (Supplementary Fig. 8C-D and 9B). Interestingly, whereas the 15mer or

18mer exon-26-targeting ASOs alone did not cause exon 26 skipping, the ASO cocktails

containing exon-24-targeting ASO caused the appearance of the triple-skipped isoform (d24-25-

26) (Fig. 2I, Supplementary Fig. 8C-D). To search for potential off-target sites on *CFTR* pre-mRNA,

we looked for sites complementary to C478 with a maximum of four nucleotide mismatches,

downstream of exon 24, and found only one site with four mismatches in intron 23. The chance

of finding an off-target with ≥ 4-nt mismatches is very low [47], suggesting that C478 is unlikely to

cause exon 25 and 26 skipping by binding to ESEs in these exons. These results suggest that an

ESE and/or the EJC in exon 24 is involved in long-range splicing regulation. Recent studies

showed that EJCs help maintain faithful splicing transcriptome-wide [48–54].

Despite the splicing alterations, the increase in the total *CFTR* mRNA by the lead ASO

cocktail resulted in increased CFTR-W1282X protein levels, compared to the scramble ASO

control (Fig. 3A-B). This result was expected, because none of the splicing changes affect the

reading frame upstream of the nonsense mutation in exon 23. Combining the lead ASO cocktail

with lumacaftor (VX-809), a corrector that improves the folding of CFTR protein [5,55], further

increased the total CFTR-W1282X protein levels (Fig. 3A-B). Three different patterns of CFTR

bands are visible on a Western blot: non-glycosylated A-band, core glycosylated B-band, and fully

mature, glycosylated C-band [56]. As shown previously [5], truncated CFTR-W1282X exists as core-

glycosylated and mature glycosylated forms (Supplementary Fig. 11A). Extensive enzymatic

deglycosylation revealed that truncated-core and fully-mature-glycosylated CFTR-W1282X

proteins are upregulated by transfection of the lead ASO cocktail in 16HBE-W1282X cells

(Supplementary Fig. 11B and C).

We next measured CFTR function in 16HBE-W1282X cells treated with the lead GAIN

ASO cocktail, with the Ussing-chamber assay [57]. FDA-approved CFTR potentiators and

correctors, such as ivacaftor (VX-770) and lumacaftor (VX-809), can enhance CFTR-W1282X

activity *in vitro*, and may potentially benefit patients with the W1282X mutation [4–6,16]. Thus, we

combined all ASO treatments for Ussing-chamber assays with VX-809 and VX-770. The 16HBE-

W1282X cells treated with the lead 15mer ASO cocktail (C478+C494+C514) showed increased

CFTR-mediated chloride current, compared to 16HBE-W1282X cells treated with scramble 15mer

ASO (Fig. 3C-E). We verified that the 16HBE-W1282X cells transfected with the lead ASO cocktail

while cultured on transwell plates for the Ussing chamber assay showed gene-specific increase

in *CFTR-W1282X* mRNA levels (Fig. 3F). Similar to the 15mer lead ASO cocktail, the 18mer lead

ASO cocktail (C24+C25+C26) significantly increased CFTR function, compared to the control

18mer scramble ASO treatment (Fig. 3G-L). This result demonstrates for the first time that gene-

specific NMD inhibition of a hypomorphic *CFTR* allele leads to an increase in CFTR-mediated

chloride current. All *CFTR-W1282X* mRNA isoforms generated by the ASO cocktail treatment

presumably terminate at the *W1282X* codon, but the contribution of each isoform to CFTR activity

may be affected by various factors, including mRNA stability, transport to the cytoplasm, and

translational efficiency.G418 is an aminoglycoside antibiotic that at high concentrations induces

translational read-through of reporters containing *CFTR* nonsense mutations, and increases

truncated CFTR-W1282X protein levels by NMD inhibition [18,58,59]. Indeed, 400 μM (0.2 mg/mL)

G418 alone increased truncated CFTR-W1282X protein levels (Supplementary Fig. 11D-E), and

200 μM (0.1 mg/mL) increased CFTR activity in 16HBE-W1282X cells (Fig. 3I and L). However,

full-length CFTR reflecting read-through activity remained below the level of detection by Western

blotting (Supplementary Fig. 11C-D). Combining 200 μM or 600 μM G418 with the lead ASO

cocktail further increased CFTR-W1282X function, compared to the respective lead ASO cocktail

treatment alone (Fig. 3J-L). This result suggests that some level of read-through at the *W1282X*

codon may occur.

Discussion

NMD severely limits the therapeutic development for CF caused by *CFTR-W1282X*

mutation. Global NMD suppression can be achieved by targeting key NMD factors by gap-mer

ASOs that induce gene knockdown or small molecules that inhibit the activity of NMD factors [17–

20]. However, targeted NMD suppression may be more desirable as it can avoid unwanted side-

effects that can be caused by non-specific NMD inhibition. Here, we demonstrate for the first time

that gene- and allele-specific NMD suppression using EJC-targeting ASO cocktails increases

truncated CFTR-W1282X protein, as well as CFTR function.

The lead GAIN ASO cocktails inhibited NMD of *CFTR-W1282X* mRNA by targeting

presumptive EJC binding sites downstream of the PTC, independently of cell type. Consistent

with the EJC-centric model of NMD [40], we achieved efficient NMD suppression only when all

downstream EJC binding sites that contribute to NMD (i.e., those on exons 24-26) were targeted

by ASOs. The lead ASO cocktails did not inhibit NMD of other endogenous NMD-sensitive

transcripts we tested, or affect the levels of NMD-insensitive *CFTR* mRNA. Likewise, the lead

ASO cocktails did not affect *CFTR* mRNA levels in cells harboring PTCs upstream of exon 23.

Thus, our results demonstrate that the lead ASO cocktails inhibit NMD of *CFTR-W1282X* mRNA

by preventing the binding of EJCs located downstream of the PTC and beyond the footprint of the

stalled ribosome. These observations rule out the possibility of global NMD suppression due to

ASO treatment, or *CFTR* mRNA stabilization by inhibition of other mRNA-degradation pathways.

The lead 15-mer and 18-mer cocktails achieved similar levels of gene-specific NMD

suppression, but had different effects on splicing. Compared to 12-mer ASOs, 18-mer ASOs tend

to have fewer off-target effects on splicing [60], but whether our lead 18-mer ASO cocktail

meaningfully reduces off-target effects and toxicity will require further investigation. We conclude

that rationally designed ASOs can modulate clinically relevant NMD, expanding the current RNA

and oligonucleotide therapeutics toolbox.

<u>Methods</u>

ASOs

All ASOs were uniformly modified with 2'-*O*-(2-methoxyethyl) (MOE) ribose, phosphorothioate

(P=S) linkages, and 5'-methylcytosine. The 15mer ASOs were obtained from Ionis

Pharmaceuticals (Carlsbad, CA) and Integrated DNA Technologies (Coralville, IA), and 18mer

ASOs were obtained from Bio-Synthesis (Lewisville, TX). All ASOs were dissolved in water and

stored at -20 ℃. Stock ASO concentrations were calculated based on the A260 measurement and

each ASO's extinction coefficient *e* (mM^{-1} x cm^{-1} @ 260 nm). The sequences of all ASOs used in

this study are listed in Table 1.

Preparation of U2OS cells expressing NMD reporters

The NMD reporters used for the ASO screening (*pW1282X-IVS23, pW1282X-IVS24, pW1282X-*

IVS25, and *pW1282X-IVS26*) were constructed from the parent NMD reporter *GFP-CFTR22-27-*

T7, which was cloned into the pCDNA5 FRT/TO plasmid (Life Technologies, Carlsbad, CA).

pCDNA5 FRT/TO allows tetracycline-inducible expression of the gene. *GFP-CFTR22-27-T7* has

the natural sequences of exons 22 to 27 of the human *CFTR* gene and shortened intervening

sequences (IVS) modified from the natural sequences of introns 22 to 26 by taking 200

nucleotides (nt) from the 5' and 3' ends of the corresponding introns. GFP and T7 cDNA

sequences were added to the 5' and 3' ends of each reporter, respectively, to facilitate gene,

transcript, and protein detection. NMD reporters *pW1282X-IVS23, pW1282X-IVS24, pW1282X-*

IVS25, and *pW1282X-IVS26* comprise only IVS23, IVS24, IVS25, or IVS26, respectively,

downstream of the PTC. The 5'ss and 3'ss of IVS25 and the 3'ss of IVS23 in *pW1282X-IVS25*

were mutated to stronger splice-site sequences (Supplementary Figure 2B) to promote proper

splicing in the minigene context.

For stable expression of NMD reporters, the reporter plasmids were co-transfected with

the pOG44 helper vector to express Flp recombinase into U2OS-TREx cells harboring a single

FRT recombination site (Life Technologies). Cells with successful NMD-reporter integration were

selected by hygromycin resistance. The expression and splicing of the transgenes were assessed

by radioactive RT-PCR, following induction with 1 µg/ml doxycycline (Research Products

International Corp, D43020-100).

CRISPR mutant DLD1 and 16HBEge cells

Using CRISPR/Cas9, we generated DLD1 cells with homozygous *CFTR-W1282X* mutation.

sgRNA against exon 23 (Table 2) was cloned downstream of the U6 promoter of the

pSpCas9(BB)-2A-GFP (PX458) plasmid (Addgene plasmid # 48138) [61], creating pSC2G-

CFTR23, which allows co-expression of the sgRNA and *Streptococcus pyogenes* Cas9

(spCas9). 4 µg of pSC2G-CFTR23 plasmid was co-transfected with 1 µM single-stranded DNA

repair template (synthesized by Sigma, Table 3) comprising the *CFTR-W1282X* and silent

protospacer adjacent motif (PAM) mutations, using lipofectamine 2000. Following transfection,

GFP+ DLD1 cells were collected using an ARIA-I cell sorter (BD), and individual clones of cells

were isolated by limiting dilution into 96-well plates. Clonal cells were expanded and passaged

until confluent in 6-well plates. We characterized 159 clones by Sanger sequencing, and

identified two heterozygous and two homozygous *W1282X* mutant clones. 16HBE14o- parental

cells gene-edited to yield 16HBEge cell lines CFF-16HBEge CFTR W1282X, F508del, G551D,

G542X, or R1162X, homozygous for *CFTR-W1282X, F508del, G551D, G542X, or R1162X*

mutation, respectively, in the endogenous loci were kindly provided by the Cystic Fibrosis

Foundation's CFFT Lab [18]. Elsewhere in the text, these cells are referred to as 16HBE-W1282X,

16HBE-F508del, 16HBE-G551D, 16HBE-G542X, or 16HBE-R1162X cells, respectively.

Tissue culture and transfection of siRNA, ASO, and plasmids

U2OS and DLD1 cells were cultured in DMEM with 10% FBS. 16HBEge cells were cultured in

MEM with 10% FBS. All cells were incubated at 37 ℃ and 5% CO_2. Cells were transfected with

ASOs and plasmids using Lipofectamine 3000 (Life Technologies, L3000015) according to the

manufacturer's protocol, and harvested 48 hrs post-transfection. Cells were transfected with

siRNA (Table 3) using Lipofectamine RNAiMax (Life Technologies, 13778075) according to the

manufacturer's protocol for transfecting short oligonucleotides, and harvested 48 hrs post-

transfection. NMD inhibition by cycloheximide was performed by treating the cells for 1 hr with

cycloheximide (Sigma, 100 µg/mL). For ASO treatment by free-uptake, 1 mM stock ASO

solutions were diluted into MEM with 10% FBS to the desired final concentrations, and the cells

were cultured for 4 days before harvesting or Ussing-chamber assays. NMD-reporter

expression was induced with 1 µg/ml doxycycline with media change, 6 hr after transfection. For

the G418-treatment group, G418 (Sigma) was added to the culture medium at the indicated final

concentrations, 24 hr before protein extraction or Ussing-chamber assays.

RNA extraction and RT-PCR

Total RNA was extracted with TRIzol (Life Technologies) according to the manufacturer's

protocol. Oligo dT(18)-primed reverse transcription was carried out with ImProm-II Reverse

Transcriptase (Roche). Semi-quantitative radioactive PCR (RT-PCR) was carried out in the

presence of ^{32}P-dCTP with AmpliTaq DNA polymerase (Thermo Fisher), and real-time

quantitative RT-PCR (RT-qPCR) was performed with Power Sybr Green Master Mix (Thermo

Fisher). Primers used for RT-PCR and RT-qPCR are listed in Table 4. RT-PCR products were

separated by 6% native polyacrylamide gel electrophoresis, detected with a Typhoon FLA7000

phosphorimager, and quantitated using MultiGauge v2.3 software (Fujifilm); RT-qPCR data

were quantitated using QuantStudio 6 Flex system.

Protein extraction, deglycosylation, and Western blotting

Cells were harvested with RIPA buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 1% NP40, 0.5%

sodium deoxycholate, and 0.1% SDS) and 2 mM EDTA + protease inhibitor cocktail (Roche) by

sonicating for 5 min at medium power using a Bioruptor (Diagenode), followed by 15-min

incubation on ice. Protein concentration was measured using the Bradford assay (Bio-Rad) with

BSA as a standard. To monitor post-translational maturation of CFTR protein, cell lysates were

incubated in 40 µg/ml PNGase F (New England Biolabs, MA) for 2 hr at 37 °C to cleave all N-

glycans before immunoblotting [62]. Cell lysates were mixed with Laemmli buffer and incubated at

37 ℃ for 30 min. The protein extracts were separated by sodium dodecyl sulfate-polyacrylamide

gel electrophoresis (SDS-PAGE) (6% Tris-chloride gels) and then transferred onto a

nitrocellulose membrane. CFTR bands C, B, and A were detected with antibody UNC-596 (J.

Riordan lab, University of North Carolina, Chapel Hill, NC). The C and B band intensities were

measured together for the quantification of CFTR protein levels. The specificity of the antibody

was confirmed by knocking down *CFTR* in WT DLD1 cells (Supplementary Fig. 11A). Na/K-

ATPase, detected with a specific antibody (Santa Cruz sc-48345), was used as a loading

control. UPF1 was detected with rabbit antibody D15G6 (Cell Signaling Technology #12040S);

mouse anti-α-tubulin antibody (Sigma T9026) was used as a loading control. IRDye 800CW or

700CW secondary antibody (LI-COR) was used for Western blotting, and the blots were imaged

and quantified using an Odyssey Infrared Imaging System (LI-COR). Statistical significance was

calculated using Student's t-test or one-way ANOVA, followed by Tukey's or Dunnett's post-test.

Ussing-chamber assay

Preparing 16HBEge cells for Ussing-chamber assay. 16HBEge cells were grown as an electrically

tight monolayer on Snapwell filter supports (Corning, cat# 3801), as described [63], and both serosal

and mucosal membranes were exposed to the ASOs for 4 days, and to CFTR correctors for 24

hrs, before the assays. The Snapwell inserts were transferred to an Ussing chamber (P2302,

Physiologic Instruments, Inc., San Diego, CA). For 16HBEge cells, the serosal side only was

superfused with 5mL of HB-PS buffer; on the mucosal side, 5 ml of CF-PS was used (137 mM

Na-gluconate; 4 mM KCl; 1.8 mM $CaCl_2$; 1 mM $MgCl_2$; 10 mM HEPES; 10 mM glucose; pH

adjusted to 7.4 with N-methyl-D-glucamine) to create a transepithelial chloride-ion gradient. After

clamping transepithelial voltage to 0 mV, the short-circuit current (I_{SC}) was measured with a

Physiologic Instruments VCC MC6 epithelial voltage clamp, while maintaining the buffer

temperature at 37 °C. Baseline activity was recorded for 20 min before agonists (final

concentrations: 10 µM forskolin (Sigma, F6886), 50 µM genistein (Sigma, G6649), and 1-10 µM

VX-770 (Selleckchem, S1144) and inhibitor (final concentration: 20 µM CFTRinh-172 (Sigma,

C2992)) were applied sequentially at 10 or 20-minute intervals, to both serosal and mucosal

surfaces. Agonists/inhibitor were added from 200x-1000x stock solutions. Data acquisition

performed using ACQUIRE & ANALYZE Revision II (Physiologic Instruments).

ESE motif analysis

Potential SR protein binding sites were analyzed by ESEfinder [46].

Statistical analyses

Statistical analyses were performed with GraphPad Prism 5. Statistical parameters are indicated

in the figures and legends. For two-tailed t-test or one-way analysis of variance (ANOVA) with

Tukey's or Dunnett's post-test, $P<0.05$ was considered significant. The Pearson correlation and

P values were calculated using R. The asterisks and hash signs mark statistical significance as

follows: n.s. $P>0.05$; */# $P<0.05$; **/## $P<0.01$; ***/### $P<0.001$.

Acknowledgments

412 We are very grateful to Martin Mense and Hermann Bihler (CFFT, Lexington, MA) for

413 generously sharing protocols and advice, and for helpful comments on the manuscript.

Author contribution

415 Y.J.K., T.N., and A.R.K. conceived the study. A.R.K. supervised the study. Y.J.K. and T.N.

416 generated the DLD1-W1282X cells. F.P. performed exon 25-targeting ASO screening using the

417 *pW1282X-IVS25* NMD reporter. Y.J.K. designed and performed all other experiments and

418 analyzed the data. Y.J.K. and A.R.K. wrote the paper, and all authors approved the manuscript.

Funding

420 This work was supported by Cystic Fibrosis Foundation grant KRAINE17GO and NIH grant

421 R37GM42699 to A.R.K. Y.K was supported by NIH grants F30HL137326-04 and

422 T32GM008444. We acknowledge assistance from Cold Spring Harbor Laboratory Shared

423 Resources, funded in part by NCI Cancer Center Support Grant 5P30CA045508.

References

1.   CFTR2. Clinical and Functional Translation of CFTR (CFTR2). https://cftr2.org/ (2020).

2.   Cystic Fibrosis Foundation. *Patient Registry Annual Data Report. Lung*

http://www.cff.org/UploadedFiles/research/ClinicalResearch/Patient-Registry-Report-

2009.pdf (2018).

3.   Abeliovich, D. *et al.* Screening for five mutations detects 97% of cystic fibrosis (CF)

chromosomes and predicts a carrier frequency of 1:29 in the Jewish Ashkenazi

population. *Am. J. Hum. Genet.* **51**, 951–6 (1992).

4.   Rowe, S. M. *et al.* Restoration of W1282X CFTR Activity by Enhanced Expression. *Am. J.*

Respir. Cell Mol. Biol. **37**, 347–356 (2007).

5.   Haggie, P. M. *et al.* Correctors and potentiators rescue function of the truncated W1282X-

CFTR translation product. *J. Biol. Chem.* **292**, jbc.M116.764720 (2016).

6.   Wang, W., Hong, J. S., Rab, A., Sorscher, E. J. & Kirk, K. L. Robust stimulation of

W1282X-CFTR channel activity by a combination of allosteric modulators. *PLoS One* **11**,

e0152232 (2016).

7.   Goor, F. Van *et al.* Rescue of CF airway epithelial cell function in vitro by a CFTR

potentiator , VX-770. *Proc Natl Acad Sci U S A* **106**, 18825–18830 (2009).

8.   Johnson, L. G. *et al.* Efficiency of gene transfer for restoration of normal airway epithelial

function in cystic fibrosis. *Nat. Genet.* **2**, 21–25 (1992).

9.   Keating, D. *et al.* VX-445-Tezacaftor-Ivacaftor in patients with cystic fibrosis and one or

two Phe508del alleles. *N. Engl. J. Med.* **379**, 1612–1620 (2018).

10.   Lentini, L. *et al.* Toward a rationale for the PTC124 (Ataluren) promoted readthrough of

premature stop codons: a computational approach and GFP-reporter cell-based assay. *Mol. Pharm.* **11**, 653–664 (2014).

11. Linde, L. *et al.* Nonsense-mediated mRNA decay affects nonsense transcript levels and governs response of cystic fibrosis patients to gentamicin. *J. Clin. Invest.* **117**, 683–692 (2007).

12. Clancy, J. P. *et al.* No detectable improvements in cystic fibrosis transmembrane conductance regulator by nasal aminoglycosides in patients with cystic fibrosis with stop mutations. *Am. J. Respir. Cell Mol. Biol.* **37**, 57–66 (2007).

13. Kerem, E. *et al.* Ataluren for the treatment of nonsense-mutation cystic fibrosis: A randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Respir. Med.* **2**, 539–547 (2014).

14. Zainal Abidin, N., Haq, I. J., Gardner, A. I. & Brodlie, M. Ataluren in cystic fibrosis: development, clinical studies and where are we now? *Expert Opin. Pharmacother.* **18**, 1363–1371 (2017).

15. Keeling, K. M. *et al.* Attenuation of nonsense-mediated mRNA decay enhances in vivo nonsense suppression. *PLoS One* **8**, e60478 (2013).

16. Mutyam, V. *et al.* Therapeutic benefit observed with the CFTR potentiator, ivacaftor, in a CF patient homozygous for the W1282X CFTR nonsense mutation. *J. Cyst. Fibros.* (2016) doi:10.1016/j.jcf.2016.09.005.

17. Usuki, F. *et al.* Inhibition of SMG-8, a subunit of SMG-1 kinase, ameliorates nonsense-mediated mRNA decay-exacerbated mutant phenotypes without cytotoxicity. *Proc. Natl. Acad. Sci.* **110**, 15037–15042 (2013).

18.   Valley, H. C. *et al.* Isogenic cell models of cystic fibrosis-causing variants in natively

expressing pulmonary epithelial cells. *J. Cyst. Fibros.* 8–15 (2018)

doi:10.1016/j.jcf.2018.12.001.

19.   Keenan, M. M. *et al.* Nonsense-mediated RNA decay pathway inhibition restores

expression and function of W1282X CFTR. *Am. J. Respir. Cell Mol. Biol.* **61**, 290–300

(2019).

20.   Huang, L. *et al.* Antisense suppression of the nonsense mediated decay factor Upf3b as

a potential treatment for diseases caused by nonsense mutations. *Genome Biol.* **19**, 1–16

(2018).

21.   Lykke-Andersen, S. & Jensen, T. H. Nonsense-mediated mRNA decay: an intricate

machinery that shapes transcriptomes. *Nat. Rev. Mol. Cell Biol.* (2015)

doi:10.1038/nrm4063.

22.   Singh, G. *et al.* The cellular EJC interactome reveals higher order mRNP structure and an

EJC-SR protein nexus. *Cell* **151**, 750–764 (2012).

23.   Le Hir, H., Saulière, J. & Wang, Z. The exon junction complex as a node of post-

transcriptional networks. *Nat. Rev. Mol. Cell Biol.* **17**, 41–54 (2016).

24.   Haque, N. & Blanchette, M. Genomewide localization of Exon-Junction-Complex (EJC) in

Drosophila. *FASEB J.* **25**, 510.1-510.1.

25.   Saulière, J. *et al.* CLIP-seq of eIF4AIII reveals transcriptome-wide mapping of the human

exon junction complex. *Nat. Struct. Mol. Biol.* **19**, 1124–1131 (2012).

26.   Hauer, C. *et al.* Exon Junction Complexes Show a Distributional Bias toward Alternatively

Spliced mRNAs and against mRNAs Coding for Ribosomal Proteins Article Exon Junction

Complexes Show a Distributional Bias toward Alternatively Spliced mRNAs and against

mRNAs Coding fo. *CellReports* 1–16 (2016) doi:10.1016/j.celrep.2016.06.096.

27. Brogna, S. & Wen, J. Nonsense-mediated mRNA decay (NMD) mechanisms. *Nat. Struct.*

Mol. Biol. **16**, 107–113 (2009).

28. Hug, N., Longman, D. & Cáceres, J. F. Mechanism and regulation of the nonsense-

mediated decay pathway. *Nucleic Acids Res.* **44**, 1483–1495 (2016).

29. Nomakuchi, T. T., Rigo, F., Aznarez, I. & Krainer, A. R. Antisense oligonucleotide-

directed inhibition of nonsense-mediated mRNA decay. *Nat Biotech* **34**, 164–166 (2016).

30. Gogtay, N. J. & Sridharan, K. Therapeutic Nucleic Acids: Current clinical status. *Br. J.*

Clin. Pharmacol. 1–14 (2016) doi:10.1111/bcp.12987.

31. Khvorova, A. & Watts, J. K. The chemical evolution of oligonucleotide therapies of clinical

utility. *Nat. Biotechnol.* **35**, 238–248 (2017).

32. Lundin, K. E., Gissberg, O. & Smith, C. I. E. Oligonucleotide therapies: the past and the

present. *Hum. Gene Ther.* **26**, 475–485 (2015).

33. Khoo, B., Roca, X., Chew, S. L. & Krainer, A. R. Antisense oligonucleotide-induced

alternative splicing of the APOB mRNA generates a novel isoform of APOB. *BMC Mol.*

Biol. **8**, 3 (2007).

34. Kole, R., Krainer, A. R. & Altman, S. RNA therapeutics: beyond RNA interference and

antisense oligonucleotides. *Nat. Rev. Drug Discov.* **11**, 125–140 (2012).

35. Shimo, T., Maruyama, R. & Yokota, T. Designing effective antisense oligonucleotides for

exon skipping. *Methods Mol. Biol.* **1687**, 143–155 (2018).

36. Shimo, T. *et al.* Design and evaluation of locked nucleic acid-based splice-switching

oligonucleotides in vitro. *Nucleic Acids Res.* **42**, 8174–8187 (2014).

37.   Harding, P. L., Fall, A. M., Honeyman, K., Fletcher, S. & Wilton, S. D. The influence of

antisense oligonucleotide length on dystrophin exon skipping. *Mol. Ther.* **15**, 157–166

(2007).

38.   Mabin, J. W. *et al.* The exon junction complex undergoes a compositional switch that

alters mRNP structure and nonsense-mediated mRNA decay activity. *Cell Rep.* **25**, 1–16

(2018).

39.   Lareau, L. F., Inada, M., Green, R. E., Wengrod, J. C. & Brenner, S. E. Unproductive

splicing of SR genes associated with highly conserved and ultraconserved DNA

elements. *Nature* **446**, 926–929 (2007).

40.   Popp, M. W.-L. & Maquat, L. E. The dharma of nonsense-mediated mRNA decay in

mammalian cells. *Mol. Cells* **37**, 1–8 (2014).

41.   Lima, W. F., Vickers, T. A., Nichols, J., Li, C. & Crooke, S. T. Defining the factors that

contribute to on-target specificity of antisense oligonucleotides. *PLoS One* **9**, e101752

(2014).

42.   Sahashi, K. *et al.* TSUNAMI: An antisense method to phenocopy splicing-associated

diseases in animals. *Genes Dev.* **26**, 1874–1884 (2012).

43.   Busch, A. & Hertel, K. J. Evolution of SR protein and hnRNP splicing regulatory factors.

Wiley Interdiscip. Rev. RNA **3**, 1–12 (2012).

44.   Sinha, R. *et al.* Antisense oligonucleotides correct the familial dysautonomia splicing

defect in IKBKAPtransgenic mice. *Nucleic Acids Res.* **46**, 4833–4844 (2018).

45.   Hua, Y., Vickers, T. A., Baker, B. F., Bennett, C. F. & Krainer, A. R. Enhancement of

534   SMN2 Exon 7 Inclusion by Antisense Oligonucleotides Targeting the Exon. *PLoS Biol.* **5**,

535   e73 (2007).

536 46. Cartegni, L., Wang, J., Zhu, Z., Zhang, M. Q. & Krainer, A. R. ESEfinder: A web resource

537   to identify exonic splicing enhancers. *Nucleic Acids Res.* **31**, 3568–3571 (2003).

538 47. Yoshida, T. *et al.* Evaluation of off-target effects of gapmer antisense oligonucleotides

539   using human cells. *Genes to Cells* **24**, 827–835 (2019).

540 48. Roignant, J. Y. & Treisman, J. E. Exon junction complex subunits are required to splice

541   Drosophila MAP kinase, a large heterochromatic gene. *Cell* **143**, 238–250 (2010).

542 49. Wang, Z. *et al.* Transcriptome-wide modulation of splicing by the exon junction complex.

543   *Genome Biol.* **15**, 551 (2014).

544 50. Blazquez, L. *et al.* Exon junction complex shapes the transcriptome by repressing

545   recursive splicing. *Mol. Cell* **72**, 496-509.e9 (2018).

546 51. Boehm, V. *et al.* Exon junction complexes suppress spurious splice sites to safeguard

547   transcriptome integrity. *Mol. Cell* **72**, 482-495.e7 (2018).

548 52. Michelle, L. *et al.* Proteins associated with the exon junction complex also control the

549   alternative splicing of apoptotic regulators. *Mol. Cell. Biol.* **32**, 954–967 (2012).

550 53. Fukumura, K. *et al.* The exon junction complex controls the efficient and faithful splicing

551   of a subset of transcripts involved in mitotic cell-cycle progression. *Int. J. Mol. Sci.* **17**,

552   (2016).

553 54. Saulière, J. *et al.* The exon junction complex differentially marks spliced junctions. *Nat.*

554   *Struct. Mol. Biol.* **17**, 1269–1271 (2010).

555 55. Ren, H. Y. *et al.* VX-809 corrects folding defects in cystic fibrosis transmembrane

conductance regulator protein through action on membrane-spanning domain 1. *Mol.*

Biol. Cell **24**, 3016–3024 (2013).

56. O'Riordan, C. R., Lachapelle, A. L., Marshall, J., Higgins, E. A. & Cheng, S. H.

Characterization of the oligosaccharide structures associated with the cystic fibrosis

transmembrane conductance regulator. *Glycobiology* **10**, 1225–1233 (2000).

57. Cozens, A. L. *et al.* CFTR expression and chloride secretion in polarized immortal human

bronchial epithelial cells. *Am. J. Respir. Cell Mol. Biol.* **10**, 38–47 (1994).

58. Xue, X. *et al.* Identification of the amino acids inserted during suppression of CFTR

nonsense mutations and determination of their functional consequences. *Hum. Mol.*

Genet. **26**, 3116–3129 (2017).

59. Mutyam, V. *et al.* Discovery of clinically approved agents that promote suppression of

cystic fibrosis transmembrane conductance regulator nonsense mutations. *Am. J. Respir.*

Crit. Care Med. **194**, 1092–1103 (2016).

60. Scharner, J. *et al.* Hybridization-mediated off-target effects of splice-switching antisense

oligonucleotides. *Nucleic Acids Res.* 1–15 (2019) doi:10.1093/nar/gkz1132.

61. Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**,

2281–2308 (2013).

62. Glozman, R. *et al.* N-glycans are direct determinants of CFTR folding and stability in

secretory and endocytic membrane traffic. *J. Cell Biol.* **184**, 847–862 (2009).

63. Zegarra-Moran, O. *et al.* Correction of G551D-CFTR transport defect in epithelial

monolayers by genistein but not by CPX or MPB-07. *Br. J. Pharmacol.* **137**, 504–512

(2002).

Figure 1. Identification of NMD-inhibiting ASOs and assessment of their specificity.

A. Schematic of NMD reporters. The numbers show the *CFTR* exons present in the NMD

reporters. The red asterisk (*) indicates the location of the W1282X mutation. **B.** Schematic of

ASO screening. 19 MOE-PS modified 15mer ASOs (yellow and magenta bars) were designed

to cover the presumptive EJC binding sites on exons 24, 25, and 26 at 1-nt resolution. **C-E.**

U2OS cells stably expressing each NMD reporter were transfected with individual ASOs

targeting EJC binding regions on *CFTR* (C) exon 24, (D) exon 25, or (E) exon 26, respectively.

Reporter mRNA levels were measured by radioactive RT-PCR, using primers listed in Table 4.

RPL32-normalized reporter expression is compared to that of the negative-control ASO (H26)

transfection, and is shown below the RT-PCR images. **F.** Effect of cycloheximide (CHX) on

CFTR expression in 16HBE-W1282X and DLD1-W1282X cells. **G.** Effect of the ASO cocktail

C478-C495-C515 (LC) on *CFTR* expression in 16HBE-W1282X cells. **H.** Comparison between

15mer and 18mer lead ASO cocktails (C478-C494-C514 or C24-C25-C26, respectively). The

15mer and 18mer scramble ASOs were used as negative controls. **I.** Effect of cycloheximide

(CHX) on *CFTR* expression in DLD1-WT, 16HBE-G551D, and 16HBE-F508del cells. **J.** *CFTR*

mRNA levels in DLD1-WT, 16HBE-G551D, and 16HBE-F508del transfected with the control

ASOs or the lead ASO cocktail at a nominal total concentration of 120 nM. **K.** Endogenous

NMD-sensitive mRNA levels in 16HBE-W1282X cells treated with cycloheximide (orange), 120

nM scramble ASO (Sc; purple) or 120 nM lead ASO cocktail (blue). All mRNA levels in F-K

were measured by RT-qPCR. *RPL32* served as internal reference for all panels except panel H,

in which *HPRT* served as internal reference. NT=No treatment; Dox: doxycycline 1 µg/mL;

Sc=Scramble ASO; CC=Control ASO cocktail C488-C507-C526; LC= lead ASO cocktail C478-

C495-C515, C478-C494-C514, or C24-C25-C26; CHX = 1-hr incubation with 100 µg/mL

cycloheximide. All error bars indicate standard deviation. For all treatments, n=3, except n=2 in

LC18-mer 120nM in panel G. For all statistical tests, n.s. $P>0.05$, *$P<0.05$, **$P<0.01$,

***$P<0.001$. Panel F, I, and K: Student's t-test. Panel G, H, and J: one-way ANOVA with Tukey's

post-test.

Figure 2. ASO-cocktail optimization and mechanism of action.

A-C. New ASOs targeting *CFTR* (H) exon 24, (I) exon 25, or (J) exon 26 were individually

screened in 16HBE-W1282X cells, in combination with two ASOs that target the other two

exons, at a total nominal concentration of 120 nM. Red bars indicate the lead ASO identified in

each combination screen. **D.** Comparison between the ASO cocktails C478-C495-C515 and

C478-C494-C514. **E-F.** *CFTR* and *eIF4A2* mRNA levels in (D) 16HBE-G542X cells and (E)

16HBE-R1162X cells transfected with ASOs at a total nominal concentration of 120 nM. **G**. The

number of required EJCs targeted by ASOs C478, C494, C515, or all together, was assessed

by transfecting 16HBE-W1282X cells with one, two, or three EJC-targeting ASOs at the same

total nominal concentration. **H**. 16HBE-W1282X cells were transfected with various

combinations of the lead ASOs C478, C495, and C515. **I.** Mean PSI of each *CFTR* isoform in

16HBE-W1282X cells transfected with various combinations of C478, C495, and C515. All

mRNA levels in A-G were measured by RT-qPCR. *RPL32* mRNA level served as an internal

reference. Error bars show standard deviation. Abbreviations are as in Figure 1. n=3 for all

treatments, except in panel C, n=5 or 6, and panel G, n=4 for NT, Sc, CC, and Lead ASO

40:40:40. For all Dunnett's post-tests, n.s. $P>0.05$, *$P<0.05$, **$P<0.01$, ***$P<0.001$. For all

Student's t-tests, #$P<0.05$, ##$P<0.01$. Panels A-H: one-way ANOVA with Dunnett's post-test

versus NT, or Student's t-test. Panel I: n=4, one-way ANOVA with Dunnett's post-test, versus

each isoform in 'lead ASO 40:40:40'.

Figure 3. Effect of the lead ASO cocktail on CFTR-W1282X protein expression and function.

A. Western blot of CFTR-W1282X protein in 16HBE-W1282X cells transfected with control or

EJC-targeting ASO cocktail C478-494-C515 and treated with VX-809 at the indicated

concentrations. The closed and open arrowheads indicate C and B bands of CFTR proteins,

respectively. **B**. Quantification of total CFTR protein in (A). **C-D**. Average traces from Ussing-

chamber assay of 16HBE-W1282X cells treated with 20 µM of (C) 15mer scramble ASO (Sc) or

(D) 15mer lead ASO cocktail C479-C494-C514. **E.** Total area under the curve in (C) and (D).

Before the assays, the cells were treated for 24 h with VX-809 (3 µM). The traces shown are the

average of the three replicates. **F**. The levels of *CFTR-W1282X* and endogenous NMD-sensitive

mRNAs in 16HBE-W1282X cells assayed in €, normalized to Sc control. *RPL32* was used as

internal reference*.* **G-K** Average traces from Ussing-chamber assay of 16HBE-W1282X cells

treated with (G) 20 µM 18mer scramble ASO, (H) 20 µM 18mer lead ASO cocktail (C24-C25-

C26), (I) 20 µM 18mer scramble ASO and 200 µM G418, (J) 20 µM 18mer lead ASO cocktail

and 200 µM G418, and (K) 20 µM 18mer lead ASO cocktail and 600 µM G418. All G418

treatments were started 24 hr prior to the assay. The traces shown are the average of three

replicates. Black error bars on each point show standard deviation. **L.** The total area under the

curve of (G-K). Error bars show standard deviations. Fsk: forskolin. n=3 for all treatments. For

all statistical tests, n.s. P>0.05, *P<0.05, **P<0.01, ***P<0.001. Panels E and F: Student's t-test.

Panels B and L: one-way ANOVA with Tukey's post-test.

TABLES

Table 1. ASOs

ASO type	ASO Name	Sequence
EJC-targeting candidate	C470	CTTGATCACTCCACT
EJC-targeting candidate	C471	TCTTGATCACTCCAC
EJC-targeting candidate	C472	TTCTTGATCACTCCA
EJC-targeting candidate	C473	TTTCTTGATCACTCC
EJC-targeting candidate	C474	ATTTCTTGATCACTC
EJC-targeting candidate	C475	TATTTCTTGATCACT
EJC-targeting candidate	C476	ATATTTCTTGATCAC
EJC-targeting candidate	C477	CATATTTCTTGATCA
EJC-targeting candidate	C478	CCATATTTCTTGATC
EJC-targeting candidate	C479	TCCATATTTCTTGAT

EJC-targeting candidate	C480	TTCCATATTTCTTGA
EJC-targeting candidate	C481	TTTCCATATTTCTTG
EJC-targeting candidate	C482	CTTTCCATATTTCTT
EJC-targeting candidate	C483	ACTTTCCATATTTCT
EJC-targeting candidate	C484	AACTTTCCATATTTC
EJC-targeting candidate	C485	CAACTTTCCATATTT
EJC-targeting candidate	C486	GCAACTTTCCATATT
EJC-targeting candidate	C487	TGCAACTTTCCATAT
EJC-targeting candidate	C488	CTGCAACTTTCCATA
EJC-targeting candidate	C489	TTCATCAAGCAGCAA
EJC-targeting candidate	C490	GTTCATCAAGCAGCA
EJC-targeting candidate	C491	GGTTCATCAAGCAGC
EJC-targeting candidate	C492	GGGTTCATCAAGCAG
EJC-targeting candidate	C493	TGGGTTCATCAAGCA
EJC-targeting candidate	C494	CTGGGTTCATCAAGC
EJC-targeting candidate	C495	ACTGGGTTCATCAAG
EJC-targeting candidate	C496	CACTGGGTTCATCAA
EJC-targeting candidate	C497	GCACTGGGTTCATCA
EJC-targeting candidate	C498	AGCACTGGGTTCATC
EJC-targeting candidate	C499	GAGCACTGGGTTCAT
EJC-targeting candidate	C500	TGAGCACTGGGTTCA
EJC-targeting candidate	C501	ATGAGCACTGGGTTC
EJC-targeting candidate	C502	AATGAGCACTGGGTT
EJC-targeting candidate	C503	AAATGAGCACTGGGT
EJC-targeting candidate	C504	CAAATGAGCACTGGG

EJC-targeting candidate	C505	CCAAATGAGCACTGG
EJC-targeting candidate	C506	TCCAAATGAGCACTG
EJC-targeting candidate	C507	ATCCAAATGAGCACT
EJC-targeting candidate	C508	TTGCTTCTATCCTGT
EJC-targeting candidate	C509	ATTGCTTCTATCCTG
EJC-targeting candidate	C510	CATTGCTTCTATCCT
EJC-targeting candidate	C511	GCATTGCTTCTATCC
EJC-targeting candidate	C512	AGCATTGCTTCTATC
EJC-targeting candidate	C513	CAGCATTGCTTCTAT
EJC-targeting candidate	C514	CCAGCATTGCTTCTA
EJC-targeting candidate	C515	TCCAGCATTGCTTCT
EJC-targeting candidate	C516	TTCCAGCATTGCTTC
EJC-targeting candidate	C517	ATTCCAGCATTGCTT
EJC-targeting candidate	C518	CATTCCAGCATTGCT
EJC-targeting candidate	C519	GCATTCCAGCATTGC
EJC-targeting candidate	C520	GGCATTCCAGCATTG
EJC-targeting candidate	C521	TGGCATTCCAGCATT
EJC-targeting candidate	C522	TTGGCATTCCAGCAT
EJC-targeting candidate	C523	GTTGGCATTCCAGCA
EJC-targeting candidate	C524	TGTTGGCATTCCAGC
EJC-targeting candidate	C525	TTGTTGGCATTCCAG
EJC-targeting candidate	C526	ATTGTTGGCATTCCA
Scramble 15mer	Scramble control based on C494	CACGCTACTTGATGC
EJC 18mer	C24-18m	TTCCATATTTCTTGATCA

EJC 18mer	C25-18m	ACTGGGTTCATCAAGCAG
EJC 18mer	C26-18m	TTCCAGCATTGCTTCTAT
Scramble 18mer	Scramble control based on C25	ACAGGCTTCTTCATGCAC
Control	H24	CTCAGGATCCACGTG
Control	H26	CAGGATCCACGTGCA
Control	H27	AGGATCCACGTGCAG

650 **Table 2. sgRNA and W1282X repair template sequences**

CFTR sgRNA Sense strand	5'-CACCGCAATAACTTTGCAACAGTGG-3'
CFTR sgRNA antisense strand	5'-AAACCCACTGTTGCAAAGTTATTGC-3'
W1282X repair template	5'-AACACTGAAGGAGAAATCCAGATCGATGGTGTGTCTTGGGATTCAATAACTTTGCAACAGTGAAGAAAAGCCTTTGGAGTGATACCACAGGTGAGCAAAAGGACTTAGCCAGAAAAAGG-3'

Table 3. siRNA

Source	Target gene	Sequence	Source
Custom-sense	UPF1	GAUGCAGUUCCGCUCCAUU	Sigma
Custom-antisense	UPF1	AAUGGAGCGGAACUGCAUC	Sigma
Predesigned	CFTR	SASI_Hs02_00302648	Sigma

Table 4. Primers for RT-PCR and RT-qPCR

Target gene	primer	seq	source
NMD-sensitive GADD45A	GADD45A_NMD_F	GAGCTCCTGCTCTTGGAGAC	*Mabin et al., 2018*
NMD-sensitive GADD45A	GADD45A_NMD_R	GCAGGATCCTTCCATTGAGA	*Mabin et al., 2018*
NMD-sensitive eIF4A2	eIF4A2_NMD_F	AGGGTCAAGTCGTGTTCTGATC	*Mabin et al., 2018*
NMD-sensitive eIF4A2	eIF4A2_NMD_R	ACCAACTGCTGCTATCGACTC	*Mabin et al., 2018*
NMD-sensitive SF3B1	SF3B1_NMD_F	AATTTCCCCAGAGCGTCTTG	*Mabin et al., 2018*
NMD-sensitive SF3B1	SF3B1_NMD_R	TTCGTGCCTTTGTCTCCATC	*Mabin et al., 2018*
NMD-sensitive C1orf37	C1orf37_NMD_F	TTGCTGCTCGAATCTCCAAG	*Mabin et al., 2018*

NMD-sensitive C1orf37	C1orf37_NMD_R	ACTTCTGCTGCCATCACAAC	*Mabin et al., 2018*
NMD-sensitive SRSF2	SRSF2_NMD_F	CCTCTTAAGAAAATGCTGCGGTCTC	Lareau et al., 2007
NMD-sensitive SRSF2	SRSF2_NMD_R	ATCAGCCAAATCAGTTAAAATCTGC	Lareau et al., 2007
NMD-sensitive SRSF4	SRSF4_NMD_F	GGATCTGAAGAACGGTCTGTTATGT	Lareau et al., 2007
NMD-sensitive SRSF4	SRSF4_NMD_R	TCACTCGTCTTTTGGTTCCCATTAG	Lareau et al., 2007
NMD-sensitive RPL12	RPL12_NMD_F	CTGGGCCTTAGCTTCTTCAC	Lareau et al., 2007
NMD-sensitive RPL12	RPL12_NMD_R	AAGTGGCACCGACTTCACCT	*Mabin et al., 2018*
CFTR-exon22	CFTR22F	CAATAAGTCCTGGCCAGAGG	
CFTR-exon26	CFTR26F	GCACAGTAATTCTCTGTGAACACAGG	
CFTR-exon27	CFTR27R	TCCTCTCGTTCAGCAGTTTCTGG	
CFTR-exon23	CFTR23F	TTGCAACAGtgaAGGAAAGCC	
CFTR-exon23	CFTR23R	AAGGCTTTCCTtcaCTGTTGC	
CFTR-exon22-splicing	CFTR-22F-splice	GCGATCTGTGAGCCGAGTC	
CFTR-exon24-splicing	CFTR-24R-splice	CTTGATCACTCCACTGTTCATAGGGATC	
RPL32	RPL32_F	AGAGGCATTGACAACAGGGTT	

RPL32	RPL32_R	GTGAGCGATCTCGGCACAG	
HPRT	HPRT_F	TGACCAGTCAACAGGGGACA	
HPRT	HPRT_R	TGCCTGACCAAGGAAAGCAA	