PULLING ON MULTIPLE THREADS OF EVIDENCE TO REVEAL TREE SPECIES COEXISTENCE

Gordon G. McNickle[1,2,3,*], Morgan V. Ritzi[1,3], Kliffi M.S. Blackstone[1,2,3], John J. Couture[2,4,5], Taylor

Nelson [2,4,5], Brady S. Hardiman[5,6], Madeline S. Montague[5], Douglass F. Jacobs[5]

[1]Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907-2054

[2]Center for Plant Biology, Purdue University, West Lafayette, IN 47907-2054

[3]Department of Biology, Purdue University, West Lafayette, IN 47907-2054

[4]Department of Entomology, Purdue University, West Lafayette, IN 47907-2054

[5]Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN 47907-2054

[6]Environmental and ecological engineering, Purdue University, West Lafayette, IN 47907-2054

* correspondence: gmcnickle@purdue.edu

ABSTRACT

Understanding drivers of species coexistence is a central challenge in ecology [1,2]. Coexistence

cannot be observed directly, and while species co-occurrence in time and space is necessary for

coexistence, it is not sufficient to prove coexistence [3,4]. Species exclusion from a region is

potentially observable, but can take decades to occur, and still might occur stochastically [5,6]. Thus,

ecologists generally use theory to identify indirect observations that are indicative of mechanisms

driving coexistence or exclusion [7-9]. Various methods have been developed to indirectly infer

coexistence, each of which requires different data, and none of which are usually conclusive on

their own (e.g. [9-12]). Here, we demonstrate agreement using three different approaches examining

coexistence of multiple hardwood species. First, in an experimental planting of three mature tree

species we found no relationship between productivity and species diversity, which could be due

to a lack of niche differences among species. Second, we used modern coexistence theory to

calculate niche and fitness differences for each pair of species, which confirmed the lack of niche

differences among species, and showed high fitness differences that could create a neutral

distribution of species in nature. Third, we used the United States Department of Agriculture

Forest Inventory and Analysis data to examine co-occurrence patterns of our species across

thousands of natural forest stands and found that indeed, these three species were distributed

randomly throughout the USA. Given that these independent methods agree, we take this as

strong evidence about a lack of coexistence.

MAIN BODY

Ecological species coexistence has a well-developed body of theory [7,8,13], but is challenging to

study empirically [4,14,15]. Strictly, coexistence occurs when the population growth trajectories of

multiple species share a stable equilibrium. Unfortunately, natural systems are never at equilibrium.

Furthermore, species may simply find themselves thrust together or apart for entirely stochastic

neutral reasons [3,4,7]. Ecologists therefore must rely on theory to identify experimental results that

might imply coexistence [9,10,16]. Generally, ecologists expect that differences among species in many

dimensions (e.g. resource use, habitat use, seasonality) control coexistence such that there is some

limit to how similar two species can be before one excludes the other. The totality of these

dimensions, describing the differences among species, is referred to as the niche of a species [1]. Here,

we combine multiple approaches to inferring species coexistence. On their own, each approach has

proven ambiguous, but when combined here they all point in the same direction providing

compelling evidence that ecologists are gaining ground on a predictive theory of coexistence.

First, one approach to indirectly infer mechanisms of coexistence is the relationship

between community productivity and species richness [5,17]. Generally, though not universally,

researchers have found that more species rich plant communities are also more productive [10,11]. One

explanation for this positive diversity-productivity correlation would be if competition within a

species was more intense than competition among species [18]. Under this interpretation, low

diversity communities grow poorly due to strong negative intraspecific feedbacks, while species rich

mixtures grow better because of diluted negative effects caused by niche differences among species.

However, this is not the only explanation. Others have argued that positive diversity productivity

relationships merely reflects sampling error with an increasing probability of having large productive

species as diversity increases [19].

To examine the relationship between productivity and diversity we planted three species of

hardwood trees in 2007: (*Prunus serotina* Ehrh., *Quercus rubra* L. and *Castanea dentata* (Marsh.)

Borkh.) in all seven possible combinations of one, two or three species, crossed with two different

planting densities (1m, or 2m between trees). The experiment has three replicate blocks, and is

located in west-central Indiana (40°26'41.9"N 87°01'46.4"W) within the historic range of these three

species (Fig S1) [20]. In the summer and fall of 2018 we took wood cores from the base of three

individual trees of each species from each plot and used dendrochronology to estimate the

productivity of these plots between 2007-2017. The tree cores give us the diameter of the base of

the tree in each year, and we used this to calculate the cross-sectional area of the tree trunk in each

year (i.e. basal area). The area of each annual growth ring is an estimate of productivity and is

calculated as the area of the tree trunk in year t minus the area in year $t-1$, also known as the

basal area increment (BAI). To estimate leaf production in 2018 and 2019, we used cylindrical litter

traps (28.6cm diameter). Finally, to estimate fine root production in 2019 we used an ingrowth core

method. In autumn 2018, a cylindrical core (7.5cm diameter, 50cm deep) was removed from the

forest floor, fine roots were sorted out and the soil replaced inside of a nylon mesh tube. After one

year, the core was retrieved in 2019, roots washed, dried and weighed.

79   We found that the diversity and productivity of leaves, wood and roots were uncorrelated in

80   these plots across all years (Fig 1; Table S1-3). Given contrasting interpretations of diversity

81   productivity relationships, the lack of a positive relationship is either: (i) an indicator of a lack of

82   niche differences among our species[21] or; (ii) simply evidence that our species grow at similar rates[19].

84   To further probe the alternative explanations of the relationship between diversity and

productivity, we used modern coexistence theory (MCET)[7,13]. MCET focuses on population growth

rates, not biomass production, and partitions coexistence into two components: stabilizing

mechanisms and fitness equalizing mechanisms [7,13]. Stabilizing mechanisms of coexistence are

mechanisms that cause one species to limit itself more than other species meaning that the negative

effects of intraspecific interactions are stronger than any effects of interspecific interactions [13]. Thus,

this allows us to directly and independently examine one explanation for the lack of correlation

between diversity and productivity. Stabilizing mechanisms arise because of niche differences such

as adaptation to different environments, or differences in biotic interactions [13]. Second, fitness

equalizing mechanisms are mechanisms that minimize average fitness differences among species [13].

If fitness is defined as per capita survival and reproduction, then large fitness differences among

species can cause spikes in births or deaths which can negate stabilizing effects. For the lack of a

positive relationship between productivity and species richness to be an indicator of non-

coexistence rather than simple species growth rates, then we hypothesized that we should find

small niche differences in demography data using MCET.

We estimated the slope of per-capita population changes from 2007-2019 as a function of

planting density for all three species either in monoculture (α_{ii}), or in each species pair (α_{ij}; Fig S2,

Table S4). Niche differences (ND) were calculated as:

$$ND = 1 - \sqrt{\left|\frac{\alpha_{ij}\alpha_{ji}}{\alpha_{ii}\alpha_{jj}}\right|},$$ Eqn. 1

and fitness differences (FD) were estimated using the competitive response ratio component of

fitness differences, calculated as:

$$FD = \sqrt{\left|\frac{\alpha_{ij}\alpha_{ii}}{\alpha_{ji}\alpha_{jj}}\right|}.$$ Eqn. 2

These formulations for ND and FD are derived elsewhere [9,22]. ND can be negative if species are

extremely similar such that $\alpha_{ij}\alpha_{ji} > \alpha_{ii}\alpha_{jj}$. In the case of negative frequency dependence, regions

of coexistence and exclusion can be demarcated by:

$$FD = \frac{1}{1-ND},$$ Eqn. 3

such that exclusion occurs above and to the right of this curve, and coexistence occurs below and to

the left of the curve [7]. In the case of positive frequency dependence [23], exclusion and priority effects

are demarcated by:

$$FD = 1 - ND,$$ Eqn. 4

where exclusion occurs above and to the right of this curve, and priority effects below and to two

the left.

Population growth was generally lower in mixture than in monoculture (Fig S2). MCET

calculations reveal that the niche differences among all combinations of the three species in our

study were too small to permit species coexistence (Fig 2). Thus, MCET provides evidence that the

lack of a diversity-productivity relationship was due to niche similarity. However, the estimated

confidence intervals for niche and fitness differences of *Q. rubra* and *P. serotina* partially overlap

with the coexistence region (Fig 2). Furthermore, most species pairs had high fitness differences that

would predict stochasticity in co-occurrence of these species across their native ranges. Therefore,

we hypothesized that if these species cannot coexist, then they should not be found together in

natural forest stands more frequently than expected due to chance. Alternatively, since each species

pair also had high fitness differences, these species might occur randomly in nature due to neutral

stochastic fitness differences.

To probe these alternative hypotheses about co-occurrence, we used the publicly available

USDA Forest Inventory and Analysis (FIA) data. The FIA data are an inventory of trees from

thousands of permanent plots across the USA, including data for hundreds of thousands of trees

allowing us to examine co-occurrence across the entire native range of our three focal species. *P.*

serotina, *Q. rubra* and *C. dentata* all have overlapping ranges in the USA (Fig 3A-C). We used the

checkerboard index of co-occurrence [3,24,25] (C-score) to quantify and statistically compare the

observed co-occurrence pattern to a null expectation that the three species are randomly

distributed throughout their range. We found that the observed negative co-occurrence pattern was

not significantly different from random (Fig 3D). This outcome means that, on average, these species

do not positively co-occur throughout their range as might be expected if coexistence was common,

but nor are they negatively associated as might be expected if exclusion was common. We conclude

that the seemingly random association of these three species across their entire range is consistent

with large fitness differences (Fig 2) that predict neutral stochastic processes are likely to control the

co-occurrence of these three hardwood species.

Here, by pulling on multiple threads of evidence, none of which was conclusive on its own,

we found strong evidence that three species of hardwood trees cannot coexist. First, the lack of a

positive relationship between productivity and species richness suggested a lack of niche differences

among species (Fig 1), but on its own could be explained by something other than coexistence [19].

Second, MCET independently confirmed the small niche differences. However, MCET also showed

large fitness differences that suggest neutral processes may control the distribution of these species.

Indeed, we found that the three study species randomly co-occur other across their entire range (Fig

3). Each of these methods has been used separately to try and infer potential mechanisms of

coexistence, but on their own are not always conclusive proof of coexistence. Predicting species

coexistence has long been a goal of ecological theory, and the confluence of multiple threads of

evidence all pointing in the same direction demonstrates that ecologists have made significant

progress in uniting theory and observation surrounding species coexistence.

ACKNOWLEDGEMENTS

This work was funded by the American Chestnut Foundation, a Purdue AgSeed award, and a

Hardwood Tree Improvement and Regeneration Centre grant to GGM, DFJ, JJC and BSH, and Hatch

awards 1010722 to GGM and ND011490 to JJC, and McIntire Stennis project IND011535 to DFJ. The

authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

MVR, KMSB and GGM performed dendrochronology analysis. MSM, DFJ and GGM performed root

ingrowth core analyses. KMSB, TN, JJC and GGM performed leaf analyses. All authors contributed to

tree tagging and survival data collection. GGM wrote the manuscript, performed MCET and C-score

calculations and all authors contributed to revisions and discussion of approaches.

DATA AVAILABILITY

The USDA FIA data are publically available from the USDA. Our site specific productivity and

demography data are publically available on DRYAD at: . All code required to reproduce analyses,

figures and tables are publically available on GitHub at: https://github.com/ggmcnickle/TreeCoexist/

REFERENCES

1   Hutchinson, G. E. Homage to santa-rosalia or why are there so many kinds of animals. *Am.*
Nat. **93**, 145-159 (1959).
2   Silvertown, J. Plant coexistence and the niche. *Trends Ecol. Evol.* **19**, 605-611, doi:DOI
10.1016/j.tree.2004.09.003 (2004).
3   Stone, L. & Roberts, A. The checkerboard score and species distributions. *Oecologia* **85**, 74-
79, doi:10.1007/bf00317345 (1990).
4   Blanchet, F. G., Cazelles, K. & Gravel, D. Co-occurrence is not evidence of ecological
interactions. *Ecol. Lett.* **23**, 1050-1063, doi:10.1111/ele.13525 (2020).
5   Tilman, D. *et al.* Diversity and productivity in a long-term grassland experiment. *Science* **294**,
843-845, doi:10.1126/science.1060391 (2001).
6   Crawley, M. J. *et al.* Determinants of species richness in the park grass experiment. *The*
American Naturalist **165**, 179-192, doi:10.1086/427270 (2005).
7   Adler, P. B., HilleRisLambers, J. & Levine, J. M. A niche for neutrality. *Ecol. Lett.* **10**, 95-104,
doi:10.1111/j.1461-0248.2006.00996.x (2007).
8   Hart, S. P., Freckleton, R. P. & Levine, J. M. How to quantify competitive ability. *J. Ecol.* **106**,
1902-1909, doi:doi:10.1111/1365-2745.12954 (2018).
9   Kraft, N. J. B., Godoy, O. & Levine, J. M. Plant functional traits and the multidimensional
nature of species coexistence. *Proceedings of the National Academy of Sciences* **112**, 797-
802, doi:10.1073/pnas.1413650112 (2015).
10   Liang, J. J. *et al.* Positive biodiversity-productivity relationship predominant in global forests.
Science **354**, 12, doi:10.1126/science.aaf8957 (2016).
11   Grace, J. B. *et al.* Integrative modelling reveals mechanisms linking productivity and plant
species richness. *Nature* **529**, 390-393, doi:10.1038/nature16524 (2016).
12   McNickle, G. G. *et al.* Checkerboard score–area relationships reveal spatial scales of plant
community structure. *Oikos* **127**, 415-426, doi:10.1111/oik.04620 (2018).
13   Chesson, P. Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**, 343-+
(2000).
14   Wright, S. J. Plant diversity in tropical forests: A review of mechanisms of species
coexistence. *Oecologia* **130**, 1-14, doi:10.1007/s004420100809 (2002).
15   Avolio, M. L. *et al.* A comprehensive approach to analyzing community dynamics using rank
abundance curves. *Ecosphere* **10**, e02881, doi:https://doi.org/10.1002/ecs2.2881 (2019).
16   Ulrich, W. & Gotelli, N. J. Pattern detection in null model analysis. *Oikos* **122**, 2-18,
doi:10.1111/j.1600-0706.2012.20325.x (2013).
17   Grime, J. P. Control of species density in herbaceous vegation. *J. Environ. Manage.* **1**, 151-
167 (1973).
18   Lotka, A. J. The growth of mixed populations: Two species competing for a common food
supply. *Journal of the Washington Academy of Sciences* **22**, 461-469 (1932).
19   Cheng, Y. X., Zhang, C. Y., Zhao, X. H. & von Gadow, K. Biomass-dominant species shape the
productivity-diversity relationship in two temperate forests. *Ann. For. Sci.* **75**, 9,
doi:10.1007/s13595-018-0780-0 (2018).
20   Gauthier, M. M., Zellers, K. E., Lof, M. & Jacobs, D. F. Inter- and intra-specific
competitiveness of plantation-grown american chestnut (castanea dentata). *For. Ecol.*
Manage. **291**, 289-299, doi:10.1016/j.foreco.2012.11.014 (2013).
21   Cardinale, B. J. *et al.* Impacts of plant diversity on biomass production increase through time
because of species complementarity. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 18123-18128,
doi:10.1073/pnas.0709069104 (2007).
22   Narwani, A., Alexandrou, M. A., Oakley, T. H., Carroll, I. T. & Cardinale, B. J. Experimental
evidence that evolutionary relatedness does not affect the ecological mechanisms of

coexistence in freshwater green algae. *Ecol. Lett.* **16**, 1373-1381, doi:10.1111/ele.12182
(2013).
23 Grainger, T. N., Letten, A. D., Gilbert, B. & Fukami, T. Applying modern coexistence theory to
priority effects. *Proceedings of the National Academy of Sciences* **116**, 6205-6210,
doi:10.1073/pnas.1803122116 (2019).
24 Diamond, J. in *Ecology and evolution of communities*   (eds ML. Cody & JM. Diamond) 342-
444 (Belknap Press, 1975).
25 Ulrich, W. & Gotelli, N. J. Null model analysis of species associations using abundance data.
Ecology **91**, 3384-3397, doi:10.1890/09-2157.1 (2010).
26 Gotelli, N. J. Null model analysis of species co-occurrence patterns. *Ecology* **81**, 2606-2621,
doi:10.1890/0012-9658(2000)081[2606:NMAOSC]2.0.CO;2 (2000).
27 Hardy, O. J. Testing the spatial phylogenetic structure of local communities: Statistical
performances of different null models and test statistics on a locally neutral community. *J.*
Ecol. **96**, 914-926, doi:10.1111/j.1365-2745.2008.01421.x (2008).
28 Vegan: Community ecology package v.2.4-1 (2016).
29 Miklós, I. & Podani, J. Randomization of presence–absence matrices: Comments and new
algorithms. *Ecology* **85**, 86-92, doi:10.1890/03-0101 (2004).
30 Ulrich, W. & Gotelli, N. J. Disentangling community patterns of nestedness and species co-
occurrence. *Oikos* **116**, 2053-2061, doi:10.1111/j.2007.0030-1299.16173.x (2007).

Fig 1: The relationship between species richness, and wood productivity estimated from leaf litter

traps from 2018-2019 (**A,B**); dendrochronology as basal area increments from 2007-2017 (BAI; **C,D**),

and fine root productivity in 2019 (**E,F**). The y-axis is on a $\log_{10}$ scale in all panels. Trees were spaced

either 1m or 2m apart (1 tree/m^2, or 0.5 trees/m^2 respectively). Points are drawn with a jitter

around richness, and the y-axis is plotted on a $\log_{10}$ scale. Lines come from a linear regression (Tables

S1-3). None of the estimates for the individual slopes of the diversity productivity relationship were

different from zero.

Fig 2: Comparison of niche and fitness differences calculated using population growth rates and

MCET for each pair of species: *P. serotina* vs *Q. rubra* (PQ); *C. dentata* vs *P. serotina* (CP) and *C.*

dentata vs *Q. rubra* (CQ). Since our propagation of error calculations could not account for

covariance this makes them larger, thus, error bars are 80% confidence intervals. The black lines

demarcate the regions of coexistence, priority effects and exclusion as labeled.

Fig 3: Distribution and population density (count) in natural forest stands of *P. serotina* **(A)** *Q. rubra*

(B) and *C. dentata* **(C)** across the FIA plots in the USA. Red represents higher density, while blue

represents low density. We compared observed C-score (red vertical line) with the null expectation

of random distribution (grey histogram) across all FIA plots **(D)**. The C score is high (relative to null)

when species negatively co-occur, and low (relative to null) when they positively co-occur. The lack

of difference compared to null (p=0.4219) suggests these three species are randomly distributed in

nature with respect to each other on average.

SUPPLEMENTARY METHODS:

Error propagation in modern coexistence theory

Since the components of modern coexistence theory are typically predicted from regression

coefficients of per-capita population growth rates (i.e. fitness) versus population density [8] that have

an associated variance-covariance matrix it is possible to use the known propagation of error

methods to create confidence intervals around the estimates of niche and fitness differences (e.g.

[23]). Here, we outline the approach we took to estimating error for niche and fitness differences. This

approach assumes no covariance between terms, which actually leads to larger errors than if we

assumed covariance, and therefore can be considered a conservative approach to estimating error.

First, to obtain the competition coefficients (α) as the slopes of the relationship between per

capita survival and population density, we must sum individual m parts of the regression model

shown in Table S2 to obtain the species and treatment specific slopes (β_i) according to:

$$f_\alpha = \sum_i^m \beta_i \qquad Eqn\ S1$$

281   Error propagates in the sum of m estimates according to:

$$\sigma_\alpha^2 = \sum_{i=1}^m \sigma_i^2. \qquad Eqn\ S2$$

283   Once the competition coefficients are estimated, the niche and fitness differences are

284   combined into ratios via multiplication and division using Eqns 1 and 2 in the main text. For the

285   purposes of this derivation, we can write this as:

$$f_r = \prod_{i=1}^l \alpha_i, \qquad Eqn\ S3$$

286   where, quotient of A by B is simply written as the product of $A \times \frac{1}{B}$. Error propagates through

287   multiplication or division, for l elements of value α_i to create variance for the internal ratios of

288   competition coefficients according to:

$$\sigma_r^2 = \sqrt{\sum_{i=1}^{l} \left(\frac{\sigma_{\alpha,i}^2}{\alpha_i}\right)^2}. \qquad\qquad Eqn\ S4$$

290   Finally, niche and fitness differences both involve taking the square root of the competition

291   coefficient ratios (f_r). The final variance after taking the square root becomes

$$\sigma_{MCET}^2 = \frac{\sigma_r^2}{2f_r}. \qquad\qquad Eqn\ S5$$

With the final variance for either niche or fitness differences in hand, the standard deviation and

standard error can be used to obtain a confidence interval for both niche and fitness differences.

Checkerboard score

The checkerboard score is named for the appearance of a perfect negative or positive co-

occurrence matrix, which takes on the appearance of a checkerboard based on the diagonal

arrangement of 0s and 1s. Here we detail how to calculate the C-score. First the number of

checkerboard units for each pair of species is given by:

$C_{ij} = (r_i - X_{ij})(r_j - X_{ij}),$ \qquad\qquad Eqn 4

where r_i is the number of times species i occurs without species j, r_j is the number of times species j

occurs without species i and X_{ij} is the number of sites where species i and j co-occur [3]. Typically,

C_{ij} is normalized by the number of possible species pairs (P) and the number of possible site pairs

(N), which using basic combinatorics are given by

$P = S(S-1)/2$ and $N = n(n-1)/2,$ \qquad\qquad Eqns 5

where S is the number of species, and n is the total number of replicate plots [3,16]. Across all sites, the

normalized C-score is given by:

$$C_n = \sum_{j=0}^{S} \sum_{i<j} \frac{C_{ij}}{PN}.$$ Eqn 6

To estimate the null expectation we used a fixed-equiprobable randomization where row

sums (i.e. species occurrences) are held constant, but the column sums (species occurrence within

sites) are allowed to vary with equiprobability [26]. A fixed-fixed randomization is more conservative

and less prone to type-I errors, but due to the rarity of *C. dentata*, it cannot be applied to the

frequent two species communities containing just *P. serotina* and *Q. rubra.* This is because retaining

the row and column sums of a two species matrix rather trivially always returns the original

occurrence matrix when there are only two species. Null community matrices were created using

the permatswap function in the vegan library in R [27,28]. For each observed matrix 1000 null matrices

were sampled with 30000 swaps between each sample and with a burn-in of 30000 swaps prior to

the first sample. We used the quasiswap method for generating null matrices, which does not

produce sequential null matrices, but instead generates a matrix at each time step that is fully

independent of previous matrices [29]. The standardized effect size of the C_n-score can be calculated

based on the observed C_n-score (C_n) and the mean (μ) and standard deviation (σ) of the expected C_n-

scores produced from the 1000 null matrices by calculating a z statistic as [25,30]:

$z = \frac{C_n - \mu}{\sigma}.$ Eqn 7

The z statistic is significant at the two-tailed p=0.05 level for -1.96<z>1.96.

TABLE S1: Regression model for leaf data shown in Fig 1A,B. The model was lm(Leaf.production~
density * diversity * year) in R. The overall regression was not significant ($F_{7,160}$ = 1.66, p=0.1217),
and critically, none of the individual slopes for the diversity productivity relationship were
significantly different from zero.

Coefficients	Estimate	SE	t	p
Intercept	260.6	51.2	5.1	<0.001
Diversity	21.3	27.7	0.77	0.443
Density(2m)	-10.5	57.3	-0.18	0.855
Year(2019)	76.2	72.4	1.0	0.295
Diversity*Density(2m)	6.0	30.9	0.19	0.847
Diversity*Year(2019)	-19.8	39.1	-0.51	0.614
Density(2m)*Year(2019)	-13.0	84.9	-0.15	0.88
Diversity*Density(2m)*Year(2019)	2.71	45.9	0.06	0.953

331
332

333 **TABLE S2:** Regression model for wood productivity shown in Fig 1C,D. The model was lm(BAI ~
334 density * diversity * year) in R. The regression was significant ($F_{43,1925}$ = 24.1, p<0.0001), though the
335 data contained a large amount of unexplained error (R^2 = 0.33). None of the individual slopes for the
336 diversity productivity relationship were significantly different from zero.

Coefficients	Estimate	SE	t	p
Intercept	0.58	2.05	0.284	0.7761
Density(2m)	0.39	2.50	0.154	0.8773
Year(2008)	0.08	2.33	0.034	0.9728
Year(2009)	0.93	2.29	0.405	0.6859
Year(2010)	1.10	2.28	0.484	0.6283
Year(2011)	1.81	2.27	0.799	0.4246
Year(2012)	2.57	2.26	1.134	0.2567
Year(2013)	2.60	2.26	1.147	0.2516
Year(2014)	4.05	2.26	1.788	0.074
Year(2015)	4.29	2.26	1.892	0.0586
Year(2016)	5.24	2.26	2.314	0.0208
Year(2017)	4.79	2.26	2.114	0.0346
Diversity	-0.08	0.94	-0.087	0.9307
Density(2m):Year(2008)	0.28	2.96	0.094	0.925
Density(2m):Year(2009)	0.25	2.88	0.086	0.9312
Density(2m):Year(2010)	0.66	2.87	0.229	0.8193
Density(2m):Year(2011)	1.52	2.86	0.533	0.5943
Density(2m):Year(2012)	2.51	2.85	0.88	0.3787
Density(2m):Year(2013)	1.68	2.85	0.592	0.5541
Density(2m):Year(2014)	1.27	2.85	0.447	0.6546
Density(2m):Year(2015)	2.61	2.85	0.918	0.3585
Density(2m):Year(2016)	1.46	2.85	0.514	0.6076
Density(2m):Year(2017)	2.14	2.85	0.75	0.4531
Diversity:Density(2m)	-0.08	1.16	-0.066	0.9472
Diversity:Year(2008)	-0.01	1.07	-0.009	0.9931
Diversity:Year(2009)	-0.17	1.05	-0.162	0.8712
Diversity:Year(2010)	-0.06	1.05	-0.057	0.9542
Diversity:Year(2011)	-0.25	1.04	-0.239	0.8108
Diversity:Year(2012)	-0.15	1.04	-0.143	0.8866
Diversity:Year(2013)	0.03	1.04	0.026	0.9791
Diversity:Year(2014)	-0.46	1.04	-0.443	0.6576
Diversity:Year(2015)	-0.40	1.04	-0.38	0.7037
Diversity:Year(2016)	-0.69	1.04	-0.659	0.51
Diversity:Year(2017)	-0.26	1.04	-0.25	0.8025
Diversity:Density(2m):Year(2008)	0.02	1.38	0.011	0.9913
Diversity:Density(2m):Year(2009)	0.07	1.34	0.052	0.9587
Diversity:Density(2m):Year(2010)	0.12	1.34	0.09	0.9284
Diversity:Density(2m):Year(2011)	-0.04	1.33	-0.034	0.9727
Diversity:Density(2m):Year(2012)	0.39	1.33	-0.295	0.7681
Diversity:Density(2m):Year(2013)	0.25	1.33	0.186	0.8523
Diversity:Density(2m):Year(2014)	0.81	1.33	0.614	0.5396
Diversity:Density(2m):Year(2015)	0.26	1.33	0.197	0.8438
Diversity:Density(2m):Year(2016)	0.80	1.33	0.606	0.5446
Diversity:Density(2m):Year(2017)	0.82	1.33	0.619	0.5358

338   **TABLE S3:** Regression model for root data shown in Fig 1E,F. The model was lm(Root.production~
339   density * diversity) in R. The overall regression was not significant ($F_{3,56}$ = 0.27, p=0.8476), and
340   critically, none of the individual slopes for the diversity productivity relationship were significantly
341   different from zero.

Coefficients	Estimate	SE	t	p
Intercept	121.5	28.5	4.3	<0.0001
Diversity	-8.3	15.1	-0.55	0.585
Density(2m)	-2.8	34.9	-0.08	0.936
Diversity*Density(2m)	6.7	18.5	0.364	0.717

TABLE S4: Regression coefficients for the logistic curves fit in Fig S2 and used to calculate
competitive ability parameters for niche and fitness differences. The model was glm(survival ~
density * species + treatment*species) in R using a binomial distribution to achieve a logistic
regression. Statistically significant coefficients are shown in boldface font. Modern coexistence
theory calculations involve ratios of competition coefficients. Thus, to avoid dividing by zero, we
dealt with non-significant regression coefficients by incorporating error into our modern coexistence
theory calculations. This allows non-significant parameters to have their appropriately high error.

Coefficients	Estimate	SE	t	p
Intercept	**2.4**	**1.08**	**2.13**	**0.0268**
Density	-0.65	1.44	-0.450	0.6528
Species(*P. serotina*)	**2.18**	**0.88**	**2.48**	**0.0133**
Species(*Q. rubra*)	**2.56**	**0.97**	**2.64**	**0.0082**
Treatment(C)	0.42	1.24	0.34	0.7343
Treatment(Q)	-1.53	1.42	-1.08	0.2797
Treatment(PQ)	**-2.44**	**0.97**	**-2.50**	**0.0124**
Treatment(CP)	-1.19	0.95	-1.25	0.2098
Treatment(CQ)	-1.69	1.16	-1.46	0.1452
Density:Species(*P. serotina*)	**-3.67**	**1.25**	**-2.94**	**0.0033**
Density:Species(*Q. rubra*)	**-4.41**	**1.47**	**-3.00**	**0.0027**
Density:treatment(C)	-0.06	1.65	-0.04	0.9696
Density:treatment(Q)	**4.79**	**1.95**	**2.46**	**0.0141**
Density:treatment(PQ)	**4.65**	**1.31**	**3.53**	**0.0004**
Density:treatment(CP)	1.87	1.13	1.65	0.0999
Density:treatment(CQ)	**3.70**	**1.61**	**2.30**	**0.0218**

FIG S1: Plot layout. **(A)** The experiment has three replicate blocks that include all possible

combinations of *C. dentata* (C, American chestnut), *P. serotina* (P, black cherry) or *Q. rubra* (Q,

northern red oak), at three different planting densities (1m, 2m or 3m between trees; 1 tree/m^2, 0.5

trees/m^2, or 0.33 trees/m^2 respectively) in a full factorial split plot design. Unfortunately, many trees

in the southern most 3m spacing plots died due to an unusual flooding event. These plots are not

included in productivity data due to inadequate sample size for analysis, however these plots are

included in demography data. Surplus chestnut trees were planted in an open space at the southern

end of the experiment because of their conservation value. **(B)** Each plot contains 56 trees that

alternate between species (indicated by 1 and 2). There is a buffer row of trees that are not

measured around each plot (dashed line) leaving 30 focal trees in the center of each plot, and 1260

individual focal trees in 1m and 2m spaced plots. (C) An aerial image of our plots showing the basic

layout. The dead 3m spaced plots, and the extra chestnut trees that are not part of the experiment,

are especially evident in this image. Field vehicles are also visible in the center of the image. Full

methodological details about the planting and experimental design can be found in: [20].

Fig S2: Per capita survival rates (our estimate of fitness) across a ten-year interval for each

monoculture, or two species mixture for all three species: *C. dentata* (C, American chestnut), *P.*

serotina (P, black cherry) or *Q. rubra* (Q, northern red oak). The slope of each line is shown above

and was our estimate of competition coefficients (α) used to calculate stabilizing and equalizing

mechanisms of coexistence. Lines represent logistic generalized linear model fits across all data

(regression coefficients given in Table S2). All y data are either 0 or 1, and points are plotted with a

jitter on x and y to aide in visualisation of the density of points at each position.