Title: Dicamba drift alters plant-herbivore interactions at the agro-ecological interface

Authors: Nia M Johnson[1] and Regina S Baucom[1]

[1]Ecology and Evolutionary Biology Department, 4034 Biological Sciences Building, University of

Michigan, Ann Arbor, MI 48109

*Corresponding author: nmjo@umich.edu

Data are provided as private-for-peer review via the following link:

https://github.com/niajohnson1/DicambaDrift_20182019.git

Abstract

Natural populations evolve in response to biotic and abiotic changes in their environment, which shape

species interactions and ecosystem dynamics. Agricultural systems can introduce novel conditions via

herbicide exposure to non-crop habitats in surrounding fields. While herbicide drift is known to produce a

variety of toxic effects in plants, little is known about its impact on non-target wildlife species

interactions. In a two-year study, we investigated the impact of herbicide drift on plant-herbivore

interactions with common weed velvetleaf (*Abutlion theophrasti*) as the focal species. The findings reveal

a significant increase in the phloem feeding silverleaf whitefly (*Bermisia tabaci*) abundance on plants

exposed to herbicide at drift rates of 0.5% and 1% of the field dose. We also identified a significant

phenotypic tradeoff between whitefly resistance and herbicide resistance in addition to whitefly resistance

and relative growth rate in the presence of dicamba drift after increasing the populations grown in year

two. In a follow-up greenhouse study, we found evidence that dicamba drift at 0.5% of the field dose

significantly increased average chlorophyll content (mg/cm^2) along with a positive correlation between

whitefly abundance and chlorophyll content. Overall, these findings suggest herbicide exposure to non-

target communities can significantly alter herbivore populations, potentially impacting biodiversity and

community dynamics of weed populations found at the agro-ecological interface.

Keywords: agroecology, eco-evolution, plant-insect interactions, herbicide drift, weed adaptations

Introduction

Biotic and abiotic interactions influence species assemblage and evolution throughout ecosystems (Thompson 1999, Klanderud et al. 2015). Often a source of atypical abiotic factors, agricultural practices are largely dependent on pesticide use for increased crop yield and labor efficiency (Gianessi 2013). Despite the economic benefits of herbicides for weed control, however, their use can lead to unintentional impacts on non-target wildlife if the herbicide 'drifts' or migrates outside of the target area during or after the initial application (Freemark and Boutin 1995, Carlsen et al. 2006a, Carlsen et al. 2006b). Though herbicide drift is known to produce a variety of toxic effects on non-target vegetation (Marrs et al. 1991, Fletcher et al. 1996, Marrs and Frost 1997, Gove et al. 2007, Boutin et al. 2014, Cederlund 2017), and can lead to significant plant phenotypic and compositional changes (Iriart, Baucom, and Ashman 2020), few studies have evaluated how herbicide drift may disrupt or influence the interactions between plants and other community members such as pollinators and herbivores.

Weeds and other non-crop plants found at the edges of crops serve as important reservoirs of insect biodiversity (Egan and Mortensen 2012), which provides crucial ecosystem services for agriculture such as pollination and pest control (Daily 1997). The plant- insect interactions that occur at the agro-ecological interface are key determinants for the movement of energy and nutrients as well as drivers of ecological and evolutionary dynamics (Ehrlich and Raven 1964, Futuyma and Agrawal 2009). However, weed communities at the agro-ecological interface can be exposed to herbicide drift rates between 0.1% and 5% of the normal field application rate of herbicide (Cessna et al. 2005), and we currently understand very little about how this drift may influence ecological interactions and potentially lead to subsequent evolutionary responses. For example, does herbicide drift change herbivore abundance or damage, and/or lead to altered patterns of plant investment in resistance to either herbicide or herbivore damage?

One herbicide known for non-target damage is dicamba (3,6-dichloro-2-methoxybenzoic acid) (Jones et al.

2019). Dicamba is a broadleaf selective synthetic auxin that is on an upwards trajectory in agriculture due to

a substantial increase in the adoption of dicamba-tolerant crops (USDA-ERA, 2019), which is evident by an

estimated 600% increase of use in the US between 2014 and 2019 (Baker, 2021). Dicamba is absorbed by

plant leaves and mimics the deformative and growth-altering effects caused by overdoses of the natural plant

auxin, indole-3- acetic acid (IAA) (Grossman 2010). Dicamba drift can cause a shift in plant and arthropod

diversity (Egan et l. 2014), delay the onset of flowering, reduce the number of flowers, and even reduce the

amount of pollinator visits (Bohneblust et al. 2016). However, we have very little information on how

dicamba drift may impact plant-herbivore interactions. A broad expectation is that exposure to a synthetic

auxin, such as dicamba, could alter plant allocation towards growth and defense, and potentially lead to

increased vulnerability to herbivores, in line with growth-defense tradeoff framework (Coley, Bryant, Chapin

1985). Recent work has shown that exposure to drift rates of herbicides like dicamba can lead to the

evolution of reduced herbicide sensitivity (Vieria et al. 2020) and recurrent selection of low dose herbicides

can lead to non-target site resistance (Busi et al. 2013), however, there is little understanding of how this

process may influence selection for allocation herbivory defense.

Here we examined the impact of dicamba drift on plant herbivory, dominated by naturally occurring

populations of the phloem feeding silverleaf whitefly (*Bermisia tabaci*) in the field, using the crop weed

velvetleaf, *Abutilon theophrasti*. First introduced to the United States from Asia in the late 17th century,

this annual species is now one of the most common broad-leaved species in and around corn and soybean

agricultural fields located in Canada, Europe, and the United States (Spencer 1984). While past work in

Amaranthus hybridus has shown that lines with evolved herbicide resistance (defined by survival given

application of herbicide at the recommended field dose) can exhibit reduced herbivory resistance

depending on light availability (Gassman 2005), there is a striking lack of information on how such a

phenotypic tradeoff evolves when populations are exposed to drift rates under natural field conditions.

We performed field experiments to address this gap in our knowledge, and specifically asked the

following questions: Does the application of dicamba drift influence herbivory in the common weed

Abutilon theophrasti (velvetleaf)? Does the amount of herbicide damage, which reflects resistance to

herbicide, influence the amount of herbivory, which reflects resistance to herbivory? (*i.e.*, is there a

trade-off between these two forms of defense in the presence of dicamba drift)? Is either herbicide or

herbivore resistance negatively correlated with relative growth rate, indicating a growth- defense

tradeoff? Is there evidence of selection acting directly on herbicide resistance, herbivory resistance, or

relative growth rate or evidence of correlative selection acting on the interactions of these focal traits? Does

herbicide drift impact leaf chlorophyll levels, and if so, does that correlate with whitefly abundance?

Methods

Field Experimental Design

We performed two consecutive field experiments to characterize the extent of herbicide drift damage and

herbivory response in *A. theophrasti*. In the first experiment, we planted replicate seeds from 23 maternal

lines originally sampled from a single population (Dexter, MI) in the summer of 2018. We increased the

number of populations from 1 to 8 for a total of 50 maternal lines in the second field experiment (SFig 1),

including the 23 maternal lines from the originally sampled population and 2-5 maternal lines from the

subsequently sampled populations. All seeds were collected in the Fall prior to each summer in and around

soybean fields located in Dexter, MI.

In the first field experiment (2018) we scarified and planted 276 total seeds in a randomized block design

(23 maternal lines x 3 treatments x 2 replications x 2 blocks) with each treatment randomized within each

block, and two replicate plants per maternal line present within each treatment/block combination. Our

three experimental treatments were two levels of a drift exposure (1% and 0.5% of the recommended field

dose of dicamba (3,6-dichloro-2-methoxybenzoic acid) (2.8 g ae/ ha dicamba, 5.6 g ae/ha dicamba) and

water as the control. For the second field experiment (2019), we scarified and planted 450 total seeds in a

randomized block design with the same treatments as the previous field experiment (50 maternal lines x 3

treatments x 3 blocks). Treatments were randomized in each block, and one replicate of each maternal

line was present within each treatment/block combination. We recorded leaf count, height, and largest

leaf width for all plants biweekly in both experiments. We recorded flower count weekly and sampled all

seeds at the end of the season for an estimate of fitness. Five weeks after seed germination, when the

average plant height was 11 cm tall, we applied 1% and 0.5% dicamba to plants in each respective

treatment with a hand-held pressurized prayer for roughly 3 seconds which ensured adequate coverage of the

entire plant.

Herbicide Damage – We assessed herbicide damage two weeks following herbicide application by

recording the proportion of leaves exhibiting visual deformation (curling or cupping), and proportion of

the plant damage was determined by evaluating the number of leaves with a cupping or curling pattern

divided by the total number of leaves.

Herbivory Damage – To estimate resistance to herbivory in 2018, we assessed both physical damage from

chewing insects and the abundance of phloem-feeding insects (dominated by silverleaf whiteflies).

Because there was little evidence of chewing insects, only whitefly abundance was assessed in 2019. We

assessed chewing herbivory damage five weeks after herbicide application using the imaging software,

Fiji ImageJ version 3.0 (Schindelin et al. 2012). To do this, we collected 3 leaves at random from each

plant and scanned them using the Canon CanoScan L110. To obtain the total surface area, we converted

each image to binary format and used the particle analysis tool which quantifies total pixel number into

centimeters. We then used a macro plug-in to place a grid with $0.1cm^2$ spacing over each image to

estimate the amount of chewing herbivore damage per leaf.

Visual estimates of whitefly abundance were sampled in both field experiments by selecting 10 leaves at

random per plant. Because most whiteflies feed and oviposit at the same location (Van Lenteren and

Noldus 1990), estimates of percentage of leaf area covered by whitefly oviposits were captured by

visually partitioning each leaf into quarters and evaluated for each section, as suggested in Johnson et al

2015. We estimated whitefly abundance on a scale from 0 – 5 (adapted from Banks 1954). A score of 0

meant no whitefly oviposits were present, 1 was 1 - 20% leaf coverage by whitefly oviposits, two was

between 21 – 40 % leaf coverage, three was 41 - 60 % leaf coverage, four representing 61 - 80% leaf

coverage, and five representing 81 - 100% leaf coverage.

Photosynthetic carbon dioxide response – To determine if there was a link between herbicide resistance

and photosynthetic response to dicamba drift, we estimated photosynthetic efficiency by examining

variation in photosynthetic carbon dioxide response curves (A-Ci Curve) using the Li-Cor 6400 portable

photosynthesis system (Open System Vers, 4.0, Li-Cor, Inc., Lincoln NE) in 2018. During gas exchange

measurements, we maintained cuvette conditions at a photosynthetic photon flux density (PPFD) of 1500

$\mu mol/ m^2 s$, air flow rate at 500 μmol/s, and leaf chamber block temperature being 30°C to match ambient

air temperature. In order to measure the gas exchange, we first set the CO_2 concentration reference at 400

μmol/mol, and we maintained the leaf under such conditions for ten minutes for adaptation and

stabilization of leaf photosynthesis. We set a loop of changing reference CO_2 concentration at 400, 300,

200, 100, and 50μmol/mol. We controlled CO_2 concentration in the cuvette with a CO_2 mixer across this

series and measurements were recorded after equilibration to a steady state. We then set another loop of

changing reference CO_2 concentration at 400, 600, 800, 1000, 1300, and 1500 μmol/mol and again logged

at each iteration. Between each group of measurements, we set the reference CO_2 concentration back to

400 μmol/mol for 5 minutes. These measurements were made four to five weeks after herbicide

application, within a ten-day period. We recorded these data on a leaf area of 2 cm^2, from 2 leaves on

each of three plants from each treatment. For plants growing in drift treatments, we included one leaf that

developed prior to herbicide application (Leaf 1), and one leaf that developed after herbicide application

which as a result exhibited a deformation in shape (Leaf 2). The purpose of taking measurements on

leaves with different times of development was to investigate if potential herbicide sequestration impacts

photosynthetic efficiency of leaves exhibiting damage, potentially impacting herbivore resistance

mechanisms for the entire plant. Measurements for each leaf per plant were taken on the same day.

Greenhouse Experimental Design

With the goals of understanding treatment effects on whitefly abundance (*i.e.* impacts on host-plant

selection), we conducted a greenhouse experiment in the summer of 2021. We planted 6 replicates of 20

maternal lines randomly selected from 5 of the populations used in the 2019 field experiment totaling 240

seeds (20 maternal lines x 6 experimental replicate seeds x 2 treatments). Half of the plants were directly

treated with dicamba at 0.5% of the field dose, while the other half was treated with water 4 weeks

following planting. In preparation for herbicide application, plants were placed in rows outside of the

greenhouse in order to prevent herbicide drift. We subsequently applied the herbicide solution in the same

manner as in the field experiments and relocated the plants back into the greenhouse once they had visually

dried. We elected to treat plants with the herbicide one week prior to the field experiment

due to observed accelerated growth in the greenhouse. Because whiteflies are common pests in many

greenhouses, and the origin of the whitefly population in the field had previously been connected to crops

transplanted from the greenhouse in plots adjacent to our own, we elected to allow whiteflies to naturally

migrate to velvetleaf plants without manipulation or introduction.

Chlorophyll Content – We estimated whitefly abundance with the same methodology as above. Since

whitefly host selection has previously been linked to leaf wavelength emissions (Mound 1962), using the

atLEAF Plus Digital Chlorophyll Meter, we measured the chlorophyll levels (mg/cm^2) of 3 leaves per plant

chosen at random and calculated the average for each plant.

Data Analysis

Herbivory Response to Drift

We conducted all statistical analyses in R studio (version 3.4.1, R Development Core Team). For each

type of herbivory measurement in the field experiments (chewing damage and whitefly abundance), we

performed analysis of variance to determine whether herbivory resistance differed in response to dicamba

application for each year separately. Prior to performing ANOVAs, we transformed response variables

using Tukey's Ladder of Powers in the rcompanion package (Mangiafico 2016) to correct for non-

normality of residuals. In 2018 (n = 118), we fit the following mixed linear model using lmer function of

the lmer4 package (Bates et al. 2011):

y = u + treatment + block + (treatment x block) + (1|maternal line) + (1|maternal line x treatment) + e

where, y, the response variable, is chewing damage or whitefly abundance, u is the intercept or mean of

the model, treatment and block are fixed-effect terms, maternal line and maternal by treatment are random

effect terms. In 2019 (n = 240), we nested maternal line into population and added population as a random

effect to the existing model: y = u + treatment + block + (treatment x block) + (1|population) + (1|population:

maternal line) + (1|population: maternal line) x treatment + e. We determined the significance of the predictor

variables using F-statistics for the fixed effects with Kenward-Roger to estimate the degrees of freedom and

used a log-likelihood ratio test to estimate χ^2 for the random effect. We then used a Welch two sample t-test

to determine if there were significant differences between the two dicamba drift treatments.

Plant Performance in Response to Drift

To determine how drift treatments influenced plant performance in the field, we ran separate mixed linear

models as described above by year. In 2018, we fit separate models for each plant trait (height, leaf count,

leaf width, and flower number): y = u + treatment + block + (treatment x block) + (1|maternal line) +

(1|maternal line x treatment) + e; where each trait was the response variable, block, treatment, and block

by treatment interactions were fixed effects, and maternal line, and maternal by treatment interactions

were random effects. In 2019, we nested maternal line into population and added population as random

effects to the existing model: y = u + treatment + block + (treatment x block) + (1|population: maternal

line) + (1|population: maternal line) x treatment + (1|population) + (1|population) x treatment + e. We

transformed each trait again using Tukey's Ladder of Powers to meet the assumptions of normality. We

performed separate analyses for each year because preliminary analysis revealed a significant difference

in plant growth between years.

Additionally, photosynthetic carbon dioxide response means were analyzed for 2018 using a Kruskal-

Wallis test with treatment as the independent variable. We performed exact Wilcoxon rank sum tests on

photosynthetic carbon dioxide response means of individuals grown in each drift treatment separately to

determine if time of leaf development impacted photosynthetic carbon dioxide response among leaves

with and without deformation by including leaf developmental time as the independent variable.

Genetic Variation for Herbicide Resistance

We examined herbicide resistance for each year separately and were explicitly interested in determining if

there was evidence for genetic variation for resistance to herbicide drift *via* a population or maternal line

by treatment interaction. We fit the following separate mixed linear models for proportion of plant

damaged from herbicide: y = u + treatment + block + (treatment x block) + (1|population: maternal line)

+ (1|population: maternal line) x treatment + (1|population) + (1|population) x treatment + e; where each

type of resistance was the response variable, block, treatment, and block by treatment interactions were

fixed effects, and maternal line, maternal by treatment interactions, population, and population by

treatment interactions were random effects. We determined if there was evidence of genetic variation for

herbicide resistance and whitefly resistance by performing log-likelihood ratio tests to estimate χ^2 for

maternal, maternal line by treatment, population, and population by treatment interactions for each

resistance type. Finally, because we are interested in potential growth-defense trade-offs, we used a

similar mixed linear model to examine genetic variation in relative growth rate.

Operational Definition of Resistance

We define herbicide resistance operationally as 1 minus the proportion of leaves exhibiting damage from

the herbicide, which was observed as yellowing and curling/deformation of leaves. Because there was

limited damage from chewing herbivores, we elected to focus on phloem-feeding herbivory, dominated

by whiteflies. Whitefly resistance was defined as 1 minus the amount of whitefly larvae coverage per

individual (as described previously).

Phenotypic Correlations

In order to test for correlations between herbicide resistance and whitefly resistance (defined

operationally as above), we performed Pearson's correlation test. We examined correlations for each

experimental year separately.

Phenotypic Selection

In accordance with the Lande–Arnold approach (e.g., Arnold et al. 2001; Arnold 2003; Hereford et al.

2004), we used phenotypic measurements to quantify natural selection on whitefly resistance, herbicide

resistance and relative growth rate. Here, we report selection gradients rather than differentials to understand

whether direct selection is acting on individual traits by controlling for any indirect selection

present. Relative fitness was calculated as the final seed count per individual divided by mean seed count

for each treatment per year (2018: n = 118; 2019: n = 240). We estimated linear (directional) selection

gradients (β) within drift environments (0.5% and 1% the field dose of dicamba) by performing multiple

linear regressions of relative fitness on whitefly resistance, herbicide resistance, relative growth rate

(calculated as change in leaf count plus change in height divided by the number of weeks spent growing)

and their interactions separately for each year. We also estimated nonlinear selection gradients (γ) in a full

model that included linear terms, quadratic terms, and the cross-product terms of focal traits. Quadratic

regression coefficients were doubled to estimate nonlinear selection gradients. Nonlinear selection

gradients examined the potential for selection on phenotypic variance of a trait (quadratic selection) or

phenotypic covariance (correlated selection) between focal traits, as there is evidence of relationships

between plant stress, defense, and growth (Huot et al. 2014, Zust and Agrawal 2017). For all selection

analysis we mean standardized focal traits (i.e., subtracted the mean and divided by the standard

deviation) and used untransformed values of relative fitness. Furthermore, we estimated linear and

nonlinear selection on whitefly resistance in the absence of herbicide drift in order to determine whether

herbicide altered the pattern of selection. We compared selection gradients between the two treatment

types using an ANCOVA to perform a regression for relative fitness on whitefly resistance, treatment,

and their interaction.

Greenhouse Experiment - Chlorophyll Content

To investigate the effects of dicamba on chlorophyll content in the greenhouse, we performed analysis of

variance on the plants grown in the greenhouse. We fit the following simple linear model: $y = u +$

treatment $+ e$; where, y, the response variable, is chlorophyll average, u is the intercept or mean of the

model, and e is the error term. In order to test for correlations between chlorophyll averages and whitefly

abundance, we performed Pearson's correlation test.

Results

Herbivory Response to Drift Exposed Host

We found no evidence that dicamba drift influenced the amount of chewing damage experienced by *A.*

theophrasti (Fig 1A; STable 1) but we did find a significant increase in whitefly abundance on dicamba

treated plants (F-value = 12.01, p-value < 0.01, Fig 1B; STable 1). This effect was also present in the

second field season in which we included five more *A. theophrasti* populations (F-value = 5.30, p-value <

0.01, Fig 1C; STable 1). Both levels of dicamba drift -- 0.5% and 1% of the suggested field dose --

exhibited higher whitefly abundance than the controls. Plants exposed to 0.5% dicamba had significantly

higher whitefly abundance than 1% dicamba treatments across both years (2018: t = 2.74, p < 0.01, Fig

IB; 2019: t = 2.06, p = 0.04, Fig 1C).

Plastic Response to Herbicide Drift on Plants

We examined several plant traits in response to dicamba drift in order to determine how drift impacts

velvetleaf growth and physiology, which may underscore herbivory responses. Across both experimental

years, we found that leaf count increased following the application of dicamba drift (Treatment effect for

2018: F-value = 28.01, p-value < 0.01, Fig 2; Stable2; 2019: F-value = 5.11, p-value < 0.01, Fig 2; STable

3), whereas leaf width decreased significantly in 2018 but not 2019 (Treatment effect for 2018: F-value =

8.25, p-value < 0.01, Fig 2; STable 2; 2019: F-value = 0.79, p-value = 0.46, Fig 2; STable 3), which may

be a result of significant population variation in 2019 for leaf width (χ^2 = 5.11, p-value = 0.02, STable 3).

Both plant height and flower count decreased as a result of dicamba drift in both experiments (Treatment

effect: Height 2018: F-value = 220.56, p-value < 0.01; Flower Count 2018: F-value = 7.28, p-value <

0.01, Fig 2; STable 2; Treatment effect: Height 2019: F-value = 2.84, p-value = 0.06; Flower Count 2019:

F-value = 6.92, p-value < 0.01, Fig 2, STable 3). Typically, plant leaf number was greater, and height and

flower count were lower in the 1% dose of dicamba drift compared to the 0.5% dose (Fig 2, STable 2).

We found evidence for block effects and block by treatment effects in the first field experiment across

most phenotypic traits, but these effects were less evident in the second field experiment (STable 3).

Finally, we found no evidence for maternal line or population effects associated with dicamba drift

exposure, indicating that at least in this sample of 50 maternal lines, there was no evidence that the plastic

response for height, leaf count, leaf width, and flower count varied genetically.

Photosynthetic carbon dioxide response (A-C_i, umol/m$_2$), which was examined in the 2018 experiment,

differed across treatments, indicating that dicamba drift significantly impacts photosynthetic carbon

dioxide response overall (x^2 = 37.605, p < 0.001; SFig 2;). Within drift treatments, we found

evidence that time of leaf development significantly impacts A-Ci curves (0.5%: W = 696, p < 0.001 SFig

2; 1%: W = 360, p < 0.001). In general, leaves exposed directly to 0.5% of the field dose of dicamba

exhibited higher A/Ci responses than their counterparts that developed after dicamba treatment (SFig 2).

In contrast to the lower dosage, leaves treated directly with 1% of the field dose of dicamba exhibited

lower A/Ci responses than their counterparts that developed after dicamba exposure (SFig 2). This result

indicates high amounts of photosynthetic plasticity in response to dicamba, potentially associated with

altered herbicide translocation and metabolism contributing to resistance to drift damage.

Chlorophyll Content

In the greenhouse, we uncovered a significant treatment effect on chlorophyll levels (mg/cm^2), where

plants exposed to dicamba at drift levels had on average higher chlorophyll levels than the controls (F =

4.56, p = 0.03, Fig 3A). We also identified a significantly positive correlation between average

chlorophyll values and whitefly abundance (r = 0.25, p < 0.001, Fig 3B), suggesting that chlorophyll

content impacts whitefly host-selection on velvetleaf.

Genetic variation for whitefly resistance, herbicide resistance, and relative growth rate

We did not detect evidence of maternal line, maternal line by treatment, population, or population by

treatment effects on whitefly resistance, herbicide resistance or relative growth rate in either year.

(2018: STable 4, 2019: STable 5). This suggests there is not significant genetic variation in the

populations we sampled from, likely due, at least in part, to the low number of maternal lines included in

the study.

Phenotypic correlations

We did not detect evidence of phenotypic correlations between any of the focal traits in 2018 (*i.e.*

herbicide resistance, whitefly resistance, and relative growth rate), but in 2019 we found evidence of a

moderately strong negative phenotypic correlation between herbicide resistance and whitefly resistance

within drift environments (r = -0.32, p > 0.001). This suggests that both types of resistance may be

indirectly impacting one another. We also found a negative correlation between whitefly resistance and

relative growth rate in the presence of drift but not in its absence in 2019 (drift environment: r = -0.22, p =

0.02; control environment: r = -0.07, p = 0.58), indicating a tradeoff between growth and herbivory

defense when exposed to herbicide drift.

Phenotypic selection

In 2018, we did not identify evidence of linear selection on herbicide resistance nor whitefly resistance

(herbicide resistance: β = -0.19, p = 0.57; whitefly resistance: β = 0.08, p = 0.30; STable 4) in the

herbicide drift environment, though we did identify selection acting on relative growth rate (β = 1.71, p <

0.001; STable 4). We further detected marginal evidence of quadratic selection acting on whitefly

resistance (γ = -0.19, p = 0.09, STable 4) and marginal correlative selection acting on whitefly resistance

and relative growth rate (γ = 0.82, p = 0.07, STable 4). In 2019, we again found no evidence of linear

selection in the herbicide drift environment and identified positive selection on relative growth rate

(whitefly resistance, β = 0.002, p = 0.99; herbicide resistance, β = 0.05, p = 0.72; relative growth rate, β =

2.17, p < 0.001; STable 5). Patterns of correlative selection differed in 2019 compared to the 2018

experiment in that we detected marginal evidence of correlative selection acting on herbicide resistance

and relative growth rate (γ = 0.66, p = 0.08, STable 5). Overall, though we found a significant negative

phenotypic correlation between resistance to herbicide and whitefly herbivory, indicating a trade-off

between the two types of resistance (at least in 2019) we found no evidence of correlative selection acting

on the two traits in either year (2018: γ = 3.36, p = 0.14; STable 4; 2019: γ = 0.29, p = 0.46; STable 5).

Discussion

Ecological effects -- dicamba drift increases whitefly abundance and alters plant morphology and

physiology

Our study provides the first evidence that dicamba drift can increase phloem feeding herbivore abundance

in the field, a finding that should be of concern to agriculture more broadly given the negative effects of

whiteflies on crops (*e.g.*, viral transmission mediated by whitefly - Hogenhout 2008, Legg et al. 2014,

Ning et al. 2015, Sundararaj et al. 2018, Moodley et al. 2019). In two separate field experiments, velvetleaf

exposed to dicamba drift showed higher whitefly abundance in comparison to their control counterparts.

These findings indicate dicamba drift can increase susceptibility to phloem feeding herbivorous insects

such as whitefly, which is aligned with previous work showing increased abundance of English green

aphids feeding on dicamba treated barley in a glasshouse experiment (Hintz 1971).

Additionally in the second year of our study, we found evidence of significant phenotypic tradeoffs between

whitefly resistance and herbicide resistance as well as between whitefly resistance and relative growth rate in

the presence of drift. Growth rate has long been linked to investment towards herbivory defense, suggesting

that when resources are limited competition favors fast-growing plant species that allocate less towards

herbivore defenses (Coley, Bryant, and Chapin 1985). In the literature, this is evident by positive correlations

between growth rate and herbivory damage (Coley 1987, Cebrian and Durate 1994, Fine et al. 2006) and

negative correlations between biomass accumulation and defense chemicals, such as salicylic acid (Meyer

et al. 2007), a plant hormone specifically involved in defense against phloem-feeding silverleaf whiteflies

(Kempema 2007). Such negative correlations suggest that pools of herbivore defenses can be depleted as

strong growth occurs under certain environmental conditions. While our results reveal no correlation

between growth rate and herbicide resistance, a significant correlation between growth rate and whitefly

resistance when under drift induced stress supports the growth-defense tradeoff expectations of depleted

reservoirs of herbivore defenses in the presence of high growth.

Because plant architecture is known to influence herbivore performance and abundance (Jaenike 1978,

Haysom and Coulson 1998, Schlinkert et al. 2015), we assessed the morphological and physiological

response of velvetleaf to dicamba drift. While our results indicate dicamba at drift levels stunts velvetleaf

height, this synthetic auxin also increased the number of leaves produced, although they were typically

smaller in size compared to non-sprayed control plants. Additionally, we examined the impact of dicamba

on A/Ci response curves, as photosynthetic capabilities are linked to natural auxin presence (McAdam et

al. 2007) and are a key driver for plant metabolism. The pattern of A/Ci response differed between drift

treatments and time of leaf development, indicating a high amount of photosynthetic plasticity which

varied over herbicide exposure and time. These findings align with a recent study that showed dicamba

exposure initially decreased photosynthesis by 22% in *Palmer amaranth,* which then improved over time

(Browne et al. 2020). Moreover, there was no evidence of maternal line nor population variation for

morphological or physiological responses to herbicide drift, suggesting a high level of plasticity among

all traits measured across individuals.

Natural auxins are involved in chlorophyll accumulation (Yuan et al. 2018). However, the addition of

synthetic auxins, such as dicamba applied at 50% of the field dose (280g ae) causes greater than 70%

reduction in chlorophyll content (Turgut 2007). Our results show that dicamba applied at drift rates (0.5%

the field dose, i.e., 2.8g ae) can *increase* the average amount of chlorophyll in plants which could impact

host- selection of whitefly. These findings are consistent with recent studies in which the greenhouse whitefly

laid more eggs on leaves with higher chlorophyll and higher nitrogen (Park et al. 2009, Tsueda 2014) due to

the highly linked nature of chlorophyll content and chloroplast thylakoids, which represent a large proportion

of leaf nitrogen content (Evans 1983, Evans 1989), but conflicts with past work reporting whitefly

attraction to yellow plants (Mound 1972, Van Lenterern and Noldus 1990). This suggests that color alone

may not be the primary indicator for whitefly host-selection with every species. Although further studies

on this dynamic are warranted, our results suggest that one mechanism behind the increase of whiteflies

on dicamba-drifted plants is due to the auxin-induced increase in chlorophyll content of leaves.

Evolutionary effects -- correlated evolution between resistance and relative growth rate

In addition to morphological and physiology responses, we also examined the potential that resistance to

both dicamba drift and whitefly damage could evolve within these populations and investigated the

potential for correlative selection between traits since we identified significant negative phenotypic

correlations between herbicide resistance and whitefly resistance along with whitefly resistance and

relative growth rate in 2019. In 2018, we detected marginal significance for correlative selection acting on

whitefly resistance and relative growth rate. Correlative selection was also marginally detected in 2019

between relative growth rate and herbicide resistance in the presence of drift. While this could imply

populations may evolve to an optimum fitness at intermediate values of relative growth rate, whitefly

resistance and herbicide resistance, we did not uncover evidence of genetic variation within these

populations. We further did not uncover significant maternal line variation for resistance traits, likely due

to high amounts of plasticity within this species and/or an insufficient number of individuals per maternal

line per treatment. Environment may also be a factor as previous studies in *Amaranthus hydrius* have

shown that the fitness cost of herbicide resistance associated with increased herbivore susceptibility can

be environmentally dependent (Gassmann 2005). Although we did not detect maternal line variation,

more work here should be done investigating selection acting on the relationship between herbicide

resistance, whitefly resistance, and relative growth rate with a greater number of maternal lines from more

distant populations and environments under consideration.

Conclusion

Herbicide drift from synthetic auxins can shift herbivorous insect host-selection at

the agro-eco interface. Such modifications of herbivore behavior have the potential to impact plant

community composition and nutrient dynamics (Schowalter 2006, Belovsky & Slade 2000) and may

likewise, reciprocally influence insect herbivore populations. Increases in insect herbivore abundance may

result in positive feedbacks where elevated consumption increases nutrient cycling and thus stimulates

insect population growth, which can directly affect the abundance of other members of the community

such predators and pollinators (Forkner & Hunter 2000, Hunter 2001, Ceulemans 2017).

Furthermore, our finding that dicamba drift significantly increases whitefly abundance on velvetleaf

populations could mean dicamba-treated weeds act as a reservoir for whitefly populations, potentially

leading to negative impacts on agricultural yields. If our results are applicable more broadly, plants

exposed to dicamba drift in nature may be preferred host for whitefly populations. As agricultural pests

that colonize more than 600 host plants globally (Byrne and Bellows 1991), whiteflies are known vectors

for transmitting over 70% of the world's plant viruses (Hogenhout 2008). Given the projected expanded

use of dicamba, there is a clear and urgent need to examine to what extent community dynamics may shift

as a result of dicamba use, especially for communities existing at the intersection of natural and managed

vegetative system.

Figures

Figure 1. Chewing damage (A) and whitefly abundance (B and C) measurements in response to dicamba drift: 0% field dose (peach), 0.5% field dose (green), and 1% field dose (blue). A) Chewing herbivory damage summarized by dicamba treatment from the first field experiment in 2018. Damage was measured as area with chewing damage/total leaf surface area. B) Treatment effect on Whitefly abundance summarized by treatment in both the B) 2018 and C) 2019 field experiments. Whitefly abundance was measured as visual estimates of percent larvae area/total leaf surface area. Each graph illustrates median values and confidence intervals.

Figure 2. Plant size measurements in response to dicamba drift environments: 0% field dose (peach), 0.5% field dose (green), and 1% field dose (blue). Shows treatment effect for both years on velvetleaf traits: A) height B) leaf count C) leaf width and D) flower count. Each graph illustrates median values and confidence intervals.

Figure 3. *Greenhouse Experiment* A) Average chlorophyll content per individual in response to dicamba

application of 0.5% of the field dose (F = 4.56, p = 0.03). B) Relationship chlorophyll average and

whitefly abundance estimated as Pearson product-moment correlation; Control treatment (peach) r = 0.34,

p = 0.002; Drift treatment (blue) r = 0.21 p = 0.01. Each graph illustrates median values and confidence

intervals.

Supplementary Tables and Figure

Treatment and Block impact on Herbivory

Herbivory Response	Chewing Damage (2018)		Whitefly Abundance (2018)		Whitefly Abundance (2019)	
Fixed Effects:	F	P	F	P	F	P
Treatment (df = 2)	0.906	0.410	**12.014**	**< 0.001**	**5.300**	**0.006**
Block 2018 (df = 1) 2019 (df = 2)	1.236	0.270	1.656	0.201	1.356	0.261
Treatment X Block 2018 (df = 2) 2019 (df = 4)	0.109	0.897	0.363	0.697	1.076	0.370
Random Effects:	$\chi2$	P	$\chi2$	P	$\chi2$	P
Population (df = 1)					0.000	1.000
Maternal Line (df = 1)	0.498	0.480	0.000	1.000	0.535	0.464
Maternal Line X Treatment (df = 1)	0.000	1.000	0.000	1.000	0.000	1.000

STable 1. Influence of herbicide treatment on chewing damage for 2018 and whitefly abundance for both

years, analyzed using F-statistics values showing effects of treatment, block, treatment by block

interactions, and likelihood ratio test statistics (x^2) showing maternal line variation on herbivory

measurements. In 2019 maternal lines are nested within populations. Significant values are expressed in

boldface.

2018 Herbicide impacts on Velvetleaf Morphology

2018								
Trait	Height		Leaf Count		Leaf Width		Flower Count	
Fixed Effects:	F	P	F	P	F	P	F	P
Treatment (df = 2)	**220.555**	**< 0.001**	**28.012**	**< 0.001**	**8.246**	**0.001**	**7.284**	**0.001**
Block (df = 1)	0.278	0.599	**7.627**	**0.007**	**6.151**	**0.014**	**5.421**	**0.022**
Treatment X Block (df = 2)	**60.804**	**< 0.001**	**15.714**	**< 0.001**	**3.554**	**0.032**	**4.287**	**0.016**
Random Effects:	$\chi2$	P	$\chi2$	P	$\chi2$	P	$\chi2$	P
Maternal Line (df = 1)	0.000	1.000	0.000	1.000	0.000	1.000	0.000	1.000
Maternal Line X Treatment (df = 1)	0.000	1.000	1.986	0.159	0.000	1.000	0.054	0.816

474   **STable 2.** 2018 influence of herbicide treatment on velvetleaf traits, analyzed using F-

475   statistics values showing the effects of treatment, block, treatment by block interactions, and likelihood

476   ratio test statistics **($\chi2$)** maternal line and maternal line by treatment interactions on variation of plant

477   phenotypes. Significant values are expressed in boldface.

478   2019 Herbicide impacts on Velvetleaf Morphology

	2019							
Trait	Height		Leaf Count		Leaf Width		Flower Count	
Fixed Effects:	*F*	P	*F*	P	*F*	P	*F*	P
Treatment (df = 2)	2.844	0.061	**5.113**	**0.007**	0.791	0.458	**6.916**	**0.001**
Block (df = 2)	0.144	0.866	0.700	0.498	2.398	0.096	1.313	0.271
Treatment X Block (df = 4)	0.551	0.699	0.506	1.697	1.763	0.188	0.470	0.758
Random Effects:	χ2	P	χ2	P	χ2	P	χ2	P
Population (df = 1)	0.000	1.000	0.632	0.427	5.112	0.024	0.397	0.528
Maternal Line (df =1)	0.000	1.000	0.000	1.000	0.000	1.000	0.166	0.684
Maternal Line X Treatment (df = 1)	0.000	1.000	0.000	1.000	0.000	1.000	0.000	0.986

STable 3. Influence of herbicide treatment on velvetleaf traits, analyzed using F-statistics

values showing the effects of treatment, block, treatment by block interactions, and likelihood ratio test

statistics **(χ2)** showing population, maternal line, population by treatment interactions, and maternal line

by treatment interactions on variation of plant phenotypes. Maternal lines were nested within populations.

Significant values are expressed in boldface.

2018 Selection Analysis

2018								
Trait	β	SE	P	γ	SE	P	*r*	P
Whitefly Resistance	0.081	0.077	0.299	-0.191	0.055	0.087	0.060	0.622
Herbicide Resistance	-0.191	0.330	0.565	3.464	1.056	0.107	-0.123	0.174
Relative Growth	**1.715**	**0.218**	**< 0.001**	**3.912**	**0.558**	**< 0.001**	**0.750**	**< 0.001**
Whitefly Resistance X Herbicide Resistance				0.712	0.334	0.291		
Whitefly Resistance X Relative Growth				0.816	0.222	0.072		
Herbicide Resistance X Relative Growth				3.364	1.109	0.136		

STable 4. 2018 Selection analysis showing direct selection on focal traits: whitefly resistance, herbicide

resistance, and relative growth rate. Linear (β) ($R^2 = 0.510$; $p < 0.001$) and quadratic (γ) (R

$^2 = 0.609$; $p < 0.001$) selection gradients with associated standard errors (SE) and *P*-values (*P*). The (r)

column represents correlation coefficients between trait and fitness, estimated as Pearson product-

moment correlations. Significant values are expressed in boldface.

2019 Selection Analysis

	2019							
Trait	β	SE	P	Y	SE	P	r	P
Whitefly Resistance	0.002	0.190	0.990	0.232	0.230	0.526	-1.56	0.098
Herbicide Resistance	0.052	0147	0.722	-0.133	0.116	0.759	-0.004	0.955
Relative Growth	**2.169**	**0.245**	**< 0.001**	0.774	0.353	0.363	**0.694**	**< 0.001**
Whitefly Resistance X Herbicide Resistance				0.285	0.190	0.455		
Whitefly Resistance X Relative Growth				-0.818	0.390	0.297		
Herbicide Resistance X Relative Growth				1.328	0.373	0.078		

STable 5. 2019 Selection analysis showing direct selection on focal traits: whitefly resistance, herbicide

resistance, and relative growth rate. Linear (β) ($R^2 = 0.453$; p < 0.001) and quadratic (γ) ($R^2 = 0.466$; p <

0.001) selection gradients, and total selection with associated standard errors (SE) and *P*-values (*P*). The

(r) column represents correlation coefficients between trait and fitness, estimated as Pearson product-

moment correlations. Significant values are expressed in boldface.

2018 Treatment and Block impact on Relative Growth Rate and Herbicide Damage

Trait	Relative Growth Rate		Herbicide Damage	
Fixed Effects:	*F*	P	*F*	P
Treatment (df = 2)	2.431	0.10	**76.464**	**< 0.001**
Block 2018 (df = 1)	**7.971**	**0.005**	107.667	1.000
Treatment X Block 2018 (df = 2)	**5.239**	**0.007**	100.918	0.340
Random Effects:	χ2	P	χ2	P
Maternal Line (df = 1)	0.000	1.000	0.000	1.000
Maternal Line X Treatment (df = 1)	0.000	1.000	0.000	1.000

507   **STable 6.** 2018 Influence of treatment and block on relative growth, and herbicide damage,

508   analyzed using F-statistics values showing the effects of treatment, block, treatment by block interactions,

509   and likelihood ratio test statistics **(χ2)** showing, maternal line, and maternal line by treatment interactions

510   on variation of plant phenotypes. Significant values are expressed in boldface.

513   2019 Treatment and Block impact on Relative Growth Rate and Herbicide Damage

Trait	Relative Growth Rate		Herbicide Damage	
Fixed Effects:	F	P	F	P
Treatment (df = 2)	0.031	0.969	**46.730**	**< 0.001**
Block (df = 2)	0.323	0.724	0.006	0.994
Treatment X Block (df = 4)	0.259	0.904	**5.982**	**< 0.001**
Random Effects:	$\chi 2$	P	$\chi 2$	P
Population (df = 1)	0.000	1.000	0.190	0.663
Population X Treatment (df = 1)	0.479	0.488	2.300	0.129
Maternal Line (df = 1)	0.000	1.000	0.000	1.000
Maternal Line X Treatment (df = 1)	0.000	1.000	0.000	1.000

STable 7. 2019 Influence of treatment and block on relative growth, and herbicide damage,

analyzed using F-statistics values showing the effects of treatment, block, treatment by block interactions,

and likelihood ratio test statistics **($\chi 2$)** showing population, maternal line, population by treatment

interactions, and maternal line by treatment interactions on variation of plant phenotypes. Significant

values are expressed in boldface.

SFigure 1. Locations of velvetleaf populations sampled and used for this study. In 2018, the field

experiment was conducted with only population one, while in 2019 the sample size was increased to all

eight populations. Populations 3, 4, 6, 7, and 8 were used in the 2021 greenhouse experiment.

SFigure 2. Photosynthetic carbon dioxide response curves by drift environment. A) A-C_i curves measured

on leaves grown without drift exposure B) Comparison of A-C_i curves measured on leaves that developed

before drift exposure (Leaf 1) and after drift exposure (Leaf 2) at 0.5% field dose. C) and at 1% field

dose.

References

Arnold, Stevan J. "Performance surfaces and adaptive landscapes." Integrative and comparative biology 43,

no. 3 (2003): 367-375.

Arnold, Stevan J., Michael E. Pfrender, and Adam G. Jones. "The adaptive landscape as a conceptual bridge

between micro-and macroevolution." Microevolution rate, pattern, process (2001): 9-32.

Belovsky, G. E., and J. B. Slade. "Insect herbivory accelerates nutrient cycling and increases plant

production." Proceedings of the National Academy of Sciences 97, no. 26 (2000): 14412-14417.

Banks, C. J. 1954. A method for estimating populations and counting large numbers of Aphis fabae Scop.

Bull. Entomol. Res. 45:751-756.

Baker, N. T. "Estimated Annual Agricultural Pesticide." 1992–2019 United States Geological Survey. (2021)

https://water.usgs.gov/nawqa/pnsp/usage/maps/show_map.php?year=2019&map=DICAMBA&hilo=L&disp

Bates D, Mächler M, Bolker B, Walker S. "Fitting Linear Mixed-Effects Models Using lme4." Journal

of Statistical Software 67 no. 1 (2015): 1–48.

Browne, Frances B., Xiao Li, Katilyn J. Price, Ryan Langemeier, Alvaro Sanz-Saez de Jauregui, J. Scott

McElroy, Yucheng Feng, and Andrew Price. "Sequential applications of synthetic auxins and glufosinate for

escaped Palmer amaranth control." Agronomy 10, no. 9 (2020): 1425.

Bohnenblust, Eric W., Anthony D. Vaudo, J. Franklin Egan, David A. Mortensen, and John F. Tooker.

"Effects of the herbicide dicamba on nontarget plants and pollinator visitation." Environmental Toxicology

and Chemistry 35, no. 1 (2016): 144-151.

Boutin, Céline, Beate Strandberg, David Carpenter, Solvejg Kopp Mathiassen, and P. J. Thomas. "Herbicide

impact on non-target plant reproduction: what are the toxicological and ecological implications?."

Environmental Pollution 185 (2014): 295-306.

Busi, Roberto, Martin M. Vila-Aiub, Hugh J. Beckie, Todd A. Gaines, Danica E. Goggin, Shiv S. Kaundun,

Myrtille Lacoste et al. "Herbicide-resistant weeds: from research and knowledge to future needs."

Evolutionary Applications 6, no. 8 (2013): 1218-1221.

Byrne, David N., and Thomas S. Bellows Jr. "Whitefly biology." Annual review of entomology 36, no. 1

(1991): 431-457.

Carlsen, S. C. K., N. H. Spliid, and B. Svensmark. "Drift of 10 herbicides after tractor spray application. 1.

Secondary drift (evaporation)." Chemosphere 64, no. 5 (2006a): 787-794.

Carlsen, S. C. K., N. H. Spliid, and B. Svensmark. "Drift of 10 herbicides after tractor spray application. 2.

Primary drift (droplet drift)." Chemosphere 64, no. 5 (2006b): 778-786.

Cebrian, J., and C. M. Duarte. "The dependence of herbivory on growth rate in natural plant communities."

Functional Ecology (1994): 518-525.

Cederlund, Harald. "Effects of spray drift of glyphosate on nontarget terrestrial plants—a critical review."

Environmental toxicology and chemistry 36, no. 11 (2017): 2879-2886.

Cessna, Allan J., Thomas M. Wolf, Gerald R. Stephenson, and Ralph B. Brown. "Pesticide movement to field

margins: routes, impacts and mitigation." Field boundary habitats: implications for weed. Insect and Disease

Management 1 (2005): 69-112.

Ceulemans, Tobias, Eva Hulsmans, Wim Vanden Ende, and Olivier Honnay. "Nutrient enrichment is

associated with altered nectar and pollen chemical composition in Succisa pratensis Moench and increased

larval mortality of its pollinator Bombus terrestris L." Plos one 12, no. 4 (2017): e0175160.

Coley, Phyllis D., John P. Bryant, and F. Stuart Chapin III. "Resource availability and plant antiherbivore

defense." Science 230, no. 4728 (1985): 895-899.

Coley, Phyllis D. "Interspecific variation in plant anti-herbivore properties: the role of habitat quality and rate

of disturbance." New Phytologist 106 (1987): 251-263.

Daily, Gretchen C. "Introduction: what are ecosystem services." Nature's services: Societal dependence on

natural ecosystems 1, no. 1 (1997).

Egan, J. Franklin, and David A. Mortensen. "A comparison of land-sharing and land-sparing strategies for

plant richness conservation in agricultural landscapes." Ecological applications 22, no. 2 (2012): 459-471.

Egan, J. Franklin, Eric Bohnenblust, Sarah Goslee, David Mortensen, and John Tooker. "Herbicide drift can

affect plant and arthropod communities." Agriculture, Ecosystems & Environment 185 (2014): 77-87.

Ehrlich, Paul R., and Peter H. Raven. "Butterflies and plants: a study in coevolution." Evolution (1964): 586-

608.

Evans, John R. "Nitrogen and photosynthesis in the flag leaf of wheat (Triticum aestivum L.)." Plant

physiology 72, no. 2 (1983): 297-302.

Evans, John R. "Photosynthesis and nitrogen relationships in leaves of C3 plants." Oecologia 78, no. 1

(1989): 9-19.

Fine, Paul VA, Zachariah J. Miller, Italo Mesones, Sebastian Irazuzta, Heidi M. Appel, M. Henry H. Stevens,

Ilari Sääksjärvi, Jack C. Schultz, and Phyllis D. Coley. "The growth–defense trade-off and habitat

specialization by plants in Amazonian forests." Ecology 87, no. sp7 (2006): S150-S162.

Fletcher, John S., Thomas G. Pfleeger, Hilman C. Ratsch, and Robert Hayes. "Potential impact of low levels

of chlorsulfuron and other herbicides on growth and yield of nontarget plants." Environmental Toxicology

and Chemistry: An International Journal 15, no. 7 (1996): 1189-1196.

Forkner, Rebecca E., and Mark D. Hunter. "What goes up must come down? Nutrient addition and predation

pressure on oak herbivores." Ecology 81, no. 6 (2000): 1588-1600.

Freemark, Kathryn, and Celine Boutin. "Impacts of agricultural herbicide use on terrestrial wildlife in

temperate landscapes: a review with special reference to North America." Agriculture, Ecosystems &

Environment 52, no. 2-3 (1995): 67-91.

Futuyma, Douglas J., and Anurag A. Agrawal. "Macroevolution and the biological diversity of plants and

herbivores." Proceedings of the National Academy of Sciences 106, no. 43 (2009): 18054-18061.

Gassmann, Aaron J. "Resistance to herbicide and susceptibility to herbivores: environmental variation in the

magnitude of an ecological trade-off." Oecologia 145, no. 4 (2005): 575-585.

Gianessi, Leonard P. "The increasing importance of herbicides in worldwide crop production." Pest

management science 69, no. 10 (2013): 1099-1105.

Gove, Benedict, Sally A. Power, G. Peter Buckley, and Jaboury Ghazoul. "Effects of herbicide spray drift

and fertilizer overspread on selected species of woodland ground flora: comparison between short-term and

long-term impact assessments and field surveys." Journal of Applied Ecology 44, no. 2 (2007): 374-384.

Grossmann, Klaus. "Auxin herbicides: current status of mechanism and mode of action." Pest Management

Science: formerly Pesticide Science 66, no. 2 (2010): 113-120.

Haysom, K. A., and J. C. Coulson. "The Lepidoptera fauna associated with Calluna vulgaris: effects of plant

architecture on abundance and diversity." Ecological entomology 23, no. 4 (1998): 377-385.

Hereford, Joe, Thomas F. Hansen, and David Houle. "Comparing strengths of directional selection: how

strong is strong?." Evolution 58, no. 10 (2004): 2133-2143.

Hintz, Sherwin David. Herbicidal influence on cereal grain aphids in North Dakota. North Dakota State

University, 1971.

Hogenhout, Saskia A., El-Desouky Ammar, Anna E. Whitfield, and Margaret G. Redinbaugh. "Insect vector

interactions with persistently transmitted viruses." Annu. Rev. Phytopathol. 46 (2008): 327-359.

Hunter, Mark D. "Effects of elevated atmospheric carbon dioxide on insect–plant interactions." Agricultural

and Forest Entomology 3, no. 3 (2001): 153-159.

Huot, Bethany, Jian Yao, Beronda L. Montgomery, and Sheng Yang He. "Growth–defense tradeoffs in

plants: a balancing act to optimize fitness." Molecular plant 7, no. 8 (2014): 1267-1287.

Iriart, Veronica, Regina S. Baucom, and Tia-Lynn Ashman. "Herbicides as anthropogenic drivers of eco-evo

feedbacks in plant communities at the agro-ecological interface." Molecular Ecology 30, no. 21 (2021): 5406-

5421.

Jaenike, John. "On optimal oviposition behavior in phytophagous insects." Theoretical population biology 14,

no. 3 (1978): 350-356.

Jones, Gordon T., Jason K. Norsworthy, Tom Barber, Edward Gbur, and Greg R. Kruger. "Off-target

movement of DGA and BAPMA dicamba to sensitive soybean." Weed Technology 33, no. 1 (2019): 51-65.

Legg, James P., Rudolph Shirima, Lensa S. Tajebe, Devid Guastella, Simon Boniface, Simon Jeremiah,

Elibariki Nsami, Patrick Chikoti, and Carmelo Rapisarda. "Biology and management of Bemisia whitefly

vectors of cassava virus pandemics in Africa." Pest management science 70, no. 10 (2014): 1446-1453.

Kempema, Louisa A., Xinping Cui, Frances M. Holzer, and Linda L. Walling. "Arabidopsis transcriptome

changes in response to phloem-feeding silverleaf whitefly nymphs. Similarities and distinctions in responses

to aphids." Plant Physiology 143, no. 2 (2007): 849-865.

Klanderud, Kari, Vigdis Vandvik, and Deborah Goldberg. "The importance of biotic vs. abiotic drivers of

local plant community composition along regional bioclimatic gradients." PloS one 10, no. 6 (2015):

e0130205.

Mangiafico, Salvatore. "rcompanion: Functions to support extension education program evaluation." R

package version 2, no. 10 (2019).

Marrs, Rob H., and Alan J. Frost. "A microcosm approach to the detection of the effects of herbicide spray

drift in plant communities." Journal of Environmental Management 50, no. 4 (1997): 369-388.

Marrs, R. H., A. J. Frost, and R. A. Plant. "Effects of herbicide spray drift on selected species of nature

conservation interest: The effects of plant age and surrounding vegetation structure." Environmental Pollution

69, no. 2-3 (1991): 223-235.

McAdam, Scott AM, Morgane P. Eléouët, Melanie Best, Timothy J. Brodribb, Madeline Carins Murphy,

Sam D. Cook, Marion Dalmais et al. "Linking auxin with photosynthetic rate via leaf venation." Plant

physiology 175, no. 1 (2017): 351-360.

Meyer, Rhonda C., Matthias Steinfath, Jan Lisec, Martina Becher, Hanna Witucka-Wall, Ottó Törjék, Oliver

Fiehn et al. "The metabolic signature related to high plant growth rate in Arabidopsis thaliana." Proceedings

of the National Academy of Sciences 104, no. 11 (2007): 4759-4764.

Moodley, Vaneson, Augustine Gubba, and Paramu L. Mafongoya. "A survey of whitefly-transmitted viruses

on tomato crops in South Africa." Crop Protection 123 (2019): 21-29.

Mound, Laurence A. "Studies on the olfaction and colour sensitivity of Bemisia tabaci (Genn.)(Homoptera,

Aleyrodidae)." Entomología experimentalis et applicata 5, no. 2 (1962): 99-104.

Ning, Wenxi, Xiaobin Shi, Baiming Liu, Huipeng Pan, Wanting Wei, Yang Zeng, Xinpei Sun et al.

"Transmission of Tomato yellow leaf curl virus by Bemisia tabaci as affected by whitefly sex and biotype."

Scientific reports 5, no. 1 (2015): 1-8.

Park, Mi Kyung, Jeong Gyu Kim, Yoo Han Song, Joon-Ho Lee, Key-il Shin, and Kijong Cho. "Effect of

nitrogen levels of two cherry tomato cultivars on development, preference and honeydew production of

Trialeurodes vaporariorum (Hemiptera: Aleyrodidae)." Journal of Asia-Pacific Entomology 12, no. 4 (2009):

227-232.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias

Pietzsch, Stephan Preibisch et al. "Fiji: an open-source platform for biological-image analysis." Nature

methods 9, no. 7 (2012): 676-682.

Schlinkert, Hella, Catrin Westphal, Yann Clough, Zoltán László, Martin Ludwig, and Teja Tscharntke. "Plant

size as determinant of species richness of herbivores, natural enemies and pollinators across 21 Brassicaceae

species." PloS one 10, no. 8 (2015): e0135928.

Schowalter, Timothy D. Insect ecology: an ecosystem approach. Academic press, 2022.

Spencer, Neal R. "Velvetleaf, Abutilon theophrasti (Malvaceae), history and economic impact in the United

States." Economic Botany 38, no. 4 (1984): 407-416.

Thompson, John N. "The evolution of species interactions." Science 284, no. 5423 (1999): 2116-2118.

USDA. "Recent trends in GE adoption." (2016).

Van Lenteren, J. C. van, and L. P. J. J. Noldus. "Whitefly-plant relationships: behavioural and ecological

aspects." Whiteflies: their bionomics, pest status and management 47 (1990): 49.

Vieira, Bruno C., Joe D. Luck, Keenan L. Amundsen, Rodrigo Werle, Todd A. Gaines, and Greg R. Kruger.

"Herbicide drift exposure leads to reduced herbicide sensitivity in Amaranthus spp." Scientific reports 10,

no. 1 (2020): 1-11.

Westra, Eric P., Scott J. Nissen, Thomas J. Getts, Philip Westra, and Todd A. Gaines. "Survey reveals

frequency of multiple resistance to glyphosate and dicamba in kochia (Bassia scoparia)." Weed Technology

33, no. 5 (2019): 664-672.

Yuan, Yujin, Lihua Mei, Mengbo Wu, Wei Wei, Wei Shan, Zehao Gong, Qian Zhang et al. "SlARF10, an

auxin response factor, is involved in chlorophyll and sugar accumulation during tomato fruit development."

Journal of experimental botany 69, no. 22 (2018): 5507-5518.

Züst, Tobias, and Anurag A. Agrawal. "Trade-offs between plant growth and defense against insect

herbivory: an emerging mechanistic synthesis." Annual review of plant biology 68 (2017): 513-534.