Host-symbiont population genomics provide insights into partner fidelity, transmission

mode and habitat adaptation in deep-sea hydrothermal vent snails

Corinna Breusing[1*], Maximilian Genetti[2], Shelbi L. Russell[3], Russell B. Corbett-Detig[2], Roxanne

A. Beinart[1*]

[1]Graduate School of Oceanography, University of Rhode Island, Narragansett, RI 02882, USA

[2]Jack Baskin School of Engineering, University of California Santa Cruz, Santa Cruz, CA 95064,

USA

[3]Department of Molecular, Cell, and Developmental Biology, University of California Santa

Cruz, Santa Cruz, CA 95064, USA

*Corinna Breusing, Roxanne A. Beinart

Email: corinnabreusing@gmail.com, rbeinart@uri.edu

Author Contributions: C.B. and R.A.B. designed the study with advice from S.L.R. and R.C.D.

C.B. performed data analyses and wrote the first draft of the paper. M.G. prepared Illumina

sequencing libraries. All authors contributed to data interpretation and writing of the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences, Evolution

Keywords: Chemosynthetic symbiosis, hydrothermal vents, host-symbiont population genomics,

habitat adaptation, symbiont transmission, symbiont specificity

This PDF file includes:

Main Text

Figures 1 to 6

Tables 1 to 1

Abstract

Abstract

Symbiont specificity, both at the phylotype and strain level, can have profound consequences for

host ecology and evolution. However, except for insights from a few model symbiosis systems,

the degree of partner fidelity and the influence of host versus environmental factors on symbiont

composition are still poorly understood. Nutritional symbioses between invertebrate animals and

chemosynthetic bacteria at deep-sea hydrothermal vents are examples of relatively selective

associations, where hosts affiliate only with particular, environmentally acquired phylotypes of

gammaproteobacterial or campylobacterial symbionts. In hydrothermal vent snails of the sister

genera *Alviniconcha* and *Ifremeria* this phylotype specificity has been shown to play a role in

habitat distribution and partitioning among different holobiont species. However, it is currently

unknown if fidelity goes beyond species level associations that might influence genetic

structuring, connectivity and habitat adaptation of holobiont populations. We used metagenomic

analyses to assess sequence variation in hosts and symbionts and identify correlations with

geographic and environmental factors. Our analyses indicate that host populations are not

differentiated across a ~800 km gradient, while symbiont populations are clearly structured

between vent locations due to a combination of neutral and selective processes. Overall, these

results suggest that host individuals flexibly associate with locally adapted strains of their

specific symbiont phylotypes, which supports a long-standing but untested paradigm of the

benefits of horizontal transmission. Strain flexibility in these snails likely enables host

populations to exploit a range of habitat conditions, which might favor wide-spread genetic

connectivity and ecological resilience unless physical dispersal barriers are present.

Significance Statement

Symbiont composition in horizontally transmitted symbioses is influenced by a combination of

host genetics, environmental conditions and geographic barriers. Yet the relative importance of

these factors and the effects of adaptive versus neutral evolutionary forces on symbiont

population structure remain unknown in the majority of marine symbioses. To address these

questions, we applied population genomic approaches in four species of deep-sea hydrothermal

vent snails that live in obligate association with chemosynthetic bacteria. Our analyses show that

host genetics plays a minor role compared to environment for symbiont strain composition

despite specificity to symbiont species and corroborate a long-standing hypothesis that vent

invertebrates affiliate with locally adapted symbiont strains to cope with the variable habitat

conditions characterizing hydrothermal vents.

Introduction

Mutualistic relationships between eukaryotes and bacterial microbes are ubiquitous in nature.

Symbionts enable hosts to gain access to novel resources and habitats, provide protection against

pathogens and predators, and can be essential for the host's diet (1, 2). For symbiotic

associations to persist over evolutionary time, hosts must successfully transmit their symbionts

from one generation to the other, either through symbiont acquisition from the environment

(horizontal transmission), direct inheritance of symbiont lineages through the host germline

(vertical transmission) or a combination of both mechanisms (mixed transmission) (3, 4). The

mode of transmission has significant implications for the composition and variation of symbionts

within and between host individuals. Vertical transmission typically results in strong genetic

coupling between host and symbiont lineages and an accompanied reduction in intra-host

symbiont diversity (3). By contrast, horizontal transmission exposes aposymbiotic hosts to a

potentially heterogenous environmental pool of symbiont lineages (3), which can promote the

formation of generalist partnerships, where multiple hosts and symbionts associate with each

other, to more specialized associations between only one or a few potential partners (1, 5). This

range is often referred to as host-symbiont specificity, which can vary in its taxonomic level for

both partners depending on the symbiotic system. The degree of partner fidelity and its effect on

symbiont composition can have dramatic impacts on holobiont functioning (6). For example,

host-symbiont specificity, both at the species and genotype level, is crucial for light production

in bioluminescent squid (7), while shifts in microbiome assemblages have been shown to affect

phytoplankton growth rates (8) and the efficiency of nitrogen fixation in nodule-forming legumes

(9–11).

Environmental transmission of obligate symbionts is particularly common in marine

ecosystems (2). However, despite its importance for host biology, the relative contributions of

host genetic, environmental and geographical factors to symbiont composition remain

understudied in most horizontally transmitted marine symbioses. It has long been hypothesized

that horizontal transmission in marine symbioses enables host organisms to associate with locally

adapted symbiont strains, conferring fitness advantages in spatially and temporally variable

marine habitats (3, 12–14). Especially for long-dispersing aposymbiotic larvae that are likely to
encounter new habitat conditions when they settle, association with a locally adapted symbiont
strain may be advantageous compared to carrying a vertically transmitted symbiont that might be
maladapted at a non-native site. This hypothesis has been indirectly supported by evidence that
marine animals with horizontally transmitted obligate microbial symbionts often host location-
specific strains (14–19). However, the influence of local adaptation relative to neutral
evolutionary processes on symbiont geographic structure and genomic traits has not been
formally evaluated.

Chemosynthetic animal-microbe symbioses are globally significant phenomena that
dominate hydrothermal vent and hydrocarbon seep ecosystems in the deep sea. In these
associations, the bacterial partner uses chemical energy from the oxidation of reduced fluid
compounds, such as hydrogen, sulfide or methane, to synthesize organic matter, which serves as
primary nutrition for the host (20). Vent animals harboring chemosynthetic symbionts are
typically highly selective in their partner choice: In the predominant number of cases host
individuals associate with only 1–2 phylotypes (species or genera) of gammaproteobacterial or
campylobacterial symbionts (20), whereas symbionts can exhibit a comparatively broad host
range. Chemosynthetic endosymbioses in deep-sea snails of the Indo-Pacific sister genera
Alviniconcha and *Ifremeria* are examples of reciprocally relatively specific partnerships, where
host species harbor only particular symbiont species or genera across their geographic
distribution. Given the absence of host-symbiont phylogenetic concordance, the symbionts are
assumed to be environmentally acquired (21), although a pseudo-vertical transmission
component is possible in *Ifremeria* given its brooding reproductive mode (22). In the Eastern
Lau Spreading Center (ELSC), previous work suggested that specificity to functionally distinct
symbiont phylotypes drives local and regional-scale habitat partitioning among four co-occurring
Alviniconcha and *Ifremeria* species (23–26). *Alviniconcha boucheti* from the ELSC contains a
campylobacterial phylotype (Epsilon) and is usually found at northern vent sites with high
concentrations of sulfide and hydrogen, while *A. kojimai* and *A. strummeri* associate with
different gammaproteobacterial phylotypes (Gamma1 and Gamma1/GammaLau, respectively)
and usually occupy mid-latitude to southern vent sites where the concentrations of these
chemical reductants are lower (23, 25–27). *Ifremeria nautilei* establishes dual symbioses with
thiotrophic and methanotrophic gammaproteobacterial endosymbionts (27) and is co-distributed

with *Alviniconcha* across their geographical range, although it typically segregates into habitat

patches with reduced fluid flow relative to its sister genus (28). While it has been well-

established that niche differentiation across hydrothermal vents is likely mediated by symbiont

phylotype specificity in *Alviniconcha* and *Ifremeria* host species, nothing is known about the

fidelity of these associations at the population level, how strain-level specificity might influence

host population structure and connectivity, and how regional adaptation is conferred

functionally.

In this study we applied population genomic methods to assess symbiont strain-level

genetic variation and patterns of host-symbiont genetic subdivision in *Alviniconcha* and

Ifremeria species from the ELSC and Tonga Volcanic Arc (Fig. 1; Table 1). Using multivariate

statistical analyses, we evaluated the impact of host traits, environment and geography on

symbiont composition in populations of both genera and assessed the effect of local adaptation

on symbiont geographic structure.

Results

Host transcriptome assemblies and population genomic structure

Host transcriptome assemblies consisted of 24,176–35,654 transcripts (totaling 20.68–28.68 Mb)

and were approximately 30.30–55.40% complete (Table S1). Mapping of host reads against the

transcriptome references and subsequent filtering of variant sites yielded 1,655–9,185 single

nucleotide polymorphisms (SNPs) per species for population genetic analyses. Irrespective of

host taxon, F_{ST} and ordination analyses revealed low genetic differentiation among host

populations sampled from different vent localities (Fig. 2; Table S2). With the exception of four

SNP sites in *A. kojimai*, no F_{ST} outliers could be detected in any host species (Table S3).

However, all species contained a number of SNPs that were moderately to highly divergent

among host populations. When analyses were constrained to these SNP subsets (*Alviniconcha*:

$F_{ST} > 0.15$; *Ifremeria*: $F_{ST} > 0.10$), population genetic structuring by vent site was observed in all

Alviniconcha species, but not *Ifremeria* (Fig. S1). For both *A. kojimai* and *A. strummeri*, the

respective SNP subsets further indicated genetic differences between Tui Malila populations that

were sampled in different years. The effect of geography on host population genetic structuring

was not significant after correction for symbiont genetic distance in partial Mantel correlations

(Table S4).

Symbiont genome assemblies and population genomic structure

Symbiont genome assemblies varied in size from 2.10–4.86 Mb and contained 2,006–6,660

predicted protein-coding genes, with GC content ranging from 34.30–59.00% (Table S2). All

assemblies were characterized by a high level of completeness (92.79–99.25%) and low amount

of contamination (0.74–5.48%) (Table S5). Symbiont populations for the Epsilon, Gamma1, Ifr-

SOX and Ifr-MOX phylotypes were largely structured by vent field or broader geographic region

based on 239–7,057 variant sites (Fig. 3, S2; Table S6). These associations were significant even

when corrected for host genetic variation based on highly differentiated SNP markers (Table S4;

$p \leqq 0.0244$, $r = 0.2665$–0.8705). Genetic differentiation between symbiont populations typically

increased with geographic distance between vent fields (Table S6). Although F_{ST} values for

GammaLau populations of *A. strummeri* were moderate when calculated between distinct vent

sites (Table S6), ordination analyses and Mantel tests provided no evidence for genetic

differentiation of this symbiont phylotype across geographic locations based on 192 marker loci

(Fig. 3, S2). The Gamma1 phylotype is associated with both *A. kojimai* and *A. strummeri* and we

therefore investigated whether populations of this phylotype varied between host species.

Ordination analyses indicated a clear clustering by host taxon that superseded the effect of

geography (Fig. 4, S3), suggesting that different Gamma1 strains associate selectively with either

A. kojimai or *A. strummeri*. By contrast, the effect of host genetics on symbiont genetic variation

within species appeared to be weak. Although partial Mantel tests suggested significant

associations of host genotype with strain composition or dominant symbiont type for the *A.*

boucheti – Epsilon, *A. kojimai* – Gamma1, *A. strummeri* – Gamma1, and *I. nautilei* – Ifr-SOX

pairs, *r* statistics were relatively low especially compared to the effect of geography, indicating

limited biological relevance (Table S4; $p \leqq 0.0364$, $r = 0.0510$–0.2270).

Gene content variation between symbiont strains

We assessed variation in gene content between symbiont strains from different vent localities or

broader geographic regions (Fig. 5, S4; Table S7). For the Gamma1 symbiont, we further

determined gene content variation between strains from different host species given that this

symbiont phylotype occurs in both *A. strummeri* and *A. kojimai* (Fig. 4, S3; Table S8).

Differentially preserved or abundant genes between symbiont strains comprised about 1–5% of

protein-coding regions. Independent of symbiont phylotype, the most common differences

among geographic or host-specific strains concerned genes of unknown function as well as a

smaller number of genes related to mobilome and anti-viral defense (Fig. 4–5, S3–4; Table S7,

S8). Geographic strains of the *A. boucheti* Epsilon symbiont further differed in the presence of an

ABC transporter, a GDP-L-fucose synthetase, a NAD(FAD)-utilizing hydrogenase and the DNA

repair protein RecN, which were conserved in strains from Kilo Moana and Tow Cam but were

absent or very lowly abundant in strains from ABE (Table S7). Within the *A. kojimai* Gamma1

phylotype, strains from ABE (but rarely Tui Malila) contained a few genes involved in DNA,

protein and cell wall metabolism as well as maturation and regulation of uptake (NiFe)

hydrogenases (*hyaC, hyaD, hoxJ*). Differences in hydrogenase-related genes were also observed

in the Gamma1 symbiont of *A. strummeri*, where an operon for a hydrogen-sensing hydrogenase

(*hupUV/hoxBC*) and genes for hydrogenase maturation and assembly proteins (*hypF, hyaF,*

hoxV/hupK, hypD, hypE, hoxX) were largely missing in strains from Tui Malila (but not Tahi

Moana) (Table S7). Between host species, Gamma1 strains notably differed in the presence of

genes for a sulfite dehydrogenase complex (*soeABC*), which was abundant in strains specific to

A. kojimai but not *A. strummeri* (Table S8). Within the *A. strummeri* GammaLau phylotype,

strains from Tui Malila contained a broad range of metabolic genes that were absent or

infrequent in strains from Tahi Moana, including genes related to macronutrient metabolism,

stress response, membrane transport and several other functional processes (Table S7). The Ifr-

SOX strains of *I. nautilei* differed in several genes related to DNA, nitrogen, lipid and amino

acid metabolism, membrane transport, cell regulation and detoxification, which were present in

strains from the ELSC but mostly missing in strains from Niua South. Within the Ifr-MOX

phylotype, strains from Tahi Moana and Tui Malila differed in genes involved in sulfide

respiration and oxidative phosphorylation, macronutrient metabolism, cation transport, motility,

cofactor, cell wall, DNA and nucleotide metabolism as well as cell regulation and stress response

(Table S7).

Adaptive variation within symbiont phylotypes

A subset of genetic variants exhibited significant associations with environmental factors in all

symbiont phylotypes except for GammaLau, with constrained ordinations explaining between

5.39% and 19.76% of the variance in the respective RDA models (Fig. 6; Table S9). Candidate

adaptive loci for each symbiont phylotype encompassed a variety of metabolic categories,

although in many cases no particular function could be assigned (Table S9). In the *A. boucheti*

Epsilon symbiont, 129 variants were significantly associated with fluid composition, while

another 47 were correlated with depth. These variants were mostly located in genes related to

protein, amino acid, and cell wall metabolism, followed by cofactor, DNA, carbon and

nucleotide metabolism, as well as membrane transport (Table S9). Several other variants were

linked to nitrogen, iron and RNA metabolism, virulence, motility, cell signaling, stress response,

respiration, cell cycle, as well as hydrogen and sulfur metabolism. In the *A. kojimai* Gamma1

symbiont, 91 variants were correlated with fluid composition and 9 were linked to year. 29 of

these variants were classified into functional categories. The most represented categories were

mobilome, cell wall and DNA metabolism, anti-viral defense, membrane transport, cofactor

metabolism and respiration. A few variants were associated with amino acid, hydrogen, protein

and RNA metabolism, and cell regulation. In the *A. strummeri* Gamma1 symbiont, 95 variants

showed associations with fluid composition, while 3 were correlated with year. 54 of these

variants could be functionally annotated and assigned to the following metabolic categories:

protein, amino acid and cell wall metabolism, membrane transport, virulence, carbon, nitrogen,

DNA/RNA and nucleoside metabolism, cofactor, fatty acid and sulfur metabolism, stress

response, cell signaling and mobilome (Table S9). The *I. nautilei* Ifr-SOX and Ifr-MOX

symbionts contained 94 and 11 putatively adaptive variants, respectively. In both symbionts,

these variants were mostly linked to depth, followed by fluid composition and year. While the

majority of variants were located in genes with unknown functions, a number of loci was linked

to mobile elements, cell cycle, anti-viral defense, DNA/RNA metabolism, and membrane

transport. In each symbiont phylotype, virtually all candidate variants were characterized by

elevated F_{ST} and gene-wide *p*N/*p*S values, further supporting their potential role in habitat

adaptation (Table S9). Several other genes exhibited increased ratios of non-synonymous to

synonymous substitution, although sites within these genes did not show significant associations

with environment (Table S10). It is possible that sites within these genes covary with other

ecological factors that could not be tested in this study or that these patterns are caused by recent

slightly deleterious mutations that were not yet purged by natural selection.

Discussion

Strain-level variation within microbial symbionts is increasingly recognized as an important driver of host ecology and evolution (6), yet its patterns, determining factors and functional implications remain poorly investigated in many non-model symbioses. In this study, we used metagenomic analyses to assess sequence and gene content differences between chemosynthetic symbionts associated with four co-occurring species of deep-sea hydrothermal vent snails and determined the impact of host genetic, geographical and environmental factors on symbiont strain composition and variation.

Despite fidelity between hosts and symbionts at the species level (23, 25), our results indicate that specificity between host genotypes and symbiont strains in *Alviniconcha* and *Ifremeria* is weak: Host populations were not partitioned across a ~800 km gradient, whereas symbiont populations were clearly structured between vent locations or broader geographic regions. Even when we correlated symbiont and host genetic distances based on highly differentiated markers, test statistics for significant associations remained low, suggesting that host genetics has a minor impact compared to environment or geography on strain composition within both *Alviniconcha* and *Ifremeria*.

These findings qualitatively agree with observations in deep-sea mussel hybrids that appear to associate with locally available symbiont strains (19), but contrast markedly with patterns in other horizontally transmitted partnerships, such as the squid-*Vibrio* symbiosis and some legume-rhizobia associations, where hosts exhibit strong strain specificity (6). Environmental uptake of locally adapted symbiont strains provides host organisms with the opportunity to optimally exploit novel habitats, but carries the risk of unsuccessful symbiont acquisition and infection by cheaters (3). Depending on the amount of partner reliance, chances of symbiont encounter and fitness variation among symbiont strains, holobionts likely find different tradeoffs between these opposing factors. Associations with chemosynthetic bacteria are obligate for hydrothermal vent animals, but are restricted to relatively ephemeral habitats that are characterized by large temporal and spatial fluctuations in environmental conditions and associated shifts in microbial communities (29–31). Strong nutritional dependency in vent symbioses combined with the uncertainty of habitat (and thus symbiont) encounter might promote specificity towards a mutualistic symbiont phylotype, while enabling enough flexibility towards different local strains of that phylotype to maximize recruitment success of host larvae at a new vent site. By contrast, bobtail squids inoculate their environment with symbiotic bacteria

and thereby ensure symbiont availability for their offspring (32), which might favor increased

strain selectivity in this association. Similarly, some legume species appear to preferentially

associate with certain rhizobial strains that are abundant in the native host range (33). Although

the factors underlying strain specificity are not fully understood, it is possible that the relative

fitness advantages provided by these locally available strains (33) coupled with the benefit of

decreased cheater invasion (11) might have selected for strong partner fidelity in these

symbioses.

Alviniconcha kojimai and *A. strummeri* represent notable exceptions to the observed

patterns, as they share identical phylotypes of one gammaproteobacterial symbiont (Gamma1),

but appear to take up different strains even in habitats where these host species co-occur. The

phylogenetic divergence between *A. kojimai* and *A. strummeri* (~25 MYA) is more recent than

that between either species and *A. boucheti* (~38 MYA) or *I. nautilei* (~113 MYA) (25). Perhaps

the closer evolutionary relationship between these two species favors associations with the same

symbiont phylotype, as has been suggested for *Bathymodiolus azoricus* and *B. puteoserpentis* on

the Mid-Atlantic Ridge as well as *B. thermophilus* and *B. antarcticus* on the East Pacific Rise

(19). However, compared to the bathymodiolin mussel system, the split between *A. kojimai* and

A. strummeri is significantly older (>17 MYR), which could indicate that the timing of phylotype

specificity evolution varies among taxonomic groups, possibly as a result of contrasting

ecological or evolutionary contexts. The fact that *A. kojimai* and *A. strummeri* nevertheless

associate with distinct symbiont strains in sympatry is potentially a mechanism to avoid

competition for niche space. Gamma1 strains of these two species notably differed in the

presence of a molybdenum-containing sulfite dehydrogenase (SoeABC), which was preserved in

strains of *A. kojimai* but not *A. strummeri*. SoeABC is the predominant enzyme for sulfite

oxidation in many purple sulfur bacteria (34) and is further involved in taurine and

dimethylsulfoniopropionate degradation in *Roseobacter* clade bacteria (35–37). Although the

physiological role of SoeABC in *A. kojimai*'s Gamma1 strain is unknown, it is possible that it

contributes to partitioning of sulfur resources among co-occurring snail holobionts.

Our data support a model of horizontal symbiont transmission in both *Alviniconcha* and

Ifremeria. While these results are expected for *Alviniconcha* which produce free-swimming

planktotrophic larvae and do not invest in their young (38), they are rather surprising for

Ifremeria which brood their offspring in a modified pouch in the female's foot (22) and would

thus have the opportunity to pseudo-vertically transmit their symbionts. It is possible that a

pseudo-vertical transmission component exists in *Ifremeria*, but that maternally acquired

symbionts get replaced or complemented by more competitive strains in the habitat where the

snail larvae settle. Our current dataset cannot distinguish this possibility from a strict horizontal

transmission mode, though future studies assessing symbiont composition in different

developmental stages of *Ifremeria* would be helpful to address this hypothesis.

While the majority of genetic variants in the symbionts did not deviate from neutral

expectations, a subset of loci showed evidence for natural selection, implying a role of both

genetic drift and local adaptation in shaping symbiont population structure. Candidate adaptive

loci spanned a surprisingly broad range of metabolic functions, many of which were probably

not causally linked to the investigated environmental predictors but other correlated variables

that we could not account for in this study. For example, in almost all symbiont phylotypes we

observed adaptive variation in genes that were related to anti-viral defense (e.g., CRISPR-

associated proteins, restriction-modification systems) or mobile elements. Although differences

in depth or geochemistry might contribute to these patterns, it is more likely that they reflect

exposure of symbiont strains to distinct viral assemblages that might covary with local habitat

conditions, as has been suggested for vestimentiferan tubeworm symbionts (39). Polymorphisms

in other genes, by contrast, are likely directly explained by variation in the analyzed

environmental factors. Geographic strains of *A. boucheti*'s Epsilon symbiont, for instance,

contained several variants under positive selection that were located in genes involved in

hydrogenase assembly, iron transport, sulfur oxidation and respiration. Likewise, the Gamma1

strains of *A. kojimai* and *A. strummeri* showed adaptive differences in variants related to

hydrogen and sulfur metabolism, respectively. Niche-specific differences in hydrogen metabolic

genes were also observed in comparisons of gene content among symbiont strains. In both *A.*

kojimai and *A. strummeri*, Gamma1 strains from Tui Malila lacked some subunits of uptake or

hydrogen-sensing hydrogenases as well as various genes for hydrogenase maturation, synthesis

and regulation. Given that H_2 concentrations at Tui Malila can drop to 35 µM in endmember

fluids and are probably lower in diffuse flow habitats (23), it is possible that hydrogen does not

constitute a major energy source for Gamma1 strains from this locality. This hypothesis is in

agreement with previous physiological experiments that revealed strikingly low hydrogen

oxidation and associated carbon fixation rates in Gamma1 symbionts from Tui Malila (26). An

alternative though mutually non-exclusive explanation for the loss of hydrogenase-related genes

at least in the *A. strummeri* strain from Tui Malila could be avoidance of intra-host competition

with co-occurring GammaLau strains, which often co-dominate in *A. strummeri* individuals at

this vent site (23). Such functional diversity is predicted to enable symbiont coexistence in a

variety of hydrothermal vent symbioses, including bathymodiolin mussels (40, 41), alvinocaridid

shrimp (42) and vestimentiferan tubeworms (43).

Compared to their symbionts, host populations were markedly less structured across the

same spatial scales. Although a proportion of genetic markers was differentiated across vent

sites, we did not find strong evidence for local adaptation in any host species, suggesting that

these patterns likely reflect random variation among localities. The limited genetic subdivision

between host populations agrees with predictions from biophysical models that indicate absence

of physical dispersal barriers for vent larvae within the Lau Back-Arc Basin (44). While

symbiont population structure, by contrast, appeared to be at least partly driven by natural

selection, it is possible that symbionts also experience stronger dispersal limitations than their

hosts, as has been hypothesized in some coral-algae symbioses (18). The environmental

distributions and life cycles of the free-living stages of *Alviniconcha* and *Ifremeria* symbionts are

currently unknown and a better understanding of these aspects will be necessary to evaluate the

relative importance of dispersal barriers on symbiont biogeography.

Overall, our findings reveal a lack of strain-level specificity in *Alviniconcha* and

Ifremeria symbioses, which possibly reflects an evolutionary strategy to cope with the transient

and dynamic nature of hydrothermal vent habitats. Strain flexibility in these associations likely

contributes to the wide-spread genetic connectivity observed among host populations, which in

turn could favor ecological resilience to natural but also anthropogenic environmental

disturbances, a relevant consideration given the increasing human pressures on hydrothermal

ecosystems worldwide (45). Our observations further support the fundamental hypothesis that

horizontal transmission in marine symbioses enables host organisms to associate with locally

adapted symbiont strains (3, 12–14). Though the genomic basis of local adaptation can be

detected in natural populations using population genomics methods, as we did here, evaluation of

the phenotypic consequences of the observed strain-level genomic trait variation will be

necessary to confirm local adaptation in these symbiont strains. Future work using organism-

based manipulative experiments will be helpful to compare the fitness of hydrothermal vent

animals hosting site-specific symbiont strains when exposed to native and foreign conditions

(46). These assessments will be critical to understand the commonality of horizontal symbiont

transmission in the marine environment, given the hypothesis that local adaptation is less

common in marine than terrestrial systems due to higher levels of gene flow (47).

Materials and Methods

Sample collection, nucleic acid extraction and sequencing

Samples of *Alviniconcha* and *Ifremeria* were collected from six vent sites (1164–2722 m) of the

Lau Basin and Tonga Volcanic Arc in 2009 and 2016 using remotely operated vehicles (Fig. 1;

Table 1). Upon recovery, animal samples were dissected, placed in RNALater™ Stabilization

Solution (Thermo Fisher Scientific, Inc.) and frozen at −80°C until further analysis. DNA was

extracted with the Quick-DNA 96 Plus extraction kit (Zymo Research, Inc.) and further purified

with the MO BIO PowerClean DNA Pro Clean-Up kit (Qiagen, Inc.). High molecular weight

(HMW) DNA for long-read sequencing was isolated with Qiagen Genomic-tips following

manufacturer's instructions.

Host transcriptome sequencing and assembly

Illumina RNAseq reads for host transcriptome assemblies were obtained from sequencing

experiments performed in (48) and (26). Adapter clipping, quality trimming, and removal of

rRNA and symbiont reads was performed as in (26). Cleaned host reads were error corrected

with RCORRECTOR (49) and filtered for uncorrectable and overrepresented sequences with the

TRANSCRIPTOMEASSEMBLYTOOLS package

(https://github.com/harvardinformatics/TranscriptomeAssemblyTools). Host transcriptome co-

assemblies for each *Alviniconcha* and *Ifremeria* species were performed with TRINITY (50) using

the PASAFLY algorithm. For each *Alviniconcha* species, additional transcripts were reconstructed

from 454 reads obtained from (51). Assembled contigs for each species were clustered with CD-

HIT-EST (52) at a 95% identity threshold to reduce transcript redundancies. Open reading frames

(ORFs) were predicted with TRANSDECODER (https://github.com/TransDecoder/TransDecoder)

considering homologies to known proteins (UniRef90) and protein domains (Pfam) as ORF

retention criteria. Transcripts that did not contain any ORF or had a non-eukaryotic origin based

on taxonomy classifications with BLOBTOOLS (53) were removed from the assembly. Final

transcriptome assemblies were evaluated for quality and completeness with BUSCO (54).

Symbiont genome sequencing and re-assembly

Illumina reads for the *A. boucheti, A. kojimai* and *Ifremeria* holobionts were obtained from

previous sequencing runs performed in (27). Raw reads were trimmed with TRIMMOMATIC (55)

and filtered for sequence contaminants through mapping against the human (GRCh38) and PhiX

reference genomes. Decontaminated reads were grouped into symbiont and host reads by

assessing best matches against draft symbiont genomes (27) with BBSPLIT

(https://sourceforge.net/projects/bbmap/). To improve contiguity of the genome assemblies we

conducted 3–4 Nanopore sequencing runs for each symbiont phylotype on a MinION device

(Oxford Nanopore Technologies) using the SQK-LSK109 ligation kit after HMW DNA

enrichment with the Circulomics Short Read Eliminator kit. Basecalling of the Nanopore reads

was done with ALBACORE (Oxford Nanopore Technologies) and adapters were clipped with

PORECHOP (https://github.com/rrwick/Porechop). Hybrid assemblies of Illumina and Nanopore

reads were constructed for each symbiont genome with SPADES (Gamma1) (56) or

METASPADES (all others) (57) choosing k-mers between 21 and 91 in 10 step increments. Raw

assemblies for the Epsilon, Gamma1 and Ifr-SOX symbionts were binned with GBTOOLS (58)

and incrementally gap filled, corrected and scaffolded with LR_GAPCLOSER (59), ra2.py (60),

and SLR (61), respectively, following recommendations by (62) The Ifr-MOX symbiont

assembly was automatically binned with METABAT2 (63) for contigs $\geqq$ 1500 bp. Shorter contigs

($\geqq$ 500 bp) were binned with MAXBIN (64). Scaffolds < 200 bp were excluded from all

assemblies. Final assemblies were polished with PILON (65), functionally annotated with RAST-

TK (66) and quality-checked with CHECKM (67) and QUAST (68).

Population-level metagenomic sequencing and variant identification in hosts and symbionts

192 barcoded high-throughput DNA sequencing libraries were prepared with a Tn5 transposase-

based protocol after (69) at the University of California Santa Cruz and then sent for 150 bp

paired-end sequencing on a NovaSeq 6000 instrument at the University of California Davis.

However, due to low read allocation 66 of the libraries were excluded from further analysis. Raw

sequence reads were trimmed, filtered and sorted by host species and symbiont phylotype as

described above. Optical duplicates were removed with PICARD's MARKDUPLICATES tool

(https://github.com/broadinstitute/picard). To resolve common alignment errors and improve

base call accuracy, we locally realigned reads around indels and recalibrated base quality scores

with LOFREQ (70).

Host population genomic variation was assessed in ANGSD (71) by inferring genotype

likelihoods based on Hardy-Weinberg equilibrium considering individual inbreeding

coefficients. To increase accuracy of the analyses, variant sites with mapping qualities < 30

(minMapQ = 30), base qualities < 20 (minQ = 20), and minimum minor allele frequencies < 0.01

(minMaf = 0.01) were excluded. We further filtered sites based on strand bias (sb_pval = 0.05),

heterozygote bias (hetbias_pval = 0.05) and probability of being variable (SNP_pval = 1e-6). In

addition, we removed spurious and improperly paired reads, adjusted mapping qualities for

excessive mismatches (C = 50) and computed per-base alignment qualities (BAQ = 1) to

disregard variants close to indel regions. Putative paralogous variants were excluded by

discarding reads with multiple mappings and by considering only sites that had a maximum

depth of 40–80. Genetic distances between individuals were inferred by calculating pairwise

covariance matrices.

Symbiont population genomic variation was determined with FREEBAYES (72) using

input parameters adjusted for the analysis of metagenomic data (-F 0.01 -C 1 -p 1 --pooled-

continuous --haplotype-length 0 --report-monomorphic). Variant calls were restricted to sites

with a minimum base quality of 20, a minimum mapping quality of 30, and a minimum coverage

of 10. To eliminate bias in variant identification and other downstream analyses due to uneven

read depth between samples, we normalized all samples to the lowest amount of coverage found

in a sample for a particular symbiont phylotype (> 10X coverage). Variants were further filtered

based on strand bias (SRP > 5 && SAP > 5 && EPP > 5), proximity to indels (5 bp) and

maximum mean depth with BCFTOOLS (73) and VCFTOOLS (74). In addition, sites and

individuals with more than 25% missing data were excluded from the analysis. Allele counts (=

symbiont strain abundances) and consensus haplotypes (= dominant symbiont strains) were

extracted with GATK's VARIANTSTOTABLE tool (75).

Genomic structure and differentiation

We performed ordination analyses with the APE and STATS packages in R (76, 77) to assess

genetic variation between symbiont and host populations. Host genetic structure was determined

through principal component analyses based on genetic covariance matrices, while symbiont

genetic structure was inferred through principal coordinate analyses based on both consensus

haplotype and allele count data transformed into Euclidean distances and Bray-Curtis

dissimilarities, respectively. For each sample, absolute allele counts were normalized to relative

counts prior to analysis. Negative eigenvalues were corrected using the method by Cailliez (78)

and final ordination plots were produced with GGPLOT2 (79). F_{ST} values between host and

symbiont populations were calculated in ANGSD and SCIKIT-ALLEL

(https://github.com/cggh/scikit-allel), respectively, following the procedure in (80). For the host

species, potential outlier loci under selective pressures were inferred with OUTFLANK (81) based

on neutral F_{ST} distributions that were obtained from quasi-independent SNP subsets determined

with PLINK (82).

Assessment of gene content variation

We used PANPHLAN (83) to investigate potential differences in gene composition between

symbiont strains from different vent localities. Downsampled symbiont reads for each species

were mapped against the corresponding symbiont reference genome using custom RAST-TK

annotations for functional categorizations. Samples were profiled for gene presence/absence

based on the following parameter thresholds: --min_coverage 1 --left_max 1.70 --right_min 0.30.

Gene content variation between symbiont strains was assessed by identifying genes that were

present/absent in 90% of samples from one or the other geographic region. To account for gene

content variation between different strains within hosts we further quantified gene abundances

with SALMON (84) using the following parameters: -l IU --meta --rangeFactorizationBins 4 --

numBootstraps 1000 --seqBias --gcBias -s -u. Abundance values were normalized with the

Trimmed Mean of M method using TRINITY's *abundance_estimates_to_matrix.pl* script (50, 85).

Gene presence/absence and differential abundance heatmaps were produced with the

COMPLEXHEATMAP package in R (86).

Assessment of symbiont variation based on environment and host genetics

We conducted partial Mantel tests with the VEGAN package in R (87) to assess relationships

between host and symbiont genetic distances and geography based on Spearman rank

correlations. As no population genetic structure could be observed in any host species based on

the full SNP datasets, we estimated host covariance matrices using only moderately to highly

differentiated SNP subsets for this analysis (*Alviniconcha*: $F_{ST} > 0.15$; *Ifremeria*: $F_{ST} > 0.10$).

Geographic distances were determined by calculating the geodesics between vent sites with the

GEOSPHERE package (88). We used redundancy analyses (RDA) following the approach in (89)

to evaluate the influence of hydrothermal fluid composition and depth on the genetic structure of

each symbiont phylotype. We further included the effect of sampling year, which encompasses

changes in hydrothermal circulation within the ELSC as indicated by the cessation in fluid flow

at the Kilo Moana vent field between 2009 and 2016. Endmember concentrations for

geochemical compounds were obtained from the literature (23, 90, 91) or unpublished data

provided by A. Diehl and J. Seewald (Table 1). Due to multi-collinearity among the chemical

species, we used the first eigenvector from principal coordinate analyses (corresponding to the

eigenvalue with the largest explanatory power) as composite value. For each symbiont

phylotype, we further assessed the strength of correlation with other predictors to exclude

variables that were highly collinear. Unless geographic location was strongly linked to

environmental predictors, we used latitude as a conditioning factor in the analyses to correct for

isolation by distance. Based on these collinearity evaluations, we tested the effect of all factors

on the two *Ifremeria* symbionts, the effect of depth and fluid composition on the *A. boucheti*

symbiont and the effect of fluid composition and year (uncorrected for geography) on the *A.*

kojimai and *A. strummeri* symbionts. SNPs were considered candidates for local adaptation if

their loadings on significant constrained RDA axes deviated more than 2.5 standard deviations

from the mean of the distribution. For each symbiont phylotype we further calculated the ratio of

non-synonymous to synonymous polymorphisms (pN/pS) with SNPEFF (92) to infer candidate

genes evolving under natural selection.

Data Accessibility

Raw Nanopore reads, host transcriptomes and symbiont genomes have been deposited in the

National Center for Biotechnology Information under BioProjects PRJNA523619,

PRJNA526236 and PRJNA741492. Annotations for new symbiont genomes are available on the

Acknowledgements

We thank the captains, crews and ROV pilots of the R/V *Thomas G. Thompson* (ROV *Jason II*) and R/V *Falkor* (ROV *Ropos*) for supporting the sample collections that made this study possible. We thank Peter Girguis for his contribution of the 2009 samples to this project, Alexander Diehl and Jeff Seewald for providing geochemical data, Michelle Hauer and Erin Frates for their assistance with sample preparation and the National Science Foundation EPSCoR Cooperative Agreement OIA-#1655221 for providing access to Brown University's high-performance computing cluster where the bioinformatic analyses were performed. We further thank the technical staff at the UC Davis Genome Center for sequencing our Illumina metagenomic libraries. This work was supported by the National Science Foundation (grant numbers OCE-1536331, 1819530 and 1736932 to R.A.B.) and the National Institute of Health (grant numbers 5K99GM135583-02 to S.L.R. and 5R35GM128932-03 to R.C.D.).

References

1.   R. M. Fisher, L. M. Henry, C. K. Cornwallis, E. T. Kiers, S. A. West, The evolution of host-symbiont dependence. *Nat. Commun.* **8**, 15973 (2017).

2.   S. L. Russell, Transmission mode is associated with environment type and taxa across bacteria-eukaryote symbioses: a systematic review and meta-analysis. *FEMS Microbiol. Lett.* **366**, fnz013 (2019).

3.   R. C. Vrijenhoek, "Genetics and evolution of deep-sea chemosynthetic bacteria and their invertebrate hosts" in *The Vent and Seep Biota: Aspects from Microbes to Ecosystems*, S. Kiel, Ed. (Springer, Dordrecht, Netherlands, 2010), pp. 15–49.

4.   M. Bright, S. Bulgheresi, A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol.* **8**, 218–230 (2010).

5.   G. Chomicki, E. T. Kiers, S. S. Renner, The evolution of mutualistic dependence. *Annu. Rev. Ecol. Evol. Syst.* **51**, 409–432 (2020).

6.   D. R. Ginete, H. Goodrich-Blair, From binary model systems to the human microbiome: factors that drive strain specificity in host-symbiont associations. *Front. Ecol. Evol.* **9**, 614197 (2021).

7.   S. C. Verma, T. Miyashiro, Quorum sensing in the squid-*Vibrio* symbiosis. *Int. J. Mol. Sci.* **14**, 16386–16401 (2013).

8.   S. L. Jackrel, J. W. Yang, K. C. Schmidt, V. J. Denef, Host specificity of microbiome
assembly and its fitness effects in phytoplankton. *ISME J.* **15**, 774–788 (2021).
9.   Q. Wang, J. Liu, H. Zhu, Genetic and molecular mechanisms underlying symbiotic
specificity in legume-*Rhizobium* interactions. *Front. Plant Sci.* **9**, 313 (2018).
10.   B. B. Allito, N. Ewusi-Mensah, V. Logah, Legume-*Rhizobium* strain specificity enhances
nutrition and nitrogen fixation in faba bean (*Vicia faba* L.). *Agronomy* **10**, 826 (2020).
11.   L. Walker, B. Lagunas, M. L. Gifford, Determinants of host range specificity in legume-
rhizobia symbiosis. *Front. Microbiol.* **11**, 585749 (2020).
12.   D. C. Smith, A. E. Douglas, *The Biology of Symbiosis*, E. Arnold, Ed. (Cambridge
University Press, 1987).
13.   A. Hilário, *et al.*, New perspectives on the ecology and evolution of siboglinid tubeworms.
PLoS ONE **6**, e16309 (2011).
14.   D. J. Thornhill, E. J. Howells, D. C. Wham, T. D. Steury, S. R. Santos, Population genetics
of reef coral endosymbionts (*Symbiodinium*, Dinophyceae). *Mol. Ecol.* **26**, 2640–2659
(2017).
15.   R. C. Vrijenhoek, M. Duhaime, W. J. Jones, Subtype variation among bacterial
endosymbionts of tubeworms (Annelida: Siboglinidae) from the Gulf of California. *Biol.*
Bull. **212**, 180–184 (2007).
16.   D. T. Pettay, T. C. LaJeunesse, Long-range dispersal and high-latitude environments
influence the population structure of a "stress-tolerant" dinoflagellate endosymbiont. *PLoS*
ONE **8**, e79208 (2013).
17.   P.-T. Ho, *et al.*, Geographical structure of endosymbiotic bacteria hosted by *Bathymodiolus*
mussels at eastern Pacific hydrothermal vents. *BMC Evol. Biol.* **17**, 121 (2017).
18.   S. W. Davies, K. N. Moreland, D. C. Wham, M. R. Kanke, M. V. Matz, *Cladocopium*
community divergence in two *Acropora* coral hosts across multiple spatial scales. *Mol.*
Ecol. **29**, 4559–4572 (2020).
19.   M. Ücker, *et al.*, Deep-sea mussels from a hybrid zone on the Mid-Atlantic Ridge host
genetically indistinguishable symbionts. *ISME J.* (2021) https:/doi.org/10.1038/s41396-
021-00927-9 (accessed August 11, 2021).
20.   N. Dubilier, C. Bergin, C. Lott, Symbiotic diversity in marine animals: the art of
harnessing chemosynthesis. *Nat. Rev. Microbiol.* **6**, 725–740 (2008).
21.   Y. Suzuki, *et al.*, Host-symbiont relationships in hydrothermal vent gastropods of the
genus *Alviniconcha* from the Southwest Pacific. *Appl. Environ. Microbiol.* **72**, 1388–1393
(2006).
22.   K. C. Reynolds, *et al.*, New molluscan larval form: Brooding and development in a
hydrothermal vent gastropod, *Ifremeria nautilei* (Provannidae). *Biol. Bull.* **219**, 7–11
(2010).
23.   R. A. Beinart, *et al.*, Evidence for the role of endosymbionts in regional-scale habitat
partitioning by hydrothermal vent symbioses. *Proc. Natl. Acad. Sci. USA* **109**, E3241–
E3250 (2012).
24.   R. A. Beinart, A. Gartman, J. G. Sanders, G. W. Luther, P. R. Girguis, The uptake and
excretion of partially oxidized sulfur expands the repertoire of energy resources
metabolized by hydrothermal vent symbioses. *Proc. R. Soc. B.* **282**, 20142811 (2015).
25.   C. Breusing, *et al.*, Allopatric and sympatric drivers of speciation in *Alviniconcha*
hydrothermal vent snails. *Mol. Biol. Evol.* **37**, 3469–3484 (2020).

26.   C. Breusing, *et al.*, Physiological dynamics of chemosynthetic symbionts in hydrothermal
606   vent snails. *ISME J.* **14**, 2568–2579 (2020).

607   27.   R. A. Beinart, C. Luo, K. T. Konstantinidis, F. J. Stewart, P. R. Girguis, The bacterial
608   symbionts of closely related hydrothermal vent snails with distinct geochemical habitats
609   show broad similarity in chemoautotrophic gene content. *Front. Microbiol.* **10**, 1818
610   (2019).

611   28.   E. Podowski, S. Ma, G. Luther, D. Wardrop, C. Fisher, Biotic and abiotic factors affecting
612   distributions of megafauna in diffuse flow on andesite and basalt along the Eastern Lau
613   Spreading Center, Tonga. *Mar. Ecol. Prog. Ser.* **418**, 25–45 (2010).

614   29.   M. Tivey, Generation of seafloor hydrothermal vent fluids and associated mineral deposits.
615   *Oceanography* **20**, 50–65 (2007).

616   30.   G. J. Dick, *et al.*, The microbiology of deep-sea hydrothermal vent plumes: ecological and
617   biogeographic linkages to seafloor and water column habitats. *Front. Microbiol.* **4**, 124
618   (2013).

619   31.   G. J. Dick, The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped
620   locally. *Nat. Rev. Microbiol.* **17**, 271–283 (2019).

621   32.   S. V. Nyholm, M. McFall-Ngai, The winnowing: establishing the squid–*Vibrio* symbiosis.
622   *Nat. Rev. Microbiol.* **2**, 632–642 (2004).

623   33.   M. Andrews, M. E. Andrews, Specificity in legume-rhizobia symbioses. *Int. J. Mol. Sci.*
624   **18**, 705 (2017).

625   34.   C. Dahl, B. Franz, D. Hensen, A. Kesselheim, R. Zigann, Sulfite oxidation in the purple
626   sulfur bacterium *Allochromatium vinosum*: identification of SoeABC as a major player and
627   relevance of SoxYZ in the process. *Microbiology* **159**, 2626–2638 (2013).

628   35.   J. M. Rinta-Kanto, *et al.*, Analysis of sulfur-related transcription by *Roseobacter*
629   communities using a taxon-specific functional gene microarray. *Environ. Microbiol.* **13**,
630   453–467 (2011).

631   36.   S. Lehmann, A. W. Johnston, A. R. Curson, J. D. Todd, A. M. Cook, "SoeABC, a novel
632   sulfite dehydrogenase in Roseobacters" in *EMBO Workshop on Microbial Sulfur
633   Metabolism*, G. Muyzer, A. J. Stams, Eds. (Sieca Repro, 2012), p. 29.

634   37.   S. Lenk, *et al.*, *Roseobacter* clade bacteria are abundant in coastal sediments and encode a
635   novel combination of sulfur oxidation genes. *ISME J.* **6**, 2178–2187 (2012).

636   38.   A. Wáren, P. Bouchet, New records, species, genera, and a new family of gastropods from
637   hydrothermal vents and hydrocarbon seeps. *Zool. Scr.* **22**, 1–90 (1993).

638   39.   M. Perez, S. K. Juniper, Insights into symbiont population structure among three
639   vestimentiferan tubeworm host species at Eastern Pacific spreading centers. *Appl. Environ.
640   Microbiol.* **82**, 5197–5205 (2016).

641   40.   T. Ikuta, *et al.*, Heterogeneous composition of key metabolic gene clusters in a vent mussel
642   symbiont population. *ISME J.* **10**, 990–1001 (2016).

643   41.   R. Ansorge, *et al.*, Functional diversity enables multiple symbiont strains to coexist in
644   deep-sea mussels. *Nat. Microbiol.* **4**, 2487–2497 (2019).

645   42.   M.-A. Cambon-Bonavita, J. Aubé, V. Cueff-Gauchard, J. Reveillaud, Niche partitioning in
646   the *Rimicaris exoculata* holobiont: the case of the first symbiotic Zetaproteobacteria.
647   *Microbiome* **9**, 87 (2021).

648   43.   J. Zimmermann, *et al.*, Dual symbiosis with co-occurring sulfur-oxidizing symbionts in
649   vestimentiferan tubeworms from a Mediterranean hydrothermal vent. *Environ. Microbiol.*
650   **16**, 3638–3656 (2014).

44. S. Mitarai, H. Watanabe, Y. Nakajima, A. F. Shchepetkin, J. C. McWilliams, Quantifying
dispersal from hydrothermal vent fields in the western Pacific Ocean. *Proc. Natl. Acad.*
Sci. USA **113**, 2976–2981 (2016).

45. L. A. Levin, D. J. Amon, H. Lily, Challenges to the sustainability of deep-seabed mining.
Nat. Sustain. **3**, 784–794 (2020).

46. F. Blanquart, O. Kaltz, S. L. Nuismer, S. Gandon, A practical guide to measuring local
adaptation. *Ecol. Lett.* **16**, 1195–1205 (2013).

47. L. M. Schiebelhut, M. N. Dawson, Correlates of population genetic differentiation in
marine and terrestrial environments. *J. Biogeogr.* **45**, 2427–2441 (2018).

48. S. L. Seston, *et al.*, Metatranscriptional response of chemoautotrophic *Ifremeria nautilei*
endosymbionts to differing sulfur regimes. *Front. Microbiol.* **7**, 1074 (2016).

49. L. Song, L. Florea, Rcorrector: efficient and accurate error correction for Illumina RNA-
seq reads. *GigaScience* **4**, 48 (2015).

50. M. G. Grabherr, *et al.*, Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

51. J. G. Sanders, R. A. Beinart, F. J. Stewart, E. F. Delong, P. R. Girguis,
Metatranscriptomics reveal differences in in situ energy and nitrogen metabolism among
hydrothermal vent snail symbionts. *ISME J.* **7**, 1556–1567 (2013).

52. W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

53. D. R. Laetsch, M. L. Blaxter, BlobTools: interrogation of genome assemblies. *F1000Res* **6**,
1287 (2017).

54. M. Seppey, M. Manni, E. M. Zdobnov, "BUSCO: Assessing Genome Assembly and
Annotation Completeness" in *Gene Prediction*, M. Kollmar, Ed. (Springer New York,
2019), pp. 227–245.

55. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

56. A. Bankevich, *et al.*, SPAdes: A new genome assembly algorithm and its applications to
single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

57. S. Nurk, D. Meleshko, A. Korobeynikov, P. A. Pevzner, metaSPAdes: a new versatile
metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

58. B. K. B. Seah, H. R. Gruber-Vodicka, gbtools: interactive visualization of metagenome
bins in R. *Front. Microbiol.* **6**, 1451 (2015).

59. G.-C. Xu, *et al.*, LR_Gapcloser: a tiling path-based gap closer that uses long reads to
complete genome assembly. *GigaScience* **8**, giy157 (2019).

60. C. T. Brown, *et al.*, Unusual biology across a group comprising more than 15% of domain
Bacteria. *Nature* **523**, 208–211 (2015).

61. J. Luo, *et al.*, SLR: a scaffolding algorithm based on long reads and contig classification.
BMC Bioinformatics **20**, 539 (2019).

62. L.-X. Chen, K. Anantharaman, A. Shaiber, A. M. Eren, J. F. Banfield, Accurate and
complete genomes from metagenomes. *Genome Res.* **30**, 315–333 (2020).

63. D. D. Kang, *et al.*, MetaBAT 2: an adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

64. Y.-W. Wu, B. A. Simmons, S. W. Singer, MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2016).

65.   B. J. Walker, *et al.*, Pilon: an integrated tool for comprehensive microbial variant detection
697   and genome assembly improvement. *PLoS ONE* **9**, e112963 (2014).
698   66.   T. Brettin, *et al.*, RASTtk: A modular and extensible implementation of the RAST
699   algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci.*
700   *Rep.* **5**, 8365 (2015).
701   67.   D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM:
702   assessing the quality of microbial genomes recovered from isolates, single cells, and
703   metagenomes. *Genome Res.* **25**, 1043–1055 (2015).
704   68.   A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for
705   genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).
706   69.   S. Picelli, *et al.*, Tn5 transposase and tagmentation procedures for massively scaled
707   sequencing projects. *Genome Res.* **24**, 2033–2040 (2014).
708   70.   A. Wilm, *et al.*, LoFreq: a sequence-quality aware, ultra-sensitive variant caller for
709   uncovering cell-population heterogeneity from high-throughput sequencing datasets.
710   *Nucleic Acids Res.* **40**, 11189–11201 (2012).
711   71.   T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation
712   Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).
713   72.   E. Garrison, G. Marth, Haplotype-based variant detection from short-read sequencing.
714   arXiv [Preprint] (2012) https://arxiv.org/abs/1207.3907 (August 11, 2021).
715   73.   H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
716   2078–2079 (2009).
717   74.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
718   (2011).
719   75.   G. A. V. de Auwera, B. D. O'Connor, *Genomics in the Cloud: Using Docker, GATK, and*
720   *WDL in Terra*, First edition (O'Reilly, 2020).
721   76.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and evolutionary
722   analyses in R. *Bioinformatics* **35**, 526–528 (2019).
723   77.   R Core Team, *R: A language and environment for statistical computing* (R Foundation for
724   Statistical Computing, Vienna, Austria, 2020).
725   78.   F. Cailliez, The analytical solution of the additive constant problem. *Psychometrika* **48**,
726   305–308 (1983).
727   79.   H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer New York, 2016).
728   80.   G. Bhatia, N. Patterson, S. Sankararaman, A. L. Price, Estimating and interpreting FST:
729   The impact of rare variants. *Genome Res.* **23**, 1514–1521 (2013).
730   81.   M. C. Whitlock, K. E. Lotterhos, Reliable detection of loci responsible for local adaptation:
731   inference of a null model through trimming the distribution of F_{ST}. *Am. Nat.* **186**, S24–S36
732   (2015).
733   82.   S. Purcell, *et al.*, PLINK: a tool set for whole-genome association and population-based
734   linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
735   83.   F. Beghini, *et al.*, Integrating taxonomic, functional, and strain-level profiling of diverse
736   microbial communities with bioBakery 3. *eLife* **10**, e65088 (2021).
737   84.   R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and
738   bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).
739   85.   M. D. Robinson, A. Oshlack, A scaling normalization method for differential expression
740   analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

86. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).
87. J. Oksanen, *et al.*, *vegan: Community Ecology Package* (2020). https://CRAN.R-
project.org/package=vegan. Deposited 28 November 2020.
88. R. J. Hijmans, *geosphere: Spherical Trigonometry* (2019). https://CRAN.R-
project.org/package=geosphere. Deposited 26 May 2019.
89. B. R. Forester, J. R. Lasky, H. H. Wagner, D. L. Urban, Comparing methods for detecting
multilocus adaptation with multivariate genotype-environment associations. *Mol. Ecol.* **27**,
2215–2233 (2018).
90. M. J. Mottl, *et al.*, Chemistry of hot springs along the Eastern Lau Spreading Center.
Geochim. Cosmochim. Acta **75**, 1013–1038 (2011).
91. G. E. Flores, *et al.*, Inter-field variability in the microbial communities of hydrothermal
vent deposits from a back-arc basin: Microbial communities of hydrothermal deposits.
Geobiology **10**, 333–346 (2012).
92. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*
strain w[1118]; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

 Figures and Tables
759

760
761 **Figure 1** Sampling map for *Alviniconcha* and *Ifremeria* species in the Eastern Lau Spreading Center and Tonga
762 Volcanic Arc.

763
764 **Figure 2** Principal component plots for *Alviniconcha* and *Ifremeria* host species based on genetic covariance
765 matrices.

766
767 **Figure 3** Principal coordinate plots for *Alviniconcha* and *Ifremeria* symbionts based on relative allele counts
768 transformed into Bray-Curtis dissimilarities. Allele counts approximate the relative proportions of different
769 symbiont strains within host individuals.

Figure 4 Principal coordinate plot based on Bray-Curtis dissimilarities (A) and presence/absence heatmap of differentially preserved genes (B) for the Gamma1 symbiont of *A. kojimai* and *A. strummeri*. Strains of this symbiont are clearly distinct between the two host species, even when these taxa co-occur (see Fig. S3). Segregation of symbiont populations by host species along the first ordination axis suggests that host affinity is a stronger predictor than geography for symbiont composition between species. Dark and light blue colors in the heatmap indicate presence and absence of genes, respectively. Dendrograms show similarities between samples based on their gene content profiles.

779
780　**Figure 5** Presence/absence heatmap for differentially preserved genes.

781
782 **Figure 6** Redundancy analysis plots of the first two ordination axes for *Alviniconcha* and *Ifremeria* symbionts.
783 Small grey dots in the center of the plot indicate genetic variants, while large colored dots represent intra-host
784 symbiont populations sampled from different vent localities. Vectors show the environmental predictors.
785 Redundancy analyses were conditioned by geography for all symbionts except for Gamma1 and GammaLau.
786 Genotype-environment associations were not significant for GammaLau.

Table 1 Sampling information for *Alviniconcha* and *Ifremeria* species from the Eastern Lau Spreading Center and Tonga Volcanic Arc. Chemical concentrations are mean endmember values for each vent field obtained from Beinart et al. [23], Mottl et al. [88], Flores et al. [89], and unpublished data provided by J. Seewald (Woods Hole Oceanographic Institution) and A. Diehl (MARUM). n.a. = not available.

Vent site	Species	Coordinates	Year	Depth [m]	H_2 [mM]	CO_2 [mM]	CH_4 [mM]	H_2S [mM]	Fe [mM]	Mn [mM]
Tonga Volcanic Arc										
Niua South	*A. boucheti* *I. nautilei*	15°09.79'S 173°34.48'W	2016	1164	0.037	n.a.	0.005	1.911	n.a.	n.a.
Eastern Lau Spreading Center										
Kilo Moana	*A. boucheti*	20°03.23'S 176°08.01'W	2009	2614	0.382	8.227	0.031	5.432	3.062	0.643
Tow Cam	*A. boucheti* *I. nautilei*	20°18.97'S 176°08.19'W	2009	2711–2722	0.150	10.648	0.045	4.795	0.316	0.389
Tahi Moana	*A. strummeri* *I. nautilei*	20°40.94'S 176°11.01'W	2016	2234–2273	0.097	7.191	0.038	3.548	0.286	0.525
ABE	*A. boucheti* *A. kojimai* *I. nautilei*	20°45.79'S 176°11.47'W	2009 2016	2130–2152	0.079	5.980	0.048	3.060	0.245	0.330
Tu'i Malila	*A. kojimai* *A. strummeri* *I. nautilei*	21°59.36'S 176°34.10'W	2009 2016	1860–1887	0.084	12.912	0.037	2.294	0.203	0.406