Host-symbiont population genomics provide insights into partner fidelity, transmission

mode and habitat adaptation in deep-sea hydrothermal vent snails

Corinna Breusing[1*], Maximilian Genetti[2], Shelbi L. Russell[3], Russell B. Corbett-Detig[2], Roxanne

A. Beinart[1*]

[1]University of Rhode Island, Graduate School of Oceanography, Narragansett, RI 02882, USA

[2]University of California Santa Cruz, Jack Baskin School of Engineering, Santa Cruz, CA 95064,

USA

[3]University of California Santa Cruz, Department of Molecular, Cell, and Developmental

Biology, Santa Cruz, CA 95064, USA

*Corresponding author: corinnabreusing@gmail.com, rbeinart@uri.edu

Running title: Host-symbiont population genomics in deep-sea hydrothermal vent snails

Keywords: Chemosynthetic symbiosis, hydrothermal vents, host-symbiont population genomics,

habitat adaptation, symbiont transmission, symbiont specificity

Abstract

Symbiont specificity, both at the phylotype and strain level, can have profound consequences for

host ecology and evolution. However, except for insights from a few model symbiosis systems,

the degree of partner fidelity and the influence of host versus environmental factors on symbiont

composition are still poorly understood. Nutritional symbioses between invertebrate animals and

chemosynthetic bacteria at deep-sea hydrothermal vents are examples of relatively selective

associations, where hosts affiliate only with particular phylotypes of gammaproteobacterial or

campylobacterial symbionts. In hydrothermal vent snails of the sister genera *Alviniconcha* and

Ifremeria this phylotype specificity has been shown to play a role in habitat distribution and

partitioning among different holobiont species. However, it is currently unknown if fidelity goes

beyond species level associations that might influence genetic structuring, connectivity and

habitat adaptation of holobiont populations. We used metagenomic analyses to assess sequence

variation in hosts and symbionts and identify correlations with geographic and environmental

factors. Our analyses indicate that host populations are not differentiated across a ~800 km

gradient, while symbiont populations are clearly structured between vent locations due to a

combination of neutral and selective processes. Overall, these results suggest that host

individuals flexibly associate with local strains of their specific symbiont phylotypes, which

confirms a model of horizontal transmission in both *Ifremeria* and *Alviniconcha*. Strain

flexibility in these snails likely enables host populations to exploit a range of habitat conditions,

which might favor wide-spread genetic connectivity and ecological resilience unless physical

dispersal barriers are present.

Introduction

Mutualistic relationships between eukaryotes and bacterial microbes are ubiquitous in nature.

Symbionts enable hosts to gain access to novel resources and habitats, provide protection against

pathogens and predators, and can be essential for the host's diet (Fisher et al. 2017; Russell

2019). For symbiotic associations to persist over evolutionary time, hosts must successfully

transmit their symbionts from one generation to the other, either through symbiont acquisition

from the environment (horizontal transmission), direct inheritance of symbiont lineages through

the host germline (vertical transmission) or a combination of both mechanisms (mixed

transmission) (Vrijenhoek 2010; Bright and Bulgheresi 2013). The mode of transmission has

significant implications for the composition and variation of symbionts within and between host

individuals. Vertical transmission typically results in strong genetic coupling between host and

symbiont lineages and an accompanied reduction in intra-host symbiont diversity (Vrijenhoek

2010). By contrast, horizontal transmission exposes aposymbiotic hosts to a potentially

heterogenous pool of free-living symbiont lineages (Vrijenhoek 2010), which can promote the

formation of generalist partnerships, where multiple hosts and symbionts associate with each

other, to more specialized associations between only one or a few potential partners (Fisher et al.

2017; Chomicki et al. 2020). This range is often referred to as host-symbiont specificity, which

can vary in its taxonomic level for both partners depending on the symbiotic system.

The degree of partner fidelity and its effect on symbiont composition can have dramatic

impacts on holobiont functioning (Ginete and Goodrich-Blair 2021). For example, host-symbiont

specificity, both at the species and genotype level, is crucial for light production in

bioluminescent squid (Verma and Miyashiro 2013), while shifts in microbiome assemblages

have been shown to affect phytoplankton growth rates (Jackrel et al. 2021) and the efficiency of

nitrogen fixation in nodule-forming legumes (Wang et al. 2018; Allito et al. 2020; Walker et al.

2020). Despite its importance for host biology, the taxonomic level of partner specificity as well

as the relative contributions of host genetic, environmental and geographical factors to symbiont

composition remain understudied in the majority of horizontally transmitted symbioses.

Chemosynthetic animal-microbe symbioses are globally significant phenomena that

dominate hydrothermal vent and hydrocarbon seep ecosystems in the deep sea. In these

associations, the bacterial partner uses chemical energy from the oxidation of reduced fluid

compounds, such as hydrogen, sulfide or methane, to convert carbon to organic matter, which

serves as primary nutrition for the host (Dubilier et al. 2008). Due to the nutritional dependence

on their bacterial symbionts, animal hosts are typically highly selective in their partner choice: In

the predominant number of cases host individuals associate with only 1–2 phylotypes of

gammaproteobacterial or campylobacterial symbionts (Dubilier et al. 2008), whereas symbionts

can exhibit a comparatively broad host range. Chemosynthetic endosymbioses in deep-sea snails

of the sister genera *Alviniconcha* and *Ifremeria* are examples of reciprocally relatively specific

partnerships, which dominate hydrothermal vent systems in the Southwest Pacific back-arc

basins. Given the absence of host-symbiont phylogenetic concordance, the symbionts are

assumed to be environmentally acquired (Suzuki et al. 2006), although a pseudo-vertical

transmission component is possible in *Ifremeria* given its brooding reproductive mode (Reynolds

et al. 2010). In the Eastern Lau Spreading Center (ELSC), previous work suggested that

specificity to functionally distinct symbiont phylotypes drives local and regional-scale habitat

partitioning among four co-occurring *Alviniconcha* and *Ifremeria* species (Beinart et al. 2012,

2015; Breusing et al. 2020a, b). *Alviniconcha boucheti* contains a campylobacterial phylotype

(Epsilon) and is usually found at northern vent sites with high concentrations of sulfide and

hydrogen, while *A. kojimai* and *A. strummeri* associate with different gammaproteobacterial

phylotypes (Gamma1 and Gamma1/GammaLau, respectively) and usually occupy mid-latitude

to southern vent sites where the concentrations of these chemical reductants are lower (Beinart et

al. 2012, 2019; Breusing et al. 2020a, b). *Ifremeria nautilei* establishes dual symbioses with

thiotrophic and methanotrophic gammaproteobacterial endosymbionts (Beinart et al. 2019) and

is co-distributed with *Alviniconcha* across their geographical range, although it typically

segregates into habitat patches with reduced fluid flow relative to its sister genus (Podowski et al.

2010). While it has been well-established that niche differentiation across hydrothermal vents is

mediated by symbiont phylotype specificity in *Alviniconcha* and *Ifremeria* host species, nothing

is known about the fidelity of these associations at the population level, nor how regional

adaptation is conferred functionally.

In this study we applied population genomic methods to assess symbiont strain-level

genetic variation and patterns of host-symbiont genetic subdivision in *Alviniconcha* and

Ifremeria species from the ELSC and Tonga Volcanic Arc. Using multivariate statistical

analyses, we evaluated the impact of host traits, environment and geography on symbiont

composition in populations of both genera and identified potential candidate genes for local

adaptation.

Materials and Methods

Sample collection, nucleic acid extraction and sequencing

Samples of *Alviniconcha* and *Ifremeria* were collected from six vent sites (1164–2722 m) of the

Lau Basin and Tonga Volcanic Arc in 2009 and 2016 using remotely operated vehicles (Fig. 1;

Table 1). Upon recovery, animal samples were dissected, placed in RNALater™ Stabilization

Solution (Thermo Fisher Scientific, Inc.) and frozen at −80°C until further analysis. DNA was

extracted with the Quick-DNA 96 Plus extraction kit (Zymo Research, Inc.) and further purified

with the MO BIO PowerClean DNA Pro Clean-Up kit (Qiagen, Inc.). High molecular weight

(HMW) DNA for long-read sequencing was isolated with Qiagen Genomic-tips following

manufacturer's instructions.

Host transcriptome sequencing and assembly

Illumina RNAseq reads for host transcriptome assemblies were obtained from sequencing

experiments performed in Seston et al. (2016) and Breusing et al. (2020b). Adapter clipping,

quality trimming, and removal of rRNA and symbiont reads was performed as in Breusing et al.

(2020b). Cleaned host reads were error corrected with RCORRECTOR (Song and Florea 2015) and

filtered for uncorrectable and overrepresented sequences with the

TRANSCRIPTOMEASSEMBLYTOOLS package

(https://github.com/harvardinformatics/TranscriptomeAssemblyTools). Host transcriptome co-

assemblies for each *Alviniconcha* and *Ifremeria* species were performed with TRINITY (Grabherr

et al. 2011) using the PASAFLY algorithm. For each *Alviniconcha* species, additional transcripts

were reconstructed from 454 reads obtained from Sanders et al. (2013). Assembled contigs for

each species were clustered with CD-HIT-EST (Li and Godzik 2006) at a 95% identity threshold

to reduce transcript redundancies. Open reading frames (ORFs) were predicted with

TRANSDECODER (https://github.com/TransDecoder/TransDecoder) considering homologies to

known proteins (UniRef90) and protein domains (Pfam) as ORF retention criteria. Transcripts

that did not contain any ORF or had a non-eukaryotic origin based on taxonomy classifications

with BLOBTOOLS (Laetsch and Blaxter 2017) were removed from the assembly. Final

transcriptome assemblies were evaluated for quality and completeness with BUSCO (Seppey et al.

2019).

Symbiont genome sequencing and re-assembly

Illumina reads for the *A. boucheti*, *A. kojimai* and *Ifremeria* holobionts were obtained from

previous sequencing runs performed in Beinart et al. (2019). Raw reads were trimmed with

TRIMMOMATIC (Bolger et al. 2014) and filtered for sequence contaminants through mapping

against the human (GRCh38) and PhiX reference genomes. Decontaminated reads were grouped

into symbiont and host reads by assessing best matches against draft symbiont genomes (Beinart

et al. 2019) with BBSPLIT (https://sourceforge.net/projects/bbmap/). To improve contiguity of the

genome assemblies we conducted 3–4 Nanopore sequencing runs for each symbiont phylotype

on a MinION device (Oxford Nanopore Technologies) using the SQK-LSK109 ligation kit after

HMW DNA enrichment with the Circulomics Short Read Eliminator kit. Basecalling of the

Nanopore reads was done with ALBACORE (Oxford Nanopore Technologies) and adapters were

clipped with PORECHOP (https://github.com/rrwick/Porechop). Hybrid assemblies of Illumina

and Nanopore reads were constructed for each symbiont genome with SPADES (Gamma1)

(Bankevich et al. 2012) or METASPADES (all others) (Nurk et al. 2017) choosing k-mers between

21 and 91 in 10 step increments. Raw assemblies for the Epsilon, Gamma1 and Ifr-SOX

symbionts were binned with GBTOOLS (Seah and Gruber-Vodicka 2015) and incrementally gap

filled, corrected and scaffolded with LR_GAPCLOSER (Xu et al. 2019), ra2.py (Brown et al.

2015), and SLR (Luo et al. 2019), respectively, following recommendations by Chen et al.

(2020). The Ifr-MOX symbiont assembly was automatically binned with METABAT2 (Kang et

al. 2019) for contigs ≧ 1500 bp. Shorter contigs (≧ 500 bp) were binned with MAXBIN (Wu et al.

2016). Scaffolds < 200 bp were excluded from all assemblies. Final assemblies were polished

with PILON (Walker et al. 2014), functionally annotated with RAST-TK (Brettin et al. 2015) and

quality-checked with CHECKM (Parks et al. 2015) and QUAST (Gurevich et al. 2013).

Population-level metagenomic sequencing and variant identification in hosts and symbionts

192 barcoded high-throughput DNA sequencing libraries were prepared with a Tn5 transposase-

based protocol after Picelli et al. (2014) at the University of California Santa Cruz and then sent

for 150 bp paired-end sequencing on a NovaSeq 6000 instrument at the University of California

Davis. However, due to low read allocation 66 of the libraries were excluded from further

analysis. Raw sequence reads were trimmed, filtered and sorted by host species and symbiont

phylotype as described above. Optical duplicates were removed with PICARD's

MARKDUPLICATES tool (https://github.com/broadinstitute/picard). To resolve common alignment

errors and improve base call accuracy, we locally realigned reads around indels and recalibrated

base quality scores with LOFREQ (Wilm et al. 2012).

Host population genomic variation was assessed in ANGSD (Korneliussen et al. 2014) by

inferring genotype likelihoods based on Hardy-Weinberg equilibrium considering individual

inbreeding coefficients. To increase accuracy of the analyses, variant sites with mapping

qualities < 30 (minMapQ = 30), base qualities < 20 (minQ = 20), and minimum minor allele

frequencies < 0.01 (minMaf = 0.01) were excluded. We further filtered sites based on strand bias

(sb_pval = 0.05), heterozygote bias (hetbias_pval = 0.05) and probability of being variable

(SNP_pval = 1e-6). In addition, we removed spurious and improperly paired reads, adjusted

mapping qualities for excessive mismatches (C = 50) and computed per-base alignment qualities

(BAQ = 1) to disregard variants close to indel regions. Putative paralogous variants were

excluded by discarding reads with multiple mappings and by considering only sites that had a

maximum depth of 40–80. Genetic distances between individuals were inferred by calculating

pairwise covariance matrices.

Symbiont population genomic variation was determined with FREEBAYES (Garrison and

Marth 2012) using input parameters adjusted for the analysis of metagenomic data (-F 0.01 -C 1

-p 1 --pooled-continuous --haplotype-length 0 --report-monomorphic). Variant calls were

restricted to sites with a minimum base quality of 20, a minimum mapping quality of 30, and a

minimum coverage of 10. To eliminate bias in variant identification and other downstream

analyses due to uneven read depth between samples, we normalized all samples to the lowest

amount of coverage found in a sample for a particular symbiont phylotype (> 10X coverage).

Variants were further filtered based on strand bias (SRP > 5 && SAP > 5 && EPP > 5),

proximity to indels (5 bp) and maximum mean depth with BCFTOOLS (Li et al. 2009) and

VCFTOOLS (Danecek et al. 2011). In addition, sites and individuals with more than 25% missing

data were excluded from the analysis. Allele counts (= symbiont strain abundances) and

consensus haplotypes (= dominant symbiont strains) were extracted with GATK's

VARIANTSTOTABLE tool (Van der Auwera and O'Connor 2020).

Genomic structure and differentiation

We performed ordination analyses with the APE and STATS packages in R (Paradis and Schliep

2019; R Core Team 2020) to assess genetic variation between symbiont and host populations.

Host genetic structure was determined through principal component analyses based on genetic

covariance matrices, while symbiont genetic structure was inferred through principal coordinate

analyses based on both consensus haplotype and allele count data transformed into Euclidean

distances and Bray-Curtis dissimilarities, respectively. For each sample, absolute allele counts

were normalized to relative counts prior to analysis. Negative eigenvalues were corrected using

the method by Cailliez (1983) and final ordination plots were produced with GGPLOT2

(Wickham 2016). F_{ST} values between host and symbiont populations were calculated in ANGSD

and SCIKIT-ALLEL (https://github.com/cggh/scikit-allel), respectively, following the procedure

by Bhatia et al. (2013). For the host species, potential outlier loci under selective pressures were

inferred with OUTFLANK (Whitlock and Lotterhos 2015) based on neutral F_{ST} distributions that

were obtained from quasi-independent subsets of single nucleotide polymorphisms (SNPs)

determined with PLINK (Purcell et al. 2007).

Assessment of gene content variation

We used PANPHLAN (Beghini et al. 2020) to investigate potential differences in gene

composition between symbiont strains from different vent localities. Downsampled symbiont

reads for each species were mapped against the corresponding symbiont reference genome using

custom RAST-TK annotations for functional categorizations. Samples were profiled for gene

presence/absence based on the following parameter thresholds: --min_coverage 1 --left_max

1.70 --right_min 0.30. Gene content variation between symbiont strains was assessed by

identifying genes that were present/absent in 90% of samples from one or the other geographic

region. To account for gene content variation between different strains within hosts we further

quantified gene abundances with SALMON (Patro et al. 2017) using the following parameters: -l

IU --meta --rangeFactorizationBins 4 --numBootstraps 1000 --seqBias --gcBias -s -u. Abundance

values were normalized with the Trimmed Mean of M method using TRINITY's

abundance_estimates_to_matrix.pl script (Robinson and Oshlack 2010; Grabherr et al. 2011).

Gene presence/absence and differential abundance heatmaps were produced with the

COMPLEXHEATMAP package in R (Gu et al. 2016).

Assessment of symbiont variation based on environment and host genetics

We conducted partial Mantel tests with the VEGAN package in R (Oksanen et al. 2020) to assess

relationships between host and symbiont genetic distances and geography based on Spearman

rank correlations. As no population genetic structure could be observed in any host species based

on the full SNP datasets, we estimated host covariance matrices using only moderately to highly

differentiated SNP subsets for this analysis (*Alviniconcha*: $F_{ST} > 0.15$; *Ifremeria*: $F_{ST} > 0.10$).

Geographic distances were determined by calculating the geodesics between vent sites with the

GEOSPHERE package (Hijmans 2019). We used redundancy analyses (RDA) following the

approach in Forester et al. (2018) (https://popgen.nescent.org/2018-03-27_RDA_GEA.html) to

evaluate the influence of hydrothermal fluid composition and depth on the genetic structure of

each symbiont phylotype. We further included the effect of sampling year, which encompasses

changes in hydrothermal circulation within the ELSC as indicated by the cessation in fluid flow

at the Kilo Moana vent field between 2009 and 2016. Endmember concentrations for

geochemical compounds were obtained from the literature or unpublished data (Mottl et al. 2011;

Beinart et al. 2012; Flores et al. 2012; Seewald, unpubl.; Diehl, unpubl.; Table 1). Due to multi-

collinearity among the chemical species, we used the first eigenvector from principal coordinate

analyses (corresponding to the eigenvalue with the largest explanatory power) as composite

value. For each symbiont phylotype, we further assessed the strength of correlation with other

predictors to exclude variables that were highly collinear. Unless geographic location was

strongly linked to environmental predictors, we used latitude as a conditioning factor in the

analyses to correct for isolation by distance. Based on these collinearity evaluations, we tested

the effect of all factors on the two *Ifremeria* symbionts, the effect of depth and fluid composition

on the *A. boucheti* symbiont and the effect of fluid composition and year (uncorrected for

geography) on the *A. kojimai* and *A. strummeri* symbionts. SNPs were considered candidates for

local adaptation if their loadings on significant constrained RDA axes deviated more than 2.5

standard deviations from the mean of the distribution. For each symbiont phylotype we further

calculated the ratio of non-synonymous to synonymous polymorphisms (pN/pS) with SNPEFF

(Cingolani et al. 2012) to infer candidate genes evolving under natural selection.

Results

Host transcriptome assemblies and population genomic structure

Host transcriptome assemblies consisted of 24,176–35,654 transcripts (totaling 20.68–28.68 Mb)

and were approximately 30.30–55.40% complete (Table S1). Mapping of host reads against the

transcriptome references and subsequent filtering of variant sites yielded 1,655–9,185 SNPs per

species for population genetic analyses. Irrespective of host taxon, F_{ST} and ordination analyses

revealed low genetic differentiation among host populations sampled from different vent

localities (Fig. 2; Table S2). With the exception of four SNP sites in *A. kojimai*, no F_{ST} outliers

could be detected in any host species (Table S3). However, all species contained a number of

SNPs that were moderately to highly divergent among host populations. When analyses were

constrained to these SNP subsets (*Alviniconcha*: $F_{ST} > 0.15$; *Ifremeria*: $F_{ST} > 0.10$), population
genetic structuring by vent site was observed in all *Alviniconcha* species, but not *Ifremeria* (Fig.
S1). For both *A. kojimai* and *A. strummeri*, the respective SNP subsets further indicated genetic
differences between Tui Malila populations that were sampled in different years. The effect of
geography on host population genetic structuring was not significant after correction for
symbiont genetic distance in partial Mantel correlations (Table S4).

Symbiont genome assemblies and population genomic structure
Symbiont genome assemblies varied in size from 2.10–4.86 Mb and contained 2,006–6,660
predicted protein-coding genes, with GC content ranging from 34.30–59.00% (Table S2). All
assemblies were characterized by a high level of completeness (92.79–99.25%) and low amount
of contamination (0.74–5.48%) (Table S5). Symbiont populations for the Epsilon, Gamma1, Ifr-
SOX and Ifr-MOX phylotypes were largely structured by vent field or broader geographic region
based on 239–7,057 variant sites (Fig. 3, S2; Table S6). These associations were significant even
when corrected for host genetic variation based on highly differentiated SNP markers (Table S4;
$p \leqq 0.0244$, $r = 0.2665–0.8705$). Genetic differentiation between symbiont populations typically
increased with geographic distance between vent fields (Table S6). Although F_{ST} values for
GammaLau populations of *A. strummeri* were moderate when calculated between distinct vent
sites (Table S6), ordination analyses and Mantel tests provided no evidence for genetic
differentiation of this symbiont phylotype across geographic locations based on 192 marker loci
(Fig. 3, S2). The Gamma1 phylotype is associated with both *A. kojimai* and *A. strummeri* and we
therefore investigated whether populations of this phylotype varied between host species.
Ordination analyses indicated a clear clustering by host taxon that superseded the effect of
geography (Fig. 4, S3), suggesting that different Gamma1 strains associate selectively with either
A. kojimai or *A. strummeri*. By contrast, the effect of host genetics on symbiont genetic variation
within species appeared to be weak. Although partial Mantel tests suggested significant
associations of host genotype with strain composition or dominant symbiont type for the *A.
boucheti* – Epsilon, *A. kojimai* – Gamma1, *A. strummeri* – Gamma1, and *I. nautilei* – Ifr-SOX
pairs, *r* statistics were relatively low especially compared to the effect of geography, indicating
limited biological relevance (Table S4; $p \leqq 0.0364$, $r = 0.0510–0.2270$).

Gene content variation between symbiont strains

We assessed variation in gene content between symbiont strains from different vent localities or

broader geographic regions (Fig. 5, S4; Table S7). For the Gamma1 symbiont, we further

determined gene content variation between strains from different host species given that this

symbiont phylotype occurs in both *A. strummeri* and *A. kojimai* (Fig. 4, S3; Table S8).

Differentially preserved or abundant genes between symbiont strains comprised about 1–5% of

protein-coding regions. Independent of symbiont phylotype, the most common differences

among geographic or host-specific strains concerned genes of unknown function as well as a

smaller number of genes related to mobilome and anti-viral defense (Fig. 4–5, S3–4; Table S7,

S8). Geographic strains of the *A. boucheti* Epsilon symbiont further differed in the presence of an

ABC transporter, a GDP-L-fucose synthetase, a NAD(FAD)-utilizing hydrogenase and the DNA

repair protein RecN, which were conserved in strains from Kilo Moana and Tow Cam but were

absent or very lowly abundant in strains from ABE (Table S7). Within the *A. kojimai* Gamma1

phylotype, strains from ABE (but rarely Tui Malila) contained a few genes involved in DNA,

protein and cell wall metabolism as well as maturation and regulation of uptake [NiFe]

hydrogenases (*hyaC, hyaD, hoxJ*). Differences in hydrogenase-related genes were also observed

in the Gamma1 symbiont of *A. strummeri*, where an operon for a hydrogen-sensing hydrogenase

(*hupUV/hoxBC*) and genes for hydrogenase maturation and assembly proteins (*hypF, hyaF,*

hoxV/hupK, hypD, hypE, hoxX) were largely missing in strains from Tui Malila (but not Tahi

Moana) (Table S7). Between host species, Gamma1 strains notably differed in the presence of

genes for a sulfite dehydrogenase complex (*soeABC*), which was abundant in strains specific to

A. kojimai but not *A. strummeri* (Table S8). Within the *A. strummeri* GammaLau phylotype,

strains from Tui Malila contained a broad range of metabolic genes that were absent or

infrequent in strains from Tahi Moana, including genes related to macronutrient metabolism,

stress response, membrane transport and several other functional processes (Table S7). The Ifr-

SOX strains of *I. nautilei* differed in several genes related to DNA, nitrogen, lipid and amino

acid metabolism, membrane transport, cell regulation and detoxification, which were present in

strains from the ELSC but mostly missing in strains from Niua South. Within the Ifr-MOX

phylotype, strains from Tahi Moana and Tui Malila differed in genes involved in sulfide

respiration and oxidative phosphorylation, macronutrient metabolism, cation transport, motility,

cofactor, cell wall, DNA and nucleotide metabolism as well as cell regulation and stress response

(Table S7).

Adaptive variation within symbiont phylotypes

A subset of genetic variants exhibited significant associations with environmental factors in all

symbiont phylotypes except for GammaLau, with constrained ordinations explaining between

5.39% and 19.76% of the variance in the respective RDA models (Fig. 6; Table S9). Candidate

adaptive loci for each symbiont phylotype encompassed a variety of metabolic categories,

although in many cases no particular function could be assigned (Table S9). In the *A. boucheti*

Epsilon symbiont, 129 variants were significantly associated with fluid composition, while

another 47 were correlated with depth. These variants were mostly located in genes related to

protein, amino acid, and cell wall metabolism, followed by cofactor, DNA, carbon and

nucleotide metabolism, as well as membrane transport (Table S9). Several other variants were

linked to nitrogen, iron and RNA metabolism, virulence, motility, cell signaling, stress response,

respiration, cell cycle, as well as hydrogen and sulfur metabolism. In the *A. kojimai* Gamma1

symbiont, 91 variants were correlated with fluid composition and 9 were linked to year. 29 of

these variants were classified into functional categories. The most represented categories were

mobilome, cell wall and DNA metabolism, anti-viral defense, membrane transport, cofactor

metabolism and respiration. A few variants were associated with amino acid, hydrogen, protein

and RNA metabolism, and cell regulation. In the *A. strummeri* Gamma1 symbiont, 95 variants

showed associations with fluid composition, while 3 were correlated with year. 54 of these

variants could be functionally annotated and assigned to the following metabolic categories:

protein, amino acid and cell wall metabolism, membrane transport, virulence, carbon, nitrogen,

DNA/RNA and nucleoside metabolism, cofactor, fatty acid and sulfur metabolism, stress

response, cell signaling and mobilome (Table S9). The *I. nautilei* Ifr-SOX and Ifr-MOX

symbionts contained 94 and 11 putatively adaptive variants, respectively. In both symbionts,

these variants were mostly linked to depth, followed by fluid composition and year. While the

majority of variants were located in genes with unknown functions, a number of loci was linked

to mobile elements, cell cycle, anti-viral defense, DNA/RNA metabolism, and membrane

transport. In each symbiont phylotype, virtually all candidate variants were characterized by

elevated F_{ST} and gene-wide pN/pS values, further supporting their potential role in habitat

adaptation (Table S9). Several other genes exhibited increased ratios of non-synonymous to

synonymous substitution, although sites within these genes did not show significant associations

with environment (Table S10). It is possible that sites within these genes covary with other

ecological factors that could not be tested in this study or that these patterns are caused by recent

slightly deleterious mutations that were not yet purged by natural selection.

Discussion

Strain-level variation within microbial symbionts is increasingly recognized as an important

driver of host ecology and evolution (Ginete and Goodrich-Blair 2021), yet its patterns,

determining factors and functional implications remain poorly investigated in many non-model

symbioses. In this study, we used metagenomic analyses to assess sequence and gene content

differences between chemosynthetic symbionts associated with four co-occurring species of

deep-sea hydrothermal vent snails and determined the impact of host genetic, geographical and

environmental factors on symbiont strain composition and variation.

Despite fidelity between hosts and symbionts at the species level (Beinart et al. 2012;

Breusing et al. 2020a), our results indicate that specificity between host genotypes and symbiont

strains in *Alviniconcha* and *Ifremeria* is weak: Host populations were not partitioned across a

~800 km gradient, whereas symbiont populations were clearly structured between vent locations

or broader geographic regions. Even when we correlated symbiont and host genetic distances

based on highly differentiated markers, test statistics for significant associations remained low,

suggesting that host genetics has a minor impact compared to environment or geography on

strain composition within both *Alviniconcha* and *Ifremeria*.

These findings qualitatively agree with observations in deep-sea mussel hybrids that

appear to associate with locally available symbiont strains (Ücker et al. 2021), but contrast

markedly with patterns in other horizontally transmitted partnerships, such as the squid-*Vibrio*

symbiosis and some legume-rhizobia associations, where hosts exhibit strong strain specificity

(reviewed in Ginete and Goodrich-Blair 2021). Environmental uptake of locally adapted

symbiont strains provides host organisms with the opportunity to optimally exploit novel

habitats, but carries the risk of unsuccessful symbiont acquisition and infection by cheaters

(Vrijenhoek 2010). Depending on the amount of partner reliance, chances of symbiont encounter

and fitness variation among symbiont strains, holobionts likely find different tradeoffs between

these opposing factors. Associations with chemosynthetic bacteria are obligate for hydrothermal

vent animals, but are restricted to relatively ephemeral habitats that are characterized by large

temporal and spatial fluctuations in environmental conditions and associated shifts in microbial

communities (Tivey 2007; Dick et al. 2013; Dick 2019). Strong nutritional dependency in vent

symbioses combined with the uncertainty of habitat (and thus symbiont) encounter might

promote specificity towards a mutualistic symbiont phylotype, while enabling enough flexibility

towards different local strains of that phylotype to maximize recruitment success of host larvae at

a new vent site. By contrast, bobtail squids inoculate their environment with symbiotic bacteria

and thereby ensure symbiont availability for their offspring (reviewed in Nyholm and McFall-

Ngai 2004), which might favor increased strain selectivity in this association. Similarly, some

legume species appear to preferentially associate with certain rhizobial strains that are abundant

in the native host range (Andrews and Andrews 2017). Although the factors underlying strain

specificity are not fully understood, it is possible that the relative fitness advantages provided by

these locally available strains (Andrews and Andrews 2017) coupled with the benefit of

decreased cheater invasion (Walker et al. 2020) might have selected for strong partner fidelity in

these symbioses.

Alviniconcha kojimai and *A. strummeri* represent notable exceptions to the observed

patterns, as they share identical phylotypes of one gammaproteobacterial symbiont (Gamma1),

but appear to take up different strains even in habitats where these host species co-occur. The

phylogenetic divergence between *A. kojimai* and *A. strummeri* (~25 MYA) is more recent than

that between either species and *A. boucheti* (~38 MYA) or *I. nautilei* (~113 MYA) (Breusing et

al. 2020a). Perhaps the closer evolutionary relationship between these two species favors

associations with the same symbiont phylotype, as has been suggested for *Bathymodiolus*

azoricus and *B. puteoserpentis* on the Mid-Atlantic Ridge as well as *B. thermophilus* and *B.*

antarcticus on the East Pacific Rise (Ücker et al. 2021). However, compared to the

bathymodiolin mussel system, the split between *A. kojimai* and *A. strummeri* is significantly

older (>17 MYR), which could indicate that the timing of phylotype specificity evolution varies

among taxonomic groups, possibly as a result of contrasting ecological or evolutionary contexts.

The fact that *A. kojimai* and *A. strummeri* nevertheless associate with distinct symbiont strains in

sympatry is potentially a mechanism to avoid competition for niche space. Gamma1 strains of

these two species notably differed in the presence of a molybdenum-containing sulfite

dehydrogenase (SoeABC), which was preserved in strains of *A. kojimai* but not *A. strummeri*.

SoeABC is the predominant enzyme for sulfite oxidation in many purple sulfur bacteria (Dahl et

al. 2013) and is further involved in taurine and dimethylsulfoniopropionate degradation in

Roseobacter clade bacteria (Rinta-Kanto et al. 2011; Lehmann et al. 2012; Lenk et al. 2012).

Although the physiological role of SoeABC in *A. kojimai*'s Gamma1 strain is unknown, it is

possible that it contributes to partitioning of sulfur resources among co-occurring snail

holobionts.

Our data support a model of horizontal symbiont transmission in both *Alviniconcha* and

Ifremeria. While these results are expected for *Alviniconcha* which produce free-swimming

planktotrophic larvae and do not invest in their young (Warén and Bouchet 1993), they are rather

surprising for *Ifremeria* which brood their offspring in a modified pouch in the female's foot

(Reynolds et al. 2010) and would thus have the opportunity to pseudo-vertically transmit their

symbionts. It is possible that a pseudo-vertical transmission component exists in *Ifremeria*, but

that maternally acquired symbionts get replaced or complemented by more competitive strains in

the habitat where the snail larvae settle. Our current dataset cannot distinguish this possibility

from a strict horizontal transmission mode, though future studies assessing symbiont

composition in different developmental stages of *Ifremeria* would be helpful to address this

hypothesis.

While the majority of genetic variants in the symbionts did not deviate from neutral

expectations, a subset of loci showed evidence for natural selection, implying a role of both

genetic drift and local adaptation in shaping symbiont population structure. Candidate adaptive

loci spanned a surprisingly broad range of metabolic functions, many of which were probably

not causally linked to the investigated environmental predictors but other correlated variables

that we could not account for in this study. For example, in almost all symbiont phylotypes we

observed adaptive variation in genes that were related to anti-viral defense (e.g., CRISPR-

associated proteins, restriction-modification systems) or mobile elements. Although differences

in depth or geochemistry might contribute to these patterns, it is more likely that they reflect

exposure of symbiont strains to distinct viral assemblages that might covary with local habitat

conditions, as has been suggested for vestimentiferan tubeworm symbionts (Perez and Juniper

2016). Polymorphisms in other genes, by contrast, are likely directly explained by variation in

the analyzed environmental factors. Geographic strains of *A. boucheti*'s Epsilon symbiont, for

instance, contained several variants under positive selection that were located in genes involved

in hydrogenase assembly, iron transport, sulfur oxidation and respiration. Likewise, the Gamma1

strains of *A. kojimai* and *A. strummeri* showed adaptive differences in variants related to

hydrogen and sulfur metabolism, respectively. Niche-specific differences in hydrogen metabolic

genes were also observed in comparisons of gene content among symbiont strains. In both *A.*

kojimai and *A. strummeri*, Gamma1 strains from Tui Malila lacked some subunits of uptake or

hydrogen-sensing hydrogenases as well as various genes for hydrogenase maturation, synthesis

and regulation. Given that H_2 concentrations at Tui Malila can drop to 35 µM in endmember

fluids and are probably lower in diffuse flow habitats (Beinart et al. 2012), it is possible that

hydrogen does not constitute a major energy source for Gamma1 strains from this locality. This

hypothesis is in agreement with previous physiological experiments that revealed strikingly low

hydrogen oxidation and associated carbon fixation rates in Gamma1 symbionts from Tui Malila

(Breusing et al. 2020b). An alternative though mutually non-exclusive explanation for the loss of

hydrogenase-related genes at least in the *A. strummeri* strain from Tui Malila could be avoidance

of intra-host competition with co-occurring GammaLau strains, which often co-dominate in *A.*

strummeri individuals at this vent site (Beinart et al. 2012). Such functional diversity is predicted

to enable symbiont coexistence in a variety of hydrothermal vent symbioses, including

bathymodiolin mussels (Ikuta et al. 2016; Ansorge et al. 2019), alvinocaridid shrimp (Cambon-

Bonavita et al. 2021) and vestimentiferan tubeworms (Zimmermann et al. 2014).

Compared to their symbionts, host populations were markedly less structured across the

same spatial scales. Although a proportion of genetic markers was differentiated across vent

sites, we did not find strong evidence for local adaptation in any host species, suggesting that

these patterns likely reflect random variation among localities. The limited genetic subdivision

between host populations agrees with predictions from biophysical models that indicate absence

of physical dispersal barriers for vent larvae within the Lau Back-Arc Basin (Mitarai et al. 2016;

Breusing et al. in review). While symbiont population structure, by contrast, appeared to be at

least partly driven by natural selection, it is possible that symbionts also experience stronger

dispersal limitations than their hosts, as has been hypothesized in some coral-algae symbioses

(Davies et al. 2020). The environmental distributions and life cycles of the free-living stages of

Alviniconcha and *Ifremeria* symbionts are currently unknown and a better understanding of these

aspects will be necessary to evaluate the relative importance of dispersal barriers on symbiont

biogeography.

Overall, our findings reveal a lack of strain-level specificity in *Alviniconcha* and

Ifremeria symbioses, which possibly reflects an evolutionary strategy to cope with the transient

and dynamic nature of hydrothermal vent habitats. Strain flexibility in these associations likely

contributes to the wide-spread genetic connectivity observed among host populations, which in

turn could favor ecological resilience to natural but also anthropogenic environmental

disturbances. Considering the impact of symbiont availability and specificity on host distribution

in a variety of vent symbioses, these results could be important for future conservation

management given the increasing human pressures on hydrothermal ecosystems worldwide

(Levin et al. 2020).

Acknowledgements

We thank the captains, crews and ROV pilots of the R/V *Thomas G. Thompson* (ROV *Jason II*)

and R/V *Falkor* (ROV *Ropos*) for supporting the sample collections that made this study

possible. We thank Pete Girguis for his contribution of the 2009 samples to this project,

Alexander Diehl and Jeff Seewald for providing geochemical data, Michelle Hauer and Erin

Frates for their assistance with sample preparation and Brown University for providing access to

their high-performance computing cluster where the bioinformatic analyses were performed. We

further thank the technical staff at the UC Davis Genome Center for sequencing our Illumina

metagenomic libraries. This work was supported by the National Science Foundation (grant

numbers OCE-1536331, 1819530 and 1736932 to R.A.B.) and the National Institute of Health

(grant numbers 5K99GM135583-02 to S.L.R. and 5R35GM128932-03 to R.C.D.).

Data Accessibility

Raw Nanopore reads, host transcriptomes and symbiont genomes have been deposited in the

National Center for Biotechnology Information under BioProjects PRJNA523619,

PRJNA526236 and PRJNA741492. Annotations for new symbiont genomes are available on the

RAST webserver (https://rast.nmpdr.org/rast.cgi) through the guest access (login: guest,

password: guest) under job IDs 6666666.685764 (Epsilon), 6666666.685765 (Ifr-SOX),

6666666.686795 (Gamma1), and 6666666.687466 (Ifr-MOX). Scripts for bioinformatic analyses

are available on GitHub under https://github.com/cbreusing/Provannid_host-symbiont_popgen.

Author's Contributions

C.B. and R.A.B. designed the study with advice from S.L.R. and R.C.D. C.B. performed data

analyses and wrote the first draft of the paper. M.G. prepared Illumina sequencing libraries. All

authors contributed to data interpretation and writing of the manuscript.

Competing Interests

The authors declare no competing interests.

References

Allito BB, Ewusi-Mensah N, Logah V. Legume-*Rhizobium* strain specificity enhances nutrition
and nitrogen fixation in faba bean (*Vicia faba* L.). Agronomy. 2020; 10(6): 826.
Andrews M, Andrews ME. Specificity in Legume-Rhizobia Symbioses. Int J Mol Sci. 2017;
18(4): 705.
Ansorge R, Romano S, Sayavedra L, Porras MÁG, Kupczok A, Tegetmeyer HE *et al.* Functional
diversity enables multiple symbiont strains to coexist in deep-sea mussels. Nat Microbiol.
2019; 4: 2487–2497.
Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M *et al.* SPAdes: a new genome
assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;
19(5): 455–477.
Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S *et al.* Integrating
taxonomic, functional, and strain-level profiling of diverse microbial communities with
bioBakery 3. Elife. 2021; 10: e65088.
Beinart RA, Sanders JG, Faure B, Sylva SP, Lee RW, Becker EL *et al.* Evidence for the role of
endosymbionts in regional-scale habitat partitioning by hydrothermal vent symbioses.
PNAS 2012; 109: E3241–3250.
Beinart RA, Gartman A, Sanders JG, Luther GW III, Girguis PR. The uptake and excretion of
partially oxidized sulfur expands the repertoire of the energy resources metabolized by
hydrothermal vent symbioses. Proc Biol Sci. 2015; 282: 20142811.
Beinart RA, Luo C, Konstantinidis K, Stewart FJ, Girguis PR. The bacterial symbionts of closely
related hydrothermal vent snails with distinct geochemical habitats show broad similarity
in chemoautotrophic gene content. Front Microbiol. 2019; 10: 1818.
Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting FST: the impact
of rare variants. Genome Res. 2013; 23(9): 1514–1521.
Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 2014; 30: 2114–2120.

Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ *et al.* RASTtk: a modular and
extensible implementation of the RAST algorithm for building custom annotation pipelines
and annotating batches of genomes. Sci Rep. 2015; 5: 8365.
Breusing C, Johnson SB, Tunnicliffe V, Clague DA, Vrijenhoek RC, Beinart RA. Allopatric and
sympatric drivers of speciation in *Alviniconcha* hydrothermal vent snails. Mol Biol Evol.
2020a; 37(12); 3469–3484.
Breusing C, Mitchell J, Delaney J, Sylva SP, Seewald JS, Girguis PR *et al.* Physiological
dynamics of chemosynthetic symbionts in hydrothermal vent snails. ISME J. 2020b; 14:
2568–2579.
Bright M, Bulgheresi S. A complex journey: transmission of microbial symbionts. Nat Rev
Microbiol. 2010 Mar; 8(3): 218–230.
Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ *et al.* Unusual biology across a group
comprising more than 15% of domain Bacteria. Nature 2015; 523: 208–211.
Cailliez F. The analytical solution of the additive constant problem. Psychometrika 1983; 48:
305–308.
Cambon-Bonavita MA, Aubé J, Cueff-Gauchard V, Reveillaud J. Niche partitioning in the
Rimicaris exoculata holobiont: the case of the first symbiotic Zetaproteobacteria.
Microbiome 2021; 9: 87.
Chen LX, Anantharaman K, Shaiber A, Eren AM, Banfield JF. Accurate and complete genomes
from metagenomes. Genome Res. 2020; 30(3): 315–333.
Chomicki G, Kiers ET, Renner SS. The evolution of mutualistic dependence. Annu Rev Ecol
Evol Syst. 2020; 51: 409–432.
Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L *et al.* A program for annotating
and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the
genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin) 2012; 6(2):
80–92.
Dahl C, Franz B, Hensen D, Kesselheim A, Zigann R. Sulfite oxidation in the purple sulfur
bacterium Allochromatium vinosum: identification of SoeABC as a major player and
relevance of SoxYZ in the process. Microbiology (Reading). 2013; 159(Pt 12): 2626–2638.
Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA *et al.* The variant call
format and VCFtools. Bioinformatics. 2011; 27(15): 2156–2158.
Davies SW, Moreland KN, Wham DC, Kanke MR, Matz MV. *Cladocopium* community
divergence in two *Acropora* coral hosts across multiple spatial scales. Mol Ecol. 2020; 29:
4559–4572.
Dick GJ, Anantharaman K, Baker BJ, Li M, Reed DC, Sheik CS. The microbiology of deep-sea
hydrothermal vent plumes: ecological and biogeographic linkages to seafloor and water
column habitats. Front Microbiol. 2013; 4: 124.
Dick GJ. The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped locally.
Nat Rev Microbiol. 2019; 17(5): 271–283.
Dubilier N, Bergin C, Lott C. Symbiotic diversity in marine animals: the art of harnessing
chemosynthesis. Nat Rev Microbiol. 2008; 6: 725–740.
Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA. The evolution of host-symbiont
dependence. Nat Commun. 2017; 8: 15973.
Flores GE, Shakya M, Meneghin J, Yang ZK, Seewald JS, Wheat GC *et al.* Inter-field variability
in the microbial communities of hydrothermal vent deposits from a back-arc basin.
Geobiology. 2012; 10(4): 333–346.

Forester BR, Lasky JR, Wagner HH, Urban DL. Comparing methods for detecting multilocus
adaptation with multivariate genotype–environment associations. Mol Ecol. 2018; 27:
2215–2233.
Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing. arXiv
preprint 2012; arXiv: 1207.3907v2 [q-bio.GN].
Ginete DR, Goodrich-Blair H. From binary model systems to the human microbiome: factors
that drive strain specificity in host-symbiont associations. Front Ecol Evol. 2021; 9: 38.
Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I *et al.* Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.
2011; 29: 644–652.
Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016; 32(18): 2847–2849.
Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 2013; 29: 1072–1075.
Hijmans RJ. geosphere: Spherical Trigonometry. R package version 1.5-10 2019.
https://CRAN.R-project.org/package=geosphere
Ikuta T, Takaki Y, Nagai Y, Shimamura S, Tsuda M, Kawagucci S *et al.* Heterogeneous
composition of key metabolic gene clusters in a vent mussel symbiont population. ISME J
2016; 10: 990–1001.
Jackrel SL, Yang JW, Schmidt KC, Denef VJ. Host specificity of microbiome assembly and its
fitness effects in phytoplankton. ISME J. 2021; 15(3): 774–788.
Kang DD, Li F, Kirton E, Thomas A, Egan R, An H *et al*. MetaBAT 2: an adaptive binning
algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ. 2019; 7: e7359.
Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation Sequencing
Data. BMC Bioinformatics. 2014; 15(1): 356.
Laetsch DR, Blaxter ML. BlobTools: Interrogation of genome assemblies [version 1; peer
review: 2 approved with reservations]. F1000Research 2017; 6: 1287.
Lehmann S, Johnston AWB, Curson ARJ, Todd JD, Cook AM. SoeABC, a novel sulfite
dehydrogenase in Roseobacters. In: Muyzer G, Stams AJM (eds). EMBO Workshop on
Microbial Sulfur Metabolism. (Sieca Repro, Delft, 2012) pp 29.
Lenk S, Moraru C, Hahnke S, Arnds J, Richter M, Kube M *et al. Roseobacter* clade bacteria are
abundant in coastal sediments and encode a novel combination of sulfur oxidation genes.
ISME J. 2012; 6(12): 2178–2187.
Levin LA, Amon DJ, Lily H. Challenges to the sustainability of deep-seabed mining. Nat Sustain
2020; 3: 784–794.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G *et al.* The Sequence
alignment/map (SAM) format and SAMtools. Bioinformatics 2009; 25: 2078–2079.
Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. Bioinformatics. 2006; 22(13): 1658–1659.
Luo J, Lyu M, Chen R, Zhang X, Luo H, Yan C. SLR: a scaffolding algorithm based on long
reads and contig classification. BMC Bioinformatics 2019; 20: 539.
Mitarai S, Watanabe H, Nakajima Y, Shchepetkin AF, McWilliams JC. Quantifying dispersal
from hydrothermal vent fields in the western Pacific Ocean. PNAS 2016; 113(11): 2976–
2981.

Mottl MJ, Seewald JS, Wheat CG, Tivey MK, Michael PJ, Proskurowski G *et al.* Chemistry of
hot springs along the Eastern Lau Spreading Center. Geochim Cosmochim Acta. 2011;
75(4): 1013–1038.
Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic
assembler. Genome Res. 2017; 27(5): 824–834.
Nyholm SV, McFall-Ngai MJ. The winnowing: establishing the squid-*Vibrio* symbiosis. Nat Rev
Microbiol. 2004; 2: 632–642.
Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D *et al.* vegan: Community
Ecology Package. R package version 2.5-7 2020. https://CRAN.R-
project.org/package=vegan
Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Bioinformatics. 2019; 35(3): 526–528.
Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality
of microbial genomes recovered from isolates, single cells, and metagenomes. Genome
Res. 2015; 25(7):1043– 1055.
Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods. 2017; 14: 417–419.
Perez M, Juniper SK. Insights into symbiont population structure among three vestimentiferan
tubeworm host species at Eastern Pacific spreading centers. Appl Environ Microbiol. 2016;
82(17): 5197–5205.
Picelli S, Björklund AK, Reinius B, Sagasser S, Winberg G, Sandberg R. Tn5 transposase and
tagmentation procedures for massively scaled sequencing projects. Genome Res. 2014;
24(12): 2033–2040.
Podowski EL, Ma S, Luther GW III, Wardrop D, Fisher CR. Biotic and abiotic factors affecting
distributions of megafauna in diffuse flow on andesite and basalt along the Eastern Lau
Spreading Center, Tonga. Mar Ecol Prog Ser. 2010; 418: 25–45.
Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D *et al.* PLINK: a toolset
for whole-genome association and population-based linkage analysis. Am J Hum Genet.
2007; 81(3): 559–575.
R Core Team. R: A language and environment for statistical computing. (R Foundation for
Statistical Computing, Vienna, Austria, 2020) https://www.R-project.org/.
Reynolds KC, Watanabe H, Strong EE, Sasaki T, Uematsu K, Miyake H *et al.* New molluscan
larval form: brooding and development in a hydrothermal vent gastropod, *Ifremeria*
nautilei (Provannidae). Biol Bull. 2010; 219(1): 7–11.
Rinta-Kanto JM, Bürgmann H, Gifford SM, Sun S, Sharma S, del Valle DA *et al.* Analysis of
sulfur-related transcription by *Roseobacter* communities using a taxon-specific functional
gene microarray. Environ Microbiol. 2011; 13(2): 453–467.
Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of
RNA-seq data. Genome Biol. 2010; 11: R25.
Russell SL. Transmission mode is associated with environment type and taxa across bacteria-
eukaryote symbioses: a systematic review and meta-analysis. FEMS Microbiol Lett. 2019;
366(3): fnz013.
Sanders J, Beinart R, Stewart F, Delong EF, Girguis PR. Metatranscriptomics reveal differences
in in situ energy and nitrogen metabolism among hydrothermal vent snail symbionts. ISME
J. 2013; 7: 1556–1567.

Seah BK, Gruber-Vodicka HR. gbtools: interactive visualization of metagenome bins in R. Front
Microbiol. 2015; 6: 1451.
Seppey M, Manni M, Zdobnov EM. BUSCO: Assessing genome assembly and annotation
completeness. Methods Mol Biol. 2019; 1962: 227–245.
Seston SL, Beinart RA, Sarode N, Shockey AC, Ranjan P, Ganesh S *et al.* Metatranscriptional
response of chemoautotrophic *Ifremeria nautilei* endosymbionts to differing sulfur
regimes. Front Microbiol. 2016; 7: 1074.
Song L, Florea L. Rcorrector: efficient and accurate error correction for Illumina RNA-seq reads.
Gigascience. 2015; 4: 48.
Suzuki Y, Kojima S, Sasaki T, Suzuki M, Utsumi T, Watanabe H *et al.* Host-symbiont
relationships in hydrothermal vent gastropods of the genus *Alviniconcha* from the
Southwest Pacific. Appl Environ Microbiol. 2006; 72: 1388–1393.
Tivey MK. Generation of seafloor hydrothermal vent fluids and associated mineral deposits.
Oceanography 2007; 20: 50–65.
Ücker M, Ansorge R, Sato Y, Sayavedra L, Breusing C, Dubilier N. Deep-sea mussels from a
hybrid zone on the Mid-Atlantic Ridge host genetically indistinguishable symbionts. ISME
J. 2021; doi: 10.1038/s41396-021-00927-9.
Van der Auwera GA, O'Connor BD. Genomics in the Cloud. (O'Reilly Media, Inc., 2020) pp
834.
Verma SC, Miyashiro T. Quorum sensing in the squid-Vibrio symbiosis. Int J Mol Sci. 2013;
14(8): 16386–16401.
Vrijenhoek RC. Genetics and evolution of deep-sea chemosynthetic bacteria and their
invertebrate hosts. In: Kiel S (ed). The Vent and Seep Biota. (Springer, Dordrecht,
Netherlands, 2010) pp 15–49.
Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S *et al.* Pilon: an integrated
tool for comprehensive microbial variant detection and genome assembly improvement.
PLoS ONE 2014; 9(11): e112963.
Walker L, Lagunas B, Gifford ML. Determinants of host range specificity in legume-rhizobia
symbiosis. Front Microbiol. 2020; 11: 585749.
Wang Q, Liu J, Zhu H. Genetic and molecular mechanisms underlying symbiotic specificity in
legume-rhizobium interactions. Front Plant Sci. 2018; 9: 313.
Warén A, Bouchet P. New records, species, genera, and a new family of gastropods from
hydrothermal vents and hydrocarbon seeps. Zool Scr. 1993; 22: 1–90.
Whitlock MC, Lotterhos KE. Reliable detection of loci responsible for local adaptation:
inference of a null model through trimming the distribution of F(ST). Am Nat. 2015;
186(Suppl 1): S24–36.
Wickham H. ggplot2: Elegant Graphics for Data Analysis. (Springer, New York, 2016) pp 260.
Wilm A, Aw PP, Bertrand D, Yeo GH, Ong SH, Wong CH *et al.* LoFreq: a sequence-quality
aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from
high-throughput sequencing datasets. Nucleic Acids Res. 2012; 40(22): 11189–11201.
Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover
genomes from multiple metagenomic datasets, Bioinformatics 2016; 32(4): 605–607.
Xu GC, Xu TJ, Zhu R, Zhang Y, Li SQ, Wang HW *et al.* LR_Gapcloser: a tiling path-based gap
closer that uses long reads to complete genome assembly. Gigascience. 2019; 8(1): giy157.

Zimmermann J, Lott C, Weber M, Ramette A, Bright M, Dubilier N *et al.* Dual symbiosis with
co-occurring sulfur-oxidizing symbionts in vestimentiferan tubeworms from a
Mediterranean hydrothermal vent. Environ Microbiol. 2014; 16(12): 3638–3656.
733

Figure Captions

Fig. 1 Sampling map for *Alviniconcha* and *Ifremeria* species in the Eastern Lau Spreading Center and Tonga Volcanic Arc.

Fig. 2 Principal component plots for *Alviniconcha* and *Ifremeria* host species based on genetic covariance matrices.

Fig. 3 Principal coordinate plots for *Alviniconcha* and *Ifremeria* symbionts based on relative allele counts transformed into Bray-Curtis dissimilarities. Allele counts approximate the relative proportions of different symbiont strains within host individuals.

Fig. 4 Principal coordinate plot based on Bray-Curtis dissimilarities (A) and presence/absence heatmap of differentially preserved genes (B) for the Gamma1 symbiont of *A. kojimai* and *A. strummeri*. Strains of this symbiont are clearly distinct between the two host species, even when these taxa co-occur (see Fig. S3). Segregation of symbiont populations by host species along the first ordination axis suggests that host affinity is a stronger predictor than geography for symbiont composition between species. Dark and light blue colors in the heatmap indicate presence and absence of genes, respectively. Dendrograms show similarities between samples based on their gene content profiles.

Fig. 5 Presence/absence heatmap for differentially preserved genes.

Fig. 6 Redundancy analysis plots of the first two ordination axes for *Alviniconcha* and *Ifremeria* symbionts. Small grey dots in the center of the plot indicate genetic variants, while large colored dots represent intra-host symbiont populations sampled from different vent localities. Vectors show the environmental predictors. Redundancy analyses were conditioned by geography for all symbionts except for Gamma1 and GammaLau. Genotype-environment associations were not significant for GammaLau.

Table 1 Sampling information for *Alviniconcha* and *Ifremeria* species from the Eastern Lau Spreading Center and Tonga Volcanic Arc. Chemical concentrations are mean endmember values for each vent field obtained from Mottl et al. (2011), Beinart et al. (2012), Flores et al. (2012), and unpublished data provided by J. Seewald (Woods Hole Oceanographic Institution) and A. Diehl (MARUM). n.a. = not available.

Vent site	Species	Coordinates	Year	Depth [m]	H_2 [mM]	CO_2 [mM]	CH_4 [mM]	H_2S [mM]	Fe [mM]	Mn [mM]
Tonga Volcanic Arc										
Niua South	*A. boucheti* *I. nautilei*	15°09.79'S 173°34.48'W	2016	1164	0.037	n.a.	0.005	1.911	n.a.	n.a.
Eastern Lau Spreading Center										
Kilo Moana	*A. boucheti*	20°03.23'S 176°08.01'W	2009	2614	0.382	8.227	0.031	5.432	3.062	0.643
Tow Cam	*A. boucheti* *I. nautilei*	20°18.97'S 176°08.19'W	2009	2711–2722	0.150	10.648	0.045	4.795	0.316	0.389
Tahi Moana	*A. strummeri* *I. nautilei*	20°40.94'S 176°11.01'W	2016	2234–2273	0.097	7.191	0.038	3.548	0.286	0.525
ABE	*A. boucheti* *A. kojimai* *I. nautilei*	20°45.79'S 176°11.47'W	2009 2016	2130–2152	0.079	5.980	0.048	3.060	0.245	0.330
Tu'i Malila	*A. kojimai* *A. strummeri* *I. nautilei*	21°59.36'S 176°34.10'W	2009 2016	1860–1887	0.084	12.912	0.037	2.294	0.203	0.406

A *A. boucheti* (1655 SNPs)

B *A. kojimai* (9185 SNPs)

C *A. strummeri* (3442 SNPs)

D *I. nautilei* (1956 SNPs)

A Epsilon (*A. boucheti*; 30 genes)

B Gamma1 (*A. kojimai*; 99 genes)

C Gamma1 (*A. strummeri*; 71 genes)

D GammaLau (*A. strummeri*; 113 genes)

E Ifr-SOX (*I. nautilei*; 100 genes)

E Ifr-MOX (*I. nautilei*; 145 genes)