Title: A Lethal Genetic Incompatibility between Naturally Hybridizing Species in Mitochondrial Complex I

Authors: Benjamin M. Moran[1,2]*, Cheyenne Y. Payne[1,2], Daniel L. Powell[1,2], Erik N. K. Iverson[3], Alex E. Donny[1], Shreya M. Banerjee[1], Quinn K. Langdon[1], Theresa R. Gunn[1], Rebecca A. Rodriguez-Soto[1], Angel Madero[1], John J. Baczenas[1], Korbin M. Kleczko[1], Fang Liu[4], Rowan Matney[4], Kratika Singhal[4], Ryan D. Leib[4], Osvaldo Hernandez-Perez[2], Russell Corbett-Detig[5,6], Judith Frydman[1,7], Casey Gifford[7], Manfred Schartl[8,9], Justin C. Havird[3], Molly Schumer[1,2,10]*

Affiliations:

[1]Department of Biology, Stanford University, Stanford, CA, USA

[2]Centro de Investigaciones Científicas de las Huastecas "Aguazarca", A.C., Calnali, Hidalgo, Mexico

[3]Department of Integrative Biology, University of Texas, Austin, TX, USA

[4]Stanford University Mass Spectrometry Core, Stanford University, Stanford, CA, USA

[5]Genomics Institute, University of California, Santa Cruz, CA, USA

[6]Department of Biomolecular Engineering, University of California, Santa Cruz, CA, USA

[7]Department of Genetics and Pediatrics, Stanford University, Stanford, CA, USA

[8]The Xiphophorus Genetic Stock Center, Texas State University, San Marcos, TX 78666

[9]Developmental Biochemistry, Biozentrum, University of Würzburg, 97070 Würzburg, Germany

[10]Hanna H. Gray Fellow, Howard Hughes Medical Institute, Stanford, CA, USA

*Correspondence: benmoran@stanford.edu, schumer@stanford.edu

Abstract

The evolution of reproductive barriers is the first step in the formation of new species and

can help us understand the diversification of life on Earth. These reproductive barriers often take

the form of "hybrid incompatibilities," where alleles derived from two different species no

longer interact properly in hybrids. Theory predicts that hybrid incompatibilities may be more

likely to arise at rapidly evolving genes and that incompatibilities involving multiple genes

should be common, but there has been sparse empirical data to evaluate these predictions. Here,

we describe a mitonuclear incompatibility involving three genes in physical contact within

respiratory Complex I in naturally hybridizing swordtail fish species. Individuals homozygous

for specific mismatched protein combinations fail to complete embryonic development or die as

juveniles, while those heterozygous for the incompatibility have reduced function of Complex I

and unbalanced representation of parental alleles in the mitochondrial proteome. We find that the

impacts of different genetic interactions on survival are non-additive, highlighting subtle

complexity in the genetic architecture of hybrid incompatibilities. We document the evolutionary

history of the genes involved, showing for the first time that an incompatibility has been

transferred between species via hybridization. This work thus provides the first glimpse into the

genetic architecture, physiological impacts, and evolutionary origin of a complex incompatibility

impacting naturally hybridizing species.

Introduction

Biologists have long been fascinated by the question of how new species are formed and what mechanisms maintain isolation between them. One key factor in the formation and maintenance of new species is the emergence of genetic incompatibilities that reduce viability or fertility in hybrids[1]. As originally described by the Dobzhansky-Muller model of hybrid incompatibility[2,3], the unique sets of mutations that accumulate in diverging species may interact poorly when they are brought together in hybrids, given that they have never been tested against one another by selection. Due to the technical challenges of identifying these interactions[4], only around a dozen genes involved in hybrid incompatibilities have been precisely mapped[5] and exploration of the functional and evolutionary causes of hybrid incompatibilities has been limited to a small number of cases in model organisms[4].

This knowledge gap leaves key predictions about the evolutionary processes that drive the emergence of hybrid incompatibilities untested. For one, theory suggests that incompatibilities should be more common within dense gene networks, both because genes involved in such interactions are expected to be tightly co-evolving and because the number of potentially incompatible genotypes explodes as the complexity of the genetic interaction increases[6,7]. Consistent with this prediction, mutagenesis experiments have highlighted the sensitivity of multi-protein interactions to changes in any of their components[6]. However, genetic interactions are notoriously difficult to detect empirically except in systems with especially powerful genetic tools[8], and this problem is exacerbated with complex genetic interactions[9,10]. Such technical challenges may explain the rarity of incompatibilities involving three or more genes in the empirical literature[6] (but see[8,11–14]).

Another open question is the degree to which the genes that become involved in hybrid

incompatibilities are predictable from their molecular or evolutionary properties. Researchers

have proposed for decades that rapid molecular evolution will increase the rate at which

incompatibilities accumulate between species[4,5,15]. While several incompatibilities identified to

date involve genes showing signatures of positive selection, it is unclear how unusual rates of

protein evolution are in genes involved in hybrid incompatibilities relative to the genomic

background[5,15]. The mitochondrial genome, in particular, has been proposed as a hotspot for the

accumulation of genetic incompatibilities[16,17], due to substitution rates up to 25× higher than the

nuclear genome in many animals[18–20] and the potential for sexually antagonistic selection driven

by its predominantly maternal inheritance[21–23], among other factors[24]. At the same time, nuclear

and mitochondrial gene products must directly interact with each other in key steps of ATP

synthesis, increasing the likelihood of coevolution between these genomes[25,26]. These molecular

and evolutionary factors suggest that interactions between mitochondrial- and nuclear-encoded

proteins could play an outsized role in the emergence of hybrid incompatibilities[16]. Crosses in

numerous species have shown that hybrid viability often depends on the identity of the maternal

species[27–30], providing indirect evidence for the prevalence of mitonuclear incompatibilities.

However, the field has struggled to move beyond these coarse-scale patterns, as few studies have

successfully mapped mitonuclear incompatibilities to the single gene level[31–34] and none of those

identified to date have been studied in species that naturally hybridize.

As we begin to identify the individual genes underlying hybrid incompatibilities, the next

frontier is evaluating the processes that drive their evolution. Over the past two decades, it has

become abundantly clear that hybridization is exceptionally common in species groups where it

was once thought to be rare[35,36]. As such, we now appreciate how frequently species derive genes

from their relatives, and in many cases are a patchwork of ancestry from several genomes due to

historical hybridization[37–39]. The impact of these dynamics on the evolution of hybrid

incompatibilities have been poorly investigated[40] since the foundational theory in this area was

developed before the ubiquity of hybridization was fully appreciated[7].

Here, we use an integrative approach to precisely map the genetic basis and physiological

impacts of a lethal mitonuclear hybrid incompatibility in swordtail fish and uncover its

evolutionary history. The sister species *Xiphophorus birchmanni* and *X. malinche* began

hybridizing in the last ~100 generations in multiple river systems[41] after premating barriers were

disrupted by habitat disturbance[42], and are a powerful system to study the emergence of hybrid

incompatibilities in young species. Despite their recent divergence[43] (~250,000 generations;

0.5% divergence per basepair), some hybrids between *X. birchmanni* and *X. malinche* experience

strong selection against incompatibilities[43,44]. One incompatibility causing melanoma has been

previously mapped in this system and population genetic patterns suggest that dozens may be

segregating in natural hybrid populations[43–46]. Moreover, the ability to generate controlled

crosses[47,48] and the development of high-quality genomic resources[46,49] makes this system

particularly tractable for identifying hybrid incompatibilities and characterizing their evolution in

natural populations. Leveraging data from controlled laboratory crosses and natural hybrid

populations, we pinpoint two nuclear-encoded *X. birchmanni* genes that are lethal when

mismatched with the *X. malinche* mitochondria in hybrids, explore the developmental and

physiological effects of this incompatibility, and trace its evolutionary history.

Admixture Mapping Reveals a Lethal Mitonuclear Incompatibility

To identify loci under selection in *X. birchmanni* × *X. malinche* hybrids, we generated ~1X low-coverage whole-genome sequence data for 943 individuals from an F_2 laboratory cross and 359 wild-caught hybrid adults, and applied a hidden Markov model to data at 629,661 ancestry-informative sites along the genome to infer local ancestry (~1 informative site per kb[45,50]; Methods, Supplementary Information 1.1.1–1.1.4). Using these results, we found evidence for a previously unknown incompatibility between the nuclear genome of *X. birchmanni* and the mitochondrial genome of *X. malinche* (Supplementary Information 1.1.5–1.1.10). Our first direct evidence for this incompatibility came from controlled laboratory crosses (Methods, Supplementary Information 1.1.1). Because the cross is largely unsuccessful in the opposite direction, all lab-bred hybrids were the offspring of F_1 hybrids generated between *X. malinche* females and *X. birchmanni* males and harbored a mitochondrial haplotype derived from the *X. malinche* parent species. Offspring of F_1 intercrosses are expected to derive on average 50% of their genome from each parent species. This expectation is satisfied genome-wide and locally along most chromosomes in F_2 hybrids (on average 50.3% *X. malinche* ancestry; Fig. S1). However, we detected six segregation distorters genome-wide[48], with the most extreme signals falling along a 6.5 Mb block of chromosome 13 and an 4.9 Mb block of chromosome 6 (Fig. 1A; Fig. 1D).

Closer examination of genotypes in the chromosome 13 region showed that almost none of the surviving individuals harbored homozygous *X. birchmanni* ancestry in a 3.75 Mb subregion (Fig. 1C; Fig. S2; 0.1% observed vs 25% expected). This pattern is unexpected even in the case of a lethal incompatibility involving only nuclear loci (where simulations indicate that

we should recover homozygous *X. birchmanni* ancestry in ~10% of surviving individuals;

Supplementary Information 1.1.1) but is consistent with a lethal mitonuclear incompatibility.

Using approximate Bayesian computation (ABC) approaches we inferred what strength of

selection against *X. birchmanni* ancestry in this region was consistent with the genotypes and

ancestry deviations observed. We estimated posterior distributions of selection and dominance

coefficients and inferred that selection on this genotype in F_2s is largely recessive and essentially

lethal (maximum a posteriori estimate $h = 0.12$ and $s = 0.996$, 95% credible interval $h = 0.010$–

0.194 and $s = 0.986$–0.999; Fig. 1B; Fig. S3; Methods; Supplementary Information 1.2.1–1.2.2).

The degree of segregation distortion observed in F_2 individuals on chromosome 6 is also

surprising (Fig. 1D). Only 3% of individuals harbor homozygous *X. birchmanni* ancestry in this

region (compared to 0.1% in the chromosome 13 region and 25% on average at other loci across

the genome; Fig 1F), which is again lower than expected for a nuclear-nuclear hybrid

incompatibility (Supplementary Information 1.1.1). ABC approaches indicate that selection on

homozygous *X. birchmanni* ancestry on chromosome 6 is also severe (maximum a posteriori

estimate $h = 0.09$ and $s = 0.91$, 95% credible interval interval $h = 0.01$–0.21 and $s = 0.87$–0.94;

Fig. 1E, Fig. S3, Supplementary Information 1.2.2). Thus, our F_2 data show that homozygous *X.*

birchmanni ancestry in regions of either chromosome 13 or chromosome 6 is almost completely

lethal in hybrids with *X. malinche* mitochondria (Fig. 1H).

To formally test for the presence of a mitonuclear incompatibility involving chromosome

13 and chromosome 6, or elsewhere in the genome, we leveraged data from natural hybrid

populations. Most naturally occurring *X. birchmanni × malinche* hybrid populations are fixed for

mitochondrial haplotypes from one parental species (Supplementary Information 1.1.2, 1.1.6).

However, a few populations segregate for the mitochondrial genomes of both parental types, and

we focused on one such population (the "Calnali Low" population, hereafter the admixture

mapping population). Admixture mapping for associations between nuclear genotype and

mitochondrial ancestry (after adjusting for expected covariance due to genome-wide ancestry[44])

revealed two genome-wide significant peaks and one peak that approached genome-wide

significance (Fig. 1G, Table S1–S3). The strongest peak of association spanned approximately

77 kb and fell within the region of chromosome 13 identified using F_2 crosses (Fig. 1G). This

peak was also replicated in another hybrid population (Fig. S4; Methods, Supplementary

Information 1.1.5) and contains only three genes: the NADH dehydrogenase ubiquinone iron-

sulfur protein 5 (*ndufs5*), E3 ubiquitin-protein ligase, and microtubule-actin cross-linking factor

1. Of these three genes, only *ndufs5* forms a protein complex with mitochondrially encoded

proteins, which along with other evidence implicates it as one of the interacting partners in the

mitonuclear incompatibility (Fig. 1C; see Supplementary Information 1.1.8 for analysis of other

genes).

Analysis of three natural hybrid populations that had fixed the mitochondrial haplotype of

one of the parental species (Fig. S5) confirmed that this region on chromosome 13 is under

selection in natural hybrid populations, with the strongest signal of selection localizing precisely

to the same three genes found under the admixture mapping peak (Fig. S6A; Supplementary

Information 1.1.6). Moreover, comparing genotypes and phenotypes in siblings allowed us to

exclude maternal effects as a driver of the chromosome 13 signal (Supplementary Information

1.1.7), and we ruled out the possibility that other confounding factors could generate the

observed patterns (Supplementary Information 1.1.9).

We also identified a peak on chromosome 6 that approached genome-wide significance

(Fig. 1G; Table S2; Supplementary Information 1.1.10) and fell precisely within the segregation

distortion region previously mapped in F$_2$ hybrids (Fig. 1D; Supplementary Information 1.1.1).

This peak contained 20 genes including the mitochondrial Complex I gene *ndufa13* (Fig. S7–S8;

Methods, Supplementary Information 1.1.10). Depletion of non-mitochondrial parent ancestry at

ndufa13 was unidirectional (Fig. 1F), consistent with selection acting only against the

combination of the *X. malinche* mitochondria with homozygous *X. birchmanni* ancestry at

ndufa13 (see Supplementary Information 1.2.3–1.2.4). Genomic analyses in natural hybrid

populations confirmed this asymmetry (Fig. S6B).

Together, these results indicate that at least two *X. birchmanni* nuclear genes cause

incompatibility when they are mismatched in ancestry with the *X. malinche* mitochondria (Fig.

1H; Supplementary Information 1.2.5). These genes, *ndufs5* and *ndufa13*, belong to a group of

proteins and assembly factors that form respiratory Complex I[51] (see Table S1 for locations of

the 51 annotated Complex I genes in the *Xiphophorus* genome). Complex I is the first component

of the mitochondrial electron transport chain that ultimately allows the cell to generate ATP.

Both nuclear proteins interface with several mitochondrially derived proteins at the core of the

Complex I structure, hinting at the possibility that physical interactions could underlie this multi-

gene mitonuclear incompatibility.

Interactions with the *X. birchmanni* mitochondria

Admixture mapping analysis also identified a strong peak of mitonuclear association on

chromosome 15, which we briefly discuss here and in Supplementary Information 1.1.10 and

1.2.1. This peak was associated with *X. birchmanni* mitochondrial ancestry (Fig. S9), indicating

that it has a distinct genetic architecture from the incompatibility involving the *X. malinche*

mitochondria and *X. birchmanni ndufs5* and *ndufa13*. Specifically, analysis of genotypes at the

admixture mapping peak indicates that the *X. birchmanni* mitochondria is incompatible with

homozygous *X. malinche* ancestry on chromosome 15 (Fig. 1C, Fig. S9). This region did not

contain any members of Complex I, but dozens of genes in this interval interact with known

mitonuclear genes (see Table S3; Supplementary Information 1.1.10). The fact that we detect

incompatible interactions with both the *X. malinche* mitochondria (at *ndufs5* and *ndufa13*) and

the *X. birchmanni* mitochondria (*ndufs5* and chromosome 15) in our admixture mapping results

supports the idea that mitonuclear interactions can act as "hotspots" for the evolution of hybrid

incompatibilities[16].

Lethal Effect of Incompatibility in Early Development

The combination of *X. birchmanni ndufs5* or *ndufa13* with the *X. malinche* mitochondria

appears to be lethal by the time individuals reach adulthood. To investigate the developmental

timing of the incompatibility, we genotyped pregnant females from the admixture mapping

population and recorded the developmental stages of their embryos[52] (swordtails are livebearing

fish; see Methods). We found a significant interaction between developmental stage and *ndufs5*

genotype, whereas *ndufa13* genotype did not affect developmental stage (Fig. S10–13;

Supplementary Information 1.3.1). While developmental asynchrony is typically on the scale of

0–2 days in pure species[53] (Supplementary Information 1.3.1), we observed much greater

variation in broods collected from the admixture mapping population where the mitochondrial

incompatibility is segregating (e.g. stages normally separated by roughly 12 days of development

found in the same brood; Supplementary Information 1.3.1; Fig. 2A–B). Genotyping results

revealed that embryos with homozygous *X. birchmanni* ancestry at *ndufs5* and *X. malinche*

mitochondria are present at early developmental stages, but that these embryos failed to develop

beyond a phenotype typical of the first seven days of gestation (the full length of gestation is 21–

28 days in *Xiphophorus*; Fig. 2A–B, Fig. 2D–E). Individuals with mismatched ancestry at *ndufs5*

whose siblings were fully developed still had a detectable heartbeat but had consumed less yolk

than their siblings and remained morphologically underdeveloped (Fig. 2D, Fig. S14–19). Unlike

other species, in *Xiphophorus* this developmental lag could itself cause mortality, since embryos

that do not complete embryonic development inside the mother fail to survive more than a few

days after birth (Supplementary Information 1.3.1; Table S4). Given that Complex I inhibition

lethally arrests development in zebrafish embryos[54,55], we also tested the effects of Complex I

inhibition on *X. birchmanni* and *X. malinche* fry, and found a similar level of sensitivity

(Supplementary Information 1.3.2).

In contrast to individuals with mismatched ancestry at *ndufs5,* those with *ndufa13*

mismatch survived embryonic development but suffered mortality in the early post-natal period

(Fig. 2C). We tracked 74 F_2 fry from 24 hours post-birth through adulthood (Supplementary

Information 1.3.3). We found that most fry with incompatible genotypes at *ndufa13* had already

suffered mortality by the time tracking began, with only 7 individuals found 24 hours post-birth

that were homozygous *X. birchmanni* at *ndufa13* (versus 19 expected; binomial $P = 0.0005$). No

natural mortality was observed between 1 day and 3 months post-birth (one individual exhibiting

distress was euthanized, and was a *ndufa13* incompatible homozygote; see Supplementary

Information 1.3.3).

Evidence for Complex Physiological and Fitness Effects in Incompatible Individuals

Our analysis of developing embryos indicates that individuals with the *ndufs5* incompatibility exhibit delayed or arrested embryonic development, whereas those with the *ndufa13* incompatibility did not. This suggests that these genes may drive lethality through partially distinct mechanisms. As such, we chose to further investigate the effects of *ndufs5* and *ndufa13* and their possible interactions using a paired genotyping and physiological approach throughout development. We sampled 235 F_2 embryos at a range of developmental stages and measured their overall rates of respiration (Supplementary Information 1.3.4–1.3.5). We also used imaging of these embryos to track cardiovascular phenotypes as these have been associated with *ndufa13* defects in mammals[56]. We found that incompatible genotypes at *ndufs5* and *ndufa13* impacted a range of phenotypes, including heart rate, length relative to compatible siblings, and length-corrected head size (Fig. S14–22, Table S5–S7). *ndufa13* mismatch has a large effect on cardiovascular phenotypes, including heart rate and the size of the sinu-atrium (an embryo-specific heart chamber, Fig. 2G–H, Fig. S22, Table S6, S8); *ndufs5* affects only heart rate (Fig. S19, Table S6). We find initial evidence that cardiac defects persist into adulthood in surviving individuals with *ndufa13* mismatch (see Supplementary Information 1.3.3; Fig. S23). By contrast, *ndufs5* mismatch has a major impact on rates of respiration and yolk consumption during development (Fig. 2F, while *ndufa13* only affects respiration through an interaction with *ndufs5* and body length, Fig. S24–27, Table S9–11).

Naively, the separable impacts of incompatible genotypes at *ndufs5* and *ndufa13* could indicate that even though these proteins are in physical contact in Complex I (see below), they represent two distinct hybrid incompatibilities. We investigated this question by taking

advantage of rare survivors of the *ndufa13* incompatibility in an expanded dataset of 1010 F_2

hybrids. Using this dataset, we were able to identify dozens of survivors of the *ndufa13*

incompatibility (3.4% of individuals) and found that genotypes at *ndufa13* and *ndufs5* were not

independent (χ^2 association test $P = 0.032$, Supplementary Information 1.2.5). Upon further

investigation, we found that the majority of survivors of the *ndufa13* incompatibility had

homozygous *X. malinche* ancestry at *ndufs5*, suggesting that harboring even one *X. birchmanni*

allele at *ndufs5* may sensitize fry to the *ndufa13* incompatibility. Indeed, we found that

individuals that had heterozygous ancestry at *ndufs5* were significantly under-represented among

surviving *ndufa13* incompatible individuals (Permutation test $P = 0.015$, Fig. 2I, Supplementary

Information 1.2.5). These findings highlight a subtle but significant non-additive effect of *ndufs5*

and *ndufa13* on survival.

Mitochondrial Biology in Adult Hybrids Heterozygous for the Incompatibility

Because few individuals homozygous for incompatible genotypes at *ndufs5* or *ndufa13*

survive past birth, our previous experiments focused on embryos. However, the small size of

Xiphophorus embryos prevents us from using assays that directly target Complex I. To further

explore the effects of the hybrid incompatibility on Complex I function *in vivo*, we turned to

adult F_1 hybrids between *X. birchmanni* and *X. malinche* (Fig. 3A). Since F_1 hybrids that derive

their mitochondria from *X. malinche* and are heterozygous for ancestry at *ndufs5* and *ndufa13* are

fully viable, we asked whether there was evidence for compensatory nuclear or mitochondrial

regulation that might be protective in F_1 hybrids. We found no evidence for significant

differences in expression of *ndufs5* or *ndufa13* (Supplementary Information 1.3.6; Fig. S28–29)

or in mitochondrial copy number (Supplementary Information 1.3.7; Fig. 3B) between F_1 hybrids

and parental species.

With no indication of a compensatory regulatory response, we reasoned that we might be

able to detect reduced mitochondrial Complex I function in hybrids heterozygous for ancestry at

ndufs5 and *ndufa13.* We quantified respiratory phenotypes in isolated mitochondria using an

Oroboros O2K respirometer in adult hybrids and both parental species (Fig. S30; Methods,

Supplementary Information 1.3.8). We found that Complex I efficiency was lower in hybrids

compared to the two parental species (Fig. 3C, Fig. S31, orthogonal contrast $t = -2.53$, $P = 0.023$,

n = 7 per genotype), and that the time required for hybrids to reach maximum Complex I-driven

respiration was ~2.5 times longer (orthogonal contrast $t = 4.303$, $P < 0.001$; Fig. 3D; Fig. S32).

On the other hand, overall levels of mitochondrial respiration were unaffected by genotype (Fig.

3E, orthogonal contrast $t = 0.078$, $P = 0.94$, n = 7 per genotype; Supplementary Information

1.3.8) as were multiple independent measures of mitochondrial integrity and function (Fig. S31–

34, Supplementary Information 1.3.8–1.3.9). Together, these data point to reduced function of

Complex I without broader phenotypic consequences in individuals heterozygous for

incompatible alleles, as previously observed in other models of Complex I dysfunction[57,58].

Given physiological evidence for some reduction in Complex I function in hybrids

heterozygous at *ndufs5* and *ndufa13*, we predicted that there might be an altered frequency of

protein complexes incorporating both *X. malinche* mitochondrial proteins and *X. birchmanni*

proteins at *ndufs5* and *ndufa13* in F_1 hybrids. To test this prediction, we took a mass

spectrometry-based quantitative proteomics approach. We used stable isotope-labeled peptides to

distinguish between the *X. birchmanni* and *X. malinche ndufs5* and *ndufa13* peptides in

mitochondrial proteomes extracted from F_1 hybrids (n = 5, see Methods, Supplementary

Information 1.4.1–1.4.4). While endogenous *ndufa13* peptides were not observed frequently

enough to quantify accurately, we found consistent deviations from the expected 50-50 ratio of

X. birchmanni to *X. malinche* peptides for *ndufs5* in F_1 hybrids, with a significant

overrepresentation of matched ancestry at *ndufs5* in the mitochondrial proteome ($t = 3.96$, $P =$

0.016; Fig. 3F; Fig. S35; Supplementary Information 1.4.5). Since we did not observe allele-

specific expression of *ndufs5* (Fig 3F; Supplementary Information 1.3.6), this result is consistent

with disproportionate degradation of *X. birchmanni*-derived *ndufs5* peptides in the mitochondrial

proteome or differences in translation of *ndufs5* transcripts derived from the two species.

Paired mitonuclear substitutions at contact sites in Complex I

While we can leverage high resolution admixture mapping to pinpoint the nuclear

components of the hybrid incompatibility, we cannot take this approach to distinguish among the

37 genes in the swordtail mitochondrial genome because they do not undergo meiotic

recombination. To begin to explore the possible mitochondrial partners of *ndufs5* and *ndufa13*,

we therefore turned to protein modeling, relying on high quality cryo-EM based structures[59–61].

Although these structures are only available for distant relatives of swordtails, the presence of

the same set of supernumerary Complex I subunits and high sequence similarity suggest that

using these structures is appropriate (Table S12–S13; Fig. S36–S38; Supplementary Information

1.4.6).

Barring a hybrid incompatibility generated by regulatory divergence (see Supplementary

Information 1.3.6), our expectation is that hybrid incompatibilities will be driven by amino acid

changes in interacting proteins[62]. We used the program RaptorX[63] to generate predicted

structures of *X. birchmanni* and *X. malinche ndufs5, ndufa13*, and nearby Complex I

mitochondrial and nuclear genes, which we aligned to a mouse cryo-EM Complex I structure[59]

(Fig. 4A; Fig. S36–S38; Methods). Using these structures, we visualized amino acid substitutions

between *X. birchmanni* and *X. malinche* at the interfaces of *ndufs5, ndufa13* and mitochondrially

encoded genes (Fig. S39–S41). While there are dozens of substitutions in the four

mitochondrially encoded genes that are in close physical proximity to *ndufs5* or *ndufa13* (Fig.

S36; *nd2, nd3, nd4l,* and *nd6*), there are only five cases where amino acid substitutions in either

nuclear gene are predicted to be close enough to contact substitutions in any mitochondrial gene,

all of which involve *nd2* or *nd6* (Table S14, Fig. 4A; see Fig. S40 for pairwise visualizations of

interacting proteins). These paired substitutions in regions of close proximity between

mitochondrial- and nuclear-encoded proteins suggest that *nd2* and *nd6* are the genes most likely

to be involved in the mitochondrial component of the hybrid incompatibility (Fig. 4A–B; Fig.

S41–S44). While there are currently no developed approaches for transgenics in livebearing fish

like *Xiphophorus* (Supplementary Information 1.3.1) and Complex I is ill-suited for *in vitro*

approaches (being comprised of 45 subunits with an intricate assembly process[64]), manipulating

these in-contact substitutions is an exciting direction for future work.

Rapid evolution of Complex I proteins

Theory predicts that hybrid incompatibilities are more likely to arise in rapidly evolving

genes[4,5,7,15]. Consistent with this hypothesis, *ndufs5* is among the most rapidly evolving genes

genome-wide between *X. birchmanni* and *X. malinche* (Fig. 4C–D). Aligning the *ndufs5* coding

sequences of *X. birchmanni, X. malinche*, and twelve other swordtail species revealed that all

four amino acid substitutions that differentiate *X. birchmanni* and *X. malinche* at *ndufs5* were

derived on the *X. birchmanni* branch (Fig. 4C). Phylogenetic tests indicate that there has been

accelerated evolution of *ndufs5* on this branch ($d_N/d_S > 99$, N = 4, S = 0, codeml branch test $P =$

0.005, Fig. 4C). Similar patterns of rapid evolution are observed at *ndufa13*, which also showed

evidence for accelerated evolution in *X. birchmanni* (Fig. 4E; $d_N/d_S = 1.2$, N = 3, S = 1, codeml

branch test $P = 0.002$). While explicit tests for adaptive evolution at *ndufs5* and *ndufa13* could

not exclude a scenario of relaxed selection (Supplementary Information 1.5.1, 1.5.2), our

comparisons across phylogenetic scales highlight strong conservation in some regions of the

proteins and rapid turnover in others, complicating our interpretation of this test (Fig. S45).

Rapid evolution of *ndufs5* and *ndufa13* could be driven by coevolution with

mitochondrial substitutions, a mechanism that has been proposed to explain the outsized role of

the mitochondria in hybrid incompatibilities[16,65]. Indeed, there is an excess of derived

substitutions in the *X. birchmanni* mitochondrial protein *nd6*, one of the proteins that physically

contacts *ndufs5* and *ndufa13* (Table S15; Fig. S42; codeml branch test $P = 0.005$). Moreover,

several of the substitutions observed in both mitochondrial and nuclear genes are predicted to

have functional consequences (Supplementary Information 1.5.1; Table S16), including ones

predicted to be in contact between *ndufs5, ndufa13, nd2*, and *nd6* (Fig. 4A, 4B; Fig. S41).

Introgression of genes underlying mitonuclear incompatibility

The presence of a mitonuclear incompatibility in *Xiphophorus* is especially intriguing,

given previous reports that mitochondrial genomes may have introgressed between species[37].

While *X. malinche* and *X. birchmanni* are sister species based on the nuclear genome, they are

mitochondrially divergent, with *X. malinche* and *X. cortezi* grouped as sister species based on the

mitochondrial phylogeny[37] (Fig. 5A–B). As we show, all *X. cortezi* mitochondria sequenced to

date are nested within *X. malinche* mitochondrial diversity (Fig. 5B, Fig. S42, S46;

Supplementary Information 1.5.3–1.5.4). Simulations indicate that gene flow, rather than

incomplete lineage sorting, drove replacement of the *X. cortezi* mitochondria with the *X.*

malinche sequence (P < 0.002 by simulation; Fig. 5C; Supplementary Information 1.5.4).

The introgression of the mitochondrial genome from *X. malinche* into *X. cortezi* raises the

possibility that other Complex I genes may have co-introgressed[66]. Indeed, the nucleotide

sequence for *ndufs5* is identical between *X. malinche* and *X. cortezi*, and the sequence of *ndufa13*

differs by a single synonymous mutation (although conservation of both genes is high throughout

Xiphophorus; Fig. S47–S48). These identical amino acid sequences suggest that hybrids between

X. cortezi and *X. birchmanni* are likely to harbor the same mitonuclear incompatibility we

observe between *X. malinche* and *X. birchmanni*, as a result of ancient introgression between *X.*

malinche and *X. cortezi* (Fig. 5D; Supplementary Information 1.5.3–1.5.5).

This inference is supported by analysis of ancestry in three contemporary *X. birchmanni*

× *X. cortezi* hybrid populations[48], two of which are completely geographically isolated,

indicating that the hybridization events were independent (Fig. S49). We find that all known *X.*

birchmanni × *X. cortezi* hybrid populations are fixed for the mitochondrial genome from *X.*

cortezi (that originated in *X. malinche*) and also show a striking depletion of non-mitochondrial

parent ancestry at *ndufs5* and *ndufa13* (Fig. 5E; Fig. S50; Supplementary Information 1.5.6).

This repeated depletion is not expected by chance (Fig. 5E; Supplementary Information 1.5.6, *P*

= 0.0001 by simulation) and instead indicates that selection has removed mismatched ancestry in

these regions of the genome. These results suggest that the mitonuclear incompatibility observed

in *X. birchmanni* × *X. malinche* is also active in hybridizing *X. birchmanni* × *X. cortezi*

populations. This exciting finding shows for the first time that genes underlying hybrid

incompatibilities can introgress together, transferring incompatibilities between related species.

Discussion

What genetic and evolutionary forces drive the emergence of hybrid incompatibilities, especially between closely related species? Theory predicts that hybrid incompatibilities involving multiple genes should be common[6,7], but with few exceptions[8,11–13], they remain virtually uncharacterized at the genic level[6]. Here, we identify incompatible interactions in mitochondrial Complex I that causes hybrid lethality in lab and wild populations. Our findings in naturally hybridizing species echoes predictions from theory and studies in lab models[8,11–13] that protein complexes may be a critical site of hybrid breakdown.

Researchers have proposed mitonuclear interactions as "hotspots" for the emergence of hybrid incompatibilities, given that mitochondrial genomes often experience higher substitution rates between species[18–20], yet must intimately interact with nuclear proteins to perform essential cellular functions[25,26]. Our findings support this prediction, identifying incompatible interactions with both the *X. malinche* and *X. birchmanni* mitochondria. We also show that there has been exceptionally rapid evolution in both mitochondrial and interacting nuclear genes in *X. birchmanni*, which we speculate may have introduced mutations that are incompatible in hybrids (Fig. 4). Whether driven by adaptation or some other mechanism, our findings support the hypothesis that the coevolution of mitochondrial and nuclear genes could drive the overrepresentation of mitonuclear interactions in hybrid incompatibilities[25,26,65]. More broadly, our results are consistent with predictions that rapidly evolving proteins are more likely to become involved in hybrid incompatibilities than their slowly evolving counterparts[4,5,15].

Characterizing the incompatibility across multiple scales of organization allowed us to explore the mechanisms through which it acts[67–69]. Our results suggest that in the case of the *X. malinche* mitochondria hybrid lethality is mediated through arrested development *in utero* of

individuals with mismatched ancestry at *ndufs5*, while individuals with *ndufa13* mismatch have

vascular defects and typically die shortly after birth. Intriguingly, individuals with *ndufa13*

mismatch that do survive are much less likely to harbor any *X. birchmanni* alleles at *ndufs5*

(even in the heterozygous state; Fig. 2I). Taken together, our results indicate that a subtle three-

way interaction overlays two strong pairwise mitonuclear incompatibilities at *ndufs5* and

ndufa13. Evolutionary biologists have been fascinated by the idea that hybrid incompatibilities

may commonly involve three or more genes following theoretical work by Orr[7] almost thirty

years ago, but this question has been challenging to address empirically. Our results highlight

how the nuances of actual fitness landscapes may defy simplifying assumptions.

Finally, this mitonuclear incompatibility provides a new case in which the same genes are

involved in incompatibilities across multiple species[38,46,70]. However, tracing the evolutionary

history of the genes that underlie it adds further complexity to this phenomenon: we found that

introgression has resulted in the transfer of genes underlying the incompatibility from *X.*

malinche to *X. cortezi*, and evidence from *X. birchmanni* × *X. cortezi* hybrid populations

indicates that the incompatibility is likely under selection in these populations as well. The

possibility that hybridization could transfer incompatibilities between species has not been

previously recognized, perhaps due to an underappreciation of the frequency of hybridization.

The impact of past hybridization on the structure of present-day reproductive barriers between

species is an exciting area for future inquiry.

Fig. 1 | Admixture mapping pinpoints mitonuclear incompatibility in *Xiphophorus*. (A) Average

ancestry of F₂ hybrids on chromosome 13 reveals segregation distortion towards *X. malinche* ancestry

across a large region of the chromosome. Blue envelope shows the 99% quantiles of *X. malinche* ancestry

at all ancestry informative sites genome wide. Dashed line represents the expected *X. malinche* ancestry

for this cross. Purple arrow points to the position of *ndufs5*. **(B)** Results of approximate Bayesian

computation (ABC) simulations estimating the strength of selection on the *X. malinche* mitochondria

when combined with *X. birchmanni* ancestry at *ndufs5*. Shown here is the posterior distribution from

accepted simulations; vertical line and inset indicates the maximum a posteriori estimate for the selection

coefficient. (**C**) Observed genotype frequencies of different genotype combinations of *ndufs5*

(chromosome 13) and mitochondrial haplotypes in the admixture mapping population. (**D**) Average

ancestry of F_2s on chromosome 6, reveals segregation distortion towards *X. malinche* ancestry across a

large region of the chromosome. Blue envelope and dashed line indicate 99% ancestry quantiles and

expected ancestry in the cross as in (**A**), red arrow points to the position of *ndufa13*. (**E**) Results of

approximate Bayesian computation (ABC) simulations estimating the strength of selection on the

combination of *X. malinche* mitochondria with *X. birchmanni ndufa13*, as in (**B**). (**F**) Observed genotype

frequencies of different genotype combinations of *ndufa13* (chromosome 6) and mitochondrial haplotypes

in the admixture mapping population. (**G**) Admixture mapping results for associations between nuclear

ancestry and mitochondrial haplotype in natural hybrids, controlling for genome-wide ancestry. Blue line

indicates the 10% false positive rate genome-wide significance threshold determined by simulations. The

peak visible on chromosome 15 is driven by interactions with the *X. birchmanni* mitochondria and an

unknown nuclear gene and is discussed in Supplementary Information 1.1.10 and 1.2.1. (**H**) Schematic of

identified interactions with the *X. malinche* mitochondrial genome from our mapping data. We detect a

subtle interaction between *ndufs5* and *ndufa13* (indicated by the dashed line; see main text, Figure 2, and

Supplementary Information 1.2.5).

Fig. 2 | Impact of the hybrid incompatibility on *Xiphophorus* hybrid embryos. (**A**) Developmental

stage and *ndufs5* genotypes of hybrid embryos with *X. malinche* mitochondria. (**B**) Lag in development of

hybrid embryos with *X. malinche* mitochondria compared to the most developed embryo in their brood, as

a function of *ndufs5* genotype. (**C**) Change in frequency of homozygous *X. birchmanni* ancestry at

ndufa13 in F_2 hybrids over development. Points and lines represent the estimate ± 2 standard errors, based

on observed frequency and sample size. (**D**) F_2 hybrid siblings showing stark difference in developmental

progression as a function of *ndufs5* genotype. Top photo shows entire length of heterozygote embryo,

while bottom photos show lagging *ndufs5*-incompatible sibling at matched scale (left) and zoomed for

detail (right). (**E**) Frequency of homozygous *X. birchmanni* ancestry along chromosome 13 in embryos

with *X. malinche* mitochondria that lagged their siblings in developmental stage by ≥1 developmental

stage (red) versus frequency of homozygous *X. birchmanni* ancestry in embryos that did not exhibit

developmental lag (blue, see Supplementary Information 1.1.7). Dashed line indicates the location of

ndufs5. Note that only 66% of embryos with developmental lag have homozygous *X. birchmanni* in this

region, indicating that there are other causes of this phenotype. For the same analysis of chromosomes 6

and 15, where we see no clear difference in average ancestry as a function of lag status, see Fig. S11–13.

(**F–H**) Respiratory and morphometric measurements in live F_2 embryos. To control for the effect of

length on other variables, residuals of each variable regressed against body length are plotted (see

Supplementary Information 1.3.4-1.3.5). Gray points denote individual measurements, colored points and

bars show mean ± 2 SE for each genotype, and brackets with asterisks denote significant differences

based on a Tukey's HSD test. (**F**) Relationship between *ndufs5* genotype and respiration in F_2 hybrid

embryos, as measured by Loligo plate respirometer. (**G**) Relationship between *ndufa13* genotype and

heart rate in F_2 hybrid embryos. (**H**) Relationship between *ndufa13* genotype and the width of the sinu-

atrium, which peristaltically pumps blood from the yolk into the embryonic heart atrium. (**I**) Frequency of

individuals heterozygous for ancestry at *ndufs5* (i.e. those with one *X. birchmanni* allele) among juveniles

and adults of varying *ndufa13* genotypes. Points and lines represent observed frequency ± 2 standard

errors. Dashed line represents expected frequency of *ndufs5* heterozygotes under a scenario of additive

selection on *ndufs5* and *ndufa13* (see Supplementary Information 1.2.5).

512
Fig. 3 | Physiological and proteomic phenotypes of viable heterozygotes harboring the hybrid

incompatibility. In all panels, colored points and whiskers show the mean ± 2 standard errors, and gray

points show individual data. (**A**) Schematic of ancestry at loci involved in the incompatibility in an *X.*

birchmanni × malinche F$_1$ hybrid. Hybrids heterozygous at both loci avoid the lethal effects of the

interaction between the *X. malinche* mitochondria and *X. birchmanni* ancestry at *ndufs5* and *ndufa13*

(Supplementary Information 1.2.2). (**B**) Results of rt-qPCR evaluating evidence for differences in

mitochondrial copy number between genotypes in liver tissue. Ratio is derived from the difference in C$_t$

between a single copy nuclear gene (*Nup43*) and the mitochondrial gene *ND1*. (**C**) Results of Oroboros

O2K respirometer assay for adult *X. birchmanni*, *X. malinche*, and hybrid individuals with *X. malinche*

mitochondria and heterozygous ancestry at *ndufs5* and *ndufa13* (n=7 per genotype) point to lower

Complex I efficiency in heterozygous hybrids (orthogonal contrast $P = 0.023$). (**D**) Time to reach the

maximum rate of Complex I-driven respiration after the addition of ADP differed between heterozygous

hybrids and parental species (orthogonal contrast $P < 0.001$). Complex I-driven respiration begins with

the addition of ADP, as the flow of electrons is previously limited by the inability of Complex V to

relieve the proton gradient in the absence of its substrate (see Fig. S32 for example time-to-peak curves).

Time to reach the peak in Complex I- and Complex II-driven respiration after the addition of succinate

did not differ across genotypes (Fig. S33). (**E**) Maximum respiration rates during full O2K protocol as a

function of genotype did not differ between groups despite significant differences in Complex I efficiency

in heterozygotes (orthogonal contrast $P = 0.94$). (**F**) Allelic balance of *ndufs5* in the F_1 hybrid proteome

and transcriptome. Left subplot shows allele-specific expression of *ndufs5* in three adult F_1 hybrids. Right

subplot shows results of quantitative mass spectrometry analysis of *ndufs5* peptides in mitochondrial

proteomes derived from five adult F_1 hybrids. Data points show the proportion of area under the spectral

curves contributed by the *X. malinche* allele in a given individual. Left column shows results for

endogenous peptides present in F_1 hybrids, and right column shows results for the control where heavy-

labeled standards of each peptide were spiked in. Inset shows the identities of heavy-labeled peptides for

each species.

Fig. 4 | Predicted structures of *Xiphophorus* respiratory Complex I and evolutionary rates of

incompatible alleles. (**A**) *Xiphophorus* respiratory Complex I structures generated by RaptorX using

alignment to a template mouse cryo-EM structure. Colored protein structures include *ndufs5, ndufa13*,

and the four mitochondrially encoded *nd* genes in contact with *ndufs5* or *ndufa13*. Inset shows the surface

of predicted contacts between these genes. Solid black lines highlight two areas predicted to be in close

contact between interspecific substitutions (alpha carbon distance ≤ 10 Angstrom for all models), while

dashed lines show three additional areas in which there was weaker evidence for a predicted contact based

on computational analyses (side chain distance ≤ 12 Angstrom in at least one model). Asterisks denote

residues with substitutions in *X. birchmanni* computationally predicted to affect protein function (Table

S16). (**B**) Detailed view of interface between *ndufs5*, *nd2*, and *nd6.* Spheres highlight substitutions

between *X. birchmanni* and *X. malinche*. For substitutions predicted to be in close proximity, residues are

labeled with letters denoting the *X. malinche* allele, the residue number, and the *X. birchmanni* allele,

respectively (see Supplementary Information 1.4.6 and Table S16 for details). (**C**) Gene tree for *ndufs5*

generated with RAxML, highlighting an excess of substitutions along the *X. birchmanni* branch. Scale bar

represents number of nucleotide substitutions per site. Derived non-synonymous substitutions are

indicated by red ticks along the phylogeny. Note that spacing between ticks is arbitrary. (**D**) Distribution

of Log_{10} d_N/d_S between *X. birchmanni* and *X. malinche* across all nuclear genes in the genome with values

for *ndufs5* and *ndufa13* highlighted. (**E**) Gene tree for *ndufa13* generated with RAxML, highlighting an

excess of substitutions along the *X. birchmanni* branch (as in C).

Fig. 5 | Phylogenetic analysis and ancestry mapping suggest that genes underlying the mitonuclear

incompatibility have introgressed from *X. malinche* into *X. cortezi*. (**A**) Nuclear phylogeny of

Xiphophorus species, showing that *X. birchmanni* and *X. malinche* are sister species[37]. (**B**) Phylogeny

constructed from whole mitochondrial genome sequences showing that *X. cortezi* mitochondria are nested

within *X. malinche* mitochondrial diversity. (**C**) Results of simulations modeling expected mitochondrial

divergence between *X. malinche* and *X. cortezi* in a scenario with no gene flow. Distributions represent

pairwise sequence divergence in two sets of simulations. The first set used the average mitochondrial

substitution rate observed between *Xiphophorus* species (red), and the second used the minimum

mitochondrial substitution rate observed (blue). The dotted line shows observed divergence between

mitochondrial haplotypes in *X. malinche* and *X. cortezi*, indicating that past mitochondrial introgression is

more consistent with the observed data than incomplete lineage sorting (Supplementary Information

1.5.4). (**D**) Clustal alignment of *ndufs5* sequences shows that *X. malinche* and *X. cortezi* have identical

amino acid sequences at *ndufs5*, hinting at possible introgression of this nuclear gene, while *X.*

birchmanni is separated from them by four substitutions. Similar patterns are observed for *ndufa13*.

Colors indicate properties of the amino acid, asterisks indicate locations where the amino acid sequences

are identical. (**E**) Non-mitochondrial parent ancestry is strongly depleted in natural *X. cortezi* × *X.*

birchmanni hybrid populations fixed for the *X. cortezi* mitochondrial haplotype (Fig. S50) at *ndufs5* and

ndufa13. Arrows show the locations of *ndufa13* and *ndufs5* on chromosome 6 and chromosome 13

respectively, which fall in the center of minor parent ancestry deserts in both independently formed

populations, as expected for strong hybrid incompatibilities. Step curves show average *X. birchmanni*

ancestry in 250 kb windows, and horizontal lines show the genome-wide average for each population.

Methods

Biological Materials

Wild parental and hybrid individuals used in this study were collected from natural populations in Hidalgo, Mexico (Permit No. PPF/DGOPA-002/19). Artificial F_1 and F_2 hybrids were generated using mesocosm tanks as described previously[47]. Caudal fin clips were used as the source for all DNA isolation and for flow cytometry, and liver tissue for RNAseq, respirometry, and proteomic assays were collected following Stanford APLAC protocol #33071.

Genotyping and local ancestry calling

Genomic DNA was extracted from fin clips and individually barcoded tagmentation based libraries were generated (Supplementary Information 1.1.3). Hybrids were genotyped with low-coverage whole genome sequencing followed by local ancestry inference across the 24 *Xiphophorus* chromosomes and the mitochondrial genome using the *ancestryinfer* pipeline[46,47,50,71] (Supplementary Information 1.1.3–1.1.4). We converted posterior probabilities for each ancestry state to hard-calls for downstream analysis, using a posterior probability threshold of 0.9, and analyzed ancestry variation across the genome.

QTL and admixture mapping

The regions interacting with the mitochondrial genome were first identified based on analysis of segregation distortion in 943 F_2 hybrids generated from F_1 crosses between *X. malinche* females and *X. birchmanni* males (Supplementary Information 1.1.1 and Langdon et al[48]). Since all hybrids in this artificial cross harbored the *X. malinche* mitochondria, we scanned for regions of exceptionally high *X. malinche* ancestry along the genome (>60% *X. malinche*

ancestry), identifying one such region on chromosome 13 and one on chromosome 6 (Fig. 1; see

also[48]). Evidence for interactions between these regions and the mitochondrial genome were

confirmed using admixture mapping in two hybrid populations that segregated for the

mitochondrial haplotype of both species (Supplementary Information 1.1.2): the Calnali Low

hybrid population (N = 359) and the Chahuaco falls hybrid population (N = 244). Briefly, we

used a partial correlation analysis to identify regions of the genome strongly associated with

mitochondrial ancestry, after regressing out genome-wide ancestry to account for covariance in

ancestry due to population structure (see [44] and Supplementary Information 1.1.5, 1.1.9).

Significance thresholds were determined using simulations (Supplementary Information 1.1.5;

see also Supplementary Information 1.1.6).

Estimates of selection on the mitonuclear incompatibility

We used an ABC approach to estimate the strength of selection against the incompatible

interaction between the *X. malinche* mitochondrial haplotype and *X. birchmanni* ancestry at the

two nuclear genes involved in the hybrid incompatibility: *ndufs5* and *ndufa13* (Supplementary

Information 1.2.2). For these simulations, we asked what selection coefficients (0–1) and

dominance coefficients (0–1) could generate the observed deviations from the expectation of 50–

50 *birchmanni–malinche* ancestry in F_2 hybrids at *ndufs5* and *ndufa13* after two generations of

selection. We performed 500,000 simulations for each interaction and accepted or rejected

simulations based on comparisons to the real data using a 5% tolerance threshold

(Supplementary Information 1.2.2). We also evaluated evidence for incompatible interactions

with the *X. birchmanni* mitochondrial haplotype (Supplementary Information 1.2.1–1.2.4).

Developmental staging and genotyping of embryos

To pinpoint when in development the incompatibility between the *X. malinche*
mitochondria and *X. birchmanni* nuclear genotypes causes lethality, we collected a dataset on the
developmental stages of embryos with different genotype combinations. Whole ovaries were
removed from pregnant females and embryos were individually dissected. Each embryo was
assigned a developmental stage ranging from 1–11 based on established protocols for poeciliid
embryos[52]. Unfertilized eggs were excluded from analysis. Following staging, individual
embryos (N = 296) were genotyped as described above and in Supplementary Information 1.3.1.
We tested for significant differences in developmental stage between siblings with compatible
and incompatible genotype combinations using a two-sided two-sample t-test (Supplementary
Information 1.3.1) and examined differences in ancestry between large groups of siblings that
varied in their developmental stages (Supplementary Information 1.1.7). We also collected data
on embryonic stage and variability between siblings in embryonic stage from both pure parental
species (Supplementary Information 1.3.1). We used a different approach to pin-point the timing
of *ndufa13* lethality given that it appeared to act postnatally (Supplementary Information 1.3.3).

Embryo Respirometery and Morphometrics

To study the mechanisms of *ndufs5*- and *ndufa13*-driven lethality, we performed oxygen
consumption measurements on F_2 embryos in a Loligo plate respirometer (Supplementary
Information 1.3.4). Embryo were dissected from mothers and transferred to wells of a 24-well
plate, where their oxygen consumption was measured over 60 minutes. The measurement was
then repeated in media dosed with 5 µM rotenone to test sensitivity to Complex I inhibition, after
which the embryos were video recorded and photographed for morphometrics in ImageJ. We

used linear models to test the effect of *ndufs5* genotype, *ndufa13* genotype, and individual

standard length on a number of variables, controlling for batch effects (Supplementary

Information 1.3.4–1.3.5).

Mitochondrial respirometry

To further evaluate mitochondrial function in individuals heterozygous for the

mitonuclear incompatibility (Supplementary Information 1.3.6–1.3.7), we conducted

respirometry assays on *X. birchmanni*, *X. malinche*, and hybrid individuals that had the *X.*

malinche mitochondria and were heterozygous for the nuclear components of the hybrid

incompatibility (N=7 of each genotype). Mitochondria were isolated from whole liver tissue and

mitochondrial respiration was quantified using the Oroboros O2K respirometry system (Fig.

S30)[72]. A step-by-step description of this protocol and methods used to calculate respiratory flux

control factors is outlined in Supplementary Information 1.3.8. We complemented the results of

these respirometry experiments with measures of mitochondrial membrane potential using a flow

cytometry-based approach (Supplementary Information 1.3.9).

Parallel reaction monitoring proteomics

For Parallel Reaction Monitoring (PRM) with mass spectrometry, we used a similar

approach to that used for respirometry to isolate whole mitochondria from five F_1 hybrids (which

harbored *X. malinche* mitochondria). This approach is described in detail in Supplementary

Information 1.4.1. Briefly, we designed heavy-labeled peptides to distinguish between the *X.*

birchmanni and *X. malinche* copies of NDUFS5 and NDUFA13, facilitating quantification of the

peptides of interest in the mitochondrial proteome (Supplementary Information 1.4.2).

Mitochondrial isolates were prepared for mass spectrometry and combined with heavy-labeled

peptides in known quantities (see Supplementary Information 1.4.3), then submitted to Orbitrap

mass spectrometry with separation with UPLC and parallel reaction monitoring (PRM) for ion

selection. The protocol for mass spectrometry and PRM is described in detail in Supplementary

Information 1.4.4.

To analyze the results, the focal peptide's spectral peak were identified based on the peak

of the heavy labeled spike-in peptide. We focused analysis on the NDUFS5 peptide

WLL[L/P]QSGEQPYK since other endogenous peptides were below the expected sensitivity

limits of our PRM protocol (Supplementary Information 1.4.5). We normalized the intensity of

the NDUFS5 peptide based on the known spike-in quantity, and quantified the proportion of

NDUFS5 in each F_1 individual derived from *X. malinche* versus *X. birchmanni* (Supplementary

Information 1.4.5). We tested whether these ratios significantly deviated from the 50-50

expectation for F_1 hybrids using a two-sided one-sample *t*-test.

Complex I protein modeling

Mapping results allowed us to identify *ndufs5* and *ndufa13* as *X. birchmanni* genes that

interact negatively with *X. malinche* mitochondrial genes. We used a protein-modeling based

approach with RaptorX (http://raptorx.uchicago.edu) to identify the mitochondrial genes most

likely to interact with *ndufs5* and *ndufa13* (see Supplementary Information 1.4.6). Using the

mouse Cryo-EM structure (PDB ID 6G2J) of Complex I, we identified proteins in contact with

ndufs5 and *ndufa13*, which included several mitochondrial (*nd2, nd3, nd4l, nd6*) and nuclear

(*ndufa1*, *ndufa8*, *ndufb5*, *ndufc2*) genes. We then used RaptorX to predict structures for both the

X. birchmanni and *X. malinche* versions of the proteins. In addition, we evaluated the robustness

of these predictions to choice of Cryo-EM template; see Supplementary Information 1.4.6.

Analysis of evolutionary rates

Comparison of predicted protein sequences from *ndufs5, ndufa13*, and mitochondrial

genes of interest (*nd2* and *nd6*) revealed a large number of substitutions between *X. birchmanni*

and *X. malinche*. We calculated d_N/d_S between *X. birchmanni* and *X. malinche* for all annotated

protein coding genes throughout the genome and found that both *ndufs5* and *ndufa13* have rapid

protein evolution (Fig. 4D; Supplementary Information 1.5.1). Examining these mutations in a

phylogenetic context revealed that many substitutions in *ndufs5, nudfa13*, and *nd6* were derived

in *X. birchmanni*. We tested for significant differences in evolutionary rates on the *X. birchmanni*

lineage and for predicted functional impacts of these substitutions; these analyses are described

in Supplementary Information 1.5.1.

Tests for ancient introgression

Previous work had indicated that the mitochondrial phylogeny in *Xiphophorus* is

discordant with the whole-genome species tree[37]. Specifically, although *X. birchmanni* and *X.*

malinche are sister species based on the nuclear genome, *X. malinche* and *X. cortezi* are sister

species based on the mitochondrial genome. We used a combination of PacBio amplicon

sequencing of 10 individuals (2 or more per species, Supplementary Information 1.5.3) and

newly available whole-genome resequencing data to confirm this result and polarize the direction

of the discordance by constructing maximum likelihood mitochondrial phylogenies with the

program RAxML[74]. We performed similar phylogenetic analyses of the nuclear genes that

interact with the *X. malinche* mitochondria (*ndufs5* and *ndufa13*; Supplementary Information

1.5.3). Combined with phylogenetic results, simulation results suggest that gene flow from *X.*

malinche into *X. cortezi* is the most likely cause of the discordance we observe between the

mitochondrial and nuclear phylogenies (Supplementary Information 1.5.3–1.5.4). Since *X.*

malinche and *X. cortezi* are not currently sympatric, this suggests ancient gene flow between

them (Supplementary Information 1.5.5).

Contemporary hybridization between *X. birchmanni* and *X. cortezi*

To investigate the possibility that hybrids between *X. birchmanni* and *X. cortezi* share the

same mitonuclear incompatibility as observed in hybrids between *X. birchmanni* and *X. malinche*

(Supplementary Information 1.5.6), we took advantage of genomic data from recently discovered

hybrid populations between *X. birchmanni* and *X. cortezi*[75]. Using data from three different *X.*

birchmanni × *X. cortezi* populations and a permutation-based approach, we asked whether

ancestry at *ndufs5* and *ndufa13* showed lower mismatch with mitochondrial ancestry than

expected given the genome-wide ancestry distribution. This analysis is described in detail in

Supplementary Information 1.5.6.

Animal Care and Use

All methods were performed in compliance with Stanford Administrative Panel on Laboratory

Animal Care protocol #33071.

Data Availability

Raw sequencing reads used in this project are available under NCBI SRA Bioprojects

PRJNA744894, PRJNA746324, PRJNA610049, and PRJNA745218. Mass spectrometry data are

available on PRIDE (accession pending), and all other datasets necessary to recreate the results

of the publication are available on Dryad (accessions pending).

Code Availability

All new customs scripts used to generate results will be made available on Github at

https://github.com/Schumerlab/mitonuc_DMI and

https://github.com/Schumerlab/Lab_shared_scripts.

References

1. Dagilis, A. J., Kirkpatrick, M. & Bolnick, D. I. The evolution of hybrid fitness during speciation. *PLOS Genetics* **15**, e1008125 (2019).

2. Dobzhansky, T. Genetic Nature of Species Differences. *Am Nat* **71**, 404–420 (1937).

3. Muller, H. Isolating mechanisms, evolution, and temperature. *Biological Symposium* **6**, 71–125 (1942).

4. Maheshwari, S. & Barbash, D. A. The Genetics of Hybrid Incompatibilities. *Annu Rev Genet* **45**, 331–355 (2011).

5. Presgraves, D. C. The molecular evolutionary basis of species formation. *Nat Rev Genet* **11**, 175–180 (2010).

6. Swamy, K. B. S., Schuyler, S. C. & Leu, J.-Y. Protein Complexes Form a Basis for Complex Hybrid Incompatibility. *Front Genet* **12**, (2021).

7. Orr, H. A. The population genetics of speciation: the evolution of hybrid incompatibilities. *Genetics* **139**, 1805–1813 (1995).

8. Boocock, J., Sadhu, M. J., Durvasula, A., Bloom, J. S. & Kruglyak, L. Ancient balancing selection maintains incompatible versions of the galactose pathway in yeast. *Science* **371**, 415–419 (2021).

9. Wei, W.-H., Hemani, G. & Haley, C. S. Detecting epistasis in human complex traits. *Nat Rev Genet* **15**, 722–733 (2014).

10. Gauderman, W. J. Sample size requirements for association studies of gene-gene interaction. *Am J Epidemiol* **155**, 478–484 (2002).

11. Phadnis, N. *et al.* An essential cell cycle regulation gene causes hybrid inviability in Drosophila. *Science* **350**, 1552–1555 (2015).

12. Tang, S. & Presgraves, D. C. Evolution of the Drosophila Nuclear Pore Complex Results in

Multiple Hybrid Incompatibilities. *Science* **323**, 779–782 (2009).

13. Cooper, J. C., Guo, P., Bladen, J. & Phadnis, N. A triple-hybrid cross reveals a new hybrid

incompatibility locus between D. melanogaster and D. sechellia. *bioRxiv* 590588 (2019)

doi:10.1101/590588.

14. Kuzmin, E. *et al.* Systematic analysis of complex genetic interactions. *Science* **360**, (2018).

15. Johnson, N. A. Hybrid incompatibility genes: remnants of a genomic battlefield? *Trends*

Genet **26**, 317–325 (2010).

16. Burton, R. S. & Barreto, F. S. A disproportionate role for mtDNA in Dobzhansky–Muller

incompatibilities? *Mol Ecol* **21**, 4942–4957 (2012).

17. Hill, G. E. Mitonuclear coevolution as the genesis of speciation and the mitochondrial DNA

barcode gap. *Ecol Evol* **6**, 5831–5842 (2016).

18. Ballard, J. W. O. & Whitlock, M. C. The incomplete natural history of mitochondria. *Mol*

Ecol **13**, 729–744 (2004).

19. Haag-Liautard, C. *et al.* Direct Estimation of the Mitochondrial DNA Mutation Rate in

Drosophila melanogaster. *PLOS Biol* **6**, e204 (2008).

20. Allio, R., Donega, S., Galtier, N. & Nabholz, B. Large Variation in the Ratio of

Mitochondrial to Nuclear Mutation Rate across Animals: Implications for Genetic Diversity

and the Use of Mitochondrial DNA as a Molecular Marker. *Mol Biol Evol* **34**, 2762–2772

(2017).

21. Barr, C. M., Neiman, M. & Taylor, D. R. Inheritance and recombination of mitochondrial

genomes in plants, fungi and animals. *New Phytol* **168**, 39–50 (2005).

22. Chase, C. D. Cytoplasmic male sterility: a window to the world of plant mitochondrial–

nuclear interactions. *Trends Genet* **23**, 81–90 (2007).

23. Case, A. L., Finseth, F. R., Barr, C. M. & Fishman, L. Selfish evolution of cytonuclear

hybrid incompatibility in Mimulus. *Proc R Soc Lond B* **283**, 20161493 (2016).

24. Lane, N. & Martin, W. The energetics of genome complexity. *Nature* **467**, 929–934 (2010).

25. Osada, N. & Akashi, H. Mitochondrial–Nuclear Interactions and Accelerated Compensatory

Evolution: Evidence from the Primate Cytochrome c Oxidase Complex. *Mol Biol Evol* **29**,

337–346 (2012).

26. Barreto, F. S. & Burton, R. S. Evidence for Compensatory Evolution of Ribosomal Proteins

in Response to Rapid Divergence of Mitochondrial rRNA. *Mol Biol Evol* **30**, 310–314

(2013).

27. Bolnick, D. I., Turelli, M., López-Fernández, H., Wainwright, P. C. & Near, T. J.

Accelerated Mitochondrial Evolution and "Darwin's Corollary": Asymmetric Viability of

Reciprocal F1 Hybrids in Centrarchid Fishes. *Genetics* **178**, 1037–1048 (2008).

28. Turelli, M. & Moyle, L. C. Asymmetric Postmating Isolation: Darwin's Corollary to

Haldane's Rule. *Genetics* **176**, 1059–1088 (2007).

29. Brandvain, Y., Pauly, G. B., May, M. R. & Turelli, M. Explaining Darwin's Corollary to

Haldane's Rule: The Role of Mitonuclear Interactions in Asymmetric Postzygotic Isolation

Among Toads. *Genetics* **197**, 743–747 (2014).

30. Tiffin, P., Olson, S. & Moyle, L. C. Asymmetrical crossing barriers in angiosperms. *Proc R*

Soc Lond B **268**, 861–867 (2001).

31. Meiklejohn, C. D. *et al.* An Incompatibility between a Mitochondrial tRNA and Its Nuclear-

Encoded tRNA Synthetase Compromises Development and Fitness in Drosophila. *PLOS*

Genetics **9**, e1003238 (2013).

32. Luo, D. *et al.* A detrimental mitochondrial-nuclear interaction causes cytoplasmic male

sterility in rice. *Nat Genet* **45**, 573–577 (2013).

33. Lee, H.-Y. *et al.* Incompatibility of Nuclear and Mitochondrial Genomes Causes Hybrid

Sterility between Two Yeast Species. *Cell* **135**, 1065–1073 (2008).

34. Hanson, M. R. & Bentolila, S. Interactions of Mitochondrial and Nuclear Genes That Affect

Male Gametophyte Development. *The Plant Cell* **16**, S154–S169 (2004).

35. Taylor, S. A. & Larson, E. L. Insights from genomes into the evolutionary importance and

prevalence of hybridization in nature. *Nat Ecol Evol* **3**, 170–177 (2019).

36. Edelman, N. B. & Mallet, J. Prevalence and Adaptive Impact of Introgression. *Annu Rev*

Genet **55**, 265–283 (2021).

37. Cui, R. *et al.* Phylogenomics reveals extensive reticulate evolution in *Xiphophorus* fishes.

Evolution **67**, 2166–2179 (2013).

38. Nelson, T. C. *et al.* Ancient and recent introgression shape the evolutionary history of

pollinator adaptation and speciation in a model monkeyflower radiation (Mimulus section

Erythranthe). *PLoS Genet* **17**, e1009095 (2021).

39. Buck, R. & Flores-Rentería, L. The Syngameon Enigma. *Plants* **11**, 895 (2022).

40. Schumer, M., Cui, R., Rosenthal, G. G. & Andolfatto, P. Reproductive Isolation of Hybrid

Populations Driven by Genetic Incompatibilities. *PLoS Genet* **11**, e1005041 (2015).

41. Culumber, Z. W. *et al.* Replicated hybrid zones of *Xiphophorus* swordtails along an

elevational gradient. *Mol Ecol* **20**, 342–356 (2011).

42. Fisher, H. S., Wong, B. B. M. & Rosenthal, G. G. Alteration of the chemical environment

disrupts communication in a freshwater fish. *Proc R Soc B* **273**, 1187–1193 (2006).

43. Schumer, M. *et al.* Natural selection interacts with recombination to shape the evolution of

hybrid genomes. *Science* **360**, 656–660 (2018).

44. Schumer, M. & Brandvain, Y. Determining epistatic selection in admixed populations. *Mol*

Ecol **25**, 2577–2591 (2016).

45. Schumer, M. *et al.* High-resolution mapping reveals hundreds of genetic incompatibilities in

hybridizing fish species. *eLife* **3**, e02535 (2014).

46. Powell, D. L. *et al.* Natural hybridization reveals incompatible alleles that cause melanoma

in swordtail fish. *Science* **368**, 731–736 (2020).

47. Powell, D. L. *et al.* The Genetic Architecture of Variation in the Sexually Selected Sword

Ornament and Its Evolution in Hybrid Populations. *Curr Biol* **31**, 923-935.e11 (2021).

48. Langdon, Q. K. *et al.* Predictability and parallelism in the contemporary evolution of hybrid

genomes. *PLoS Genet* **18**, e1009914 (2022).

49. Schartl, M. *et al.* The genome of the platyfish, Xiphophorus maculatus, provides insights into

evolutionary adaptation and several complex traits. *Nat Genet* **45**, 567–572 (2013).

50. Schumer, M., Powell, D. L. & Corbett-Detig, R. Versatile simulations of admixture and

accurate local ancestry inference with mixnmatch and ancestryinfer. *Mol Ecol Resour* **20**,

1141–1151 (2020).

51. Sharma, L. K., Lu, J. & Bai, Y. Mitochondrial Respiratory Complex I: Structure, Function

and Implication in Human Diseases. *Curr Med Chem* **16**, 1266–1277 (2009).

52. Haynes, J. L. Standardized Classification of Poeciliid Development for Life-History Studies.

Copeia **1995**, 147 (1995).

53. Tavolga, W. N. Embryonic development of the platyfish (Platypoecilus), the swordtail

(Xiphophorus), and their hybrids. Bulletin of the AMNH ; v. 94, article 4. *Embryonic*

development in fish (1949).

54. Byrnes, J. *et al.* Pharmacologic modeling of primary mitochondrial respiratory chain

dysfunction in zebrafish. *Neurochem Int* **117**, 23–34 (2018).

55. Pinho, B. R. *et al.* How mitochondrial dysfunction affects zebrafish development and

cardiovascular function: an in vivo model for testing mitochondria-targeted drugs. *Br J*

Pharmacol **169**, 1072–1090 (2013).

56. González-Quintana, A. *et al.* Novel NDUFA13 Mutations Associated with OXPHOS

Deficiency and Leigh Syndrome: A Second Family Report. *Genes (Basel)* **11**, 855 (2020).

57. McElroy, G. S. *et al.* NAD+ Regeneration Rescues Lifespan, but Not Ataxia, in a Mouse

Model of Brain Mitochondrial Complex I Dysfunction. *Cell Metab* **32**, 301-308.e6 (2020).

58. McElroy, G. S. *et al.* Reduced expression of mitochondrial complex I subunit Ndufs2 does

not impact healthspan in mice. *Sci Rep* **12**, 5196 (2022).

59. Agip, A.-N. A. *et al.* Cryo-EM structures of complex I from mouse heart mitochondria in

two biochemically defined states. *Nat Struct Mol Biol* **25**, 548–556 (2018).

60. Kampjut, D. & Sazanov, L. A. The coupling mechanism of mammalian respiratory complex

I. *Science* **370**, (2020).

61. Fiedorczuk, K. *et al.* Atomic structure of the entire mammalian mitochondrial complex I.

Nature **538**, 406–410 (2016).

62. Mack, K. L. & Nachman, M. W. Gene Regulation and Speciation. *Trends Genet* **33**, 68–80

(2017).

63. Källberg, M. *et al.* Template-based protein structure modeling using the RaptorX web server.

Nature Protocols **7**, 1511–1522 (2012).

64. Formosa, L. E., Dibley, M. G., Stroud, D. A. & Ryan, M. T. Building a complex complex:

Assembly of mitochondrial respiratory chain complex I. *Semin Cell Dev Biol* **76**, 154–162

(2018).

65. Hill, G. E. Mitonuclear Compensatory Coevolution. *Trends Genet* **36**, 403–414 (2020).

66. Sloan, D. B., Havird, J. C. & Sharbrough, J. The on-again, off-again relationship between

mitochondrial genomes and species boundaries. *Mol Ecol* **26**, 2212–2236 (2017).

67. Ellison, C. K. & Burton, R. S. Disruption of Mitochondrial Function in Interpopulation

Hybrids of Tigriopus Californicus. *Evolution* **60**, 1382–1391 (2006).

68. Ellison, C. K., Niehuis, O. & Gadau, J. Hybrid breakdown and mitochondrial dysfunction in

hybrids of Nasonia parasitoid wasps. *J Evol Biol* **21**, 1844–1851 (2008).

69. Olson, J. R., Cooper, S. J., Swanson, D. L., Braun, M. J. & Williams, J. B. The Relationship

of Metabolic Performance and Distribution in Black-Capped and Carolina Chickadees.

Physiol Biochem Zool **83**, 263–275 (2010).

70. Lu, Y. *et al.* Oncogenic allelic interaction in Xiphophorus highlights hybrid incompatibility.

PNAS **117**, 29786–29794 (2020).

71. Corbett-Detig, R. & Nielsen, R. A Hidden Markov Model Approach for Simultaneously

Estimating Local Ancestry and Admixture Time Using Next Generation Sequence Data in

Samples of Arbitrary Ploidy. *PLOS Genetics* **13**, e1006529 (2017).

72. Chung, D. J., Bryant, H. J. & Schulte, P. M. Thermal acclimation and subspecies-specific

effects on heart and brain mitochondrial performance in a eurythermal teleost (Fundulus

heteroclitus). *J Exp Biol* **220**, 1459–1471 (2017).

73. Ng, P. C. & Henikoff, S. SIFT: predicting amino acid changes that affect protein function.

Nucleic Acids Res **31**, 3812–3814 (2003).

74. Stamatakis, A. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. *Bioinformatics* **22**, 2688–2690 (2006).

75. Powell, D. L. *et al.* Two new hybrid populations expand the swordtail hybridization model

system. *Evolution* **75**, 2524–2539 (2021).

Acknowledgements

We thank Peter Andolfatto, Stepfanie Aguillon, Yaniv Brandvain, Jenn Coughlan, Hunter Fraser, Yuki Haba, Nitin Phadnis, Molly Przeworski, Ken Thompson and members of the Schumer lab for helpful discussion and/or feedback on earlier versions of this manuscript. We thank Alexa Pollock for help performing rotenone trials. We thank the Federal Government of Mexico for permission to collect fish. Stanford University and the Stanford Research Computing Center provided computational support for this project. We thank the Vincent Coates Foundation Mass Spectrometry Laboratory, Stanford University Mass Spectrometry (RRID:SCR_017801) for technical and experimental support. **Funding:** This work was supported by a Knight-Hennessy Scholars fellowship and NSF GRFP 2019273798 to B. Moran, a CEHG fellowship and NSF PRFB (2010950) to Q. Langdon, NIH P30 CA124435 in utilizing the Stanford Cancer Institute Proteomics/Mass Spectrometry Shared Resource, NIH grant 1R35GM142836 to J. Havird, and a Hanna H. Gray fellowship, Sloan Fellowship, and NIH grant 1R35GM133774 to M. Schumer.

Author Contributions: B.M.M., D.L.P, and M. Schu, designed the project; B.M.M., C.Y.P., E.N.K.I., S.M.B, A.E.D, A.M, J.J.B, K.K., F.L., R.M., K.S., O.H.-P., J.C.H., A. M. and M. Schu. collected data; B.M.M., C.Y.P., Q.L.K., F.L., J.C.H., A.M., and M. Schu. performed analyses; D.L.P., T.R.G., R.D.L., C.G., R.C.-D., J.F. and M. Scha. provided expertise and technical support.

Competing Interests: the authors declare no competing interests.

941   **Materials & Correspondence:** correspondence and requests for materials should be directed to

942   B.M.M. (benmoran@stanford.edu) or M. Schu (schumer@stanford.edu).