Modeling miRNA-driven post-transcriptional regulation by using exon-intron split analysis (EISA) in pigs

Emilio Mármol-Sánchez[1*], Susanna Cirera[2], Laura M. Zingaretti[3], Mette Juul Jacobsen[2], Yuliaxis Ramayo-Caldas[4], Claus B. Jørgensen[2], Merete Fredholm[2], Tainã Figueiredo Cardoso[1†], Raquel Quintanilla[4], Marcel Amills[1,5]

[1]Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain.

[2]Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, 1871 Frederiksberg C, Denmark.

[3]Universidad Nacional de Villa María, Villa María, Córdoba, Argentina.

[4]Animal Breeding and Genetics Program, Institute for Research and Technology in Food and Agriculture (IRTA), Torre Marimon, 08140 Caldes de Montbui, Barcelona, Spain.

[5]Departament de Ciència Animal i dels Aliments, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain.

*Emilio Mármol-Sánchez current address: Stockholm University, The Wenner-Gren Institute, Department of Molecular Biosciences, SciLifelab.

†Tainã Figueiredo Cardoso current address: Embrapa Pecuária Sudeste, Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA), 13560-970, São Carlos, SP, Brazil.

24   **Corresponding author:** Emilio Mármol-Sánchez. Stockholm University, The Wenner-

25   Gren Institute, Department of Molecular Biosciences, SciLifelab. Email:

26   emilio.marmol.sanchez@gmail.com

Abstract

Bulk sequencing of RNA transcripts has typically been used to quantify gene expression

levels in different experimental systems. However, linking differentially expressed (DE)

mRNA transcripts to gene expression regulators, such as miRNAs, remains challenging,

as miRNA-mRNA interactions are commonly identified *post hoc* after selecting sets of

genes of interest, thus biasing the interpretation of underlying gene regulatory networks.

In this study, we aimed at disentangling miRNA-driven post-transcriptional signals

linked to porcine muscle and adipose tissue energy homeostasis. For this purpose, we

performed an exon-intron split analysis (EISA) on muscle and fat RNA-seq data from

two independent pig populations. One of these populations was subjected to fasting-

feeding conditions, while the other represented divergent fatness profiles. After running

EISA, protein-coding mRNA genes with downregulated exonic fractions and high post-

transcriptional signals were significantly enriched for binding sites of DE upregulated

miRNAs. Moreover, these downregulated genes showed an increased expression

covariation for the exonic fraction compared to the intronic fraction. On the contrary,

they did not show enrichment for binding sites of non-DE highly expressed or

downregulated DE miRNAs. Among the set of loci displaying miRNA-driven post-

transcriptional regulatory signals, we observed genes related to glucose homeostasis

(*DKK2*, *PDK4*, *IL18*, *NR4A3*, *CHRNA1*, *TET2*), cell differentiation (*PBX1*, *BACH2*)

and adipocytes metabolism (*SESN3*, *ESRRG*, *SAMD4*, *LEP*, *PTGFR*, *SERPINE2*,

RNF157, *GPLD1*, *NCF2*, *OSBPL10*, *PRSS23*). Our results highlighted mRNA genes

showing post-transcriptional miRNA-driven downregulation using the exonic and intronic fractions of RNA-seq datasets from muscle and adipose tissues in pigs.

Keywords: Exon-intron split analysis, microRNA, pigs, energy homeostasis.

1. Introduction

Messenger RNA (mRNA) expression and turnover in metabolic processes are subjected to complex yet poorly characterized regulatory mechanisms that contribute to shaping fine-tuned biological responses to different stimuli [1]. Cellular metabolic changes are hence a direct manifestation of intricate interactions between expressed transcripts and other regulatory elements that modify their abundance, localization, fate and degradation rate. MicroRNAs (miRNAs) are primarily engaged in the post-transcriptional control of gene expression through inhibition of translation and/or destabilization of target mRNAs by poly(A) shortening and subsequent degradation [2]. Changes in the abundance of mRNAs targeted by miRNAs can be inferred through covariation analysis. Such approach can help to unravel direct or indirect molecular interactions connecting and regulating biological networks.

In order to disentangle regulatory functions driven by miRNAs, research studies typically focus on specific sets of genes of interest, showing significant expression changes and harboring binding sites for DE miRNAs [3–9]. This approach, however, is biased by the fact that genes are selected after differential expression analysis and by the performance of an *ad hoc* search of predicted interactions between the 3'-UTRs of mRNAs and the seed regions of miRNAs. Besides, one of the main limitations of differential expression analysis is that it does not discriminate whether changes in expression take place either at the transcriptional or post-transcriptional levels. Such

distinction is essential to understand at which level of the mRNA life-cycle the regulation is taking place.

To address this issue, Gaidatzis *et al*. [10] proposed that the magnitude of the post-transcriptional component can be deduced by comparing the amounts of exonic and intronic reads. Based on this, these authors devised a methodology denoted *exon-intron split analysis* (EISA), which separates the transcriptional and post-transcriptional components of gene regulation. By considering that intronic reads are mainly derived from heterogeneous nuclear RNAs (unprocessed mRNAs or pre-mRNAs), they assessed the magnitude of the transcriptional regulation. Such assumption is based on early reports describing intronic expression as a proxy of nascent transcription and co-transcriptional splicing events [11–13]. In this way, a gene showing similar levels of intronic reads in two different states but a strong downregulation of exonic reads after applying a certain treatment or challenge (nutrition, infection, temperature etc.), could be indicative of an induced inhibition at the post-transcriptional level [10,14,15].

A high number of differential expression studies have been performed in pigs during the last decade [16–24], but to the best of our knowledge, in none of these studies the transcriptional and post-transcriptional components of gene regulation have been independently analyzed. In the present study, we aimed to dissect the contribution of miRNAs to post-transcriptional regulation in pigs by applying the EISA methodology using two independent porcine experimental models.

2. Materials and methods

2.1. Experimental design, sampling and processing

In this study, we have used two independent experimental systems:

(i) Duroc pigs: Twenty-three gilts divided in two fasting/feeding regimes, i.e., 11 gilts

(*AL-T0*) slaughtered in fasting conditions and 12 gilts (*AL-T2*) slaughtered immediately

after 7 h with access to *ad libitum* feed intake [7,22,25]. Immediately after slaughtering,

gluteus medius (GM) skeletal muscle samples were collected and snap-frozen at -80°C.

(ii) Duroc-Göttingen minipig F_2 inter-cross: Ten individuals with divergent fatness

profiles for their body mass index (BMI) metric (5 *lean* and 5 *obese*) were selected from

the UNIK resource population [26,27], as described in Jacobsen et al. 2019 [28].

Retroperitoneal adipose tissue was collected at slaughter and mature adipocytes were

subsequently isolated following the protocol of Decaunes et al. 2011 [29] with

modifications reported in [28].

Further details about RNA-seq and small RNA-seq expression data generated from both

experimental designs have been previously described [7,22,28]. Sequencing reads

generated in the RNA-Seq and small RNA-Seq datasets from both pig resources were

trimmed with the Cutadapt software [30]. Reads were mapped against the Sscrofa11.1

porcine assembly [31] with the HISAT2 aligner [32] and default parameters for RNA-

Seq reads. In contrast, the Bowtie Alignment v.1.2.1.1 software [33] with small

sequence reads specifications (*bowtie -n 0 -l 25 -m 20 -k 1 --best --strata*) was used to

align small RNA-Seq reads to the Sscrofa11.1 porcine reference assembly [31].

2.2. Exon/Intron quantification

We generated exonic and intronic-specific annotations spanning all genes available

using the gtf formatted Sscrofa.11.1 v.103 gene annotation file (Ensembl repositories:

http://ftp.ensembl.org/pub/release-103/gtf/sus_scrofa/). Overlapping intronic/exonic

regions, as well as singleton positions were removed [34]. Each intronic region was

trimmed by removing 10 nucleotides on both ends to avoid exonic reads mapping close

to exon/intron junctions. We then used the featureCounts function included in the

Rsubread package [35] to quantify gene expression profiles based on exonic and intron

expression patterns for each gene, independently. MiRNA expression profiles were

estimated using the Sscrofa11.1 v.103 mature miRNA annotation with the

featureCounts software tool [36] in single-end mode and with default parameters.

2.3. Exon/intron split analysis (EISA).

We applied EISA to infer post-transcriptional gene regulation in our two independent

porcine datasets. For this purpose, we separately estimated the exonic and intronic

abundance of each annotated mRNA gene using the Sscrofa11.1 v.103 exon/intron

custom annotation generated as described above. Only genes showing average

expression values above 1 count-per-million in at least 50% of animals were retained

for further analyses. Normalization was performed independently for exon and intron

counts by multiplying each i^{th} gene expression in each j^{th} sample by the corresponding

mean gene expression and dividing by the total number of quantified counts per sample

[10]. Exonic and intronic gene abundances were subsequently transformed to a $\log_2$

scale, adding a pseudo-count of 1 and averaged within each considered treatment groups

(*AL-T0* and *AL-T2* for GM tissues and *lean* and *obese* for adipocyte isolates).

Only genes with successful exonic and intronic quantified read counts were considered

in our analyses. The transcriptional component (Tc) contribution to the observed

differences in each i^{th} gene was expressed as the increment of intronic counts in fed

(*AL-T2*) and *obese* animals with respect to fasting (*AL-T0*) and *lean* animals (ΔInt =

$\text{Int}_{2i} - \text{Int}_{1i}$), respectively. The increment of exonic counts (ΔEx) was also calculated,

and the post-transcriptional component (PTc) effect was expressed as ΔEx $-$ ΔInt =

$(\text{Ex}_{2i} - \text{Ex}_{1i}) - (\text{Int}_{2i} - \text{Int}_{1i})$. Both components were z-scored to represent comparable

ranges between ΔEx and ΔInt estimates. All implemented analyses have been

summarized in **Fig. S1**. A ready-to-use modular pipeline for running EISA is publicly

available at https://github.com/emarmolsanchez/EISACompR.

2.4. Post-transcriptional signal prioritization

In order to obtain a prioritized list of genes showing relevant signals of post-

transcriptional regulation, the top 5% genes with the highest negative PTc scores were

retrieved. We only focused on genes showing strongly reduced exonic fraction (ΔEx)

values of at least 2-folds for post-transcriptional signals in both experimental systems.

2.5. Differential expression analyses and significance of PTc scores

Differential expression analyses were carried out with the *edgeR* package [37] by

considering the exonic fraction of mRNAs, as well as miRNA expression profiles from

RNA-Seq and small RNA-Seq datasets, respectively, and in the two experimental

systems under study. Expression filtered raw counts for exonic reads were normalized

with the trimmed mean of M-values normalization (TMM) method [38] and the

statistical significance of mean expression differences was tested with a quasi-likelihood

F-test [37]. Correction for multiple hypothesis testing was implemented with the

Benjamini-Hochberg false discovery rate approach [39]. Messenger RNAs were

considered as differentially expressed (DE) when the absolute value of the fold-change

(FC) was higher than 2 ($|FC| > 2$) and q-value < 0.05. For miRNAs, $|FC| > 1.5$ and q-

value < 0.05 were used instead. This more flexible threshold was motivated by the fact

that miRNAs are commonly lowly expressed and show more stable and subtle

expression changes compared to mRNAs [7,40]. The statistical significance of the post-

transcriptional (PTc) scores was evaluated by incorporating the intronic quantification

as an interaction effect for exonic abundances [10]. The fasting pigs (*AL-T0*) from

Duroc skeletal muscle, as well as *obese* pigs from Duroc-Göttingen adipocyte

expression profiles were considered to be the baseline controls, i.e., any given

upregulation in ΔEx or ΔInt values represents and overexpression in fed (*AL-T2*) Duroc

gilts and *lean* Duroc-Göttingen minipigs with respect to their fasting (*AL-T0*) and *obese*

counterparts.

2.6. miRNA target prediction

Putative interactions between the seed regions of expressed miRNAs (small RNA-seq

datasets) and the 3'-UTRs of expressed protein-coding mRNA genes (RNA-seq

datasets) were predicted on the basis of sequence identity using the Sscrofa11.1

reference assembly. The annotated 3'-UTRs longer than 30 nts from porcine mRNAs

were retrieved from the Sscrofa11.1 v.103 available at BioMart

(http://www.ensembl.org/biomart) and miRBase [41] databases. Redundant seeds from

mature porcine microRNAs were removed. The seedVicious v1.1 tool [42] was used to

infer miRNA-mRNA interactions. MiRNA-mRNA 8mer, 7mer-m8 and 7mer-A1

interactions were considered as the most relevant among the full set of canonical and

non-canonical interactions [2,43,44].

Based on the study of Grimson et al. [43], the *in silico*-predicted miRNA-mRNA

interactions matching any of the following criteria were removed: (i) Binding sites

located in 3'-UTRs at less than 15 nts close to the end of the open reading frame (and

the stop codon) or less than 15 nts close to the terminal poly(A) tail (E criterion), (ii)

binding sites located in the middle of the 3'-UTR in a range comprising 45-55% of the

central region of the non-coding sequence (M criterion), and (iii) binding sites lack AU-

rich elements in their immediate upstream and downstream flanking regions comprising

30 nts each (AU criterion).

Covariation patterns between miRNAs and their predicted mRNA targets were assessed

by computing Spearman's correlation coefficients (ρ) with the TMM normalized and

$\log_2$ transformed expression profiles of the exonic fractions of mRNA and miRNA

genes. To determine the contribution of miRNAs to post-transcriptional regulation in

the two experimental systems under study, only miRNA-mRNA predicted pairs

comprising DE upregulated miRNAs (FC > 1.5; q-value < 0.05) and mRNA genes with

relevant PTc scores (see post-transcriptional signal prioritization section) were taken

into consideration.

2.7. miRNA target enrichment analyses

We sought to determine if the overall number of mRNA genes with high post-

transcriptional signals (the ones with the top 5% negative PTc scores and reduced ΔEx

values > 2-folds) were significantly enriched to be targeted by at least one upregulated

miRNA (FC > 1.5; q-value < 0.05), compared with the whole set of expressed mRNAs

genes with available 3'-UTRs from both datasets. Enrichment analyses were carried out

using the Fisher's exact test implemented in the *fisher.test* R function. Significance

level was set at nominal P-value < 0.05.

We also tested whether these genes were enriched for binding sites of the top 5% most

highly expressed miRNA genes, excluding significantly upregulated miRNAs, as well

as for binding sites of significantly downregulated miRNAs (FC < -1.5; q-value < 0.05).

Given the relatively low statistical significance of DE miRNAs observed in the UNIK

Duroc-Göttingen minipigs (l*ean* vs *obese*), we considered that miRNAs were

significantly upregulated in this particular data set when FC >1.5 and *P*-value < 0.01.

Further details can be found in **Supplementary Methods**.

2.8. Gene covariation network and covariation enrichment score

We computed pairwise correlation coefficients among the whole set of DE mRNA

genes in the *AL-T0* vs *AL-T2* (*q*-value < 0.05, N = 454) and *lean* vs *obese* (*q*-value <

0.05, N = 299). These correlations were compared with those corresponding to the set of

genes with relevant post-transcriptional signals and putatively targeted by DE

upregulated miRNAs. Normalized exonic and intronic estimates in the log_2 scale

obtained from EISA analyses were used to compute Spearman's correlation coefficients

(ρ) for each potential pair of DE mRNA genes plus those with post-transcriptional

signals but without significant DE. Self-correlation pairs were excluded. Significant

correlations were identified with the Partial Correlation with Information Theory (PCIT)

network inference algorithm [45] implemented in the *pcit* R package [46]. Non-

significant covarying pairs were set to zero, while a value of 1 was assigned to the

significant ones with both positive or negative coefficients $|\rho| > 0.6$.

The potential contribution of miRNAs to shape the observed covariation patterns was

assessed by calculating a covariation enrichment score (CES) following Tarbier et al.

2020 [47]. If multiple genes are downregulated by any upregulated miRNAs in a

coordinated manner, we would expect to observe a reduced abundance in their mature

spliced mRNA forms but not in the corresponding primary transcripts, i.e., we would

detect a covariation only for the exonic fractions (but not for the intronic ones). In other

words, the intronic fraction, eventually spliced and degraded in the nucleus, should not

reflect any posterior post-transcriptional regulatory effects in the cytoplasm, so little or

null covariation might be expected. Significant differences among the set of exonic,

intronic and control CES values were tested with a non-parametric approach using a

Mann-Whitney U non-parametric test [48].

As a control randomized test for enrichment analyses between miRNAs and mRNAs

with high post-transcriptional downregulatory signals, we implemented a bootstrap

corrected iteration to generate 100 random sets of 10 expressed mature miRNA genes

without seed redundancy, which were used as input for miRNA target prediction with

the sets of mRNA genes with the high post-transcriptional signals (top 5% PTc scores

and at least 2-folds ΔEx reduction). The distribution of odds ratios obtained after

iterating over each random set of miRNAs (N = 100) were then compared with odds

ratios obtained with the set of significantly upregulated miRNAs.

The *P*-value for the significance of the deviation of observed odds ratios against the

bootstrapped odds ratios distribution was defined as, $P - value = 1 - \frac{r+1}{k+1}$, where r

is the number of permuted odds ratios with values equal or higher than the observed

odds ratio for enrichment analyses with the set of upregulated miRNAs, and k is the

number of defined permutations (N = 100).

2.9. Estimating the expression levels of miRNAs and several of their predicted

mRNA targets by qPCR

The same total RNA extracted from adipocytes and used for sequencing was

subsequently employed for cDNA synthesis and qPCR verification. Five mRNAs (*LEP,*

OSBLP10, PRSS23, RNF157 and *SERPINE2*) among the top 5% negative PTc scores

and showing at least 2-folds reduction in their ΔEx values were selected for qPCR

profiling. Two reference genes (*TBP* and *ACTB*, as defined by Nygard et al. 2007 [49])

were used for normalization. Accordingly, three of the most DE miRNAs were selected

for qPCR profiling (ssc-miR-92b-3p, ssc-miR-148a-3p and ssc-miR-214-3p), plus two

highly expressed non-DE miRNAs for normalization (ssc-let-7a and ssc-miR-23a-3p)

from the *lean* vs *obese* small RNA-Seq dataset. Further details about qPCR

experimental procedures are available in **Supplementary Methods**. All primers for

mRNA and miRNA expression profiling are available at **Table S2**. Raw Cq values for

each assay are available at **Table S3**.

277   3. Results

278   3.1. The analysis of post-transcriptional regulation in muscle samples from fasting
279   and fed Duroc gilts

Identification of genes predicted to be post-transcriptionally regulated by miRNAs

After the processing, mapping and quantification of mRNA and miRNA expression

levels in GM skeletal muscle samples from Duroc gilts, an average of 45.2 million reads

per sample (~93%) were successfully mapped to 31,908 genes annotated in the

Sscrofa11.1 v.103 assembly (including protein coding and non-coding genes). Besides,

an average of 2.2 million reads per sample (~42%) mapped to 370 annotated porcine

miRNA genes.

A total of 30,322 (based on exonic reads) and 22,769 (based on intronic reads) genes

were successfully quantified after splitting the reference Sscrofa11.1 v.103 assembly

between exonic and intronic features. The exonic fraction displayed an average of

1,923.94 estimated counts per gene, whereas the intronic fraction showed an average of

83.02 counts per gene. In other words, exonic counts were ~23 fold more abundant than

those corresponding to intronic regions.

Differential expression analyses based on exonic fractions identified 454 DE genes.

Among those, only genes with |FC| > 2 were retained, making a total of 52 upregulated

and 80 downregulated genes in the *AL-T0* vs *AL-T2* comparison (**Table S4, Fig. S2A**).

Besides, differential expression analyses on small RNA-seq data revealed 16 DE

miRNAs, of which 8 were upregulated in *AL-T2* pigs, representing 6 unique miRNA

seeds (**Table S5**). These non-redundant seeds of significantly upregulated miRNAs in

fed animals (ssc-miR-148a-3p, ssc-miR-7-5p, ssc-miR-30-3p, ssc-miR-151-3p, ssc-

miR-374a-3p and ssc-miR-421-5p) were selected as potential post-transcriptional

regulators of GM muscle mRNA expression in response to nutrient supply. EISA results

made possible to detect 133 genes with significant effects (**Table S6**), of which three

had > 2-folds reduced ΔEx fractions and two of them had significantly negative PTc

scores (**Table 1**).

To detect mRNA genes putatively downregulated by miRNAs at the post-transcriptional

level, mRNA genes displaying the top 5% negative PTc scores with at least 2-folds ΔEx

reduction were selected as putative miRNA-targets (**Fig. S2B**). With this approach, 26

mRNA genes were selected (**Table 1**). One of them (ENSSSCG00000049158) did not

have a properly annotated 3´-UTR so it was excluded from further analyses. Among this

set of genes with high post-transcriptional signals, 18 appeared as significantly

downregulated (FC < -2; *q*-value < 0.05) in the differential expression analysis

considering their exonic fractions (**Table 1** and **Table S4**).

Context-based pruning of predicted miRNA-mRNA interactions removes spurious

unreliable target events

As a first step to determine if genes with highly negative PTc scores and showing a

marked reduction in exonic fractions were repressed by upregulated DE miRNAs

(**Table S5**), we investigated the accuracy and reliability of *in silico* predictions

regarding miRNA binding sites in their 3'-UTRs (**Table S7**). We evaluated the presence

of enriched binding sites over a random background of expressed genes with no

context-based removal of predicted binding sites, applying each one of the three

selection criteria reported in Methods independently, as well as by combining them

pairwise or altogether. As depicted in **Fig. S3A** and **S3B**, introducing additional

context-based filtering criteria for removing spurious unreliable binding site predictions

resulted in an overall increased enrichment of miRNA targeted genes within the top 1%

(**Fig. S3A**) and 5% (**Fig. S3B**) negative PTc scores and displaying > 2-folds ΔEx

reduction. This significantly increased enrichment was more evident when using the AU

criterion, as shown in **Fig. S3A**. However, we also detected a slight increment when

adding the other two context-based removal criteria (M and E). These findings were less

evident when taking into consideration the list of the top 5% genes (**Fig. S3B**).

Nevertheless, an increased enrichment for targeted mRNAs by DE upregulated miRNAs

was detectable for all combined filtering criteria, especially for 7mer-A1 binding sites,

and probably at the expense of the scarcer and more efficient 8mer binding sites. Based

on these results, we decided to apply the three joint criteria (AU, M and E) for

enrichment analyses between the set of mRNA genes with high post-transcriptional

signals and putatively targeted by miRNAs.

Genes with relevant post-transcriptional signals are enriched for putative miRNA

binding sites in their 3'-UTRs

Target prediction and context-based pruning of miRNA-mRNA interactions for mRNA

genes displaying the top 5% negative PTc scores and at least 2-folds reductions in the

ΔEx exonic fraction (N = 25 after excluding ENSSSCG00000049158; **Fig. 1A**) made

possible to detect 11 8mer, 21 7mer-m8 and 22 7mer-A1 miRNA binding sites (**Table**

S7) corresponding to the 6 non-redundant seeds of DE miRNAs upregulated in *AL-T2*

gilts. This set of 25 mRNA genes showing putative post-transcriptional repression by

miRNAs showed a significant enrichment in 8mer, 7mer-m8 and 7mer-A1 sites (**Fig.**

1B), and this was especially relevant when combining these three types of binding sites.

More importantly, 21 out of these 25 genes were predicted as putative targets of

miRNAs upregulated in the GM muscle samples from *AL-T2* fed gilts (**Table 1** and

Table S5).

We also evaluated the enrichment of the mRNA genes with the top 5% negative PTc

scores and at least 2-folds ΔEx reduction (N = 25, **Table 1**) to be targeted by the

following sets of miRNAs: (i) Non-redundant miRNAs downregulated in *AL-T2* fed

gilts (ssc-miR-1285, ssc-miR-758, ssc-miR-339, sc-miR-22-3p, ssc-miR-296-5p, ssc-

miR-129a-3p, ssc-miR-181c and ssc-miR-19b, **Table S5**), (ii) the top 5% most

expressed miRNAs, excluding those being upregulated (ssc-miR-1, ssc-miR-133a-3p,

ssc-miR-26a, ssc-miR-10b, ssc-miR-378, ssc-miR-99a-5p, ssc-miR-27b-3p, ssc-miR-

30d, ssc-miR-486 and ssc-let-7f-5p), and (iii) for an iteration (N = 100) of random sets

of 10 expressed miRNAs, irrespective of their DE and abundance status, as a control

test. None of these additional analyses recovered a significant enrichment for any type

of the three considered miRNA target subtypes (**Fig. 1B**).

The mRNA with the highest and significant PTc score was the Dickkopf WNT

Signaling Pathway Inhibitor 2 (*DKK2*), and thus, this was the gene with the highest

putative miRNA-driven downregulation and no additional transcriptional influence

(**Table 1**). Moreover, the DKK2 locus was the only gene harboring two miRNA 8mer

binding sites (**Table S7**), although it was not among the most highly DE mRNA genes

according to differential expression analyses (**Table S4**). As depicted in **Table 1**, only

suggestive significant differences (*P*-value < 0.01) in mean expression were found for

DKK2 transcripts. Besides, among the set of mRNA genes displaying the top 5% PTc

scores and at least 2-folds ΔEx reduction (**Table 1**), 5 were not DE, and 3 of them only

showed suggestive DE, while the rest of DE mRNAs displayed a wide range of

significance (**Table S4**). The divergence between EISA and differential expression

results is also shown in **Fig. 1C**, where not all the mRNA genes highlighted by EISA

(**Fig. 1A**) were DE downregulated loci.

The miRNAs with the highest number of significant miRNA-mRNA interactions (in

terms of correlations between their expression levels) were ssc-miR-30a-3p and ssc-

miR-421-5p, which showed 9 and 8 significant miRNA-mRNA interactions, followed

by ssc-miR-148-3p with 4 significant interactions with mRNA genes showing

significant post-transcriptional signals (**Table S7**).

Genes showing post-transcriptional regulatory signals predominantly covary at the

exonic level

To further elucidate whether genes displaying the top 5% PTc scores are strong

candidates to be regulated by miRNAs according to *in silico* predictions (N = 21), we

evaluated the covariation patterns among them and with the whole set of 454 mRNA

genes DE between *AL-T0* and *AL-T2* gilts.

By calculating CES values (see Methods) for the 21 genes putatively targeted by DE

upregulated miRNAs, we obtained an estimation of the fold change in their observed

covariation with respect to other 435 DE mRNAs. CEs values were measured for both

their exonic and intronic fractions. Our analyses revealed that 19 out of these 21 genes

showed an increased covariation of approximately 2-folds in their exonic fractions when

compared to their intronic fractions (**Table S8**, **Fig. 1D**). When we iteratively analyzed

the observed fold change in covariation for random sets of genes (N = 1,000), they

displayed CES ≈ 1, indicative of no covariation (**Fig. 1D**). The observed CES

distributions of exonic, intronic and control sets were significantly different (*P*-value =

3.663E-06) after running non-parametric tests (**Fig. 1D**), thus supporting that the genes

displaying the top 5% PTc scores are probably repressed by DE upregulated miRNAs.

3.2. Studying post-transcriptional signals in fat metabolism using an independent

Duroc-Göttingen minipig population

After pre-processing and filtering of sequenced reads from adipocytes samples, we were

able to retrieve ~98.1 and ~0.87 million mRNA and small RNA reads per sample, and

~96.5% and ~73.4% of these reads mapped to annotated porcine mRNA and miRNA

genes, respectively. Differential expression analyses revealed a total of 299 DE

mRNAs, of which 52 were downregulated and 95 were upregulated, respectively (**Table**

S9). Regarding miRNAs, only one gene (ssc-miR-92b-3p) was significantly upregulated

in *lean* pigs, while 7 additional miRNAs showed suggestive differential expression (*P*-

value < 0.01), of which 4 were downregulated, and one (ssc-miR-92a) had the same

seed sequence as ssc-miR-92b-3p (**Table S10**).

After running EISA on the mRNA expression profiles for exonic and intronic fractions,

a total of 44 downregulated mRNAs in *lean* pigs displayed the top 5% PTc scores with

reduced ΔEx > 2-folds (**Table S11, Fig. 2A**). One of them (ENSSSCG00000016928)

did not have a properly annotated 3′-UTR so it was excluded from further analyses. The

whole set of mRNA genes from EISA results is available at **Table S12**. Among this set

of genes with high post-transcriptional signals, 13 appeared as significantly

downregulated (FC < -2; *q*-value < 0.05) and 9 had suggestive repression (*P*-value <

0.01) in the differential expression analysis considering their exonic fractions (**Table S9**

and **S11**). In this dataset, the sestrin 3 (*SESN3*) locus showed the second highest

negative PTc score, but was the only one among the rest of highlighted genes where a

significant post-transcriptional signal was observed (**Table S11**), i.e., equivalent to the *DKK2* gene in Duroc pigs, this was the only locus showing post-transcriptional regulation with no additional co-occurring transcriptional signal. Such strong downregulation was also evidenced in differential expression analyses, where it was the most significantly DE gene (**Table S9**).

From the set of downregulated genes, 25 out of the 43 mRNA genes analyzed for miRNA binding sites were classified as putative targets of the set of miRNAs upregulated in *lean* pigs (ssc-miR-92b-3p, ssc-miR-148a-3p, ssc-miR-204 and ssc-miR-214-3p; **Table S10**). Target prediction and context-based pruning of miRNA-mRNA interactions for these 43 mRNA genes made it possible to detect 8 8mer, 21 7mer-m8 and 24 7mer-A1 miRNA binding sites (**Table S13**) corresponding to the non-redundant seeds of selected upregulated miRNAs (N = 4) in *lean* minipigs (**Table S10**). The *SESN3* gene showed the highest number of predicted putative miRNA target sites in its 3´-UTR (**Table S13**).

Enrichment analyses for the set of putative miRNA target genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction (N = 25, **Table S11**) revealed no significant enrichment for the three types of miRNA target sites considered, although a slight increase when considering 8mer + 7mer-m8 binding sites and all three types together could be observed (**Fig. 2B**). Among this set of 25 genes, 7 appeared as significantly downregulated (FC < -2; *q*-value < 0.05) and 6 had suggestive repression (*P*-value < 0.01) in the differential expression analysis considering their exonic fractions (**Table S9** and **S11**, **Fig. 2C**).

In agreement with results obtained for the first studied skeletal muscle expression dataset, the exonic fraction of the mRNA genes putatively targeted by upregulated miRNAs in lean pigs showed approximately 2-folds significantly increased covariation

(*P*-value = 2.703E-02) with regard to their intronic fraction (**Fig. 2D**). Besides, 18 out

of these 25 mRNA genes showed an overall increased covariation in their exonic

fractions compared with their intronic fractions, expressed as the increment in their CES

values (ΔCES = exonic CES – intronic CES, **Table S14**).

Because adipose tissue samples were available, qPCR analyses were performed to

assess whether mRNAs among the top 5% negative PTc scores and with at least 2-folds

ΔEx reduction together with upregulated miRNAs displayed patterns of expression

consistent with those obtained in the RNA-seq and small RNA-seq experiments. To this

end, we selected five mRNAs (*LEP*, *OSBPL10*, *PRSS23*, *RNF157* and *SERPINE2*) and

3 miRNAs (ssc-miR-148a-30, ssc-miR-214-3p and ssc-miR-92b-3p) for qPCR

verification. All the analyzed mRNA genes showed a reduced expression in *lean* pigs

compared with their *obese* counterparts (**Fig. 2E**) and the *LEP* gene was the most

significantly downregulated gene (log$_2$FC = -1.953; *P*-value = 1.120E-03). This result

was in agreement with the strong downregulation observed in differential expression

analyses based on RNA-Seq data ((log$_2$FC = -1.957; *q*-value = 3.443E-03, **Table S9**).

With regard to miRNAs, the opposite pattern of expression was observed, with all the

three profiled miRNA genes being upregulated in *lean* pigs. Moreover, as reported in

Table S10, ssc-miR-92b-3p was the miRNA with the most significant upregulation as

evidenced in qPCR analyses (*P*-value = 3.57E-02, **Fig. 2F**).

4. Discussion

In this study, we undertook the inference of functional miRNA-mRNA interactions

using EISA for post-transcriptional signals prioritization, combined with target

prediction of upregulated miRNAs, as well as enrichment and covariation analyses.

We observed that the majority of mRNA genes with highly negative PTc scores, i.e., predominantly downregulated at their exonic fractions, also had a coordinated downregulatory effect of their intronic fractions, taken as a proxy of transcriptional repression. This was evidenced by the overall low significance of post-transcriptional signals within the mRNA genes with the top 5% negative PTc scores and reduced ΔEx in both analyzed experimental conditions. Only two genes (*DKK2* and *NAV2*) in *AL-T0* vs *AL-T2* and one gene (*SESN3*) in the *lean* vs *obese* contrasts showed significant PTc scores, revealing that, overall, a coordinated downregulatory effect at both transcriptional and post-transcriptional levels was present, which is in agreement with previous studies using EISA [14,15]. However, it is worth noting that we did not consider the significance of PTc scores as a relevant criterion for prioritizing putative post-transcriptionally downregulated genes, as these will appear as significant when the post-transcriptional activity is the only mechanism modulating the target gene expression profile. Only co-occurring yet opposite transcriptional and post-transcriptional events or single post-transcriptional signals would arise as significant, excluding genes with both coordinated downregulation at the transcriptional and post-transcriptional level.

We decided to use the intronic fraction of expressed mRNAs as a proxy of their transcriptional activity, a source of yet unspliced mRNA transcripts leading to the accumulation of intronic sequences prior to their debranching and degradation by exonucleases. This allows the use of RNA-seq datasets to apply EISA without the need of further experimental procedures, and it can also be applied to investigate transcriptional regulatory signals [15]. Previous reports have also explored the use of specific techniques to capture nascent mRNA transcripts before they are spliced [50–52], and these have been used to account for the transcriptional activity in a similar

approach to EISA [53]. Although more advanced methodologies measuring nascent

transcription of mRNAs might provide a better resolution for future experimental

designs [54], EISA would still be useful to explore the large amount of available RNA-

seq data where additional experiments are no longer possible.

Since the efficacy of miRNA-based regulation on mRNA targets depends on the context

of the target site within the 3'-UTR [43], we have described the usefulness of

introducing context-based filtering criteria for removing spurious *in silico*-predicted

target sites for miRNAs. Using enrichment analyses, we were able to link the

downregulated mRNAs at their exonic fractions to upregulated miRNAs that were

putatively targeting them and triggering their observed decay in differential expression

analyses. The influence of other non-DE highly expressed miRNAs or downregulated

miRNAs was discarded by the lack of predicted targeted mRNA genes with high post-

transcriptional downregulatory signals for such miRNAs. Overall, the increase in

enrichment significance shown for targeted mRNAs with post-transcriptional signals by

upregulated miRNAs revealed the ability of context-based filtering criteria to

discriminate and remove weak or false positive target sites located within unfavored

regions of the 3'-UTR. However, highly efficient target sites such as those of type 8mer,

although scarcer than 7mer-m8 sites, might still be functional even at unfavored

positions [43,55,56]. This may partially explain the relative lack of 8mer sites found in

the top post-transcriptionally regulated mRNA genes in both experimental setups.

Although miRNAs were the major post-transcriptional regulators that we considered in

this study, it is important to remark that other additional post-transcriptional

modifications and interactions might be responsible for the observed downregulation of

mRNAs. For instance, long non-coding RNAs [57], circular RNAs [58,59], RNA

methylation [60] or RNA binding proteins [61–63] can all act as post-transcriptional

regulators of targeted mRNAs. We further hypothesized that genes showing relevant post-transcriptional downregulatory effects might be regulated by the same set of significantly upregulated miRNAs, which could induce shared covariation in their expression profiles at the exonic level. On the contrary, their intronic fractions would be mainly unaffected, as introns would have been excised prior to any given miRNA-driven downregulation, if occurring. In this way, an increased gene covariation might be detectable within the sets of commonly targeted mRNA genes with relevant post-transcriptional signals at the exon but not at the intron level, as opposed to covariation events of these set of genes with the rest of DE genes. Our results revealed an increased degree of covariation between genes with high post-transcriptional signals at their exonic fractions, highlighting a putative coordinated downregulation by the set of significantly upregulated miRNAs.

From the analysis of top mRNA genes showing the strongest post-transcriptional downregulatory effects in fasted vs fed gilts, several biological functions putatively regulated by miRNAs were revealed. The *DKK2* gene was the one showing the highest negative PTc score, and its post-transcriptional regulatory signal was also significant, indicating that no additional coordinated transcriptional downregulation was found for this particular gene. Moreover, this gene also showed the strongest covariation difference in its exonic fraction compared with its intronic fraction. This consistent post-transcriptional regulatory effect might be explained by the presence of two miRNA target sites of type 8mer in its 3′-UTR for ssc-miR-421-5p and ssc-miR-30a-3p, two highly DE and upregulated miRNAs. Besides, ssc-miR-30e-3p, a miRNA sharing its seed and regulatory effect with ssc-miR-30a-3p, was also upregulated in fed (*AL-T2*) gilts, which would reinforce the repression of their targeted mRNA transcripts. The *DKK2* gene is a member of the dickkopf family that inhibits the Wnt signaling pathway

through its interaction with the LDL-receptor related protein 6 (LRP6). Its repression has been associated with reduced blood-glucose levels and improved glucose uptake [64], as well as with improved adipogenesis [65] and inhibition of aerobic glycolysis [66]. These results agreed with the increased glucose usage and triggered adipogenesis in muscle tissue after nutrient supply. Other additional relevant post-transcriptionally downregulated mRNA genes found by EISA were: (i) pyruvate dehydrogenase kinase 4 (*PDK4*), a mitochondrial enzyme that inhibits pyruvate to acetyl-CoA conversion and hinders glucose utilization promoting fatty acids oxidation in energy-deprived cells under fasting conditions [67,68]; (ii) interleukin 18 (*IL18*), involved in controlling energy homeostasis in the muscle by inducing AMP-activated protein kinase (AMPK) [69], a master metabolic regulator that is suppressed upon nutrient influx in cells [70]; (iii) nuclear receptor subfamily 4 group A member 3 (*NR4A3*), which activates both glycolytic and glycogenic factors [71], as well as β-oxidation in muscle cells [72]; (iv) acetylcholine receptor subunit α (*CHRNA1*) of muscle cells, that is linked to the inhibition of nicotine-dependent *STAT3* upregulation [73] that results in protection against insulin resistance in muscle [74]; (v) PBX homeobox 1 (*PBX1*), a regulator of adipocyte differentiation [75]; (vi) Tet methylcytosine dioxygenase 2 (*TET2*), linked to glucose-dependent AMPK phosphorylation [76]; and (vii) BTB domain and CNC homolog (*BACH2*), associated with mTOR complex 2 (mTORC2) glucose-dependent activation [77,78] and the repression of forkhead box protein O1 (*FOXO1*) [79] and *PDK4* in a coordinated manner [7,80]. Overall, the highlighted downregulated genes in the muscle of gilts after nutrient supply pointed towards a common regulatory function of miRNAs in modulating glucose uptake and energy homeostasis pathways of the skeletal myocytes.

Alternative post-transcriptional regulatory factors other than miRNAs might explain the presence of non-miRNA targets within the top post-transcriptional signals related to regulatory mechanisms not directly involved in energy homeostasis or glucose usage. For instance, three circadian clock-related mRNA genes that showed high post-transcriptional signals were the circadian associated repressor of transcription (*CIART*), period 1 (*PER1*) and salt inducible kinase 1 (*SIK1*), yet the first two were not detected as targets of DE miRNAs. As previously reported for this experimental design [22], the presence of several genes showing abundance differences might reflect a tight feedback interplay among them, where their expression and accumulation are coordinately regulated.

Regarding EISA results in RNA-seq profiles of adipocytes from *lean* vs *obese* Duroc-Göttingen minipigs, several of the mRNA genes that showed high post-transcriptional repression are involved in the regulation of lipid metabolism and energy homeostasis. The gene showing the highest post-transcriptional signal was the estrogen related receptor γ (*ESRRG*), which modulates oxidative metabolism and mitochondrial function in adipose tissue and imposes downregulation of adipocyte differentiation when repressed [81]. The second locus highlighted by EISA was sestrin 3 (*SESN3*), an activator of mTORC2 and PI3K/AKT signaling pathway [82] that protects against insulin resistance and promotes lipolysis when inhibited [83]. This gene showed the most significant downregulation in *lean* pigs, and gathered multiple putative binding sites for all the four upregulated miRNAs analyzed. Other genes showing significant post-transcriptional regulation were the following: The sterile α motif domain containing 4A (*SAMD4A*) that has been linked to the inhibition of preadipocyte differentiation and leanness phenotype in knockdown experiments [84,85]. The prostaglandin F2- receptor protein (PTGFR), for which overexpression has been

associated with hypertension and obesity risk [86], and its repression appears to

improve insulin sensitivity and glucose homeostasis [87]. The expression of serpin E1

and E2 (*SERPINE1*, *SERPINE2*) is linked to obesogenic states and diabetic symptoms

[88], while their inhibition improved glucose metabolism [89]. The serine protease 23

(PRSS23), which regulates insulin sensitivity and cytokine expression in adipose tissue,

and its downregulation confers protective effects against inflammation and reduced

fasting glucose level improving insulin resistance [90]. A high expression of ring finger

protein 157 (*RNF157*) has been linked to high fatness profiles and increased autophagy

in adipose tissue [91]. Silencing of ORP10 protein, encoded by the *OSBLP10* gene,

promotes low-density lipoprotein (LDL) synthesis and inhibits lipogenesis [92]. The

serum levels of glycosylphosphatidylinositol phospholipase 1 (GPLD1) are regulated by

insulin and glucose metabolism [93] and linked to the development of insulin resistance

and metabolic syndrome [94]. Overexpression of neutrophil cytosolic 2 (*NCF2*), the

gene showing the highest increase in covariation at the exonic fraction, was described in

obese humans [95]. The repression of RAP1 GTPase activating protein (RAP1GAP)

promotes RAP1 activity, which protects against obesity and insulin and glucose

resistance [96,97]. Finally, leptin production was also decreased in lean pigs. This key

adipokine is mainly produced in adipose tissue [98] and regulates appetite, energy

expenditure and body weight [99,100]. In summary, similar to what we found for

glucose metabolism and energy homeostasis in the first experimental population, we

were also able to describe a set of post-transcriptionally downregulated genes in Duroc-

Göttingen minipigs tightly related to adipose tissue metabolism regulation.

5. Conclusions

In this study we have implemented an exon/intron split analysis of RNA-seq data from

skeletal muscle and adipose tissue of pigs in two independent populations. In this way,

we were able to identify a set of genes putatively regulated by miRNAs at the post-

transcriptional level. Many of these genes were involved in the regulation of energy

homeostasis and development of both skeletal muscle and adipose tissues analyzed in

the present study, which were in agreement with metabolic modifications in response to

nutrient supply and the fatness profile of pigs. Overall, the use of EISA for post-

transcriptional signals prioritization combined with target prediction of upregulated

miRNAs, as well as enrichment analyses on the set of selected targeted genes and their

coordinated covariation, reinforced the usefulness of this approach to infer functional

miRNA-mRNA interactions using exonic and intronic fractions of commonly available

RNA-seq datasets.

Support

The present research work was funded by grants AGL2013-48742-C2-1-R and

AGL2013-48742-C2-2-R awarded by the Spanish Ministry of Economy and

Competitivity. E. Mármol-Sánchez was funded with a PhD fellowship FPU15/01733

awarded by the Spanish Ministry of Education and Culture (MECD). YRC is recipient

of a Ramon y Cajal fellowship (RYC2019-027244-I) from the Spanish Ministry of

Science and Innovation.

Data availability

641 The RNA-seq and small RNA-seq datasets from skeletal muscle tissue used in the

642 current study are available at the Sequence Read Archive (SRA) database with

643 BioProject codes PRJNA386796 and PRJNA595998, respectively. For the adipose

tissue samples, RNA-seq and small RNA-seq datasets are available at PRJNA563583 and PRJNA759240.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

The authors would like to thank the Department of Veterinary Animal Sciences in the Faculty of Health and Medical Sciences of the University of Copenhagen for providing sequencing data and their facilities and resources for qPCR experiments. We also acknowledge Selección Batallé S.A. for providing animal material and the support of the Spanish Ministry of Economy and Competitivity for the Center of Excellence Severo Ochoa 2020–2023 (CEX2019-000902-S) grant awarded to the Centre for Research in Agricultural Genomics (CRAG, Bellaterra, Spain). Thanks also to the CERCA Programme of the Generalitat de Catalunya for their support.

References

[1]   N.J. Martinez, A.J.M. Walhout, The interplay between transcription factors and microRNAs in genome-scale regulatory networks, BioEssays. 31 (2009) 435–445. doi:10.1002/bies.200800212.

[2]   D.P. Bartel, Metazoan microRNAs, Cell. 173 (2018) 20–51. doi:10.1016/j.cell.2018.03.006.

[3]   C.M.J. Mentzel, C. Anthon, M.J. Jacobsen, P. Karlskov-Mortensen, C.S. Bruun, C.B. Jørgensen, J. Gorodkin, S. Cirera, M. Fredholm, Gender and obesity specific microRNA expression in adipose tissue from lean and obese pigs., PLoS One. 10 (2015) e0131650. doi:10.1371/journal.pone.0131650.

669 [4] H. Han, S. Gu, W. Chu, W. Sun, W. Wei, X. Dang, Y. Tian, K. Liu, J. Chen,

670   miR-17-5p regulates differential expression of *NCOA3* in pig intramuscular and

671   subcutaneous adipose tissue, Lipids. 52 (2017) 939–949. doi:10.1007/s11745-

672   017-4288-4.

673 [5] W. Wei, B. Li, K. Liu, A. Jiang, C. Dong, C. Jia, J. Chen, H. Liu, W. Wu,

674   Identification of key microRNAs affecting drip loss in porcine longissimus dorsi

675   by RNA-Seq, Gene. 647 (2018) 276–282. doi:10.1016/j.gene.2018.01.005.

676 [6] S. Xie, X. Li, L. Qian, C. Cai, G. Xiao, S. Jiang, B. Li, T. Gao, W. Cui, L.L.

677   Guan, An integrated analysis of mRNA and miRNA in skeletal muscle from

678   myostatin-edited Meishan pigs, Genome. 62 (2019) 305–315. doi:10.1139/gen-

679   2018-0110.

680 [7] E. Mármol-Sánchez, Y. Ramayo-Caldas, R. Quintanilla, T.F. Cardoso, R.

681   González-Prendes, J. Tibau, M. Amills, Co-expression network analysis predicts

682   a key role of microRNAs in the adaptation of the porcine skeletal muscle to

683   nutrient supply, J. Anim. Sci. Biotechnol. 11 (2020) 10. doi:10.1186/s40104-019-

684   0412-z.

685 [8] M.A. Iqbal, A. Ali, F. Hadlich, M. Oster, H. Reyer, N. Trakooljul, V.

686   Sommerfeld, M. Rodehutscord, K. Wimmers, S. Ponsuksili, Dietary phosphorus

687   and calcium in feed affects miRNA profiles and their mRNA targets in jejunum

688   of two strains of laying hens, Sci. Rep. 11 (2021) 13534. doi:10.1038/S41598-

689   021-92932-3.

690 [9] A. Ali, E. Murani, F. Hadlich, X. Liu, K. Wimmers, S. Ponsuksili, In utero fetal

691   weight in pigs is regulated by microRNAs and their target genes, Genes. 12

692   (2021) 1264. doi:10.3390/genes12081264.

693 [10] D. Gaidatzis, L. Burger, M. Florescu, M.B. Stadler, Analysis of intronic and

exonic reads in RNA-seq data characterizes transcriptional and post-
transcriptional regulation, Nat. Biotechnol. 33 (2015) 722–729.
doi:10.1038/nbt.3269.

[11] A. Ameur, A. Zaghlool, J. Halvardson, A. Wetterbom, U. Gyllensten, L.
Cavelier, L. Feuk, Total RNA sequencing reveals nascent transcription and
widespread co-transcriptional splicing in the human brain., Nat. Struct. Mol.
Biol. 18 (2011) 1435–40. doi:10.1038/nsmb.2143.

[12] J.M. Gray, D.A. Harmin, S.A. Boswell, N. Cloonan, T.E. Mullen, J.J. Ling, N.
Miller, S. Kuersten, Y.-C. Ma, S.A. McCarroll, S.M. Grimmond, M. Springer,
SnapShot-Seq: a method for extracting genome-wide, in vivo mRNA dynamics
from a single total RNA sample., PLoS One. 9 (2014) e89673.
doi:10.1371/journal.pone.0089673.

[13] G.-J. Hendriks, D. Gaidatzis, F. Aeschimann, H. Großhans, Extensive oscillatory
gene expression during C. elegans larval development., Mol. Cell. 53 (2014)
380–92. doi:10.1016/j.molcel.2013.12.013.

[14] J. Cursons, K.A. Pillman, K.G. Scheer, P.A. Gregory, M. Foroutan, S. Hediyeh-
Zadeh, J. Toubia, E.J. Crampin, G.J. Goodall, C.P. Bracken, M.J. Davis,
Combinatorial targeting by microRNAs co-ordinates post-transcriptional control
of EMT, Cell Syst. 7 (2018) 77-91.e7. doi:10.1016/j.cels.2018.05.019.

[15] K.A. Pillman, K.G. Scheer, E. Hackett-Jones, K. Saunders, A.G. Bert, J. Toubia,
H.J. Whitfield, S. Sapkota, L. Sourdin, H. Pham, T.D. Le, J. Cursons, M.J. Davis,
P.A. Gregory, G.J. Goodall, C.P. Bracken, Extensive transcriptional responses
are co-ordinated by microRNAs as revealed by Exon–Intron Split Analysis
(EISA), Nucleic Acids Res. 47 (2019) 8606-8619. doi:10.1093/nar/gkz664.

[16]   D. Pérez-Montarelo, A. Fernández, C. Barragán, J.L. Noguera, J.M. Folch, M.C.

720   Rodríguez, C. Óvilo, L. Silió, A.I. Fernández, Transcriptional characterization of

721   porcine leptin and leptin receptor genes, PLoS One. 8 (2013) e66398.

722   doi:10.1371/journal.pone.0066398.

723   [17]   C. Ovilo, R. Benítez, A. Fernández, Y. Núñez, M. Ayuso, A.I. Fernández, C.

724   Rodríguez, B. Isabel, A.I. Rey, C. López-Bote, L. Silió, Longissimus dorsi

725   transcriptome analysis of purebred and crossbred Iberian pigs differing in muscle

726   characteristics., BMC Genomics. 15 (2014) 413. doi:10.1186/1471-2164-15-413.

727   [18]   A. Puig-Oliveras, Y. Ramayo-Caldas, J. Corominas, J. Estellé, D. Pérez-

728   Montarelo, N.J. Hudson, J. Casellas, J.M. Folch, M. Ballester, Differences in

729   muscle transcriptome among pigs phenotypically extreme for fatty acid

730   composition., PLoS One. 9 (2014) e99720. doi:10.1371/journal.pone.0099720.

731   [19]   C.M. Pilcher, C.K. Jones, M. Schroyen, A.J. Severin, J.F. Patience, C.K. Tuggle,

732   J.E. Koltes, Transcript profiles in longissimus dorsi muscle and subcutaneous

733   adipose tissue: A comparison of pigs with different postweaning growth rates, J.

734   Anim. Sci. 93 (2015) 2134–2143. doi:10.2527/jas.2014-8593.

735   [20]   M. Ayuso, A. Fernández, Y. Núñez, R. Benítez, B. Isabel, A.I. Fernández, A.I.

736   Rey, A. González-Bulnes, J.F. Medrano, Á. Cánovas, C.J. López-Bote, C. Óvilo,

737   Developmental stage, muscle and genetic type modify muscle transcriptome in

738   pigs: Effects on gene expression and regulatory factors involved in growth and

739   metabolism, PLoS One. 11 (2016) e0167858. doi:10.1371/journal.pone.0167858.

740   [21]   T.F. Cardoso, A. Cánovas, O. Canela-Xandri, R. González-Prendes, M. Amills,

741   R. Quintanilla, RNA-seq based detection of differentially expressed genes in the

742   skeletal muscle of Duroc pigs with distinct lipid profiles, Sci. Rep. 7 (2017)

743   40005. doi:10.1038/srep40005.

[22] T.F. Cardoso, R. Quintanilla, J. Tibau, M. Gil, E. Mármol-Sánchez, O. González-

Rodríguez, R. González-Prendes, M. Amills, Nutrient supply affects the mRNA

expression profile of the porcine skeletal muscle, BMC Genomics. 18 (2017)

603. doi:10.1186/s12864-017-3986-x.

[23] J. Horodyska, K. Wimmers, H. Reyer, N. Trakooljul, A.M. Mullen, P.G. Lawlor,

R.M. Hamill, RNA-seq of muscle from pigs divergent in feed efficiency and

product quality identifies differences in immune response, growth, and

macronutrient and connective tissue metabolism, BMC Genomics. 19 (2018)

791. doi:10.1186/s12864-018-5175-y.

[24] R. Benítez, N. Trakooljul, Y. Núñez, B. Isabel, E. Murani, E. De Mercado, E.

Gómez-Izquierdo, J. García-Casco, C. López-Bote, K. Wimmers, C. Óvilo,

Breed, diet, and interaction effects on adipose tissue transcriptome in iberian and

duroc pigs fed different energy sources, Genes. 10 (2019) 589.

doi:10.3390/genes10080589.

[25] M. Ballester, M. Amills, O. González-Rodríguez, T.F. Cardoso, M. Pascual, R.

González-Prendes, N. Panella-Riera, I. Díaz, J. Tibau, R. Quintanilla, Role of

AMPK signalling pathway during compensatory growth in pigs, BMC Genomics.

19 (2018) 682. doi:10.1186/s12864-018-5071-5.

[26] L.J.A. Kogelman, H.N. Kadarmideen, T. Mark, P. Karlskov-Mortensen, C.S.

Bruun, S. Cirera, M.J. Jacobsen, C.B. Jørgensen, M. Fredholm, An F2 pig

resource population as a model for genetic studies of obesity and obesity-related

diseases in humans: Design and genetic parameters, Front. Genet. 4 (2013) 29.

doi:10.3389/fgene.2013.00029.

[27] S.D. Pant, P. Karlskov-Mortensen, M.J. Jacobsen, S. Cirera, L.J.A. Kogelman,

C.S. Bruun, T. Mark, C.B. Jørgensen, N. Grarup, E.V.R. Appel, E.A.A.

Galjatovic, T. Hansen, O. Pedersen, M. Guerin, T. Huby, P. Lesnik, T.H.E.

Meuwissen, H.N. Kadarmideen, M. Fredholm, Comparative analyses of QTLs

influencing obesity and metabolic phenotypes in pigs and humans, PLoS One. 10

(2015) e0137356. doi:10.1371/journal.pone.0137356.

[28] M.J. Jacobsen, J.H. Havgaard, C. Anthon, C.M.J. Mentzel, S. Cirera, P.M.

Krogh, S. Pundhir, P. Karlskov-Mortensen, C.S. Bruun, P. Lesnik, M. Guerin, J.

Gorodkin, C.B. Jørgensen, M. Fredholm, R. Barrès, Epigenetic and

transcriptomic characterization of pure adipocyte fractions from obese pigs

identifies candidate pathways controlling metabolism, Front. Genet. 10 (2019)

1268. doi:10.3389/fgene.2019.01268.

[29] P. Decaunes, D. Estève, A. Zakaroff-Girard, C. Sengenès, J. Galitzky, A.

Bouloumié, Adipose-derived stromal cells: cytokine expression and immune cell

contaminants., Methods Mol. Biol. 702 (2011) 151–161. doi:10.1007/978-1-

61737-960-4_12.

[30] M. Martin, Cutadapt removes adapter sequences from high-throughput

sequencing reads, EMBnet.Journal. 17 (2011) 10. doi:10.14806/ej.17.1.200.

[31] A. Warr, N. Affara, B. Aken, H. Beiki, D.M. Bickhart, K. Billis, W. Chow, L.

Eory, H.A. Finlayson, P. Flicek, C.G. Girón, D.K. Griffin, R. Hall, G. Hannum,

T. Hourlier, K. Howe, D.A. Hume, O. Izuogu, K. Kim, S. Koren, H. Liu, N.

Manchanda, F.J. Martin, D.J. Nonneman, R.E. O'Connor, A.M. Phillippy, G.A.

Rohrer, B.D. Rosen, L.A. Rund, C.A. Sargent, L.B. Schook, S.G. Schroeder, A.S.

Schwartz, B.M. Skinner, R. Talbot, E. Tseng, C.K. Tuggle, M. Watson, T.P.L.

Smith, A.L. Archibald, An improved pig reference genome sequence to enable

pig genetics and genomics research, Gigascience. 9 (2020) 1–14.

doi:10.1093/gigascience/giaa051.

[32]   D. Kim, J.M. Paggi, C. Park, C. Bennett, S.L. Salzberg, Graph-based genome

795   alignment and genotyping with HISAT2 and HISAT-genotype, Nat. Biotechnol.

796   37 (2019) 907–915. doi:10.1038/s41587-019-0201-4.

797   [33]   B. Langmead, C. Trapnell, M. Pop, S.L. Salzberg, Ultrafast and memory-efficient

798   alignment of short DNA sequences to the human genome, Genome Biol. 10

799   (2009) R25. doi:10.1186/gb-2009-10-3-r25.

800   [34]   M. Lawrence, W. Huber, H. Pagès, P. Aboyoun, M. Carlson, R. Gentleman, M.T.

801   Morgan, V.J. Carey, Software for computing and annotating genomic ranges,

802   PLoS Comput. Biol. 9 (2013) e1003118. doi:10.1371/journal.pcbi.1003118.

803   [35]   Y. Liao, G.K. Smyth, W. Shi, The R package Rsubread is easier, faster, cheaper

804   and better for alignment and quantification of RNA sequencing reads, Nucleic

805   Acids Res. 47 (2019) e47–e47. doi:10.1093/nar/gkz114.

806   [36]   Y. Liao, G.K. Smyth, W. Shi, featureCounts: an efficient general purpose

807   program for assigning sequence reads to genomic features, Bioinformatics. 30

808   (2014) 923–930. doi:10.1093/bioinformatics/btt656.

809   [37]   M.D. Robinson, D.J. McCarthy, G.K. Smyth, edgeR: a Bioconductor package for

810   differential expression analysis of digital gene expression data., Bioinformatics.

811   26 (2010) 139–40. doi:10.1093/bioinformatics/btp616.

812   [38]   M.D. Robinson, A. Oshlack, A scaling normalization method for differential

813   expression analysis of RNA-seq data, Genome Biol. 11 (2010) R25.

814   doi:10.1186/gb-2010-11-3-r25.

815   [39]   Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: a practical and

816   powerful approach to multiple testing, J. R. Stat. Soc. Ser. B. 57 (1995) 289–300.

817   doi:10.2307/2346101.

[40]   Y. Guo, J. Liu, S.J. Elfenbein, Y. Ma, M. Zhong, C. Qiu, Y. Ding, J. Lu,

Characterization of the mammalian miRNA turnover landscape, Nucleic Acids

Res. 43 (2015) 2326–2341. doi:10.1093/nar/gkv057

[41]   A. Kozomara, M. Birgaoanu, S. Griffiths-Jones, MiRBase: From microRNA

sequences to function, Nucleic Acids Res. 47 (2019) D155–D162.

doi:10.1093/nar/gky1141.

[42]   A. Marco, SeedVicious: Analysis of microRNA target and near-target sites, PLoS

One. 13 (2018) e0195532. doi:10.1371/journal.pone.0195532.

[43]   A. Grimson, K.K.H. Farh, W.K. Johnston, P. Garrett-Engele, L.P. Lim, D.P.

Bartel, MicroRNA targeting specificity in mammals: determinants beyond seed

pairing, Mol. Cell. 27 (2007) 91–105. doi:10.1016/j.molcel.2007.06.017.

[44]   R.C. Friedman, K.K.H. Farh, C.B. Burge, D.P. Bartel, Most mammalian mRNAs

are conserved targets of microRNAs, Genome Res. 19 (2009) 92–105.

doi:10.1101/gr.082701.108.

[45]   A. Reverter, E.K.F. Chan, Combining partial correlation and an information

theory approach to the reversed engineering of gene co-expression networks,

Bioinformatics. 24 (2008) 2491–2497. doi:10.1093/bioinformatics/btn482.

[46]   N.S. Watson-Haigh, H.N. Kadarmideen, A. Reverter, PCIT: an R package for

weighted gene co-expression networks based on partial correlation and

information theory approaches, Bioinformatics. 26 (2010) 411–413.

doi:10.1093/bioinformatics/btp674.

[47]   M. Tarbier, S.D. Mackowiak, J. Frade, S. Catuara-Solarz, I. Biryukova, E. Gelali,

D.B. Menéndez, L. Zapata, S. Ossowski, M. Bienko, C.J. Gallant, M.R.

Friedländer, Nuclear gene proximity and protein interactions shape transcript

covariations in mammalian single cells, Nat. Commun. 11 (2020) 5445.

doi:10.1038/s41467-020-19011-5.

[48] H.B. Mann, D.R. Whitney, On a test of whether one of two random variables is

stochastically larger than the other, Ann. Math. Stat. 18 (1947) 50–60.

doi:10.1214/aoms/1177730491.

[49] A.B. Nygard, C.B. Jørgensen, S. Cirera, M. Fredholm, Selection of reference

genes for gene expression studies in pig tissues using SYBR green qPCR, BMC

Mol. Biol. 8 (2007) 67. doi:10.1186/1471-2199-8-67.

[50] L.J. Core, J.J. Waterfall, J.T. Lis, Nascent RNA sequencing reveals widespread

pausing and divergent initiation at human promoters, Science. 322 (2008) 1845–

1848. doi:10.1126/science.1162228.

[51] D.B. Mahat, H. Kwak, G.T. Booth, I.H. Jonkers, C.G. Danko, R.K. Patel, C.T.

Waters, K. Munson, L.J. Core, J.T. Lis, Base-pair resolution genome-wide

mapping of active RNA polymerasesusing precision nuclear run-on (PRO-seq),

Nat. Protoc. 11 (2016) 1455. doi:10.1038/nprot.2016.086.

[52] A. Blumberg, Y. Zhao, Y.-F. Huang, N. Dukler, E.J. Rice, A.G. Chivu, K.

Krumholz, C.G. Danko, A. Siepel, Characterizing RNA stability genome-wide

through combined analysis of PRO-seq and RNA-seq data, BMC Biol. 19 (2021)

30. doi:10.1186/s12915-021-00949-x.

[53] R.K. Patel, J.D. West, Y. Jiang, E.A. Fogarty, A. Grimson, Robust partitioning of

microRNA targets from downstream regulatory changes, Nucleic Acids Res. 48

(2020) 9724–9746. doi:10.1093/nar/gkaa687.

[54] E.M. Wissink, A. Vihervaara, N.D. Tippens, J.T. Lis, Nascent RNA analyses:

tracking transcription and its regulation, Nat. Rev. Gen. 20 (2019) 705-723.

doi:10.1038/s41576-019-0159-6.

[55] R. Denzler, S.E. McGeary, A.C. Title, V. Agarwal, D.P. Bartel, M. Stoffel,

Impact of microRNA levels, target-site complementarity, and cooperativity on

competing endogenous RNA-regulated gene expression, Mol. Cell. 64 (2016)

565-579. doi:10.1016/j.molcel.2016.09.027.

[56]   S.E. McGeary, K.S. Lin, C.Y. Shi, T.M. Pham, N. Bisaria, G.M. Kelley, D.P.

Bartel, The biochemical basis of microRNA targeting efficacy, Science. 366

(2019) eaav1741. doi:10.1126/science.aav1741.

[57]   R.Z. He, D.X. Luo, Y.Y. Mo, Emerging roles of lncRNAs in the post-

transcriptional   regulation   in   cancer,   Genes   Dis.   6   (2019)   6–15.

doi:10.1016/j.gendis.2019.01.003.

[58]   S. Memczak, M. Jens, A. Elefsinioti, F. Torti, J. Krueger, A. Rybak, L. Maier,

S.D. Mackowiak, L.H. Gregersen, M. Munschauer, A. Loewer, U. Ziebold, M.

Lanthaler, C. Kocks, F. le Noble, N. Rajewsky, Circular RNAs are a large class

of animal RNAs with regulatory potency, Nature. 495 (2013) 333–338.

doi:10.1038/nature11928.

[59]   P.G. Maass, P. Glažar, S. Memczak, G. Dittmar, I. Hollfinger, L. Schreyer, A.V.

Sauer, O. Toka, A. Aiuti, F.C. Luft, N. Rajewsky, A map of human circular

RNAs in clinically relevant tissues, J. Mol. Med. (Berl). 95 (2017) 1179–1189.

doi:10.1007/S00109-017-1582-9.

[60]   B.S. Zhao, I.A. Roundtree, C. He, Post-transcriptional gene regulation by mRNA

modifications,   Nat.   Rev.   Mol.   Cell   Biol.   18   (2017)   31.

doi:10.1038/nrm.2016.132.

[61]   T. Glisovic, J.L. Bachorik, J. Yong, G. Dreyfuss, RNA-binding proteins and

post-transcriptional   gene   regulation,   FEBS   Lett.   582   (2008)   1977–1986.

doi:10.1016/j.febslet.2008.03.004.

[62]   M.W. Hentze, A. Castello, T. Schwarzl, T. Preiss, A brave new world of RNA-

binding   proteins,   Nat.   Rev.   Mol.   Cell   Biol.   19   (2018)   327–341.

doi:10.1038/nrm.2017.130.

[63]   A. Velázquez-Cruz, B. Baños-Jaime, A. Díaz-Quintana, M.A.D. la Rosa, I. Díaz-

Moreno, Post-translational control of RNA-binding proteins and disease-related

dysregulation, Front. Mol. Biosci. 8 (2021) 658852.

doi:10.3389/fmolb.2021.658852.

[64]   X. Li, J. Shan, W. Chang, I. Kim, J. Bao, H.-J. Lee, X. Zhang, V.T. Samuel, G.I.

Shulman, D. Liu, J.J. Zheng, D. Wu, Chemical and genetic evidence for the

involvement of Wnt antagonist Dickkopf2 in regulation of glucose metabolism,

Proc. Natl. Acad. Sci. 109 (2012) 11402–11407. doi:10.1073/pnas.1205015109.

[65]   J. Yang, B. yin Shi, Dickkopf (Dkk)-2 is a beige fat-enriched adipokine to

regulate adipogenesis, Biochem. Biophys. Res. Commun. 548 (2021) 211–216.

doi:10.1016/j.bbrc.2021.02.068.

[66]   F. Deng, R. Zhou, C. Lin, S. Yang, H. Wang, W. Li, K. Zheng, W. Lin, X. Li, X.

Yao, M. Pan, L. Zhao, Tumor-secreted dickkopf2 accelerates aerobic glycolysis

and promotes angiogenesis in colorectal cancer, Theranostics. 9 (2019) 1001-

1014. doi:10.7150/thno.30056.

[67]   J.Y. Jeong, N.H. Jeoung, K.-G. Park, I.-K. Lee, Transcriptional regulation of

pyruvate dehydrogenase kinase, Diabetes Metab. J. 36 (2012) 328–35.

doi:10.4093/dmj.2012.36.5.328.

[68]   S. Zhang, M.W. Hulver, R.P. McMillan, M.A. Cline, E.R. Gilbert, The pivotal

role of pyruvate dehydrogenase kinases in metabolic flexibility, Nutr. Metab. 11

(2014) 10. doi:10.1186/1743-7075-11-10.

[69]   B. Lindegaard, V.B. Mathews, C. Brandt, P. Hojman, T.L. Allen, E. Estevez,

M.J. Watt, C.R. Bruce, O.H. Mortensen, S. Syberg, C. Rudnika, J. Abildgaard, H.

Pilegaard, J. Hidalgo, S. Ditlevsen, T.J. Alsted, A.N. Madsen, B.K. Pedersen,

M.A. Febbraio, Interleukin-18 activates skeletal muscle AMPK and reduces

weight gain and insulin resistance in mice, Diabetes. 62 (2013) 3064–3074.

doi:10.2337/DB12-1095.

[70] P. Jiang, L. Ren, L. Zhi, Z. Yu, F. Lv, F. Xu, W. Peng, X. Bai, K. Cheng, L,

Quang, X. Zhang, X. Wang, Y. Zhang, D. Yang, X. Hu, R.P. Xiao, Negative

regulation of AMPK signaling by high glucose via E3 ubiquitin ligase MG53,

Mol. Cell. 81 (2021) 629-637.e5. doi:10.1016/j.molcel.2020.12.008.

[71] C. Zhang, B. Zhang, X. Zhang, G. Sun, X. Sun, Targeting orphan nuclear

receptors NR4As for energy homeostasis and diabetes, Front. Pharmacol. 11

(2020) 587457. doi:10.3389/fphar.2020.587457.

[72] M.A. Pearen, J.M. Goode, R.L. Fitzsimmons, N.A. Eriksson, G.P. Thomas, G.J.

Cowin, S.C. Wang, Z.K. Tuong, G.E. Muscat, Transgenic muscle-specific Nor-1

expression regulates multiple pathways that effect adiposity, metabolism, and

endurance, Mol. Endocrinol. 27 (2013) 1897–1917. doi:10.1210/me.2013-1205.

[73] S. Xu, H. Ni, H. Chen, Q. Dai, The interaction between STAT3 and nAChRα1

interferes with nicotine-induced atherosclerosis via Akt/mTOR signaling

cascade, Aging. 11 (2019) 8120–8138. doi:10.18632/aging.102296.

[74] L. Zhang, Z. Chen, Y. Wang, D.J. Tweardy, W.E. Mitch, Stat3 activation induces

insulin resistance via a muscle-specific E3 ubiquitin ligase Fbxo40, Am. J.

Physiol. Endocrinol. Metab. 318 (2020) E625–E635.

doi:10.1152/ajpendo.00480.2019.

[75] M.C. Monteiro, M. Sanyal, M.L. Cleary, C. Sengenès, A. Bouloumié, C. Dani, N.

Billon, PBX1: a novel stage-specific regulator of adipocyte development, Stem

Cells. 29 (2011) 1837–1848. doi:10.1002/stem.737.

[76] D. Wu, D. Hu, H. Chen, G. Shi, I.S. Fetahu, F. Wu, K. Rabidou, R. Fang, L. Tan,

S. Xu, H. Liu, C. Argueta, L. Zhang, F. Mao, G. Yan, J. Chen, Z. Dong, R. Lv,

Y. Xu, M. Wang, Y. Ye, S. Zhang, D. Duquette, S. Geng, C. Yin, C.G. Lian, G.F.

Murphy, G.K. Adler, R. Garg, L. Lynch, P. Yang, Y. Li, F. Lan, J. Fan, Y. Shi,

Y.G. Shi, Glucose-regulated phosphorylation of TET2 by AMPK reveals a

pathway linking diabetes to cancer, Nature. 559 (2018) 637–641.

doi:10.1038/s41586-018-0350-5.

[77]   T. Tamahara, K. Ochiai, A. Muto, Y. Kato, N. Sax, M. Matsumoto, T. Koseki, K.

Igarashi, The mTOR-Bach2 cascade controls cell cycle and class switch

recombination during B cell differentiation, Mol. Cell. Biol. 37 (2017) e00418-

17. doi:10.1128/mcb.00418-17.

[78]   G. Leprivier, B. Rotblat, How does mTOR sense glucose starvation? AMPK is

the usual suspect, Cell Death Discov. 6 (2020) 27. doi:10.1038/S41420-020-

0260-9.

[79]   A. Itoh-Nakadai, M. Matsumoto, H. Kato, J. Sasaki, Y. Uehara, Y. Sato, R.

Ebina-Shibuya, M. Morooka, R. Funayama, K. Nakayama, K. Ochiai, A. Muto,

K. Igarashi, A Bach2-Cebp gene regulatory network for the commitment of

multipotent hematopoietic progenitors, Cell Rep. 18 (2017) 2401–2414.

doi:10.1016/j.celrep.2017.02.029.

[80]   K. Gopal, B. Saleme, R. Al Batran, H. Aburasayn, A. Eshreif, K.L. Ho, W.K.

Ma, M. Almutairi, F. Eaton, M. Gandhi, E.A. Park, G. Sutendra, J.R. Ussher,

FoxO1 regulates myocardial glucose oxidation rates via transcriptional control of

pyruvate dehydrogenase kinase 4 expression, Am. J. Physiol. Circ. Physiol. 313

(2017) H479–H490. doi:10.1152/ajpheart.00191.2017.

[81]   M. Kubo, N. Ijichi, K. Ikeda, K. Horie-Inoue, S. Takeda, S. Inoue, Modulation of

adipogenesis-related gene expression by estrogen-related receptor γ during

adipocytic differentiation, Biochim. Biophys. Acta - Gene Regul. Mech. 1789

(2009) 71–77. doi:10.1016/j.bbgram.2008.08.012.

[82]   X. Huang, G. Liu, J. Guo, Z. Su, The PI3K/AKT pathway in obesity and type 2

diabetes, Int. J. Biol. Sci. 14 (2018) 1483. doi:10.7150/ijbs.27173.

[83]   R. Tao, X. Xiong, S. Liangpunsakul, X.C. Dong, Sestrin 3 protein enhances

hepatic insulin sensitivity by direct activation of the mTORC2-Akt signaling,

Diabetes. 64 (2015) 1211–1223. doi:10.2337/db14-0539.

[84]   Z. Chen, W. Holland, J.M. Shelton, A. Ali, X. Zhan, S. Won, W. Tomisato, C.

Liu, X. Li, E.M.Y. Moresco, B. Beutler, Mutation of mouse *Samd4* causes

leanness, myopathy, uncoupled mitochondrial respiration, and dysregulated

mTORC1 signaling, Proc. Natl. Acad. Sci. 111 (2014) 7367–7372.

doi:10.1073/pnas.1406511111.

[85]   Y. Liu, H. Liu, Y. Li, R. Mao, H. Yang, Y. Zhang, Y. Zhang, P. Guo, D. Zhan, T.

Zhang, Circular RNA *SAMD4A* controls adipogenesis in obesity through the

miR-138-5p/EZH2   axis,   Theranostics.   10   (2020)   4705–4719.

doi:10.7150/thno.42417.

[86]   B. Xiao, S.M. Gu, M.J. Li, J. Li, B. Tao, Y. Wang, Y. Wang, S. Zuo, Y. Shen, Y.

Yu, D. Chen, G. Chen, D. Kong, J. Tang, Q. Liu, D.R. Cheng, Y. Liu, S. Alberti,

M. Dovizio, R. Landolfi, L. Mucci, P.-Z. Miao, P. Gao, D.-L. Zhu, J. Wang, B.

Li, P. Patrignani, Y. Yu, Rare SNP rs12731181 in the miR-590-3p target site of

the Prostaglandin F2α receptor gene confers risk for essential hypertension in the

Han Chinese population, Arterioscler. Thromb. Vasc. Biol. 35 (2015) 1687–

1695. doi:10.1161/atvbaha.115.305445.

[87]   Y. Wang, S. Yan, B. Xiao, S. Zuo, Q. Zhang, G. Chen, Y. Yu, D. Chen, Q. Liu,

Y. Liu, Y. Shen, Y. Yu, Prostaglandin F 2α facilitates hepatic glucose production

through CaMKIIγ/p38/FOXO1 signaling pathway in fasting and obesity, Diabetes. 67 (2018) 1748–1760. doi:10.2337/db17-1521.

[88] P. Kaur, M.D. Reis, G.R. Couchman, S.N. Forjuoh, J.F. Greene, Jr, A. Asea, SERPINE 1 links obesity and diabetes: a pilot study, J. Proteomics Bioinform. 3 (2010) 191. doi:10.4172/jpb.1000139.

[89] G.M. Coudriet, J. Stoops, A.V. Orr, B. Bhushan, K. Koral, S. Lee, D.M. Previte, H.H. Dong, G.K. Michalopoulos, W.M. Mars, J.D. Piganelli, A Noncanonical role for plasminogen activator inhibitor type 1 in obesity-induced diabetes, Am. J. Pathol. 189 (2019) 1413–1422. doi:10.1016/j.ajpath.2019.04.004.

[90] C.-S. Kuo, J.-S. Chen, L.-Y. Lin, G.W. Schmid-Schönbein, S. Chien, P.-H. Huang, J.-W. Chen, S.-J. Lin, Inhibition of serine protease activity protects against high fat diet-induced inflammation and insulin resistance, Sci. Reports. 10 (2020) 1–11. doi:10.1038/s41598-020-58361-4.

[91] J. Kosacka, M. Nowicki, S. Paeschke, P. Baum, M. Blüher, N. Klö ting, Up-regulated autophagy: as a protective factor in adipose tissue of WOKW rats with metabolic syndrome, Diabetol. Metab. Syndr. 10 (2018) 13. doi:10.1186/S13098-018-0317-6.

[92] J. Perttilä, K. Merikanto, J. Naukkarinen, I. Surakka, N.W. Martin, K. Tanhuanpää, V. Grimard, M.-R. Taskinen, C. Thiele, V. Salomaa, A. Jula, M. Perola, I. Virtanen, L. Peltonen, V.M. Olkkonen, OSBPL10, a novel candidate gene for high triglyceride trait in dyslipidemic Finnish subjects, regulates cellular lipid metabolism, J. Mol. Med. 2009 878. 87 (2009) 825–835. doi:10.1007/S00109-009-0490-Z.

[93] R.F. Bowen, N.S. Raikwar, L.K. Olson, M.A. Deeg, Glucose and insulin regulate glycosylphosphatidylinositol-specific phospholipase D expression in islet beta

cells, Metabolism. 50 (2001) 1489–1492. doi:10.1053/meta.2001.28087.

[94]   S. Ussar, O. Bezy, M. Blüher, C.R. Kahn, Glypican-4 enhances insulin signaling

via interaction with the insulin receptor and serves as a novel adipokine,

Diabetes. 61 (2012) 2289–2298. doi:10.2337/db11-1395.

[95]   J. Xu, X. Bao, Z. Peng, L. Wang, L. Du, W. Niu, Y. Sun, Comprehensive

analysis of genome-wide DNA methylation across human polycystic ovary

syndrome ovary granulosa cell, Oncotarget. 7 (2016) 27899.

doi:10.18632/oncotarget.8544.

[96]   F. Yeung, C.M. Ramírez, P.A. Mateos-Gomez, A. Pinzaro, G. Ceccarini, S.

Kabir, C. Fernández-Hernando, A. Sfeir, Nontelomeric role for Rap1 in

regulating metabolism and protecting against obesity, Cell Rep. 3 (2013) 1847–

1856. doi:10.1016/j.celrep.2013.05.032.

[97]   P. Martínez, G. Gómez-López, F. García, E. Mercken, S. Mitchell, J.M. Flores,

R. de Cabo, M.A. Blasco, RAP1 protects from obesity through its extratelomeric

role regulating gene expression, Cell Rep. 3 (2013) 2059–2074.

doi:10.1016/j.celrep.2013.05.030.

[98]   W. Yang, T. Kelly, J. He, Genetic epidemiology of obesity, Epidemiol. Rev. 29

(2007) 49–61. doi:10.1093/epirev/mxm004.

[99]   Y. Zhou, L. Rui, Leptin signaling and leptin resistance, Front. Med. 7 (2013)

207–222. doi:10.1007/s11684-013-0263-5.

[100] A.G. Izquierdo, A.B. Crujeiras, F.F. Casanueva, M.C. Carreira, Leptin, obesity,

and leptin resistance: where are we 25 years later?, Nutrients. 11 (2019) 2704.

doi:10.3390/nu11112704.

1042

Table 1: mRNA genes with the top 5% post-transcriptional (PTc) scores and at least 2-folds exonic fraction (ΔEx) reduction (equivalent to -1 in the $\log_2$ scale) from *gluteus medius* skeletal muscle expression profiles of fasting (*AL-T0*, N = 11) and fed (*AL-T2*, N = 12) Duroc gilts.

ID	Gene	$\log_2$FC	ΔEx	PT_c	*P*-value	*q*-value	DE	miRNA target
ENSSSCG00000032094	*DKK2*	-2.010	-1.431	-4.738	1.654E-05	3.830E-03	•	x
ENSSSCG00000015334	*PDK4*	-2.108	-5.250	-4.698	4.693E-03	1.330E-01	x	x
ENSSSCG00000015037	*IL18*	-1.655	-1.191	-3.682	4.787E-03	1.340E-01	•	x
ENSSSCG00000005385	*NR4A3*	-1.337	-3.082	-3.646	4.038E-02	4.098E-01	x	x
ENSSSCG00000003766	*DNAJB4*	-1.391	-1.008	-3.348	8.358E-03	1.905E-01		x
ENSSSCG00000015969	*CHRNA1*	-1.561	-1.339	-3.341	2.606E-03	9.406E-02	x	x
ENSSSCG00000039419	*SLCO4A1*	-1.055	-2.279	-3.180	2.820E-02	3.544E-01	x	x
ENSSSCG00000049158		-1.107	-1.096	-3.164	3.182E-02	3.735E-01		x
ENSSSCG00000004347	*FBXL4*	-1.298	-1.126	-3.133	1.422E-03	6.520E-02	x	x
ENSSSCG00000004979	*MYO9A*	-1.239	-1.003	-3.043	7.296E-03	1.731E-01		x
ENSSSCG00000013351	*NAV2*	-1.163	-1.196	-2.863	2.605E-04	2.301E-02	x	x
ENSSSCG00000032741	*TBC1D9*	-0.913	-1.061	-2.736	1.534E-02	2.583E-01	•	x
ENSSSCG00000031728	*ABRA*	-1.238	-1.393	-2.704	1.295E-03	6.116E-02	x	x
ENSSSCG00000006331	*PBX1*	-0.891	-1.039	-2.480	1.135E-02	2.177E-01	x	x
ENSSSCG00000035037	*SIK1*	-1.357	-1.289	-2.475	3.999E-03	1.212E-01	x	x
ENSSSCG00000038374	*CIART*	-1.027	-1.321	-2.052	1.543E-02	2.587E-01	x	
ENSSSCG00000023806	*LRRN1*	-0.776	-1.013	-1.983	1.580E-01	7.074E-01		x
ENSSSCG00000009157	*TET2*	-0.381	-1.123	-1.792	4.880E-01	9.582E-01		x
ENSSSCG00000011133	*PFKFB3*	-0.022	-2.256	-1.785	9.712E-01	9.987E-01	x	x
ENSSSCG00000002283	*FUT8*	-0.578	-1.286	-1.784	9.887E-02	6.059E-01	x	x
ENSSSCG00000023133	*OSBPL6*	-0.432	-1.088	-1.772	3.835E-01	9.108E-01	x	
ENSSSCG00000017986	*NDEL1*	-0.767	-1.644	-1.759	1.006E-02	2.081E-01	x	x
ENSSSCG00000031321	*NR4A1*	-0.630	-1.328	-1.720	6.298E-02	5.006E-01	x	
ENSSSCG00000035101	*KLF5*	-0.519	-1.487	-1.708	2.942E-01	8.488E-01	x	x
ENSSSCG00000004332	*BACH2*	-0.714	-2.105	-1.705	9.089E-02	5.861E-01	x	x
ENSSSCG00000017983	*PER1*	-0.773	-1.073	-1.627	3.000E-02	3.662E-01	x	

[a]$\log_2$FC: estimated $\log_2$ fold change for mean exonic fractions from *gluteus medius* expression profiles of fasted *AL-T0* and fed *AL-T2* Duroc gilts; [b]ΔEx: exonic fraction increment ($Ex_2 - Ex_1$) when comparing exon abundances in *AL-T0* (Ex_1) vs *AL-T2* (Ex_2) Duroc gilts; [c]PTc: post-transcriptional signal ($\Delta Ex - \Delta Int$) in z-score scale; [d]q-value: q-value calculated with the false discovery rate (FDR) approach [39]. The "x" symbols represent differentially expressed (DE) genes (FC < -2; *q*-value < 0.05) according to their exonic fractions, as well as those targeted by at least one of the upregulated miRNAs excluding redundant seeds (N = 6, **Table S5**). The "•" symbol represents suggestive differential expression (*P*-value < 0.01, **Table S4**).

Figure legends

Figure 1: (**A**) Scatterplot depicting mRNA genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction according to exonic (ΔEx) and PTc (ΔEx – ΔInt) values (in purple) and putatively targeted by DE upregulated miRNAs (FC > 1.5; *q*-value < 0.05) from *gluteus medius* skeletal muscle expression profiles of fasted (*AL-T0*, N = 11) and fed (*AL-T2*, N = 12) Duroc gilts. (**B**) Enrichment analyses of the number of mRNA genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction putatively targeted by DE upregulated miRNAs (FC > 1.5; *q*-value < 0.05), DE downregulated miRNAs (FC < -1.5; *q*-value < 0.05) and the top 5% most highly expressed miRNAs, excluding DE upregulated miRNAs. (**C**) Scatterplot depicting DE upregulated (in green) and downregulated (in red) mRNA genes (|FC| > 2; *q*-value < 0.05) according to exonic (ΔEx) and PTc (ΔEx – ΔInt) values. (**D**) Covariation enrichment scores (CES) for the exonic and intronic fractions of mRNA genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction, putatively targeted by upregulated miRNAs from *gluteus medius* skeletal muscle expression profiles of fasted (*AL-T0*, N = 11) and fed (*AL-T2*, N = 12) Duroc gilts. The control set of CES values were generated by permuted (N = 1,000) random sets of exonic and intronic profiles of genes with the same size as those used before (N = 21). Significant differences were assessed using a Mann-Whitney U non-parametric test [47].

Figure 2: (**A**) Scatterplot depicting mRNA genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction according to exonic (ΔEx) and PTc (ΔEx – ΔInt) values (in purple) and putatively targeted by upregulated miRNAs (FC > 1.5; *P*-value < 0.01) from adipocyte expression profiles of *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index (BMI). (**B**) Enrichment analyses

of the number of mRNA genes with the top 5% negative PTc scores and at least 2-folds

ΔEx reduction putatively targeted by DE upregulated miRNAs (FC > 1.5; *q*-value <

0.05), DE downregulated miRNAs (FC < -1.5; *q*-value < 0.05) and the top 5% most

highly expressed miRNAs, excluding DE upregulated miRNAs. (**C**) Scatterplot

depicting DE upregulated (in green) and downregulated (in red) mRNA genes (|FC| > 2;

q-value < 0.05) according to exonic (ΔEx) and PTc (ΔEx – ΔInt) values. (**D**)

Covariation enrichment scores (CES) for the exonic and intronic fractions of mRNA

genes with the top 5% negative PTc scores and at least 2-folds ΔEx reduction,

putatively targeted by upregulated miRNAs from adipocyte expression profiles of *lean*

(N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs. The control set of CES values

were generated by permuted (N = 1,000) random sets of exonic and intronic profiles of

genes with the same size as those used before (N = 25). Significant differences were

assessed using a Mann-Whitney U non-parametric test [47]. (**E**) Barplots depicting

qPCR $\log_2$ transformed relative quantities (Rq) for *LEP, OSBPL10, PRSS23, RNF157*

and *SERPINE2* mRNA transcripts measured in adipocytes from the retroperitoneal fat

of *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs. (**F**) Barplots depicting

qPCR $\log_2$ transformed relative quantities (Rq) for ssc-miR-148a-3p, ssc-miR-214-3p

and ssc-miR-92b-3p miRNA transcripts measured in the isolated adipocytes from the

retroperitoneal fat of *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs.

Supplementary Tables

Table S1: Phenotype values for selected Duroc-Göttingen minipigs from the F2-UNIK

resource population according to their body mass indexes.

1105

1106 **Table S2:** Primers for qPCR validation of selected mRNA and miRNA genes according

1107 to EISA results in the F2-UNIK Duroc-Göttingen minipig population comparing *lean*

1108 (N = 5) and *obese* (N = 5) individuals.

1109

1110 **Table S3:** Raw Cq values from qPCR analyses after efficiency correction measuring

1111 adipocyte expression profiles of selected mRNAs and miRNAs from *lean* (N = 5) and

1112 *obese* (N = 5) Duroc-Göttingen minipigs.

1113

1114 **Table S4:** Genes detected by *edgeR* tool as differentially expressed when comparing

1115 *gluteus medius* expression profiles of fasted *AL-T0* (N = 11) and fed *AL-T2* (N = 12)

1116 Duroc gilts.

1117

1118 **Table S5:** microRNAs detected by *edgeR* tool as differentially expressed when

1119 comparing *gluteus medius* expression profiles of fasted *AL-T0* (N = 11) and fed *AL-T2*

1120 (N = 12) Duroc gilts.

1121

1122 **Table S6:** EISA results for post-transcriptional signals (PTc) detected in *gluteus medius*

1123 skeletal muscle expression profiles of fasted (*AL-T0*, N = 11) and fed (*AL-T2*, N = 12)

1124 Duroc gilts.

1125

1126 **Table S7:** Binding sites for differentially upregulated miRNAs found in mRNA genes

1127 with the top 5% negative PT_c scores and at least 2-folds reduction in the exonic fraction

1128 (ΔEx) of *gluteus medius* skeletal muscle expression profiles from fasting (*AL-T0*, N =

1129 11) and fed (*AL-T2*, N = 12) Duroc gilts.

1130

Table S8: Covariation enrichment scores (CES) for the exonic and intronic fractions of mRNA genes with the top 5% negative post-transcriptional signals (PTc) and at least 2-folds reduction in their exonic (ΔEx) fraction, that were putatively targeted by DE upregulated miRNAs from *gluteus medius* skeletal muscle expression profiles *AL-T0 vs AL-T2* Duroc gilts.

Table S9: Genes detected by *edgeR* tool as differentially expressed when comparing adipocyte expression profiles from *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index.

Table S10: microRNA genes detected by *edgeR* tool as differentially expressed when comparing adipocyte expression profiles from *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index.

Table S11: mRNA genes with the top 5% post-transcriptional signals (PTc) and at least 2-fold exonic fraction (ΔEx) reduction (equivalent to -1 in the log2 scale) from adipocyte expression profiles of *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index.

Table S12: EISA results for post-transcriptional signals (PTc) detected in adipocyte expression profiles of *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index.

1153

Table S13: Binding sites for differentially upregulated miRNAs found in mRNA genes with the top 5% negative PTc scores and at least 2-folds reduction in the exonic fraction (ΔEx) of adipocyte expression profiles from *lean* (N = 5) and *obese* (N = 5) Duroc-Göttingen minipigs according to their body mass index.

Table S14: Covariation enrichment scores (CES) for the exonic and intronic fractions of mRNA genes with the top 5% negative post-transcriptional signals (PTc) and at least 2-folds reduction in their exonic (ΔEx) fraction, that were putatively targeted by DE upregulated miRNAs from adipocyte expression profiles of *lean* vs *obese* Duroc-Göttingen minipigs.

Supplementary Figures

Figure S1: Diagram depicting the consecutive steps implemented for studying miRNA-driven post-transcriptional regulatory signals applying the EISA approach and additional enrichment and covariation analyses.

Figure S2: Scatterplots depicting the exonic (ΔEx) and intronic (ΔInt) fractions from *gluteus medius* skeletal muscle expression profiles of fasting (*AL-T0*, N = 11) and fed (*AL-T2*, N = 12) Duroc gilts. (**A**) mRNA genes with the top 5% post-transcriptional (PTc) negative scores and at least 2-folds reduced exonic (ΔEx) fractions (equivalent to -1 in the $\log_2$ scale), suggestive of miRNA-driven post-transcriptional regulation. (**B**) mRNA genes differentially expressed showing upregulation (FC > 2; *q*-value < 0.05, in green) and downregulation (FC < -2, *q*-value < 0.05, in red) in fed (*AL-T2*, N = 12) Duroc gilts with respect to their fasted (*AL-T0*, N = 11) counterparts.

1179

Figure S3: Enrichment analyses of the number of genes with the (**A**) top 1% and (**B**)

top 5% negative PTc scores and at least 2-fold reduced exonic fractions (ΔEx)

putatively targeted by upregulated miRNAs (FC > 1.5; *q*-value < 0.05) from *gluteus*

medius skeletal muscle expression profiles of fasting (*AL-T0*, N = 11) and fed (*AL-T2*,

N = 12) Duroc gilts. Results show the change in enrichment significance when

incorporating context-based pruning of miRNA binding sites of type 8mer, 7mer-m8

and 7mer-A1. R: Raw enrichment analyses without any additional context-based

pruning. AU: Enrichment analyses removing miRNA binding sites without AU-rich

flanking sequences (30 nts upstream and downstream). M: Enrichment analyses

removing miRNA binding sites located in the middle of the 3'-UTR sequence (45-

55%). E: Enrichment analyses removing miRNA binding sites located too close (< 15

nts) to the beginning or the end of the 3'-UTR sequences. The black dashed line

represents a *P*-value = 0.05.

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204