Gut microbiome dysbiosis during COVID-19 is associated with increased risk for bacteremia and microbial translocation.

Mericien Venzon[*1,2], Lucie Bernard-Raichon[*2], Jon Klein[*3], Jordan E. Axelrad[*4], Chenzhen Zhang[*5], Grant A. Hussey[5], Alexis P. Sullivan[5], Arnau Casanovas-Massana[6], Maria G. Noval[7], Ana M. Valero-Jimenez[7], Juan Gago[1,8], Gregory Putzel[7,12], Alejandro Pironti[7,12], Evan Wilder[4], Yale IMPACT Research Team[11], Lorna E. Thorpe[8,12], Dan R. Littman[2,9], Meike Dittmann[7], Kenneth A. Stapleford[7], Bo Shopsin[7,10,12], Victor J. Torres[7,12], Albert I. Ko[6], Akiko Iwasaki[3,9], Ken Cadwell[2,4,7,12]†, Jonas Schluter[5,7,12]†

1. Vilcek Institute of Graduate Biomedical Sciences, New York University Grossman School of Medicine, New York, NY, USA
2. Kimmel Center for Biology and Medicine at the Skirball Institute, New York University Grossman School of Medicine, New York, NY, USA
3. Department of Immunobiology, Yale University School of Medicine, New Haven, CT, USA
4. Division of Gastroenterology, Department of Medicine, New York University Grossman School of Medicine, New York, NY, USA
5. Institute for Systems Genetics, New York University Grossman School of Medicine, New York, NY, USA
6. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT, USA.
7. Department of Microbiology, New York University Grossman School of Medicine, New York, NY, USA
8. Department of Population Health, New York University Grossman School of Medicine, New York, NY, USA.
9. Howard Hughes Medical Institute, Chevy Chase, MD.
10. Department of Medicine, Division of Infectious Diseases, New York University Grossman School of Medicine, New York, NY, USA
11. IMPACT team members listed in the appendix.
12. Antimicrobial-Resistant Pathogens Program, New York University School of Medicine, New York, NY, USA.

*) These authors contributed equally

Correspondence:

† ken.cadwell@nyulangone.org

† jonas.schluter@nyulangone.org

Abstract

The microbial populations in the gut microbiome have recently been associated with COVID-19 disease severity. However, a causal impact of the gut microbiome on COVID-19 patient health has not been established. Here we provide evidence that gut microbiome dysbiosis is associated with translocation of bacteria into the blood during COVID-19, causing life-threatening secondary infections. Antibiotics and other treatments during COVID-19 can potentially confound microbiome associations. We therefore first demonstrate in a mouse model that SARS-CoV-2 infection can induce gut microbiome dysbiosis, which correlated with alterations to Paneth cells and goblet cells, and markers of barrier permeability. Comparison with stool samples collected from 96 COVID-19 patients at two different clinical sites also revealed substantial gut microbiome dysbiosis, paralleling our observations in the animal model. Specifically, we observed blooms of opportunistic pathogenic bacterial genera known to include antimicrobial-resistant species in hospitalized COVID-19 patients. Analysis of blood culture results testing for secondary microbial bloodstream infections with paired microbiome data obtained from these patients indicates that bacteria may translocate from the gut into the systemic circulation of COVID-19 patients. These results are consistent with a direct role for gut microbiome dysbiosis in enabling dangerous secondary infections during COVID-19.

Main text

A better understanding of factors contributing to the pathology of coronavirus disease 2019 (COVID-19) is an urgent global priority. Previous reports have demonstrated that severe COVID-19 is frequently associated with specific inflammatory immune phenotypes, lymphopenia, and a generally disproportionate immune response leading to systemic organ failure[1,2]. Even in mild cases, gastrointestinal symptoms are reported frequently, and recent studies reported that COVID-19 patients lose commensal taxa of the gut microbiome during hospitalization[3–5]. Differences in gut bacterial populations relative to healthy controls were observed in all COVID-19 patients, but most strongly in patients who were treated with antibiotics during their hospitalization[4]. Most recently, COVID-19 patients treated with broad spectrum antibiotics at admission were shown to have increased susceptibility to multi-drug resistant infections and nearly double the mortality rate from septic shock[6,7]. Furthermore, although initially estimated to be low (6.5%)[8], more recent studies have detected bacterial secondary infections in as much as 12-14% of COVID-19 patients[9,10]. However, the causal direction of the relationship between disease symptoms and gut bacterial populations is not yet clear.

Complex gut microbiota ecosystems can prevent the invasion of potentially pathogenic bacteria[11,12]. Conversely, when the gut microbiota incurs damage, such as through antibiotics treatment, competitive exclusion of pathogens is weakened[13–15] and potentially dangerous blooms of antibiotic resistant bacterial strains can occur[16,17]. In immunocompromised cancer patients, blooms of Enterococcaceae and Gram-negative proteobacteria can lead to gut dominations by few or single species[18–21]. Such gut domination events are dangerous to these patients because they are associated with increased risk of translocation of antibiotic resistant bacteria from the gut into the blood stream[18]. Bacterial co-infection can also cause life-threatening complications in patients with severe viral infections[7,8,22]; therefore, antibacterial agents were administered empirically to nearly all critically ill suspected COVID-19 patients since the incidence of bacterial superinfection was unknown early during the pandemic[4,23]. However, it is now known that nosocomial infection during prolonged hospitalization is the primary threat to patients with COVID-19[24], rather than bacterial co-infection upon hospital admission[9,25–27]. Evidence from immunocompromised cancer patients suggests that indiscriminate administration of broad-spectrum antibiotics may, counter-intuitively, increase

83   nosocomial bloodstream infection (nBSI) rates by causing gut dominations of resistant microbes

84   that can translocate into the blood[18,28]. Thus, empiric antimicrobial use, i.e. without direct

85   evidence for a bacterial infection, in patients with severe COVID-19 may be especially

86   pernicious because it may select for antimicrobial resistance and could promote gut

87   translocation-associated nBSI.

88   The role of the gut microbiome in respiratory viral infections in general[29–31], and in

COVID-19 patients in particular, is only beginning to be understood. Animal models of

influenza virus infection have uncovered mechanisms by which the microbiome influences

antiviral immunity[32–34], and in turn, the viral infection was shown to disrupt the intestinal barrier

of mice by damaging the gut microbiota[35,36]. Hence, we hypothesized that gut dysbiosis during

COVID-19 may be associated with BSIs. To test this, we first determined whether SARS-CoV-2

infection could directly cause gut dysbiosis independently of hospitalization and treatment. K18-

hACE2 mice (*K18-ACE2tg* mice), express human *ACE2* driven by the *cytokeratin-18* promoter

(*K18-ACE2tg* mice). Although the overexpression of ACE2 prevents investigation of long term

consequences of infection due to potential non-specific disease, which is a major limitation of

the model, an advantage of these mice is that they develop severe respiratory disease in a virus

dose-dependent manner, partially mirroring what is observed in COVID-19 patients[37–40]. Daily

changes in fecal bacterial populations were monitored following intranasal inoculation of mice

with a range of doses (10, 100, 1000, and 10000 PFU) of SARS-CoV-2 or mock-treatment (**Fig.**

1a, Extended Data Fig. S1). Although we detected viral RNA in the lungs of mice infected

with doses as low as 100 PFU (**Extended Data Fig. S1c**), mice inoculated with doses lower than

10000 PFU displayed minimal or no signs of disease (**Extended Data Fig. S1a,b**), and as

expected based on this outcome, shifts in their microbiome were inconsistent (**Extended Data**

Fig. S2). Thus, we focused on findings from the 10000 PFU inoculum.

Mice infected with 10000 PFU displayed weight loss and other signs of disease around

day 4 (**Extended Data Fig. S1a,b, S2e,f**), alongside microbiome changes characterized by a

significant loss of alpha diversity (inverse Simpson index, **Fig. 1b**) corresponding to shifts in the

bacterial community composition (**Fig. 1c,d**). We performed time series analyses on bacterial

family abundances, contrasting their trajectories in infected (10000 PFU) and uninfected mice.

This revealed that the strongest shift over time in infected mice was characterized by significant

increases of Akkermansiaceae ($p<0.0002$, **Fig. 1d**). Ranking all bacterial family trajectories by

their estimated changes over time in infected mice showed that this increase in Akkermansiaceae

was accompanied by significant losses of Clostridiaceae 1, a family of obligate anaerobe

bacteria, and of Erysipelotrichiaceae (**Fig. 1e**). These results demonstrated that SARS-CoV-2

infection leads to gut microbiome dysbiosis over time in a mouse model.

We then determined if this dysbiosis was also associated with intestinal defects that could

enable translocation of bacteria into the blood. While several of the infected mice displayed signs

of barrier dysfunction the observed differences in plasma concentrations of Fluorescein

isothiocyanate (FITC)-dextran following its administration by gavage, or other markers of

intestinal barrier permeability, fatty acid-binding protein (iFABP), Lipopolysaccharide-binding

protein (LBP), and citrulline did not reach significance (**Extended Data Fig. S3a,b**). The

reduced colon lengths as well as reductions in the villus lengths in the duodenum or ileum, i.e.

markers of overt inflammation, that we observed were also non-significant compared with

control mice (**Extended Data Fig. S3c,d**). However, infected mice that had incurred the most

severe microbiome injury in the form of diversity loss showed the most evidence of gut

permeability–the highest FITC-dextran concentrations in the blood of mice detected across all

samples came from the two out of the four mice with the most extreme dysbiosis and highest

levels of Akkermansiaceae, a family of mucin-degrading bacterial species (**Extended Data Fig.**

S4).

Interestingly, we also detected a significant increase in the number of mucus-producing

goblet cells and a decrease in the number of Paneth cells in the ileum (but not in the duodenum)

of infected mice (**Fig. 2a,c and Extended Data Fig. S3e**). The decrease in Paneth cells was

accompanied by structural abnormalities, most notably deformed or misplaced granules (**Fig.**

2b). These morphological abnormalities in Paneth cells were reminiscent of observations in the

ileum of patients with inflammatory bowel disease (IBD) as well as in a virally-triggered animal

model of IBD, where such structures were indicative of defects in packaging and secretion of the

granule protein lysozyme[41–43]. Thus, to quantify the Paneth cell granule defect, we performed

lysozyme immunofluorescence and found a significant increase in the proportion of Paneth cells

displaying abnormal staining patterns compared with the controls (**Fig. 2b,c**). We then

investigated if these physiological defects were associated with dysbiosis in the microbiome. The

most severely sick mice also had the most striking shifts in their microbiome composition and

the lowest microbiota diversity at the end of the experiment (**Extended Data Fig. S4a,b**). To

associate the observed physiological defects with microbiome dysbiosis, we plotted the numbers

of goblet cells per crypt-villus unit and Paneth cells per crypt, as well as the percentage of

abnormal Paneth cells against bacterial alpha diversity and the $\log_{10}$-relative abundance of

Akkermansiaceae (**Fig. 2d,e**). Goblet cell counts per crypt-villus unit were negatively correlated

with alpha diversity, and, conversely, positively correlated with Akkermansiaceae. While no

statistically significant association was found between diversity, Akkermansiaceae abundance

and Paneth cell counts per crypt, we observed a striking positive correlation between the

percentage of abnormal Paneth cells and Akkermansiaceae, and a corresponding negative

correlation with diversity. We were unable to reliably detect viral RNA in intestinal samples

(**Extended Data Fig. S1c**), raising the possibility that systemic immune responses rather than

direct cytotoxicity from local viral infection mediate these changes. Altogether, these results

show that the gut microbiome dysbiosis observed in K18-hACE2 mice infected with a high dose

of SARS-CoV-2 are associated with alterations in key epithelial cells, and signs of barrier

permeability in the mice displaying the greatest disruption in microbiome diversity.

159   To investigate the microbiome in COVID-19 patients, we profiled the bacterial

composition of the fecal microbiome in 130 samples (**Fig. 3a**) obtained from SARS-CoV-2

infected patients treated at NYU Langone Health (NYU, 67 samples from 60 patients) and Yale

New Haven Hospital (YALE, 63 samples from 36 patients, **Supplementary Table 1**). Analysis

of metagenomic data obtained from sequencing of the 16S rRNA genes revealed a wide range of

bacterial community diversities, as measured by the inverse Simpson index, in samples from

both centers (NYU: [1.0, 32.3], YALE: [1.5, 29.3], **Fig. 3b**); on average, samples from NYU

were less diverse (-4, $p<0.01$, two-tailed T-test, **Fig. 3c**), and as reported previously, samples

from patients admitted to the ICU had reduced diversity (-3.9, $p<0.05$, two-tailed T-test,

Extended Data Fig. S5a). However, the composition in samples between the two centers did not

show systematic compositional differences (**Fig. 3d,e,f**). On average, in both centers, members

of the phyla Firmicutes and Bacteroidetes represented the most abundant bacteria, followed by

Proteobacteria (**Fig. 3d**). The wide range of bacterial diversities was reflected in the high

variability of bacterial compositions across samples (**Fig. 3e,f**). In samples from both centers,

microbiome dominations, defined as a community where a single genus reached more than 50%

of the population, were observed frequently (NYU: 21 samples, YALE: 12 samples),

representing states of severe microbiome injury in COVID-19 patients (**Fig. 3g, Extended Data**

Fig. S5a,b). Strikingly, samples associated with a BSI, defined here as a positive clinical blood

culture test result, had strongly reduced bacterial α-diversities (mean difference: -5.2, CI$_{BEST}$[-

8.2, -2.2], **Fig. 3h**).

The lower diversity associated with samples from 25 patients (15 NYU, 10 Yale) with

BSIs led us to investigate their bacterial taxon compositions and the potential that gut dysbiosis

was associated with BSI events. Importantly, BSI patients had received antibiotic treatments

during hospitalization (**Extended Data Fig. S6, Supplementary Table 2**), which could

exacerbate COVID-19 induced shifts in microbiota populations[16,17,20], and may indeed be

administered in response to a suspected or confirmed BSI. We noted that most BSI patients

received antibiotics prior to their BSI, with 6 out of 25 patients receiving antibiotics only after

detection of BSI. Principal coordinate analysis of all stool samples indicated that the BSI-

associated samples spanned a broad range of compositions (**Fig. 3h**). To identify bacterial

abundance patterns that consistently distinguished BSI from non-BSI-associated samples, we

performed a Bayesian logistic regression. The model estimated the association of the 10 most

abundant bacterial genera with BSI cases, i.e. it identified enrichment or depletion of bacterial

genera in BSI associated samples (**Fig. 3i**). This analysis revealed that the genus

Faecalibacterium was negatively associated with BSI (OR: -0.5, CI:[-0.86, -0.15]), which was

also observed when we included microbiome domination as an additional factor in the model

(**Extended Data Fig. S7a**). However, our analysis also included stool samples that were taken

only after a positive blood culture was obtained, calling into question the plausibility of gut

translocation; a complementary analysis only using stool samples obtained prior or on the same

day of a positive blood culture also identified *Faecalibacterium* as most negatively associated

with BSI (**Extended Data Fig. S7b**). Furthermore, a higher-resolution analysis using amplicon

sequencing variant (ASV) relative abundances as predictors of BSI (**Extended Data Fig. S7c,d**),

identified an ASV of the *Faecalibacterium* genus as most negatively associated with BSI, in

agreement with our main analysis. *Faecalibacterium* is an immunosupportive, short-chain fatty

acid producing genus that is a prominent member of the human gut microbiome[44–46], and its

reduction is associated with disruption to intestinal barrier function[47,48], perhaps via ecological

network effects[48].

To evaluate the effect size of the association between *Faecalibacterium* and BSIs, we

performed a counterfactual posterior predictive check. Using the average genus composition

found across all samples, we first computed the distribution of predicted BSI risks (**Fig. 3j**), and

compared this risk distribution with a hypothetical bacterial composition which increased

Faecalibacterium by 10% points. The predicted risk distributions associated with these two

compositions differed strongly (mean difference 15%, CI: [1%, 32%], **Fig. 3j**). Domination

states of the microbiome increase the risk for BSIs in immunocompromised cancer patients[18];

such dominations imply high relative abundances of single taxa, and therefore a low diversity.

Consistent with this, *Faecalibacterium* abundance was positively correlated with diversity (R:

0.55, p<10^{-10}, **Extended Data Fig. S8**) in our data set and as reported previously[44].

We therefore next investigated a direct association between the bacteria populating the

gut microbiome and the organisms identified in the blood of patients. Visualizing the bacterial

composition in stool samples from patients alongside the BSI microorganisms (**Extended Data**

Fig. S9a) suggested a correspondence with the respective taxa identified in the blood: high

abundances of the BSI-causing microbes were found in corresponding stool samples. A rank

abundance analysis matching the organisms identified in clinical blood cultures to the

composition of bacteria in corresponding stool samples indicated enrichment of taxa belonging

to the same bacterial orders as BSI causing organisms (**Extended Data Fig. S9b**), suggesting

translocation of bacteria from the gut into the blood stream.

To further investigate evidence for translocation of gut bacteria into the blood, we next

performed shotgun metagenomic sequencing on a subset of BSI-associated samples with

sufficient remaining material in order to match the organism identified in clinical blood cultures

at the species level with reads obtained from stool samples (**Fig. 3k, Supplementary Table 3**).

In four cases of positive blood cultures of *Staphylococcus* species, no reads matching the

clinically identified species were detected (**Supplementary Table 3**). This may explain why the

rank analysis suggested that Staphylococcales were not generally enriched in BSIs by

Staphylococcus (**Extended Data Fig. S9a,b**). In all investigated cases of positive blood cultures

by other organisms, the species identified in clinical blood cultures had corresponding reads in

the stool samples. Furthermore, the relative abundances of matched species tended to be larger

than the average abundances of matched species across all samples (**Supplementary Table 3**).

Consistent with this, in one case of a *S. aureus* BSI where corresponding stool relative

abundances of *Staphylococcus* were low, reads from shotgun sequencing did not match the

genomes of isolates obtained from the same patient better than *S. aureus* genomes from other

isolates (**Extended Data Fig. S9d**). Strikingly, shotgun metagenomic reads matched the genome of isolates well in another case where relative abundances of *Staphylococcus* were enriched in the stool (**Extended Data Fig. S9c**), providing evidence that here, the same strains were found in stool and blood of the same patient.

Collectively, these results reveal an unappreciated link between SARS-CoV-2 infection, gut microbiome dysbiosis, and a severe complication of COVID-19, BSIs. The loss of diversity and immunosupportive *Faecalibacterium* in patients with BSIs mirrored a similar loss of diversity in the most severely sick mice deliberately infected with SARS-CoV-2, and as observed by other labs and other model systems[49–51]. Notably, a recent study reproduced these changes in the microbiome in an antibiotics-naïve cohort[52], suggesting that the viral infection causes gut dysbiosis, either through gastrointestinal infection[53–57] or through a systemic inflammatory response[2,4]. Furthermore, the pronounced increase in Akkermansiaceae in mice was also observed in our patient samples, and has been reported previously in patients and in K18-hACE2 mice[49,58]. However, the dysbiosis in patients with COVID-19 exceeded the microbiota shifts observed in the mouse experiments, including microbiome dominations by single taxa, which was not seen in the mouse experiments. It is possible that in our experiment, mice were sacrificed before perturbations to the gut microbial populations reached a maximum. hACE2 knock-in mice, which display reduced disease[37], were not tested in the scope of this study but could provide additional insights in the future. However, it is also plausible that the frequently administered antibiotic treatments that hospitalized COVID-19 patients receive exacerbated SARS-CoV-2 induced microbiome perturbations. Additionally, unlike the controlled environment experienced by laboratory mice, hospitalized patients are uniquely exposed to antimicrobial-resistant infectious agents present on surfaces and shed by other patients.

Despite these limitations of the mouse model, we observed that SARS-CoV-2 infection led to alteration of intestinal epithelial cells with established roles in intestinal homeostasis and gastrointestinal disease[59,60]. Microbiome ecosystem shifts are likely both cause and consequence of these epithelial cell alterations, since epithelial secretions are predicted to affect overall community structure disproportionately strongly[61,62]. For example, disruption of Paneth cell-derived antimicrobials including lysozyme are sufficient to impact microbiome composition[63–65], and, conversely, *Akkermansia*, which was increased in infected mice, can have epithelium remodeling properties[66].

Our observation that the type of bacteria that entered the bloodstream was enriched in the

associated stool samples is a well characterized phenomenon in cancer patients[18], especially

during chemotherapy induced leukocytopenia when patients are severely

immunocompromised[16,44]. COVID-19 patients are also immunocompromised and frequently

incur lymphopenia, rendering them susceptible to secondary infections[67]. Our data suggests

dynamics in COVID-19 patients may be similar to those observed in cancer patients: BSI-

causing organisms may translocate from the gut into the blood, potentially due to loss of gut

barrier integrity, through tissue damage downstream of antiviral immunity rather than

chemotherapy. Consistent with this possibility, soluble immune mediators such as TNFα and

interferons produced during viral infections, including SARS-CoV-2, damage the intestinal

epithelium to disrupt the gut barrier, especially when the inflammatory response is sustained as

observed in patient with severe COVID-19[43,68,69]. Indeed, blood plasma in severely sick COVID-

19 patients are enriched for markers of disrupted barrier integrity and higher levels of

inflammation markers[70], suggesting microbial translocation. Our data supports this model with

direct evidence because we were able to match sequencing reads from stool samples to genomes

of species detected in the blood of patients.

One limitation of our data is temporal ordering of samples. Occasionally stool samples

were collected after observation of BSI, and this mismatch in temporal ordering is counter

intuitive for gut-to-blood translocation and a causal interpretation of our associations. However,

the reverse direction, that blood infection populates and changes the gut community, is unlikely

for the organisms identified in the blood, and if our associations were not causal, we would

expect no match between BSI organisms and stool compositions.

Taken together, our findings support a scenario in which gut-to-blood translocation of

microorganisms following microbiome dysbiosis, a known issue for chronic conditions such as

cancer, leads to dangerous BSIs during COVID-19. We suggest that investigating the underlying

mechanism behind our observations will inform the judicious application of antibiotics and

immunosuppressives in patients with respiratory viral infections and increase our resilience to

pandemics.

Materials and Methods

Bioethics statement

The collection of COVID-19 human biospecimens for research has been approved by the

NYUSOM Institutional Review Board under il8-01121 Inflammatory Bowel Disease and Enteric

Infection at NYU Langone Health. The data presented in this study were also approved by Yale

Human Research Protection Program Institutional Review Boards (FWA00002571, protocol ID

2000027690). Informed consent was obtained from all enrolled patients.

Mouse experiments

<u>Cells & virus</u>

Vero E6 (CRL-1586; American Type Culture Collection) were cultured Dulbecco's Modified

Eagle's Medium (DMEM, Corning) supplemented with 10% fetal bovine serum (FBS, Atlanta

Biologics) and 1% nonessential amino acids (NEAA, Corning). SARS-CoV-2, isolate USA-

WA1/2020 19 (BEI resources #NR52281), a gift from Dr. Mark Mulligan at the NYU Langone

Vaccine Center was amplified once in Vero E6cells. All experiments with SARS-CoV-2 were

conducted in the NYU Grossman School of Medicine ABSL3 facility in accordance with its

Biosafety Manual and Standard Operating Procedures, by personnel equipped with powered air-

purifying respirators.

<u>Mice</u>

Heterozygous K18-hACE2 C57BL/6J mice (strain: 2B6.Cg-Tg(K18-ACE2)2Prlmn/J) were

obtained from The Jackson Laboratory. Several were paired with C57BL/6J mice to generate

additional heterozygous mice for subsequent experiments and the remaining were used to

perform initial experiments. Animals from the same breeder pool (i.e., littermates) were

randomly assigned and housed in cages according to the experimental groups and fed standard

chow diets. Cage bedding was mixed prior to infection in a subset of experiments to further

reduce possible cage effect. All animal studies were performed according to protocols approved

by the NYU School of Medicine Institutional Animal Care and Use Committee (IACUC

n°170209 and 180802) and in the ABSL3 facility of NYU Grossman School of Medicine (New

York, NY), in accordance with its Biosafety Manual and Standard Operating Procedures. 12-

week-old or 24-week-old K18-hACE2 males were administered either 10-10000 PFU SARS-

CoV-2 diluted in 50μL PBS (Corning) or 50μL PBS (non-infected, 0) via intranasal

administration under xylazine-ketamine anesthesia (AnaSedR AKORN Animal Health,

KetathesiaTM Henry Schein Inc). Viral titer in the inoculum was verified by plaque assay in

Vero E6 cells. Following infection, mice were monitored daily for weight loss, temperature loss

and signs of disease. A disease score was calculated as the sum of scores obtained for each of the

following criteria: ruffled fur (no= 0, yes=1), hunched back (no=0, slightly=1, exacerbated=2),

heavy breathing (no=0, yes=1), altered mobility (no=1, reduced activity=1, no mobility=2). Stool

samples were collected and stored at -80°C.

<u>Measurement of viral load</u>

Whole lungs and 1cm of proximal duodenum, terminal ileum and proximal colon were collected

five-to seven days after infection. Intestinal pieces were wash with PBS and all organs were

transferred in Eppendorf tubes containing 500µl of PBS and a 5mm stainless steel bead (Qiagen)

and h C57BL/6J mice omogenized using the Qiagen TissueLyser II. Homogenates were cleared

for 5 min at $5,000 \times g$, and the viral supernatant or nasal wash was diluted 4X in TRIzol reagent

(Invitrogen) and frozen at -80°C for titration by qRT-PCR. RNA was extracted from the TRIzol

homogenates using chloroform separation and isopropanol precipitation, followed by additional

purification using RNeasy spin columns with DNase treatment according to the manufacturer's

instructions (Rneasy Mini Kit; RNAse-Free DNase Set; QIAGEN). RNA was reverse-

transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems).

qPCR was performed using Applied Biosystems TaqMan RNA-to-CT One-Step Kit (Fisher-

Scientific), 500nM of the primers (Fwd 5'-ATGCTGCAATCGTGCTACAA-3', Rev 5'-

GACTGCCGCCTCTGCTC-3') and 100nM of the N probe (5'-/56-

FAM/TCAAGGAAC/ZEN/AACATTGCCAA/3IABkFQ/-3'). Serial dilutions of *in-vitro* transcribed

RNA of the SARS-CoV-2 Nucleoprotein (generated as previously described[71]) were used to

generate a standard curve and calculate copy numbers per ug of RNA in the samples.

<u>Microscopy</u>

5cm of proximal duodenum, distal ileum and entire colon were flushed with 10% acetate

buffered formalin (Fisher scientific), cut open along the length, pinned on black wax and fixed

with formalin for 72hrs at RT. 2 cm strips of intestinal tissues were embedded in low melting

point agarose (Promega) to enrich for well-oriented crypt-villus units. Paraffin embedding,

sectioning, and staining were performed by the NYU Experimental Pathology Research

Laboratory. 5um sections were stained with hematoxylin and eosin (H&E) and imaged using

brightfield wholeslide scanning. Lysozyme staining was performed using anti-lysozyme

(ab108508, Abcam) and DAPI immunostaining and analyzed using a Zeiss AxioObserver.Z1

with Axiocam 503 Mono operated with Zen Blue software. 50 small intestinal villi per mouse

were measured for villi length. Goblets cell were quantified from 50 villus-crypt units (one villus

+ half of the 2 surrounding crypts) per mouse. Paneth cells numbers and lysozyme staining

patterns were quantified from 50 crypts per mouse. Previously defined criteria were used to

quantify the proportion of Paneth cells displaying abnormal lysozyme staining[43]. Mean values

were calculated for each mouse and used as individual data points.

<u>Measurement of intestinal permeability</u>

Mice were fasted for 4hrs before oral gavage with 200uL of fluorescein isothiocyanate (FITC)-

dextran (3-5 kDa, Sigma-Aldrich) dissolved in sterile PBS (60mg/ml). After 4 hrs, mice were

euthanized and blood was collected by cardiac puncture. FITC-dextran in plasma was quantified

using a plate reader (excitation, 485 nm; emission, 530 nm). Citrulline, intestinal fatty acid-

binding protein, and lipopolysaccharide (LPS)-binding protein were quantified in the plasma by

enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions

(MyBioSource, CA).

<u>Time series analyses of bacterial family abundances</u>

We $\log_{10}$-transformed bacterial relative abundances adding a pseudo count to fill zeros ($2*10^{-6}$,

as done before). We then analyzed the time series with the following model that included fixed

effects for the intercepts and slopes of the treatment (i.e. indicator variables for uninfected (0

PFU), and infected (10000PFU), and random effects per cage and per mice to account for cage

effects and repeated measurements from the same individual mouse, respectively. The model

was implemented in the R programming language using the lmer function of the lme4 library

with the following model formula:

389            $\log X_i \sim 1 + \text{PFU} + \text{time:PFU} + (1|\text{cage}) + (\text{time}|\text{mouseid})$

Human study

<u>Study population and data collection</u>

This study involved 96 patients with laboratory-confirmed SARS-CoV-2 infection. SARS-CoV-

2 infection was confirmed by a positive result of real-time reverse transcriptase-polymerase

chain reaction assay on a nasopharyngeal swab. 60 patients were seen at NYU Langone Health,

New York, between January 29, 2020 – July 2, 2020. In order to be eligible for inclusion in the

study, stool specimens needed to be from individuals >18 years of age. Data including

demographic information, clinical outcomes, and laboratory results were extracted from the

electronic medical records in the NYU Langone Health clinical management system. Blood and

stool samples were collected by hospital staff. OmnigeneGut kits were used on collected stool. In

parallel, 36 patients were admitted to YNHH with COVID-19 between 18 March 2020 and 27

May 2020 as part of the YALE IMPACT cohort described at length elsewhere[2]. Briefly,

participants were enrolled after providing informed consent and paired blood and stool samples

were collected longitudinally where feasible for duration of hospital admission. No statistical

methods were used to predetermine sample size for this cohort. Demographic information of

patients was aggregated through a systematic and retrospective review of the EHR and was used

to construct **Supplementary Table 1**. Symptom onset and etiology were recorded through

standardized interviews with patients or patient surrogates upon enrolment in our study, or

alternatively through manual EHR review if no interview was possible owing to clinical status at

enrolment. The clinical data were collected using EPIC EHR and REDCap 9.3.6 software. At the

time of sample acquisition and processing, investigators were blinded to patient clinical status.

<u>DNA extraction and bacterial 16S rRNA sequencing</u>

For bacterial DNA extraction 700µL of SL1 lysis buffer (NucleoSpin Soil kit, Macherey-Nagel)

was added to the stool samples and tubes were heated at 95°C for 2h to inactivate SARS-CoV-2.

Samples were then homogenized using the FastPrep-24TM instrument (MP Biomedicals) and

extraction was pursued using the NucleoSpin Soil kit according to the manufacturer's

instructions. DNA concentration was assessed using a NanoDrop spectrophotometer. Samples

with too low DNA concentration were excluded. DNA from human samples was extracted with

PowerSoil Pro (Qiagen) on the QiaCube HT (Qiagen), using Powerbead Pro (Qiagen) plates with

0.5mm and 0.1mm ceramic beads. For mouse samples, the variable region 4 (V4) of the 16S

rRNA gene was amplified by PCR using primers containing adapters for MiSeq sequencing and

single-index barcodes. All PCR products were analyzed with the Agilent TapeStation for quality

control and then pooled equimolar and sequenced directly in the Illumina MiSeq platform using

the 2x250 bp protocol. Human samples were prepared with a protocol derived from [72], using

KAPA HiFi Polymerase to amplify the V4 region of the 16S rRNA gene. Libraries were

sequenced on an Illumina MiSeq using paired-end 2x250 reads and the MiSeq Reagent Kitv2.

Bioinformatic processing and taxonomic assignment

Amplicon sequence variants (ASVs) were generated via dada2 v1.16.0 using post-QC FASTQ

files. Within the workflow, the paired FASTQ reads were trimmed, and then filtered to remove

reads containing Ns, or with maximum expected errors >= 2. The dada2 learn error rate model

was used to estimate the error profile prior to using the core dada2 algorithm for inferring the

sample composition. Forward and reverse reads were merged by overlapping sequence, and

chimeras were removed before taxonomic assignment. ASV taxonomy was assigned up to genus

level using the SILVAv.138 database with the method described in [73] and a minimum

boostrapping support of 50%. Species-level taxonomy was assigned to ASVs only with 100%

identity and unambiguous matching to the reference.

Shotgun metagenomic sequencing

DNA was quantified with Qiant-iT Picogreen dsDNA Assay (Invitrogen). Libraries were

prepared with a procedure adapted from the Nextera Library Prep kit (Illumina), and sequenced

on an Illumina NovaSeq using paired-end 2x150 reads (Illumina) aiming for 100M read depth.

DNA sequences were filtered for low quality (Q-Score < 30) and length (< 50), and adapter

sequences were trimmed using cutadapt. Fastq files were converted a single fasta using shi7.

Sequences were trimmed to a maximum length of 100 bp prior to alignment. DNA sequences

were taxonomically classified using the MetaPhlAn2 analysis tool

(http://huttenhower.sph.harvard.edu/metaphlan2). MetaPhlAn2 maps reads to clade-specific

marker genes identified from ~17,000 reference genomes and estimates clade abundance within a

sample from these mappings.

Mapping shotgun reads to whole genome sequences of clinical isolates

Quality-controlled reads were re-classified using Kraken2 (Minikraken2 v2 database, available

on https://ccb.jhu.edu/software/kraken2/index.shtml). Reads that were classified by Kraken2 as

Staphylococcus aureus (or a strain thereof) were further mapped using Bowtie2 separately to

each of a collection of *Staphylococcus aureus* isolates. The collection was composed of all NCBI

RefSeq assemblies as of 11/17/2021, in addition to *Staphylococcus aureus* isolates that were

isolated from our subjects. Bowtie2 mapped reads were then further filtered, keeping only reads

that mapped without mismatches. A neighbor-joining (NJ) tree was produced from this

collection of genomes using Snippy (https://github.com/tseemann/snippy).

Compositional analyses

<u>α-Diversity</u>

We calculated the inverse Simpson (*IVS*) index from relative ASV abundances (*p*) with *N* ASVs

in a given sample, $IVS = \frac{1}{\Sigma_i^N p_i^2}$.

<u>Principal Coordinate Analyses</u>

Bray-Curtis distances were calculated from the filtered ASV table using QIIME 1.9.1 and

principal components of the resulting distance matrix were calculated using the scikit-learn

package for the Python programming language, used to embed sample compositions in the first

two principal coordinates.

<u>Average compositions and manipulation of compositions</u>

To describe the average composition of a set of samples we calculated the central tendency of a

compositional sample [74]. For counter factual statistical analyses that require changes to a

composition, e.g. an increase in a specific taxon, we deployed the perturbation operation ($\oplus$),

which is the compositional analogue to addition in Euclidean space[74]. A sample *x* containing the

original relative taxon abundances is perturbed by a vector *y*,

$y: \ x \oplus y = \left[\frac{x_1 y_1}{\sum_{i=1}^{D} x_i y_i}, \frac{x_2 y_2}{\sum_{i=1}^{D} x_i y_i}, \ldots, \frac{x_D y_D}{\sum_{i=1}^{D} x_i y_i} \right] \forall x, y \in S^D$

where S^D represents the D-part simplex.

<u>Bayesian t-test</u>

To compare diversity measurements between different sample groups, e.g. different clinical

status, we performed a Bayesian estimation of group differences (BEST, [75]) , implemented using

the pymc3 package for the Python programming language; with priors ($\sim$) and deterministic

calculations (=) to assess differences in estimated group means as follows:

$\qquad g_1 \sim \text{Normal}(\mu = 15, \sigma = 15)$

$\qquad g_2 \sim \text{Normal}(\mu = 15, \sigma = 15)$

$\qquad \sigma_{g1} \sim \text{Uniform}(\text{low} = 1e{-}4, \text{high} = 30)$

$\qquad \sigma_{g2} \sim \text{Uniform}(\text{low} = 1e{-}4, \text{high} = 30)$

$\qquad \nu \sim \text{Exponential}(1/15) + 1$

$\qquad \lambda_1 = \sigma_{g1}^{-2}$

$\qquad \lambda_2 = \sigma_{g2}^{-2}$

$\qquad G1 \sim \text{StudentT}(\text{nu} = \nu, \text{mu} = g_1, \text{lam} = \lambda_1)$

$\qquad G2 \sim \text{StudentT}(\text{nu} = \nu, \text{mu} = g_2, \text{lam} = \lambda_2)$

$\qquad \Delta = G1 - G2$

Bayesian inference was performed using "No U-turn sampling"[76]. Highest density intervals

(HDI) of the posterior estimation of group differences (Δ) were used to determine statistical

certainty (***: 99% HDI >0 or <0, **: 95%HDI, *:90% HDI). The BEST code was implemented

following the pymc3 documentation.

<u>Cross-validated logistic regression to associate BSI cases with ASV composition</u>

We first removed ASVs with low prevalence (present in fewer than 5% of all samples), and low

abundances (maximum observed relative abundance <0.01) leaving 269 ASVs. We then scaled

the ASV relative abundances between 0 and 1 (min-max scaling) and performed logistic

regressions, relating ASV abundances to BSI status (1: BSI, 0: non-BSI) using the

sklearn.linear_model. LogisticRegressionCV module for the Python programming language with

an L1 (lasso) penalty, iterating over a range of regularization strengths ([0.01,0.1, 1., 10., 100.,

1000.]) using the "liblinear" solver. We retained the inferred ASV association coefficients with

non-zero values for each tested regularization strength to visualize the cross-validation path.

<u>Bayesian logistic regression</u>

We performed a Bayesian logistic regression to distinguish compositional differences between

infection-associated samples and samples from patients without secondary infections. We

modeled the infection state of patient sample i, y_i with a Binomial likelihood:

$y_i \sim \text{Binomial}(n = 1, p = p)$

$p = \text{inverse logistic}(\alpha + X_i\beta)$

$\alpha \sim \text{Normal}(\mu = 0, \sigma = 1)$

$\beta \sim \text{Normal}(\mu = 0, \sigma = 1)$

Where prior distributions are indicated by $\sim$; α is the intercept of the generalized linear model, β

is the coefficient vector for the $\log_{10}$-relative taxon abundances X_i in sample i or, in some cases,

the binary indicator variable for gut microbiome domination.

Data Availability

The raw sequencing data have been deposited on the Sequencing Reads Archive (SRA), and SRA

accession numbers are available for two bioprojects corresponding to the mouse sequencing data

PRJNA745367 (**Supplementary Table 4**) and the human stool samples PRJNA746322

(**Supplementary Table 5**).

Figures

Fig. 1. SARS-CoV-2 infection causes gut microbiome alterations in mice. **a** Timelines of fecal microbiota composition measured by 16S rRNA gene sequencing in mice infected with 0 or 10^4 PFU of SARS-CoV-2 ; time of infection=Day 1. Bars represent the composition of the 15 most abundant bacterial families per sample, blocks of samples correspond to an individual mouse's time course (x-axis label indicate experiment id, PFU, and mouse id). **b** α-diversity (inverse Simpson index) per infection group in the beginning (t_{start}) and at the end (t_{end}) of the experiment (n.s.: non-significant, **: p<0.01, one-tailed, paired t-test). **c** Principal coordinate plot of bacterial compositions in samples from the start (top) and end (bottom) of the experiment. **d** log_{10}-relative family abundances at the final time point; boxplots show median and interquartile ranges, whiskers extend to 1.5 times max- and min- quartile values, n.s.: not significant; *: p-value < 0.05; **: p-value < 0.01; ***: p-value < 0.001; Wilcoxon rank-sum tests. **e** Regression coefficients of the estimated changes in family abundances per day in mice infected with 10^4 PFU obtained from linear mixed effects models with varying effects per mouse and per cage (only significant coefficient results shown, abbreviations and colors as per the bacterial family legend).

Fig. 2 SARS-CoV-2 infection causes abnormalities in the gut epithelium of mice. a.
Representative H&E-stained section of the ileum depicting crypt-villus axes from K18-hACE2
mice on day 5-6 post intranasal inoculation with 10000 PFU SARS-CoV-2 or mock treatment.
Green arrows indicate goblet cells, scale bars correspond to 25μm. Bottom panels show high
magnification images of the indicated crypt with black arrowheads pointing at Paneth cells, scale
bars correspond to 10μm. **b.** Representative anti-lysozyme immunofluorescence images of the
ileal crypt (two images per group). White and orange doted circles delineate normal and
abnormal Paneth cells, respectively. Abnormality is characterized by distorted, depleted, or
diffuse lysozyme distribution patterns in Paneth cells. Lysozyme = red, DAPI = blue, scale bars
correspond to 10 m. **c.** Quantification of goblet cell number per villus (left), Paneth cells per
crypt (middle) based on H&E staining, and frequency of normal versus abnormal Paneth cell
lysozyme distribution pattern based on the immunofluorescence staining as depicted in b. Dots
represent the mean cell number per crypt-villus unit in each mouse, 50 units were counted per
mouse. Results were pooled from 3 independent experiments with n=3-5 mice per group for each
experiment. Boxplots indicate median and interquartile ranges (ns=non-significant, p>0.05; **,
p<0.01; ***, p<0.001; ****, p<0.0001 Mann-Whitney U-test). **d.** Correlation of Goblet cell
number per villus (left, Pearson correlation r=-0.48, p=0.015), Paneth cells per crypt (middle,
r=0.14, p-value=0.483) and frequency of abnormal Paneth cell lysozyme distribution pattern
(right, r=-0.5528, p=0.014) for the mice shown in c with α-diversity (inverse Simpson) of the gut
microbiome measured at the last day before sacrifice. **e.** Correlation of Goblet cell number per
villus (left, r=0.63, p<0.001), Paneth cells per crypt (middle, r=-0.29, p=0.149) and frequency of
abnormal Paneth cell lysozyme distribution pattern (right, r=0.65, p-value=0.003) for the mice
shown in c with $\log_{10}$-relative abundances of *Akkermansia* in fecal samples from the last day
before sacrifice; lines: univariate linear regression, shaded region: 95% CI.

Fig. 3. The dysbiotic gut microbiome in COVID-19 in patients from NYU Langone Health
(n=60) and Yale New Haven Hospital (n=36) is associated with secondary bloodstream
infections. **a** Bacterial family composition in stool samples (Yale, n = 63 samples; NYU, n = 67)
identified by 16S rRNA gene sequencing; bars represent the relative abundances of bacterial
families; red circles indicate samples with single taxa >50%. Samples are sorted by center and
bacterial α-diversity (inverse Simpson index, **b**). **c** α-diversity in samples from NYU Langone
Health and Yale New Haven Hospital; **p<0.01, two-sided T-test. **d** Average phylum level
composition per center. **e-g** Principal coordinate plots of all samples shown in **a**, labeled by center
(**e**), most abundant bacterial family (**f**) and domination status of the sample (**g**), and BSI status;
inset: boxplot of inverse Simpson index diversity by BSI (**h**). **i** Coefficients from a Bayesian
logistic regression with most abundant bacterial genera as predictors of BSI status. **j**
Counterfactual posterior predictions of BSI risk based on bacterial composition contrasting the
predicted risk of the average composition across all samples (red) with the risk predicted from a
composition where *Faecalibacterium* was increased by 10% (blue). **k** shotgun metagenomic reads
matched the species identified in clinical blood cultures in 70% of all investigated cases; the
histogram shows the distribution of $\log_{10}$-ratios of relative abundances of matched species in
corresponding stool samples to their corresponding mean abundances across all samples.

Acknowledgments

We thank René Niehus for helpful discussions on the implementation of the various Bayesian analyses. We thank the NYU Langone's Genome Technology Center, the NYU Langone's Experimental Pathology Research Laboratory and the NYU Langone's Microscopy Laboratory supported in part by NYU Langone Health's Laura and Isaac Perlmutter Cancer Center Support (grant P30CA016087) from the National Cancer Institute Langone and by the NIH S10 OD021747 grant for use of their instruments and technical assistance. We also thank the Office of Science & Research High-Containment Laboratories at NYU Grossman School of Medicine for their support in the completion of this research.

Yale IMPACT Team

Abeer Obaid, Alice Lu-Culligan, Allison Nelson, Anderson Brito, Angela Nunez, Anjelica Martin, Annie Watkins, Bertie Geng, Chaney Kalinich, Christina Harden, Codruta Todeasa, Cole Jensen, Daniel Kim, David McDonald, Denise Shepard, Edward Courchaine, Elizabeth B. White, Eric Song, Erin Silva, Eriko Kudo, Giuseppe DeIuliis, Harold Rahming, Hong-Jai Park, Irene Matos, Jessica Nouws, Jordan Valdez, Joseph Fauver, Joseph Lim, Kadi-Ann Rose, Kelly Anastasio, Kristina Brower, Laura Glick, Lokesh Sharma, Lorenzo Sewanan, Lynda Knaggs, Maksym Minasyan, Maria Batsu, Mary Petrone, Maxine Kuang, Maura Nakahata, Melissa Campbell, Melissa Linehan, Michael H. Askenase, Michael Simonov, Mikhail Smolgovsky, Nicole Sonnert, Nida Naushad, Pavithra Vijayakumar, Rick Martinello, Rupak Datta, Ryan Handoko, Santos Bermejo, Sarah Prophet, Sean Bickerton, Sofia Velazquez, Tara Alpert, Tyler Rice, William Khoury-Hanold, Xiaohua Peng, Yexin Yang, Yiyun Cao & Yvette Strong

Author contributions

LBR performed the mouse experiments with help from MGN, AMVJ. CZ performed mouse microbiome analyses with help from LBR, MV and KC. MV, JEA and JS prepared the samples from NYU. MV, JEA prepared the clinical data from NYU with help from JG, EW, BS. JK provided the data from Yale with help from ACM and the IMPACT team, AIK and AI. JS designed and performed the analyses with CZ, and help by GAH and APS. JS and KC designed the research question with support from VJT and BS. JS and KC wrote the manuscript with help

by LBR, MV and CZ. All other authors contributed materials, scientific feedback and

commented on the manuscript.

Funding

This work was in part funded by NYU Grossman School of Medicine startup research funds and

NIH/NIAID DP2 award (DP2AI164318) to JS, and the Yale School of Public Health and the

Beatrice Kleinberg Neuwirth Fund, as well as NIH grants to KC (DK093668, AI121244,

HL123340, AI130945, AI140754, DK124336), a Faculty Scholar grant from the Howard Hughes

Medical Institute (KC), Crohn's & Colitis Foundation (KC), Kenneth Rainin Foundation (KC),

Judith & Stewart Colton Center of Autoimmunity (KC). Further funding was provided by grants

from the NIH/NIAID to MD (R01AI143639 and R21AI139374), from the NIH to MV

(5T32AI100853), by Jan Vilcek/David Goldfarb Fellowship Endowment Funds to AMVJ, by

The G. Harold and Leila Y. Mathers Charitable Foundation to MD, and by NYU Grossman

School of Medicine Startup funds to MD and KAS, and the NYU Grossman School of Medicine

COVID-19 seed research funds to VJT, and funds from the NYU Langone Health Antimicrobial-

Resistant Pathogens Program to BS, AP, and VJT. KC and VJT also receive support from NIH

grant OT2HL161847. MN was supported by the American Heart Association Postdoctoral

Fellowship 19-A0-00-1003686. IMPACT received support from the Yale COVID-19 Research

Resource Fund. AI and DRL are Investigators of the Howard Hughes Medical Institute. AIK

received support from the Beatrice Kleinberg Neuwirth Fund, Bristol Meyers Squibb Foundation

and COVID-19 research funds from the Yale Schools of Public Health and Medicine.

Conflicts

KC has received research support from Pfizer, Takeda, Pacific Biosciences, Genentech, and

Abbvie; consulted for or received an honoraria from Puretech Health, Genentech, and Abbvie;

and holds U.S. patent 10,722,600 and provisional patents 62/935,035 and 63/157,225. JS is

cofounder of Postbiotics Plus Research LLC.

Bibliography

1. Fajgenbaum, D. C. & June, C. H. Cytokine Storm. *N. Engl. J. Med.* **383,** 2255–2273 (2020).
2. Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* **584,** 463–469 (2020).
3. Zuo, T. *et al.* Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization. *Gastroenterology* **159,** 944–955.e8 (2020).
4. Yeoh, Y. K. *et al.* Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19. *Gut* **70,** 698–706 (2021).
5. Gu, S. *et al.* Alterations of the gut microbiota in patients with coronavirus disease 2019 or H1N1 influenza. *Clin. Infect. Dis.* **71,** 2669–2678 (2020).
6. Nori, P. *et al.* Bacterial and fungal coinfections in COVID-19 patients hospitalized during the New York City pandemic surge. *Infect. Control Hosp. Epidemiol.* **42,** 84–88 (2021).
7. Grasselli, G. *et al.* Hospital-Acquired Infections in Critically Ill Patients With COVID-19. *Chest* (2021). doi:10.1016/j.chest.2021.04.002
8. Yu, D. *et al.* Low prevalence of bloodstream infection and high blood culture contamination rates in patients with COVID-19. *PLoS One* **15,** e0242533 (2020).
9. Langford, B. J. *et al.* Bacterial co-infection and secondary infection in patients with COVID-19: a living rapid review and meta-analysis. *Clin. Microbiol. Infect.* **26,** 1622–1629 (2020).
10. Shafran, N. *et al.* Secondary bacterial infection in COVID-19 patients is a stronger predictor for death compared to influenza patients. *Sci. Rep.* **11,** 12703 (2021).
11. Buffie, C. G. *et al.* Precision microbiome reconstitution restores bile acid mediated resistance to Clostridium difficile. *Nature* **517,** 205–208 (2015).
12. Buffie, C. G. & Pamer, E. G. Microbiota-mediated colonization resistance against intestinal pathogens. *Nat. Rev. Immunol.* **13,** 790–801 (2013).
13. Modi, S. R., Collins, J. J. & Relman, D. A. Antibiotics and the gut microbiota. *J. Clin. Invest.* **124,** 4212–4218 (2014).
14. Shimasaki, T. *et al.* Increased Relative Abundance of Klebsiella pneumoniae Carbapenemase-producing Klebsiella pneumoniae Within the Gut Microbiota Is Associated With Risk of Bloodstream Infection in Long-term Acute Care Hospital Patients. *Clin. Infect. Dis.* **68,** 2053–2059 (2019).
15. Kim, S., Covington, A. & Pamer, E. G. The intestinal microbiota: Antibiotics, colonization resistance, and enteric pathogens. *Immunol. Rev.* **279,** 90–105 (2017).
16. Morjaria, S. *et al.* Antibiotic-Induced Shifts in Fecal Microbiota Density and Composition during Hematopoietic Stem Cell Transplantation. *Infect. Immun.* **87,** (2019).
17. Niehus, R. *et al.* Quantifying antibiotic impact on within-patient dynamics of extended-spectrum beta-lactamase resistance. *Elife* **9,** (2020).
18. Taur, Y. *et al.* Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin. Infect. Dis.* **55,** 905–914 (2012).
19. Taur, Y. *et al.* Reconstitution of the gut microbiota of antibiotic-treated patients by autologous fecal microbiota transplant. *Sci. Transl. Med.* **10,** (2018).
20. Liao, C. *et al.* Compilation of longitudinal microbiota data and hospitalome from hematopoietic cell transplantation patients. *Sci. Data* **8,** 71 (2021).
21. Peled, J. U. *et al.* Microbiota as Predictor of Mortality in Allogeneic Hematopoietic-Cell Transplantation. *N. Engl. J. Med.* **382,** 822–834 (2020).

22. McCullers, J. A. The co-pathogenesis of influenza viruses with bacteria in the lung. *Nat.*
Rev. Microbiol. **12,** 252–262 (2014).
23. Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel
Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **323,** 1061–1069 (2020).
24. Westblade, L. F., Simon, M. S. & Satlin, M. J. Bacterial coinfections in coronavirus disease
2019. *Trends Microbiol.* **29,** 930–941 (2021).
25. Sepulveda, J. *et al.* Bacteremia and Blood Culture Utilization during COVID-19 Surge in
New York City. *J. Clin. Microbiol.* **58,** (2020).
26. Lansbury, L., Lim, B., Baskaran, V. & Lim, W. S. Co-infections in people with COVID-19:
a systematic review and meta-analysis. *J. Infect.* **81,** 266–275 (2020).
27. Sieswerda, E. *et al.* Recommendations for antibacterial therapy in adults with COVID-19 -
an evidence based guideline. *Clin. Microbiol. Infect.* **27,** 61–66 (2021).
28. Zhai, B. *et al.* High-resolution mycobiota analysis reveals dynamic intestinal translocation
preceding invasive candidiasis. *Nat. Med.* **26,** 59–64 (2020).
29. Haak, B. W. *et al.* Impact of gut colonization with butyrate-producing microbiota on
respiratory viral infection following allo-HCT. *Blood* **131,** 2978–2986 (2018).
30. Deriu, E. *et al.* Influenza Virus Affects Intestinal Microbiota and Secondary Salmonella
Infection in the Gut through Type I Interferons. *PLoS Pathog.* **12,** e1005572 (2016).
31. Yildiz, S., Mazel-Sanchez, B., Kandasamy, M., Manicassamy, B. & Schmolke, M. Influenza
A virus infection impacts systemic microbiota dynamics and causes quantitative enteric
dysbiosis. *Microbiome* **6,** 9 (2018).
32. Steed, A. L. *et al.* The microbial metabolite desaminotyrosine protects from influenza
through type I interferon. *Science* **357,** 498–502 (2017).
33. Abt, M. C. *et al.* Commensal bacteria calibrate the activation threshold of innate antiviral
immunity. *Immunity* **37,** 158–170 (2012).
34. Ichinohe, T. *et al.* Microbiota regulates immune defense against respiratory tract influenza A
virus infection. *Proc. Natl. Acad. Sci. USA* **108,** 5354–5359 (2011).
35. Sencio, V. *et al.* Influenza infection impairs the gut's barrier properties and favors secondary
enteric bacterial infection through reduced production of short-chain fatty acids. *Infect.*
Immun. (2021). doi:10.1128/IAI.00734-20
36. Wang, J. *et al.* Respiratory influenza virus infection induces intestinal immune injury via
microbiota-mediated Th17 cell-dependent inflammation. *J. Exp. Med.* **211,** 2397–2410
(2014).
37. Winkler, E. S. *et al.* SARS-CoV-2 Causes Lung Infection without Severe Disease in Human
ACE2 Knock-In Mice. *J. Virol.* **96,** e0151121 (2022).
38. Yinda, C. K. *et al.* K18-hACE2 mice develop respiratory disease resembling severe
COVID-19. *PLoS Pathog.* **17,** e1009195 (2021).
39. Zheng, J. *et al.* COVID-19 treatments and pathogenesis including anosmia in K18-hACE2
mice. *Nature* **589,** 603–607 (2021).
40. Golden, J. W. *et al.* Human angiotensin-converting enzyme 2 transgenic mice infected with
SARS-CoV-2 develop severe and fatal respiratory disease. *JCI Insight* **5,** (2020).
41. Cadwell, K. *et al.* A key role for autophagy and the autophagy gene Atg16l1 in mouse and
human intestinal Paneth cells. *Nature* **456,** 259–263 (2008).
42. Cadwell, K. *et al.* Virus-plus-susceptibility gene interaction determines Crohn's disease
gene Atg16L1 phenotypes in intestine. *Cell* **141,** 1135–1145 (2010).
43. Matsuzawa-Ishimoto, Y. *et al.* Autophagy protein ATG16L1 prevents necroptosis in the

intestinal epithelium. *J. Exp. Med.* **214,** 3687–3705 (2017).

44. Schluter, J. *et al.* The gut microbiota is associated with immune cell dynamics in humans.
Nature **588,** 303–307 (2020).

45. Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1 immunotherapy
in melanoma patients. *Science* **359,** 97–103 (2018).

46. Diefenbach, C. S. *et al.* Microbial dysbiosis is associated with aggressive histology and
adverse clinical outcome in B-cell non-Hodgkin lymphoma. *Blood Adv.* **5,** 1194–1198
(2021).

47. Sokol, H. *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium
identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci. USA*
105, 16731–16736 (2008).

48. Wrzosek, L. *et al.* Bacteroides thetaiotaomicron and Faecalibacterium prausnitzii influence
the production of mucus glycans and the development of goblet cells in the colonic
epithelium of a gnotobiotic model rodent. *BMC Biol.* **11,** 61 (2013).

49. Seibert, B. *et al.* Mild and Severe SARS-CoV-2 Infection Induces Respiratory and Intestinal
Microbiome Changes in the K18-hACE2 Transgenic Mouse Model. *Microbiol. Spectr.* **9,**
e0053621 (2021).

50. Sencio, V. *et al.* Alteration of the gut microbiota following SARS-CoV-2 infection
correlates with disease severity in hamsters. *Gut Microbes* **14,** 2018900 (2022).

51. Sokol, H. *et al.* SARS-CoV-2 infection in nonhuman primates alters the composition and
functional activity of the gut microbiota. *Gut Microbes* **13,** 1–19 (2021).

52. Zhang, F. *et al.* Prolonged Impairment of Short-Chain Fatty Acid and L-Isoleucine
Biosynthesis in Gut Microbiome in Patients With COVID-19. *Gastroenterology* **162,** 548–
561.e4 (2022).

53. Gaebler, C. *et al.* Evolution of antibody immunity to SARS-CoV-2. *Nature* **591,** 639–644
(2021).

54. Park, S.-K. *et al.* Detection of SARS-CoV-2 in Fecal Samples From Patients With
Asymptomatic and Mild COVID-19 in Korea. *Clin. Gastroenterol. Hepatol.* **19,** 1387–
1394.e2 (2021).

55. Xiao, F. *et al.* Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology*
158, 1831–1833.e3 (2020).

56. Cheung, K. S. *et al.* Gastrointestinal Manifestations of SARS-CoV-2 Infection and Virus
Load in Fecal Samples From a Hong Kong Cohort: Systematic Review and Meta-analysis.
Gastroenterology **159,** 81–95 (2020).

57. Lamers, M. M. *et al.* SARS-CoV-2 productively infects human gut enterocytes. *Science* **369,**
50–54 (2020).

58. Cao, J. *et al.* Integrated gut virome and bacteriome dynamics in COVID-19 patients. *Gut*
Microbes **13,** 1–21 (2021).

59. Klag, T., Stange, E. F. & Wehkamp, J. Defective antibacterial barrier in inflammatory bowel
disease. *Dig. Dis.* **31,** 310–316 (2013).

60. Ramanan, D. & Cadwell, K. Intrinsic defense mechanisms of the intestinal epithelium. *Cell*
Host Microbe **19,** 434–441 (2016).

61. Schluter, J. & Foster, K. R. The evolution of mutualism in gut microbiota via host epithelial
selection. *PLoS Biol.* **10,** e1001424 (2012).

62. McLoughlin, K., Schluter, J., Rakoff-Nahoum, S., Smith, A. L. & Foster, K. R. Host
selection of microbiota via differential adhesion. *Cell Host Microbe* **19,** 550–559 (2016).

63. Fernandez-Castañer, M. *et al.* Evaluation of B-cell function in diabetics by C-peptide
determination in basal and postprandial urine. *Diabete Metab* **13,** 538–542 (1987).
64. Yu, S. *et al.* Paneth Cell-Derived Lysozyme Defines the Composition of Mucolytic
Microbiota and the Inflammatory Tone of the Intestine. *Immunity* **53,** 398–416.e8 (2020).
65. Salzman, N. H. *et al.* Enteric defensins are essential regulators of intestinal microbial
ecology. *Nat. Immunol.* **11,** 76–83 (2010).
66. van der Lugt, B. *et al.* Akkermansia muciniphila ameliorates the age-related decline in
colonic mucus thickness and attenuates immune activation in accelerated aging Ercc1-/Δ7
mice. *Immun. Ageing* **16,** 6 (2019).
67. Wang, L. *et al.* An observational cohort study of bacterial co-infection and implications for
empirical antibiotic therapy in patients presenting with COVID-19 to hospitals in North
West London. *J. Antimicrob. Chemother.* **76,** 796–803 (2021).
68. Labarta-Bajo, L. *et al.* Type I IFNs and CD8 T cells increase intestinal barrier permeability
after chronic viral infection. *J. Exp. Med.* **217,** (2020).
69. Karki, R. *et al.* Synergism of TNF-α and IFN-γ Triggers Inflammatory Cell Death, Tissue
Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock Syndromes. *Cell*
184, 149–168.e17 (2021).
70. Giron, L. B. *et al.* Plasma Markers of Disrupted Gut Permeability in Severe COVID-19
Patients. *Front. Immunol.* **12,** 686240 (2021).
71. Xie, X. *et al.* An Infectious cDNA Clone of SARS-CoV-2. *Cell Host Microbe* **27,** 841–
848.e3 (2020).
72. Gohl, D. M. *et al.* Systematic improvement of amplicon marker gene methods for increased
accuracy in microbiome studies. *Nat. Biotechnol.* **34,** 942–949 (2016).
73. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*
73, 5261–5267 (2007).
74. Pawlowsky-Glahn, V., Egozcue, J. J. & Tolosana-Delgado, R. *Modelling and analysis of*
compositional data. (John Wiley & Sons, Ltd, 2015). doi:10.1002/9781119003144
75. Kruschke, J. K. Bayesian estimation supersedes the t test. *J. Exp. Psychol. Gen.* **142,** 573–
603 (2013).
76. Homan, M. D. & Gelman, A. The No-U-Turn Sampler: Adaptively Setting Path Lengths in
Hamiltonian Monte Carlo. *J. Mach. Learn. Res.* **15,** 1593–1623 (2014).
821
822

Supporting Information

824
825

Extended Data Fig. S1 SARS-CoV-2 infection in K18-hACE2 mice.

a-b. Following inoculation with 0, 10, 100, 1000 or 10000 PFU of SARS-CoV-2 or mock

infection, mice were monitored daily for weight loss (a) and signs of disease quantified by a

composite score based on ruffled fur, hunched back, heavy breathing and absence of mobility

(b). Median and interquartile range determined for each group at each time point are depicted.

Results are pooled from 1-3 independent experiments. For each group, the total number of mice

is indicated. c. Viral burden in lung or intestinal tissue of K18-hACE2 mice was analyzed at 5-6

days after infection with 100, 1000, 10000 PFU of SARS-CoV-2 or mock infection by qRT-

PCR. Dots represent the copy number of N RNA per μg of RNA calculated for each mouse.

Results were pooled from 1 (100 and 1000 PFU doses) or 2 (mock and 10000 PFU) independent

experiments with n=2-5 mice per group for each experiment. The median and interquartile range

are depicted for each experimental group. The dotted line depicts the limit of detection.

Extended Data Fig. S2 Inconsistent microbiomes dynamics in mice with lower infection

doses. a Bars represent bacterial family compositions in stool samples collected from mice over

time, mouse time courses grouped as indicated by boxes. **b** bacterial alpha diversity in first (t_{start})

and last (t$_{end}$) samples collected. **c** principal coordinate plots of bacterial compositions in first and

last samples colored by infection dose (in PFU). **d** bacterial family abundances by infection dose

at the final sample collected. **E** diversity, weight and temperature z-scores (calculated from all

data points) over time per mouse as shown in a and Fig. 1. **F** untransformed diversity, weights

and temperatures relative to the beginning of the experiment.

Extended Data Fig. S3 Some intestinal parameters are not modified during SARS-CoV-2

infection. K18-hACE2 mice were analyzed on day 5-6 post intranasal inoculation with 10000 PFU

SARS-CoV-2 or mock treatment. **a.** Quantification of fluorescence intensity in the blood following

oral administration of FITC-dextran. **B.** Intestinal fatty acid-binding protein (iFABP), LPS-binding

protein (LBP), and citrulline concentration in plasma. **C.** Quantification of colon length. **d.**

Quantification of villus length in the duodenum (left) and ileum (right) based on H&E staining. **E.**

Quantification of goblet cell number (left) and Paneth cell number (middle) per crypt-villus unit
in the proximal duodenum based on H&E staining and calculation of goblet cell per Paneth cell
ratio based on these quantifications (right). Individual mice, represented by the circles as well as
the median and interquartile ranges are depicted. In d, e, each circle shows the mean for each
mouse of the cell number counted per crypt-villus unit on 50 units. Results were pooled from 2
(for a) or 3 independent experiments with n=3-5 mice per group for each experiment. Significant
differences were determined using the Mann-Whitney U test (ns=non-significant, $p > 0.05$; **,
$p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$).

867
868

Extended Data Fig. S4 Strongest gut dysbiosis is correlated with markers of defects in the

intestinal barrier and epithelium. A Reproduction of Fig. 1 showing bacterial compositions in

mice infected with 10^4 PFUs, highlighting four mice time courses of mice with lowest diversity

872 and highest disease scores at the end of the experiment (**b**). **c-d** Correlations between alpha

873 diversity (**c**) (inverse Simpson) and $\log_{10}$ relative *Akkermansia* abundances (**d**) at the end of the

874 experiment with epithelium phenotypes and gut barrier integrity markers measured in the blood

875 of mice (data from mice highlighted in **a** with circles in corresponding colors, lines: linear

876 regression, shaded region: 95%CI).

Extended Data Fig. S5 a Samples from patients requiring ICU transfer have lower diversity on average (p=0.005, Wilcoxon rank-

sum); bars as in Fig. 1 with ICU status of patients and domination state of samples indicated. **b** Genus abundances in samples with a

single genus >50% relative abundance.

Extended Data Fig. S6 Patients with a positive clinical blood culture result (BSI) received

antibiotics, prior or on the day of blood culture results (cross symbol: first recorded antibiotic

administration, blue: sequenced stool sample, diamond: positive blood culture result (BSI)).

Extended Data Fig. S7 a Posterior coefficient estimates from a Bayesian logistic regression

regressing $\log_{10}$ relative abundances of the top 10 most abundant bacterial genera on BSI status

using only BSI cases with associated stool samples taken prior or on the day of a confirmed

positive blood culture. **b** Posterior coefficient estimates from a Bayesian logistic regression

regressing $\log_{10}$ relative abundances of the top 10 most abundant bacterial genera on BSI status

with domination status of the microbiome as an additional predictor (domination: >50% of the

composition by one taxon). **c** ASVs associated with samples from patients with BSI. Coefficients

from a cross-validated, L1-penalized logistic regression correlating the binary outcome (BSI)

with $\log_{10}$-transformed relative ASV abundances. **d** Cross-validation paths; for all regularization

strengths (L1-penalty) used, a *Faecalibacterium* ASV was most negatively associated with BSI-

positive samples.

Extended Data Fig. S8 *Faecalibacterium* relative abundance is positively correlated with

bacterial alpha diversity. Log10 transformed relative abundances of the genus

Faecalibacterium in stool samples from patients are correlated with the inverse Simpson

diversity index; line from linear regression, shaded region: 95%CI.

Extended Data Fig. S9 Bacteria in stool of COVID-19 patients match taxa identified blood

cultures. a Organisms identified in blood cultures together with bars representing the bacterial

family compositions in stool samples; multiple samples belonging to the same patient grouped

by a white box. Two samples with matching whole genome sequenced (WGS) blood isolates

indicated. **b** Rank analysis of abundance patterns in stool samples from different BSI categories;

a filled circle indicates the calculated rank of the focal BSI category (row) in terms of the

corresponding taxon stool abundance relative to samples from other BSI categories (Lact:

Lactobacillales, Enbct: Enterobacterales; Pseu: Pseudomonadales, Bact: Bacteroidales, Staph:

Staphylococcales. Only 5 out of 7 BSI categories are shown because fungal BSIs and the

uninfected category have no corresponding bacterial stool abundances). **c,d** left: neighbor-joining

tree constructed from all NCBI RefSeq assemblies of *Staphylococcus aureus* genomes in

addition to isolates that were isolated from subjects highlighted in **a**. right: counts of perfect read

matches of shotgun metagenomic reads from stool samples, red: stool sample sequencing read

matches to WGS of isolates from the same patient, black: matches to other genomes.

Supplementary Table 1: Clinical characteristics of patients with confirmed COVID-19 at NYU Langone Health and Yale New Haven Hospital

	NYU, N = 60	YALE, N = 36
Age (years)	51 ± 17.5	62.52 ± 19.72
Sex (F | M)	42% | 58%	39% | 61%
Hospital course and Outcomes		
ICU Admission	53%	65%
Pneumonia	42%	77%
Diarrhea	13%	32%
Intubation	36%	41%
Sepsis	23%	18%
Encephalopathy	12%	3%
Death	5%	21%
Length of stay (median, IQR)	37 (10-86)	27 (11-35.25)
Risk Factors		
Cancer within 1 year	7%	4%
Chronic Heart Disease	18%	36%
Hypertension	38%	64%
Chronic Lung Disease	7%	20%
Immunosuppression	17%	4%

927 **Supplementary Table 2: Clinical characteristics of COVID-19 patients at NYU Langone**

928 **Health and Yale New Haven Hospital with and without positive blood culture results (BSI).**

	BSI, N = 26	non-BSI N = 53
Hospital course and Outcomes		
ICU Admission	69%	64%
Pneumonia	73%	53%
Diarrhea	31%	64%
Intubation	58%	36%
Sepsis	35%	21%
Encephalopathy	19%	6%
Death	15%	9%
Length of stay (median, IQR)	59 (23-91.5)	22 (6-51)

930
931

933　**Supplementary Table 3: Shotgun metagenomic reads mapped to species identified in clinical blood cultures**. Dark grey shading:

934　no sequencing reads from stool samples matched the species identified in clinical blood samples, light grey shading: species of the

935　same genus but not the same species had non-zero read counts in stool samples. The relative abundance of identified species were

936　contrasted with their mean abundances (log10 ratio).

Organism identified in blood	species identified in stool sample	Log ratio
Bacteroides thetaiotaomicron	Bacteroides_thetaiotaomicron_14-106904-2	2.92
Enterococcus faecalis Group D	Enterococcus_faecalis_LD33	1.8
Escherichia coli	Escherichia_coli_K-12_substr._W3110	2.2
Escherichia coli	Escherichia_coli_IAI39	1.6
Escherichia coli	Escherichia_coli_536	2.8
Klebsiella pneumoniae	Klebsiella_pneumoniae_KPNIH27	-1.7
Lactobacillus species	Lactobacillus_curvatus_WiKim38	3.9
Pseudomonas aeruginosa	Pseudomonas_aeruginosa_SJTD-1	3.3
Serratia marcescens	Serratia_marcescens_CAV1492	-0.2
Staphylococcus aureus	Staphylococcus_aureus_RF122	1.9
Proteus mirabilis	Proteus_mirabilis;t__Proteus_mirabilis_BB2000	0.56
Acinetobacter lwolfii	Acinetobacter_calcoaceticus_EGD_AQ_BF14	-0.6
Staphylococcus	Staphylococcus_sp._HMSC063G01_HMSC063G01	0.9
Staphylococcus	Staphylococcus_epidermidis_W23144	3.3
Staphylococcus aureus	not found	
Staphylococcus hominis	not found	
Staphylococcus capitis	not found	
Staphylococcus epidermidis, hominis	Staphylococcus_pseudintermedius_063228	2.2
Staphylococcus epidermidis, hominis ssp hominis	not found	
Staphylococcus epidermidis	Staphylococcus_aureus_JKD6008	2.7
Staphylococcus epidermidis	Staphylococcus_epidermidis_DAR1907	1.1
Staphylococcus capitis	Staphylococcus_sp._HMSC067F07_HMSC067F07	3.7
Staphylococcus epidermidis, hominis ssp hominis	Staphylococcus_hominis_793_SHAE	0.8
Staphylococcus epidermidis	Staphylococcus_sp._HMSC070D05_HMSC070D05	3.8
Staphylococcus hominis, epidermidis	Staphylococcus_hominis_MMP2	1.0
Staphylococcus hominis, epidermidis	Staphylococcus_epidermidis_ATCC12228 GCF7645.1	0.2

Supplementary Table 4: SRA accession numbers for the bioproject PRJNA745367

corresponding to the mouse sequencing data.

(Excel sheet)

Supplementary Table 5: SRA accession numbers for the bioproject PRJNA746322

corresponding to the human stool samples sequencing data.

(Excel sheet)

946