Title: Ammonium regulates the development of pine roots through hormonal crosstalk and
differential expression of transcription factors in the apex

Authors: Francisco Ortigosa[1], César Lobato-Fernández[1], Hitomi Shikano[2], Concepción
Ávila[1], Shu Taira[2], Francisco M. Cánovas[1], Rafael A. Cañas[1,3]*.

Affiliation: [1]Grupo de Biología Molecular y Biotecnología, Departamento de Biología
Molecular y Bioquímica, Universidad de Málaga, Campus Universitario de Teatinos,
29071-Málaga, Spain
[2]Faculty of Food and Agricultural Sciences, Fukushima University, Kanayagawa,
Fukushima 960-1296, Japan.
[3]Integrative Molecular Biology Lab, Universidad de Málaga, Campus Universitario de
Teatinos, 29071-Málaga, Spain

*Correspondence: Rafael A. Cañas, Departamento de Biología Molecular y Bioquímica,
Facultad de Ciencias, Universidad de Málaga, Campus Universitario de Teatinos s/n, E-
29071 Málaga, Spain. e-mail: rcanas@uma.es Phone number: +34-925-131673 Fax
number: +34-952-132041.

ORCID: FO (0000-0002-7517-7594); CLF (0000-0001-9576-7236); HS (0000-0001-8973-
3355); CA (0000-0001-8817-7529); ST (0000-0002-9243-0395); FMC (0000-0002-4914-
2558) and RAC (0000-0001-9727-5585)

ABSTRACT

Ammonium is a prominent source of inorganic nitrogen for plant nutrition, but excessive amounts can be toxic for many species. However, most conifers are tolerant to ammonium, a relevant physiological feature of this ancient evolutionary lineage. For a better understanding of the molecular basis of this trait, ammonium-induced changes in the transcriptome of maritime pine (*Pinus pinaster* Ait.) root apex have been determined by laser capture microdissection and RNA sequencing. Ammonium promoted changes in the transcriptional profiles of multiple transcription factors, such as *SHORT-ROOT*, and phytohormone-related transcripts, such as *ACO*, involved in the development of the root meristem. Nano-PALDI-MSI and transcriptomic analyses showed that the distributions of IAA and CKs were altered in the root apex in response to ammonium nutrition. Taken together, the data suggest that this early response is involved in the increased lateral root branching and principal root growth, which characterize the long-term response to ammonium supply in pine. All these results suggest that ammonium induces changes in the root system architecture through the IAA-CK-ET phytohormone crosstalk and transcriptional regulation.

KEYWORDS: Transcriptomics; conifer; *Pinus pinaster*; nitrogen nutrition; laser capture microdissection; root development; phytohormones

INTRODUCTION

Nitrogen (N) is a vitally important nutrient for all living organisms because it is a constituent of essential biomolecules such as nucleic acids, proteins, amino acids, porphyrins and hormones, among others (Miller & Cramer, 2005). This nutrient is indispensable for the proper growth and development of plants, and it can be assimilated from different kinds of sources, including organic (peptides, amino acids, and urea) and inorganic (nitrate and ammonium) forms (Näsholm et al., 1998, Hachiya & Sakakibara, 2017). Together with nitrate (NO_3^-), ammonium (NH_4^+) is one of the main forms of inorganic N available for plants, and the relative proportions of these elements in soil can vary depending on biological and climate conditions (Bijlsma et al., 2000).

Experimental evidences strongly suggest that NH_4^+ is perceived and recognized by plant cells as a signal that promotes physiological and morphological changes in plants (Liu and von Wirén, 2017). However, NH_4^+ at millimolar levels usually causes toxicity in most plants. The effects caused by an excessive NH_4^+ availability include decreased plant growth, leaf chlorosis, decreased root/shoot ratios, decreased root gravitropism and altered root system architecture (RSA) (Esteban et al., 2016). Regarding the changes in RSA, the inhibition of root elongation, enhancement of lateral root (LR) branching and impaired root gravitropism are commonly observed (Y. Liu & von Wirén, 2017).

Root elongation involves three interconnected biological processes: cell division, cell expansion and cell differentiation (Youssef et al., 2018). Both cell division and cell expansion are affected by NH_4^+ through two independent determinants: i) a decreased capacity for protein N-glycosylation, and ii) an increased production of reactive oxygen species (ROS) (Jia & von Wirén, 2020). In *Arabidopsis,* root growth inhibition in response to NH_4^+ represses root cell production by decreasing the meristem size and the number of dividing cells without altering the cell division rate (Y. Liu et al., 2013). Furthermore, NH_4^+ also decreases the number of root cap cells (Y. Liu et al., 2013). In this sense, several transcription factors (TFs) have been described to play key roles in root cap development, such as BEARSKIN (BRN) and SOMBRERO (SMB) (Bennett et al., 2010, 2014; Kamiya et al., 2016). However, the transcriptional pathway regulating NH_4^+ root cap development has not yet been deciphered.

In addition, auxins (IAAs) play a prominent role in cell division and cell expansion (Wang & Ruan, 2013). IAAs can be synthesized in the root meristem and transported toward

upstream adjacent regions through specific transporters, such as AUXIN RESPONSE 1 (AUX1) and PINFORMED 1-7 (PIN1-7) (Grunewald & Friml, 2010). IAAs induce the expression of PLETHORA 1-4 (PLT1-4) TFs in the root meristem, which are responsible for cell proliferation maintenance (Galinha et al., 2007). Other TFs have been described to play key roles in the stem cell niche and the quiescent center (QC) maintenance to produce all tissues required to form a mature root, including the TFs SHORT-ROOT (SHR) and SCARECROW (SCR) (Sablowski, 2011). *SHR* expression is localized in a zone of the stele that constitutes the central part of the root and stem (Miyashima et al., 2011; Kim et al., 2020). When *SHR* transcripts are translated, SHR proteins move into adjacent cells (pericycle cells, endodermis, QC and phloem pole) to activate the expression of SCR (Helariutta et al., 2000; Nakajima et al., 2000; Sena et al., 2004; Cui et al., 2007). SHR also plays key roles in phloem development, controlling the asymmetric cell division process for sieve element development by the regulation of NARS1 and SND2 NAC-type TFs (Kim et al., 2020). This fact is relevant because these cell types constitute the phloem (Greb, 2020), which is the main means of IAA transport (Chapman & Estelle, 2009). However, Y. Liu et al. (2013) showed that NH_4^+ does not affect the IAA content at QC, where the maximum level of this phytohormone in roots is localized (Zhou et al., 2010).

As it was mentioned above, LR branching is enhanced by NH_4^+ supply (Araya et al., 2016). When the quadruple ammonium transporter (AMT) knockout lines, that exhibit a severe reduction in LR branching, were complemented with *AtAMT1.1* or *AtAMT1.3*, only *AtAMT1.3*-complemented plants were able to restore the LR branching phenotype induced by NH_4^+ (Lima et al., 2010), suggesting that the NH_4^+-induced LR branching signaling events are dependent on AtAMT1.3 (Lima et al., 2010). Recently, it has been demonstrated that under NH_4^+ supply, RSA changes mediated by IAAs transport are highly linked to root acidification (Jia et al., 2020; Meier et al., 2020), which seems to be the molecular basis for the AMT transporter effect on LR branching since NH_4^+ uptake is accompanied by pH imbalance. In addition, IAAs have been suggested to be involved in this signaling event due to the repression of PIN2 under a supply of NH_4^+ (Y. Liu et al., 2013; Zou et al., 2013), which is provoked by the PIN2 hyperphosphorylation (Ötvös et al., 2021). Regarding root agravitropic response, NH_4^+ promotes the downregulation of AUX1 and PIN2, which are two pivotal IAA transporters and are subject to the antagonistic action between PIN2 and ARG1. ARG1 is involved in the transduction of the root gravity signal and required for

normal AUX1 expression and basipetal IAA transport in root apices, promoting an
asymmetric auxin flow (Y. Liu et al., 2013; Zou et al., 2013). All these evidences highlight
the importance of IAA in orchestrating the root system configuration in response to this
nutritional stimulus. However, the transcriptional regulatory mechanisms that control this
response remain unclear.

Together with IAAs, cytokinins (CKs) have been previously established to be important in
root plant growth and vascular development (Kamada-Nobusada et al., 2013; Miyashima et
al., 2019; Mao et al., 2020). CK biosynthesis and activity in plants are closely related to N
availability (Takei et al., 2004; Kamada-Nobusada et al., 2013). In the roots of rice, NH_4^+
nutrition leads to the accumulation of different CKs and CK-derived compounds (Kamada-
Nobusada et al., 2013). During active growth of *Arabidopsis*, the content of CKs first
increased in the vascular system, thus reflecting CKs transport from roots to shoots
(Shtratnikova et al., 2015). Recently, it has been reported that CK signaling in the early
protophloem-sieve-element cell files of *Arabidopsis* root procambial tissue promotes the
expression of several DOF TFs (Miyashima et al., 2019). Together with the IAA-
responsive dependent HD-ZIP III proteins, these TFs compose a transcriptional network
that integrates spatial information of the hormonal domains and miRNA gradients, which
are essential for root vascular development (Miyashima et al., 2019).

Previous works in *Arabidopsis*, based on root protoplast generation and cell-sorting using
FACS coupled to expression studies, revealed cell-specific responses to different processes
such as N nutrition (Gifford et al., 2008; Walker et al., 2017) or plant immunity (Rich-
Griffin et al 2020). Regarding N nutrition, co-expression network approaches revealed that
within the root cortex N controlled a wide range of processes including phytohormone
responses, while in pericycle regulatory networks were related to formation of new organs,
organ structure development, and establishment of cell localization (Walker et al., 2017)
showing that phytohormones are candidates signaling for cell-specific responses to N
(Gifford et al., 2008).

Conifers are an ancient group of gymnosperms that cover vast regions in the Northern
Hemisphere, and they have exceptional ecologic and socioeconomic importance (Farjon et
al., 2018). Since conifers represent a differentiated evolutionary lineage of plants, they
exhibit substantial differences in N metabolism with respect to angiosperms such as
Arabidopsis. For instance, conifers lack glutamine biosynthesis in the plastid of

photosynthetic cells (Cánovas et al., 2007), and most species prefer NH_4^+ over NO_3^- as the
main source of inorganic N (Kronzucker et al., 1997; Hawkins & Robbins, 2010). This is
the case for maritime pine (*Pinus pinaster* Aiton) (Warren & Adams, 2002; Ortigosa et al.,
2020), a southwestern Mediterranean conifer employed as a model in studies of N nutrition
and metabolism and for which a large body of genomics resources are available (Cañas et
al., 2017; Ortigosa et al., 2020; 2021). Previous transcriptomic studies of pine seedling
tissues provided an overview of the gene expression distribution in different pine tissues
but also it was possible to highlight the relationships between gene expression patterns and
function in a tissue-dependent manner (Cañas et al., 2014; 2017).

Maritime pine seedlings under NH_4^+ supply accumulate more biomass than those fed with
NO_3^- what is related with a higher uptake of NH_4^+ than NO_3^- (Ortigosa et al., 2020). In
maritime pine roots, a long-term supply of NH_4^+ induced the expression of transcripts
related to defense, such as *antimicrobial peptide 1* (*PpAMP1*) (Canales et al., 2010), which
is not found in dicots and can regulate the NH_4^+ uptake (Canales et al., 2011). There also
was a close relationship between NH_4^+-responsive genes and genes involved in amino acid
metabolism, particularly those involved in asparagine biosynthesis and utilization (Canales
et al., 2010). Additionally, NH_4^+ promotes changes in the epitranscriptome that mainly
regulate the translational response and growth, including the repression of 1-
aminocyclopropane-1-carboxylic acid (ACC) oxidase (ACO), the last enzyme in the
ethylene (ET) biosynthesis pathway (Ortigosa et al., 2021). Considering this background,
the main goal of the present work was to decipher the molecular mechanisms underlying
the early response to NH_4^+ in the root apex and its relationship with root development in
maritime pine.

MATERIALS AND METHODS

Plant material

Maritime pine seeds (*P. pinaster* Aiton) from "Sierra Segura y Alcaraz" (Albacete, Spain) were provided by the *Área de Recursos Genéticos Forestales* of the Spanish *Ministerio de Agricultura, Pesca y Alimentación*. Seed germination was carried out following the protocol described elsewhere (Cañas et al., 2006). Seedlings were grown in vermiculite in plant growth chambers (Aralab Fitoclima 1200, Rio de Mouro, Portugal) under 16 h light photoperiod, a luminal intensity of 125 $\mu mol\ m^{-2}\ s^{-1}$, a constant temperature of 23 °C, 50% relative humidity and watered twice a week with distilled water. One-month-old maritime pine seedlings were used for the experiments. Pine seedlings were randomly subdivided into three different groups, relocated into forestall seedbeds and irrigated with 80 mL of a solution that contains macro- and micronutrients without any N source (1.16 mM KCl; 0.63 mM KH_2PO_4; 0.35 mM $MgSO_4 \cdot 7H_2O$; 0.17 mM $CaCl_2 \cdot H_2O$; 80 μM EDTA-$FeSO_4$; 25.9 μM H_3BO_3; 10.2 μM $MnCl_2 \cdot 4H_2O$; 1.3 μM $ZnSO_4 \cdot 7H_2O$; 0.7 μM $CuSO_4 \cdot 5H_2O$; 0.1 μM $Na_2MoO_4 \cdot 2H_2O$). After three days of acclimation, the control group was irrigated with 80 mL of water and the experimental group was supplied with 80 mL of 3 mM NH_4Cl. Root samples were collected 24 h after the treatment and immediately frozen in liquid N (Figure 1).

Ten seedlings were collected and pooled per each sample for RT-qPCR validation. The same experiment was carried out three independent times. The screening condition and adequate development of each experiment was verified through the gene expression analysis by RT-qPCR of control genes following previous results (Ortigosa et al., 2021) (Figure 1). To perform laser capture microdissection (LCM), seedling's root apexes were cut, and tissues (5-6 mm) were imbided in a specimen holder with Tissue-Tek optimal cutting temperature embedding medium (Sakura Finetek, Alphen aan den Rijn, The Netherlands) and immediately frozen in liquid N for cryostat sectioning. Frozen samples were stored at -80°C until use.

Root phenotyping in response to nitrogen supply

For the root phenotypes under different nitrogen sources (NH_4^+ and NO_3^-), the pine seedlings were cultivated in a growth chamber with a 16/8 h light/dark photoperiod, light intensity of 125 $\mu mol\ m^{-2}\ s^{-1}$, constant temperature of 23°C, and 50% relative humidity

(Aralab Fitoclima 1200, Rio de Mouro, Portugal). One-month-old seedlings (n = 30) were

randomly transplanted into seedbeds for each experimental condition: distilled water, 3 mM

KNO_3 and 3 mM NH_4Cl. Each experimental condition group was irrigated twice per week

with 50 mL of the corresponding N solution or 50 mL of distilled water for 74 days. After

treatment, seedling roots were dissected and whole roots, primary root (PR) and LRs were

weighted. The PR length and number of LRs was manually assessed (Figure S1).

Whole root RNA isolation

201   Total RNA from roots of maritime pine seedlings was isolated following the protocol

202   described by Liao et al. (2004) and modified by Canales et al. (2012). RNA concentration

203   and purity were determined spectrophotometrically using a Nanodrop ND-1000 (Thermo

204   Scientific, Waltham, MA, USA). Purity of the preparation was assessed by determining the

205   260/280 and 260/230 ratios. The integrity was assessed by electrophoresis.

Laser capture microdissection, RNA isolation and low-input RNA-seq

LCM procedure was carried through as previously described (Cañas et al., 2014). Full step

protocol is described in Methods S1. Four different tissue areas were isolated by

microdissection corresponding to the root cap (RC), meristem (RM), developing cortex

(RDC) and developing vessels (RDV) areas (Video S1).

All RNA extractions from the microdissection procedure were carried out using

manufacturer's instruction protocol (non-LCM) for the RNAqueous-Micro RNA Isolation

Kit (Ambion, Austin, TX, USA). RNA quality, DNA contamination and first quantification

were performed via RNA Pico Assay for the 2100 Bioanalyzer (Agilent, Santa Clara, CA,

USA). Quantification was verified via a Qubit RNA BR (Broad-Range) Assay Kit

(Invitrogen, Paisley, UK). RNA samples with RNA integrity number (RIN) higher than 7

were used for subsequent RNA sequencing, mRNA amplification and cDNA synthesis 1 ng

of RNA for each sample.

The low input RNA-seq was carried out by Novogen (Hong Kong). The cDNA synthesis

and amplification, and the library preparation was made with the SMART-Seq™ v4

Ultra™ Low Input RNA Kit for Sequencing (Takara, Mountain View, CA, USA) following

the manufacturer's instructions. RNA sequencing was made in a NovaSeq 6000 sequencer

according to the manufacturer's instructions for paired-end reads (Illumina, San Diego, CA,

USA). The 24 samples were sequenced producing paired-end reads of 150 bp length. The

sequencing output is shown in the Table S1.

The raw reads were trimmed (quality and contamination) using SeqTrimBB software

(https://github.com/rafnunser/seqtrimbb). Only the pairs in which both reads passed the

quality test were further analyzed (Q>20). Trimmed reads are shown in the Table S1. These

reads were assembled using Trinity 2.11 (Haas et al., 2013). Contigs lower than 400 pb

were eliminated, for the rest of contigs the redundancy was reduced using CD-HIT-EST

software (Fu et al., 2012). This transcriptome shotgun assembly project has been deposited

at DDBJ/EMBL/GenBank under the accession GJFX00000000. The version described in

this paper is the first version, GJFX01000000. The final transcriptome was used as the

reference for the read mapping that was performed with BWA using the MEM option (Li

and Durbin, 2009). The read count was obtained with the phyton script *sam2counts*

(https://github.com/vsbuffalo/sam2counts). Differentially expressed (DE) transcripts were

identified using the edgeR package for R, the transcripts were normalized by cpm and

filtered; 2 cpm in at least 2 samples (Robinson et al., 2010). Each sample was from a single

seedling in a different experimental replicate. The samples were basically grouped by tissue

and nutritional condition (Figure S2). For the tissue-treatment interaction only the

transcripts with FDR < 0.05 and the three experimental replicates with the same expression

sense than the final logFC (positive or negative) were considered as differentially expressed

(DE). For the tissue analysis same parameters have been considered. The DE transcripts

were used to construct a gene co-expression network. An unsigned network has been

carried out using the R package WGCNA soft-thresholding power value of 9 (Langfelder &

Horvath, 2008). From the network, the 10% of the transcripts with more connections in

each module were considered hub genes.

These RNA-Seq data have been deposited in the NCBI's Gene Expression Omnibus (Edgar

et al., 2002) and are accessible through GEO Series with the accession number GSE175587

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE175587). Additionally, RNA-

seq and network results are accessible through a database in html format that can be

installable   with   R   packages   and   downloaded   from   GitHub

(https://github.com/ceslobfer/Rootapp).

Total RNA (1 ng) was retrotranscribed and amplified to verify the expression analyses for

several DE genes by RT-qPCR. The cDNA synthesis and amplification protocol was

carried out using the Conifer RNA Amplification (CRA+) protocol previously described by
Cañas et al. (2014). Full step protocol is described in Methods S1. The amplification
process was monitored using the ERCC RNA Spike-in kit (Thermo Scientific, Waltham,
MA, USA) according to manufacturer's instructions. The primers used for cDNA synthesis
and amplification are listed in the Table S2.

Functional annotation and enrichment analyses
The assembled transcriptome was functionally annotated with BLAST2GO (Götz et al.,
2008) using DIAMOND software with *blastx* option (Buchfink et al., 2015) against the
NCBI's plants-*nr* database (NCBI Resource Coordinators, 2016). Blast results were
considered valid with e-value < 1.0E-6. Singular enrichment analysis (SEA) of the GO
terms was made in the AGRIGO v2.0 web tool under standard parameters using as GO
term reference the whole assembled transcriptome annotation (Tian et al., 2017).
Representative enriched GO was determined using REVIGO with 0.5 as dispensability
cutoff value (Supek et al., 2011).

RT-qPCR
The cDNA synthesis was performed using 1 µg of total RNA and iScript™ cDNA
Synthesis Kit (Bio-Rad, Hercules, CA, USA) following manufacturer's instructions. The
qPCR primers were designed following the MIQE guidelines (Bustin et al., 2009). The
primers are listed in the Table S2. qPCRs were carried out using 10 ng of cDNA and 0.4
mM of primers and 2X SsoFast™ EvaGreen® Supermix (Bio-Rad, Hercules, CA, USA) in
a total volume of 10 µL. Relative quantification of gene expression was performed using
thermocycler CFX 384™ Real-Time System, Bio-Rad, Hercules, CA, USA). The qPCR
program was as follows: 3 min at 95 °C (1 cycle), 1 s at 95 °C and 5 s at 60 °C (50 cycles)
and a melting curve from 60 to 95 °C, to generate the dissociation curve to confirm the
specific amplification of each individual reaction. The analyses were carried out as
described by Cañas et al. (2014) using the MAK3 model in the R package *qpcR* (Ritz &
Spiess, 2008). Normalization for gene expression of experimental design viability was
performed using geometric mean of two reference genes, a *Saposin-like aspartyl protease*
(pp_199988) and *Myosin heavy chain-related* (pp_58489) that were previously tested for
maritime pine (Granados et al., 2016). Normalization for gene expression of LCM isolated

tissues was performed using as reference gene a *SKP1/ASK1 family protein* (pp_18128) that

was previously tested for maritime pine LCM samples (Granados et al., 2016). For the RT-

qPCR analysis, three technical replicates of each sample and three biological replicates

were made.

Imaging of phytohormones in roots sections

The main phytohormones were localized in root apex cuts using nano-particle assisted laser

desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). The elongation

zones, which corresponded to the region from 2.5 - 5 mm of the root apex, were embedded

in super cryoembedding medium (SCEM, Leica Biosystems, Wetzlar, Germany) and frozen

in liquid nitrogen. The specimen block was cut into 10 µm sections using a cryostat (NX-

70, Thermo Scientific, Waltham, MA, USA) set at -23°C in the chamber and at -25°C on

the object holder. The sections were gently mounted on slides coated with indium tin oxide

(ITO) (Bruker Daltonik GmbH, Billerica, MA, USA). Optical images of the sections were

obtained by a virtual slide scanner (Nanozmmer-SQ, Hamamatsu Photonics, Shizuoka,

Japan) before analysis by Nano-PALDI-MSI.

For Nano-PALDI-MSI, iron oxide-based nanoparticles (Fe-NPs) were prepared by stirring

aqueous solutions of $FeCl_2 \cdot 4H_2O$ (5 mL, 100 mM; FUJIFILM Wako Pure Chemical), and

3-aminopropyltriethoxysilane (5 mL; γ-APTES; Shin-Etsu Chemical, Tokyo, Japan) at

room temperature for 1 h. The resulting precipitate was washed several times with ultrapure

water, resuspended in methanol (Moritake et al., 2007).

One milligram of Fe-NPs was resuspended in 1 mL methanol and sprayed on pine root

tissue sections on ITO-coated glass slides with an airbrush (nozzle caliber, 0.2 mm). To

obtain images, each data point on the section were irradiated with 200 laser shots in the

positive ion detection mode of the mass spectrometer. Only signals between 80 and 800 *m/z*

were analyzed to detect the corelated IAA (*m/z* 176.3), cytokinin (*m/z* 221.3), ACC (*m/z*

102.8), salicylic acid (*m/z* 139.4), JA (*m/z* 211.0) and ABA (*m/z* 265.0) as protonated ions,

respectively. For each section, approximately 101,800 data points were obtained, 5 µm

apart. The MS image was reconstructed from the obtained MS spectra with a mass bin

width of m/z ± 0.1 from the exact mass using flexImaging 4.0 (Bruker Daltonik GmbH).

The accurate mass of the ions was used for image generation, and mass accuracy and root-

mean- square error (RMSE) were automatically calculated by the imaging software to avoid

false-positive signals (Shiono and Taira, 2020). Comparisons of MS images were derived
from the relative intensity for each signal normalized by the highest intensity spot on the
slide. The peak intensity value of the spectra was normalized by dividing them with the
total ion current (TIC) to achieve semi-quantitative analysis between control- and NH_4^+-
treated roots.
326
Phylogenetic analyses
Evolutionary analyses were performed in MEGA7 (Kumar et al., 2016). The protein
sequence alignment was made with Muscle (Edgar, 2004). The evolutionary history was
inferred using the Neighbor-Joining method (Saitou & Nei, 1987). The bootstrap consensus
tree inferred from 1000 replicates is taken to represent the evolutionary history of the taxa
analyzed (Felsenstein, 1985).

RESULTS

Tissue-specific transcriptomic response to NH$_4^+$ supply

In this study, tissue-specific transcriptome changes triggered by NH$_4^+$ nutrition were analyzed in the maritime pine root apex by a combination of laser capture microdissection and high-throughput RNA sequencing. Prior proceeding with tissue isolation by LCM, each experimental replicate was validated through the expression analysis of transcripts that were expected to be upregulated by NH$_4^+$ supply at 24 h post-irrigation (Figure 1) (Canales et al., 2011; Ortigosa et al., 2021). As expected, NH$_4^+$ induced the accumulation of transcripts coding for glutamine synthetase 1b (*PpGS1b*), antimicrobial peptide 1 (*PpAMP1*) and S-adenosyl methionine synthase (*PpSAMS*). Although not significant, the NADH glutamate synthase (*PpNADH-GOGAT*) expression also increased with NH$_4^+$ supply.

As described in the Materials and methods section, root apexes from seedlings that were irrigated with 3 mM NH$_4^+$ and harvested at 24 h were used for the isolation of four different tissues by LCM, namely, RC, RM, RDC and RDV (Video S1). The differential expression results are shown in Dataset S1. A total of 295 DE transcripts were identified in the low-input RNA-seq analysis (Figure 2A,B), of which 182 DE transcripts were downregulated (Figure 2A) and 113 were upregulated (Figure 2B). Among the isolated tissues, RDC showed the highest number of DE transcripts (107 upregulated and 70 downregulated) while RDV had the lowest response when seedlings were treated with NH$_4^+$ (12 upregulated transcripts). Interestingly, only 2 genes were upregulated in all root tissues, namely, two splicing isoforms from a common gene with unknown function.

The analysis of the DE transcripts revealed 8 expression patterns (Figure 2C). Among the groups with the most significant differences, the cluster 2 had transcripts repressed in RC by NH$_4^+$, and it contained several TFs, such as *PpSHR*, *PpDOF11* and *PpGATA20-like,* and transcripts related to phytohormones, such as *PpVAN3/FKD2* (Figure 2C; Dataset S2). In cluster 3, transcripts related to root development, such as *PpBAM1*, and phytohormones such as *flavanone 3-dioxygenase*, were found to be repressed by NH$_4^+$ in RC and RDC (Figure 2C; Dataset S2). In RDC and RM, NH$_4^+$ induced the transcripts framed into cluster 4, which are mainly related to defense, such as chitinases of class I and IV (Figure 2C; Dataset S2).

To validate our transcriptomic analysis, the expression value of several DE transcripts was

corroborated by RT-qPCR (Figure 2D; Dataset S1). The downregulation of *PpSHR*

(pp_346622) and *PpNPF3.1* (pp_58258) in RC; *PpDOF12* (pp_85381) and *PpACO*

(pp_202153) in RM; and *PpBLISTER-like* (pp_170664) in RDC was confirmed. The

upregulation of transcripts coding a *cycloeucalenol cycloisomerase* (*PpCPI1*; pp_189071)

in RC and *PpAMP1* (pp_580007) in RDC was also corroborated.

Functional enrichment analysis of tissue-specific transcriptomic response to NH$_4^+$

supply

The functional study of DE transcripts, including the SEA results, is summarized in Figure

3. In the most responsive tissues (RC, RM and RDC), NH$_4^+$ induced a general alteration in

the amounts of transcripts coding TFs involved in development (most of them

downregulated) and transcripts related to different phytohormones (Figure 3A-C). In the

less responsive tissue (RDV) but also in RM and RDC, NH$_4^+$ promoted the accumulation of

defense-related transcripts (Figure 3B-D). In the SEA results, no significant GO terms were

found for DE transcripts upregulated in RC and RDV and downregulated in RM and RDV.

The complete results of the SEA analyses are shown in Dataset S3. The largest number of

SEA results were for transcripts downregulated in RC and upregulated in RDC. The

downregulated transcripts in RC were significantly enriched in GO terms in the biological

process (BP) category, such as "biological regulation" (GO:0065007), "gene expression"

(GO:0010467), "response to hormone" (GO:0009725), "regulation of hormone levels"

(GO:0010817) and "regulation of transcription, DNA-templated" (GO:0006355) (Figure

3E). The terms enriched in the molecular function (MF) category included "transcription

factor activity" (GO:0003700), "protein binding" and "DNA binding" (GO:0005515 and

GO:0003677) (Figure 3F). In the BP category for the downregulated transcripts in RDC

only the "cytoskeleton organization" term (GO:0007010) was significantly enriched;

however, for the upregulated transcripts in this tissue, numerous enriched functions were

significant, such as those involved in defense responses such as "induction of programmed

cell death" (GO:0012502), "cell killing" (GO:0001906), "defense response" (GO:0006952)

and "cell wall macromolecule catabolic process" (GO:0016998) and "amino sugar catabolic

process" (GO:0046348) (Figure 3E). Regarding the MF category for the upregulated

transcripts in RDC, there were significant GO terms related to the response to biotic stress,

such as "chitinase activity" (GO:0004568), "chitin binding" (GO:0008061), "mannose
binding" (GO:0005537) and "hydrolase activity, acting on glycosyl bonds" (GO:0016798)
(Figure 3F). Interestingly, the SEA results of the upregulated transcripts in RM were
similar but limited for defense responses, which was also found for the upregulated
transcripts in RDC (Figure 3E,F). Finally, the only enriched GO terms for the cellular
component category were "extracellular region" (GO:0005576) and "extracellular space"
(GO:0005615) for the transcripts upregulated in RDC tissue (Dataset S3).

RNA-seq analysis in function of the root tissue
Although it has been studied more in deep in a previously work (Cañas et al., 2017), the
transcriptional changes underlying different tissues that form the root apex of maritime pine
(RC, RM, RDC and RDV) have been briefly analyzed (Dataset S1). The comparison of the
DE transcripts showed a set of characteristic genes of each tissue. This gene core was
composed by 1014, 2, 529 and 2692 DE transcripts for RC, RM, RDC and RDV,
respectively (Figure S3A). The functional study of the core DE transcripts revealed the
main functions for these tissues (Dataset S3). Some of the enriched GO terms for RC are
"aminoglycan metabolic process" (GO:0006022), "cell wall organization or biogenesis"
(GO:0071554), "dehiscence" (GO:0009900) or "polysaccharide catabolic process"
(GO:0000272) (Figure S3B). For RDC gene core, GO terms such as "anatomical structure
formation involved in morphogenesis" (GO:0048646), "catabolic process" (GO:0009056)
or "response to stimulus" (GO:0050896) were observed (Figure S3B). On the other hand,
RDV was the tissue that showed the highest number of DE transcripts when compared to
the rest of the tissues, which presented enrichment GO terms such as "auxin influx"
(GO:0060919), "carbohydrate metabolic process" (GO:0005975), "fluid transport"
(GO:0042044) or "phloem or xylem histogenesis" (GO:0010087), among others (Figure
S3B). Interestingly, in RM only 2 DE transcripts were identified as characteristic and both
transcripts are splicing variants of the same gene encoding a bHLH TF, similar to the
Arabidopsis FAMA TF (AT3G24140) (Figure S3C).

Transcription factors affected by NH_4^+ in the apex of maritime pine roots
The DE transcripts were individually analyzed looking to identify different kinds of
regulators, such as TFs and transcripts involved in the phytohormone response. From a total

of 295 DE transcripts, 31 TFs were identified (Figure 4, Table 1, Dataset S1). All 10 FTs

identified in RC were repressed in the presence of NH_4^+. However, in RM and RDC, 6 and

4 TF transcripts were downregulated while 5 and 6 TF transcripts were upregulated,

respectively. Interestingly, a high abundance of TFs related to root growth and development

was observed (Table 1), such as the strong downregulation of *PpSHR* in RC, *INCURVATA-*

like and homeobox-leucine zipper protein *HAT-like* both in RM and involved in meristem

developmental regulation, and the TF identified as *zinc finger C2H2 SHOOT*

GRAVITROPISM in RC related to the gravitropism response. *PpSHR* was the most highly

repressed TF (-11 logFC). Furthermore, the induction of two transcriptional repressors of

different plant developmental processes was observed, with *PpNAC31* identified as an

SMB-like NAC TF in RM and an *OFP17*-like repressor in RDC. In addition, novel TFs in

maritime pine were identified. Two of them belong to the DOF-family and were named as

PpDOF11 and *PpDOF12*, and they were both downregulated in RC and RM, respectively.

Phylogenetic analysis of these two new DOF-type TFs revealed that *PpDOF11* is grouped

with *AtDOF1.4* (AT1G28310) and *OsDOF7.2* (LOC_Os07g32510) into subfamily E and

that *PpDOF12* is grouped with members of subfamily A of *P. pinaster*: *PpDOF4*,

PpDOF7, *PpDOF8* and *PpDOF10* (Figure S4, Table S3). Additionally, a member of the

NAC family (*PpNAC38)* that was upregulated in RM tissue was identified. This TF has

homology to *BRN1* (AT1G33280) (e-value: 7e-96) and *BRN2* (AT4G10350) (e-value: 3e-

97) of *Arabidopsis thaliana* and is framed within the NAC subfamily C (Figure S5, Table

S4).

A gene co-expression network was constructed to identified hub TFs and putatively

determine the processes controlled by them. Thirteen transcripts were identified as hubs

corresponding to 8 different TFs: *PpDOF1*, *PpDOF2*, *PpDOF11*, *PpGATA20-like*,

PpINCURVATA-like, *PpMyb149*, *PpSGR/IDD16* and *PpSHR*. All of them were DE for

NH_4^+ treatment in different tissues except RDV but all of them were more expressed in

RDV than in the rest of tissues except *PpGATA20-like* (Dataset S1). The functions and

expression localization of the most correlated transcripts (>|0,9|) were analyzed (Figure S6;

Dataset S4). Most of the transcripts were correlated with *DOF* TFs, *PpINCURVATA-like*

and *PpSGR/IDD16* but also belonged to the RDV gene core (1879 from 2362) (Figure

S6A). Thus, the enriched functions were mainly related with RDV functions such as "auxin

influx" (GO:0060919), "carbohydrate metabolic process" (GO:0005975), "fluid transport"

(GO:0042044) or "phloem or xylem histogenesis" (GO:0010087) (Figure S6B; Dataset S5).

Interestingly, transcripts correlated with *PpGATA20-like* were enriched in functions such as

"gene expression" (GO:0010467) and "ribosome biogenesis" (GO:0042254) (Figure S6B;

Dataset S5).

Phytohormone-related genes

Twenty-six DE transcripts related to phytohormone pathways were identified (Table 1), of

which 19 were upregulated and 7 were downregulated (Figure 4, Dataset S1). Most of these

DE transcripts were related to IAA (Table 1). Some interesting examples are the

downregulation of transcripts encoding proteins required for PIN transporters localization,

such as VAN3/FKD2 Auxin canalis/PH2 (pp_60103) in RC and the upregulation of a

PpCPI1 (pp_189071) in RC, which is related to PIN transporters endocytosis. In addition,

in RDC, the repression of transcripts coding for the IAA receptor TIR1/AFB (pp_238715)

and for a sulfotransferase (pp_208269) with a high similarity degree to SULT202B1

(AT3G45070) (e-value: 5e-72) was observed.

CK-related genes were the second most represented phytohormone-related transcripts, and

strong repression of CK riboside 5'-monophosphate phosphoribohydrolase (pp_222714) (-

6.9 logFC), a CK-activating enzyme, was observed in RC. However, upregulation of a

transcript coding for a CK dehydrogenase (pp_215376), a CK-inactivating enzyme, was

also observed in RDC. The repression of transcripts related to brassinosteroids and ethylene

(ET) biosynthesis, e.g., cytochrome P450 90B1 (pp_247190) in RDC and *PpACO*

(pp_202153) in RM was also observed. In addition, transcripts related to the gibberelin

(GA) response and abscisic acid (ABA) biosynthesis were upregulated, such as a GASA

gibberellin-regulated cysteine-rich protein (pp_111379) in RDC and two ABA2 coding

transcripts that might be involved in ABA biosynthesis (pp_198945, and pp_209084) in

RM and RDC.

Phytohormone detection in maritime pine root apex

To corroborate whether NH_4^+ nutrition altered the spatial allocation of several

phytohormones as the transcriptomic data suggest, the distribution of multiple

phytohormones was determined by Nano-PALDI-MSI in the apex of maritime pine roots

(Shiono & Taira, 2020). In comparison to control seedlings, changes in the patterns of

IAAs, CK (tZ, *trans*-Zeatin), ACC, ABA, salicylic acid (SA) and jasmonic acid (JA) were
observed in the presence of NH_4^+ (Figures 5 and S5). In the control plants, an IAA gradient
was observed, the maximum level was detected in the most distal area of the principal root
apex (Figure 5A); however, in the NH_4^+-treated roots, the IAA maximum was detected in a
more distant zone from the root tip (Figure 5A). CK (tZ) showed a wide distribution in the
roots of control plants (Figure 5B). In the NH_4^+-treated seedlings, CK was distributed in the
outer tissues of the root mainly below the root tip (Figure 5B). Regarding ET, its precursor
ACC was found in a very restricted area of the root tip in control and NH_4^+-treated plants,
showing no changes when plants were supplied with NH_4^+ (Figure 5C). ABA was detected
mainly in the RDC tissue in both control and treated seedlings, and it was higher under
NH_4^+ supply and distributed below the root apex (Figure 5D). SA and JA phytohormone
distributions were also affected by the presence of NH_4^+ (Figure S7). SA was located at the
end of the root tip, while JA was also detected in the elongation zone in control plants.
When the seedlings were supplied with NH_4^+, SA showed a vaguer spatial distribution with
respect to the observed distribution in the control plants (Figure S7A) while JA tended to
slightly increase in the outermost parts of the root apex ends and disappeared from the
elongation zone (Figure S7B).
509

Maritime pine root phenotype
Maritime pine root morphological studies showed no differences between treatments
regarding whole root biomass accumulation (Figure 6A). When the weight of the principal
roots (PRs) was measured, NH_4^+-fed seedlings exhibited a significant reduction in PR
weight compared to water- and NO_3^--treated plants (Figure 6B) and a statistically higher
root length when pine seedlings were supplied with either inorganic nitrogen form (Figure
6C). Moreover, NH_4^+ promoted an increase in LR number and weight compared to the
water and NO_3^- treatments (Figure 6D,E), although no statistically significant differences
were observed regarding LR density (LRD) between treatments (Figure 6F).

DISCUSSION

Despite the obvious environmental and economic importance of conifers, little is known about their response to diverse nutrients at the molecular and developmental levels. Since they represent an ancient lineage of gymnosperms, conifers are of particular interest from an evolutionary perspective because they have common but also differential responses with angiosperm models. The aim of the present work was to explore the early response of maritime pine to NH_4^+ nutrition in the different tissues of the root apex. For this purpose, tissue isolation was performed by LCM and combined with low-input RNA-seq to determine the local transcriptomic response in the growth area of the root. This strategy avoids the dilution effect of transcripts with low and much localized expression, revealing new genes not previously annotated (Cañas et al., 2017), such as *PpDOF11-12* and *PpNAC38*. Clearly a dilution effect can be observed when compared the expression of marker genes in the whole roots (Figure 1) and in the specific tissue transcriptomics (Figure S8A). Supporting this, from 47,118 transcripts used in the tissue specific RNA-seq 8,491 transcripts were not previously identified in whole root transcriptomic experiments (Dataset S1). A similar number of DE transcripts was obtained in the present work (295) when compared to the same NH_4^+ nutrition treatment over the whole root (350) (Ortigosa et al., 2021) (Figure S8B). However, only 3 transcripts with the same identifiers are shared, although there are different transcripts from the same gene that are DE in both analyses, for example *PpAMP1* (RDC - pp_58007 / whole root - pp_58005) (Figure S8C).

The response to NH_4^+ in RDC involved the expression of defense-related genes as previously described in maritime pine and other model plants (Figure 3C,E,F) (Canales et al., 2010; Patterson et al., 2010; Ravazzolo et al., 2020; Ortigosa et al., 2021). Obviously, this response might be linked to the peripheral localization of RDC in the roots since the defense proteins encoded by the expressed transcripts develop their functions in the extracellular media (Dataset S3). Some examples are the upregulation of different *ginkbilobin-like/embryo abundant protein* transcripts (e.g. pp_252647), *chitinases* (e.g. pp_117806) and the *PpAMP1* (pp58007), which are consistent with results previously reported in maritime pine roots (Canales et al., 2010; Ortigosa et al., 2021).

In angiosperm model plants, NH_4^+ alters RSA by inhibiting root elongation, stimulating LR branching and affecting root hair development (Y. Liu & von Wirén, 2017). RSA principally targets root system processes, such as root elongation, root gravitropism and LR

branching, appearing to occur in the root tip (Li et al., 2014). Based on the presented

transcriptomic analyses, early root exposure to NH_4^+ caused a wide impact on the

expression of TFs related to root growth and development in pine (Figure 2, Table 1). Thus,

NH_4^+ caused a strong repression of *PpSHR* (pp_246622) in RC tissue, a GRAS-type TF

involved in the regulation and coordination of root development, including phloem

differentiation (Kim et al., 2020). Interestingly, in RDC tissue, it was observed that

PpBAM-like (pp_19744) transcripts were also severely affected by NH_4^+ (Dataset S1),

which is consistent with previous cell-specific transcriptomic profiles observed in

Arabidopsis (Brady et al., 2007). BAM1/2 kinase receptors are required for SHR-dependent

formative divisions in roots and for the proper CYCD6;1 expression, which is required to

promote division in cortex endodermal initial daughter cells (Crook et al., 2020).

Additionally, a progressive reduction in the abundance of IAA efflux carriers (PINs) in

Arabidopsis shr mutants has been described (Lucas et al., 2011), and NH_4^+ negatively

affects the expression of PIN2 and AUX1 coding transcripts (Y. Liu et al., 2013). The

expression of genes coding for PIN transporters was not affected in pine roots. Recently, it

has been linked IAA signaling pathway to acidification (Jia et al., 2020; Meier et al., 2020).

However, based on our results, no significant changes were observed in the expression of

pH-related genes. However, it was observed that the expression of coding transcripts for

proteins involved in IAA transporter localization and components of the IAA signaling

pathway were affected by NH_4^+. This is the case for *PpCPI1* (pp_189071), which is

upregulated in RC and required for PIN2 endocytosis (Men et al., 2008), and the strong

repression of *VAN3/PH2/FKD2* (pp_60103) in RC and RM tissues (Table 2), which is

crucial in *Arabidopsis* for the vascular leaf pattern formation by the proper PIN transporter

localization (Hou et al., 2010). These results suggest that NH_4^+ triggers an alteration in

polar auxin transport (PAT) involving a highly coordinated transcriptomic response

between tissues of the pine root. The Nano-PALDI-MSI data confirmed that under a supply

of NH_4^+, PAT was impaired, thus promoting a putative local increase in IAA presumably in

RC and/or RM tissues where cambial development takes place based on IAA-related

transcripts and TF expression (Figures 5A and 7), which could suggest a different

regulatory mechanism of those pH-dependent described in rice and *Arabidopsis* (Jia et al.,

2020; Meier et al., 2020). One possible alternative for this PAT alteration could be IAA

conjugation with sugars/amino acids as previously described (Tamura et al., 2010; Di et al.,

2021). However, our data revealed non-significant expression changes of IAA conjugating
genes (Dataset S1). This suggests that the NH_4^+ concentration used in this work is not
excessive for maritime pine compared with those applied before (>5 mM NH_4^+) in rice and
Arabidopsis (Tamura et al., 2010; Di et al., 2021), which is supported by the non-inhibition
of PR growth under NH_4^+ supply (Figure 6C). Therefore, other mechanisms could be acting
during the observed PAT process in the root apex of maritime pine.

In addition, our results differ from those previously described in *Arabidopsis* where the
presence of NH_4^+ does not seem to affect the distribution of IAAs in the root tip (Y. Liu et
al., 2013). This finding could be related to root vascular development and root cell
differentiation since adjustments to IAA flux are involved in root cambium development
(Marhava et al., 2018), and SCR, a partner of SHR, is involved in the expression and polar
localization of PINs (Xu et al., 2006).

Following this line of argument, it is interesting to mention that four maritime pine DOFs
were significantly downregulated: *PpDOF11* (pp_58878) and *PpDOF12* (pp_85381) in RC
and RM tissues respectively, and *PpDOF1* (pp_233816) and *PpDOF2* (pp_119525) in
RDC (Figure 2), which were preferentially expressed in tissues close to vascular vessels
(Figures S9-S12). Therefore, it is tempting to suggest that these DOFs may be involved in
the development of root vessels, as described for PHLOEM EARLY DOF (PEAR)
proteins, which are regulated by CK levels (Miyashima et al., 2019). This is also supported
by the network analysis and the functions of the most correlated transcripts to these DOFs
(Figure S6B). Interestingly, the CK distribution under NH_4^+ supply (Figure 5B) correlates
with the repression of *CK riboside 5'-monophosphate phosphoribohydrolase* (pp_222714),
which is a CK activator, in RC (Kuroha et al., 2009) and the upregulation of a *CK*
dehydrogenase (pp_215376), which is involved in CK degradation (Werner et al., 2003), in
RDC (Figure 3A,C). All these results suggest that NH_4^+ could promote an alteration of CK
spatial distribution in maritime pine roots, which could be the reason for the wide
repression of DOF TFs in RC and RM tissues and its possible role as regulators of
cambium development (Figure 7). Phytohormone imaging showed that CK and IAA were
distributed in different zones under a supply of NH_4^+ (Figure 5A,B), which was possibly
due to the opposite roles of these phytohormones in the root meristem since CK negatively
regulates QC specification and functions by modulating the IAA response (Zhang et al.,
2013).

Additionally, homeobox-leucine zipper (HD-Zip) and NAC-domain TFs are described to

take part during development processes (Ochando et al., 2006; Bennett et al., 2010). In

Arabidopsis, several HATs play important roles in development including the regulation of

meristem and procambial maintenance and/or formation (Prigge et al., 2005; Roodbarkelari

& Groot, 2017). Accordingly, two HD-Zip coding transcripts identified as *HAT-like*

(pp_141831) and *INCURVATA-like* (pp_85012) were downregulated in RM tissue (Table

1). These findings are consistent with the repression of TFs related to the root vascular

development process and suggest that NH_4^+ could affect PR vascular development during

early exposure (Figure 7). Nevertheless, long-term NH_4^+ nutrition promotes PR growth but

causes a decrease in PR weight (Figure 6B,C), suggesting that root radial growth could be

affected by this transcriptional network. In future studies, cell division and elongation

should be analyzed in the root these zones. Unfortunately, the exact anatomy of root

meristem with the quiescent center position has not yet been defined in conifers.

Interestingly, in RM, the upregulation of two NAC TFs, *PpNAC31* and *PpNAC38,* similar

to the *Arabidopsis SMB* and *BRN* genes (Table 1, Figure S4), is consistent with the

expression patterns of *SMB*, *BRN1* and *BRN2*, which largely overlap in *Arabidopsis* roots

(Bennett et al., 2010). SMB represses stem cell-like divisions in the root cap daughter cells

(Bennet et al., 2014; Kamiya et al., 2016), and together with BRN1 and BRN2, it regulates

cellular maturation of the root cap in *Arabidopsis* (Bennett et al., 2010). These findings are

consistent with the reduction in the number of root cap cells under NH_4^+, as described by Y.

Liu et al., (2013), thus providing a possible explanation for this observation in *Arabidopsis*.

In rice, the endogenous ABA content alleviates NH_4^+ rice toxicity, promoting NH_4^+

assimilation by stimulating the GS/GOGAT cycle and enhancing antioxidant activities

(superoxide dismutase, ascorbate peroxidase and catalase) (Sun et al., 2020). In the

maritime pine root apex under NH_4^+ supply, ABA levels increased (Figure 5D), and

transcripts encoding ABA2 enzymes (e.g. pp_198956) involved in ABA biosynthesis were

upregulated in RM and RDC, and several putative ABA-induced TFs (e.g. pp_233345)

were upregulated in RC and RM (Dataset S1). However, the GS/GOGAT cycle was

apparently not induced, although it was observed in the whole root response to NH_4^+ supply

(Ortigosa et al., 2021) (Figure 1). In general, there was no expression changes in genes

involved in N uptake and assimilation at the root apex of pine in response to NH_4^+ supply.

In angiosperms, the upregulation of GS/GOGAT cycle has been described in specific

tissues such as root epidermis around root hair zone in rice (Ishiyama et al., 1998; Ishiyama

et al., 2004a) and *Arabidopsis* (Ishiyama et al., 2004b; Konishi et al. 2017). This raises the

question of whether conifers might have a different response to NH_4^+ availability at the root

apex than angiosperms.

Regarding antioxidant metabolism, different transcripts involved in ascorbate production

were downregulated (Dataset S1), including *PpBLISTER-like* TF (pp_170664) in RDC

tissue (Purdy et al., 2011) and two *L- gulonolactone oxidases* (pp_183087 and pp_234650)

in RC and RDC tissues (Aboobucker et al., 2017). This finding suggests that the

glutathione-ascorbate cycle is not operative in maritime pine roots exposed to NH_4^+, which

is consistent with the levels of oxidized glutathione observed under long-term experiments

(Ortigosa et al., 2020).

Finally, ET is involved in NH_4^+-responsive pathways in rice (Sun et al., 2017) and in

cambial development in poplar trees (Love et al., 2009). In *Arabidopsis,* the ET levels

increased under a supply of NH_4^+, which reduced lateral root formation because of the

repression of auxin transporter 1 (AUX1) (Li et al., 2014). In contrast, the results in

maritime pine suggest a reduction in ET biosynthesis following NH_4^+ application. A

transcript encoding an ACO was repressed in RM, which is consistent with proteomic and

epitranscriptomic results in maritime pine whole roots (Ortigosa et al., 2021). This finding

is consistent with the downregulation of *BLISTER-like* TF and *L- gulonolactone oxidases*

since ascorbate is one of the ACO enzyme substrates (Dilley et al., 2013). However, ACC

was not affected by NH_4^+ treatment (Figure 5C). Accordingly, NH_4^+ stimulated the LR

components (number and weight) and PR length (Figure 5) in maritime pine, which is

consistent with the increased number of root apexes under NH_4^+ nutrition compared to

NO_3^- nutrition in *Pinus massoniana* Lamb. (Ren et al., 2020). Taken together, the results

suggest that maritime pine reduces ET biosynthesis under NH_4^+ supply, which could be

linked to the PAT and CK alteration observed (Figure 4A,B) as well as to the expression

patterns observed for TFs related to cambium development (Table 1) due to the importance

of IAA-ET-CK crosstalk during root development (J. Liu et al., 2017).

CONCLUSIONS

NH_4^+ nutrition is a complex process in plants since NH_4^+ is a nutrient that can be toxic

when supplied in excess. However, conifers in general are quite tolerant to NH_4^+.

Therefore, the identifying of the differential mechanisms underlying NH_4^+ tolerance in
conifers would be enormously helpful for improving NH_4^+ tolerance not only in these trees
but also in other crops. In contrast to other plant models, supplying NH_4^+ to pine represses
ET and ascorbate production and alters IAA and CK patterns, thus regulating a
transcriptional network in the short term. Consequently, the growth and development of PR
and LRs are stimulated in pine.

The expression profiles of genes related to root cambium development and phytohormones
suggest a molecular mechanism underlying changes in the RSA phenotype that includes
IAA-CK-ET crosstalk and a transcriptional network at least during the early root response
to NH_4^+ supply. The results reported in the present study tentatively link *SHR*, and other
TFs to NH_4^+ nutrition and its phenotypic effect on RSA, which likely affect early vascular
development (Figure 7).

This work provides new and valuable data to unravel the mechanisms involved in the
response of maritime pine to NH_4^+ nutrition. However, further research efforts are required
to reach a full understanding of the molecular basis of NH_4^+ tolerance.

ACKNOWLEDGEMENTS
Acknowledgments: This research was funded by Spanish *Ministerio de Ciencia e*
Innovación, grant numbers BIO2015-73512-JIN MINECO/AEI/FEDER, UE, RTI2018-
094041-B-I00 and EQC2018-004346-P. FO was supported by grants from the *Universidad*
de Málaga (*Programa Operativo de Empleo Juvenil vía SNJG, UMAJI11, FEDER, FSE,*
Junta de Andalucía) and BIO-114*, Junta de Andalucía*.

Author Contributions
FO, HS and ST have performed the experiments; CLF has performed transcriptomic data
analysis; FO and RAC have wrote the manuscript; CA and FMC made additional
contributions and edited the manuscript; CA, FMC and RAC have acquired the funds; FO
and RAC have planned and designed the research.

Data Accessibility Statement

The data that support the findings of this study are available from different databases,

supporting information and from the corresponding author upon reasonable request. RNA-

Seq data have been deposited in the NCBI's Gene Expression Omnibus and are accessible

through GEO Series with the accession number GSE175587

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE175587). The transcriptome

shotgun assembly project has been deposited at DDBJ/EMBL/GenBank under the

accession GJFX00000000. The version described in this paper is the first version,

GJFX01000000.

Conflict of Interest

The authors have no conflict of interest.

REFERENCES

Aboobucker, S. I., Suza, W. P., & Lorence, A. (2017). Characterization of two *Arabidopsis*

L-gulono-1, 4-lactone oxidases, AtGulLO3 and AtGulLO5, involved in ascorbate

biosynthesis. Reactive Oxygen Species (Apex, NC), *4*(12), 389.

doi.org/10.20455/ros.2017.861.

Antolín-Llovera, M., Leivar, P., Arró, M., Ferrer, A., Boronat, A., & Campos, N. (2011).

Modulation of plant HMG-CoA reductase by protein phosphatase 2A: positive and negative

control at a key node of metabolism. Plant Signaling & Behavior, *6*(8), 1127-1131. doi:

10.4161/psb.6.8.16363.

Araya, T., Kubo, T., von Wirén, N., & Takahashi, H. (2016). Statistical modeling of

nitrogen‐dependent modulation of root system architecture in *Arabidopsis thaliana*. Journal

of Integrative Plant Biology, *58*(3), 254-265. doi.org/10.1111/jipb.12433.

Bak, S., Beisson, F., Bishop, G., Hamberger, B., Höfer, R., Paquette, S., & Werck-

Reichhart, D. (2011). Cytochromes P450. The Arabidopsis Book/American Society of

Plant Biologists, *9*. doi: 10.1199/tab.0144.

Bennett, T., van den Toorn, A., Sanchez-Perez, G. F., Campilho, A., Willemsen, V., Snel,

B., & Scheres, B. (2010). SOMBRERO, BEARSKIN1, and BEARSKIN2 regulate root cap

maturation in *Arabidopsis*. The Plant Cell, *22*(3), 640-654.

doi.org/10.1105/tpc.109.072272.

Bennett, T., van den Toorn, A., Willemsen, V., & Scheres, B. (2014). Precise control of

plant stem cell activity through parallel regulatory inputs. Development, *141*(21), 4055-

4064. doi: 10.1242/dev.110148.

Bijlsma, R. J., Lambers, H., & Kooijman, S. A. L. M. (2000). A dynamic whole-plant

model of integrated metabolism of nitrogen and carbon. 1. Comparative ecological

implications of ammonium-nitrate interactions. Plant and Soil, *220*(1), 49-69.

Brady, S. M., Orlando, D. A., Lee, J. Y., Wang, J. Y., Koch, J., Dinneny, J. R., ... &

Benfey, P. N. (2007). A high-resolution root spatiotemporal map reveals dominant

expression patterns. Science, *318*(5851), 801-806. doi.org/10.1126/science.1146265.

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using

DIAMOND. Nature Methods, *12*(1), 59-60. doi.org/10.1038/nmeth.3176.

Bustin, S. A., Benes, V., Garson, J. A., Hellemans, J., Huggett, J., Kubista, M., ... &

Wittwer, C. T. (2009). The MIQE guidelines: minimum information for publication of

quantitative Real-Time PCR experiments. Clinical Chemistry, *55*(4):611-22.

doi.org/10.1373/clinchem.2008.112797.

Canales, J., Flores-Monterrosso, A., Rueda-López, M., Avila, C., & Cánovas, F. M. (2010).

Identification of genes regulated by ammonium availability in the roots of maritime pine

trees. Amino Acids, *39*(4), 991-1001. doi.org/10.1007/s00726-010-0483-9.

Canales, J., Avila, C., & Cánovas, F. M. (2011). A maritime pine antimicrobial peptide

involved in ammonium nutrition. Plant, Cell & Environment, *34*(9), 1443-1453.

doi.org/10.1111/j.1365-3040.2011.02343.x.

Canales, J., Rueda-López, M., Craven-Bartle, B., Avila, C., & Cánovas, F. M. (2012).

Novel insights into regulation of asparagine synthetase in conifers. Frontiers in Plant

Science, *3*, 100. doi.org/10.1007/s00726-010-0483-9.

Cánovas, F. M., Avila, C., Canton, F. R., Canas, R. A., & de la Torre, F. (2007).

Ammonium assimilation and amino acid metabolism in conifers. Journal of Experimental

Botany, *58*(9), 2307-2318. doi.org/10.1093/jxb/erm051.

Canas, R. A., de la Torre, F., Canovas, F. M., & Canton, F. R. (2006). High levels of

asparagine synthetase in hypocotyls of pine seedlings suggest a role of the enzyme in re-

allocation of seed-stored nitrogen. Planta, *224*(1), 83-95. doi.org/10.1007/s00425-005-
0196-6.

Cañas, R. A., Canales, J., Gómez-Maldonado, J., Ávila, C., & Cánovas, F. M. (2014).
Transcriptome analysis in maritime pine using laser capture microdissection and 454
pyrosequencing. Tree Physiology, *34*(11), 1278-1288. doi.org/10.1093/treephys/tpt113.

Canas, R. A., Li, Z., Pascual, M. B., Castro-Rodríguez, V., Avila, C., Sterck, L., ... &
Canovas, F. M. (2017). The gene expression landscape of pine seedling tissues. The Plant
Journal, *91*(6), 1064-1087. doi.org/10.1111/tpj.13617.

Chaban, C., Waller, F., Furuya, M., & Nick, P. (2003). Auxin responsiveness of a novel
cytochrome P450 in rice coleoptiles. Plant Physiology, *133*(4), 2000-2009.doi:
10.1104/pp.103.022202.

Chapman, E. J., & Estelle, M. (2009). Mechanism of auxin-regulated gene expression in
plants. Annual Eeview of Genetics, *43*, 265-285. doi.org/10.1146/annurev-genet-102108-
134148.

Coll-Garcia, D., Mazuch, J., Altmann, T., & Müssig, C. (2004). EXORDIUM regulates
brassinosteroid-responsive genes. FEBS Letters, *563*(1-3), 82-86. doi: 10.1016/S0014-
5793(04)00255-8.

Corrales, A. R., Carrillo, L., Lasierra, P., Nebauer, S. G., Dominguez-Figueroa, J.,
Renau-Morata, B., ... & Medina, J. (2017). Multifaceted role of cycling DOF factor 3
(CDF3) in the regulation of flowering time and abiotic stress responses in *Arabidopsis*.
Plant, Cell & Environment, *40*(5), 748-764. doi: 10.1111/pce.12894.

Crook, A. D., Willoughby, A. C., Hazak, O., Okuda, S., VanDerMolen, K. R., Soyars, C.
L., ... & Nimchuk, Z. L. (2020). BAM1/2 receptor kinase signaling drives CLE peptide-
mediated formative cell divisions in *Arabidopsis* roots. Proceedings of the National
Academy of Sciences, *117*(51), 32750-32756. doi.org/10.1073/pnas.2018565117.

Cui, H., Levesque, M. P., Vernoux, T., Jung, J. W., Paquette, A. J., Gallagher, K. L., ... &

Benfey, P. N. (2007). An evolutionarily conserved mechanism delimiting SHR movement

defines a single layer of endodermis in plants. Science, *316*(5823), 421-425.

doi.org/10.1126/science.1139531.

Cui, D., Zhao, J., Jing, Y., Fan, M., Liu, J., Wang, Z., ... & Hu, Y. (2013). The *Arabidopsis*

IDD14, IDD15, and IDD16 cooperatively regulate lateral organ morphogenesis and

gravitropism by promoting auxin biosynthesis and transport. PLoS Genetics, *9*(9),

e1003759. doi: 10.1371/journal.pgen.1003759.

Di, D.-W., Li, G., Sun, L., Wu, J., Wang, M., Kronzucker, H.J., …& Shi, W. (2021). High

ammonium inhibits root growth in *Arabidopsis thaliana* by promoting auxin conjugation

rather than inhibiting auxin biosynthesis. Journal of Plant Physiology, *261*, 153415. doi:

10.1016/j.jplph.2021.153415.

Dilley, D. R., Wang, Z., Kadirjan-Kalbach, D. K., Ververidis, F., Beaudry, R., &

Padmanabhan, K. (2013). 1-Aminocyclopropane-1-carboxylic acid oxidase reaction

mechanism and putative post-translational activities of the ACCO protein. AoB Plants, *5*.

doi.org/ 10.1093/aobpla/plt031.

Domínguez-Figueroa, J., Carrillo, L., Renau-Morata, B., Yang, L., Molina, R. V., Marino,

D., ... & Medina, J. (2020). The *Arabidopsis* transcription factor CDF3 is involved in

nitrogen responses and improves nitrogen use efficiency in tomato. Frontiers in Plant

Science, *11*, 1825. doi: 10.3389/fpls.2020.601558.

Edgar, R., Domrachev, M., & Lash, A. E. (2002). Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. Nucleic Acids Research, *30*(1), 207-

210. doi.org/10.1093/nar/30.1.207.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Research, *32*(5), 1792-1797. doi.org/10.1093/nar/gkh340.

Esteban, R., Ariz, I., Cruz, C., & Moran, J. F. (2016). Mechanisms of ammonium toxicity
and the quest for tolerance. Plant Science, *248*, 92-101.
doi.org/10.1016/j.plantsci.2016.04.008.

Farjon, A. (2018). The Kew review: conifers of the world. Kew Bulletin, *73*: 8,
doi.org/10.1007/s12225-018-9738-5.

Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap.
Evolution. *39*, 783-791, doi.org/10.1111/j.1558-5646.1985.tb00420.x.

Fendrych, M., Van Hautegem, T., Van Durme, M., Olvera-Carrillo, Y., Huysmans, M.,
Karimi, M., ... & Nowack, M. K. (2014). Programmed cell death controlled by
ANAC033/SOMBRERO determines root cap organ size in *Arabidopsis*. Current Biology,
24(9), 931-940. doi: 10.1016/j.cub.2014.03.025.

Fujita, S., Ohnishi, T., Watanabe, B., Yokota, T., Takatsuto, S., Fujioka, S., ... & Mizutani,
M. (2006). *Arabidopsis* CYP90B1 catalyses the early C-22 hydroxylation of C27, C28 and
C29 sterols. The Plant Journal, *45*(5), 765-774. doi: 10.1111/j.1365-313X.2005.02639.x.

Fukaki, H., Wysocka-Diller, J., Kato, T., Fujisawa, H., Benfey, P. N., & Tasaka, M. (1998).
Genetic evidence that the endodermis is essential for shoot gravitropism in *Arabidopsis*
thaliana. The Plant Journal, *14*(4), 425-430. doi: 10.1046/j.1365-313x.1998.00137.x.

Galinha, C., Hofhuis, H., Luijten, M., Willemsen, V., Blilou, I., Heidstra, R., & Scheres, B.
(2007). PLETHORA proteins as dose-dependent master regulators of *Arabidopsis* root
development. Nature, *449*(7165), 1053-1057. doi.org/10.1038/nature06206.

Gifford, M. L., Dean, A., Gutierrez, R. A., Coruzzi, G. M., & Birnbaum, K. D. (2008).
Cell-specific nitrogen responses mediate developmental plasticity. Proceedings of the
National Academy of Sciences, 105(2), 803-808. doi.org/10.1073/pnas.0709559105.

Götz, S., García-Gómez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., ...
& Conesa, A. (2008). High-throughput functional annotation and data mining with the
Blast2GO suite. Nucleic Acids Research, *36*(10), 3420-3435. doi.org/10.1093/nar/gkn176.

Granados, J. M., Ávila, C., Cánovas, F. M., & Cañas, R. A. (2016). Selection and testing of
reference genes for accurate RT-qPCR in adult needles and seedlings of maritime pine.
Tree Genetics & Genomes, *12*(3), 1-15. doi.org/10.1007/s11295-016-1018-7.

Greb, T. (2020). Plant Development: how phloem patterning occurs. Current Biology, *30*:
217-219, doi.org/10.1016/j.cub.2020.01.021.

Grunewald, W., & Friml, J. (2010). The march of the PINs: developmental plasticity by
dynamic polar targeting in plant cells. The EMBO Journal, *29*(16), 2700-2714.
doi.org/10.1038/emboj.2010.181.

Guo, Y., Qin, G., Gu, H., & Qu, L. J. (2009). Dof5. 6/HCA2, a Dof transcription factor
gene, regulates interfascicular cambium formation and vascular tissue development in
Arabidopsis. The Plant Cell, *21*(11), 3518-3534. doi: 10.1105/tpc.108.064139.

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., ... &
Regev, A. (2013). De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. Nature Protocols, *8*(8), 1494-1512.
doi: 10.1038/nprot.2013.084.

Hachiya, T., & Sakakibara, H. (2017). Interactions between nitrate and ammonium in their
uptake, allocation, assimilation, and signaling in plants. Journal of Experimental Botany,
68(10), 2501-2512. doi.org/10.1093/jxb/erw449.

Hawkins, B. J., & Robbins, S. (2010). pH affects ammonium, nitrate and proton fluxes in
the apical region of conifer and soybean roots. Physiologia Plantarum, *138*(2), 238-247.
doi.org/10.1111/j.1399-3054.2009.01317.x.

Helariutta, Y., Fukaki, H., Wysocka-Diller, J., Nakajima, K., Jung, J., Sena, G., ... &
Benfey, P. N. (2000). The SHORT-ROOT gene controls radial patterning of the
Arabidopsis root through radial signaling. Cell, *101*(5), 555-567. doi.org/10.1016/S0092-
8674(00)80865-X.

Hong, Z. H., Qing, T., Schubert, D., Kleinmanns, J. A., & Liu, J. X. (2019). BLISTER-
regulated vegetative growth is dependent on the protein kinase domain of ER stress
modulator IRE1A in *Arabidopsis thaliana*. PLoS Genetics, *15*(12), e1008563. doi:
10.1371/journal.pgen.1008563.

Hou, H., Erickson, J., Meservy, J., & Schultz, E. A. (2010). FORKED1 encodes a PH
domain protein that is required for PIN1 localization in developing leaf veins. The Plant
Journal, *63*(6), 960-973. doi.org/10.1111/j.1365-313X.2010.04291.x.

Houben, M., & Van de Poel, B. (2019). 1-Aminocyclopropane-1-carboxylic acid oxidase
(ACO): the enzyme that makes the plant hormone ethylene. Frontiers in Plant Science, *10*,
695. doi: 10.3389/fpls.2019.00695.

Ishiyama, K., Hayakawa, T. & Yamaya, T. (1998). Expression of NADH-dependent
glutamate synthase protein in the epidermis and exodermis of rice roots in response to the
supply of ammonium ions. Planta, *204*(3), 288-294. doi: 10.1007/s004250050258.

Ishiyama, K., Inoue, E., Tabuchi, M., Yamaya, T. & Takahashi, H. (2004a). Biochemical
background and compartmentalized functions of cytosolic glutamine synthetase for active
ammonium assimilation in rice roots. Plant and Cell Physiology, *45*(11), 1640-1647. doi:
10.1093/pcp/pch190.

Ishiyama, K., Inoue, E., Watanabe-Takahashi, A., Obara, M., Yamaya, T. & Takahashi, H.
(2004b). Kinetic properties and ammonium-dependent regulation of cytosolic isoenzymes
of glutamine synthetase in *Arabidopsis*. Journal of Biological Chemistry, *279*(16), 16598-
16605. doi: 10.1074/jbc.M313710200.

Jia, Z., & von Wirén, N. (2020). Signaling pathways underlying nitrogen-dependent
changes in root system architecture: from model to crop species. Journal of Experimental
Botany, *71*(15), 4393-4404. https://doi.org/10.1093/jxb/eraa033

Jia, L.T., Xie, Y.M., Wang, Z., Luo, L., Zhang, C., Pelissier, P.M, … & Xuan, W. (2020)
Rice plants respond to ammonium stress by adopting a helical root growth pattern. The
Plant Journal, *104*(4), 1023-1037. doi: 10.1111/tpj.14978.

Kamada-Nobusada, T., Makita, N., Kojima, M., & Sakakibara, H. (2013). Nitrogen-
dependent regulation of de novo cytokinin biosynthesis in rice: the role of glutamine
metabolism as an additional signal. Plant and Cell Physiology, *54*(11), 1881-1893.
doi.org/10.1093/pcp/pct127.

Kamiya, M., Higashio, S. Y., Isomoto, A., Kim, J. M., Seki, M., Miyashima, S., &
Nakajima, K. (2016). Control of root cap maturation and cell detachment by BEARSKIN
transcription factors in *Arabidopsis*. Development, *143*(21), 4063-4072.doi:
10.1242/dev.142331.

Kim, H., Zhou, J., Kumar, D., Jang, G., Ryu, K. H., Sebastian, J., ... & Lee, J. Y. (2020).
SHORTROOT-mediated intercellular signals coordinate phloem development in
Arabidopsis roots. The Plant Cell, *32*(5), 1519-1535. doi.org/10.1105/tpc.19.00455.

Konishi, N., Ishiyama, K., Beier, M.P., Inoue, E., Kanno, K., Yamaya, T., Takahashi, H. &
Kojima, S. (2017). Contribution of two glutamine synthetase isozymes to ammonium
assimilation in *Arabidopsis* roots. Journal of Experimental Botany, *68*(3), 613-625. doi:
10.1093/jxb/erw454.

Kronzucker, H. J., Siddiqi, M. Y., & Glass, A. D. (1997). Conifer root discrimination
against soil nitrate and the ecology of forest succession. Nature, *385*(6611), 59-61.
doi.org/10.1038/385059a0.

Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: molecular evolutionary genetics

analysis version 7.0 for bigger datasets. Molecular Biology and Evolution, *33*(7), 1870-

1874.doi.org/10.1093/molbev/msw054.

Kuroha, T., Tokunaga, H., Kojima, M., Ueda, N., Ishida, T., Nagawa, S., ... & Sakakibara,

H. (2009). Functional analyses of LONELY GUY cytokinin-activating enzymes reveal the

importance of the direct activation pathway in *Arabidopsis*. The Plant Cell, *21*(10), 3152-

3169. doi.org/10.1105/tpc.109.068676.

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation

network analysis. BMC Bioinformatics, *9*, 559. https://doi.org/10.1186/1471-2105-9-559.

Liao, Z., Chen, M., Guo, L., Gong, Y., Tang, F., Sun, X., & Tang, K. (2004). Rapid

isolation of high-quality total RNA from *Taxus* and *Ginkgo*. Preparative Biochemistry and

Biotechnology, *34*(3), 209-214. doi.org/10.1081/PB-200026790.

Lima, J. E., Kojima, S., Takahashi, H., & von Wirén, N. (2010). Ammonium triggers lateral

root branching in *Arabidopsis* in an AMMONIUM TRANSPORTER1; 3-dependent

manner. The Plant Cell, *22*(11), 3621-3633.doi.org/10.1105/tpc.110.076216.

Liu, Y., Lai, N., Gao, K., Chen, F., Yuan, L., & Mi, G. (2013). Ammonium inhibits

primary root growth by reducing the length of meristem and elongation zone and

decreasing elemental expansion rate in the root apex in *Arabidopsis thaliana*. PloS One,

8(4), e61031. doi.org/10.1371/journal.pone.0061031.

Liu, Y., & von Wirén, N. (2017). Ammonium as a signal for physiological and

morphological responses in plants. Journal of Experimental Botany, *68*(10), 2581-2592.

doi.org/10.1093/jxb/erx086.

1002

1003   Love, J., Björklund, S., Vahala, J., Hertzberg, M., Kangasjärvi, J., & Sundberg, B. (2009).

1004   Ethylene is an endogenous stimulator of cell division in the cambial meristem of *Populus*.

1005   Proceedings of the National Academy of Sciences, *106*(14), 5984-5989.
1006   doi.org/10.1073/pnas.0811660106.

1007

1008   Lucas, M., Swarup, R., Paponov, I. A., Swarup, K., Casimiro, I., Lake, D., ... & Bennett,
1009   M. J. (2011). Short-Root regulates primary, lateral, and adventitious root development in
1010   *Arabidopsis*. Plant Physiology, *155*(1), 384-398. doi.org/10.1104/pp.110.165126.

1011

1012   Mao, C., He, J., Liu, L., Deng, Q., Yao, X., Liu, C., ... & Ming, F. (2020). OsNAC2
1013   integrates auxin and cytokinin pathways to modulate rice root development. Plant
1014   Biotechnology Journal, *18*(2), 429-442. doi.org/10.1111/pbi.13209.

1015

1016   Marhava, P., Bassukas, A. E. L., Zourelidou, M., Kolb, M., Moret, B., Fastner, A., ... &
1017   Hardtke, C. S. (2018). A molecular rheostat adjusts auxin flux to promote root protophloem
1018   differentiation. Nature, *558*(7709), 297-300. doi.org/10.1038/s41586-018-0186-z.

1019

1020   Meier, M., Liu, Y., Lay-Pruitt, K.S., Takahashi, H. & von Wirén, N. (2020) Auxin-
1021   mediated root branching is determined by the form of available nitrogen. Nature Plants,
1022   *6*(9), 1136-1145. doi: 10.1038/s41477-020-00756-2.

1023

1024   Men, S., Boutté, Y., Ikeda, Y., Li, X., Palme, K., Stierhof, Y. D., ... & Grebe, M. (2008).
1025   Sterol-dependent endocytosis mediates post-cytokinetic acquisition of PIN2 auxin efflux
1026   carrier polarity. Nature Cell Biology, *10*(2), 237-244. doi.org/10.1038/ncb1686.

1027

1028   Miller, A.J. & Cramer, M.D. (2005). Root nitrogen acquisition and assimilation. In Root
1029   physiology: From gene to function. Springer, Dordrecht, 1-36, doi.org/10.1007/1-4020-
1030   4099-7_1.

1031

1032   Miyashima, S., Koi, S., Hashimoto, T., & Nakajima, K. (2011). Non-cell-autonomous
1033   microRNA165 acts in a dose-dependent manner to regulate multiple differentiation status
1034   in the *Arabidopsis* root. Development, *138*(11), 2303-2313. doi: 10.1242/dev.060491.

1035

1036 Miyashima, S., Roszak, P., Sevilem, I., Toyokura, K., Blob, B., Heo, J. O., ... & Helariutta,
1037 Y. (2019). Mobile PEAR transcription factors integrate positional cues to prime cambial
1038 growth. Nature, *565*(7740), 490-494. doi.org/10.1038/s41586-018-0839-y.

1039

1040 Moreno-Risueno, M. Á., Martínez, M., Vicente-Carbajosa, J., & Carbonero, P. (2007). The
1041 family of DOF transcription factors: from green unicellular algae to vascular plants.
1042 Molecular Genetics and Genomics, *277*(4), 379-390. doi.org/10.1007/s00438-006-0186-9.

1043

1044 Moritake, S., Taira, S., Ichiyanagi, Y., Morone, N., Song, S. Y., Hatanaka, T., ... & Setou,
1045 M. (2007). Functionalized nano-magnetic particles for an in vivo delivery system. Journal
1046 of Nanoscience and Nanotechnology, *7*(3), 937-944. doi.org/10.1166/jnn.2007.216.

1047

1048 Nakajima, K., Sena, G., Nawy, T., & Benfey, P. N. (2001). Intercellular movement of the
1049 putative transcription factor SHR in root patterning. Nature, *413*(6853), 307-
1050 311.doi.org/10.1038/35095061.

1051

1052 Näsholm, T., Ekblad, A., Nordin, A., Giesler, R., Högberg, M., & Högberg, P. (1998).
1053 Boreal forest plants take up organic nitrogen. Nature, *392*(6679), 914-
1054 916.doi.org/10.1038/31921.
1055 NCBI Resource Coordinators. (2016). Database resources of the national center for
1056 biotechnology information. Nucleic Acids Research, *44*(D1), D7-D19.

1057

1058 Ochando, I., Jover-Gil, S., Ripoll, J. J., Candela, H., Vera, A., Ponce, M. R., ... & Micol, J.
1059 L. (2006). Mutations in the microRNA complementarity site of the INCURVATA4 gene
1060 perturb meristem function and adaxialize lateral organs in *Arabidopsis*. Plant Physiology,
1061 *141*(2), 607-619. doi.org/10.1104/pp.106.077149.

1062

1063 Ötvös, K., Marconi, M., Vega, A., O'Brien, J., Johnson, A., Abualia, R., … & Benková, E.
1064 (2021). Modulation of plant root growth by nitrogen source-defined regulation of polar
1065 auxin transport. The EMBO Journal, *40*(3), e106862.
1066 https://doi.org/10.15252/embj.2020106862

1067

1068   Ortigosa, F., Valderrama-Martín, J. M., Urbano-Gámez, J. A., García-Martín, M. L., Ávila,
1069   C., Cánovas, F. M., & Cañas, R. A. (2020). Inorganic nitrogen form determines nutrient
1070   allocation and metabolic responses in maritime pine seedlings. Plants, *9*: 481,
1071   doi.org/10.3390/plants9040481.

1072

1073   Ortigosa, F., Lobato-Fernández, C., Pérez-Claros, J. A., Cantón, F. R., Ávila, C., Cánovas,
1074   F. M., & Cañas, R. A. (2021). Epitranscriptome changes triggered by ammonium nutrition
1075   regulate the proteome response of maritime pine roots. bioRxiv.
1076   doi.org/10.1101/2021.04.20.440618.

1077

1078   Patterson, K., Cakmak, T., Cooper, A., Lager, I. D. A., Rasmusson, A. G., & Escobar, M.
1079   A. (2010). Distinct signalling pathways and transcriptome response signatures differentiate
1080   ammonium-and nitrate-supplied plants. Plant, Cell & Environment, *33*(9), 1486-
1081   1501.doi.org/10.1111/j.1365-3040.2010.02158.x.

1082

1083   Prigge, M. J., Otsuga, D., Alonso, J. M., Ecker, J. R., Drews, G. N., & Clark, S. E. (2005).
1084   Class III homeodomain-leucine zipper gene family members have overlapping,
1085   antagonistic, and distinct roles in *Arabidopsis* development. The Plant Cell, *17*(1), 61-76.
1086   doi.org/10.1105/tpc.104.026161.

1087

1088   Purdy, S. J., Bussell, J. D., Nelson, D. C., Villadsen, D., & Smith, S. M. (2011). A nuclear-
1089   localized protein, KOLD SENSITIV-1, affects the expression of cold-responsive genes
1090   during prolonged chilling in *Arabidopsis*. Journal of Plant Physiology, *168*(3), 263-269.
1091   doi.org/10.1016/j.jplph.2010.07.001.

1092

1093   Ravazzolo, L., Trevisan, S., Forestan, C., Varotto, S., Sut, S., Dall'Acqua, S., ... &
1094   Quaggiotti, S. (2020). Nitrate and ammonium affect the overall maize response to nitrogen
1095   availability by triggering specific and common transcriptional signatures in roots.
1096   International Journal of Molecular Sciences, *21*(2), 686. doi.org/10.3390/ijms21020686.

1097

1098   Ren, Q., Zhou, Y., & Zhou, X. (2020). Combined Transcriptome and Proteome Analysis of
1099   Masson Pine (Pinus massoniana Lamb.) Seedling root in response to nitrate and ammonium

1100 supplementations. International Journal of Molecular Sciences, *21*(20), 7548.
1101 doi.org/10.3390/ijms21207548.

1102

1103 Rich-Griffin, C., Eichmann, R., Reitz, M. U., Hermann, S., Woolley-Allen, K., Brown, P.
1104 E., ... & Schäfer, P. (2020). Regulation of cell type-specific immunity networks in
1105 Arabidopsis roots. Plant Cell, 32(9), 2742-2762. doi.org/10.1105/tpc.20.00154.

1106

1107 Ritz C, Spiess AN. 2008. Ritz, C., & Spiess, A. N. (2008). qpcR: an R package for
1108 sigmoidal model selection in quantitative real-time polymerase chain reaction analysis.
1109 Bioinformatics, *24*(13), 1549-1551. doi.org/10.1093/bioinformatics/btn227.

1110

1111 Roodbarkelari, F., & Groot, E. P. (2017). Regulatory function of homeodomain‐leucine
1112 zipper (HD‐ZIP) family proteins during embryogenesis. New Phytologist, *213*(1), 95-104.
1113 doi.org/10.1111/nph.14132.

1114

1115 Rubio, V., Linhares, F., Solano, R., Martín, A. C., Iglesias, J., Leyva, A., & Paz-Ares, J.
1116 (2001). A conserved MYB transcription factor involved in phosphate starvation signaling
1117 both in vascular plants and in unicellular algae. Genes & Development, *15*(16), 2122-2133.
1118 doi: 10.1101/gad.204401.

1119

1120 Sablowski, R. (2011). Plant stem cell niches: from signalling to execution. Current Opinion
1121 in Plant Biology, *14*(1), 4-9. doi.org/10.1016/j.pbi.2010.08.001.

1122

1123 Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for
1124 reconstructing phylogenetic trees. Molecular Biology and Evolution, *4*(4), 406-425.
1125 doi.org/10.1093/oxfordjournals.molbev.a040454.

1126

1127 Schatlowski, N., Stahl, Y., Hohenstatt, M. L., Goodrich, J., & Schubert, D. (2010). The
1128 CURLY LEAF interacting protein BLISTER controls expression of polycomb-group target
1129 genes and cellular differentiation of *Arabidopsis thaliana*. The Plant Cell, *22*(7), 2291-
1130 2305. doi: 10.1105/tpc.109.073403.

1131

1132   Schoenbohm, C., Martens, S., Eder, C., Forkmann, G., & Weisshaar, B. (2000).

1133   Identification of the Arabidopsis thaliana flavonoid 3'-hydroxylase gene and functional

1134   expression of the encoded P450 enzyme. Biological Chemistry, *381*(8):749-53. doi:

1135   10.1515/BC.2000.095.

1136

1137   Schwarz, R. & Dayhoff. M. (1979). Matrices for detecting distant relationships. In Dayhoff

1138   M., editor, Atlas of protein sequences:353-58. National Biomedical Research Foundation.

1139

1140   Sena, G., Jung, J. W., & Benfey, P. N. (2004). A broad competence to respond to SHORT

1141   ROOT revealed by tissue-specific ectopic expression. Development, *131*(12):2817-26 doi:

1142   10.1242/dev.01144.

1143

1144   Shiono, K., & Taira, S. (2020). Imaging of multiple plant hormones in roots of rice (Oryza

1145   sativa) using nanoparticle-assisted laser desorption/Ionization mass spectrometry. Journal

1146   of Agricultural and Food Chemistry, *68*(24), 6770-6775. doi: 10.1021/acs.jafc.0c00749.

1147

Shtratnikova, V. Y., Kudryakova, N. V., Kudoyarova, G. R., Korobova, A. V., Akhiyarova,

G. R., Danilova, M. N., ... & Kulaeva, O. N. (2015). Effects of nitrate and ammonium on

growth of Arabidopsis thaliana plants transformed with the ARR5:: GUS construct and a

role for cytokinins in suppression of disturbances induced by the presence of ammonium.

Russian Journal of Plant Physiology, *62*(6), 741-752.

doi.org/10.1134/S1021443715060151.

1154

1155   Smit, M. E., Llavata-Peris, C. I., Roosjen, M., van Beijnum, H., Novikova, D., Levitsky,

1156   V., ... & Weijers, D. (2020). Specification and regulation of vascular tissue identity in the

1157   Arabidopsis embryo. Development, *147*(8), dev186130. doi: 10.1242/dev.186130. PMID:

1158   32198154.

1159

1160   Sun, L., Di, D., Li, G., Kronzucker, H. J., & Shi, W. (2017). Spatio-temporal dynamics in

1161   global rice gene expression (*Oryza sativa* L.) in response to high ammonium stress. Journal

1162   of plant physiology, *212*, 94-104. doi.org/10.1016/j.jplph.2017.02.006.

1163

1164 Sun, L., Di, D. W., Li, G., Kronzucker, H. J., Wu, X., & Shi, W. (2020). Endogenous ABA
1165 alleviates rice ammonium toxicity by reducing ROS and free ammonium via regulation of
1166 the SAPK9–bZIP20 pathway. Journal of Experimental Botany, *71*(15), 4562-
1167 4577.doi.org/10.1093/jxb/eraa076.

1168

1169 Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO summarizes and
1170 visualizes long lists of gene ontology terms. PloS One, *6*(7), e21800. doi:
1171 10.1371/journal.pone.0021800.

1172

1173 Takei, K., Ueda, N., Aoki, K., Kuromori, T., Hirayama, T., Shinozaki, K., ... & Sakakibara,
1174 H. (2004). AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in
1175 *Arabidopsis*. Plant and Cell Physiology, *45*(8), 1053-1062. doi.org/10.1093/pcp/pch119.

1176

1177 Tamura, W., Hidaka, Y., Tabuchi, M., Kojima, S., Hayakawa, T., Sato, T., …& Yamaya, T.
1178 (2010) Reverse genetics approach to characterize a function of NADH-glutamate synthase1
1179 in rice plants. Amino Acids, *39*(4), 1003-1012. doi: 10.1007/s00726-010-0531-5.

1180

1181 Tan, W., Zhang, D., Zhou, H., Zheng, T., Yin, Y., & Lin, H. (2018). Transcription factor
1182 HAT1 is a substrate of SnRK2. 3 kinase and negatively regulates ABA synthesis and
1183 signaling in *Arabidopsis* responding to drought. PLoS Genetics, *14*(4), e1007336. doi:
1184 10.1371/journal.pgen.1007336.

1185

1186 Tian, T., Liu, Y., Yan, H., You, Q., Yi, X., Du, Z., ... & Su, Z. (2017). agriGO v2. 0: a GO
1187 analysis toolkit for the agricultural community, 2017 update. Nucleic Acids Research,
1188 *45*(W1), W122-W129. doi.org/10.1093/nar/gkx382.

1189

1190 Walker, L., Boddington, C., Jenkins, D., Wang, Y., Grønlund, J. T., Hulsmans, J., &
1191 Gifford, M. L. (2017). Changes in gene expression in space and time orchestrate
1192 environmentally mediated shaping of root architecture. The Plant Cell, 29(10), 2393-2412.
1193 doi.org/10.1105/tpc.16.00961.

1194

1195   Wang, S., Chang, Y., Guo, J., Zeng, Q., Ellis, B. E., & Chen, J. G. (2011). *Arabidopsis*
1196   ovate family proteins, a novel transcriptional repressor family, control multiple aspects of
1197   plant   growth   and   development.   PLoS   One,   *6*(8),   e23896.   doi:
1198   10.1371/journal.pone.0023896.

1199

1200   Ward, J. M., Cufr, C. A., Denzel, M. A., & Neff, M. M. (2005). The Dof transcription
1201   factor OBP3 modulates phytochrome and cryptochrome signaling in *Arabidopsis*. The Plant
1202   Cell, *17*(2), 475-485. doi: 10.1105/tpc.104.027722.

1203

1204   Warren, C. R., & Adams, M. A. (2002). Possible causes of slow growth of nitrate-supplied
1205   *Pinus pinaster*. Canadian Journal of Forest Research, *32*(4), 569-580. doi.org/10.1139/x01-
1206   225.

1207

1208   Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H., & Schmu☐lling, T.
1209   (2003). Cytokinin-deficient transgenic *Arabidopsis* plants show multiple developmental
1210   alterations indicating opposite functions of cytokinins in the regulation of shoot and root
1211   meristem activity. The Plant Cell, *15*(11), 2532-2550. doi.org/10.1105/tpc.014928.

1212

1213   Willemsen, V., Bauch, M., Bennett, T., Campilho, A., Wolkenfelt, H., Xu, J., ... & Scheres,
1214   B. (2008). The NAC domain transcription factors FEZ and SOMBRERO control the
1215   orientation of cell division plane in *Arabidopsis* root stem cells. Developmental Cell, *15*(6),
1216   913-922. doi: 10.1016/j.devcel.2008.09.019.

1217

1218   Xu, J., Hofhuis, H., Heidstra, R., Sauer, M., Friml, J., & Scheres, B. (2006). A molecular
1219   framework   for   plant   regeneration.   Science,   *311*(5759),   385-388.
1220   doi.org/10.1126/science.1121790.

1221

1222   Youssef, C., Bizet, F., Bastien, R., Legland, D., Bogeat-Triboulot, M. B., & Hummel, I.
1223   (2018). Quantitative dissection of variations in root growth rate: a matter of cell
1224   proliferation or of cell expansion?. Journal of Experimental Botany, *69*(21), 5157-5168.
1225   doi.org/10.1093/jxb/ery272.

1226

1227   Zhang, D., Ye, H., Guo, H., Johnson, A., Zhang, M., Lin, H., & Yin, Y. (2014).
1228   Transcription factor HAT 1 is phosphorylated by BIN 2 kinase and mediates
1229   brassinosteroid repressed gene expression *in Arabidopsis*. The Plant Journal, *77*(1), 59-70.
1230   doi: 10.1111/tpj.12368.

1231

1232   Zhang, S., & Wang, X. (2017). One new kind of phytohormonal signaling integrator: Up-
1233   and-coming GASA family genes. Plant Signaling & Behavior, *12*(2), e1226453. doi:
1234   10.1080/15592324.2016.1226453.

1235

1236   Zhang, W., Swarup, R., Bennett, M., Schaller, G. E., & Kieber, J. J. (2013). Cytokinin
1237   induces cell division in the quiescent center of the *Arabidopsis* root apical meristem.
1238   Current Biology, *23*(20), 1979-1989. doi.org/10.1016/j.cub.2013.08.008.

1239

1240   Zhou, W., Wei, L., Xu, J., Zhai, Q., Jiang, H., Chen, R., ... & Li, C. (2010). *Arabidopsis*
1241   tyrosylprotein sulfotransferase acts in the auxin/PLETHORA pathway in regulating
1242   postembryonic maintenance of the root stem cell niche. The Plant Cell, *22*(11), 3692-3709.
1243   doi.org/10.1105/tpc.110.075721.

1244

1245   Zou, N., Li, B., Chen, H., Su, Y., Kronzucker, H. J., Xiong, L., ... & Shi, W. (2013).
1246   GSA-1/ARG 1 protects root gravitropism in *Arabidopsis* under ammonium stress. New
1247   Phytologist, *200*(1), 97-111. doi.org/10.1111/nph.12365.

1248

1249   Zúñiga-Mayo, V. M., Marsch-Martínez, N., & de Folter, S. (2012). The class II HD-ZIP
1250   JAIBA gene is involved in meristematic activity and important for gynoecium and fruit
1251   development in *Arabidopsis*. Plant Signaling & Behavior, *7*(11), 1501-1503. doi:
1252   10.4161/psb.21901.

1253

1254 **Table 1.** Most representative DE transcripts coding for TFs and hormone related proteins.

TF in maritime pine	Gene ID	Tissue	Transcript in *A.thaliana*	Gene ID	Process involved	References
PpDOF11	*pp_58878*	RC	*HCA2/ DOF5.6*	*AT5G62940*	Root tip vascular proliferation: radial growth of protophloem sieve elements	Guo et al., 2009 Miyashima et al.,2019
PpSHR	*pp_246622*	RC	*SHR*	*AT4G37650*	- Quiescent center cells specification - Asymmetric cell division related to rootradial pattern formation - Phloem development	Fukaki et al., 1998 Helariutta et al., 2000 Cui et al., 2007 Kim et al., 2020
PpGATA20-like	*pp_204819*	RC	*GATA20*	*AT2G18380*	Root tip vascular development	Smit et al., 2020
PpSGR/IDD 16	*pp_174625*	RC	*IDD14*	*AT1G68130*	- Lateral organ morphogenesis and gravitropic responses - Establishment of auxin gradients (auxin biosynthesis and transport)	Cui et al., 2013
PpPHR1-like	*pp_64350*	RD C	*PHR1*	*AT4G28610*	Phosphate starvation	Rubio et al., 2001
PpOFP17	*pp_80824*	RD C	*OFP17*	*AT2G30395*	Plant growth and development	Wang et al., 2011
PpBLISTER-like	*pp_170664*	RD C	*BLISTER*	*AT3G23980*	Plant vegetative growth	Schatlowskiet al., 2010 Hong et al., 2019

PpDOF2	*pp_119525*	RDC	*PEAR1/ DOF2.4*	*AT2G37590*	Root tip vascular proliferation: radial growth ofprotophloem sieve elements	Miyashima et al.,2019
PpDOF12	*pp_85381*	RM	*CDF3*	*AT3G47500*	Involved in abiotic stress tolerance and nitrogen response	Corrales et al., 2017 Domínguez-Figueroaet al., 2020
PpDOF1	*pp_233816*	RM	*OPB3*	*AT3G55370*	Regulator of phyB and cry1 signaling pathways	Ward et al., 2005
PpHAT-like	*pp_141831*	RM	*HAT1*	*AT4G17460*	- Inflorescence meristem regulation, floralmeristem determination, gynoecium medial domain development - Repressor of BR responsive genes and ABA biosynthesis	Zúñiga-Mayo et al., 2012 Zhang et al., 2014 Tan et al., 2018
PpINCURV ATA-like	*pp_85012*	RM	*ATBH-15*	*AT1G52150*	Root meristem patterning and putatively in the regulation of procambial and vascular tissue formation or maintenance.	Ochando et al., 2006
PpNAC31	*pp_168080*	RM	*SMB*	*AT1G79580*	Cellular maturation of root cap regulation	Willemsen et al., 2008 Bennett et al., 2010 Fendrych et al., 2014
PpNAC38	*pp_72282*	RM	*BRN1/BRN2*	*AT1G33280*	Cellular maturation of root cap regulation	Bennett et al., 2010
PpVAN3/FKD2	*pp_60103*	RC and RM	*VAN3/ FORKED2*	*AT5G13300*	Plant vascular tissue formation by PIN transporters localization	Steynen & Schultz, 2003 Naramoto et al., 2009 Hou et al., 2010
Cytokinin riboside 5'-monophosphate phosphoribohydrolase	*pp_222714*	RC	*LOG3*	*AT2G37210*	Cytokinin activation	Kuroha et al., 2009

Cycloeucalenol cycloisomerase	*pp_189071*	RC	*CPII*	*AT5G50375*	PIN2 endocytosis	Men et al., 2008
Sulfotransferase	*pp_208269*	RDC	*SULT20B1*	*AT3G45070*	Flavonoids sulfation	Hashiguchi et al., 2013
PpABA2	*pp_198945*	RDC and RM	*ABA2*	*AT1G52340*	Biosynthesis of abscisic acid	González-Guzmán et al., 2002
PpCYP90B1	*pp_247190*	RDC	*CYP90B1*	*AT3G50660*	Biosynthesis of brassinosteroids	Fujita et al., 2006
PpTIR1	*pp_238715*	RDC	*TIR1*	*AT3G62980*	Auxin signalling pathway	Dharmasiri et al., 2005 Salehin et al., 2015
PpCYP87A3	*pp_81114*	RDC	*CYP87A2*	*AT1G12740*	Putatively involved in hormone biosynthesis and/or auxin signaling	Chaban et al., 2003 Bark et al., 2011
GASA gibberellin regulated cysteine richprotein	*pp_111379*	RDC	*GASA10*	*AT5G59845*	Putatively involved in response tophytohormones	Zhang & Wang, 2017
HMG-CoA reductase	*pp_144669*	RDC	*HMGR*	*AT2G17370*	Regulatory enzyme of the cytokinin and brassinosteroid biosynthesis pathways	Antolín-Llovera et al., 2011
PpABA2	*pp_209084*	RDC	*ABA2*	*AT1G52340*	Biosynthesis of abscisic acid	González-Guzmán etal., 2002

Cytokinin dehydrogenase	pp_215376	RDC	CKX1	AT2G41510	Catalyzes the oxidation of cytokinins	Werner et al., 2003
PpACO	pp_202153	RM	ACO	AT2G19590	Ethylene biosynthesis	Houben et al., 2019
Flavonoid 3'-monooxygenase	pp_203885	RM	F3'H	AT5G07990	Involved in flavonoids biosynthesispathway from naringenin to eriodictyol and dihydrokaempferol to dihydroquercetin	Schoenbohm et al., 2000
EXORDIUM	pp_129078	RM	EXO	AT4G08950	Putatively involved in brassinosteroids resopnse in roots	Coll-Garcia et al., 2004

1255

 Figure legends

 Figure 1. Experimental design. Timeline from germination to harvest. RT-qPCR data for
 the validation of the experimental design are included. *Glutamine synthetase 1b* (*PpGS1b*);
 NADH-dependent glutamate synthase (*PpNADH-GOGAT*); *antimicrobial peptide 1*
 (*PpAMP1*); *S-adenosyl methionine synthase* (*PpSAMS*) Significant differences were
 determined with a *t*-test (* *P*-value < 0.05; ** *P*-value < 0.01). Error bars show SE with n =
 3. Isolated regions from the root apex: cap (RC), meristem (RM), developing cortex (RDC)
 and developing vessels (RDV).

 Figure 2. Low-input RNA-seq results in the different tissues of maritime pine root tip. (**A**)
 Venn diagram of down-regulated DE transcripts. (**B**) Venn diagram of up-regulated DE
 transcripts. (**C**) Heatmap and hierarchical clustering of DE transcripts. Expression values
 correspond to median CPM between biological/experimental replicates. The hierarchical
 clustering was made employing the Ward's minimum variance method. (**D**) Experimental
 validation of low-input RNA-seq results for 8 DE transcripts through RT-qPCR. RC, the
 root cap; RM, root meristem; RDC, root developing cortex; and RDV, root developing
 vessels. *SHORT-ROOT* (*PpSHR*); *nitrate transporter 1 / peptide transporter family 3.1*
 (*PpNPF3.1*); *cycloeucalenol cycloisomerase* (*PpCPI1*); *DOF transcription factor 12*
 (*PpDOF12*); *NAC transcription factor 38* (*PpNAC38*); *aminocyclopropane-1-carboxylic*
 acid oxidase (*PpACO*); *antimicrobial peptide 1* (*PpAMP1*); *BLISTER-like transcription*
 factor (*PpBLISTER-like*).

 Figure 3. Resume of the main functional results obtained in the low-input RNA-Seq
 analysis from LCM isolated samples. Functions and DE transcripts in (**A**) the root cap
 (RC), (**B**) root meristem (RM), (**C**) root developing cortex (RDC) and (**D**) root developing
 vessels (RDV). Blue arrows indicate downregulated transcripts, red arrows upregulated
 ones. (**E**) Significant GO terms from Biological Processes category after a SEA analysis.
 (**F**) Significant GO terms from Molecular Function category after a SEA analysis.

 Figure 4. Expression heatmaps of differentially expressed transcriptional regulators and
 hormone related transcripts. (A) Transcription factors and other putative regulators. (B)

1288   Transcripts involved in hormone metabolism, transport and signaling. CPM expression

1289   values were normalized by transcript.

1290

Figure 5. Phytohormone localization in root apex sections. The main phytohormones were

localized in root apex cuts of control and 3 mM NH_4^+ treated seedlings. The phytohormone

identification and imaging was made using nano-particle assisted laser

desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). (**A**) Auxin / Indole-

3-acetic acid (IAA); (**B**) cytokinin (CK); (**C**) aminocyclopropane-1-carboxylic acid (ACC);

(**D**) abscisic acid (ABA). Size bars correspond to 500 µm. Column graphs show relative

hormone accumulations respect to control samples in the approximate LCM areas. Dashed

lines mark hormone accumulation in the control roots.

1299

Figure 6. Root growth and architecture parameters. Pine seedlings were cultivated for 74

days and irrigated once a week with 50 mL of water, 3 mM KNO_3 and 3 mM NH_4Cl,

respectively. Six different parameters were measured after harvesting: (**A**) weight of whole

roots; (**B**) weight of primary root; (**C**) primary root length; (**D**) number of lateral roots; (**E**)

weight of lateral roots; (**F**) lateral root density. Significant differences were determined

with a one-way ANOVA. Letters above the conditions show significant differences based

on a Turkey post-hoc test ($P < 0.05$) with n = 30.

1307

Figure 7. Schematic representation of the transcriptional response induced by NH_4^+ supply

in maritime pine seedling roots 24 h after irrigation. Transcripts in red letters are

upregulated. Transcripts in blue letters are downregulated. The root tip tissues are the root

cap (RC) root meristem (RM) and root developing cortex (RDC). Results for root

developing vessels (RDV) are not shown since they were very limited.

Supporting information

Video S1. LCM cutting procedure.

Table S1. LCM low-input RNA sequencing results.

Table S2. List of used primers.

Table S3. Protein sequences of DOF transcription factors in maritime pine, *Arabidopsis*
and rice.

Table S4. Protein sequences of NAC transcription factors in maritime pine and
Arabidopsis.

Method S1. Complete LCM and cDNA amplification procedures.

Figure S1. Aspect of root architecture after long-term nutritional treatments.

Figure S2. MDS plot of the LCM low-input RNA-seq samples.

Figure S3. RNA-seq differential expression analysis between tissues. (**A**) Venn diagrams
shows the amounts of tissue-specific DE transcripts. (**B**) Functions of the transcripts that
were always significantly expressed in a specific tissue. Significant GO terms from
Biological Processes category after a SEA analysis with a 0.4 cut-off value for
dispensability. (**C**) Expression of transcription factor PpFAMA-like (pp_179856) in the
different isolated tissues of the root apex. PpFAMA-like was differentially expressed in the
RM respect to the rest of tissues but not in the nutritional treatment comparison.

Figure S4. Phylogenetic tree of DOF transcription factor family in maritime pine,
Arabidopsis and rice. The protein sequences employed to infer the phylogenetic tree are
presented in the Table S3. The protein sequence alignment was made with Muscle (Edgar,
2004). The evolutionary history was inferred using the Neighbor-Joining method (Saitou
and Nei, 1987). The bootstrap consensus tree inferred from 1000 replicates is taken to
represent the evolutionary history of the taxa analyzed (Felsenstein, 1985). Branches
corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed.
The evolutionary distances were computed using the Dayhoff matrix-based method
(Schwarz and Dayhoff, 1979) and are in the units of the number of amino acid substitutions
per site. The analysis involved 78 amino acid sequences. All positions containing gaps and
missing data were eliminated. There were 54 positions in the final dataset mainly
corresponding to the DOF domain. Evolutionary analyses were conducted in MEGA7
(Kumar et al., 2016). DOFs from maritime pine have a red dot and their names are in bold.
Abbreviations: At, *Arabidopsis thaliana*; Pp, *Pinus pinaster*; Os, *Oryza sativa*.

Figure S5. Phylogenetic tree of NAC transcription factor family in maritime pine and
Arabidopsis. The protein sequences employed to infer the phylogenetic tree are presented
in the Table S4. The protein sequence alignment was made with Muscle (Edgar, 2004). The
evolutionary history was inferred using the Neighbor-Joining method (Saitou and Nei,
1987). The bootstrap consensus tree inferred from 1000 replicates is taken to represent the
evolutionary history of the taxa analyzed (Felsenstein, 1985). Branches corresponding to
partitions reproduced in less than 50% bootstrap replicates are collapsed. The evolutionary
distances were computed using the Dayhoff matrix-based method (Schwarz and Dayhoff,
1979) and are in the units of the number of amino acid substitutions per site. The analysis
involved 151 amino acid sequences. All positions with less than 80% site coverage were
eliminated. That is, fewer than 20% alignment gaps, missing data, and ambiguous bases
were allowed at any position. There were 261 positions in the final dataset. Evolutionary
analyses were conducted in MEGA7 (Kumar et al., 2016). NACs from maritime pine have
a red dot and their names are in bold.
Figure S6. Gene expression correlation network and functions. (**A**) Significant
transcription factors for ammonium treatment that are hubs in a WGCNA gene correlation
network and their connections with their correlated transcripts. The correlation cutoff value
was |0,9|. Edges in red correspond to positive correlations. Edges in blue correspond to
negative correlations. Red nodes are the hub TFs. Blue nodes are transcripts that always
were significantly expressed in RDV. Some nodes were annotated with significant GO
terms: turquoise nodes with GO:0022622 (Root system development); yellow nodes with
GO:0010467 (Gene expression); orange nodes with GO:0042254 (Ribosome biogenesis);
and pink nodes with GO:0010467 (Gene expression) and GO:0042254 (Ribosome
biogenesis). (**B**) Functions of the correlated transcripts with hub TFs. Significant GO terms
from Biological Processes category after a SEA analysis with a 0.4 cut-off value for
dispensability.
Figure S7. Salicylic acid and jasmonic acid localization in root tip sections. The
phytohormones were localized in root apex cuts of control and 3 mM NH_4^+ treated
seedlings. The phytohormone identification and imaging was made using nano-particle
assisted laser desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). (**A**)
Salicylic acid (SA); (**B**) jasmonic acid (JA). Size bars correspond to 500 µm. Column

graphs show relative hormone accumulations respect to control samples in the approximate

LCM areas. Dashed lines mark hormone accumulation in the control roots.

Figure S8. Comparison between LCM and whole root transcriptomics data obtained from

ammonium supply experiment with pine seedlings. DE transcripts were obtained from

Ortigosa et al., 2021 (doi: 10.1101/2021.04.20.440618). (**A**) Expression of marker genes

for whole root response to NH_4^+ supply in the tissues isolated through LCM. (**B**) Venn

diagram with DE transcripts shared between LCM and whole root analyses. (**C**) LogFC

data for shared DE transcripts between LCM and whole root approaches including the

antimicrobial peptide 1 (*PpAMP1*) and its identifiers at LCM dataset (pp_58007) and

whole root (pp_58005).

Figure S9. Expression atlas of *PpDof1* in the tissues of one-month old seedlings (Cañas et

al., 2017).

Figure S10. Expression atlas of *PpDof2* in the tissues of one-month old seedlings (Cañas et

al., 2017).

Figure S11. Expression atlas of *PpDof11* in the tissues of one-month old seedlings (Cañas

et al., 2017).

Figure S12. Expression atlas of *PpDof12* in the tissues of one-month old seedlings (Cañas

et al., 2017).

Dataset S1. LCM low-input RNA-seq differential expression results. DE transcripts up-

regulated are in highlighted red. DE transcripts down-regulated are in highlighted blue.

Dataset S2. LCM low-input RNA-seq clustering results of DE transcripts.

Dataset S3. LCM low-input RNA-seq functional enrichment of DE transcripts. Only

conditions with significant functions are shown.

Dataset S4. Selected correlated transcripts with hub transcription factors for ammonium

treatment.

Dataset S5. Functional enrichment of correlated transcripts with hub transcription factors

for ammonium treatment.

Fig. 1. Experimental design. Timeline from germination to harvest. RT-qPCR data for the validation of the experimental design are included. *Glutamine synthetase 1b* (*PpGS1b*); *NADH-dependent glutamate synthase* (*PpNADH-GOGAT*); *antimicrobial peptide 1* (*PpAMP1*); *S-adenosyl methionine synthase* (*PpSAMS*) Significant differences were determined with a *t*-test (* *P*-value < 0.05; ** *P*-value < 0.01). Error bars show SE with n = 3. Isolated regions from the root apex: cap (RC), meristem (RM), developing cortex (RDC) and developing vessels (RDV).

Fig. 2. Low-input RNA-seq results in the different tissues of maritime pine root tip. (**A**) Venn diagram of down-regulated DE transcripts. (**B**) Venn diagram of up-regulated DE transcripts. (**C**) Heatmap and hierarchical clustering of DE transcripts. Expression values correspond to median CPM between biological/experimental replicates. The hierarchical clustering was made employing the Ward's minimum variance method. (**D**) Experimental validation of low-input RNA-seq results for 8 DE transcripts through RT-qPCR. RC, the root cap; RM, root meristem; RDC, root developing cortex; and RDV, root developing vessels. *SHORT-ROOT* (*PpSHR*); *nitrate transporter 1 / peptide transporter family 3.1* (*PpNPF3.1*); *cycloeucalenol cycloisomerase* (*PpCPI1*); *DOF transcription factor 12* (*PpDOF12*); *NAC transcription factor 38* (*PpNAC38*); *aminocyclopropane-1-carboxylic acid oxidase* (*PpACO*); *antimicrobial peptide 1* (*PpAMP1*); *BLISTER-like transcription factor* (*PpBLISTER-like*).

Fig. 3. Resume of the main functional results obtained in the low-input RNA-Seq analysis from LCM isolated samples. Functions and DE transcripts in (**A**) the root cap (RC), (**B**) root meristem (RM), (**C**) root developing cortex (RDC) and (**D**) root developing vessels (RDV). Blue arrows indicate downregulated transcripts, red arrows upregulated ones. (**E**) Significant GO terms from Biological Processes category after a SEA analysis. (**F**) Significant GO terms from Molecular Function category after a SEA analysis.

Fig. 4. Expression heatmaps of differentially expressed transcriptional regulators and hormone related transcripts. (**A**) Transcription factors and other putative regulators. (**B**) Transcripts involved in hormone metabolisms, transport and signaling. CPM expression values were normalized by transcript.

Fig. 5. Phytohormone localization in root apex sections. The main phytohormones were localized in root apex cuts of control and 3 mM NH_4^+ treated seedlings. The phytohormone identification and imaging was made using nano-particle assisted laser desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). (**A**) Auxin / Indole-3-acetic acid (IAA); (**B**) cytokinin (CK); (**C**) aminocyclopropane-1-carboxylic acid (ACC); (**B**) abscisic acid (ABA). Size bars correspond to 500 μm. Column graphs show relative hormone accumulations respect to control samples in the approximate LCM areas. Dashed lines mark hormone accumulation in the control roots.

Fig. 6. Root growth and architecture parameters. Pine seedlings were cultivated for 74 days and irrigated once a week with 50 mL of water, 3 mM KNO_3 and 3 mM NH_4Cl, respectively. Six different parameters were measured after harvesting: (**A**) weight of whole roots; (**B**) weight of primary root; (**C**) primary root length; (**D**) number of lateral roots; (**E**) weight of lateral roots; (**F**) lateral root density. Significant differences were determined with a one-way ANOVA. Letters above the conditions show significant differences based on a Turkey post-hoc test ($P < 0.05$) with a n = 30.

Fig. 7. Schematic representation of the transcriptional response induced by NH_4^+ supply in maritime pine seedling roots 24 h after irrigation. Transcripts in red letters are upregulated. Transcripts in blue letters are downregulated. The root tip tissues are the root cap (RC) root meristem (RM) and root developing cortex (RDC). Results for root developing vessels (RDV) are not shown since they were very limited.