Gambling environment exposure increases temporal discounting but improves model-based control in regular slot-machine gamblers.

Ben Wagner [1,2], David Mathar [1] & Jan Peters [1]

[1]Department of Psychology, Biological Psychology, University of Cologne, Cologne, Germany

[2]Faculty of Psychology, Chair of Neuroimaging, Technical University of Dresden, Dresden, Germany

Correspondence

Ben Wagner (ben_jonathan.wagner@tu-dresden.de), Jan Peters (jan.peters@uni-koeln.de)

Funding

This work was funded by Deutsche Forschungsgemeinschaft (grant PE1627/5-1 to J.P.).

Abstract

Gambling disorder is a behavioral addiction that negatively impacts personal finances, work, relationships and mental health. In this pre-registered study (https://osf.io/5ptz9/?view_only=62956a2afcd1495984db4be18c54b50a) we investigated the impact of real-life gambling environments on two computational markers of addiction, temporal discounting and model-based reinforcement learning. Gambling disorder is associated with increased temporal discounting and reduced model-based learning. Regular gamblers (n = 30, DSM-5 score range 3-9) performed both tasks in a neutral (café) and a gambling-related environment (slot-machine venue) in counterbalanced order. Data were modeled using drift diffusion models for temporal discounting and reinforcement learning via hierarchical Bayesian estimation. Replicating previous findings, gamblers discounted rewards more steeply in the gambling-related context. This effect was positively correlated with gambling related cognitive distortions (pre-registered analysis). In contrast to our pre-registered hypothesis, model-based reinforcement learning was improved in the gambling context. Here we show that temporal discounting and model-based reinforcement learning are modulated in opposite ways by real-life gambling cue exposure. Results challenge aspects of habit theories of addiction, and reveal that laboratory-based computational markers of psychopathology are under substantial contextual control.

Introduction

Gambling disorder is a behavioral addiction that can have detrimental effects on quality of life including personal finances, work, relationships and overall mental health (Blaszczynski & Nower, 2002; Muggleton et al., 2021). Despite these negative consequences, many gamblers are motivated to continue to play, and praise the temporary excitement and pleasure (Fauth-Bühler et al., 2017). Accumulating evidence suggests similarities of gambling disorder and substance-use-disorders both on behavioral, cognitive and neural levels (Balodis & Potenza, 2020; Leeman & Potenza, 2012; Lobo et al., 2015; N. M. Petry, 2010; Singer et al., 2020). In light of these similarities, the fifth edition of the "Diagnostic and Statistical Manual of Mental Disorders" categorizes gambling disorder in the category of "Substance-related and Addictive Disorders" (Association, 2013). In contrast to substance-use-disorders, differences in behavioral and/or neural effects between gamblers and controls are unlikely to be confounded by chronic or acute drug effects (Clark et al., 2019; Peters & Büchel, 2011; Wiehler & Peters, 2015) . Gambling disorder has thus been termed a "pure addiction" (Mark Dixon, Ghezzi, et al., 2006).

Recently, categorical definitions of mental illness have increasingly been called into question. The National Institute for Mental Health of the United States proposed the Research Domain Criteria (RDoC) to foster characterization of the dimensions underlying psychiatric disorders. According to this approach, research in cognitive science should focus on the identification of continuous neuro-cognitive dimensions that might go awry in disease, i.e. trans-diagnostic markers (Nelson et al., 2016). Here we focus on two promising candidates for such trans-diagnostic processes that are affected across a range of psychiatric conditions, including gambling disorder: temporal discounting, i.e. the devaluation of delayed rewards (Bickel et al., 2019; Lempert et al., 2019; Peters & Büchel, 2011), and model-based (MB) control during reinforcement learning (Daw et al., 2011). MB control refers to computationally more expensive goal-directed strategies that utilize models of the environment, contrasting with model-free (MF) control that operates on stimulus-response associations (Balleine & O'Doherty, 2010; Daw et al., 2011; Doll et al., 2012; Valerie Voon et al., 2017).

Steep discounting has been consistently observed in substance use disorders and gambling disorder (Bickel et al., 2012; Bickel et al., 2019; MacKillop et al., 2011; Reynolds, 2006). Moreover, alterations in temporal discounting occur in a range of other disorders, including depression, bipolar disorder, schizophrenia and borderline personality disorder (Amlung et al., 2019), underlining the trans-diagnostic nature of this process. Changes in the contributions of MF and MB control have likewise been reported across multiple disorders,

including gambling disorder (Wyckmans et al., 2019), schizophrenia (Culbreth et al., 2016),

obsessive compulsive disorder (Gillan et al., 2020) and substance use disorders (Sebold et al.,

2014). Reduced MB control is also reflected in sub-clinical psychiatric symptom severity

(Gillan et al., 2016).

Addiction is known to be under substantial contextual control. Addiction-related cues

and environments are powerful triggers of subjective craving, drug use and relapse. Incentive

sensitization theory (T. Robinson & Berridge, 1993; Terry E. Robinson & Berridge, 2008)

provides a theoretical framework that links such effects to a highly sensitized dopamine system

that responds to drugs and addiction-related cues. Increased responses of the dopamine system

to addiction-related cues ("cue-reactivity") has been consistently observed in neuroimaging

studies of human addicts (Courtney et al., 2016; Moeller & Paulus, 2018), and there is evidence

that trans-diagnostic behavioral traits are likewise under contextual control. For example,

regular gamblers discount delayed rewards substantially more steeply when tested in a

gambling-related environment as compared to a neutral environment (Mark. Dixon, Jacobs, &

Sanders, 2006). Similar effects have been observed in laboratory tasks that include gambling-

related cues (Dale et al., 2019; Genauck et al., 2020; Miedl et al., 2014) but whether other

putative trans-diagnostic traits such as MB control are under similar contextual control is

unclear. Beyond, it is unclear whether gambling severity or maladaptive control beliefs (Raylu

& Oei, 2004) modulate such effects.

Though rarely examined in naturalistic settings, contextual effects on trans-diagnostic

dimensions of decision-making are of substantial clinical and scientific interest. Settings with

high ecological validity might provide more informative insights into the central drivers of

maladaptive behavior than laboratory-based studies (Anderson & Brown, 1984). If such trans-

diagnostic traits are further exacerbated in e.g. addiction-related environments, this could

constitute a mechanism underlying the maintenance and/or escalation of maladaptive behavior.

Second, traits such as temporal discounting can be modulated (Bickel et al., 2011; Bickel et al.,

2019; Lempert & Phelps, 2016) and could thus serve as a potential treatment target (Lempert

et al. 2019).

The present pre-registered study thus had the following aims. First, we aimed to

replicate the findings by Dixon et al. (Mark Dixon, Ghezzi, et al., 2006), who observed

increased temporal discounting in gambling-related environments in regular gamblers,

compared to neutral environments. Second, we extended their approach by including a modified

version of the prominent 2-step sequential decision task (Daw et al., 2011) to test whether

model-based control of behavior is likewise under contextual control. Reduced model-based

control has been linked to a range of psychiatric conditions (see above) including gambling

disorder (Wyckmans et al., 2019). Third, we directly tested for associations of contextual effects

with gambling severity and working memory capacity. Finally, our tasks allowed for

comprehensive computational modelling of choices and response time (RT) distributions.

Analyses of reinforcement learning and decision-making have recently been shown to

substantially benefit from an incorporation of RTs (Fontanesi et al., 2019; Pedersen et al., 2017;

Peters & D'Esposito, 2020; Shahar et al., 2019; Wagner et al., 2020) via the application of

sequential sampling models such as the drift diffusion model (DDM) (Forstmann et al., 2016).

Such analyses yield additional insights into the latent processes underlying decision-making

(Wagner et al., 2020) and can improve parameter stability (Shahar et al., 2019). To account for

these recent developments, we complemented our pre-registered analyses with additional

analyses of temporal discounting and reinforcement learning drift diffusion models (RLDDM).

Methods

Preregistration

This   study   was   preregistered   via   the   open   science   framework

(https://osf.io/5ptz9/?view_only=62956a2afcd1495984db4be18c54b50a). We deviated from

the pre-registered study design in the following ways. First, it was initially planned to use a lab-

setting for the neutral (non-gambling) testing environment. However, this was changed

following pre-registration to a café, which we felt was more similar to the gambling

environment in terms of the presence of social cues and the overall level of distraction. Second,

we initially aimed to include gamblers fulfilling at least one DSM-5 criterion for gambling

disorder. This was adjusted to a stricter inclusion criterion of at least three DSM-5 criteria. Both

of these changes were implemented before testing began.

To further account for recent developments in computational modelling we also

complemented the pre-registered computational modeling analyses with additional analyses of

RT distributions via temporal discounting and reinforcement learning DDMs (Fontanesi et al.,

2019; Pedersen et al., 2017; Peters & D'Esposito, 2020; Wagner et al., 2020). As a model-free

measure of intertemporal choice we used choice proportions of larger-but-later values instead

of the area under the empirical discounting curve (AUC) (Myerson et al., 2001).

A-priori sample size was calculated based on results by Dixon et al. (2006) observed an

effect size of $d = .5$ for the effect of gambling environments on temporal discounting in regular

gamblers. Power analysis (Faul et al., 2007) yielded a minimum sample size of n = 26 with

alpha error probability of .05 and power of .80. We then pre-registered a target sample size of
n = 30.

Participants

Participants were recruited via advertisements posted online and in local gambling venues.
First, they were screened via a telephone interview to verify that they show evidence for
problematic gambling behavior, with a primary gambling mode of electronic slot machines.
Further inclusion criteria were age in the range of nineteen to fifty, no illegal drug use, and no
history of neuropsychiatric disorders, current medication or a history of cardiovascular disease.
The ethics committee of the University of Cologne Medical Center approved all study
procedures.

Forty-two participants were then invited to a first appointment, were they provided
written informed consent and completed a questionnaire assessment and a set of working
memory tasks (see section on *background screening* below). Five participants dropped out
during or after the first appointment. Four additional participants were excluded after the first
appointment because they fulfilled less than three DSM-V criteria for gambling disorder. Two
participants dropped out after the first experimental testing session, and one participant was
excluded because he fell asleep twice during one testing session. Due to technical problems, we
obtained complete datasets for thirty participants for the intertemporal choice task and twenty-
nine participants for the two-step task, with twenty-eight participants overlapping.

Overall procedure

Participants were invited to three appointments. At the first appointment (*baseline screening*;

see below) participants were invited to our lab and performed a questionnaire assessment and

four working memory tasks. Participants were randomly assigned to one of the two locations

(café vs. casino) on the first experimental appointment (pseudorandomized location [first

session neutral or gambling] and task-version; see section on tasks below). We label the café

environment as neutral because no gambling associated cues were present. In both locations,

the delay discounting task was completed first, followed by the 2-step task. Appointments were

made on an individual basis but spaced within 7+-2 days and around the same time of day +- 2

hours. The café environment was an ordinary café serving non-alcoholic drinks and snacks and

furnished with 10 tables and approximately 50 m^2 of size. Testing occurred while the café was

in business as usual and experimenter and participant sat at a table next to a wall to assure some

privacy. The café was usually moderately attended and testing occurred at the same spot for all

participants, with only a few exceptions when this seat was taken. The gambling environment

was a common slot-machine venue operated by a German gambling conglomerate. The

experimenter and participant were seated at a table placed next to a wall in sight of the electronic

gaming machines (EGMs). In total there were four EGMs in direct sight of the participant and

a total of ten in the room (hidden by eye protection walls). The density of gambling related cues

varied as a function of people playing at EGMs, background sounds e.g. sounds of winning or

money dropping were all depended on regularly customers. However, in nearly all cases other

people were playing EGMs in direct sight of the participants. The experimenter was granted

permission to conduct research in two local gambling venues. Two chairs and a table to use for

the experimental session were provided. In both locations, subjects were placed in such a way

that neither experimenter nor customers could view their screen. Both tasks ran on a 15inch

Laptop using the Psychopysics toolbox (Kleiner et al., 2007) running in Matlab (The

MathWorks ©).

Background screening

Participants filled out a battery of questionnaires regarding gambling related cognition (GRCS)

(Raylu & Oei, 2004) and symptom severity (DSM-5;KFG,SOGS) (Falkai, 2015; Lesieur &

Blume, 1987; J. Petry & Baulig, 1996), demographic evaluation and standard psychiatric

diagnostic tools (see Supplemental Tables S1 and S2).

We assessed working memory capacity using a set of four working memory paradigms.

First, in an Operation Span Task (Redick et al., 2012) subjects were required to memorize a

sequence of letters while being distracted by math-operations. Second, in a Listening Span Task

(adapted from the German version of the Reading Span Test developed by van den Noort et al.

(van den Noort et al., 2008) subjects were required to listen to a series of sentences and had to

recall the last word of each sentence. Last, subjects performed two different versions of a Digit

Span Task (forward/backward) that were adopted from the Wechsler Adult Intelligence Scale

(Wechsler, 2008). Here, participants listened to a series of numerical digits which they had to

recall as a series in regular or reverse order. All working memory scores were z-transformed

and averaged to obtain a single compound working memory score (z-score).

Temporal discounting task

Participants performed 140 trials of a temporal discounting task where on each trial they made

a choice between a smaller-but-sooner (SS) immediate reward, and a larger-but-later (LL)

reward delivered after a specific delay. SS and LL rewards were randomly displayed on the left

and right sides of the screen, and participants were free to make their choice at any time. While

SS rewards were held constant at 20€. LL rewards were computed as multiples of the SS reward

(task version 1: 1.05, 1.055, 1.15, 1.25, 1.35, 1.45, 1.55, 1.65, 1.85, 2.05, 2.25, 2.55, 2.85, 3.05,

3.45, 3.85; task version 2: 1.025, 1.08, 1.2, 1.20, 1.33, 1.47, 1.5, 1.70, 1.83, 2.07, 2.3, 2.5, 2.80,

3.10, 3.5, 3.80. Each LL reward from one version was then combined with each delay option

for this version (in days): (either: 1, 7, 13, 31, 58, 122, or v: 2, 6 15, 29, 62, 118) yielding 140

trials in total. The mean larger LL magnitude was the same across task versions and the order

was counterbalanced across subjects and session (neutral/gambling).

At the end of each session, one decision was randomly selected and paid out in the form

of a gift certificate for a large online store, either immediately (in the case of an SS choice) or

via email/text message after the respective delay (in the case of a LL choice).

2-step task

Participants performed a slightly modified version of the 2-step task, a sequential

reinforcement-learning (Daw et al., 2011). Based on more recent suggestions (Kool et al., 2016)

we modified the outcome stage by replacing the fluctuating reward probabilities (reward vs. no

reward) with fluctuating reward magnitudes (Gaussian random walks with reflecting

boundaries at 0 and 100, and standard deviation of 2.5). In total the task comprised 300 trials.

Each trial consisted of two successive stages: In the 1^{st} stage (S1), participants chose between

two fractals embedded in grey boxes. After taking an S1 action, participants transitioned to one

of two possible 2^{nd} stages (S2) with fixed transition probabilities of 70% and 30%. In S2,

participants chose between two new fractals each providing a reward outcome in points

(between 0-100) that fluctuated over time. To achieve optimal performance, participants had

to learn two aspects of the task. They had to learn the transition structure, that is, which S1

stimulus preferentially (70%) leads to which pair of S2 stimuli. Further, they had to infer the

fluctuating reward magnitudes associated with each S2 stimulus. .

In both versions, the tasks differed in the S1 and S2 stimuli, and in the fluctuating

rewards in S2. However both task versions reward walks were equal in variance and mean and

were presented in counterbalanced order per session (neutral/gambling). Participants were

instructed about the task structure and performed 40 practice trials (with different random walks

and symbols) at the first appointment (*Baseline screening*). Following task completion, points

(*0.25) were converted to € and participants could win a bonus of up to 4.50€ that was added

to the baseline reimbursement of 10€/h.

Computational modeling and Statistical Analysis

Temporal discounting model

We applied a single-parameter hyperbolic discounting model to describe how subjective value

changes as a function of LL reward height and delay (Mazur, 1987; Green and Myerson,

2004):

245
$$SV(LL_t) = \frac{A_t}{1 + \exp(k + s_k * I_t) * D_t} \quad (1)$$

Here, A_t is the reward height of the LL option on trial t, D_t is the LL delay in days on trial t and

I_t is an indicator variable that takes on a value of 1 for trials from the gambling context and 0

for trials from the neutral condition. The model has two free parameters: k is the hyperbolic

discounting rate (modeled in log-space) and s_k is a weighting parameter that models the degree

of change in discounting in the gambling compared with the neutral context condition.

Softmax action selection

Softmax action selection models choice probabilities as a sigmoid function of value differences

(Sutton and Barto, 1998):

256
$$P(LL)_t = \frac{\exp\left[(\beta + s_\beta * I_t) * SV(LL_t)\right]}{\exp\left[(\beta + s_\beta * I_t) * SV(SS_t)\right] + \exp\left[(\beta + s_\beta * I_t) * SV(LL_t)\right]} \quad (2)$$

Here, *SV* is the subjective value of the larger but later reward according to Eq. 1 and β is an

inverse temperature parameter, modeling choice stochasticity (for $\beta = 0$, choices are random

and as β increases, choices become more dependent on the option values). $SV(SS_t)$ was fixed at

at 20 and I_t is again the dummy-coded context regressor, and s_β models the context effect on β.

Temporal discounting drift diffusion models

To more comprehensively examine environmental effects on choice dynamics, we additionally

replaced softmax action selection with a series of drift diffusion model (DDM)-based choice

rules. In the DDM, choices arise from a noisy evidence accumulation process that terminates

as soon as the accumulated evidence exceeds one of two response boundaries. In the present

setting, the upper boundary was defined as selection of the LL option, whereas the lower

boundary was defined as selection of the SS option.

RTs for choices of the SS option were multiplied by -1 prior to model fitting. We

furthermore used a percentile-based cut-off, such that for each participant the fastest and

slowest 2.5 percent of trials were excluded from the analysis. We then first examined a null

model (DDM$_0$) without any value modulation. Here, the RT on each trial t ($t \in 1{:}140$) is

distributed according to the Wiener First Passage Time (*wfpt*):

276
$$RT_t \sim wfpt(\alpha + s_\alpha * I_t, \tau + s_\tau * I_t, z + s_z * I_t, v + s_v * I_t) \quad (3)$$

The parameter α models the boundary separation (i.e. the amount of evidence required before

committing to a decision), τ models the non-decision time (i.e., components of the RT related

to motor preparation and stimulus processing), z models the starting point of the evidence

accumulation process (i.e., a bias towards one of the response boundaries, with $z>.5$ reflecting

a bias towards the LL boundary, and $z<.5$ reflecting a bias towards the SS boundary) and v

models the rate of evidence accumulation. Note that for each parameter x, we also include a

parameter s_x that models the change in that parameter from the neutral context to the gambling

context (coded via the dummy-coded condition regressor I_t).

As in previous work (Pedersen et al., 2017; Fontanesi et al., 2019; Peters and D'Esposito, 2020,

Wagner et al. 2020), we then set up temporal discounting diffusion models with modulation of

drift rates by the difference in subjective values between choice options. First, we set up a

version with linear modulation of drift-rates (DDM$_{lin}$) (Pedersen et al., 2017):

290
$$v_t = \left(v_{coeff} + s_{v_{coeff}} * I_t \right) * \left(SV(LL_t) - SV(SS_t) \right) \quad (4)$$

Here, the drift rate on trial t is calculated as the scaled value difference between the subjective

LL and SS rewards. As noted above, RTs for SS options were multiplied by -1 prior to model

estimation, such that this formulation predicts SS choices whenever SV(SS)>SV(LL) (the trial-

wise drift rate is negative), and predicts longest RTs for trials with the highest decision-conflict

(i.e., in the case of SV(SS)= SV(LL) the trial-wise drift rate is zero). We next examined a DDM

with non-linear trial-wise drift rate scaling (DDM$_S$) that has recently been reported to account

for the value-dependency of RTs better than the DDM$_{lin}$ (Fontanesi et al., 2019; Peters &

D'Esposito, 2020; Wagner et al., 2020). In this model, the scaled value difference from Eq. 4 is

additionally passed through a sigmoid function with asymptote v_{max}:

301
$$v_t = S \left[\left(v_{coeff} + s_{v_{coeff}} * I_t \right) * \left(SV(LL_t) - SV(SS_t) \right) \right] \quad (5)$$

303
$$S(m) = \frac{2 * \left(v_{max} + s_{v_{max}} * I_t \right)}{1 + \exp(-m)} - \left(v_{max} + s_{v_{max}} * I_t \right) \quad (6)$$

All parameters including v_{coeff} and v_{max} were again allowed to vary according to the context,

such that we included s_x parameters for each parameter x that were multiplied with the dummy-

coded condition predictor I_t.

Reinforcement Learning model

Hybrid model

We first applied a slightly modified version of the hybrid RL model (Daw et al., 2011) to

analyze the strength of model-free and model-based RL strategies. The model updates MF state-

action values (Q_{MF}-values, Eq. 7, 8) in both stages through prediction errors (Eq. 9, 10). In

stage 1, MB state-action values (Q_{MB}) are then computed from the transition and reward

estimates using the Bellman Equation (Eq. 11).

$$Q_{MF,S1}(a_{j,t}) = Q_{MF,S1}(a_{j,t}) + (\eta_1 + s_{\eta 1} * I_t)\delta_{s1,t} + (\eta_2 + s_{\eta 2} * I_t)\delta_{S2,t} \quad (7)$$

$$Q_{MF,S2}(s_{2i,t}, a_{j,t}) = Q_{MF,S2}(s_{2i,t}, a_{j,t}) + (\eta_2 + s_{\eta 2} * I_t)\delta_{S2,t} \quad (8)$$

$$\delta_{S1,t} = Q_{MF,S2}(s_{2i,t}, a_{j,t}) - Q_{MF,S1}(a_{j,t-1}) \quad (9)$$

$$\delta_{S2,t} = r_{2\,t} - Q_{MF,S2}(s_{2i,t-1}, a_{j,t-1}) \quad (10)$$

$$Q_{MB}(a_{j,t}) = P(s_{21}|s_1, a_j) * \max_{a \in \{a_1, a_2\}} Q_{MF,S2}(s_{21}, a) + P(s_{22}|s_1, a_j) * \max_{a \in \{a_1, a_2\}} Q_{MF,S2}(s_{22}, a) \quad (11)$$

Here, i indexes the two different second stages (S_{21}, S_{22}), j indexes actions a (a_1, a_2) and t

indexes the trials. Further, η_1 and η_2 denote the learning rate for S1 and S2, respectively. S2 MF

Q-values are updated by means of reward $(r_{2,t})$ prediction errors ($\delta_{S2,t}$) (Eq. 8, 10). To model

S1 MF Q-values we allow for reward prediction errors at the 2nd-stage to influence 1st-stage

Q-values (Eq. 7, 9).

In addition, as proposed by Toyama et al. (Toyama et al., 2017, 2019) Q-values of all unchosen

stimuli were assumed to decay with decay-rate η_{decay} and centered to the mean of reward walks

(0.5). A decay of Q-values over time accounts for the fact that participants know that reward

walks fluctuate over time. The decay was implemented according to Eq. 12 and 13:

$$Q_{unchosen}(s_{k,t}, a_{j,t}) = Q_{unchosen}(s_{k,t-1}, a_{j,t-1}) * (\eta decay_S) + (1 - (\eta decay_S) * 0.5 \quad (12)$$

where

$$\eta decay_S = \eta_{decay} + s_{\eta decay} * I_t \quad (13)$$

and $k \in \{1, 21, 22\}$, that is, k indexes the three task stages.

S1 action selection is then modelled via weighting S1 MF and MB Q-values through a softmax

action-selection. S2 stage action selection is likewise modelled as a function of MF Q-value

differences. Separate 'inverse temperature' parameters β model subjects' weights of MF and

MB Q-Values (Eq. 14 and Eq. 15). The additional parameter ρ captures 1st-stage choice

perseveration, and is set to 1 if the previous S1 choice was the same and is zero otherwise.

346
$$p\left(a_{j,t} = a \middle| s_{1,t}\right) = \frac{\exp\left(\beta_{MB_s}*Q_{MB}(a) + \beta_{MF_s}*Q_{MF,S1}(a) + \rho_s*rep(a)\right)}{\sum_{a'} exp(\beta_{MB_s}*Q_{MB}(a') + \beta_{MF_s}*Q_{MF,S1}(a') + \rho_s*rep(a'))}, \quad (14)$$

347
$$p\left(a_{j,t} = a \middle| s_{2,t}\right) = \frac{\exp\left(\beta_{2_s}*Q_{MF,S2}(a)\right)}{\sum_{a'} \exp(\beta_{2_s}*Q_{MF,S2}(a'))}, \quad (15)$$

where:

$\beta_{MB_s} = \beta_{MB} + s_{\beta_{MB}} * I_t$

$\beta_{MF_s} = \beta_{MF} + s_{\beta_{MF}} * I_t$

$\rho_s = \rho + s_\rho * I_t$

$\beta_{2_s} = \beta_2 + s_{\beta_2} * I_t$

Hybrid model with drift diffusion action selection

As in our analysis of temporal discounting we replaced softmax action selection with a DDM

choice rule (Shahar et al., 2019), leaving the reinforcement learning equations unchanged. For

each stage of the task, the upper boundary was defined as selection of one stimulus, whereas

the lower boundary was defined as selection of the other stimulus. We modelled each stage of

the task using separate non-decision time (τ), boundary separation (α) and drift- rate (v)

parameters. The bias (z) was fixed to 0.5. All parameters including *vcoeff*$_{MF}$, *vcoeff*$_{MB}$ and *v*$_{max}$

were again allowed to vary according to the context, such that we included s_x parameters for

each parameter *x* that were multiplied with the dummy-coded condition predictor I_t (see above).

Data were filtered using a percentile-based cut-off, such that for each participant the fastest and

slowest 2.5 percent of RTs/trials were excluded from further analysis. In addition, an absolute

cutoff of > 150ms was applied. We then first examined a null model (DDM$_0$; Eq. 3) without

any value modulation followed by two value-informed models where the drift-rate (v) is a linear

(Eq. 16 and 17) or sigmoid (Eq. 18) function of MF and MB Q-value weights. For the linear

version, the drift rate in S1 is

370
$$v_{S1,t} = vcoeff_{MB_s} * (Q_{MB[2]} - Q_{MB[1]}) + vcoeff_{MF_s} * (Q_{MF[2]} - Q_{MF[1]}) + p_s * rep(a') \quad (16)$$

372 and the drift rate in S2 is calculated as

$$v_{S2,t} = vcoeff_{S2} * (Q_{MF_{S2[2]}} - Q_{MF_{S2[1]}}) \ (17)$$

376 For the non-linear version, the linear drift rate from equations 16 and 17 are additionally passed

377 through a sigmoid:

$$v_{Si,t} = \frac{2 * v_{max_{Si_S}}}{1 + \exp(-m)} - v_{max_{Si_S}} \ (18)$$

380 where

$$vcoeff_{MB_S} = vcoeff_{MB} + s_{v_{MB}} * I_t$$
$$vcoeff_{MF_S} = vcoeff_{MF} + s_{v_{MF}} * I_t$$
$$vcoeff_{S2_S} = vcoeff_{S2} + s_{S2} * I_t$$
$$vmax_{Si_S} = vmax_{Si} + s_{Si} * I_t$$

Hierarchical Bayesian models

Softmax models were fit to all trials from all participants using a hierarchical Bayesian

modeling approach with separate group-level distributions for all baseline parameters for the

neutral context and shift parameters (*s_x*) for the gambling context.

For the intertemporal choice model fitting was performed using Markov Chain Monte

Carlo (MCMC) sampling as implemented in the JAGS (Version 4.3) software package (Martyn

Plummer, 2003) in combination with the Wiener module (Wabersich and Vandekerckhove,

2014). Model estimation was done in R (Version 4.0.3) using the corresponding R2Jags

package (Version 0.6-1). For baseline group-level means, we used uniform priors defined over

numerically plausible parameter ranges (see code and data availability section for details). For

all *s_x* parameters modeling context effects on model parameters, we used Gaussian priors with

means of 0. For group-level precisions, we used gamma distributed priors (.001, .001). We

initially ran 2 chains with a varying burn-in period and thinning of two until convergence. Chain

convergence was then assessed via the Gelman-Rubinstein convergence diagnostic $\hat{R}$ and

sampling was continued until $1 \le \hat{R} \le 1.02$ for all group-level and individual-subject

parameters. 20k additional samples were then retained for further analysis.

For the 2-step task, model estimation was performed using MCMC sampling as

implemented in STAN (Stan Development Team, 2020) via R (Version 4.0.3) and the rSTAN

package (Version 2.21.0).

For baseline group-level means, we used uniform and normal priors defined over numerically

plausible parameter ranges (see code and data availability section for details). For all s_x

parameters modeling context effects on model parameters, we used Gaussian priors with means

of 0. For group-level standard deviations we used cauchy (0, 2.5) distributed priors. We initially

ran 2 chains with a burn-in period of 1000 and retained 2000 samples for further analysis. Chain

convergence was then assessed via the Gelman-Rubinstein convergence diagnostic $\hat{R}$ and

sampling was continued until $1 \leq \hat{R} \leq 1.02$. This threshold was not met for one participant ($\hat{R}$

< 1.4).

For both tasks, relative model comparison was performed via the *loo*-package in R

(Version 2.4.1) using the Widely-Applicable Information Criterion (WAIC) where lower values

reflect a superior fit of the model (Vehtari et al., 2017). We then show posterior group

distributions for all parameters of interest as well as their 85% and 95% highest density

intervals. For group comparisons we report Bayes Factors for directional effects for s_x

hyperparameter distributions of $s_x > 0$ (gambling context > neutral context), estimated via kernel

density estimation using R via the RStudio (Version 1.3) interface. These are computed as the

ratio of the integral of the posterior difference distribution from 0 to $+\infty$ vs. the integral from 0

to $-\infty$. Using common criteria (Beard et al. 2016), we considered Bayes Factors between 1 and

3 as anecdotal evidence, Bayes Factors above 3 as moderate evidence and Bayes Factors above

10 as strong evidence. Bayes Factors above 30 and 100 were considered as very strong and

extreme evidence respectively, whereas the inverse of these reflect evidence in favor of the

opposite hypothesis.

Posterior Predictive checks

We carried out posterior predictive checks to examine whether models reproduced key patterns in the data, in particular the value-dependency of RTs (Peters & D'Esposito, 2020; Wagner et al., 2020) and participant's choices. For the intertemporal choice task, we binned trials of each individual participant into five bins, according to the absolute difference in subjective larger-later vs. smaller-sooner value ("decision conflict", computed according to each participant's median posterior log(k) parameter from the DDM_S, and separately for the neutral and gambling context. For each participant and context, we then plotted the mean observed RTs as a function of decision conflict, as well as the mean RTs across 10k data sets simulated from the posterior distributions of the DDM_0, DDM_{lin} and DDM_S. For the 2-step task, we extracted mean posterior parameter estimates and simulated 1000 datasets in R (Version 4.0.3) using the Rwiener package (Version 1.3.3). We then show RTs as a function of S2 reward difference of observed data and the mean RTs across 1000 simulated datasets for of all DDMs. We further show that our models capture the relationship of S2 reward differences and optimal (max[reward]) choices.

Model free analysis

As a model-agnostic measure of temporal discounting, we examined arcsine-square-root transformed proportions of LL choices as a function context (neutral vs. gambling) with order (neutral vs. gambling session first) as fixed and subject as random effect using a hierarchical generalized linear model (HGLM). For the 2-step task we likewise use a HGLM approach and modeled 2nd-stage RTs as a function of transition (common vs. rare) and context (neutral vs. gambling) as fixed and subject as random effect. In line with our modelling analyses, data were filtered so that implausibly fast RTs were excluded (see Methods). A standard analysis of stay probabilities (Daw et al., 2011) adapted to our task version is reported in the Supplement (Supplemental Table S5).

Subjective Craving Rating

On each testing day, participants rated their subjective craving ("How much do you desire to gamble right now?") on a visual-analogue scale ranging from 0 to 100, both at the beginning of the testing session, and at the end following task completion. We then used paired t-tests to examine whether subjective craving differed between the testing environments (neutral vs. gambling).

Results

Subjective craving

Craving was assessed on a visual-analogue-scale before and after task performance. Due to

technical problems, ratings of the first eight participants were lost. Another two participants did

not complete post-task ratings. In the remaining n=22 participants, craving was substantially

higher in the gambling-related environment compared to the neutral environment (paired t-test

pre-task: t_{23} = -3.13; p = 0.0048, Cohen's *d*: 0.75; post-task: t_{21} = -4.32, p = 0.0003, Cohen's *d*

= 0.68; see Figure 1).

Figure 1. Subjective craving was assessed at the beginning (A) and at the end (B) of each
testing session via a visual-analogue scale rating. Craving was significantly higher in the
gambling environment, both at the start of the session (p = 0.0048) and at the end of the session
(p = 0.0003).

Temporal discounting

Model-agnostic analysis temporal discounting task

Raw proportions of larger-but-later (LL) choices are plotted in Figure 2A for each context. A

hierarchical linear model on arcsine-square-root transformed proportion values with context

(gambling vs. neutral) and order (gambling first vs. neutral first) as fixed effects and subject as

random effect confirmed a significant main effect of context (F_{28} = 13.33, p = 0.01) such that

participants made more LL selections in the neutral vs. the gambling-related environment.

There was no effect of order on choice proportions. Overall response time (RT) distributions

are plotted in Figure 2B with choices of the LL option coded as positive RTs and choices of the

smaller-sooner option coded as negative RTs.

Figure 2. Behavioral data from the temporal discounting task. A: raw proportions of larger-later (LL) choices in each context. B: Overall response time distributions with choices of the LL option coded as positive RTs and choices of the smaller-sooner option coded as negative RTs; Note, this was done to add choice coding to the computational model.

Softmax choice rule

We first modeled the data using standard softmax action selection. This analysis revealed an overall context effect on log(k), such that discounting was substantially steeper in the gambling context compared to the neutral context (Figure 3B, 95% HDI > 0). Examination of Bayes Factors indicated that an increase in log(k) in the gambling context (s_k) was about 116 times more likely than a decrease (see Figure 3 and Table 3). There was no evidence for a change in choice stochasticity (softmax[β]; Figure 3C/D).

Figure 3. Softmax model; Posterior distributions of mean hyperparameter distributions for the neutral baseline context (blue) and the corresponding shift in the gambling context (pink). A, discount-rate log (k); B, shift in discount-rate (s_k); C, softmax β; D, shift in softmax β; Thin (thick) horizontal line denote 95% (85%) highest posterior density intervals

Temporal discounting drift diffusion models (DDMs)

Model comparison of temporal discounting DDMs revealed the same model ranking in each context (Supplemental Table S3) such that the data were best accounted for by a temporal discounting DDM with non-linear drift rate scaling. This model accounted for around 90% of decisions (Supplemental Table S4, Supplemental Figure S1) and posterior predictive checks confirmed that it reproduced individual-participant RTs (Supplemental Figure S2).

We next examined the posterior distributions of model parameters of the best-fitting TD-DDM model (DDMs with sigmoid drift rate scaling; we further report model comparison, binary choice predictions and posterior predictive checks in the corresponding *Model comparison and validation* section in the supplement). Results are plotted in Figure 4 and Figure 5 and Bayes Factors for all context-effects are listed in Table 1. There was a consistent positive association between trial-wise drift rates and value differences in the neutral context (Figure 4E, the 95% HDI for the drift rate coefficient parameter did not include 0). Likewise, there was a numerical bias towards the smaller-sooner option in the baseline condition (85% HDI < 0.5, see Figure 4F). The non-decision time was numerically smaller in the gambling context (85 % HDI < 0, Figure 5B, Table 1), amounting to on average a 50ms faster non-

525   decision time. The maximum drift-rate was substantially higher in the gambling context (95%

526   HDI > 0, Figure 5D).

527   As in the softmax model (Figure 3), we observed a substantial increase in the discount

528   rates log(k) in the gambling context (95% HDI > 0, see Figure 5A, Table 1).

Figure 4. Temporal discounting drift diffusion model results: posterior distributions for hyperparameter means from the neutral context. A: discount-rate log(k), B: non-decision time τ, C: boundary separation α, D: maximum drift-rate v_{max}, E: drift-rate coefficient v_{coeff}, F: starting-point z. Thin (thick) horizontal line denote 95% (85%) highest posterior density intervals

Figure 5. Temporal discounting drift diffusion model results: posterior distributions for hyperparameter means for context shift (s_x) parameters modeling changes from the neutral to the gambling context. A: shift in discount-rate (s_k), B: shift in non-decision time s_τ, C: shift in boundary separation s_α, D: shift in maximum drift-rate v_{max}, E: shift in drift-rate coefficient v_{coeff}, F: shift in starting-point s_z. Thin (thick) horizontal line denote 95% (85%) highest posterior density intervals.

546 **Table 1**. Overview of overall context differences. For group comparisons we report Bayes
547 Factors for directional effects for s_x hyperparameter distributions of $s_x > 0$ (gambling context
548 > neutral context).

Model parameter (change in gambling context)	Softmax Model		DDM$_S$	
	Mean	*dBF*	*Mean*	*dBF*
s_k (discount-rate)	0.77	1688.53	0.40	54.20
s_β (softmax beta)	0.025	2.27	-	-
s_{vcoeff} (drift-rate coeff.)	-	-	-0.012	0.25
s_τ (non-decision time)	-	-	-0.05	0.10
s_α (boundary separation)	-	-	0.10	4.40
s_z (starting point bias)	-	-	0.02	13.64
s_{vmax} (max drift-rate)	-	-	0.33	39490.71

Temporal discounting and gambling-related questionnaire data

As preregistered, we next examined whether the increased in discount-rate s_k in the gambling
context was associated with symptom severity or gambling related cognition. We therefore
computed a compound symptom severity *z*-score of DSM-5 (Falkai, 2015), SOGS (Lesieur &
Blume, 1987) and KFG (J. Petry & Baulig, 1996) scores. Gambling context-related changes in
temporal discounting were not significantly associated with symptom severity (rho = -0.05, p
= 0.78) but were positively associated with Gambling Related Cognition Scale (Raylu & Oei,
2004) (rho = 0.39; p = 0.03); see Figure 6). There were no significant correlations between
changes in craving and changes in discounting or working memory capacity and temporal
discounting (Supplemental Results 1).

Figure 6. Relationship of total scores from the gambling-related cognition scale (GRCS) (Raylu & Oei, 2004) and changes in discount-rate from neutral to gambling environment (s_k)[softmax model].

2-step reinforcement learning task

Model-agnostic analysis 2-step task

Participants earned significantly more points in the gambling context (t-test: t_{28} = -2.44, *p* = 0.02, Cohen's *d* = 0.22). For S2 RTs, we observed a significant main effect of transition (Supplemental Table S7 and Supplemental Figure S3) and a trend for a transition x context interaction (p = 0.07; see Supplemental Table S7), reflecting increased model-based control (Otto et al., 2015; Shahar et al., 2019).

An analysis of stay probabilities adapted to the present 2-step task version is shown in Supplemental Table S5. In each context, we observed main effects of reward (reflecting model-free RL) and reward x transition interaction (reflecting model-based RL). The reward x transition x context interaction was not significant.

Hybrid model with softmax choice rule

We first examined a hybrid model (Daw et al., 2011) with extensions by ourselves and Otto et al. (Otto et al., 2015) using a standard softmax choice rule (see Methods; Figure 7). This model included separate parameters for S1 and S2 learning rates, model-free and model-based β weights for S1 and a β weight for S2 *Q*-value differences. We confirmed substantial contributions of both MB and MF values to S1 choices (Figure 7B,C). There was an increase in the S2 learning-rate η (95% HDI > 0, Figure 7F) in the gambling context. Furthermore, there was a strong decrease in MF β weights (95% HDI < 0, Figure 7H) such that participants showed substantially less MF behavior in the gambling environment compared to the neutral

environment. BFs for directional effects indicate that an increase in MB reinforcement learning
is 4 times more likely than a decrease. For examination of Bayes Factors see Table 2.

Figure 7. Hybrid model with softmax choice rule posterior distributions (top row: neutral
context, bottom row: parameter changes in gambling context) of all group level means. A, S1
and S2 learning-rates. B, MB β weight. C, MF β weight. D, S2 β weight. E, perseveration
parameter ρ. F, shift in S1 and S2 learning rates. G, shift in MB β. H, shift in MF β. I, shift in
S2 β. J, shift in stickiness parameter ρ. Thin (thick) horizontal line denote 95% (85%) highest
posterior density intervals.

Hybrid model with drift diffusion choice rule

We next combined the hybrid model with a DDM choice-rule (Shahar et al., 2019) and likewise

compared DDMs that varied in the way that they accounted for the influence of *Q*-value

differences on trial-wise drift rates in both task stages. Model comparison yielded the same

model ranking in each context, such that the data were best accounted for by an RLDDM with

non-linear drift rate scaling (Supplemental Table S8). This model accounted for around 73% of

S1 choices, and around 81% of S2 choices (Supplemental Table S9). Posterior predictive checks

confirmed that this model reproduced the observed RTs (Supplemental Figure S4) and choice

proportions (Supplemental Figure S5).

Posterior distributions for the best-fitting RLDDM are shown in Figure 8 (neutral context

parameters) and Figure 9 (gambling context changes). We observed positive associations

between trial-wise drift rates and *Q*-value differences (Figure 8F-J, all 95% HDIs above 0).

Likewise, as in the softmax model, beta weights were positive, indicating MB and MF effects

on behavior (Figure 8E-G, all 95% HDIs > 0). In the gambling context, we observed a decrease

in the MF component (85% HDI < 0) and a robust increase in MB contributions (95% HDI >

0). BFs for directional effects are provided in Table 2. Overall, these results suggest decreased

MF and increased MB reinforcement learning due to gambling context exposure.

Model parameter (shift)	Softmax Model		DDMs	
	Mean	*dBF*	*Mean*	*dBF*
$s_{\eta S1}$ (learning-rate S1)	0.44	3.29	0.0801	1.186
$s_{\eta S2}$ (learning-rate S2)	0.40	92.3	0.280	14.658
$s_{\tau S1}$ (non-decision times S1)	-	-	0.001	0.8454
$s_{\tau S2}$ (non-decision times S2)	-	-	0.001	1.161
s_{ρ} (Stickiness S1)	0.04	1.946	0.05	2.365
$s_{\alpha S1}$ (boundary separation S1)	-	-	-0.002	0.9354
$s_{\alpha S2}$ (boundary separation S2)	-	-	0.0149	2.026
β_{MF} / $S_{vcoeffMF}$ (MF beta/ drift-rate coeff.)	-1.14	0.010	-0.93	0.083
β_{MB} / $S_{vcoeffMB}$ (MB beta/ drift-rate coeff.)	1.08	4.00	4.01	169.62
β_{S2} / $S_{vcoeffS2}$ (S2 beta / drift-rate coeff.)	-0.44	0.428	-0.64	0.271
s_{vmaxS1} (max drift-rate S1)	-	-	-0.19	0.296
s_{vmaxS2} (max drift-rate S2)	-	-	0.41	15.83

Figure 8. RL-DDM. Posterior distributions of all hyperparameters for the neutral baseline
condition. A: S1 and S2 learning rates η. B: S1 and S2 non-decision time τ. C: S1 and S2
boundary separation α. D: S1 and S2 drift-rate maximum v_{max}. E: MF drift-rate coefficient
$vcoeff_{MF}$. F: MB drift-rate coefficient $vcoeff_{MB}$. G: S2 drift-rate coefficient $vcoeff_{S2}$. H:
stickiness parameter ρ. Thin (thick) horizontal line denote 95% (85%) highest posterior density
intervals.

Figure 9. RL-DDM. Posterior distributions of all hyperparameters shift-parameters modelling the change from neutral to gambling condition. A, shift in Stage 1 and Stage 2 learning rates η. B, shift in S1 and S2 non-decision time τ. C, shift in S1 and S2 boundary separation α. D, shift in S1 and S2 drift-rate maximum v_{max}. E, shift in S1 MF drift-rate coefficient $vcoeff_{MF}$. F, shift in S1 MB drift-rate coefficient $vcoeff_{MB}$. G, shift in S2 drift-rate coefficient $vcoeff_{S2}$. H, shift in stickiness parameter ρ. Thin (thick) horizontal line denote 95% (85%) highest posterior density intervals.

Reinforcement learning and gambling related questionnaire data

As preregistered, we examined associations between ρ (perseveration/stickyness) and gambling severity (average *z*-score across SOGS (Lesieur & Blume, 1987), KFG (J. Petry & Baulig, 1996) and DSM-5 criteria). The association was non-significant ρ (r = -0.10, p = 0.59). There were no significant correlations between changes in craving and changes in MB behavior, nor between MB behavior and working memory capacity (Supplemental Results 2).

Discussion

Here we comprehensively examined the contextual modulation of two putatively trans-diagnostic markers implicated in addiction, temporal discounting (Bickel et al., 2019; Lempert et al., 2019) and model-based control (Gillan et al., 2016; Gillan et al., 2020) in a pre-registered study. We studied regular slot machine gamblers, a group previously characterized by high levels of temporal discounting (Wiehler & Peters, 2015) and reduced model-based control (Wyckmans et al., 2019). Following a seminal study by Dixon et al. (Mark. Dixon, Jacobs, & Sanders, 2006), regular gamblers were tested in gambling environments (slot-machine venues) and neutral control environments. Gambling cue exposure modulated temporal discounting and model-based control in gamblers in opposite ways: replicating Dixon et al., (2006), discounting

substantially increased in a gambling context. In contrast, model-based (MB) control improved (increased). This differential modulation of two prominent trans-diagnostic traits in (behavioral) addiction has important theoretical and clinical implications.

Theoretical accounts highlight the central role of addiction-related cues and environments in drug addiction (T. Robinson & Berridge, 1993). Similar mechanisms have been suggested to underlie gambling disorder (M. J. F. Robinson et al., 2016). Because terrestrial slot machine gambling is directly linked to specific locations, gambling disorder is uniquely suited to investigate the impact of cue exposure on behavior. We replicated the finding of Dixon et al. (2006) of steeper discounting in gambling vs. neutral environments in gamblers. This effect was observed across model agnostic analyses (proportion of LL choices) and computational modeling (softmax, drift diffusion models [DDM]). We additionally extended these earlier results in the following ways. First, we observed an association of this effect with maladaptive control beliefs (GRCS) (Raylu & Oei, 2004) suggesting that such beliefs contribute to increased temporal discounting in gambling environments. Second, in a subset of participants, we confirmed that exposure to gambling environments substantially increases subjective craving. Third, comprehensive modeling via DDMs revealed additional effects on latent decision processes. The gambling context-related attenuation in non-decision time mirrors previous effects of pharmacological enhancement of dopamine transmission (Wagner et al. 2020). In contrast to these earlier pharmacological results, we observed a substantial *increase* in maximum drift rate (V_{max}) in the gambling context, reflecting increased value sensitivity of RTs. Lastly, our results complement cue-reactivity designs showing increased impulsive and/or risky choice in gamblers during exposure to gambling cues in laboratory studies (Dale et al., 2019; Genauck et al., 2020; Miedl et al., 2014). However, effect sizes during naturalistic cue exposure (e.g. the present study and Dixon et al., 2006) were substantially larger than during lab-based exposure in these previous studies.

In addition to temporal discounting, we included a 2-step sequential decision-making task designed to dissociate model-based (MB) from model-free (MF) contributions to behavior (Daw et al., 2011). Reductions in MB control are associated with compulsivity-related disorders (Gillan et al., 2016; Gillan et al., 2020; V. Voon et al., 2015b). We observed increased MB learning and reduced MF learning in gamblers in the gambling context, a pattern of results consistent between softmax and DDM models. These findings were again corroborated by model-agnostic analyses. First, participants earned more points in the gambling context, an effect linked to MB learning (Kool et al., 2016). Second, the slowing of RTs following rare transitions, an indirect measure for MB learning (Otto et al., 2015) tended to be more

pronounced in the gambling vs. neutral context. The MF effect correlated with gambling severity in an exploratory analysis, such that higher gambling severity was associated with a greater reduction in MF reinforcement learning in the gambling context. Together, these findings converge on the picture of *decreased* MF and *increased* MB control in gamblers when tested in gambling-related environments.

The latter result contrast with our pre-registered hypothesis of *reduced* MB control, which was based on findings of reduced MB control in populations with extensive habit formation (Gillan et al., 2016; Gillan et al., 2020; V. Voon et al., 2015a). Addiction is likewise thought to be inherently associated with pathological habits (Barry J Everitt & Trevor W Robbins, 2005; Robbins & Everitt, 1999) which are thought to be triggered by exposure to environmental cues (Antons et al., 2020). We thus hypothesized gambling environments would likewise trigger increased MF behavior and reduced MB behavior on the 2-step task. However, critics of habit theory have emphasized that addiction might in contrast be associated with excessive goal-directed behavior, in particular in the presence of addiction-related cues (Hogarth 2020). Our findings are more in line with this latter perspective. This interpretation is compatible with incentive sensitization theory (T. Robinson & Berridge, 1993; Terry E. Robinson & Berridge, 2008), which proposes that addiction-related environments exert their influence on behavior in part via a potentiation in dopamine release (Anselme & Robinson, 2013; Berridge, 2016; T. E. Robinson & Berridge, 2001). Earlier studies observed increased MB control following increases in DA neurotransmission (Sharp et al., 2016; Wunderlich et al., 2012), which could contribute to the present findings regarding 2-step task performance. Furthermore, our results are compatible with decreased MF control under L-Dopa (Kroemer et al., 2019). The gambling context might thus enhance goal-directed control via an improved construction and/or utilization of the task transition structure. This interpretation further resonates with other perspectives on DA function including a regulation of outcome sensitivity or precision (FitzGerald et al., 2015; Shiner et al., 2012), or the general motivation to exert (cognitive) effort (Berke, 2018). The observed increase in S2 learning rates could likewise be mediated in part by increases in DA transmission (Frank & O'Reilly, 2006).

If the effects of gambling environments on 2-step task performance are (at least in part) driven by increases in DA, then the question arises why gamblers at the same time exhibited substantially increased temporal discounting. The literature on DA effects on temporal discounting is a mixed bag (D'Amour-Horvat & Leyton, 2014) with some studies showing reduced discounting (van Gaalen et al., 2006; Wagner et al., 2020), some increased discounting (Pine et al., 2010) and others suggesting baseline-dependent effects (Petzold et al., 2019).

Given that DA was neither measured nor directly manipulated here, these issues cannot

be directly resolved. However, our data might nonetheless provide some insights. Effects of

DA on decision-making might depend on both task and context (Mikhael et al., 2021). Under

this view, DA signals average reward in the environment (context) and its effects on

performance further differ as a function of task controllability [see (Mikhael et al., 2021) for

details]. DA might thus facilitate cognitive control (Ott & Nieder, 2019; Westbrook et al., 2020)

when cognitive effort requirements are high, and there is control over the outcome (e.g. 2-step

task). In contrast, DA might facilitate impulsive choice for cognitively less demanding tasks

(e.g. temporal discounting task) that are performed in an addiction-related context (Antons et

al., 2020; Terry E. Robinson & Berridge, 2008) signaling high reward (Mikhael et al., 2021).

A further mechanism known to modulate temporal discounting is episodic future thinking or

future prospection (Gershman & Bhui, 2020; Peters & Büchel, 2010). Future prospection has

been shown to attenuate temporal discounting in a range of settings (Rösch et al., 2021) and

might be attenuated at gambling venues. Participants might be generally focused on the present

in the presence of cues or contexts endowed with high levels of incentive salience (Flagel et al.,

2009).

Our results show that two prominent (potentially trans-diagnostic) computational

processes, temporal discounting and MB control, are differentially modulated by addiction-

related environments in regular slot machine gamblers. This provides a computational

psychiatry perspective on factors that contribute to the understanding of this disorder. The

substantial contextual effects on temporal discounting further highlight the potential clinical

relevance of this process (Amlung et al., 2019; Lempert et al., 2019). Gambling disorder is

reliably associated with increased temporal discounting (Mark. Dixon et al., 2003; Mark.

Dixon, Jacobs, & Sanders, 2006; MacKillop et al., 2011; Miedl et al., 2012; Wiehler & Peters,

2015) . This trait-like behavior then appears to be further exacerbated during exposure to

gambling-related environments, potentially contributing to the maintenance of maladaptive

behavior. In contrast, MB control improved (increased) in a gambling context, despite the fact

that an earlier study reported reduced MB control in gamblers (Wyckmans et al., 2019). In

general these findings are further compatible with a greater tendency for pattern matching

(Wilke et al., 2014) or enhanced cause-effect associations that might translate into increased

MB control (Joukhador et al., 2004) and studies suggesting that DA increases the willingness

to spend cognitive effort (Westbrook et al., 2020; Westbrook et al., 2021). 2-step task transitions

are not random, but can be learned and exploited. An increased tendency to seek for patterns

during gambling context exposure might facilitate this behavior. Our findings suggest that

gamblers do generally show MB control, which contrasts in parts with one recent study
(Wyckmans et al., 2019). This is supported by the robust RTs increases observed following rare
transitions (Supplemental Table S7, Supplemental Figure S3) and the positive MB parameters
observed across models, somewhat contrasting with the findings of Wyckmans et al.
(Wyckmans et al., 2019), although different 2-step task versions have been used in these
studies.

We also extended previous studies on this topic via a recent class of value-based
decision models based on the DDM (Fontanesi et al., 2019; Pedersen et al., 2017; Peters &
D'Esposito, 2020; Shahar et al., 2019; Wagner et al., 2020). Comprehensive RT-based analysis
revealed that standard DDM parameters were largely unaffected by context, suggesting that
primarily MF and MB contributions to evidence accumulation were affected by gambling
environments (Figure 10.). Posterior predictive checks showed that a DDM with non-linear
trial-wise drift rate scaling captured the relationship of decision conflict (SS-LL value
difference) and RTs, replicating prior findings (Peters & D'Esposito, 2020; Wagner et al.,
2020). We previously reported good parameter recovery of such temporal discounting DDMs
(Peters & D'Esposito, 2020; Wagner et al., 2020).

A number of limitations need to be acknowledged. First, as in the original study (Mark.
Dixon, Jacobs, & Sanders, 2006) we did not test a non-gambling control group. However, the
observed associations between experimental effects and gambling severity / gambling-related
cognition suggests that these effects are at least in part driven by the underlying problem
gambling symptoms. Second, MB and MF effects in the 2-step task might be affected by
instructions (da Silva & Hare, 2019). Participants in our study were well instructed in written
and verbal form and completed extensive training trials. Moreover, due to the counter-balanced
within-subject design, the observed context-dependent changes in MB/MF behavior cannot be
attributed to overall instruction effects. Third, MB control might more generally be related to
attentional or motivational processes. For example, incentives boost 2-step task performance
(Patzelt et al., 2019). However, we ensured that mean and variance of reward walks as well as
incentives were identical in both contexts. Fourth, although participants were tested in the same
venues, the number of customers present varied across participants, affecting e.g. noise levels
and auditory gambling cues (slot machine sounds etc.). A trade-off between the control of such
variables and ecological validity is unavoidable when testing in naturalistic settings. Finally,
DA neurotransmission was obviously not assessed, rendering our interpretation of the effects
in terms of the incentive sensitization theory speculative. But the substantial increase in

subjective craving supports the idea that cue exposure had subjective effects predicted by

incentive sensitization.

To conclude, here we show that two computational trans-diagnostic markers with high

relevance for gambling disorder in particular and addiction more generally are modulated in

opposite ways by exposure to real gambling environments. Gamblers showed increased

temporal discounting in a gambling context, and this effect was modulated by maladaptive

control beliefs. In contrast, MB control improved, a finding that posits a challenge for

habit/compulsion theories of addiction. Ecologically valid testing settings such as those

investigated here can thus yield novel insights into environmental drivers of maladaptive

behavior underlying mental disorders.

Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft (PE1627/5-1 to JP)

Author contributions

JP conceived the idea and acquired the funding. JP and BJW designed the study. BJW acquired

the data. BJW analyzed the data and performed the modeling. DM contributed analytical

tools/software. BJW wrote the first draft of the paper. BJW and JP wrote the paper. DM

provided revisions. JP supervised the project.

References

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental*

disorders (5th ed.). Author.

Amlung, M., Marsden, E., Holshausen, K., Morris, V., Patel, H., Vedelago, L., Naish, K. R.,

Reed, D. D., & McCabe, R. E. (2019). Delay Discounting as a Transdiagnostic Process in

Psychiatric Disorders: A Meta-analysis. *JAMA Psychiatry*, *76*(11), 1176–1186.

https://doi.org/10.1001/jamapsychiatry.2019.2102

Anderson, G., & Brown, R. I. (1984). Real and laboratory gambling, sensation-seeking and

arousal. *British Journal of Psychology (London, England : 1953)*, *75 (Pt 3)*, 401–410.

https://doi.org/10.1111/j.2044-8295.1984.tb01910.x

Anselme, P., & Robinson, M. J. F [Mike J. F.] (2013). What motivates gambling behavior?

Insight into dopamine's role. *Frontiers in Behavioral Neuroscience*, *7*, 182.

https://doi.org/10.3389/fnbeh.2013.00182

Antons, S., Brand, M., & Potenza, M. N. (2020). Neurobiology of cue-reactivity, craving, and

inhibitory control in non-substance addictive behaviors. *Journal of the Neurological*

Sciences, *415*, 116952. https://doi.org/10.1016/j.jns.2020.116952

Association, A. P. (2013). *Diagnostic and Statistical Manual of Mental Disorders*. American

Psychiatric Association. https://doi.org/10.1176/appi.books.9780890425596

Balleine, B. W., & O'Doherty, J. P. (2010). Human and rodent homologies in action control:

Corticostriatal determinants of goal-directed and habitual action.

Neuropsychopharmacology : Official Publication of the American College of

Neuropsychopharmacology, *35*(1), 48–69. https://doi.org/10.1038/npp.2009.131

Balodis, I. M., & Potenza, M. N. (2020). Common neurobiological and psychological

underpinnings of gambling and substance-use disorders. *Progress in Neuro-*

Psychopharmacology & Biological Psychiatry, *99*, 109847.

https://doi.org/10.1016/j.pnpbp.2019.109847

Barry J Everitt, & Trevor W Robbins (2005). Neural systems of reinforcement for drug

addiction: From actions to habits to compulsion. *Nature Neuroscience*, *8*(11), 1481–1489.

https://doi.org/10.1038/nn1579

Beck, A., Steer, R., & Brown, G. (1996). Manual for the Beck Depression Inventory-II (BDI-

II).

Berke, J. D. (2018). What does dopamine mean? *Nature Neuroscience*, *21*(6), 787–793.

https://doi.org/10.1038/s41593-018-0152-y

Berridge, K. C [K. C.]. (2016). Incentive Motivation and Incentive Salience☆. In J. Stein

(Ed.), *Reference module in neuroscience and biobehavioral psychology*. Elsevier.

https://doi.org/10.1016/B978-0-12-809324-5.00342-4

Bickel, W. K., Athamneh, L. N., Basso, J. C., Mellis, A. M., DeHart, W. B., Craft, W. H., &

Pope, D. (2019). Excessive discounting of delayed reinforcers as a trans-disease process:

Update on the state of the science. *Current Opinion in Psychology*, *30*, 59–64.

https://doi.org/10.1016/j.copsyc.2019.01.005

Bickel, W. K., Jarmolowicz, D. P., Mueller, E. T., Koffarnus, M. N., & Gatchalian, K. M. (2012). Excessive discounting of delayed reinforcers as a trans-disease process contributing to addiction and other disease-related vulnerabilities: Emerging evidence. *Pharmacology & Therapeutics*, *134*(3), 287–297. https://doi.org/10.1016/j.pharmthera.2012.02.004

Bickel, W. K., Yi, R., Landes, R. D., Hill, P. F., & Baxter, C. (2011). Remember the future: Working memory training decreases delay discounting among stimulant addicts. *Biological Psychiatry*, *69*(3), 260–265. https://doi.org/10.1016/j.biopsych.2010.08.017

Blaszczynski, A., & Nower, L. (2002). A pathways model of problem and pathological gambling. *Addiction (Abingdon, England)*, *97*(5), 487–499. https://doi.org/10.1046/j.1360-0443.2002.00015.x

Clark, L., Boileau, I., & Zack, M. (2019). Neuroimaging of reward mechanisms in Gambling disorder: An integrative review. *Molecular Psychiatry*, *24*(5), 674–693. https://doi.org/10.1038/s41380-018-0230-2

Courtney, K. E., Schacht, J. P., Hutchison, K., Roche, D. J. O., & Ray, L. A. (2016). Neural substrates of cue reactivity: Association with treatment outcomes and relapse. *Addiction Biology*, *21*(1), 3–22. https://doi.org/10.1111/adb.12314

Culbreth, A. J., Westbrook, A [Andrew], Daw, N. D., Botvinick, M., & Barch, D. M. (2016). Reduced model-based decision-making in schizophrenia. *Journal of Abnormal Psychology*, *125*(6), 777–787. https://doi.org/10.1037/abn0000164

da Silva, C. F., & Hare, T. A. (2019). *Humans are primarily model-based and not model-free learners in the two-stage task* (Vol. 20). https://doi.org/10.1101/682922

Dale, G., Rock, A. J., & Clark, G. I. (2019). Cue-Reactive Imagery Mediates the Relationships of Reward Responsiveness with Both Cue-Reactive Urge to Gamble and

Positive Affect in Poker-Machine Gamblers. *Journal of Gambling Studies.* Advance online

publication. https://doi.org/10.1007/s10899-019-09864-x

D'Amour-Horvat, V., & Leyton, M. (2014). Impulsive actions and choices in laboratory

animals and humans: Effects of high vs. Low dopamine states produced by systemic

treatments given to neurologically intact subjects. *Frontiers in Behavioral Neuroscience*, *8*,

432. https://doi.org/10.3389/fnbeh.2014.00432

Daw, N., Gershman, S., Seymour, B., Dayan, P [Peter], & Dolan, R. J. (2011). Model-Based

Influences on Humans' Choices and Striatal Prediction Errors. *Neuron*, *69*(6), 1204–1215.

https://doi.org/10.1016/j.neuron.2011.02.027

Dixon, M [Mark], Ghezzi, P., Lyons, C., & Wilson, G. (2006). *Gambling: Behavior Theory,*

Research, and Application. New Harbinger Publications.

Dixon, M [Mark.], Jacobs, E. A., & Sanders, S. (2006). CONTEXTUAL CONTROL OF

DELAY DISCOUNTING BY PATHOLOGICAL GAMBLERS. *Journal of Applied*

Behavior Analysis, *39*(4), 413–422. https://doi.org/10.1901/jaba.2006.173-05

Dixon, M [Mark.], Marley, J., & Jacobs, E. A. (2003). Delay discounting by pathological

gamblers. *Journal of Applied Behavior Analysis*, *36*(4), 449–458.

https://doi.org/10.1901/jaba.2003.36-449

Doll, B. B., Simon, D. A., & Daw, N. D. (2012). The ubiquity of model-based reinforcement

learning. *Current Opinion in Neurobiology*, *22*(6), 1075–1081.

https://doi.org/10.1016/j.conb.2012.08.003

Falkai, P. (2015). *Diagnostisches und statistisches Manual psychischer Störungen: DSM-5*.

Hogrefe.

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*power 3: A flexible statistical

power analysis program for the social, behavioral, and biomedical sciences. *Behavior*

Research Methods, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Fauth-Bühler, M., Mann, K., & Potenza, M. N. (2017). Pathological gambling: A review of

the neurobiological evidence relevant for its classification as an addictive disorder.

Addiction Biology, *22*(4), 885–897. https://doi.org/10.1111/adb.12378

FitzGerald, T. H. B., Dolan, R. J., & Friston, K. (2015). Dopamine, reward learning, and

active inference. *Frontiers in Computational Neuroscience*, *9*, 136.

https://doi.org/10.3389/fncom.2015.00136

Flagel, S. B., Akil, H., & Robinson, T. E [Terry E.] (2009). Individual differences in the

attribution of incentive salience to reward-related cues: Implications for addiction.

Neuropharmacology, *56 Suppl 1*, 139–148.

https://doi.org/10.1016/j.neuropharm.2008.06.027

Fontanesi, L., Gluth, S., Spektor, M. S., & Rieskamp, J. (2019). A reinforcement learning

diffusion decision model for value-based decisions. *Psychonomic Bulletin & Review*,

26(4), 1099–1121. https://doi.org/10.3758/s13423-018-1554-2

Forstmann, B. U., Ratcliff, R., & Wagenmakers, E.-J. (2016). Sequential Sampling Models in

Cognitive Neuroscience: Advantages, Applications, and Extensions. *Annual Review of*

Psychology, *67*, 641–666. https://doi.org/10.1146/annurev-psych-122414-033645

Frank, M. J [Michael J.], & O'Reilly, R. C. (2006). A mechanistic account of striatal

dopamine function in human cognition: Psychopharmacological studies with cabergoline

and haloperidol. *Behavioral Neuroscience*, *120*(3), 497–517. https://doi.org/10.1037/0735-

7044.120.3.497

Genauck, A., Andrejevic, M., Brehm, K., Matthis, C., Heinz, A., Weinreich, A.,

Kathmann, N., & Romanczuk-Seiferth, N. (2020). Cue-induced effects on decision-making

distinguish subjects with gambling disorder from healthy controls. *Addiction Biology*,

25(6), e12841. https://doi.org/10.1111/adb.12841

Gershman, S. J., & Bhui, R. (2020). Rationally inattentive intertemporal choice. *Nature*

Communications, *11*(1), 3365. https://doi.org/10.1038/s41467-020-16852-y

Gillan, C. M., Kalanthroff, E., Evans, M., Weingarden, H. M., Jacoby, R. J., Gershkovich, M.,

Snorrason, I., Campeas, R., Cervoni, C., Crimarco, N. C., Sokol, Y., Garnaat, S. L.,

McLaughlin, N. C. R., Phelps, E. A., Pinto, A., Boisseau, C. L., Wilhelm, S., Daw, N. D.,

& Simpson, H. B [H. B.] (2020). Comparison of the Association Between Goal-Directed

Planning and Self-reported Compulsivity vs Obsessive-Compulsive Disorder Diagnosis.

JAMA Psychiatry, *77*(1), 77–85. https://doi.org/10.1001/jamapsychiatry.2019.2998

Gillan, C. M., Kosinski, M., Whelan, R., Phelps, E. A., & Daw, N. D. (2016). Characterizing

a psychiatric symptom dimension related to deficits in goal-directed control. *ELife*, *5*.

https://doi.org/10.7554/eLife.11305

Hautzinger, M., Keller, F., & Kühner, C. (2009). BDI-II. Beck-Depressions-Inventar.

Revision. 2, Auflage.

Heatherton, T. F., Kozlowski, L. T., Frecker, R. C., & Fagerstrom, K. O. (1991). The

Fagerström Test for Nicotine Dependence: A revision of the Fagerstrom Tolerance

Questionnaire. *British Journal of Addiction*, *86*(9), 1119–1127.

Joukhador, J., Blaszczynski, A., & Maccallum, F. (2004). Superstitious beliefs in gambling

among problem and non-problem gamblers: Preliminary data. *Journal of Gambling*

Studies, *20*(2), 171–180. https://doi.org/10.1023/B:JOGS.0000022308.27774.2b

Kleiner, M., Brainard, D., & Pelli, D. (2007). *What's new in Psychtoolbox-3?*

https://pure.mpg.de/rest/items/item_1790332/component/file_3136265/content

Kool, W., Cushman, F. A., & Gershman, S. J. (2016). When Does Model-Based Control Pay

Off? *PLOS Computational Biology*, *12*(8), e1005090.

https://doi.org/10.1371/journal.pcbi.1005090

Kroemer, N. B., Lee, Y., Pooseh, S., Eppinger, B., Goschke, T., & Smolka, M. N. (2019). L-

DOPA reduces model-free control of behavior by attenuating the transfer of value to

action. *NeuroImage*, *186*, 113–125. https://doi.org/10.1016/j.neuroimage.2018.10.075

Leeman, R. F., & Potenza, M. N. (2012). Similarities and differences between pathological

gambling and substance use disorders: A focus on impulsivity and compulsivity.

Psychopharmacology, *219*(2), 469–490. https://doi.org/10.1007/s00213-011-2550-7

Lempert, K. M., & Phelps, E. A. (2016). The Malleability of Intertemporal Choice. *Trends in*

Cognitive Sciences, *20*(1), 64–74. https://doi.org/10.1016/j.tics.2015.09.005

Lempert, K. M., Steinglass, J. E., Pinto, A., Kable, J. W., & Simpson, H. B [Helen Blair]

(2019). Can delay discounting deliver on the promise of RDoC? *Psychological Medicine*,

49(2), 190–199. https://doi.org/10.1017/S0033291718001770

Lesieur, H. R., & Blume, S. B. (1987). The South Oaks Gambling Screen (SOGS): A new

instrument for the identification of pathological gamblers. *American Journal of Psychiatry*,

144(9).

Lobo, D. S. S., Aleksandrova, L., Knight, J., Casey, D. M., el-Guebaly, N., Nobrega, J. N., &

Kennedy, J. L. (2015). Addiction-related genes in gambling disorders: New insights from

parallel human and pre-clinical models. *Molecular Psychiatry*, *20*(8), 1002.

https://doi.org/10.1038/mp.2014.113

MacKillop, J., Amlung, M. T., Few, L. R., Ray, L. A., Sweet, L. H., & Munafò, M. R. (2011).

Delayed reward discounting and addictive behavior: A meta-analysis.

Psychopharmacology, *216*(3), 305–321. https://doi.org/10.1007/s00213-011-2229-0

Martyn Plummer. (2003). Jags: A Program for Analysis of Bayesian Graphical Models Using

Gibbs Sampling. In M. Plummer (Ed.), *JAGS: a program for analysis of Bayesian*

graphical models using Gibbs sampling.: Proceedings of the 3rd International Workshop

on Distributed Statistical Computing (DSC 2003). 2003.

Miedl, S. F., Büchel, C., & Peters, J. (2014). Cue-induced craving increases impulsivity via

changes in striatal value signals in problem gamblers. *The Journal of Neuroscience : The*

Official Journal of the Society for Neuroscience, *34*(13), 4750–4755.

https://doi.org/10.1523/JNEUROSCI.5020-13.2014

Miedl, S. F., Peters, J., & Büchel, C. (2012). Altered neural reward representations in

pathological gamblers revealed by delay and probability discounting. *Archives of General*

Psychiatry, *69*(2), 177–186. https://doi.org/10.1001/archgenpsychiatry.2011.1552

Mikhael, J. G., Lai, L., & Gershman, S. J. (2021). Rational inattention and tonic dopamine.

PLOS Computational Biology, *17*(3), e1008659.

https://doi.org/10.1371/journal.pcbi.1008659

Moeller, S. J., & Paulus, M. P. (2018). Toward biomarkers of the addicted human brain:

Using neuroimaging to predict relapse and sustained abstinence in substance use disorder.

Progress in Neuro-Psychopharmacology & Biological Psychiatry, *80*(Pt B), 143–154.

https://doi.org/10.1016/j.pnpbp.2017.03.003

Muggleton, N., Parpart, P., Newall, P., Leake, D., Gathergood, J., & Stewart, N. (2021). The

association between gambling and financial, social and health outcomes in big financial

data. *Nature Human Behaviour.* Advance online publication.

https://doi.org/10.1038/s41562-020-01045-w

Myerson, J., Green, L., & Warusawitharana, M. (2001). Area under the curve as a measure of

discounting. *Journal of the Experimental Analysis of Behavior*, *76*(2), 235–243.

https://doi.org/10.1901/jeab.2001.76-235

Nelson, L. D., Strickland, C., Krueger, R. F., Arbisi, P. A., & Patrick, C. J. (2016).

Neurobehavioral Traits as Transdiagnostic Predictors of Clinical Problems. *Assessment*,

23(1), 75–85. https://doi.org/10.1177/1073191115570110

Ott, T., & Nieder, A. (2019). Dopamine and Cognitive Control in Prefrontal Cortex. *Trends in*

Cognitive Sciences, *23*(3), 213–234. https://doi.org/10.1016/j.tics.2018.12.006

Otto, A. R., Skatova, A., Madlon-Kay, S., & Daw, N. D. (2015). Cognitive control predicts

use of model-based reinforcement learning. *Journal of Cognitive Neuroscience*, *27*(2),

319–333. https://doi.org/10.1162/jocn_a_00709

Patzelt, E. H., Kool, W., Millner, A. J., & Gershman, S. J. (2019). Incentives Boost Model-

Based Control Across a Range of Severity on Several Psychiatric Constructs. *Biological*

Psychiatry, *85*(5), 425–433. https://doi.org/10.1016/j.biopsych.2018.06.018

Pedersen, M. L., Frank, M. J [Michael J.], & Biele, G. (2017). The drift diffusion model as the

choice rule in reinforcement learning. *Psychonomic Bulletin & Review*, *24*(4), 1234–1251.

https://doi.org/10.3758/s13423-016-1199-y

Peters, J., & Büchel, C. (2010). Episodic future thinking reduces reward delay discounting

through an enhancement of prefrontal-mediotemporal interactions. *Neuron*, *66*(1), 138–

148. https://doi.org/10.1016/j.neuron.2010.03.026

Peters, J., & Büchel, C. (2011). The neural mechanisms of inter-temporal decision-making:

Understanding variability. *Trends in Cognitive Sciences*, *15*(5), 227–239.

https://doi.org/10.1016/j.tics.2011.03.002

Peters, J., & D'Esposito, M. (2020). The drift diffusion model as the choice rule in inter-

temporal and risky choice: A case study in medial orbitofrontal cortex lesion patients and

controls. *PLOS Computational Biology*, *16*(4), e1007615.

https://doi.org/10.1371/journal.pcbi.1007615

Petry, J., & Baulig, T. (1996). KFG: Kurzfragebogen zum Glücksspielverhalten.

Psychotherapie Der Glücksspielsucht, 300–302.

Petry, N. M. (2010). Pathological gambling and the DSM-V. *International Gambling Studies*,

10(2), 113–115. https://doi.org/10.1080/14459795.2010.501086

Petzold, J., Kienast, A., Lee, Y., Pooseh, S., London, E. D., Goschke, T., & Smolka, M. N.

(2019). Baseline impulsivity may moderate L-DOPA effects on value-based decision-

making. *Scientific Reports*, *9*(1), 5652. https://doi.org/10.1038/s41598-019-42124-x

Pine, A., Shiner, T., Seymour, B., & Dolan, R. J. (2010). Dopamine, time, and impulsivity in

humans. *The Journal of Neuroscience : The Official Journal of the Society for

Neuroscience*, *30*(26), 8888–8896. https://doi.org/10.1523/JNEUROSCI.6028-09.2010

Raylu, N., & Oei, T. P. S. (2004). The Gambling Related Cognitions Scale (GRCS):

Development, confirmatory factor validation and psychometric properties. *Addiction

(Abingdon, England)*, *99*(6), 757–769.

Redick, T. S., Broadway, J. M., Meier, M. E., Kuriakose, P. S., Unsworth, N., Kane, M. J., &

Engle, R. W. (2012). Measuring Working Memory Capacity With Automated Complex

Span Tasks. *European Journal of Psychological Assessment*, *28*(3), 164–171.

https://doi.org/10.1027/1015-5759/a000123

Reynolds, B. (2006). A review of delay-discounting research with humans: Relations to drug

use and gambling. *Behavioural Pharmacology*, *17*(8), 651–667.

https://doi.org/10.1097/FBP.0b013e3280115f99

Robbins, T. W., & Everitt, B. J. (1999). Drug addiction: Bad habits add up. *Nature*,

398(6728), 567–570. https://doi.org/10.1038/19208

Robinson, M. J. F [M. J. F.], Fischer, A. M., Ahuja, A., Lesser, E. N., & Maniates, H. (2016).

Roles of "Wanting" and "Liking" in Motivating Behavior: Gambling, Food, and Drug

Addictions. *Current Topics in Behavioral Neurosciences*, *27*, 105–136.

https://doi.org/10.1007/7854_2015_387

Robinson, T., & Berridge, K. C [Kent C.] (1993). The neural basis of drug craving: An

incentive-sensitization theory of addiction. *Brain Research Reviews*, *18*(3), 247–291.

https://doi.org/10.1016/0165-0173(93)90013-P

Robinson, T. E., & Berridge, K. C [K. C.] (2001). Incentive-sensitization and addiction.

Addiction (Abingdon, England), *96*(1), 103–114. https://doi.org/10.1046/j.1360-

0443.2001.9611038.x

Robinson, T. E [Terry E.], & Berridge, K. C [Kent C.] (2008). Review. The incentive

sensitization theory of addiction: Some current issues. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *363*(1507), 3137–3146.

https://doi.org/10.1098/rstb.2008.0093

Rösch, S. A., Stramaccia, D. F., & Benoit, R. G. (2021). *Promoting farsighted decisions via*

episodic future thinking: A meta-analysis. https://doi.org/10.31234/osf.io/53ju2

Saunders, J. B., Aasland, O. G., Babor, T. F., La Fuente, J. R. de, & Grant, M. (1993).

Development of the alcohol use disorders identification test (AUDIT): WHO collaborative

project on early detection of persons with harmful alcohol consumption-II. *Addiction*

(Abingdon, England), *88*(6), 791–804.

Sebold, M., Deserno, L., Nebe, S [Stephan], Nebe, S [Stefan], Schad, D. J., Garbusow, M.,

Hägele, C., Keller, J., Jünger, E., Kathmann, N., Smolka, M. N., Smolka, M., Rapp, M. A.,

Schlagenhauf, F., Heinz, A., & Huys, Q. J. M. (2014). Model-based and model-free

decisions in alcohol dependence. *Neuropsychobiology*, *70*(2), 122–131.

https://doi.org/10.1159/000362840

Shahar, N., Hauser, T. U., Moutoussis, M., Moran, R., Keramati, M., & Dolan, R. J. (2019).

Improving the reliability of model-based decision-making estimates in the two-stage

decision task with reaction-times and drift-diffusion modeling. *PLoS Computational*

Biology, *15*(2), e1006803. https://doi.org/10.1371/journal.pcbi.1006803

Sharp, M. E., Foerde, K., Daw, N. D., & Shohamy, D. (2016). Dopamine selectively

remediates 'model-based' reward learning: A computational approach. *Brain : A Journal of*

Neurology, *139*(Pt 2), 355–364. https://doi.org/10.1093/brain/awv347

Shiner, T., Seymour, B., Wunderlich, K., Hill, C., Bhatia, K. P., Dayan, P [Peter], &
Dolan, R. J. (2012). Dopamine and performance in a reinforcement learning task: Evidence
from Parkinson's disease. *Brain : A Journal of Neurology*, *135*(Pt 6), 1871–1883.
https://doi.org/10.1093/brain/aws083

Singer, B. F., Anselme, P., Robinson, M. J. F [Mike J. F.], & Vezina, P. (2020). An overview
of commonalities in the mechanisms underlying gambling and substance use disorders.
Progress in Neuro-Psychopharmacology & Biological Psychiatry, *101*, 109944.
https://doi.org/10.1016/j.pnpbp.2020.109944

Stan Development Team. (2020). *Stan* (Version 2.26) [Computer software]. https://mc-
stan.org

Toyama, A., Katahira, K., & Ohira, H. (2017). A simple computational algorithm of model-
based choice preference. *Cognitive, Affective & Behavioral Neuroscience*, *17*(4), 764–783.
https://doi.org/10.3758/s13415-017-0511-2

Toyama, A., Katahira, K., & Ohira, H. (2019). Reinforcement Learning With Parsimonious
Computation and a Forgetting Process. *Frontiers in Human Neuroscience*, *13*, 153.
https://doi.org/10.3389/fnhum.2019.00153

van den Noort, M., Bosch, P., Haverkort, M., & Hugdahl, K. (2008, January 21). *A Standard
Computerized Version of the Reading Span Test in Different Languages.* Hogrefe & Huber
Publishers. https://econtent.hogrefe.com/doi/abs/10.1027/1015-5759.24.1.35

van Gaalen, M. M., van Koten, R., Schoffelmeer, A. N. M., & Vanderschuren, L. J. M. J.
(2006). Critical involvement of dopaminergic neurotransmission in impulsive decision
making. *Biological Psychiatry*, *60*(1), 66–73.
https://doi.org/10.1016/j.biopsych.2005.06.005

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using

leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

https://doi.org/10.1007/s11222-016-9696-4

Voon, V [V.], Derbyshire, K., Rück, C., Irvine, M. A., Worbe, Y., Enander, J.,

Schreiber, L. R. N., Gillan, C., Fineberg, N. A., Sahakian, B. J., Robbins, T. W.,

Harrison, N. A., Wood, J., Daw, N. D [N. D.], Dayan, P [P.], Grant, J. E., &

Bullmore, E. T. (2015a). Disorders of compulsivity: A common bias towards learning

habits. *Molecular Psychiatry*, *20*(3), 345–352. https://doi.org/10.1038/mp.2014.44

Voon, V [V.], Derbyshire, K., Rück, C., Irvine, M. A., Worbe, Y., Enander, J.,

Schreiber, L. R. N., Gillan, C., Fineberg, N. A., Sahakian, B. J., Robbins, T. W.,

Harrison, N. A., Wood, J., Daw, N. D [N. D.], Dayan, P [P.], Grant, J. E., &

Bullmore, E. T. (2015b). Disorders of compulsivity: A common bias towards learning

habits. *Molecular Psychiatry*, *20*(3), 345–352. https://doi.org/10.1038/mp.2014.44

Voon, V [Valerie], Reiter, A., Sebold, M., & Groman, S. (2017). Model-Based Control in

Dimensional Psychiatry. *Biological Psychiatry*, *82*(6), 391–400.

https://doi.org/10.1016/j.biopsych.2017.04.006

Wagner, B., Clos, M., Sommer, T., & Peters, J. (2020). Dopaminergic Modulation of Human

Intertemporal Choice: A Diffusion Model Analysis Using the D2-Receptor Antagonist

Haloperidol. *The Journal of Neuroscience : The Official Journal of the Society for*

Neuroscience, *40*(41), 7936–7948. https://doi.org/10.1523/JNEUROSCI.0592-20.2020

Westbrook, A [A.], Frank, M [MJ.], & Cools, R. (2021). A mosaic of cost-benefit control

over cortico-striatal circuitry. *Trends in Cognitive Sciences.* Advance online publication.

https://doi.org/10.1016/j.tics.2021.04.007

Westbrook, A [A.], van den Bosch, R., Määttä, J. I., Hofmans, L., Papadopetraki, D.,

Cools, R., & Frank, M. J [M. J.] (2020). Dopamine promotes cognitive effort by biasing

the benefits versus costs of cognitive work. *Science*, *367*(6484), 1362–1366.

https://doi.org/10.1126/science.aaz5891

Wiehler, A., & Peters, J. (2015). Reward-based decision making in pathological gambling:

The roles of risk and delay. *Neuroscience Research*, *90*, 3–14.

https://doi.org/10.1016/j.neures.2014.09.008

Wilke, A., Scheibehenne, B., Gaissmaier, W., McCanney, P., & Barrett, H. C. (2014).

Illusionary pattern detection in habitual gamblers. *Evolution and Human Behavior*, *35*(4),

291–297. https://doi.org/10.1016/j.evolhumbehav.2014.02.010

Wunderlich, K., Smittenaar, P., & Dolan, R. J. (2012). Dopamine enhances model-based over

model-free choice behavior. *Neuron*, *75*(3), 418–424.

https://doi.org/10.1016/j.neuron.2012.03.042

Wyckmans, F., Otto, A. R., Sebold, M., Daw, N., Bechara, A., Saeremans, M., Kornreich, C.,

Chatard, A., Jaafari, N., & Noël, X. (2019). Reduced model-based decision-making in

gambling disorder. *Scientific Reports*, *9*(1), 19625. https://doi.org/10.1038/s41598-019-

56161-z

1133

1134

1135

1136

1137

1138

1139

1140   **Supplemental Information**

1141   **Supplemental Table S1.** Baseline screening questionnaires.

	Reference	Measure
AUDIT	(Saunders et al., 1993)	Alcohol-Use-Disorders Identification Text
BDI	(Beck et al., 1996; Hautzinger et al., 2009)	Beck Depression Inventory II
DSM-5	(American Psychiatric Association, 2013; Falkai, 2015)	Diagnostic criteria for gambling disorder
FTND	(Heatherton et al., 1991)	Fagerström Test for Nicotine Dependence
GRCS	(Raylu & Oei, 2004)	Gambling-related cognition scale
KFG	(J. Petry & Baulig, 1996)	Kurzfragebogen zum Glücksspielverhalten
SOGS	(Lesieur & Blume, 1987)	South Oaks Gambling Screen

1142

1143

1144   **Supplemental Table S2.** Summary of demographics and clinical information (n = 30).

	Mean	SD	Range
Age	30.87	7.94	20-47
School years	11.83	1.93	8-17
Income	1284.73	590.71	300-2500
FTND	2.5	3.01	0-9
AUDIT	7.6	7.10	0-26
DSM-5	5.9	2.02	3-9
KFG	27.54	10.31	7-54
SOGS	9.5	3.87	3-16
BDI	17.27	10.13	0-45
GRCS	17.60	4.54	9.57-30.4

1145

Temporal discounting drift diffusion models (DDMs)

Model comparison and validation

We compared three versions of the drift diffusion model (DDM) that varied in the way that they

accounted for the influence of value differences on trial-wise drift rates, based on model-fit

(WAIC). To verify comparable model ranking across conditions, we first carried out a model

comparison separately for each environment (see Supplemental Table S3). In both

environments, a DDM with nonlinear drift-rate scaling (DDM_s) (Fontanesi et al., 2019; Peters

& D'Esposito, 2020; Wagner et al., 2020) accounted for the data best when compared to a DDM

with linear scaling (DDM_{lin}) (Pedersen et al., 2017) and a null model without value modulation

(DDM_0).

1156   We then build a full model with group level distributions for the baseline condition

(neutral context) and s_x parameters for each model parameter x, modeling the change from the

neutral to the gambling context. S_x parameters where modeled with Gaussian priors with means

of zero (see methods section). Model ranking was confirmed for the full model (Supplemental

Table S3). We next compared the DDMs and the softmax model with respect to the proportion

of binary choices (LL vs. SS selections) that they correctly accounted for. As can be seen from

Supplemental Table S4, the DDM_S and DDM_{lin} performed numerically on par with the softmax

model, whereas the DDM_0 performed substantially worse (see Supplemental Figure S1,

Supplemental Table S4). Posterior predictive checks for the winning model showed that it

accurately captured the effect of decision conflict (value difference) on RTs (see section *Posterior Predictive Checks* below and Supplemental Figure S2). Parameter recovery for this model was reported in our prior papers (Peters & D'Esposito, 2020; Wagner et al., 2020).

Supplemental Table S3 Temporal discounting DDM model comparison using the Watanabe-Akaike Information Criterion (WAIC) revealed the same model ranking for each context (neutral vs. gambling) and the full model. Scores are WAIC (SE).

	Neutral	**Gambling**	**Full model**
DDM_0	12037.7 (150.1)	11754.9 (157.7)	23792.4 (217.4)
DDM_{lin}	9304.6 (155.5)	9174.7 (158.5)	18949.6 (219.0)
DDM_s	8982.3 (155.6)	8744.6 (157.6)	17656.0 (220.8)

Supplemental Table S4 Proportions of correctly predicted binary choices (mean [range]) for the temporal discounting models (neutral vs. gambling context; see Supplemental Figure S1).

	Neutral	**Gambling**
Softmax	0.89 [0.57-0.98]	0.90 [0.66-1.00]
DDM_0	0.74 [0.52-0.93]	0.76 [0.55-0.99]
DDM_{lin}	0.89 [0.58-0.98]	0.90 [0.65-0.99]
DDM_s	0.90 [0.60-0.99]	0.91 [0.70-1.00]

1176

Supplemental Figure S1. Proportions of correctly predicted binary choices for the softmax
model (A) and the drift diffusion model with non-linear drift rate scaling (B, DDMs) in both
contexts (neutral [blue], gambling [pink]).

Posterior predictive checks

Temporal discounting

We carried out posterior predictive checks to visualize if our computational analysis captures

key aspect in the data, in particular the value-dependency of RTs (Peters & D'Esposito, 2020;

Wagner et al., 2020) . For the temporal discounting task, we binned trials per participant into

five bins according to the absolute difference in larger-later vs. smaller-sooner value ("decision

conflict", computed according to each participant's median posterior $\log(k)$ parameter from the

DDM_S, and separately for the neutral and gambling context conditions). We then plotted the

mean observed RTs as a function of decision conflict per participant and context, as well as the

mean RTs across 10.000 data sets simulated from the posterior distributions of the DDM_0,

DDM_{lin} and DDM_S (see Supplemental Figure S2).

1191

Supplemental Figure S2. Posterior predictive checks for temporal discounting drift diffusion
models. For each participant and condition (Gambling vs. Neutral), trials were binned into five
equal sized bins according to the absolute difference between subjective LL and SS option
values (decision conflict bin). Plotted are mean observed RTs per bin (data) as well model-
generated RTs (blue: DDM_0, red: DDM_{lin}, orange: DDM_S) averaged over 10,000 datasets
simulated from the respective posterior distributions of the hierarchical models.

1198

Supplemental Table S5. Model agnostic analysis of stay probability via a hierarchical general
linear model (HGLM). HGLMs were estimated for each context separately using reward and
transition as fixed and subject as random effects. The full model with stay probability as
dependent variable included the predictors reward, transition (rare vs. common) and context
(gambling vs. neutral) as fixed effects and subject as random effect. As a model-agnostic
performance measure, the probability of choosing the same S1 option as in the previous trial
(stay-probability) is typically analyzed as a function of reward, transition, and their interaction
(Daw et al., 2011). Since the 2-step task version employed here utilized continuous payoffs,
every trial was rewarded. The "reward" in S2 can thus not be used to directly predict stay
probabilities, as done in previous work. Therefore, here the "reward" factor is computed relative
to a moving average of recent rewards as a reference. Specifically, we categorized a reward R_{t-1}
as positive "R+" if R_{t-1} was higher than the mean of last 7 rewards ($R_t > \text{mean}[R_{t-1: t-7}]$) and as
negative "R-" if $R_t < \text{mean}(R_{t-1:t-7})$.

Neutral Context			
	Estimate	z-Value	p
Reward	0.27350	4.127	3.68e-05***
Transition	0.29908	3.455	0.00055***
Reward*Transition	-0.51326	-4.190	2.79e-05***
Gambling Context			
Reward	0.38419	5.889	3.89e-09***
Transition	0.39439	4.653	3.27e-06***
Reward*Transition	-0.71324	-5.927	3.08e-09***
Full Model			
Reward	0.27361	4.131	3.60e-05***
Transition	0.29924	3.460	0.00054***
Context	-0.12701	-0.426	0.67015
Reward*Transition	-0.51377	-4.199	2.68e-05***
Reward*Context	0.11057	1.190	0.23423
Transition*Context	0.09502	0.785	0.43249
Reward*Transition*Context	-0.19892	-1.160	0.24614

1212

Model free analysis of Stage 1 RTs

S1 RTs were modeled as a function of categorized reward in the previous trial (see previous
section for how this was defined) and context as fixed effects and trial and subject as random
effects. Previous reward significantly increased RTs (t = -2.431, p = 0.015, see Supplemental
Table S6). We also observed a reward * context interaction (see Supplemental Table S6),
reflecting Also reaction times were slower, when previously rewarded in the gambling context
when contrasted to the neutral context indicating an increased effect of reward on S1 response
caution in the gambling context.

1221

1222 **Supplemental Table S6.** Hierarchical general linear model results of S1 RTs with reward and
1223 context as fixed effects and subject as random effect.

S1 RT Model			
	Estimate	t-Value	p
Reward	-0.025	-2.431	0.015*
Context	-0.0005	-0.044	0.97
Reward*Context	0.029	2.050	0.04*

1224

1225

1226 *Model free analysis of Stage 2 RTs*

1227 **Supplemental Table S7**. Model agnostic analysis of S2 RTs. HGLM with transition and
1228 context as fixed effects and subject as random effect.

S2 RT Model			
	Estimate	**T-statistic**	p
Transition	0.13	17.227	< 2e-16***
Context	0.0004	0.07	0.94
Transition*Context	0.02	1.804	0.07

1229

1230

1231

1232 **Supplemental Figure S3** Model free analysis of S2 RTs. RTs were substantially slower
1233 following rare transitions, both in the neutral (A) and the gambling context (B), see also Table
1234 4.

1235

Hybrid model with drift diffusion choice rule

Model comparison and validation

Model comparison based on the WAIC (Vehtari et al., 2017) (see Supplemental Table S8)

revealed that in the neutral context, a DDM with nonlinear drift-rate scaling DDM_s (Fontanesi

et al., 2019; Peters & D'Esposito, 2020; Wagner et al., 2020) accounted for the data best when

compared to a DDM with linear drift rate scaling (DDM_{lin}) (Pedersen et al., 2017) and a null-

model without learning (DDM0) (see Supplemental Table S8). The same ranking held for the

gambling context.

We next build a full model with group level distributions for the baseline condition

(neutral context) and additional s_x parameters for each model parameter x, modeling the change

in from the neutral to the gambling context. These s_x parameters where modeled with Gaussian

priors with means of zero (see methods section). The full model reproduced the model ranking

(see Supplemental Table S8). We then compared the three DDMs and the softmax model with

respect to the proportion of binary choices that they correctly accounted for. As can be seen

from see Supplemental Table S9, the DDM_S and DDM_{lin} performed numerically on par with

the softmax model, whereas the DDM_0 performed substantially worse. Posterior predictive

checks showed that the final model accurately captured the effect of reward differences on

second stage RTs and reproduced choice behavior (see Supplemental Figure S4 and S5 below).

1254

1255 **Supplemental Table S8**. Reinforcement learning DDM model comparison using the Widely-
1256 Applicable Information Criterion (WAIC) revealed the same model ranking for each condition
1257 (neutral or gambling context) as well as for the full model. Scores are WAIC (SE).

	Neutral	Gambling	Full model
DDM0	11808.5 (1549.3)	10942.1 (1569.2)	22764.0 (2193.7)
DDMlin	4616.2 (1897.0)	4429.0 (1729.9)	15519.7 (3984.4)
DDMs	4357.2 (1935.1)	4197.2 (1749.7)	8800.3 (2670.0)

1258

1259

1260 **Supplemental Table S9**. 2-step task models. Proportions of correctly predicted binary choices
1261 (mean [range]) for all models.

	Neutral		Gambling	
	Stage 1	Stage 2	Stage 1	Stage 2
DDM$_0$	0.63 [0.49-1.00]	0.62 [0.50-0.78]	0.56 [0.46-0.99]	0.63 [0.51-0.79]
DDM$_{lin}$	0.74 [0.51-1.00]	0.80 [0.53-0.96]	0.74 [0.50-0.99]	0.81 [0.59-0.95]
DDM$_s$	0.74 [0.49-1.00]	0.80 [0.55-0.96]	0.72 [0.47-0.99]	0.81 [0.59-0.95]
Softmax	0.72 [0.42-1.00]	0.79 [0.49-0.96]	0.72 [0.45-0.99]	0.81 [0.56-0.96]

1262

1263

Posterior predictive checks

We conducted posterior predictive checks to evaluate if our different hierarchical models
capture both the relationship of RTs and reward differences and the relationship of reward
differences and optimal choices. An optimal choice here is defined as a choice for the random
walk with highest payout. To this end, we binned all trials into four bins, according to the
absolute max(reward) differences in stage 2. For each reward difference bin we then plot the
mean observed RTs, as well as the mean simulated RTs across 1000 datasets simulated using
our mean parameter estimates for the posterior distributions of the DDM$_0$, DDM$_{lin}$, and DDM$_S$.
We further show the mean observed optimal choices (max[reward]) vs. the mean simulated
optimal choices given our mean parameter estimates for the posterior distribution of each
model. These results are shown in Supplemental Figures S4 and S5. As can be seen, the DDM$_s$
provided the best account of how RTs vary as a function of reward differences. This model
outperformed the other models in capturing the relationship of reward differences and optimal
choices (Supplemental Figure S5).

1278

1279

Supplemental Figure S4. Group level posterior predictive checks. Trials were binned into four
equal sized bins according to the absolute difference in *reward* values given S2 reward walks.
Plotted are mean observed RTs per bin (data; dashed line) as well model-generated RTs (blue
represents DDM_0; orange represents DDM_{lin}; red represents DDM_S) averaged 1000 simulated
datasets simulated from the mean parameter estimates the posterior distribution of each
hierarchical model.

1286

Supplemental Figure S5. Group level posterior predictive checks. Trials were binned into four
equal sized bins according to the absolute difference in *reward* values given S2 reward walks.
Plotted are mean optimal choices in % of all choices per bin (data; dashed line) as well model-
generated RTs (blue represents DDM_0; orange represents DDM_{lin}; red represents DDM_S)
averaged 1000 simulated datasets simulated from the mean parameter estimates for the
posterior distribution of each hierarchical model.

1293

1294 **Supplemental Results 1: Working memory, subjective craving and temporal discounting**

1295 Preregistered analysis:

1296 We hypothesized a positive relationship of decision noise parameter and working memory z-

1297 score, which we confirmed in each context (neutral condition: $r = 0.45$, $p = 0.013$); gambling

1298 condition: $r = 0.45$ $p =$ $p = 0.013$).

1299

Exploratory analysis:

There was no significant correlation between discount-rate and working memory ($r = -0.03$, p

$= 0.89$). A further exploratory analysis revealed an association of working memory and drift-

rate coefficient. Here, higher working memory capacity was associated with higher drift rate

coefficients (neutral condition: $r = 0.42$, $p = 0.02$; gambling condition: $r = 0.37$, $p = 0.048$).

The difference in subjective craving from the neutral to the gambling environment, was not

significantly associated with the change in the discount-rate (pre-task craving rating: $r = 0.28$,

$p = 0.18$; post-task craving rating: 0.09, $p = 0.68$).

1308

Supplemental Results 2: Working memory, subjective craving and 2-step task

performance

Preregistered analysis:

WM z-score was positively but non-significant correlated with MB RL (neutral context: $r =$

0.27, $p = 0.16$, gambling context: $r = 0.31$, $p = 0.10$). The change in MB RL from neutral to

gambling context was not significantly correlated with WM ($r = 0.27$, $p = 0.16$).

1315

Exploratory analysis:

We further explored the association of WM capacity and S2 learning rates. This analysis

revealed that overall WM capacity was positively associated with baseline (neutral context) (r

$= 0.57$, $p = 0.001$) and gambling context ($r = 0.37$, $p = 0.048$) S2 learning rates. The change in

subjective craving from the neutral to the gambling environment, was not significantly

associated with the change in MB drift-rate weight (pre-task craving rating: -0.01, $p = 0.98$;

post-task craving rating: $r = 0.33$, $p = 0.14$).

1323

1324

1325 **Supplemental Figure S6** Gambling severity (y-axis; average z-score across DSM, KFG and
1326 SOGS) was associated with a greater gambling context related decrease in MF drift-rate weights
1327 ($S_{vcoeffMF}$, r = -0.48, p = 0.009).

1328

1329