Feedbacks between size and density determine rapid eco-phenotypic dynamics

ABSTRACT

1. Body size is a fundamental trait linked to many ecological processes—from individuals to
ecosystems. Although the effects of body size on metabolism are well-known, the potential
reciprocal effects of body size and density are less clear. Specifically, 1) whether changes in
body size or density more strongly influence the other and 2) whether coupled rapid changes in
body size and density are due to plasticity, rapid evolutionary change, or a combination of both.
30
2. Here, we address these two issues by experimentally tracking population density and mean
body size in the protist *Tetrahymena pyriformis* as it grows from low density to carrying
capacity. We then use *Convergent Cross Mapping* time series analyses to infer the direction,
magnitude, and causality of the link between body size and ecological dynamics. We confirm the
results of our analysis by experimentally manipulating body size and density while keeping the
other constant. Last, we fit mathematical models to our experimental time series that account for
purely plastic change in body size, rapid evolution in size, or a combination of both, to gain
insight into the processes that most likely explain the observed dynamics.
39
40    3. Our results indicate that changes in body size more strongly influence changes in density than
41    the other way around, but also show that there is reciprocity in this effect (i.e., a *feedback*). We
42    show that a model that only accounts for purely plastic change in size most parsimoniously
43    explains observed, coupled phenotypic and ecological dynamics.
44
4. Together, these results suggest 1) that body size can shift dramatically through plasticity, well
within ecological timescales, 2) that rapid changes in body size may have a larger effect on
ecological dynamics than the reverse, but 3) phenotypic and ecological dynamics influence each
as populations grow. Overall, we show that rapid plastic changes in functional traits like body
size may play a fundamental –but currently unrecognized– role in familiar ecological processes
such as logistic population growth.

51
52
53
54
55    Key Words: Plasticity, Eco-evolutionary Dynamics, Traits, Phenotypes, Phenotypic change

INTRODUCTION

Body size influences organismal energetic demands (Gillooly *et al.* 2001; Brown *et al.*

2004), diet breadth (Wasserman & Mitter 1978; Gravel *et al.* 2013), the strength of ecological

interactions (Berlow *et al.* 2009), trophic cascades (DeLong *et al.* 2015), and food web structure

(Brose *et al.* 2006; Riede *et al.* 2011; Gravel *et al.* 2013; Gibert & DeLong 2014), all of which

have ecosystem-level consequences (Anderson-Teixeira, Vitousek & Brown 2008; Trebilco *et al.*

2013). Because of its myriad ecological consequences—from individuals (e.g., (Glasheen &

McMahon 1996; Hurlbert, Ballantyne IV & Powell 2008; Pavković-Lučić & Kekić 2013)) to

ecosystems (e.g., (Brown *et al.* 2004; Riede *et al.* 2011; Trebilco *et al.* 2013))—body size is one

of the most important functional traits.

Body size can fuel or limit population growth through well-known relationships with

demographic parameters like carrying capacity (K) and the intrinsic growth rate (r) (Damuth

1981; Savage *et al.* 2004; DeLong *et al.* 2015). For example, smaller organisms typically

reproduce faster and can reach higher carrying capacities than larger ones, but also have higher

mortality rates, which leads to faster population turnover (e.g., (Huryn & Benke 2007)). As a

consequence, body size often determines how populations grow, and hence, ecological

dynamics. On the flip side, ecological dynamics can themselves influence body size, although

these effects are not well understood. For example, individual size is under physiological

regulatory control (Davidowitz, Roff & Nijhout 2005; Chelini, Delong & Hebets 2019) and

therefore responds to changes in resource levels (Marañón *et al.* 2013). While resource levels can

vary independently of population density (Holt 2008; Fey, Gibert & Siepielski 2019; Nguyen,

Lara-Gutiérrez & Stocker 2020), high density leads to crowding and increasing intra-specific

competition (Gavina *et al.* 2018). Stronger competition in turn reduces resource availability at

the individual level, which may result in stunted growth and smaller body sizes (e.g., (Vanni *et*

*al.* 2009)). Consequently, while body size can, and often does, influence ecological dynamics,

ecological dynamics can also influence body size. Which one more strongly influences the other,

however, is not well understood, among other reasons because both population density and body

size can change dramatically over time (DeLong, Hanley & Vasseur 2014; Clements & Ozgul

2016).

Due to either natural or sexual selection (Preziosi & Fairbairn 2000; Chelini, Delong &

Hebets 2019), body size can change evolutionarily (Hairston *et al.* 2005; DeLong *et al.* 2016).

For example, predation selects for smaller but faster-growing prey in pitcher plant inquiline

communities (terHorst, Miller & Levitan 2010) and for shorter developmental times that result in

smaller individuals in mayflies (Peckarsky *et al.* 2001). Selection can act on traits rather quickly

(Thompson 1998; Hairston *et al.* 2005) and rapid evolutionary change has been shown to

influence ecological dynamics as they unfold (e.g., (Becks *et al.* 2012; Rudman *et al.* 2018;

Schaffner *et al.* 2019)), while ecological dynamics, in turn, influence the pace and direction of

evolutionary change (e.g., (Cortez 2016; DeLong & Gibert 2016; Frickel, Sieber & Becks 2016;

Gibert & Yeakel 2019)). Therefore, selection imposed by ecological dynamics may influence

body size, whose rapid change can alter ecological dynamics. Because studying rapid

evolutionary change in body size is unfeasible for most organisms, it is unclear how pervasive

these processes are in nature.

In addition to evolution, body size can change within generations through plasticity

(David, Legout & Moreteau 2006; Ghosh, Testa & Shingleton 2013; Lafuente, Duneau &

Beldade 2018; Cameron *et al.* 2020). For example, organisms grow in size throughout ontogeny

and the environment often influences those ontogenetic trajectories, leading to plastic variation

in body size (Lafuente, Duneau & Beldade 2018; Chelini, Delong & Hebets 2019). Epigenetic

DNA modifications can also result in rapid phenotypic change from one generation to the next in

response to shifts in biotic or abiotic conditions (e.g., maternal effects, (Roach & Wulff 1987;

Galloway & Etterson 2007)). Teasing apart the contributions of plastic and evolutionary

processes on body size has been the subject of great scientific interest (Amarillo-Suarez, Stillwell

& Fox 2011; Walczyńska, Franch-Gras & Serra 2017; Lafuente, Duneau & Beldade 2018;

Yengo *et al.* 2018; Cox *et al.* 2019). However, whether plasticity or evolution more strongly

influences rapid changes in body size, especially when this is coupled to rapidly shifting

ecological dynamics, is not well understood (e.g.,  (DeLong, Hanley & Vasseur 2014)). Among

other reasons, this is because teasing apart plastic and evolutionary change over short periods of

time is challenging, even when sufficiently long time series are available (Ellner, Geber &

Hairston 2011). As body size mediates ecological interactions and processes, and can change on

ecological timescales, it is important to understand, track, and predict such change.

Here, we address these gaps by answering the following questions: 1) Do rapid changes

in body size more strongly influence population dynamics (i.e., changes in density over time), or

do population dynamics more strongly influence changes in body size? and 2) Are observed

changes in body size most consistent with a model of plasticity, rapid evolution, or one that

accounts for both plastic and evolutionary change? To address the first question, we track

changes in the density and average body size of multiple experimental populations of the protist

*Tetrahymena pyriformis*, then use time series analysis to infer causality. In protists, we expect

changes in body size to be at least partly caused by plasticity because reproduction (cell division)

is tightly linked to ontogenetic changes in body size (cells grow then divide when a critical size

is attained). However, *T. pyriformis* also reproduces extremely fast (~4 generations per day) and

exhibits wide standing variation in body size (Wieczynski *et al.* 2021), so rapid evolutionary

change is also possible. Therefore, to distinguish the impacts of plasticity and evolution on

changes in body size, we fit alternative mathematical models (Plasticity Model, Eco-

Evolutionary Model, and Plasticity + Eco-Evo Model) to our experimental time series and use

model selection to infer which one best explains our data. Our results show that rapid, purely

plastic changes in body size more strongly influence changes in density than the other way

around, thus suggesting that plastic phenotypic change may be integral to ecological dynamics.

METHODS

*Microcosm growth assays*

We tracked changes in abundance and body size in the protist *Tetrahymena pyriformis* for 14

days. To do so, we set up 6 experimental microcosms in 250 mL autoclaved borosilicate jars

filled with 200 mL of Carolina protist pellet media (1L of autoclaved DI water per pellet)

inoculated with pond bacteria from Duke Forest (Gate 9/Wilbur pond, Lat=36.02°, Long=-

78.99°, Durham, NC) and a wheat seed as a carbon source. Microcosms were initialized at

densities of 10 ind/mL and incubated in temperature (22°C) and humidity-controlled (65%

humidity) growth chambers (Percival AL-22L2, Percival Scientific, Perry, Iowa) on a 12hr

night/day cycle. Densities (ind/mL) and trait dynamics were tracked daily for two weeks through

fluid imaging of 1 mL subsamples of each microcosm (Fig 1a, FlowCam, Fluid Imaging

Technologies, Scarborough, ME, USA). The FlowCam can image particles ranging from 5-10

μm to 2mm in length. Cell images were automatically sorted and measured by the FlowCam's

proprietary software yielding individual-level data on 150k cells over 14 days, giving our

experiment unparalleled insight into how density and body size changed together over the course

of this experiment. Using these data, density was quantified as a simple cell count per volume

sampled and body size was quantified as the volume of a spheroid, in $\mu$m$^3$. Last, we quantified

changes in total biomass, measured as the sum of the mass of all individual cells in a sample (in

grams, *g*, obtained by converting protist volumes estimated by the FlowCam from $\mu$m$^3$ to cm$^3$

and assuming that the density of protists equals that of water, i.e., 1g/cm$^3$). Neither water nor

nutrients were replaced throughout the course of this experiment.

*Time-series analysis*

To assess whether change in body size more strongly influenced changes in density, or

vice versa, we used *Convergent Cross Mapping* (or CCM, (Sugihara *et al.* 2012; Rogers *et al.*

2020)) on the density and body size time series. Conceptually, CCM quantifies the degree to

which one time series causally influences another one by estimating how much information of

the one is contained in the other (Takens 1981; Sugihara *et al.* 2012). If a variable X causally

influences another variable Y, but Y does not influence X, Y should contain information about

X, but not the other way around. CCM does that by quantifying whether variable X can be

predicted from the time series of Y (and vice-versa) for subsets of the time series of increasing

length (this procedure is called 'cross-mapping'). If X more strongly influences changes in Y

than the other way around, it also means Y responds to X more strongly than X responds to Y. If

the effect of X on Y is causal –as opposed to there being simple correlation with an unobserved

variable Z– the ability to predict Y from X should increase with library size, while the error

associated with that prediction should decline. If predictability does not change with library size,

there is correlation, but not causation (e.g., variable X and Y could be responding to a third

unobserved variable Z instead of to each other, leading to spurious correlation between the two,

(Sugihara *et al.* 2012)).

To perform the analysis, we used R package multispatialCCM v1.0 (Clark *et al.* 2015),

which works on replicated time series. In a nutshell, the procedure operates as follows: first, the

algorithm does state-space reconstruction using 'delay-embedding' (Sugihara *et al.* 2012). That

is, it attempts to reconstruct the manifold of the system (i.e., the collection of all states taken by

all variables for all time points) using lagged versions of each variable, one at a time. The

number of lagged versions of each time series needed for this reconstruction is called the

'Embedding Dimension', $E$ (Sugihara & May 1990; Sugihara *et al.* 2012). A value of E much

larger than the number of observed variables suggests effects of other non-observed variables on

the dynamics of the system (Sugihara *et al.* 2012). Second, the 'multispatialCCM' version of the

CCM algorithm takes bootstrap pseudo-replicates (n=800) of varying size (i.e., library size,

ranging from E to E*n, where n is the number of replicated time series) across replicates (Clark

*et al.* 2015). Third, it uses those bootstrapped time series to 'cross-map', that is, to predict the

values of one state variable based on exponentially weighted values of the reconstructed

manifold of the system using the other state variable (i.e., 'predicting X based on information

contained in Y'). By quantifying the correlation coefficient (or the *predictability* coefficient) of

the observed and predicted values, CCM produces a measure of how strong of a causal effect one

variable has on the other (Sugihara *et al.* 2012), if such an effect exists and is indeed causal.

Multiple previous studies have already shown how well CCM infers causation in

different ecological systems and environmental conditions (Sugihara *et al.* 2012; Clark *et al.*

2015; Karakoç, Clark & Chatzinotas 2020; Kondoh *et al.* 2020; Rogers *et al.* 2020).

 *Experimental manipulations of size and density*

In addition to our time series analysis, we experimentally manipulated density and size,

while controlling for the other variable, to assess whether possible effects of size on density and

vice-versa could be detected experimentally.

*Experiment 1: effect of density on body size.* To manipulate density while keeping body

size constant, we started 30 microcosms (as detailed in *Microcosm growth assays* subsection)

populated with *T. pyriformis* at an initial density of 10ind/mL, at Day -2. At Day 0, we filtered

half of the volume (100 mL) in 15 of those microcosms using Whatman GF/A filters, which have

a pore size small enough to filter out the protists, but large enough to allow the bacteria protists

feed on to pass through. The original microcosms were then replenished with the filtered water

with bacteria (but no protists). This procedure halved the density of 15 out of the initial 30 jars

(Fig 1b) while keeping the size distribution of *T. pyriformis*, growth medium, and number of

bacteria, the same, in 'low' (jars with half filtered, half unfiltered growth medium) and 'high'

density treatments (jars with unfiltered growth medium).

*Experiment 2: effect of body size on density.* To manipulate body size while keeping

density constant, we started 15 microcosms (as described before) at Day -7 ('Day -7 jars'

henceforth), and another 15 microcosms at Day -2 ('Day -2 jars' henceforth). At Day 0, we

removed and filtered, as before, half of the volume (100 mL) in all jars. We then replenished all

jars with filtered medium from jars in the other group, so all jars contained a mixture of equal

parts medium and bacteria (resources) from Day -2 and Day -7 jars (Fig 1c). Simultaneously, we

filtered an additional amount of medium from Day -7 jars to ensure that the cell density in this

group, once all growth medium had been added, matched that of day -2 jars (Fig 1c). This was

done by calculating the volume of growth medium of the original Day -7 jars that needed to be

filtered based on the observed density (after manipulation) in Day -2 jars. Because *T. pyriformis*

decays in size as it grows to carrying capacity (Fig 2a, b), Day -7 jars contained, on average,

smaller individuals, Day -2 jars contained relatively larger individuals (Fig 1c), but both groups

had the same population density, medium, and bacterial density (resources).

We used the FlowCam to quantify body size and density in Day 0 (the day the

manipulations where made) and Day 2 (i.e., two days later). In both experiments, we expected

density and size to change over time (from Day 0 to Day 2), meaning that statistically speaking,

we expect time to influence both density and size. However, if either size or density influence the

other, we also expect the interaction between time and density (experiment 1), or time and size

(experiment 2), to be significant. In that case, the interaction term indicates by how much a

difference in starting density or size influences the change in the response variable as time

elapses. A large (small) interaction term would indicate a large (small) effect of the initial

difference in either size or density on the response variable. We tested for these interactions

using linear models in R v3.6.1 (R Core Team 2013) with either 1) body size as a response

variable and density, day, and their interaction as predictors (experiment 1) or 2) density as a

response variable and body size, day, and their interaction as predictors (experiment 2). To

compare the effect sizes of density on size, and size on density, we standardized all variables in

R package *effectsize* v0.6.01 (Ben-Shachar, Lüdecke & Makowski 2020) by re-centering and re-

scaling variables to a normal distribution with mean equal to zero and standard deviation equal to

1 (Gelman 2008).

*Mathematical models*

While others have argued that laboratory cultures should have low levels of heritable genetic

variation (DeLong, Hanley & Vasseur 2014), *T. pyriformis* has mechanisms to maintain larger

than expected levels of genetic variation (Dimond & Zufall 2016). Because of this, and to assess

whether observed changes in body size were more likely due to plasticity or rapid evolution, we

fitted two possible models that track change in the abundance and average body size of a

population, *N*, as it grows logistically towards a carrying capacity *K* with intrinsic growth rate *r*.

Following previous work (Abrams 1977; Abreu *et al.* 2019; Lax, Abreu & Gore 2020;

Wieczynski *et al.* 2021), we included an additional mortality term in the ecological dynamics,

$mN$, to account for regular loss of individuals from the population through sampling. This

additional mortality term has been shown to better describe the ecological dynamics of a

microbial microcosm with frequent sampling, like ours (Abreu *et al.* 2019; Lax, Abreu & Gore

2020). Furthermore, we assumed that the intrinsic growth rate and the carrying capacity were

functions of (average) body size, *M*, of the forms $r = aM^\alpha$ and $K = bM^\gamma$, following well-known

allometric relationships (Damuth 1981; Savage *et al.* 2004). Taken together, the ecological

baseline model was thus written as:

$$\frac{dN}{dt} = aM^\alpha \left(1 - \frac{N}{bM^\gamma}\right) - mN. \qquad \text{eq [1]}$$

To incorporate coupling between ecological dynamics (changes in N) and both plastic

and evolutionary changes in *M*, we used three alternative model formulations to track changes in

*M*: the first assumed that only plastic change in *M* could occur (Plasticity Model), the second

assumed that only evolutionary change in *M* could occur (Eco-Evolutionary Model), while the

third allowed both processes to occur simultaneously (Plasticity + Eco-Evo Model).

The Plasticity Model modifies the existing Supply-Demand model for body size

dynamics (DeLong 2012; DeLong & Walsh 2015) and assumes that the rate of change in body

size, $\frac{dM}{dt}$, increases with energy intake (supply), and decreases with energy loss (demand).

Following a recent study (DeLong 2020), we assumed that the supply in a species growing

logistically depends on the ratio of the carrying capacity, K, and the abundance of the species, N,

times a conversion rate constant $e$, that converts the supply to units of $M$. When the population is

small, the supply approaches infinity, and it approaches $e$ when N grows to K. Following

previous studies (DeLong, Hanley & Vasseur 2014), we also assumed that the demand was the

metabolic cost of the organism, which is known to increase allometrically with body size, as

$cM^{\delta}$. Taken together, the equation controlling changes in body size M was:

$$\frac{dM}{dt} = \frac{eK}{N} - cM^{\delta}. \qquad \text{eq [2]}$$

For the Eco-Evolutionary Model, we followed previous studies on eco-evolutionary

dynamics (e.g., (Abrams, Harada & Matsuda 1993; Abrams & Matsuda 1997; Jones *et al.* 2009;

Ellner & Becks 2010; Jones & Gomulkiewicz 2012; Cortez 2016; Cortez 2018)) to incorporate

evolution in $M$ at a rate that equals the product of the total amount of additive heritable variation

in body size (*i.e.*, the product of the total phenotypic variance, $\sigma^2$, and the narrow sense

heritability, $h^2$), and the selection gradient (*i.e.,* the change in fitness, $F$, with respect to a change

in $M$, which represents the strength of selection acting on $M$). Taken together, the equation

controlling the change in M over time under these assumptions was:

$$\frac{dM}{dt} = \sigma^2 h^2 \frac{\partial F(M)}{\partial M}, \qquad \text{eq [3]}$$

were $F(M) = \frac{1}{N}\frac{dN}{dt}$ (Lande 1976; Schreiber, Bürger & Bolnick 2011).

Last, the Plasticity + Eco-Evo Model assumes that both Supply-Demand (plastic) and

Eco-Evo (rapid evolution) contributions can simultaneously influence the rate of change of $M$,

resulting in the following model for body size dynamics:

$$\frac{dM}{dt} = \frac{eK}{N} - cM^\delta + \sigma^2 h^2 \frac{\partial F(M)}{\partial M}.$$ eq [4]

This model does not account for possible interplay between plastic and evolutionary change,

such as plasticity facilitating evolution, plasticity impeding evolution, or evolving plasticity, all

of which can occur and have been reviewed elsewhere (Diamond & Martin 2016).

None of our models accounts for shifts in age or size structure because here we were are

specifically interested in mathematically tracking changes in mean body size, not changes in the

entire trait distribution, which also requires model formulations that are not amenable to ODE

fitting (Chen, Baños & Buceta 2018; Nieto-Acuña *et al.* 2019).

*Model fitting, parameter uncertainty, and model selection*

We fitted the models in Eqs 1-4 to the *T. pyriformis* time series using R package FME

v1.3.6.1 (Soetaert & Petzoldt 2010). However, non-linear model fitting tends to get stuck in sub-

optimal maxima/minima during residual minimization (or similar procedures) and it is often

impossible to simultaneously estimate all model parameters, in which case the model is said to

be non-identifiable (Motulsky & Christopoulos 2004; Miao *et al.* 2011). The conversion

parameters $a$ and $b$ of our models –which convert from units of $M^\alpha$ and $M^\gamma$ into units of $r$ and

$K$, respectively– were not identifiable (i.e., the fitting procedure could not simultaneously

estimate them and all other model parameters without yielding negative or other non-sensical

parameter values). Because initial model fits suggested values close to 1 and -1 for the allometric

parameter $\alpha$ and $\gamma$, respectively, we estimated $a$ and $b$ from our data by solving $aM^\alpha$ and

$bM^\gamma$ using the observed intrinsic growth rates for the first two days of growth (r ~ 3.20 day-1),

the observed average $K \sim 6400$ ind/mL, average $M$ ($\sim 10^4 \mu m^3$) obtained from the FlowCam from

day 0 to day 4, and setting $\alpha = 1$ and $\gamma = -1$. Doing so resulted in initial parameter values of

$\sim 10^{-4}$ for $a$ and $\sim 10^{7}$ for $b$, which were then optimized during preliminary model fitting (i.e., the

iterative process of providing initial parameter guesses and assessing model fit to increase the

chance that the fitting procedure succeeds). Because parameters $a$ and $b$ do not play an important

biological role –they convert units of body size (to the power of $\alpha$ or $\gamma$) into units of $r$ or $K$–

these parameters should not be expected to change across models, and were thus assumed to be

equal for all models and set constant during model parameter and uncertainty estimation of all

remaining parameters. Last, we assumed that the scaling parameter of the metabolic cost, $\delta$,

equaled 1, as has been shown to be the case for protists (DeLong *et al.* 2010), despite it being

closer to ¾ for metazoans (Gillooly *et al.* 2001; Brown *et al.* 2004; DeLong *et al.* 2010).

Fitting of ordinary differential equations in package FME relies on the Levenberg-

Marquardt algorithm for parameter estimation, and a Metropolis-Hastings MCMC procedure for

estimation of parameter uncertainty (Soetaert & Petzoldt 2010). Model comparison was done

using Akaike Information Criterion (Burnham & Anderson 2002) as:

$$AIC_i = 2k_i + n_N \ln(SSE_{i,N}) + n_M \ln(SSE_{i,M}) \ , \hspace{3cm} \text{eq[4]}$$

where $k_i$ is the number of parameters of model $i$, $n_N$ is the number of datapoints considered in

the density time series, $n_M$ is the number of datapoints considered in the body size time series,

$SSE_{i,N}$ is the sum of squared errors of model $i$ with respect to the density time series, and $SSE_{i,M}$

is the sum of squared errors of model $i$ with respect to the body size time series. An AIC

difference >2 indicates that one model is significantly better than the alternative (Burnham &

Anderson 2002). Akaike weights were calculated as $e^{-0.5\Delta AIC}$, where $\Delta AIC = AIC_i -$

$\min(AIC)$. Weights are bound between 0 and 1 and models with larger weights can be

interpreted as having larger relative likelihoods (Burnham & Anderson 2002). All data and code

are available at https://anonymous.4open.science/r/Tetra_Rapid_BodySize_Shifts-5FD7.

RESULTS

*Time series analyses*

Tetrahymena abundances increased to carrying capacity roughly 4 days after microcosms

where initialized (Fig 2a, Fig S1a). Body size increased over the first day, then decreased more

or less continuously for 12-13 days (Fig 2b, Fig S1b). The CCM analysis showed large

predictability values for densities using changes in body size as a predictor, but smaller

predictability for body size using densities as the predictor (Fig 2c). The stronger effect of

change in body size on density was found to be causal (sharp increase in predictability with

library size and decrease in standard deviation, Fig 2c). The effect of density on body size

seemed to only be weakly causal (slow convergence of predictability and little change in

standard deviation, Fig 2c). This indicated that while changes in body size more strongly

influenced density changes, both seemed to have at least some level of influence on each other.

*Experimental results*

Our density manipulation resulted in a roughly two-fold difference in density among

experimental jars at Day 0 ($p < 10^{-5}$, Appendix Fig S2), while the manipulation of body size

resulted in a 20% size difference in average body size among experimental jars ($p < 10^{-8}$,

Appendix Fig S3). A Tukey post-hoc test indicated no statistically significant differences in body

size between high- and low-density jars at Day 0 (p=0.15, Fig 3a), and no statistically significant

differences in density between large- and small-size treatments at Day 0 (p=0.64, Fig 3b).

Together, these results indicate that we correctly manipulated density and size while keeping the

other variable constant. Overall, density increased, and size decreased from Day 0 to Day 2

(Table 1, Fig 3), consistent with trends observed in our time series (Fig 2).

The imposed initial differences in density and size resulted in significant interactions with

time (Table 1, Fig 3), indicating significant reciprocal effects of size on density and of density on

size. Consistent with results from our CCM analysis (Fig 2c), body size had a larger effect on

density (in magnitude) than the other way around (Table 1). However, in standardized units

(units of standard deviations, SD), it could be possible for the imposed initial difference in

density to be smaller relative to that in size, which could have led to a smaller overall effect of

density on size than the other way around. To control for that effect, we divided the observed

effect size of density on body size by the imposed (standardized) initial differences in density

between low- and high-density jars, and also divided the observed effect size of body size on

density by the imposed (standardized) initial difference in body size between small and large size

jars. The resulting number could then be interpreted as the magnitude of the effect the predictor

variable had on the change observed in the response variable from Day 0 to Day 2 (in units of

SD), per unit difference in the initial treatment (also in SD). This resulted in a standardized effect

size of body size on density that, while much closer, was still larger in magnitude than that of the

effect of density on size (in absolute values, density→size = 0.57, size→density =0.64), still

consistent with our CCM results.

*Mathematical models*

All models fitted the empirical data remarkably well (Fig 4, Table 2). MCMC chains

converged for all fitted parameters (Appendix Fig S4-S9) and model parameters were free of

correlations for 25 out of 31 total parameter pairs across the three models (except for $\gamma$ and *m,*

and c and e in the Plasticity Model, $\alpha$ and $m$ in the eco-evo model, and c, e and $\sigma^2 h^2$, in the

Plasticity+Eco-Evo model, Fig S5, S7, and S9). The Plasticity and Plasticity+Eco-Evo Models fit

the abundance data slightly better than the Eco-Evo Model (notice departures of Eco-Evo Model

in the early time steps, Fig 4a, c, e), while the Plasticity and Eco-Evo Models fit the body size

data better than the Plasticity + Eco-Evo Model (Fig 4b, d, f). All models arrived at very similar

fitted values for model parameters they had in common (Table 2), indicating good agreement

between them all. Consistent with the literature, all models identified the scaling of K and M, $\gamma$,

as a negative number close to -¾ (Table 2). However, all models identified the scaling between r

and M, $\alpha$, as positive and close to 1 (Table 2), while the literature pins that value—across

species—to -¼ (Savage *et al.* 2004).

Despite all models fitting the data well, model selection through AIC indicated very large

differences in model likelihood, with the Plasticity Model being –by far– the most likely

(AIC$_{weight}$ ~ 1 for the Plasticity Model, but effectively zero for the other two models, Table 2).

The Plasticity + Eco-Evo Model was the least likely of all fitted models, perhaps owing to a

larger number of model parameters, which are penalized by AIC (Table 2). This result thus

suggests that the observed coupled density and body-size dynamics were mostly driven by

plasticity, while rapid evolution or a combination of plasticity and rapid evolution are less likely

to explain the observed dynamics.

DISCUSSION

Because of the myriad ecological consequences of body size (e.g. (Gillooly *et al.* 2001;

Brown *et al.* 2004; DeLong *et al.* 2010)), it is important to understand how changes in body

size—plastic or evolutionary—may influence, or be influenced by, ecological dynamics. We

show that changes in body size more strongly influence changes in density than the other way

around (Fig 2c, 3b), but that density also influences changes in body size (Fig 2c, 3a). This

suggests the existence of a (possibly asymmetric) feedback between the two. Additionally, a

model that accounts for rapid plastic change in body size provides the most parsimonious

explanation for the observed, coupled ecological and phenotypic dynamics (Fig 4, Table 2).

Previous results indicated that phenotypic change often lags ecological change (e.g., (DeLong *et*

*al.* 2016)) but that, under certain conditions, very rapid shifts in body size may precede important

changes in ecological dynamics and can thus be used as early warning signals of state shifts

(Clements and Ozgul 2015). Our results add to this literature by showing that phenotypic change

not only occurs well within ecological timescales and responds to ecological dynamics, but may

even causally influence those dynamics (Figs 2, 3).

Understanding the mechanisms of this possible feedback between size and density

dynamics is central to gain insights as to how coupled eco-phenotypic dynamics may occur in

the wild. In many unicellular organisms, cell growth (increase in body size) at the individual

level and cell division are intimately intertwined: cells grow until a critical size is reached, which

triggers DNA synthesis and eventual division (Baserga 1968). Larger cells are closer to the

critical size threshold that triggers cell division (Jorgensen & Tyers 2004), leading to faster cell

division (reproduction) in the next generation, which ultimately results in faster population

growth. This link between size and cell division provides a possible explanation for why our

results identify changes in size as important drivers of changes in density (Fig 2c, Fig 3). In line

with this argument, all models predicted a positive relationship between body size and the

intrinsic growth rate, r ($\alpha \sim 1$, Table 2). This result stands in contrast to empirical data across

species and theoretical expectations, which show lower intrinsic growth rates for larger

organisms (or $\alpha < 0$, (Savage *et al.* 2004)). Within species, however, larger individuals typically

reproduce more and die at lower rates (Peters 1983), leading to higher *r*, due to lower mortality

and higher reproduction (Kingsolver & Huey 2008). This positive relationship between body size

and the demographic processes that fuel intrinsic growth rates are well understood within species

(Peters 1983; Kingsolver & Huey 2008) –emphasizing how inter-species and intra-species body

size scaling may often differ (Rall *et al.* 2012)– but also providing a plausible mechanism

through which changes in size may be causally influencing changes in density, as our results

show.

On the flip side, cells can enter and exit the cell division cycle depending on internal and

external cues, such as nutrient availability (Baserga 1968; Fukada *et al.* 2007). As the *T.*

*pyriformis* population reaches carrying capacity, low resource levels likely cue cells to exit the

cell division cycle, resulting, in turn, in stunted growth and reduced average body sizes (because

cells grow to reproduce) thus providing a possible explanation as to how density may influence

body size. If that is the case, then observed total standing phenotypic variation in our population

should be largely non-heritable, as also suggested in a recent study (Jacob & Legrand 2021).

Surprisingly, the Eco-Evolutionary and Plasticity + Eco-Evo Models support this idea, as they

indicate that the total amount of standing heritable variation in body size is rather small ($\sigma^2 h^2 =$

0.992 with high confidence for the Eco-Evo model and $\sigma^2 h^2 = 0.878$ for Plasticity + Eco-Evo

although with very low confidence, Table 2). For comparison, the total amount of phenotypic

variation (heritable or not) in our initial population was 94 (units of $\mu m^3$ squared), so the

heritable portion of that total phenotypic variation would be on the order of 1%. Shifts in *T.*

*pyriformis* phenotype have also been shown to occur differentially across environmental

conditions (DeLong *et al.* 2017; Weber de Melo *et al.* 2020), again suggesting the occurrence of

plasticity and very low heritability in this species, and providing support for the above

mechanism of response of body size on density.

While the Eco-Evo and Plasticity + Eco-Evo Models were found to be less parsimonious,

our results do not rule out the possibility of rapid evolution in this system. Indeed, plastic change

often precedes evolutionary change and occurs mostly along the axes of variation with the largest

amount of heritable variation (the classic evolutionary 'path of least resistance' (Lande 1976;

Lande 1979; Lande & Arnold 1983)), thus setting the stage for evolution to occur along those

axes (Noble, Radersma & Uller 2019). As we state in our methods, our current models do not

account for such a scenario, which could very well be at play here. Indeed, despite low

heritability, the less parsimonious Eco-Evo and Plasticity + Eco-Evo models suggest that short

term selection imposed by density-dependence may be strong enough to consistently shift body

size over time, which in turn influences population dynamics. Our own data indicate that *T.*

*pyriformis* reproduces at a rate of 3.5-4 new individuals per individual per day. This extremely

fast population growth may eventually allow for evolutionary change in body size—provided

that selection is strong enough, because of low heritability—even if it lags behind plastic change

(Chevin, Lande & Mace 2010; Fox *et al.* 2019).

Interestingly, neither water nor nutrients were replenished during our experiment; both

were limited and were likely consistently lost from the system through sampling and respiration.

This nutrient impoverishment should lead to a strong decline in carrying capacity over time.

Such a decline was not, or was only very weakly, observed (Fig 4). However, shifting the focus

from abundances to total biomass shows a different picture: biomass increased with abundance

in the first few days, but then declined over time (Appendix Fig S10) likely due to density being

roughly constant after day 4 but average body size declining consistently after day 2 (Fig 2b, Fig

4, Fig S10). As resources wane from the system, a rapid decline in nutrient concentration may

therefore be selecting for smaller body size (Vanni *et al.* 2009), which results in lower total

biomass, but also lower competition (through a reduction in metabolic needs associated with

smaller size (Brown *et al.* 2004)), ultimately allowing the population to remain at very high

densities despite waning resources and increasingly lower biomass. Our data thus suggest

interesting ways in which a rapid plastic changes in body size may allow organisms to regulate

population growth and density-dependent factors, even as nutrients become increasingly limited.

*Caveats and concluding remarks*

Both the CCM and the experimental results agreed that changes in size had a larger effect

on changes in density than the other way around (Fig 2c, Fig 3). Yet, they differed on how much

stronger this effect of size on density is. This difference may be due to a couple of reasons. First,

CCM infers the magnitude of causal effects of one variable on the other throughout the entire

time series. It does not inform at what time, exactly, the effect of one variable is larger than the

effect of the other variable (Sugihara *et al.* 2012). So, it could very well be that the effect of

density is much larger in the first few days –when density changes the most– but then declines

over time. Second, our manipulations of size and density likely cannot be extrapolated beyond

the first few days of the ecological dynamics (Days 0 to 7), and those first few days coincide

with the time span over which larger changes in density were observed. So, while the CCM may

underestimate large, temporally localized effects of density (by looking at the entirety of the time

series), our experimental work may be overestimating the overall effects of density (as it focuses

on their possibly larger effects in the first few days of the dynamics) even though, taken together,

both results agree qualitatively.

Because neither CCM nor the experimental work should be used to infer how the effect

of size and density on one another may change over time, it is entirely possible, even likely, that

both effects change in magnitude over time. This likely explains why, despite body size clearly

having an effect on density (Fig 2, 3), density changed little after Day 6 while body size declined

from Day 2 to Day 14 (Fig 2b). Indeed, as cells are cued into exiting the cell division cycle due

to lack of nutrients, fewer and fewer of them remain reproductive and the proposed mechanism

through which cell size may influence density (i.e., through its effect on reproduction) may

decline as the population remains at high density and resource scarcity sets in. Taken together,

while our results indicate the existence of a feedback between changes in size and changes in

density, they also suggest the possibility that the magnitude –and perhaps even the direction of

that feedback– may change over time, certainly a promising avenue for future research.

Overall, our study shows that feedbacks between rapid plastic change in body size and

change in density are likely integral to the process of population growth itself. This study sheds

light on the ecological and evolutionary constraints that regulate population growth and provides

new insights about how organisms cope with the negative effects of density-dependence. Our

results also emphasize the need to further study and understand the ecological consequences of

rapid plastic phenotypic change (Yamamichi, Yoshida & Sasaki 2011; Tariel, Plénet & Luquet

2020), as plasticity, particularly in body size, may play a crucial role in determining the fate of

networks of species interactions in a warming world (Barbour & Gibert 2021; Jacob & Legrand

2021).

Table 1: ANOVA results from experimental manipulations of density and size

|  |  |  | Standardized Estimates (95% C.I.) | **p-val** |
|---|---|---|:---:|:---:|
| **(a)** | **Model** | Body Size ~ Density*Time | — | — |
|  | **Parameters** | Intercept | 0.96 (0.76, 1.16) | **<$10^{-16}$** |
|  |  | Density (Low Density) | -0.17 (-0.45, 0.12) | 0.244 |
|  |  | Time | -2.21 (-2.49, -1.19) | **<$10^{-16}$** |
|  |  | Density*Time | 0.90 (0.51, 1.30) | **$2.92*10^{-5}$** |
| **(b)** | **Model** | Density ~ Body Size*Time | — | — |
|  | **Parameters** | Intercept | -0.63 (-0.82, -0.45) | **<$10^{-16}$** |
|  |  | Density | -0.28 (-0.54, -0.02) | **0.036** |
|  |  | Time | 2.13 (1.87, 2.39) | **<$10^{-16}$** |
|  |  | Body Size*Time | -1.17 (-1.53, -0.80) | **$3.79*10^{-8}$** |

Table 2: Parameter estimates, parameter uncertainty, and model selection for all fitted models.

| | | $r$ scaling with body size ($\alpha$) | $K$ scaling with body size ($\gamma$) | Mortality rate ($m$) | Supply-Demand conversion parameter ($e$) | Supply-Demand loss parameter ($c$) | Heritable Variation ($\sigma^2 h^2$) | | Allometric Intercept for $r$ ($a$) | Allometric Intercept for $K$ ($b$) | AIC ($\Delta$AIC) | AIC$_{weight}$ |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **(a) Plasticity Model** | Estimate | 1.022 | -0.721 | 0.969 | 24.295 | 0.087 | — | Initial value | $\sim 10^{-4}$ | $\sim 10^{7}$ | **3242.5 (0)** | **~1** |
| | 95% Credible Interval | (1.010, 1.023) | (-0.735, -0.720) | (0.770, 0.991) | (18.400, 26.714) | (0.077, 0.091) | — | Post-Optimization | $2.2*10^{-4}$ | $8.2*10^{6}$ | | |
| **(b) Eco-Evo Model** | Estimate | 1.056 | -0.748 | 0.897 | — | — | 0.922 | Initial value | As above | As above | **3287.65 (+45.15)** | **1.57*10$^{-10}$** |
| | 95% Credible Interval | (1.048, 1.064) | (-0.753, -0.743) | (0.770, 1.026) | — | — | (0.825, 1.193) | Post-Optimization | As above | As above | | |
| **(c) Plasticity + Eco-Evo Model** | Estimate | 1.012 | -0.726 | 0.860 | 24.394 | 0.086 | 0.878 | Initial value | As above | As above | **3301.2 (+58.72)** | **1.77*10$^{-13}$** |
| | 95% Credible Interval | (1.008, 1.023) | (-0.734, -0.722) | (0.766, 0.994) | (14.603, 32.223) | (0.073, 0.096) | (-0.498, 0.384) | Post-Optimization | As above | As above | | |

528 FIGURES

Fig 1: (a) Depiction of experimental procedure: microcosms were initialized at 10ind/mL and
each microcosm was replicated six times. Population density and cell size were recorded daily
for 14 days thereafter using fluid imaging. (b) Experimental setup to manipulate density while
controlling for body size. This involved filtering the Tetrahymena out of 100mL of media in half
of the experimental jars. (c) Experimental setup to manipulate body size while controlling for
density. This involved the filtration of 100mL of media of Day-2 (D-2) and Day-7 (D-7) jars,
which was then added to jars of the other group (ensuring equal resources and media). An extra
filtration step ensured that the density in 100mL of Day-2 jars equaled the density in 100mL of
Day-7 jars. To find the volume ($x$) to be filtered from 100mL of unfiltered Day-7 media (then
returned to Day-7 jars), we noticed that $Dens_{D-2} = Dens_{D-7}$ if $\frac{Count_{D-2}}{Vol_{D-2}} = \frac{Count_{D-7}}{Vol_{D-7}}$, and that
$Vol_{D-2} = 100mL$ and $Vol_{D-7} = 100mL - x$. Solving for $x$ yielded how much volume had to
be filtered and returned to the same jars. This extra filtering step was done first and the filtrate
was set aside before the other two steps.

545    Fig 2: (a) Time series of Tetrahymena density. (b) Time series of mean Tetrahymena body size.

546    (c) Convergent Cross Mapping predictability plot against library size (i.e., length of the time

547    series used for analysis), using body size to predict density (purple) or density to predict body

548    size (yellow), repeated 1000 times. Solid line indicates mean values and dashed lines indicate

549    standard deviation of the mean.

Fig 3: (a) Boxplot of the response of body size over two days across high- and low-density treatments. Solid black lines connecting the bodies of the boxplots are the to help see the interactive effect of the predictor variable and time. (b) Boxplot of the response of density over two days across large- and small-body-size treatments. Solid lines as in (b). N.S. indicates that the manipulation of density (a) and size (b) did not significantly alter the other (size in (a) and density in (b)) at Day 0.

576 Fig 4: (a) Density data (grey points) and Plasticity Model fit (solid blue). Uncertainty is

577 represented as 700 model predictions (transparent lines) whose parameters were sampled from

578 posterior distributions for each model parameter, estimated during model fitting (Table 2). (b) As

579 in (a), but for body size data. (c–d, e–f) As in (a–b) but for the Eco-Evo Model fit (c–d) or the

580 Plasticity + Eco-Evo Model fit (e–f).

REFERENCES

Abrams, P.A. (1977) Density-Independent Mortality and Interspecific Competition: A Test of
Pianka's Niche Overlap Hypothesis. *The American Naturalist,* **111,** 539-552.
Abrams, P.A., Harada, Y. & Matsuda, H. (1993) On the Relationship between Quantitative
Genetic and Ess Models. *Evolution,* **47,** 982-985.
Abrams, P.A. & Matsuda, H. (1997) Prey Adaptation as a Cause of Predator-Prey Cycles.
*Evolution,* **51,** 1742-1750.
Abreu, C.I., Friedman, J., Andersen Woltz, V.L. & Gore, J. (2019) Mortality causes universal
changes in microbial community composition. *Nature Communications,* **10,** 2120.
Amarillo-Suarez, A.R., Stillwell, R.C. & Fox, C.W. (2011) Natural selection on body size is
mediated by multiple interacting factors: a comparison of beetle populations varying
naturally and experimentally in body size. *Ecology and Evolution,* **1,** 1-14.
Anderson-Teixeira, K.J., Vitousek, P.M. & Brown, J.H. (2008) Amplified temperature
dependence in ecosystems developing on the lava flows of Mauna Loa , Hawai ' i.
*Proceedings of the National Academy of Sciences,* **105,** 228-233.
Barbour, M.A. & Gibert, J.P. (2021) Genetic and plastic rewiring of food webs under climate
change. *Journal of Animal Ecology.*
Baserga, R. (1968) Biochemistry of the cell cycle: a review *Cell Tissue Kinetics,* **1,** 167-191.
Becks, L., Ellner, S.P., Jones, L.E. & Hairston, N.G. (2012) The functional genomics of an eco-
evolutionary feedback loop: Linking gene expression, trait evolution, and community
dynamics. *Ecology Letters,* **15,** 492-501.
Ben-Shachar, M., Lüdecke, D. & Makowski, D. (2020) effectsize: Estimation of Effect Size Indices
and Standardized Parameters. *Journal of Open Source Software,* **5**.
Berlow, E.L., Dunne, J.A., Martinez, N.D., Stark, P.B., Williams, R.J. & Brose, U. (2009) Simple
prediction of interaction strengths in complex food webs. *Proceedings of the National
Academy of Sciences of the United States of America,* **106,** 187-191.
Brose, U., Jonsson, T., Berlow, E.L., Warren, P.H., Banasek-Richter, C., Bersier, L.-F., Blanchard,
J.L., Brey, T., Carpenter, S.R., Cattin, M.-F., Cushing, L., Hassan, A.D., Dell, A.I., Edwards,
F., Harper-Smith, S., Jacob, U., Ledger, M.E., Martinez, N.D., Memmott, J., Mintenbeck,
K., Pinnegar, J.K., Rall, B.C., Rayner, T.S., Reuman, D.C., Ruess, L., Ulrich, W., Williams,
R.J., Woodward, G. & Cohen, J.E. (2006) Consumer–resource body-size relationships in
natural food webs. *Ecology,* **87,** 2411-2417.
Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004) Toward a Metabolic
Theory of Ecology. *Ecology,* **85,** 1771-1789.
Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference: A practical
information-theoretic approach,* 2nd ed. edn. Springer Science, New York.
Cameron, M.D., Grant, V.H., Joly, K., Schmidt, J.H., Gustine, D.D., Mangipane, L.S., Mangipane,
B. & Sorum, M.S. (2020) Body size plasticity in North American black and brown bears.
*Ecosphere,* **e03235,** 1-11.
Chelini, M.C., Delong, J.P. & Hebets, E.A. (2019) Ecophysiological determinants of sexual size
dimorphism: integrating growth trajectories, environmental conditions, and metabolic
rates. *Oecologia,* **191,** 61-71.

Chen, Y., Baños, R. & Buceta, J. (2018) A Markovian Approach towards Bacterial Size Control
and Homeostasis in Anomalous Growth Processes. *Scientific Reports,* **8,** 9612.

Chevin, L.M., Lande, R. & Mace, G.M. (2010) Adaptation, plasticity, and extinction in a changing
environment: towards a predictive theory. *PLoS Biology,* **8,** e1000357.

Clark, A.T., Ye, H., Isbell, F., Deyle, E.R., Cowles, J., Tilman, G.D. & Sugihara, G. (2015) Spatial
convergent cross mapping to detect causal relationships from short time series. *Ecology,*
**96,** 1174-1181.

Clements, C.F. & Ozgul, A. (2016) Including trait-based early warning signals helps predict
population collapse. *Nature Communications,* **7**.

Cortez, M.H. (2016) How the Magnitude of Prey Genetic Variation Alters Predator-Prey Eco-
Evolutionary Dynamics. *The American Naturalist,* **188,** 329-341.

Cortez, M.H. (2018) Genetic variation determines which feedbacks drive and alter predator–
prey eco-evolutionary cycles. *Ecological Monographs,* **88,** 353-371.

Cox, S.L., Ruff, C.B., Maier, R.M. & Mathieson, I. (2019) Genetic contributions to variation in
human stature in prehistoric Europe. *Proceedings of the National Academy of Sciences,*
**116,** 21484-21492.

Damuth, J. (1981) Population density and body size in mammals. *Nature,* **290,** 699-700.

David, J.R., Legout, H. & Moreteau, B. (2006) Phenotypic plasticity of body size in a temperate
population of Drosophila melanogaster: when the temperature-size rule does not apply.
*Journal of genetics,* **85,** 9-23.

Davidowitz, G., Roff, D.A. & Nijhout, H.F. (2005) A Physiological Perspective on the Response of
Body Size and Development Time to Simultaneous Directional Selection. *Integrative and*
*Comparative Biology,* **45,** 525-532.

DeLong, J.P. (2012) Experimental demonstration of a ' rate – size ' trade-off governing body size
optimization. *Evolutionary Ecology Research,* **14,** 343-352.

DeLong, J.P. (2020) Detecting the Signature of Body Mass Evolution in the Broad-Scale
Architecture of Food Webs. *The American Naturalist,* **196,** 443-453.

DeLong, J.P., Brassil, C.E., Erickson, E.K., Forbes, V.E., Moriyama, E. & Rieckhof, W.R. (2017)
Dynamic thermal reaction norms and body size oscillations challenge explanations of
the temperature–size rule. *Evolutionary Ecology Research,* **18,** 293-303.

DeLong, J.P., Forbes, V.E., Galic, N., Gibert, J.P., Laport, R.G., Phillips, J.S. & Vavra, J.M. (2016)
How fast is fast? Eco-evolutionary dynamics and rates of change in populations and
phenotypes. *Ecology and Evolution,* **6,** 573-581.

DeLong, J.P. & Gibert, J.P. (2016) Gillespie eco-evolutionary models (GEMs) reveal the role of
heritable trait variation in eco-evolutionary dynamics. *Ecology and Evolution,* **6,** 935-
945.

DeLong, J.P., Gilbert, B., Shurin, J.B., Savage, V.M., Barton, B.T., Clements, C.F., Dell, A.I., Greig,
H.S., Harley, C.D.G., Kratina, P., McCann, K.S., Tunney, T.D., Vasseur, D.A. & O'Connor,
M.I. (2015) The Body Size Dependence of Trophic Cascades. *The American Naturalist,*
**185,** 354-366.

DeLong, J.P., Hanley, T.C. & Vasseur, D.A. (2014) Predator-prey dynamics and the plasticity of
predator body size. *Functional Ecology,* **28,** 487-493.

DeLong, J.P., Okie, J.G., Moses, M.E., Sibly, R.M. & Brown, J.H. (2010) Shifts in metabolic scaling,
production, and efficiency across major evolutionary transitions of life. *Proceedings of*
*the National Academy of Sciences of the United States of America,* **107,** 12941-12945.
DeLong, J.P. & Walsh, M. (2015) The interplay between resource supply and demand
determines the influence of predation on prey body size. *Canadian Journal of Fisheries*
*and Aquatic Sciences,* **72,** 1-7.
Diamond, S.E. & Martin, R.A. (2016) The interplay between plasticity and evolution in response
to human-induced environmental change. *F1000 Research,* **5,** 2835.
Dimond, K.L. & Zufall, R.A. (2016) Hidden genetic variation in the germline genome of
Tetrahymena thermophila. *Journal of Evolutionary Biology,* **29,** 1284-1292.
Ellner, Stephen P. & Becks, L. (2010) Rapid prey evolution and the dynamics of two-predator
food webs. *Theoretical Ecology***,** 133-152.
Ellner, S.P., Geber, M.A. & Hairston, N.G. (2011) Does rapid evolution matter? Measuring the
rate of contemporary evolution and its impacts on ecological dynamics. *Ecology Letters,*
**14,** 603-614.
Fey, S.B., Gibert, J.P. & Siepielski, A.M. (2019) The consequences of mass mortality events for
the structure and dynamics of biological communities. *Oikos***,** 1-12.
Fox, R.J., Donelson, J.M., Schunter, C., Ravasi, T. & Gaitan-Espitia, J.D. (2019) Beyond buying
time: the role of plasticity in phenotypic adaptation to rapid environmental change.
*Philosophical Transactions of the Royal Society B: Biological Sciences,* **374,** 20180174.
Frickel, J., Sieber, M. & Becks, L. (2016) Eco-evolutionary dynamics in a coevolving host-virus
system. *Ecology Letters,* **19,** 450-459.
Fukada, S., Uezumi, A., Ikemoto, M., Masuda, S., Segawa, M., Tanimura, N., Yamamoto, H.,
Miyagoe-Suzuki, Y. & Takeda, S. (2007) Molecular signature of quiescent satellite cells in
adult skeletal muscle. *Stem Cells,* **25,** 2448-2459.
Galloway, L.F. & Etterson, J.R. (2007) Transgenerational plasticity is adaptive in the wild. *Science*
*(New York, N.Y.),* **318,** 1134-1136.
Gavina, M.K.A., Tahara, T., Tainaka, K.I., Ito, H., Morita, S., Ichinose, G., Okabe, T., Togashi, T.,
Nagatani, T. & Yoshimura, J. (2018) Multi-species coexistence in Lotka-Volterra
competitive systems with crowding effects. *Scientific Reports,* **8,** 1198.
Gelman, A. (2008) Scaling regression inputs by dividing by two standard deviations. *Statistics in*
*Medicine,* **27,** 2865-2873.
Ghosh, S.M., Testa, N.D. & Shingleton, A.W. (2013) Temperature-size rule is mediated by
thermal plasticity of critical size in Drosophila melanogaster. *ZooKeys,* **298**.
Gibert, J.P. & DeLong, J.P. (2014) Temperature alters food web body-size structure. *Biology*
*Letters,* **10,** 20140473-20140473.
Gibert, J.P. & Yeakel, J.D. (2019) Eco-Evolutionary Origins of Diverse Abundance , Biomass , and
Trophic Structures in Food Webs. *Frontiers in Ecology and Evolution,* **7,** 1-11.
Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001) Effects of size and
temperature on metabolic rate. *Science (New York, N.Y.),* **293,** 2248-2251.
Glasheen, J. & McMahon, T. (1996) Size-dependence of water-running ability in basilisk lizards
(Basiliscus basiliscus). *The Journal of Experimental Biology,* **199,** 2611-2618.

Gravel, D., Poisot, T., Albouy, C., Velez, L. & Mouillot, D. (2013) Inferring food web structure
from predator-prey body size relationships. *Methods in Ecology and Evolution,* **4,** 1083-
1090.
Hairston, N.G., Ellner, S.P., Geber, M.a., Yoshida, T., Fox, J.a., Hairston Jr, N.G., Ellner, S.P.,
Geber, M.a., Yoshida, T. & Fox, J.a. (2005) Rapid evolution and the convergence of
ecological and evolutionary time. *Ecology Letters,* **8,** 1114-1127.
Holt, R.D. (2008) Theoretical perspectives on resource pulses. *Ecology,* **89,** 671-681.
Hurlbert, A.H., Ballantyne IV, F. & Powell, S. (2008) Shaking a leg and hot to trot: the effects of
body size and temperature on running speed in ants. *Ecological Entomology,* **33,** 144-
154.
Huryn, A. & Benke, A. (2007) Relationship between biomass turnover and body size for stream
communities *Body Size: The Structure and Function of Aquatic Ecosystems* (eds A.
Hildrew, D. Raffaelli & R. Edmonds-Brown), pp. 55-76. Cambridge University Press
Cambridge.
Jacob, S. & Legrand, D. (2021) Phenotypic plasticity can reverse the relative extent of intra- and
interspecific variability across a thermal gradient. *Proceedings of the Royal Society B,*
**288,** 20210428.
Jones, E.I. & Gomulkiewicz, R. (2012) Biotic interactions, rapid evolution, and the establishment
of introduced species. *The American Naturalist,* **179,** E28-36.
Jones, L.E., Becks, L., Ellner, Stephen P., Hairston Jr, N.G., Yoshida, T. & Fussmann, G.F. (2009)
Rapid contemporary evolution and clonal food web dynamics. *Philosophical transactions*
*of the Royal Society of London. Series B, Biological sciences,* **364,** 1579-1591.
Jorgensen, P. & Tyers, M. (2004) How cells coordinate growth and division. *Current Biology,* **14,**
R1014-1027.
Karakoç, C., Clark, A.T. & Chatzinotas, A. (2020) Diversity and coexistence are influenced by
time-dependent species interactions in a predator-prey system. *Ecology Letters,* **23,** 983-
993.
Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature, and fitness: three rules. *Evolutionary*
*Ecology Research,* **10,** 251-268.
Kondoh, M., Kawatsu, K., Osada, Y. & Ushio, M. (2020) Theoretical Ecology, concepts, and
applications: A data-driven approach to complex ecological systems. *Theoretical*
*Ecology, Concepts and Applications* (eds K.S. McCann & G. Gellner), pp. 117-133. Oxford
University Press, Oxford.
Lafuente, E., Duneau, D. & Beldade, P. (2018) Genetic basis of thermal plasticity variation in
Drosophila melanogaster body size. *PLoS Genetics,* **14,** e1007686.
Lande, R. (1976) Natural Selection and Random Genetic Drift in Phenotypic Evolution. *Evolution,*
**30,** 314-334.
Lande, R. (1979) Quantitative genetic analysis of multivariate evolution, applied to brain: body
size allometry. *Evolution,* **33,** 402-416.
Lande, R. & Arnold, S.J. (1983) The Measurement of Selection on Correlated Characters.
*Evolution,* **37,** 1210-1226.
Lax, S., Abreu, C.I. & Gore, J. (2020) Higher temperatures generically favour slower-growing
bacterial species in multispecies communities. *Nature Ecology and Evolution.*

Marañón, E., Cermeno, P., Lopez-Sandoval, D.C., Rodriguez-Ramos, T., Sobrino, C., Huete-
Ortega, M., Blanco, J.M. & Rodriguez, J. (2013) Unimodal size scaling of phytoplankton
growth and the size dependence of nutrient uptake and use. *Ecology Letters,* **16,** 371-
379.

Miao, H., Xia, X., Perelson, A.S. & Wu, H. (2011) On Identifiability of Nonlinear Ode Models and
Applications in Viral Dynamics. *SIAM Review,* **53,** 3-39.

Motulsky, H. & Christopoulos, A. (2004) *Fitting Models to Biological Data using Linear and*
*Nonliear Regression: A Practical Guide to Curve Fitting.* Oxford University Press, Oxford,
UK.

Nguyen, J., Lara-Gutiérrez, J. & Stocker, R. (2020) Environmental fluctuations and their effects
on microbial communities, populations, and individuals. *FEMS Microbiology Reviews,*
**fuaa068,** 1-16.

Nieto-Acuña, C.A., Vargas-Garcia, C.A., Singh, A. & Pedraza, J.M. (2019) Efficient computation of
stochastic cell-size transient dynamics. *BMC Bioinformatics,* **20,** 647.

Noble, D.W.A., Radersma, R. & Uller, T. (2019) Plastic responses to novel environments are
biased towards phenotype dimensions with high additive genetic variation. *Proceedings*
*of the National Academy of Sciences,* **116,** 13452-13461.

Pavković-Lučić, S. & Kekić, V. (2013) Developmental temperature, body size and male mating
success in fruit flies, Drosophila melanogaster (Diptera : Drosophilidae). *European*
*Journal of Entomology,* **110,** 31-37.

Peckarsky, B.L., Taylor, B.W., McIntosh, A.R., McPeek, Mark A. & Lytle, D.A. (2001) Variation in
Mayfly Size at Metamorphosis as a Developmental Response to Risk of Predation.
*Ecology,* **82,** 740-757.

Peters, R.H. (1983) *The Ecological Implications of Body Size.* Cambridge University Press,
Cambridge, UK.

Preziosi, R.D. & Fairbairn, D.J. (2000) Lifetime selection on adult body size and components of
body size in a waterstrider: opposing selection and maintenance of sexual size
dimorphism. *Evolution,* **54,** 558-566.

Rall, B.C., Brose, U., Hartvig, M., Kalinkat, G., Schwarzmüller, F., Vucic-Pestic, O. & Petchey, O.L.
(2012) Universal temperature and body-mass scaling of feeding rates. *Philosophical*
*transactions of the Royal Society of London. Series B, Biological sciences,* **367,** 2923-
2934.

Riede, J.O., Brose, U., Ebenman, B., Jacob, U., Thompson, R.M., Townsend, C.R. & Jonsson, T.
(2011) Stepping in Elton's footprints: a general scaling model for body masses and
trophic levels across ecosystems. *Ecology Letters,* **14,** 169-178.

Roach, D.a. & Wulff, R.D. (1987) Maternal Effects in Plants. *Annual Review of Ecology and*
*Systematics,* **18,** 209-235.

Rogers, T.L., Munch, S.B., Stewart, S.D., Palkovacs, E.P., Giron-Nava, A., Matsuzaki, S.S. &
Symons, C.C. (2020) Trophic control changes with season and nutrient loading in lakes.
*Ecology Letters,* **23,** 1287-1297.

Rudman, S.M., Barbour, M.A., Csilléry, K., Gienapp, P., Guillaume, F., Hairston Jr, N.G., Hendry,
A.P., Lasky, J.R., Rafajlović, M., Räsänen, K., Schmidt, P.S., Seehausen, O., Therkildsen,
N.O., Turcotte, M.M. & Levine, J.M. (2018) What genomic data can reveal about eco-
evolutionary dynamics. *Nature Ecology & Evolution,* **2,** 9-15.

Savage, V.M., Gillooly, J.F., Brown, J.H. & Charnov, E.L. (2004) Effects of body size and
temperature on population growth. *The American Naturalist,* **163,** 429-441.
Schaffner, L.R., Govaert, L., De Meester, L., Ellner, S.P., Fairchild, E., Miner, B.E., Rudstam, L.G.,
Spaak, P. & Hairston, N.G. (2019) Consumer-resource dynamics is an eco-evolutionary
process in a natural plankton community. *Nature Ecology and Evolution,* **3,** 1351-1358.
Schreiber, S.J., Bürger, R. & Bolnick, D.I. (2011) The community effects of phenotypic and
genetic variation within a predator population. *Ecology,* **92,** 1582-1593.
Soetaert, K. & Petzoldt, T. (2010) Inverse Modelling, Sensitivity and Monte Carlo Analysis in R
Using Package FME. *Journal Of Statistical Software,* **33,** 1-28.
Sugihara, G., May, R., Ye, H., Hsieh, C.H., Deyle, E., Fogarty, M. & Munch, S. (2012) Detecting
causality in complex ecosystems. *Science,* **338,** 496-500.
Sugihara, G. & May, R.M. (1990) Nonlinear forecasting as a way to distinguishing chaos from
measurement errir in time series. *Science,* **344,** 734-741.
Takens, F. (1981) Detecting strange attractors in turbulence. *Dynamical Systems and*
*Turbulence.* . Springer, Berlin, Heilderberg.
Tariel, J., Plénet, S. & Luquet, É. (2020) Transgenerational Plasticity in the Context of Predator-
Prey Interactions. *Frontiers in Ecology and Evolution,* **8**.
Team, R.C. (2013) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. R Foundation for Statistical Computing, Vienna,
Austria.
terHorst, C.P., Miller, T.E. & Levitan, D.R. (2010) Evolution of prey in ecological time reduces the
effect size of predators in experimental microcosms. *Ecology,* **91,** 629-636.
Thompson, J.N. (1998) Rapid evolution as an ecological process. *Trends in Ecology & Evolution,*
**13,** 329-332.
Trebilco, R., Baum, J.K., Salomon, A.K. & Dulvy, N.K. (2013) Ecosystem ecology: size-based
constraints on the pyramids of life. *Trends in Ecology & Evolution,* **28,** 423-431.
Vanni, M.J., Duncan, J.M., González, M.J. & Horgan, M.J. (2009) Competition Among Aquatic
Organisms. *Encyclopedia of Inland Waters* (ed. G.E. Likens), pp. 395-404. Academic
Press.
Walczyńska, A., Franch-Gras, L. & Serra, M. (2017) Empirical evidence for fast temperature-
dependent body size evolution in rotifers. *Hydrobiologia,* **796,** 191-200.
Wasserman, S.S. & Mitter, C. (1978) The relationship of body size to breadth of diet in some
Lepidoptera. *Ecological Entomology,* **3,** 155-160.
Weber de Melo, V., Lowe, R., Hurd, P.J. & Petchey, O.L. (2020) Phenotypic responses to
temperature in the ciliate Tetrahymena thermophila. *Ecology and Evolution,* **10,** 7616-
7626.
Wieczynski, D.J., Singla, P., Doan, A., Singleton, A., Han, Z.Y., Votzke, S., Yammine, A. & Gibert,
J.P. (2021) Linking species traits and demography to explain complex temperature
responses across levels of organization. *Proceedings of the National Academy of*
*Sciences,* **118**.
Yamamichi, M., Yoshida, T. & Sasaki, A. (2011) Comparing the effects of rapid evolution and
phenotypic plasticity on predator-prey dynamics. *The American Naturalist,* **178,** 287-
304.

Yengo, L., Sidorenko, J., Kemper, K.E., Zheng, Z., Wood, A.R., Weedon, M.N., Frayling, T.M.,
Hirschhorn, J., Yang, J., Visscher, P.M. & Consortium, G. (2018) Meta-analysis of
genome-wide association studies for height and body mass index in approximately
700000 individuals of European ancestry. *Human Molecular Genetics,* **27,** 3641-3649.
845