Rapid plastic shifts in body size precede and determine population growth

Jean P. Gibert[1*], Ze-Yi Han[1], Daniel J. Wieczynski[1], Andrea Yammine[1]

[1]Department of Biology, Duke University, Durham NC, USA

*Corresponding author: jean.gibert@duke.edu

ACKNOWLEDGMENTS

We are indebted to John DeLong for insightful discussions on the Supply-Demand model. AY, DJW, and JPG, were supported by a U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science Program Grant to JPG, under Award Number DE-SC0020362.

CONFLICTS OF INTERESTS

None

AUTHOR CONTRIBUTIONS

JPG conceived the study and designed the experimental work with AY. AY collected the experimental data. JPG analyzed the data with inputs from ZYH and DJW. JPG wrote the first version of the manuscript and all authors contributed to subsequent versions. All authors approved the final version for submission and agree to be held accountable for all information within.

DATA AVAILABILITY

All data and code is already available at https://github.com/JPGibert/Tetra_Rapid_BodySize_Shifts and will be archived in Zenodo and/or Dryad upon publication.

ABSTRACT

1. Body size is a fundamental trait linked to many ecological processes—from individuals to
ecosystems. Although the effects of body size on metabolism are well-known, how body size
influences, and is influenced by, population growth and density is less clear. Specifically, 1)
whether body size, or population dynamics, more strongly influences the other, and, 2) whether
observed changes in body size are due to plasticity or rapid evolutionary change, are not well
understood.
2. Here, we address these two issues by experimentally tracking population density and mean
body size in the protist *Tetrahymena pyriformis* as it grows from low density to carrying
capacity. We then use state-of-the-art time series analyses to infer the direction, magnitude, and
causality of the link between body size and ecological dynamics. Last, we fit two alternative
dynamical models to our empirical time series to assess whether plasticity or rapid evolution
better explains changes in mean body size.
3. Our results clearly indicate that changes in body size precede and determine changes in
population density, not the other way around. We also show that a model assuming that size
changes via plasticity more parsimoniously explains these observed coupled phenotypic and
ecological dynamics than one that assumes rapid evolution drives changes in size.
4. Together these results suggest that rapid, plastic phenotypic change not only occurs well
within ecological timescales but may even precede—and causally influence—ecological
dynamics. Furthermore, large individuals may be favored and fuel high population growth rates
when population density is low, but smaller individuals may be favored once populations reach
carrying capacity and resources become scarcer. Thus, rapid plastic changes in functional traits
may play a fundamental and currently unrecognized role in familiar ecological processes like
logistic population growth.
55
56
57

59   Key Words: Eco-evolutionary Dynamics, Traits, Phenotypes, Phenotypic change, Ecological
60   dynamics

INTRODUCTION

Because of its myriad ecological consequences—from individuals (e.g., (Glasheen & McMahon 1996; Hurlbert, Ballantyne IV & Powell 2008; Pavković-Lučić & Kekić 2013)) to ecosystems (e.g., (Brown *et al.* 2004; Riede *et al.* 2011; Trebilco *et al.* 2013))—body size is one of the most important functional traits. Body size controls energetic demands (Gillooly *et al.* 2001; Brown *et al.* 2004), diet breadth (Wasserman & Mitter 1978; Gravel *et al.* 2013), the strength of ecological interactions (Berlow *et al.* 2009), trophic cascades (DeLong *et al.* 2015), and food web structure (Brose *et al.* 2006; Riede *et al.* 2011; Gravel *et al.* 2013; Gibert & DeLong 2014), all of which have ecosystem-level consequences (Anderson-Teixeira, Vitousek & Brown 2008; Trebilco *et al.* 2013). Body size can fuel or limit population growth through well-known relationships with demographic parameters like carrying capacity (K) and the intrinsic growth rate (r) (Damuth 1981; Savage *et al.* 2004; DeLong *et al.* 2015). For example, smaller organisms typically reproduce faster and can reach higher carrying capacities than larger ones, but also have higher mortality rates, which leads to faster population turnover (e.g., (Huryn & Benke 2007)). As a consequence, body size often determines how populations grow, and hence, ecological dynamics.

However, ecological dynamics can themselves influence body size, although these effects are less well understood. For example, individual size is under physiological regulatory control (Davidowitz, Roff & Nijhout 2005; Chelini, Delong & Hebets 2019) and therefore responds to changes in resource levels (Marañón *et al.* 2013). While resource levels can vary independently of population abundance (Holt 2008; Fey, Gibert & Siepielski 2019; Nguyen, Lara-Gutiérrez & Stocker 2020), high abundance leads to crowding and increasing intra-specific competition (Gavina *et al.* 2018). Stronger competition in turn effectively reduces resource availability at the

individual level, which often results in stunted growth and smaller body sizes (e.g., (Vanni *et al.*

2009)). Consequently, while body size can, and often does, influence ecological dynamics,

ecological dynamics can also influence body size. Which one more strongly influences the other,

however, is not known.

Body size can change over time through multiple mechanisms, both plastic (David,

Legout & Moreteau 2006; Ghosh, Testa & Shingleton 2013) and evolutionary (Hairston *et al.*

2005; DeLong *et al.* 2016), and body size is well-known to be under both natural and sexual

selection (Preziosi & Fairbairn 2000; Chelini, Delong & Hebets 2019). For example, predation

selects for smaller but faster growing prey in pitcher plant inquiline communities (terHorst,

Miller & Levitan 2010) and for shorter developmental times that result in smaller individuals in

mayflies (Peckarsky *et al.* 2001). Selection can act on traits rather quickly (Thompson 1998;

Hairston *et al.* 2005), and rapid evolutionary change has been shown to influence ecological

dynamics as they unfold (e.g., (Becks *et al.* 2012; Rudman *et al.* 2018; Schaffner *et al.* 2019)).

However, ecological dynamics in turn influence the pace and direction of evolutionary change

(e.g., (Cortez 2016; DeLong & Gibert 2016; Frickel, Sieber & Becks 2016; Gibert & Yeakel

2019)). In such eco-evolutionary feedback loops, selection imposed by ecological dynamics may

influence body size, whose rapid change can in turn alter ecological dynamics. However,

studying rapid evolutionary change in body size in unfeasible for most organisms and it is

therefore unclear how pervasive these processes are in nature.

While body size can change through evolution, rapid or otherwise, it also can change

within generations through plasticity (Lafuente, Duneau & Beldade 2018; Cameron *et al.* 2020).

For example, organisms grow in size throughout ontogeny and the environment often influences

ontogenetic trajectories, leading to variation in body size (Chelini, Delong & Hebets 2019),

while epigenetic DNA modifications can result in rapid phenotypic change from one generation

to the next in response to shifts in biotic or abiotic conditions (e.g., maternal effects, (Roach &

Wulff 1987; Galloway & Etterson 2007; Barbour & Gibert 2021)). Whether plastic change or

rapid evolutionary responses more strongly contribute to observed changes in body size over

short periods of time, however, is not well understood (e.g., (DeLong, Hanley & Vasseur

2014)). Among other reasons, this is because teasing apart plastic and evolutionary change can

be challenging even when sufficiently long time series exist (Ellner, Geber & Hairston 2011).

However, because body size mediates ecological interactions and processes, and can change on

ecological timescales, it is important to understand, track, and predict such change.

Here, we address these gaps by answering the following questions: 1) do rapid changes in

body size more strongly influence (and precede) population growth, or instead, does population

growth more strongly influence (and precede) changes in body size?, and, 2) are observed

changes in body size more consistent with a model of plasticity or rapid evolution? To address

the first question, we track changes in the density and body size of multiple experimental

populations of the protist *Tetrahymena pyriformis*, then use state-of-the art time series analysis to

assess causality. In protists, we expect changes in body size to be at least partly caused by

plasticity because reproduction (cell division) is tightly linked to ontogenetic changes in body

size (cells grow then divide when a critical size is attained). However, *T. pyriformis* also

reproduces extremely fast (4+ generations per day) and exhibits wide standing variation in body

size (Wieczynski *et al.* 2021), so rapid evolutionary change is also possible. Therefore, to

address our second question, we compare two alternative mathematical models: one that assumes

only plasticity drives change in body size (plasticity model), and one that assumes only rapid

evolution drives change in body size (eco-evolutionary model). We then fit the proposed models

to abundance and body size time series from our experiment and use model comparison to infer

what mechanism—plasticity or rapid evolution—most likely explains our data. Our results show

that rapid plastic changes in body size precede and determine changes in population density over

time, thus suggesting that plastic phenotypic change may be integral to ecological dynamics.

METHODS

Microcosm experiments

We tracked changes in abundance and body size in the protist *Tetrahymena pyriformis* for 14

days. To do so, we set up 6 experimental microcosms in 250 mL autoclaved borosilicate jars

filled with 200 mL of Carolina protist pellet media (1L of autoclaved DI water per pellet)

inoculated with pond bacteria from Duke Forest (Gate 9/Wilbur pond, Lat=36.02°, Long=-

78.99°, Durham, NC, Fig 1). Microcosms were initialized at densities of 10 ind/mL and were

incubated in temperature (22°C) and humidity-controlled (65% humidity) growth chambers

(Percival AL-22L2, Percival Scientific, Perry, Iowa) on a 12hr night/day cycle. Densities

(ind/mL) and trait dynamics were tracked daily through fluid imaging of 1 mL subsamples of

each microcosm through fluid imaging (Fig 1, FlowCam, Fluid Imaging Technologies,

Scarborough, ME, USA). In doing so, we imaged over 150k individual cells, thus giving our

experiment unparalleled insight into how density and body size changed together over the course

of this experiment. Last, neither water nor nutrients were replaced throughout the course of this

experiment.

Time-series analysis

To assess whether change in body size more strongly influenced changes in density, or

vice versa, we used *Convergent Cross Mapping* (or CCM, (Sugihara *et al.* 2012; Rogers *et al.*

2020)) on the density and body size time series. CCM quantifies the degree to which one time

series causally influences another one by estimating how much information of the one is

contained in the other (Sugihara *et al.* 2012): if a variable X causally influences another variable

Y, but Y does not influence X, we should expect Y to contain information about X, but not the

other way around. CCM assesses this by quantifying whether variable X at time t can be used to

predict variable Y at time t+1 for subsets of the time series of increasing length (the length of

these subsets is called the 'library size'). If X more strongly influences changes in Y than the

other way around, it also means Y responds to X more strongly than X responds to Y. If the

effect of X on Y is causal, as opposed there being simple correlation, the ability to predict Y

from X should increase with library size, while the error associated with that prediction should

decline. If predictability that does not change with library size, there is correlation but not

causation. If X causes Y, it also implies that change in X precedes change in Y, to which Y

responds to. The analysis produces an estimate of how well one time series predicts the other

(i.e., the *predictability* coefficient): the larger this value, the stronger the link—causal or

correlative—between one time series and the other (Sugihara *et al.* 2012). To perform the

analysis, we used R package multispatialCCM v1.0 (Clark *et al.* 2015), which allowed us to use

replicated time series to increase power. Previous studies have shown that CCM allows for the

inference of causation in different ecological systems and environmental conditions for which

long enough time series exist (Sugihara *et al.* 2012; Clark *et al.* 2015; Karakoç, Clark &

Chatzinotas 2020; Kondoh *et al.* 2020; Rogers *et al.* 2020).

Mathematical models

To assess whether observed changes in body size were more likely due to plasticity or rapid

evolution, we fitted two possible models that track change in the abundance and average body

size of a population, *N*, as it grows logistically towards a carrying capacity *K* with intrinsic

growth rate *r*. Following previous work (Abrams 1977; Abreu *et al.* 2019; Lax, Abreu & Gore

2020; Wieczynski *et al.* 2021), we included an additional mortality term in the ecological

dynamics, mN, to account for regular loss of individuals from the population (through

sampling). This additional mortality term has been shown to better describe the ecological

dynamics of a microbial microcosm with frequent sampling, like ours (Abreu *et al.* 2019; Lax,

Abreu & Gore 2020). Furthermore, we assumed that the intrinsic growth rate and the carrying

capacity were functions of body size, *M*, of the forms $r = aM^\alpha$ and $K = bM^\gamma$, following well-

known allometric relationships (Damuth 1981; Savage *et al.* 2004). Taken together, the

ecological baseline model was thus written as:

197
$$\frac{dN}{dt} = aM^\alpha \left(1 - \frac{N}{bM^\gamma}\right) - mN. \qquad \text{eq [1]}$$

To incorporate the coupling between changes in N and changes in *M*, we used two

alternative model formulations to track changes in *M*: the first assumed that only plastic change

in *M* could occur (plasticity model), the second assumed that only rapid evolutionary change in

M could occur (eco-evolutionary model). The plasticity model modifies the existing Supply-

Demand model for body size dynamics (DeLong 2012; DeLong & Walsh 2015) and assumes

that the rate of change in body size, $\frac{dM}{dt}$, increases with energy intake (supply), and decreases

with energy loss (demand). Following a recent study (DeLong 2020), we assumed that the supply

in a species growing logistically depends on the ratio of the carrying capacity, K, and the

abundance of the species, N, times a conversion rate constant *e*, that converts the supply to units

of M. When the population is small, the supply approaches infinity, and it approaches e when N

grows to K. Following previous studies (DeLong, Hanley & Vasseur 2014), we also assumed

that the demand was the metabolic cost of the organism, which is known to increase

allometrically with body size, as cM^δ. Taken together, the equation controlling changes in body

size M was:

$$\frac{dM}{dt} = \frac{eK}{N} - cM^\delta.$$ eq [2]

For the evolutionary model, we followed previous studies on eco-evolutionary dynamics

(e.g., (Abrams, Harada & Matsuda 1993; Abrams & Matsuda 1997; Jones *et al.* 2009; Ellner &

Becks 2010; Jones & Gomulkiewicz 2012; Cortez 2016; Cortez 2018)) to incorporate evolution

in M at a rate that equals the product of the total amount of additive heritable variation in body

size (*i.e.*, the product of the total phenotypic variance, σ^2, and the narrow sense heritability, h^2),

and the selection gradient (*i.e.,* the change in fitness, F, with respect to a change in M, which

represents the strength of selection acting on M). Taken together, the equation controlling the

change in M over time under these assumptions was:

$$\frac{dM}{dt} = \sigma^2 h^2 \frac{\partial F(M)}{\partial M},$$ eq [3]

were $F(M) = \frac{1}{N}\frac{dN}{dt}$ (Lande 1976; Schreiber, Bürger & Bolnick 2011).

Model fitting, parameter uncertainty, and model selection

To assess how well the models in Eqs 1-3 reproduced empirical body size and density

dynamics from experimental microcosms, we fitted the models to the *T. pyriformis* time series

using R package FME v1.3.6.1 (Soetaert & Petzoldt 2010). The allometric parameters a and b

for r and K respectively, should not differ between the two models, so they were optimized

during preliminary model fitting, then set constant for both models during final model fitting and

estimation of parameter uncertainty (Table 1). We assumed that the scaling parameter of the

metabolic cost, δ, equaled 1, as has been shown to be the case for protists (DeLong *et al.* 2010),

despite it being closer to ¾ for metazoans (Gillooly *et al.* 2001; Brown *et al.* 2004; DeLong *et al.*

2010). Fitting of ordinary differential equations using package FME relies on the Levenberg-

Marquardt algorithm for parameter estimation, and a Metropolis-Hastings MCMC procedure for

estimation of parameter uncertainty (Soetaert & Petzoldt 2010). Model comparison was done

using Akaike Information Criterion (Burnham & Anderson 2002) as:

$$AIC_i = 2k_i + n_N \ln\left(SSE_{i,N}\right) + n_M \ln\left(SSE_{i,M}\right) \ , \qquad \text{eq[4]}$$

Where k_i is the number of parameters of model i, n_N is the number of datapoints considered in

the density time series, n_M the number of datapoints considered in the body size time series,

$SSE_{i,N}$ the sum of squared errors of model i with respect to the density time series, and $SSE_{i,M}$ is

the sum of squared errors of model i with respect to the body size time series. All data and code

are available at https://github.com/JPGibert/Tetra_Rapid_BodySize_Shifts.

RESULTS

Tetrahymena abundances increased to carrying capacity roughly 4 days after microcosms where

initialized (Fig 2a). Body size increased over the first day, then decreased continuously for 12-13

days (Fig 2b). Results from CCM analysis showed large predictability values for densities using

changes in body size as a predictor, but smaller predictability values for body size using densities

as the predictor (Fig 2c). Moreover, the stronger effect of body size on densities was found to be

causal (sharp increase in predictability with library size and decrease in standard deviation, Fig

2c), but that of densities on body size is either not, or only weakly causal (slow convergence of

predictability and little change in standard deviation, Fig 2c). This indicates that changes in body

size influence density changes over time more strongly than density influences body size, and

because size predicts density but not the other way around, it also indicates that changes in body

size precede changes in population density.

Both models (plasticity and eco-evolutionary) fit the empirical data remarkably well (Fig

3a-d, Table 1). MCMC chains converged for all fitted parameters (Appendix Fig S1, S2) and

model parameters were identifiable and free of correlations for all but three parameter pairs

across the two models (α and m, and c and e in the plasticity model, and γ and m in the eco-evo

model, Fig S3, S4). The plasticity model did a slightly better job at fitting the abundance data

than the eco-evolutionary model (notice departures of eco-evo model during initial dynamics,

Fig 3e), and both did a good job at fitting the body size data (Fig 3f). Both models arrived at very

similar fitted values for the three model parameters they had in common (Table 1), indicating

good agreement between the two. Consistent with existing literature, both models identified the

scaling of K and M (γ), as a negative number close to -¾ (Table 1). However, both models

predicted the scaling between r and M (α) as positive and close to 1 (Table 1) while the literature

pins that value—across species—to -¼ (Savage *et al.* 2004). Last, model selection through AIC

indicated that the plasticity model was the most parsimonious of the two (Table 1), likely

indicating that observed coupled density and body size dynamics were mostly driven by

plasticity.

DISCUSSION

Because of the myriad ecological consequences of body size (e.g. (Gillooly *et al.* 2001;

Brown *et al.* 2004; DeLong *et al.* 2010)), it is important to understand how changes in body

size—plastic or evolutionary—may influence or be influenced by ecological dynamics. We show

that changes in body size more strongly influence changes in densities than the other way around

(Fig 2), and that a model that accounts for rapid plastic change in body size provides the most

parsimonious explanation for the observed paired ecological and body size dynamics (Fig 3,

Table 1). While previous studies found that phenotypic change often lags behind ecological rates

of change (e.g., (DeLong *et al.* 2016)), our results clearly show that phenotypic change not only

occurs well within ecological timescales, but may even precede—and causally influence—those

dynamics (Fig 2c). Our results thus emphasize the need to further study and understand the

ecological consequences of rapid plastic phenotypic change (Yamamichi, Yoshida & Sasaki

2011; Tariel, Plénet & Luquet 2020), as plasticity may play a crucial role in determining the fate

of networks of species interactions in a warming world (Barbour & Gibert 2021; Jacob &

Legrand 2021).

Cells can enter and leave the cell division cycle depending on cell size and nutrient

availability (Baserga 1968; Fukada *et al.* 2007). As the *T. pyriformis* population reaches carrying

capacity, low resource levels likely cue cells into existing the cell division cycle, in turn resulting

in stunted growth (cells grow to reproduce (Baserga 1968)) and reduced average body sizes.

Observed total standing phenotypic variation is thus likely largely non-heritable, as also

suggested in a recent study (Jacob & Legrand 2021). Surprisingly, the eco-evolutionary model

itself supports this idea, as it indicates that the total amount of standing heritable variation in

body size is rather small ($\sigma^2 h^2 = 0.992$, Table 1). For comparison, the total amount of

phenotypic variation (heritable or not) in our initial population was 94 (units of μm^3 squared), so

the heritable portion of that total phenotypic variation would be of the order of ~1%, in

agreement with our model selection results. Shifts in *T. pyriformis* phenotype have also been

shown to occur differentially across environmental conditions (DeLong *et al.* 2017; Weber de

Melo *et al.* 2020), again suggesting the occurrence of plasticity.

While the rapid evolution model is less parsimonious, our results do not rule out the

possibility of rapid evolution. Indeed, plastic change often precedes evolutionary change and

occurs mostly along the axes of variation with the largest amount of heritable variation (the

classic evolutionary 'path of least resistance' (Lande 1976; Lande 1979; Lande & Arnold 1983)),

thus setting the stage for evolution to occur along those axes (Noble, Radersma & Uller 2019). A

similar process may be at play here. Indeed, despite low heritability, the less parsimonious eco-

evolutionary model suggests that short term selection imposed by density-dependence may be

strong enough to consistently shift body size over time, which in turn influences population

dynamics (Fig 2). Our own data indicate that *T. pyriformis* reproduces at a rate of 3.5-4 new

individuals per individual per day. This extremely fast population growth may eventually allow

for evolutionary change in body size—if selection is strong enough, as heritability is low, but not

nil—even if it likely lags behind plastic change (Chevin, Lande & Mace 2010; Fox *et al.* 2019).

Interestingly, neither water nor nutrients were replenished during our experiment; both

were limited and were likely consistently lost from the system over time through sampling. This

nutrient impoverishment should lead to a strong decline in K. However, a strong decline in K

was not only not observed, but the most parsimonious model suggests that such a decline either

does not occur, or does so only weakly (Fig 3). Because carrying capacity declines with body

size (Damuth 1981), a rapid decline in nutrient concentrations may thus select for smaller body

size (Vanni *et al.* 2009), which could in turn counter the effect of negative density-dependence

overall. These results suggest a way in which rapid plastic change in body size may allow

organisms to regulate population growth and density-dependent factors, even as nutrients

become increasingly limited.

Last, both the eco-evolutionary and plasticity models predicted a positive relationship

between body size and the intrinsic growth rate (Fig 4a). This result stands in contrast to

empirical data across species, which show lower intrinsic growth rates for larger organisms (or

$\alpha < 0$, REFS). Within species, however, larger individuals typically reproduce more and die at

lower rates (Peters 1983), leading to higher r due to lower mortality and higher reproduction

(Kingsolver & Huey 2008). This positive relationship between body size and the demographic

processes that fuel intrinsic growth rates are well understood within species (Peters 1983;

Kingsolver & Huey 2008), emphasizing how inter-species and intra-species body size scaling

may often differ (Rall *et al.* 2012).

Overall, our study shows that rapid plastic change in phenotypes more strongly

influences ecological dynamics than the other way around, suggesting that rapid changes in body

size are likely integral to the process of population growth itself. This study sheds light on the

ecological and evolutionary constraints that regulate population growth and provides new

insights about how organisms cope with the negative effects of density-dependence.

Table 1: Parameter estimates and uncertainty for both fitted models.

		r scaling with body size (α)	K scaling with body size (γ)	Mortality rate (m)	Supply-Demand conversion parameter (e)	Supply-Demand loss parameter (c)	Heritable Variation ($\sigma^2 h^2$)		Allometric Intercept for r (a)	Allometric Intercept for K (b)	AIC (ΔAIC)
(a) Plasticity Model	Estimate	-0.721	1.022	0.969	24.295	0.0873	—	Set value	~10^{-4}	~10^6	**3246.97 (0)**
	95% Credible Interval	(-0.735, -0.720)	(1.010, 1.023)	(0.770, 0.991)	(18.400, 26.714)	(0.077, 0.091)	—	Post-Optimization	$2.2*10^{-4}$	$8.2*10^6$	
(b) Eco-Evo Model	Estimate	-0.748	1.056	0.897	—	—	0.992	Initial value	As above	As above	**3287.66 (+40.69)**
	95% Credible Interval	(-0.753, -0.743)	(1.048, 1.064)	(0.770, 1.026)	—	—	(0.825, 1.193)	Post-Optimization	As above	As above	

346   FIGURES

348   Fig 1: Depiction of experimental procedure: microcosms were initialized at 10ind/mL and each

349   microcosm was replicated six times. Population density and cell size were recorded daily for 14

350   days thereafter using fluid imaging.

357 Fig 2: (a) Time series of Tetrahymena density. (b) Time series of mean Tetrahymena body size.

358 (c) Convergent Cross Mapping predictability plot against library size (i.e., length of the time

359 series used for analysis), using body size to predict density (purple) or density to predict body

360 size (yellow), repeated 1000 times. Solid line indicates mean values and dashed lines indicate

361 standard deviation of the mean.

Fig 3: (a) Density data (grey points) and plasticity model fit (solid blue). Uncertainty is represented as 700 model predictions whose parameters were sampled from posterior distributions for each model parameter, estimated during model fitting (transparent blue, Table 1). (b) As in (a), but for body size data. (c–d) As in (a–b) but for the eco-evolutionary model fit. (e) Comparison of the plasticity (blue) and evo-evolutionary (green) model fits to the density data. The plasticity model fits the data slightly better, especially at the beginning of the experiment. (f) As in (e) but for body size data. The eco-evolutionary model fits the data slightly better at first, although both models fit well after day 2.

REFERENCES

Abrams, P.A. (1977) Density-Independent Mortality and Interspecific Competition: A Test of
Pianka's Niche Overlap Hypothesis. *The American Naturalist,* **111,** 539-552.
Abrams, P.A., Harada, Y. & Matsuda, H. (1993) On the Relationship between Quantitative
Genetic and Ess Models. *Evolution,* **47,** 982-985.
Abrams, P.A. & Matsuda, H. (1997) Prey Adaptation as a Cause of Predator-Prey Cycles.
Evolution, **51,** 1742-1750.
Abreu, C.I., Friedman, J., Andersen Woltz, V.L. & Gore, J. (2019) Mortality causes universal
changes in microbial community composition. *Nature Communications,* **10,** 2120.
Anderson-Teixeira, K.J., Vitousek, P.M. & Brown, J.H. (2008) Amplified temperature
dependence in ecosystems developing on the lava flows of Mauna Loa , Hawai ' i.
Proceedings of the National Academy of Sciences, **105,** 228-233.
Barbour, M.A. & Gibert, J.P. (2021) Genetic and plastic rewiring of food webs under climate
change. *Journal of Animal Ecology*.
Baserga, R. (1968) Biochemistry of the cell cycle: a review *Cell Tissue Kinetics,* **1,** 167-191.
Becks, L., Ellner, S.P., Jones, L.E. & Hairston, N.G. (2012) The functional genomics of an eco-
evolutionary feedback loop: Linking gene expression, trait evolution, and community
dynamics. *Ecology Letters,* **15,** 492-501.
Berlow, E.L., Dunne, J.A., Martinez, N.D., Stark, P.B., Williams, R.J. & Brose, U. (2009)
Simple prediction of interaction strengths in complex food webs. *Proceedings of the*
National Academy of Sciences of the United States of America, **106,** 187-191.
Brose, U., Jonsson, T., Berlow, E.L., Warren, P.H., Banasek-Richter, C., Bersier, L.-F.,
Blanchard, J.L., Brey, T., Carpenter, S.R., Cattin, M.-F., Cushing, L., Hassan, A.D., Dell,
A.I., Edwards, F., Harper-Smith, S., Jacob, U., Ledger, M.E., Martinez, N.D., Memmott,
J., Mintenbeck, K., Pinnegar, J.K., Rall, B.C., Rayner, T.S., Reuman, D.C., Ruess, L.,
Ulrich, W., Williams, R.J., Woodward, G. & Cohen, J.E. (2006) Consumer–resource
body-size relationships in natural food webs. *Ecology,* **87,** 2411-2417.
Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004) Toward a Metabolic
Theory of Ecology. *Ecology,* **85,** 1771-1789.
Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference: A practical*
information-theoretic approach, 2nd ed. edn. Springer Science, New York.
Cameron, M.D., Grant, V.H., Joly, K., Schmidt, J.H., Gustine, D.D., Mangipane, L.S.,
Mangipane, B. & Sorum, M.S. (2020) Body size plasticity in North American black and
brown bears. *Ecosphere,* **e03235,** 1-11.
Chelini, M.C., Delong, J.P. & Hebets, E.A. (2019) Ecophysiological determinants of sexual size
dimorphism: integrating growth trajectories, environmental conditions, and metabolic
rates. *Oecologia,* **191,** 61-71.
Chevin, L.M., Lande, R. & Mace, G.M. (2010) Adaptation, plasticity, and extinction in a
changing environment: towards a predictive theory. *PLoS Biology,* **8,** e1000357.
Clark, A.T., Ye, H., Isbell, F., Deyle, E.R., Cowles, J., Tilman, G.D. & Sugihara, G. (2015)
Spatial convergent cross mapping to detect causal relationships from short time series.
Ecology, **96,** 1174-1181.
Cortez, M.H. (2016) How the Magnitude of Prey Genetic Variation Alters Predator-Prey Eco-
Evolutionary Dynamics. *The American Naturalist,* **188,** 329-341.

Cortez, M.H. (2018) Genetic variation determines which feedbacks drive and alter predator–prey
eco-evolutionary cycles. *Ecological Monographs,* **88,** 353-371.
Damuth, J. (1981) Population density and body size in mammals. *Nature,* **290,** 699-700.
David, J.R., Legout, H. & Moreteau, B. (2006) Phenotypic plasticity of body size in a temperate
population of Drosophila melanogaster: when the temperature-size rule does not apply.
Journal of genetics, **85,** 9-23.
Davidowitz, G., Roff, D.A. & Nijhout, H.F. (2005) A Physiological Perspective on the Response
of Body Size and Development Time to Simultaneous Directional Selection. *Integrative*
and Comparative Biology, **45,** 525-532.
DeLong, J.P. (2012) Experimental demonstration of a ' rate – size ' trade-off governing body
size optimization. *Evolutionary Ecology Research,* **14,** 343-352.
DeLong, J.P. (2020) Detecting the Signature of Body Mass Evolution in the Broad-Scale
Architecture of Food Webs. *The American Naturalist,* **196,** 443-453.
DeLong, J.P., Brassil, C.E., Erickson, E.K., Forbes, V.E., Moriyama, E. & Rieckhof, W.R.
(2017) Dynamic thermal reaction norms and body size oscillations challenge
explanations of
the temperature–size rule. *Evolutionary Ecology Research,* **18,** 293-303.
DeLong, J.P., Forbes, V.E., Galic, N., Gibert, J.P., Laport, R.G., Phillips, J.S. & Vavra, J.M.
(2016) How fast is fast? Eco-evolutionary dynamics and rates of change in populations
and phenotypes. *Ecology and Evolution,* **6,** 573-581.
DeLong, J.P. & Gibert, J.P. (2016) Gillespie eco-evolutionary models (GEMs) reveal the role of
heritable trait variation in eco-evolutionary dynamics. *Ecology and Evolution,* **6,** 935-
945.
DeLong, J.P., Gilbert, B., Shurin, J.B., Savage, V.M., Barton, B.T., Clements, C.F., Dell, A.I.,
Greig, H.S., Harley, C.D.G., Kratina, P., McCann, K.S., Tunney, T.D., Vasseur, D.A. &
O'Connor, M.I. (2015) The Body Size Dependence of Trophic Cascades. *The American*
Naturalist, **185,** 354-366.
DeLong, J.P., Hanley, T.C. & Vasseur, D.A. (2014) Predator-prey dynamics and the plasticity of
predator body size. *Functional Ecology,* **28,** 487-493.
DeLong, J.P., Okie, J.G., Moses, M.E., Sibly, R.M. & Brown, J.H. (2010) Shifts in metabolic
scaling, production, and efficiency across major evolutionary transitions of life.
Proceedings of the National Academy of Sciences of the United States of America, **107,**
12941-12945.
DeLong, J.P. & Walsh, M. (2015) The interplay between resource supply and demand
determines the influence of predation on prey body size. *Canadian Journal of Fisheries*
and Aquatic Sciences, **72,** 1-7.
Ellner, Stephen P. & Becks, L. (2010) Rapid prey evolution and the dynamics of two-predator
food webs. *Theoretical Ecology***,** 133-152.
Ellner, S.P., Geber, M.A. & Hairston, N.G. (2011) Does rapid evolution matter? Measuring the
rate of contemporary evolution and its impacts on ecological dynamics. *Ecology Letters,*
14, 603-614.
Fey, S.B., Gibert, J.P. & Siepielski, A.M. (2019) The consequences of mass mortality events for
the structure and dynamics of biological communities. *Oikos***,** 1-12.
Fox, R.J., Donelson, J.M., Schunter, C., Ravasi, T. & Gaitan-Espitia, J.D. (2019) Beyond buying
time: the role of plasticity in phenotypic adaptation to rapid environmental change.
Philosophical Transactions of the Royal Society B: Biological Sciences, **374,** 20180174.

Frickel, J., Sieber, M. & Becks, L. (2016) Eco-evolutionary dynamics in a coevolving host-virus system. *Ecology Letters,* **19,** 450-459.

Fukada, S., Uezumi, A., Ikemoto, M., Masuda, S., Segawa, M., Tanimura, N., Yamamoto, H., Miyagoe-Suzuki, Y. & Takeda, S. (2007) Molecular signature of quiescent satellite cells in adult skeletal muscle. *Stem Cells,* **25,** 2448-2459.

Galloway, L.F. & Etterson, J.R. (2007) Transgenerational plasticity is adaptive in the wild. *Science (New York, N.Y.),* **318,** 1134-1136.

Gavina, M.K.A., Tahara, T., Tainaka, K.I., Ito, H., Morita, S., Ichinose, G., Okabe, T., Togashi, T., Nagatani, T. & Yoshimura, J. (2018) Multi-species coexistence in Lotka-Volterra competitive systems with crowding effects. *Scientific Reports,* **8,** 1198.

Ghosh, S.M., Testa, N.D. & Shingleton, A.W. (2013) Temperature-size rule is mediated by thermal plasticity of critical size in Drosophila melanogaster. *ZooKeys,* **298**.

Gibert, J.P. & DeLong, J.P. (2014) Temperature alters food web body-size structure. *Biology Letters,* **10,** 20140473-20140473.

Gibert, J.P. & Yeakel, J.D. (2019) Eco-Evolutionary Origins of Diverse Abundance , Biomass , and Trophic Structures in Food Webs. *Frontiers in Ecology and Evolution,* **7,** 1-11.

Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001) Effects of size and temperature on metabolic rate. *Science (New York, N.Y.),* **293,** 2248-2251.

Glasheen, J. & McMahon, T. (1996) Size-dependence of water-running ability in basilisk lizards (Basiliscus basiliscus). *The Journal of Experimental Biology,* **199,** 2611-2618.

Gravel, D., Poisot, T., Albouy, C., Velez, L. & Mouillot, D. (2013) Inferring food web structure from predator-prey body size relationships. *Methods in Ecology and Evolution,* **4,** 1083-1090.

Hairston, N.G., Ellner, S.P., Geber, M.a., Yoshida, T., Fox, J.a., Hairston Jr, N.G., Ellner, S.P., Geber, M.a., Yoshida, T. & Fox, J.a. (2005) Rapid evolution and the convergence of ecological and evolutionary time. *Ecology Letters,* **8,** 1114-1127.

Holt, R.D. (2008) Theoretical perspectives on resource pulses. *Ecology,* **89,** 671-681.

Hurlbert, A.H., Ballantyne IV, F. & Powell, S. (2008) Shaking a leg and hot to trot: the effects of body size and temperature on running speed in ants. *Ecological Entomology,* **33,** 144-154.

Huryn, A. & Benke, A. (2007) Relationship between biomass turnover and body size for stream communities *Body Size: The Structure and Function of Aquatic Ecosystems* (eds A. Hildrew, D. Raffaelli & R. Edmonds-Brown), pp. 55-76. Cambridge University Press Cambridge.

Jacob, S. & Legrand, D. (2021) Phenotypic plasticity can reverse the relative extent of intra- and interspecific variability across a thermal gradient. *Proceedings of the Royal Society B,* **288,** 20210428.

Jones, E.I. & Gomulkiewicz, R. (2012) Biotic interactions, rapid evolution, and the establishment of introduced species. *The American Naturalist,* **179,** E28-36.

Jones, L.E., Becks, L., Ellner, Stephen P., Hairston Jr, N.G., Yoshida, T. & Fussmann, G.F. (2009) Rapid contemporary evolution and clonal food web dynamics. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences,* **364,** 1579-1591.

Karakoç, C., Clark, A.T. & Chatzinotas, A. (2020) Diversity and coexistence are influenced by time-dependent species interactions in a predator-prey system. *Ecology Letters,* **23,** 983-993.

Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature, and fitness: three rules. *Evolutionary Ecology Research,* **10,** 251-268.

Kondoh, M., Kawatsu, K., Osada, Y. & Ushio, M. (2020) Theoretical Ecology, concepts, and applications: A data-driven approach to complex ecological systems. *Theoretical Ecology, Concepts and Applications* (eds K.S. McCann & G. Gellner), pp. 117-133. Oxford University Press, Oxford.

Lafuente, E., Duneau, D. & Beldade, P. (2018) Genetic basis of thermal plasticity variation in Drosophila melanogaster body size. *PLoS Genetics,* **14,** e1007686.

Lande, R. (1976) Natural Selection and Random Genetic Drift in Phenotypic Evolution. *Evolution,* **30,** 314-334.

Lande, R. (1979) Quantitative genetic analysis of multivariate evolution, applied to brain: body size allometry. *Evolution,* **33,** 402-416.

Lande, R. & Arnold, S.J. (1983) The Measurement of Selection on Correlated Characters. *Evolution,* **37,** 1210-1226.

Lax, S., Abreu, C.I. & Gore, J. (2020) Higher temperatures generically favour slower-growing bacterial species in multispecies communities. *Nature Ecology and Evolution*.

Marañón, E., Cermeno, P., Lopez-Sandoval, D.C., Rodriguez-Ramos, T., Sobrino, C., Huete-Ortega, M., Blanco, J.M. & Rodriguez, J. (2013) Unimodal size scaling of phytoplankton growth and the size dependence of nutrient uptake and use. *Ecology Letters,* **16,** 371-379.

Nguyen, J., Lara-Gutiérrez, J. & Stocker, R. (2020) Environmental fluctuations and their effects on microbial communities, populations, and individuals. *FEMS Microbiology Reviews,* **fuaa068,** 1-16.

Noble, D.W.A., Radersma, R. & Uller, T. (2019) Plastic responses to novel environments are biased towards phenotype dimensions with high additive genetic variation. *Proceedings of the National Academy of Sciences,* **116,** 13452-13461.

Pavković-Lučić, S. & Kekić, V. (2013) Developmental temperature, body size and male mating success in fruit flies, Drosophila melanogaster (Diptera : Drosophilidae). *European Journal of Entomology,* **110,** 31-37.

Peckarsky, B.L., Taylor, B.W., McIntosh, A.R., McPeek, Mark A. & Lytle, D.A. (2001) Variation in Mayfly Size at Metamorphosis as a Developmental Response to Risk of Predation. *Ecology,* **82,** 740-757.

Peters, R.H. (1983) *The Ecological Implications of Body Size*. Cambridge University Press, Cambridge, UK.

Preziosi, R.D. & Fairbairn, D.J. (2000) Lifetime selection on adult body size and components of body size in a waterstrider: opposing selection and maintenance of sexual size dimorphism. *Evolution,* **54,** 558-566.

Rall, B.C., Brose, U., Hartvig, M., Kalinkat, G., Schwarzmüller, F., Vucic-Pestic, O. & Petchey, O.L. (2012) Universal temperature and body-mass scaling of feeding rates. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences,* **367,** 2923-2934.

Riede, J.O., Brose, U., Ebenman, B., Jacob, U., Thompson, R.M., Townsend, C.R. & Jonsson, T. (2011) Stepping in Elton's footprints: a general scaling model for body masses and trophic levels across ecosystems. *Ecology Letters,* **14,** 169-178.

Roach, D.a. & Wulff, R.D. (1987) Maternal Effects in Plants. *Annual Review of Ecology and Systematics,* **18,** 209-235.

Rogers, T.L., Munch, S.B., Stewart, S.D., Palkovacs, E.P., Giron-Nava, A., Matsuzaki, S.S. & Symons, C.C. (2020) Trophic control changes with season and nutrient loading in lakes. *Ecology Letters,* **23,** 1287-1297.

Rudman, S.M., Barbour, M.A., Csilléry, K., Gienapp, P., Guillaume, F., Hairston Jr, N.G., Hendry, A.P., Lasky, J.R., Rafajlović, M., Räsänen, K., Schmidt, P.S., Seehausen, O., Therkildsen, N.O., Turcotte, M.M. & Levine, J.M. (2018) What genomic data can reveal about eco-evolutionary dynamics. *Nature Ecology & Evolution,* **2,** 9-15.

Savage, V.M., Gillooly, J.F., Brown, J.H. & Charnov, E.L. (2004) Effects of body size and temperature on population growth. *The American Naturalist,* **163,** 429-441.

Schaffner, L.R., Govaert, L., De Meester, L., Ellner, S.P., Fairchild, E., Miner, B.E., Rudstam, L.G., Spaak, P. & Hairston, N.G. (2019) Consumer-resource dynamics is an eco-evolutionary process in a natural plankton community. *Nature Ecology and Evolution,* **3,** 1351-1358.

Schreiber, S.J., Bürger, R. & Bolnick, D.I. (2011) The community effects of phenotypic and genetic variation within a predator population. *Ecology,* **92,** 1582-1593.

Soetaert, K. & Petzoldt, T. (2010) Inverse Modelling, Sensitivity and Monte Carlo Analysis in R Using Package FME. *Journal Of Statistical Software,* **33,** 1-28.

Sugihara, G., May, R., Ye, H., Hsieh, C.H., Deyle, E., Fogarty, M. & Munch, S. (2012) Detecting causality in complex ecosystems. *Science,* **338,** 496-500.

Tariel, J., Plénet, S. & Luquet, É. (2020) Transgenerational Plasticity in the Context of Predator-Prey Interactions. *Frontiers in Ecology and Evolution,* **8**.

terHorst, C.P., Miller, T.E. & Levitan, D.R. (2010) Evolution of prey in ecological time reduces the effect size of predators in experimental microcosms. *Ecology,* **91,** 629-636.

Thompson, J.N. (1998) Rapid evolution as an ecological process. *Trends in Ecology & Evolution,* **13,** 329-332.

Trebilco, R., Baum, J.K., Salomon, A.K. & Dulvy, N.K. (2013) Ecosystem ecology: size-based constraints on the pyramids of life. *Trends in Ecology & Evolution,* **28,** 423-431.

Vanni, M.J., Duncan, J.M., González, M.J. & Horgan, M.J. (2009) Competition Among Aquatic Organisms. *Encyclopedia of Inland Waters* (ed. G.E. Likens), pp. 395-404. Academic Press.

Wasserman, S.S. & Mitter, C. (1978) The relationship of body size to breadth of diet in some Lepidoptera. *Ecological Entomology,* **3,** 155-160.

Weber de Melo, V., Lowe, R., Hurd, P.J. & Petchey, O.L. (2020) Phenotypic responses to temperature in the ciliate Tetrahymena thermophila. *Ecology and Evolution,* **10,** 7616-7626.

Wieczynski, D.J., Singla, P., Doan, A., Singleton, A., Han, Z., Votzke, S., Yammine, A. & Gibert, J.P. (2021) Simple traits predict complex temperature responses across ecological scales.

Yamamichi, M., Yoshida, T. & Sasaki, A. (2011) Comparing the effects of rapid evolution and phenotypic plasticity on predator-prey dynamics. *The American Naturalist,* **178,** 287-304.