Epigenetic modulation of selected immune response genes and altered functions of T lymphocytes and macrophages collectively contribute to autoimmune diabetes protection

Sundararajan Jayaraman[a,b*], Maria Arianas[a], Arathi Jayaraman[a],

[a]Dept. of Surgery, the University of Illinois at Chicago, Chicago, IL 60612

[b]Current address: Dept. of Surgery, University of Illinois College of Medicine at Peoria, IL 61603; Telephone: 630-209-9685.

*Correspondence: anue2468@uic.edu

Abbreviations:

HDAC, histone deacetylase; qRT-PCR, quantitative reverse transcriptase-mediated polymerase chain reaction; T1D, type 1 diabetes; Th, T helper, TSA, Trichostatin A.

ABSTRACT

We have previously demonstrated that treatment of female NOD mice with the histone deacetylase inhibitor Trichostatin A (TSA) bestowed irreversible protection against autoimmune diabetes. Herein we show that drug treatment diminished the infiltration of the pancreas with CD4$^+$ and CD8$^+$ T cells and Ly-6C$^+$ monocytes. Significantly, TSA administration selectively repressed the expression of a set of genes exaggerated during diabetes and constitutively expressed primarily in the spleen and rarely in the pancreas. These genes encode lymphokines, macrophage-associated determinants, and transcription factors. Although the copy numbers of many histone deacetylases increased during diabetes in the spleen and pancreas, only those upregulated in the spleen were rendered sensitive to repression by TSA treatment. The T lymphocytes derived from drug-treated donors displayed diminished diabetogenic potential following transfer into immunodeficient NOD.*scid* mice. In the immunocompromised recipients, diabetes caused by the transfer of activated T lymphocytes from untreated diabetic mice was hampered by the co-transfer of highly purified splenic Ly-6C$^+$ macrophages from drug-treated mice. However, the transfer of Ly-6C$^+$ macrophages from drug-treated mice failed to block ongoing diabetes in wild-type NOD mice. These data demonstrate that the modified gene expression and functional alteration of T lymphocytes and macrophages collectively contribute to diabetes protection afforded by the histone modifier in female NOD mice.

Keywords: Epigenetics, Genes, Histone deacetylases, Lymphocytes, Macrophages, Trichostatin A, Type 1 Diabetes.

1. Introduction

Type 1 diabetes (T1D), a T cell-mediated autoimmune disease, occurs at an alarming rate in genetically susceptible individuals [1-3]. Although genome-wide association studies implicated more than 40 different genetic loci, the causal genes of T1D remain unknown [4]. Lack of familial history and robust concordance among identical twins in developing T1D implied the involvement of environmental factors. Epigenetics provides a mechanism by which external factors can produce a variety of phenotypic variations with identical genotypes. Epigenetic mechanisms include DNA methylation, post-translational modifications of histones, and microRNA-mediated gene regulation [5]. Methylation of cytosine residues in gene promoters results in transcriptional silencing of some [6] but not all genes [7]. Reversible acetylation of the e-amino group of lysine in the histone tails by histone acetyltransferases and deacetylation by histone deacetylases (HDACs) are the best-characterized post-translational modifications of histones [8]. Transcriptional permissiveness correlates with histone acetylation in contrast to deacetylation resulting in gene repression. Non-selective, small molecule HDAC inhibitors can induce chromatin modifications and thereby affect gene expression. Trichostatin A (TSA) derived from *Streptomyces platensis* is the most potent inhibitor of class I, class IIa, class IIb, and class IV HDACs, which increases histone acetylation resulting in up- and down-regulation of genes *in vitro* [9] and *in vivo* [10-16].

Previously, we have demonstrated the utility of TSA to mitigate autoimmune diseases such as T1D [10-13] and experimental autoimmune encephalomyelitis [14-16]. The protection against T1D afforded by the histone modifier was accompanied by reduced inflammation of the pancreatic islets and regeneration of insulin-producing β-cells [10]. Significantly, treatment with TSA abrogated the ability of splenocytes to adoptively transfer T1D into immunodeficient

NOD.*scid* mice, indicating the effect of the epigenetic drug on diabetogenic T lymphocytes [11]. Transcriptome analysis of *uninduced* splenocytes unraveled the exaggerated expression of inflammatory genes, including the macrophage selective *Cela3b* (elastase 3) in diabetic mice repressed by TSA. These results are consistent with the possibility that chromatin remodeling by TSA can retard the induction and progression of T1D by altering the transcriptional programs of multiple cell types. In the current investigation, we analyzed the effects of the epigenetic drug on T lymphocytes and macrophages, respectively implicated in induction and manifestation of T1D [17-19], by quantitative reverse transcriptase-mediated polymerase chain reaction (qRT-PCR) and an adoptive transfer model. Our study unraveled that a select set of both exaggerated and constitutively expressed genes encoding cytokines, cell surface determinants, transcription factors, and HDACs was modulated by the histone modifier more predominantly in the spleen. Our data also demonstrate the distinct effects of TSA on the function of T-cells and macrophages. These data lend novel insights into the mechanisms of T1D and its possible manipulation by epigenetic reprogramming.

2. Materials and materials

2.1. Diabetes treatment

The Office of Animal Care and Institutional Biosafety of the University of Illinois at Chicago approved the animal protocol. Experiments were conducted following the NIH guide for the care and use of laboratory animals.

Six to eight weeks old female NOD/ShiLtj (H-2^{g7}) mice purchased from The Jackson Laboratory (Bar Harbor, ME) were maintained under specific pathogen-free conditions and provided with animal chow and water *ad libitum*. Mice were injected s.c. with 500 µg/Kg body

weight of TSA or DMSO at weekly intervals between 16 and 24 wks of age [10-11]. Non-fasting blood glucose level exceeding 250 mg/dL on two consecutive weekly determinations was considered as diabetic.

2.2 Confocal microscopy

The following antibodies were used for staining pancreatic sections: guinea pig anti-insulin Ab (Zymed Laboratories, South San Francisco, CA), tetramethylrhodamine iso-thiocyanate–rabbit antisera raised against guinea pig Ig (Sigma Aldrich, St. Louis, MO), anti-Ly-6C-FITC (clone HK1.4, eBiosciences, San Diego, CA), anti-Ly-6G-FITC (clone 1AB, BD Pharmingen, San Diego, CA), rat monoclonal anti-CD4 (clone RM4-5, eBiosciences-ThermoFisher Scientific, Waltham, MA), DyLight™ 488 goat anti-rat IgG (catalog # 405409, BioLegend, San Diego, CA) and anti-CD8-FITC (clone 53-6.7, eBiosciences). Confocal images were acquired using a Zeiss LSM510 laser scanning microscope and processed using the Zeiss LSM Image browser (4.0 version; Zeiss, Oberkochen, Germany) and Adobe Photoshop Elements version 9.0, as described [20].

2.3 Gene expression analysis

Spleens were harvested from 20-24-wks old diabetic mice and 26-28 wks old mice treated with TSA from 16 to 24-wks of age. Total RNA was extracted from the spleen and pancreas of five *individual mice* per group/experiment, treated with Turbo DNase, and converted to cDNA using High-capacity cDNA Reverse Transcription kit (Applied Biosystems), as described [10-11, 14-16, 20]. Real-time qRT-PCR was performed in *triplicate* on Applied Biosystems ViiA7 Real-time PCR system using 1 µl of cDNA equivalent to 100 ng of total RNA and 2X SYBR Green master mix. The primers used in this study were designed and validated

previously [10-11, 14-16, 20] and synthesized at the Integrated DNA Technologies (Coralville,

IA). The level of specific gene expression in each sample was ascertained using *Gapdh* as the

normalizer and the $2^{-\Delta\Delta C}T$ method.

2.4 Adoptive transfer of diabetes

Spleens were harvested from 20-24-wks old diabetic mice and from 26-28 wks old mice

treated with TSA between 16 to 24-wks of age. Splenocytes (10×10^6/ml) were cultured with

five µg of Concanavalin A (Sigma Aldrich) for two days as described [19]. More than 95% of

these cells were CD3$^+$, as determined by flow cytometry. Splenocytes were also stained with PE-

Cyanine 5 conjugated anti-CD11b (clone M1/70, eBiosciences) and anti-Ly-6C-FITC antibodies

and sorted on a MoFlo Astrios cell sorter, which yielded 2.27 % of the input with > 98%

CD11b$^+$Ly-6C$^+$ cells. Both male and female NOD.*scid* mice were injected i.v. with activated 2 x

10^6 T lymphocytes with or without sorted 2×10^5 CD11b$^+$Ly-6C$^+$ cells or purified monocytes

alone. Splenocytes were also stained with anti-Ly-6C-FITC and incubated with anti-FITC

antibody and purified on a magnetic column (Miltenyi Biotec Inc., Auburn, CA). This procedure

yielded 2 % of the input with >95% Ly-6C$^+$ cells, as determined by flow cytometry. Female

wild-type NOD mice received two injections of purified 2×10^5 Ly-6C$^+$ cells at 13 and 21-wk of

age. Numbers of mice tested are given in the figure legends.

2.5 Statistics

The difference in diabetes incidence between controls (untreated/DMSO-treated mice)

and those treated with TSA was analyzed for statistical significance using the Wilcoxon Signed

Rank Test. Gene expression data were analyzed for statistical significance between indicated

groups by two-way ANOVA or two-tailed unpaired *t*-test using GraphPad Prism (6.0) software and indicated in individual figure legends. A *P*-value of <0.05 was considered significant.

3. Results

3.1 Protection from T1D is associated with constrained infiltration of inflammatory cells into the islets.

We have consistently observed that 80-100% of female NOD mice procured from The Jackson Laboratory and housed in our animal facility developed T1D by 18 weeks of age [10-13, 20]. Weekly treatment with TSA starting at 16 wk and until 24 wks of age substantially attenuated T1D (Supplementary Fig. 1). Injection of the vehicle DMSO failed to alter the course or intensity of the disease [10-13]. We have also documented that DMSO treatment did not change gene expression compared to untreated diabetic mice. Since DMSO-treated mice did not have noticeable changes in disease course and gene expression compared to untreated diabetic mice, we compared the litter mates of sex- and age-matched untreated diabetic mice with TSA-treated mice in all experiments. Treatment with TSA reduced the invasive cellular infiltration of the islets and reversed moderate to a severe loss of insulin-producing beta cells [10]. However, the nature of cellular infiltration was not previously characterized. To this end, we performed confocal analysis, which indicated the distribution of Ly-6C$^+$ macrophages, Ly-6G$^+$ neutrophils, CD4$^+$, and CD8$^+$ T-cells across the entire area of the diabetic pancreas (Fig. 1). Treatment with TSA significantly reduced the infiltration of primarily CD4$^+$ cells and, to some extent, CD8$^+$ T-cells implicated in T1D [17]. The influx of Ly-6C$^+$ macrophages but not neutrophils was also reduced by TSA treatment, consistent with a role for macrophages in diabetes [18-19]. Flow cytometric analysis confirmed that the numbers of CD11b$^+$Ly-6C$^+$ monocytes but not

CD11b$^+$F4/80$^+$ macrophages or CD11c$^+$ dendritic cells were also diminished in the spleen of TSA-treated mice (Supplementary Fig. 2). These data indicated that reduced influx of CD4$^+$ and CD8$^+$ T cell subsets and Ly-6C$^+$ macrophages into the islets serves as a mechanism of TSA-mediated amelioration of autoimmune diabetes.

3.2 TSA treatment selectively repressed the expression of Th1 and Th17 lymphokines, primarily in the spleen.

Our previous transcriptome analysis of *uninduced* splenocytes indicated the downregulation of a few inflammatory genes, including *Cela3b* (elastase) by TSA treatment [11]. In the current investigation, we analyzed a total of 43 genes selected for their putative roles in T1D. These included genes encoding 11 lymphokines, 13 accessory cell-associated determinants, seven transcription factors, a chemokine, and 11 genes encoding HDACs. Total RNA was extracted from diabetic and TSA-treated animals, converted to cDNA, and used as the template in the qRT-PCR assay as described before [10-16]. The genes critical for Th17 cell development, namely *Il17a* and *Il23* [21-22], and the Th1-specific *Ifng* [23] were significantly increased in the spleen of diabetic mice compared to the prediabetic mice displaying normal levels of blood glucose (Fig. 2a). Significantly, TSA treatment diminished the levels of these genes. Although *Il4*, *Il18*, and *Il27p28* were not exaggerated during diabetes, they were nevertheless repressed by treatment with the histone modifier. However, other genes such as *Il22*, *Il27ebi3*, *Il10*, *Il12*, and *Tgfb* were neither upregulated during diabetes nor modulated by the histone modifier, indicating that they are unlikely to contribute to diabetogenesis in this model.

Preliminary data indicated that the expression of the genes analyzed herein remained comparable in the pancreas of prediabetic and overtly diabetic mice. Therefore, the gene expression levels in the pancreas of diabetic and TSA-treated mice were compared. Unlike the spleen, TSA treatment increased the level of *Il4* but repressed the transcription of *Il22* and *Il23* in the pancreas (Fig. 2b). Collectively, these results suggest that TSA treatment differentially regulated the expression of both inducible and constitutively expressed immune response-related genes in the spleen and pancreas, and inversely correlated with normoglycemia.

3.3 Histone modifier regulates constitutive expression of macrophage-associated cytokines primarily in the spleen.

The chemokine CCL2, implicated in the mobilization of myeloid cells from the bone marrow causing insulitis [24], was upregulated in the spleen of diabetic mice, which was repressed by TSA treatment (Fig. 3a). Although macrophage-selective genes such as *Csf2*, *Mmp12*, and *Ym1* were not exaggerated during diabetes and yet were reduced by the histone modifier. Significantly, *Cela3b* expression was lower in TSA-treated spleens, as shown earlier [11]. Only the steady-state level of *Cd39* was subject to regulation by the HDAC inhibitor in the pancreas (Fig. 3b). These data suggest that TSA treatment predominantly diminished the constitutive expression of many macrophage-selective genes in the spleen compared to the pancreas.

3.4 Epigenetic drug impacts the expression of transcription factors sparingly in the spleen.

Notably, the transcription factor *Ahr* involved in nitric oxide and arginine production, and alteration of M1/M2 macrophage polarization [25] was transcriptionally increased in the spleen of diabetic mice and reduced following TSA treatment (Fig. 4a). Only the constitutive expression

of *Rorgt*, crucial for the transcription of IL-17A [26], was repressed by TSA treatment.

Surprisingly, *Tbet* and *Gata3*, respectively involved in the transcription of IFN-γ and IL-4, and other transcription factors such as *Eomes*, *Dec1*, and *Foxp3* examined were neither transcriptionally upregulated during diabetogenesis nor subject to epigenetic regulation. Interestingly, the *Foxp3* transcript level was lower in the TSA-treated pancreas (Fig. 4b). These data suggest that TSA-mediated protection from diabetes is associated with selective repression of IL-17A transcription in the lymphoid organ.

3.5 Epigenetic modifier differentially modulated HDACs in the spleen and pancreas.

Although *in vivo* treatment of NOD mice with TSA induced the hyperacetylation of histone H3 in the spleen and pancreas of NOD mice [11], it is unclear whether it can differentially affect the expression of class I (*Hdac1*, *Hdac2*, *Hdac3*, and *Hdac8*), class IIa (*Hdac4*, *Hdac5*, *Hdac7*, and *Hdac9*), class IIb (*Hdac6* and *Hdac10*) or class IV (*Hdac11*) HDAC genes in the lymphoid system and the target organ pancreas. In the spleen of prediabetic, normoglycemic, 12-wk old female NOD mice, the basal levels of the *Hdac5* and *Hdac8* were minimal (Fig. 5a). During diabetes manifestation, the transcription of *Hdac4*, *Hdac8*, and *Hdac9* genes was increased, which was repressed by TSA treatment. Interestingly, *Hdac1*, *Hdac3*, *Hdac4*, *Hdac8*, and *Hdac11* were upregulated in the pancreas of diabetic mice (Fig. 5b). Whereas most of these genes were not repressed by the histone modifier, *Hdac4* was upregulated in drug-treated mice. It is noteworthy that *Hdac2*, *Hdac5*, *Hdac7*, *Hdac9*, and *Hdac10* genes could not be reliably detected in the pancreas of prediabetic, overtly diabetic, or TSA-treated mice by qRT-PCR in repeated experiments (n=25-30 mice). Thus, TSA-mediated protection against T1D was accompanied by reduction of the exaggerated expression of *Hdac4*, *Hdac8*, and *Hdac9* genes in the spleen, while *Hdac4* was upregulated in the pancreas.

3.6 TSA treatment differentially affected the function of T lymphocytes and macrophages.

We next determined whether epigenetically modulated genes could impact the diabetogenic potential of T lymphocytes and the presumptive effector functions of macrophages in diabetes [18-19]. The data shown in Fig. 6a indicate that the transfer of Concanavalin A-activated T lymphocytes from diabetic mice mediated the disease in 100% of the immunodeficient NOD.*scid* mice by nine wks. However, similarly activated T lymphocytes derived from TSA-treated mice failed to induce the disease in NOD.*scid* recipients, consistent with the data obtained using whole splenocytes [11]. Interestingly, the adoptive transfer of sorted CD11b$^+$Ly-6C$^+$ macrophages from diabetic or TSA-treated mice did not cause diabetes in the immunodeficient mice, indicating that the macrophages are not the effectors of T1D in contrast to a previous observation indicating the beta-cell cytotoxicity *in vitro* by macrophages derived from diabetic mice [19]. Surprisingly, the co-transfer of CD11b$^+$Ly-6C$^+$ macrophages purified from the TSA-treated but not diabetic mice suppressed the T-cell mediated diabetes in NOD.*scid* recipients. In contrast to the data obtained in an adoptive transfer model, two consecutive injections of splenic Ly6-C$^+$ macrophages purified from TSA-treated or diabetic mice failed to impede the ongoing disease in wild-type female NOD mice (Fig. 6b), inconsistent with diabetes suppression mediated by polarized M2-type macrophages reported previously [27]. These data suggest that the epigenetic modulation of macrophages resulted in distinct outcomes dictated by the recipients' immune status.

4. Discussion

Although many intervention strategies have been previously reported to delay the diabetes onset in female NOD mice, the success of translating these studies into T1D treatment

remains extremely slim [28]. We have reported that TSA treatment just prior to the onset of

diabetes provided irreversible protection against diabetes in female NOD mice [10-13].

Protection was associated with increased histone H3 hyperacetylation in both the spleen and

pancreas [10]. We now show that diabetes protection is associated with reduced infiltration of

$CD4^+$, $CD8^+$, and Ly6-C^+ cells in the pancreas (Fig. 1), indicating specific action of the

chromatin modifier on specific immune cell types. This is consistent with reduced transcription

and expression of IFN-γ and IL-17A in the peripheral lymphoid organ following TSA

treatment.* These data are congruent with the notion that the protective effects of the histone

modifier against autoimmune diabetes involves reduction in the generation of T helper subsets

and compromised homing of the macrophages and T-cell subsets to the islets of Langerhans.

For the first time, we show herein that diabetogenesis involves exaggerated expression of

Hdac4, *Hdac8*, and *Hdac9* genes in the secondary lymphoid organ, which were repressed by the

histone modifier (Fig. 5a). Although many *Hdac* genes were also exaggerated in the pancreas of

diabetic mice, surprisingly, they were resistant to TSA treatment (Fig. 5b). This was not due to

the lack of access of TSA to the pancreas since *Hdac4* was enhanced in the pancreas of drug

treated mice. Thus, the *Hdac* genes are differentially regulated by histone hyperacetylation in the

lymphoid tissue *vs.* the target organ, consistent with the active role of T-cells in diabetes

induction. Interestingly, we noted that the *Hdac3* gene was not repressed either in the spleen or

pancreas by extremely low doses of TSA (Fig. 5a-b), which is at odds with the finding that

administering substantial amounts of the HDAC3 selective inhibitor BRD3308 conferred

marginal protection against diabetes in NOD mice [29]. It is unclear whether the BRD3308

inhibitor can exert immunosuppressive effects as TSA [10-16, Fig. 1-6]. However, BRD3308

reduced cytokine-induced apoptosis of dispersed primary islets *in vitro* and improved glycemia

and insulin secretion in obese diabetic rats [30-32]. However, evidence for suppressing

autoimmune responses by BRD3308 is lacking. Interestingly, TSA blocked the inflammatory

cytokine-mediated beta-cell toxicity *in vitro* more effectively than the BRD3308 inhibitor [32].

Since the most potent HDAC inhibitor, TSA is endowed with both immunomodulatory [10-16,

Fig. 1-6] and beta-cell protective effects [32], it seems to be an ideal drug of choice for treating

autoimmune diabetes with an extremely low dose regimen compared to a large amount of the

HDAC3 selective BRD3308 inhibitor required to exert marginal protection [29].

An important finding is that out of 32 immune response-related genes investigated, only a

few were exaggerated during diabetes and regulated by TSA treatment. Notably, *Ifng* and *Il17a*,

respectively the signature genes of Th1 and Th17 subsets, were magnified during diabetes (Fig.

2a). Although IFN-γ-producing Th1 cells were shown to mediate diabetes in neonatal NOD mice

[33], the genetic absence of IFN-γ delayed but did not prevent diabetes [34]. Paradoxically, a

protective role of IFN-γ in diabetes was also reported [35-36]. Thus, the role of IFN-γ in diabetes

remains controversial. Whereas IL-17A was shown to contribute to diabetes [37], others raised

questions about the independence of Th17 cells to induce diabetes [38-39]. Interestingly, T1D

onset was shown to depend on both IL-17 and IFN-γ signaling [40]. Our data showing the

repression of both *Il17a* and *Ifng* in TSA-treated mice that attained normoglycemia supports this

contention. A closer analysis unraveled that the TSA treatment prevented the upregulation of

Il23, required for the generation of Th17 cells [22] (Fig. 2) and *Ccl2* implicated in diabetes [24]

(Fig. 3). A novel finding is that the gene encoding the transcription factor *Ahr* [25] was

accentuated in the spleen during diabetes and repressed by the histone modifier (Fig. 4),

indicating a crucial for this transcription factor in autoimmune diabetes manifestation. In

addition to these upregulated genes, the steady-state levels of several genes implicated in T1D

such as *Il4* [41], *Il18* [42], and *Il27p28* [43] were also diminished in TSA-treated spleen (Fig. 2).

Similarly, the basal level expression of the transcription factor *Rorgt*, crucial for IL-17A

transcription [26], was suppressed in the spleen of drug-treated mice. Interestingly, *Il22*, *Il23*,

and *Cd39* implicated in diabetes [44] were similarly repressed in the pancreas of mice cured of

diabetes. Interestingly, *Foxp3*, crucial for immunoregulation [45], was decreased in the pancreas

after TSA treatment (Fig. 4), indicating that the regulation of T1D is independent of

CD4$^+$FoxP3$^+$ cells, as we reported earlier [11]. This is similar to a model of T1D delay mediated

by the administration of the migration inhibitory factor antagonist that repressed Foxp3

expression [46]. Overall, our data provide supporting evidence for the contribution of previously

known genes such as *Ifng*, *Il17a*, and *Ccl2*, and importantly implicate novel roles of *Il23* and *Ahr*

in diabetes manifestation, more so in lymphoid tissue than in the pancreas. Some of these genes

upregulated during diabetogenesis may be the effect rather than the cause of diabetes. Further

work is necessary to parse the causative role of these genes in autoimmune diabetes.

The diminished diabetogenic potential of whole splenocytes [10] and the enriched T

lymphocytes (Fig. 6a) by epigenetic reprogramming reiterates the importance of T-cells in

diabetes*. In addition to T lymphocytes, macrophages had been proposed to play a crucial role

during the effector phase of T1D [18-19]. Both antigen presentation to cytotoxic CD8$^+$ T cells

[18] and direct cytotoxicity on dispersed islet cells *in vitro* [19] have been attributed to

macrophages. However, direct evidence for the actual participation of macrophages in

diabetogenesis *in vivo* has been difficult to establish. We observed that the CD11b$^+$Ly-6C$^+$

macrophages from diabetic mice failed to transfer diabetes to immunodeficient mice, indicating

that the macrophages do not serve as effectors of T1D (Fig. 6a), unlike proposed previously [19].

Unexpectedly, we found that TSA treatment endowed the CD11b$^+$Ly-6C$^+$ macrophages with the

ability to suppress disease induction upon co-transfer with diabetogenic T cells into NOD.*scid*

recipients (Fig. 6a). However, the transfer of Ly-6C$^+$ macrophages derived from TSA-treated

mice failed to suppress ongoing diabetes in wild-type NOD mice (Fig. 6b). This paradoxical

finding appears similar to the failure of polarized Th2 [41] and Th17 cells [38] derived from the

BDC2.5 TCR transgenic mice to transfer diabetes into neonatal wild-type NOD mice while

readily transferring the disease to immunodeficient NOD.*scid* mice. These intriguing findings

were attributed to the lack of endogenous immunoregulatory mechanisms in immunodeficient

mice facilitating the action of transferred cells without opposition, unlike the wild-type NOD

mice. Our data are also different from diabetes suppression in NOD mice mediated by the

injection of M2-type macrophages polarized *in vitro* from bone-marrow-derived monocytes

using a cocktail of IL-4/IL-10/TGF-β [27]. Apparently, TSA treatment did not convert

macrophages into M2-type macrophages. Therefore, this strategy, transfer of macrophages from

TSA-treated mice may not be useful for T1D treatment. Although histone hyperacetylation has

been proposed to impact monocytes and macrophages central to tissue homeostasis and immune

responses [47-48], little evidence exists about the functional modulation of macrophages under

diabetic conditions. Previously, we observed that NOD mice treated with TSA and cured of T1D

displayed a lower level of the pro-inflammatory gene *Cela3b* (elastase 3) [11], which was

validated herein (Fig. 3a). Moreover, TSA treatment downregulated macrophage-specific genes

such as *Csf2*, *Mmp12*, and *Ym1*. Further studies are required to determine whether these changes

can contribute to the suppressive function of macrophages observed in the lymphopenic

environment, NOD.*scid* mouse.

Conclusions

The data presented herein and elsewhere [10-13*] indicate that the histone modifier mediated protection against T1D involves the modified expression of a select set of immune response-related genes such as *Il17a*, *Ifng*, *Ccl2*, and *Ahr* in the spleen and altered functions of T cells and macrophages. However, changes in the expression of substantial numbers of immune response genes analyzed in the pancreas of TSA-treated mice remained minimal compared to the lymphoid tissue. This is consistent with a more active role of lymphoid cells in the induction and manifestation of autoimmune diabetes. Although we have analyzed 32 immune response-related genes in the pancreas of diabetic mice and their manipulation by TSA treatment, genes pertinent to the exocrine and endocrine tissues remain to be scrutinized. Since T1D induction is dependent on T lymphocytes, our epigenetic analysis has focused on the Th1 and Th17 subsets, allowing the parsing of the roles of, respectively, *Ifng* and *Il17a*.* The disease protection afforded by TSA treatment is inversely correlated with the levels of a small number of genes mainly in the lymphoid tissue, suggesting that they could serve as biomarkers of T1D and targets for manipulation of the disease. We believe that similar analysis of multiple immune response-related genes and inflammatory genes in T1D patients may provide significant insights into disease pathogenesis.

Acknowledgements

Mark Holterman is acknowledged for the support of this work. The expert help of the Imaging Core and Flow Cytometry Facility at the University of Illinois at Chicago in acquiring confocal images and high-speed sorting of macrophages, respectively is acknowledged.

Author contributions

CRediT roles:

Sundararajan Jayaraman: Conceptualization, data curation, formal analysis, funding acquisition, methodology, project administration, resources, supervision, validation, writing original draft, review & editing.

Maria Arianas: Methodology, review & editing.

Arathi Jayaraman: Methodology, review & editing.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Figure legends

Fig. 1 Cellular infiltration in the pancreas was retarded by TSA treatment

Diabetic mice and those treated with TSA during 16 and 24-wk of age were sacrificed when they were 26-28-wks old. Sections of the pancreas were stained with guinea pig antisera against insulin followed by TRITC labeled rabbit anti-guinea pig antibody (red), along with FITC-conjugated anti-Ly-6C, anti-Ly-6G, anti-CD4 or anti-CD8 antibodies (green). After counterstaining with Hoechst (blue), the sections were analyzed on a confocal microscope. Representative images from multiple experiments are shown.

Fig. 2 Differential modulation of lymphokine genes by the histone modifier

Spleen was harvested from 12-wk old prediabetic mice and 25-26-wks old diabetic mice. Pancreas was collected from 25-26-wks old diabetic mice. Mice treated with TSA between 16

and 24-wks of age were sacrificed when they were 25-26-wks old and spleen and pancreas were harvested. Five mice per group were tested in three different experiments. Total RNA was extracted from the spleen (**a**) and pancreas (**b**) of five individual mice was analyzed, converted to cDNA, and used as the template in qRT-PCR. Data indicate the mean +/- SD. Representative data from three independent experiments are depicted. Asterisks indicate statistical significance in the spleen samples from prediabetic, diabetic, and TSA-treated mice, as assessed by two-way ANOVA (**a**) and by two-tailed unpaired *t*-test between diabetic and TSA-treated pancreas (**b**) ($P<0.05$). Increased values between prediabetic and diabetic mice were indicated by double asterisks whereas a single asterisk denoted the decrease in values in TSA-treated mice compared to diabetic mice.

Fig. 3 Differential regulation of accessory cell-associated determinants following TSA treatment

The expression levels of genes associated with accessory cells were analyzed in (**a**) the spleen and (**b**) pancreas. Total RNA was extracted and gene expression was analyzed by qRT-PCR using the same cDNA preparations analyzed in Fig. 2. Data indicate arithmetic mean +/- SD of three determinations. Representative data from three independent experiments are depicted (*n*=5 mice per group/experiment). Asterisks indicate statistical significance in the spleen samples from prediabetic, diabetic, and TSA-treated mice, as assessed by two-way ANOVA (**a**) and by two-tailed unpaired *t*-test between diabetic and TSA-treated pancreas (**b**) ($P<0.05$). Increased values between prediabetic and diabetic mice were indicated by double asterisks, whereas the decrease in values in TSA-treated mice compared to diabetic mice was denoted by a single asterisk.

Fig. 4 Selective regulation of the transcription factor genes by TSA

The expression levels of genes encoding various transcription factors were analyzed in (**a**) the spleen and (**b**) pancreas by qRT-PCR using the same cDNA preparations analyzed in Fig. 2 and 3. Data indicate the mean +/- SD of three determinations. Representative data from three independent experiments are depicted (*n*=5 mice per group/experiment). Double asterisks indicate a statistically significant increase between prediabetic and diabetic mice (*P<0.05), whereas a single asterisk denotes a decrease in the value between diabetic and TSA-treated groups (**P<0.01). Statistical significance was assessed respectively by two-way ANOVA (**a**) and two-tailed unpaired *t*-test between diabetic and TSA-treated pancreas (**b**) as directed by the GraphPad Prism software.

Fig. 5 Differential modulation of *Hdac* genes by the epigenetic drug

The expression levels of *Hdac* genes in the spleen (**a**) and pancreas (**b**) were determined in cDNA samples illustrated in Fig. 2-4. Data indicate the mean +/- SD of three determinations. Representative data from three independent experiments are depicted (*n*=5 mice per group/experiment). An increase in the values between prediabetic and diabetic mice was indicated by double asterisks. The decrease in values in TSA-treated mice compared to diabetic mice was denoted by a single asterisk. Statistical significance was assessed using two-way ANOVA (**a**) and by a two-tailed unpaired *t*-test between diabetic and TSA-treated pancreas (**b**). *P<0.05 and **P<0.01.

Fig. 6 Epigenetic regulation of T cells and macrophages

(**a**) Diabetes was monitored in NOD.*scid* mice (10-12-wks old) injected i.v. with Concanavalin A-activated 2×10^6 T lymphocytes from diabetic or TSA-treated NOD mice (26-28-wks of age) (P=0.0027). Mice were also co-transferred with diabetic T-cells and 2×10^5 sorted CD11b$^+$Ly-6C$^+$ macrophages from the spleens of diabetic mice or those treated with TSA and diabetes monitored (P=0.003). Some mice received only macrophages isolated from diabetic mice or TSA-treated mice. Four mice per group were tested. Representative data from two independent experiments are shown. Statistical significance between specified groups was assessed using a two-tailed, unpaired *t*-test. (**b**) Wild-type female NOD mice were injected with purified 2×10^5 Ly6-C$^+$ macrophages from diabetic mice or those treated with TSA at 13 and 21-wks of age, as indicated by the arrows. Diabetes was monitored weekly. Five mice per group were tested. Data shown are from two independent experiments.

5. References

[1]   American Diabetes Association, Diagnosis and classification of diabetes mellitus, Diabetes Care, 30 (2007) Suppl 1: S42-47. https://doi.org/10.2337/dc07-S042.

[2]   V. Hyttinen, J. Kaprio, L. Kinnunen, M. Koskenvuo, J. Tuomilehto, Genetic liability of type 1 diabetes and the onset age among 22,650 young Finnish twin pairs: a nationwide follow-up study. Diabetes, 52 (2003)1052-1055. https://doi.org/10.2337/diabetes.52.4.1052.

[3]   J.E. Shaw, R.A. Sicree, P.Z. Zimmet PZ, Global estimates of the prevalence of diabetes for 2010 and 2030. Diabetes Res. Clin. Pract. 87 (2010) 4-14. https://doi.org/10.1016/j.diabres.2009.10.007.

[4]   J.A. Todd, Etiology of type 1 diabetes, Immunity, 32 (2010) 457–467. https://doi.org/10.1016/j.immuni.2010.04.001.

[5]   C.D. Allis, T. Jenuwein, The molecular hallmarks of epigenetic control, Nat. Rev. Genet. 17 (2016) 487-500. https://doi.org/10.1038/nrg.2016.59.

[6]   A. Bird, Genetic determinants of the epigenome in development and cancer, Swiss Med. Wkly. 147 (2017) w14523. https://doi.org/10.4414/smw.2017.14523.

[7]   A. Jayaraman, S. Jayaraman, DNA hypermethylation does not negatively impact the transcription of the TNF-α gene in an acute T-cell leukemia, Epigenomics, 11 (2019) 1753-1763. https://doi.org/10.2217/epi-2019-0015.

[8]   C. Micelli, G. Rastelli, Histone deacetylases: structural determinants of inhibitor selectivity, Drug Discov. Today, 20 (2015) 718-735. https://doi.org/10.1016/j.drudis.2015.01.007.

[9]   J. de Ruijter, A.H. van Gennip, N.H. Caron, S. Kemp, A.B. van Kuilenburg, Histone

deacetylases (HDACs): characterization of the classical HDAC family, Biochem. J. 370

(2003) 737–749. https://doi.org/10.1042/BJ20021321.

[10]   T. Patel, V. Patel, R. Singh, S. Jayaraman, Chromatin remodeling resets the immune

system to protect against autoimmune diabetes in mice, Immunol. Cell Biol. 89 (2011)

640–649. https://doi.org/10.1038/icb.2010.144.

[11]   S. Jayaraman, A. Patel, A. Jayaraman, V. Patel, M. Holterman, B. Prabhakar B,

Transcriptome analysis of epigenetically modulated genome indicates signature genes in

manifestation of type 1 diabetes and its prevention in NOD mice. PLoS One, 8 (2013)

e55074. https://doi.org/10.1371/journal.pone.0055074.

[12]   S. Jayaraman, Novel methods of type 1 diabetes treatment, Discov. Med. 17 (2014), 347-

355. PMID: 24979255.

[13]   S. Jayaraman, Type 1 diabetes and epigenetics, in: R. Zhang (Ed.), The epigenetics of

autoimmunity. Translational Epigenetics Series, Volume 5. Elsevier Inc., Amsterdam,

The Netherlands, 2018, pp. 188-205.

[14]   A. Jayaraman, A. Soni, B.S. Prabhakar, M. Holterman, S. Jayaraman, The epigenetic

drug Trichostatin A ameliorates experimental autoimmune encephalomyelitis via T cell

tolerance induction and impaired influx of T cells into the spinal cord, Neurobiol. Dis.

108 (2017) 1-12. https://doi.org/10.1016/j.nbd.2017.07.015.

[15]   A. Jayaraman, M. Sharma, B. Prabhakar, M. Holterman, S. Jayaraman, Amelioration of

progressive autoimmune encephalomyelitis by epigenetic regulation involves selective

repression of mature neutrophils during the preclinical phase, Exp. Neurol. 304 (2018)

14-20. https://doi.org/10.1016/j.expneurol.2018.02.008.

[16]   A. Jayaraman, K. Avgush, R. Kulam, A. Soni, A. Khan, M. Kerdjoudj, S. Jayaraman, Treatment of autoimmune encephalomyelitis with a histone deacetylase inhibitor. Analyzing the role of immune-response genes, Free Neuropathology (2020) 1:19. https://doi.org/10.17879/freeneuropathology-2020-2819.

[17]   S.W. Christianson, L.D. Shultz, E.H. Leiter, Adoptive transfer of diabetes into immunodeficient NOD-scid/scid mice. Relative contributions of CD4+ and CD8+ T-cells from diabetic versus prediabetic NOD.NON-Thy-1a donors, Diabetes, 42 (1993) 44-55. https://doi.org/10.2337/diab.42.1.44.

[18]   H.S. Jun, C.S, Yoon, L. Zbytnuik, N. van Rooijen, J.W. Yoon, The role of macrophages in T cell-mediated autoimmune diabetes in nonobese diabetic mice. J. Exp. Med. 189 (1999) 347□358. https://doi.org/10.1084/jem.189.2.347.

[19]   B. Calderon, A. Suri, E.R. Unanue, In CD4+ T-cell-induced diabetes, macrophages are the final effector cells that mediate islet beta-cell killing: studies from an acute model, Am. J. Pathol. 169, (2006) 2137–2147. https://doi.org/10.2353/ajpath.2006.060539.

[20]   S. Jayaraman, T. Patel, V. Patel, S. Ajani, R. Garza, A. Jayaraman, S. Kwon, R. Singh, D. Rondelli, B.S. Prabhakar, M. Holterman, Transfusion of nonobese diabetic mice with allogeneic newborn blood ameliorates autoimmune diabetes and modifies the expression of selected immune response genes, J. Immunol. 184 (2010) 3008-3015. https://doi.org/10.4049/jimmunol.0903615.

[21]   S.L. Gaffen, Recent advances in the IL-17 cytokine family, Curr. Opin. Immunol. 23, (2011) 613□619. https://doi.org/10.1016/j.coi.2011.07.006.

[22]   M.J. McGeachy, D.J. Cua, The link between IL-23 and Th17 cell-mediated immune

        pathologies, Semin. Immunol. 19 (2007) 372□376.

        https://doi/10.1016/j.smim.2007.10.012.

[23]   A. Skapenko, H. Schulze-Koops, Analysis of Th1/Th2 T-cell subsets, Methods Mol.

        Med. 136 (2007) 87□96. https://doi.org/10.1007/978-1-59745-402-5_7.

[24]   A.P. Martin, S. Rankin, S. Pitchford, I.F. Charo, G.C. Furtado, S.A. Lira, Increased

        expression of CCL2 in insulin-producing cells of transgenic mice promotes mobilization

        of myeloid cells from the bone marrow, marked insulitis, and diabetes, Diabetes, 57

        (2008) 3025-3033. https://doi.org/10.2337/db08-0625.

[25]   S. Climaco-Arvizu, O. Domínguez-Acosta, M.A. Cabañas-Cortés, M. Rodríguez-

        Sosa, F.J. Gonzalez, L. Vega, G. Elizondo, Aryl hydrocarbon receptor influences nitric

        oxide and arginine production and alters M1/M2 macrophage polarization, Life Sci. 155,

        (2016) 76–84. https://doi.org/10.1016/j.lfs.2016.05.001.

[26]   I.I. Ivanov, B.S. McKenzie, L. Zhou, C.E. Tadokoro, A. Lepelley, J.J. Lafaille, D.J.

        Cua, D.R. Littman, The orphan nuclear receptor RORgammat directs the differentiation

        program of proinflammatory IL-17+ T helper cells, Cell, 126 (2006) 1121–1133.

        https://doi.org/10.1016/j.cell.2006.07.035.

[27]   R. Parsa, P. Andresen, A. Gillett, S. Mia, X.M. Zhang, S. Mayans, D.Holmberg, R.A.

        Harris, Adoptive transfer of immunomodulatory M2 macrophages prevents type 1

        diabetes in NOD mice, Diabetes, 61 (2012) 2881–2892. https://doi.org/10.2337/db11-

        1635.

[28]   L.K. Shoda, D.L. Young, S. Ramanujan, C.C. Whiting, M.A. Atkinson, J.A. Bluestone,

        G.S. Eisenbarth, D. Mathis, A.A. Rossini, S.E. Campbell, R. Kahn, H.T. Kreuwel, A

comprehensive review of inventions in the NOD mouse and implications for translation, Immunity, 23 (2005) 115–126. https://doi.org/10.1016/j.immuni.2005.08.002.

[29]   E. Dirice, R.W.S. Ng, R. Martinez, J. Hu, F.F. Wagner, E.B. Holson, B.K. Wagner, R.N. Kulkarni, Isoform-selective inhibitor of histone deacetylase 3 (HDAC3) limits pancreatic islet infiltration and protects female nonobese diabetic mice from diabetes, J. Biol. Chem. 292 (2017)17598–17608. https://doi.org/10.1074/jbc.M117.804328.

[30]   M. Lundh, D.P. Christensen, M. Damgaard Nielsen, S.J. Richardson, M.S. Dahllo☐f, T. Skovgaard, J. Berthelsen, C.A. Dinarello, A. Stevenazzi, P. Mascagni, L.G. Grunnet, N.G. Morgan, T. Mandrup-Poulsen, Histone deacetylases 1 and 3 but not 2 mediate cytokine-induced beta cell apoptosis in INS-1 cells and dispersed primary islets from rats and are differentially regulated in the islets of type 1 diabetic children, Diabetologia, 55 (2012) 2421–2431. https://doi.org/10.1007/s00125-012-2615-0.

[31]   M. Lundh, T. Galbo, S.S. Poulsen, T. Mandrup-Poulsen, Histone deacetylase 3 inhibition improves glycaemia and insulin secretion in obese diabetic rats, Diabetes Obes. Metab. 17 (2015) 703-707. https://doi.org/10.1111/dom.12470.

[32]   F.F. Wagner, M. Lundh, T. Kaya, P. McCarren, Y.L. Zhang, S. Chattopadhyay, J.P. Gale, T. Galbo, S.L. Fisher, B.C. Meier, A. Vetere, S. Richardson, N.G. Morgan, D.P. Christensen, T.J. Gilbert, J.M. Hooker, M. Leroy, D. Walpita, T. Mandrup-Poulsen, B.K. Wagner, E.B. Holson, An isochemogenic set of inhibitors to define the therapeutic potential of histone deacetylases in β-cell protection, ACS Chem. Biol. 11 (2016) 363-374. https://doi.org/10.1021/acschembio.5b00640.

[33]   J.D. Katz, C. Benoist, D. Mathis, T helper cell subsets in insulin-dependent diabetes, Science, 268 (1995) 1185-1188. https://doi.org/10.1126/science.7761837.

[34] B. Hultgren, X. Huang, N. Dybdal, T.A. Stewart, Genetic absence of gamma-interferon delays but does not prevent diabetes in NOD mice, Diabetes, 45 (1996) 812-817. https://doi.org/10.2337/diab.45.6.812.

[35] S. Trembleau, G. Penna, S. Gregori, N. Giarratana, L. Adorini, IL-12 administration accelerates autoimmune diabetes in both wild-type and IFN-gamma-deficient nonobese diabetic mice, revealing pathogenic and protective effects of IL-12-induced IFN-gamma, J. Immunol. 170 (2003) 5491-5501. https://doi.org/10.4049/jimmunol.170.11.5491.

[36] J. Zhang, Z. Huang, R. Sun, Z. Tian, H. Wei, IFN-γ induced by IL-12 administration prevents diabetes by inhibiting pathogenic IL-17 production in NOD mice, J. Autoimmun. 38 (2012) 20-28. https://doi.org/10.1016/j.jaut.2011.11.017.

[37] S.M. Bellemore, E. Nikoopour, O. Krougly, E. Lee-Chan, L.A. Fouser, B. Singh, Pathogenic T helper type 17 cells contribute to type 1 diabetes independently of interleukin-22, Clin. Exp. Immunol. 183 (2016) 380-388. https://doi.org/10.1111/cei.12735.

[38] N. Martin-Orozco, Y. Chung, S.H. Chang, Y.H. Wang, C. Dong, Th17 cells promote pancreatic inflammation but only induce diabetes efficiently in lymphopenic hosts after conversion into Th1 cells, Eur. J. Immunol. 39 (2009) 216-224. https://doi.org/10.1002/eji.200838475.

[39] D. Bending, H. De la Peña, M. Veldhoen, J.M. Phillips, C. Uyttenhove, B. Stockinger, A. Cooke A, Highly purified Th17 cells from BDC2.5NOD mice convert into Th1-like cells in NOD/SCID recipient mice, J. Clin. Invest. 119 (2009) 565-572. https://doi.org/10.1172/JCI37865.

[40]   G. Kuriya, T. Uchida, S. Akazawa, M. Kobayashi, K. Nakamura, T. Satoh, I. Horie, E. Kawasaki, H. Yamasaki, L. Yu, Y. Iwakura, H. Sasaki, Y. Nagayama, A. Kawakami, N. Abiru, Double deficiency in IL-17 and IFN-γ signalling significantly suppresses the development of diabetes in the NOD mouse, Diabetologia, 56 (2013) 1773-1780. https://doi.org/10.1007/s00125-013-2935-8.

[41]   S.V. Pakala, M.O. Kurrer, J.D. Katz, T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186 (1997) 299-306. https://doi.org/10.1084/jem.186.2.299.

[42]   A.M. Marleau, N.E. Sarvetnick, IL-18 is required for self-reactive T cell expansion in NOD mice, J. Autoimmun. 36 (2011) 263–277. https://doi.org/10.1016/j.jaut.2011.02.005.

[43]   A.E. Ciecko, B. Foda, J.Y. Barr, S. Ramanathan, M.A. Atkinson, D.V. Serreze, A.M. Geurts, S.M. Lieberman, Y.G. Chen, Interleukin-27 is essential for type 1 diabetes development and Sjögren syndrome-like Inflammation, Cell Rep. 29 (2019) 3073–3086.e5. https://doi.org/10.1016/j.celrep.2019.11.010.

[44]   J.S. Chia, J.L. McRae, P.J. Cowan, K.M. Dwyer, The CD39-adenosinergic axis in the pathogenesis of immune and nonimmune diabetes, J. Biomed. Biotechnol. 2012 (2012) 320495. https://doi.org/10.1155/2012/320495.9.

[45]   S. Hori, T. Nomura, S. Sakaguchi, Control of regulatory T cell development by the transcription factor Foxp3, Science, 299 (2003)1057-1061. https://doi.org/10.1126/science.1079490.

[46]   H. Korf, L. Breser, J. Van Hoeck, J. Godoy, D.P. Cook, B. Stijlemans, E. De Smidt, C. Moyson, J.P. Monteiro Carvalho Mori Cunha, V. Rivero, C. Gysemans, C. Mathieu, MIF

inhibition interferes with the inflammatory and T cell-stimulatory capacity of NOD

macrophages and delays autoimmune diabetes onset, PLoS One, 12 (2017) e0187455.

https://doi.org/10.1371/journal.pone.0187455.

[47]   K. Das Gupta, M.R. Shakespear, A. Iyer, D.P. Fairlie, M.J. Sweet, Histone deacetylases

in monocyte/macrophage development, activation and metabolism: refining HDAC

targets for inflammatory and infectious diseases, Clin. Transl. Immunology, 5 (2016) e62.

http://doi.org/10.1038/cti.2015.46.

[48]   T. Roger, J. Lugrin, D. Le Roy, G. Goy, M. Mombelli, T. Koessler, X.C. Ding, A.L.

Chanson, M.K. Reymond, I. Miconnet, J. Schrenzel, P. François, T. Calandra, Histone

deacetylase inhibitors impair innate immune responses to 43 Toll-like receptor agonists

and to infection, Blood, 117 (2011) 1205–1217. https://doi.org/10.1182/blood-2010-05-

284711.

*V. Patel, A. Jayaraman, S. Jayaraman, Epigenetic reprogramming ameliorates type 1 diabetes

by decreasing the generation of Th1 and Th17 subsets and restoring self-tolerance in CD4[+] T

cells, under consideration of publication. 2021.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6