Temperature and Guanidine Hydrochloride effects on the folding thermodynamics of WW domain and variants

Meng Qin,[†,‡] Zhenxing Liu,[¶] Wei Wang,[‡] and D. Thirumalai[§]

[†]*Biophysics Program, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742*

[‡]*National Laboratory of Solid State Microstructure and Department of Physics, Nanjing University, and Collaborative Innovation Center of Advanced Microstructures, Nanjing 210093, China*

[¶]*Department of Physics, Beijing Normal University, Beijing 100875, China*

[§]*Department of Chemistry, The University of Texas at Austin, Austin, TX 78712*

E-mail:

Abstract

We use simulations based on an all atom Go model to calculate the folding temperatures (T_fs) and free energies (ΔGs) of two variants of the WW domain, which is a small all β-sheet protein. The results, without adjusting any parameter, are in good agreement with experiments, thus validating the simulations. We then used the Molecular Transfer Model to predict the changes in their ΔG and T_fs as Guanidine Hydrochloride concentration is varied. The predictions can be readily tested in experiments.

Introduction

In this paper, we study thermal denaturation and effects of guanidine hydrochloride (GdmCl) on the thermodynamics of folding of the well-characterized small all β WW domain[1–3] and two variants using all atom Go (AAGO) model.[4–6] Denaturation effects are accounted for using the successful Molecular Transfer Model (MTM).[7–10] The rationale for using the Go model using lattice[11–14] or off-lattice[15,16] Go and AAGO models is that non-native interactions only play a sub-dominant role in determining the folding mechanisms of globular proteins.[17,18] The computations based on the Go model provided support to theory,[19–22] which ushered in the notion that the gradient in the funneled energy landscape drives folding predominantly to the native state under folding conditions.

The studies cited above and subsequent developments (see perspectives by several authors[23–29]) have given tool sets, which have firmly established that small single domain proteins fold by parallel routes. Most of the computations, which have led to far reaching conclusions, have been done by triggering folding by changing the temperature. Experiments, on the other hand, typically investigate folding by altering the concentration of denaturants, such as urea or GdmCl at a fixed temperature. Therefore, practical ways of including denaturant effects are needed so that quantitative comparison between simulations and experiments can be made. We introduced the MTM to accomplish this goal.[7,30] Currently, MTM is the only method that can be used to obtain fairly accurate results for denaturant-induced folding

(see for example[8–10]), regardless of the size of topology of the folded state.

Our goal here is to calculate the thermodynamic properties of the all β sheet protein using AAGO and then predict how they change as function of GdmCl. The folded state of the wild-type Pin WW domain (PDB entry:1PIN) has a twisted triple-stranded antiparallel β-sheets spanning Lys6 to Gly39 (Figure 1). There are two loops, loop1 with 6 residues ($S_{16}R_{17}S_{18}S_{19}G_{20}R_{21}$), and a shorter loop 2 with 4 residues. Another important member of the WW domain family, the Formin Binding Protein 28 (FBP28) WW domain (PDB entry: 1E0L) with residues from Ser6 to Pro33, with a structure that is similar to the 1PIN WW domain. Except for the difference in the 5 residue loop1 ($T_{13}A_{14}D_{15}G_{16}K_{17}$), FBP28 is homologous to the 1PIN sequence. Not surprisingly, they have a similar folding behavior. If the 6-residual loop1 in 1PIN WW domain is mutated with the 5-residual FBP28 loop1 creates a variant (PDB entry:2F21), which is has an improved thermodynamic stability with the melting temperature that increases from from 332K (1PIN) to 350.5K.[1] Here, we focus on 1PIN and 2F21 domains. The Table of Contents graphics shows a folding trajectory for the FBP28 domain.

In all of our previous applications of the MTM we used a coarse-grained Self-Organized Polymer (SOP) model[31] in which each residue in the protein was represented using two beads, one centered at the α-carbon and the other at the center of mass of the side chain. Here, we developed the AAGO model, following previous studies,[4–6] to model the WW domains. We used the MTM in concert with AAGO to investigate the effect of GdmCl on the folding of WW domain and two variants, after ensuring that the simulations in the absence of denaturants reproduce the experimental results quantitatively.[1,2] We then predicted the thermodynamic properties for the two variants as a function of GdmCl. The current work shows that the tools used here are sufficient to anticipate denaturant effects on the thermodynamics of globular proteins.

Model and Simulation Methods

All-atom Go model (AAGO) In the AAGO model all heavy (non-hydrogen) atoms are included, and bond lengths, bond angles, dihedral bonds are maintained as harmonic potentials. As is customary in Go models, there is an attractive potential between non-bonded atom pairs (heavy atoms i and j, where $i > j + 2$) that are in contact in the native state. All other non-local interactions are repulsive, which ensures that the atoms have a defined excluded volume.

Native non-bonded interactions separated by exactly three bonds (1-4 interaction) are reduced by a scale factor 0.5, in calculating the dihedral potentials. We used the standard functional form of the potential, given by,

$$V = \sum_{bonds} K_r(r - r_{eq})^2 + \sum_{angles} K_\theta(\theta - \theta_{eq})^2 + \sum_{dihedrals} \frac{V_n}{2}(1 + \cos(n\phi - \gamma)) +$$
$$\sum_{i<j-2} [(A_{ij}/R_{ij})^{12} - (B_{ij}/R_{ij})^{10}] + \sum_{i<j}(\sigma_{nonij}/R_{ij})^{12} \tag{1}$$

In the above equation, the atom types, parameters for bond angle and dihedral interactions are similar to those employed in the amber ff94 force field.[32] The equilibrium values of r_{eq}, θ_{eq} values are taken from X-ray structures, and K_θ value was taken from vibrational analysis of small molecules containing the specific atoms and standard normal mode analysis, as previously described.[32] The force constants associated with the (ϕ, ψ) angles are chosen, following the Amber force field ff99SB,[33] to achieve a better balance of secondary structure elements. In the above equation, V_n is the amplitude of the dihedral angle potential, n is dihedral periodicity, and γ is the phase of the dihedral angle, ϕ. The Fourier series is approximated using a small number of terms. The form of the improper torsions is the same as the dihedral angles.

In order to reduce the simulation time, the side chain of each amino acid is assumed to be rigid. This simplification means that the relative positions of these heavy atoms in one side

chain are rigidly fixed during the simulations. The movement of these atoms in rigid body can be decomposed into the translational of the center of mass and the rotational movement of the rigid body. This approximation, which does not compromise accuracy, eliminates the need to calculate the bond stretch, bond angle, and dihedral angle potentials.

Simulations

Extensive thermodynamic sampling of the protein WW domain is achieved using low friction Langevin dynamics[34] simulations. The equations of motion for the position of a heavy atom, r_i, are integrated using, $m\ddot{\vec{r}} = -\zeta\dot{\vec{r}} + F_c + \Gamma$. Here, m is the mass of the atom in each amino acid, $F_c = -\partial V/\partial r_i$, and $\Gamma(t)$ is a random force with a white noise spectrum satisfying the autocorrelation function in the discretized form $< \Gamma(t)\Gamma(t+h) >= (2\zeta k_B T/h)\delta(0,k)$, where $\delta_{0,k}$ is the Kronecker delta function, h is the integration step size, and $n = 0, 1, 2, \ldots$.

We used the Verlet leapfrog algorithm to integrate the equations of motion. The velocity, v_i, at time $t + \frac{h}{2}$ and the position, r_i, at time $t + h$ of an atom are given by,

$$V_i(t + \frac{h}{2}) = \frac{2m - h\zeta}{2m + h\zeta}v(t - \frac{h}{2}) + \frac{2h}{2m + h\zeta}[F(t) + \Gamma(t)], \qquad (2)$$

and $r_i(t + h) = r_i(t) + v_i(t + \frac{h}{2})$. The value of the time step used to integrate above two equations is $h = 0.02\tau_L$, τ_L is the unit of time, and a low friction (a tenth of the viscosity in water) is used to obtain enhanced sampling and converged equilibrium thermodynamic data.

Molecular Transfer Model (MTM)

In order to include denaturant effects, we used the MTM, which first involves computing the transfer free energy. which has been remarkably successful in studying the folding of a variety of proteins.[8,35] The free energy of transferring a protein conformation, described by the set of coordinates r_is, from water $[C] = 0$) to aqueous denaturant solution ($[C] \neq 0$) is

taken to be as $\delta G(r_i, [C]) = \sum_k \delta g(k, [C]) \alpha_k(r_i) / \alpha_{Gly-k-Gly}$, where the sum is over backbone and side chain, $\delta g(k, [C]) = m_k[C] + b_k$ (m_k and b_k, and $\alpha_{Gly-k-Gly}$[8] are the transfer free energy of group k, $\alpha_k(r_i)$ is the SASA of k, and $\alpha_{Gly-k-Gly}$ is the SASA of the k^{th} group in the tripeptide Gly-k-Gly, respectively. Thus, the effective free energy function for a protein at $[C] \neq 0$ is $G_P(r_i, [C]) = V(r_i) + \delta G(r_i, [C])$. We computed $\alpha_{Gly-k-Gly}$ by summing the contribution of all the SASA of the heavy atoms in the side chain and backbone.[36] Note that MTM does not assume whether the protein folds by a two-state or multi-state mechanism. In several previous studies,[8,10,35,37] we have shown that MTM and a coarse-grained model of a protein with beads accurately describes the thermodynamics and kinetics of folding of a number of proteins.

Weighted Histogram Method

The heat capacity, C_v and the free-energy are calculated using the weighted histogram analysis method, which gives optimal estimate of the density of states of the system, allowing us to calculate the relevant thermodynamic quantities accurately. We calculated C_v and the free energy are calculated based on 25 independent long-time runs at different temperatures from 300K to 450K.

Results

Folding trajectories: Figure1 shows the folding trajectories of 1PIN and 2F21 domains. We plot the radius of gyration from an unfolded state for the two variants as a function of time, t. There is a precipitous decrease in R_g in $\approx 5\mu s$ as the protein folds from a fully denatured state. In the process of reaching the folded state, this WW domain samples a few intermediate states. The folding trajectory for the more stable 2F21 is different. In this case, only prior to reaching the folded state, the protein is compact. It is clear that in 2F21 folding, compaction occurs on similar time scales. In addition, the folding rate of 2F21

is about an order of magnitude greater than for the 1PIN. The ensemble of conformations sampled by 2F21 during the folding process is conformationally more heterogeneous than found for the less stable 1PIN.

Heat capacity and the folding temperature (T_f): From the plots of C_v as a function of T, we find there is a single peak for both 1PIN and 2F21 variants. The width of the transition is somewhat narrower for 2F21 than for 1PIN. The peaks in the C_v plots, which we associate with the folding temperature, are in excellent agreement with experiments. This finding is particularly gratifying given that we did not *adjust* a single parameter to obtain agreement with experiments.

Temperature-dependent Energies: Heat capacity C_v as a function of T shows that 1PIN and 2F21 domains fold in a two state manner. Therefore, we could compute the difference between the folded and unfolded states using, $\Delta G = -k_B T \frac{P_f}{(1-P_f)}$ where P_f is the probability of being in the folded state. Comparison of the calculated and experimental values for ΔG as a function of T for both the variants shows very good agreement (Figure 3).

GdmCl effects on folding thermodynamics: The results in Figures 2 and 3 show that the all atom Go model reproduces the folding thermodynamics as a function of temperature, thus validating our model. We used the ensemble of conformations generated at various temperatures to calculate the GdmCl-dependent folding temperature and free energies using the MTM model. Plots of C_v at various values of the GdmCl concentration are displayed in Figure 4. Two observations can be made: (1) The dependence of the folding temperature on GdmCl changes linearly as the concentration of the denaturant increases. (2) At all values of the denaturant concentrations T_f for 2F21 is greater than for 1PIN, which is refection of the differences in the stabilities. A similar conclusion may be reached from the plot of ΔG as a function of GdmCl concentration at the indicated temperatures (Figure 5).

Conclusion

We have reported the folding thermodynamics of two variants of a small all β-sheet protein. The good agreement, without adjusting any parameter to fit measurements, between simulations and experiments for thermal folding shows that AAGO is a reasonable starting point for investigating the folding of globular proteins. The results for GdmCl effects, shown in Figures 4 and 5, are genuine predictions, which could be readily tested.

We end with a couple of remarks. (1) The all atom Go model by definition ignores non-native interactions. This is a limitation of the model even though we have reproduced the experimental results accurately (Figures 2 and 3). It is necessary to include non-native interactions for some test cases to assess the error in calculate folding thermodynamics. (2) Hydration effects are completely neglected and could also be important in describing the equilibrium aspects of folding. One way of accounting for hydration effects is to simulate folding of all atom Go model in explicit solvent. If this protocol is successful then the role of water can be directly addressed. (3) Given the success of the MTM, we believe it will be interesting to use the conformations generated using atomic detailed simulations for WW domain[3] and use it in conjunction with MTM to obtain the effects of denaturants (GdmCl or urea) on the folding thermodynamics.

Acknowledgement

We thank Natalia Denesyuk for help in the initial stages in developing the All Atom Go Model. Most of the work was done while MQ and DT were at the University of Maryland. This work was supported in part by a grant from National Science Foundation (CHE 19-00093) and the Collie-Welch Chair (F-0019) administered through the Welch Foundation. ZL acknowledges financial support from the National Natural Science Foundation of China (11104015,11675017,11735005).

References

(1) Jager, M.; Zhang, Y.; Bieschke, J.; H, N.; M, D.; Bowman, M.; Noel, J.; Gruebele, M.; Kelley, J. Structure-function-folding relationship in a WW domain. *Proc. Natl. acad. Sci.* **2006**, *103*, 10648–10653.

(2) Liu, F.; Du, D.; Fuller, A.; Davoren, J.; Wipf, P.; Gruebele, M. An experimental survey of the transition between two-state and downhill protein folding scenarios. *Proc. Natl. Acad. Sci.* **2008**, *105*, 2369–2374.

(3) Freddolino, P.; F, L.; M, G.; Schulten, K. Ten-Microsecond Molecular Dynamics Simulation of a Fast-Folding WW Domain. *Biophys. J.* **2008**, L75–L77.

(4) Shimada, J.; Kussell, E.; Shakhnovich, E. The folding thermodynamics and kinetics of crambin using an all-atom Monte Carlo simulation. *J. Mol. Biol.* **2001**, 79–95.

(5) Shimada, J.; Shakhnovich, E. The ensemble folding kinetics of protein G from an all-atom Monte Carlo simulation. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 11175.

(6) Clementi, C.; Garcia, A.; Onuchic, J. Interplay among tertiary contacts, secondary structure formation and side-chain packing in the protein folding mechanism: all-atom representation study of protein L. *J. Mol. Biol.* **2003**, *326*, 933–954.

(7) O'Brien, E. P.; Ziv, G.; Haran, G.; Brooks, B. R.; Thirumalai, D. Effects of denaturants and osmolytes on proteins are accurately predicted by the molecular transfer model. *Proceedings of the National Academy of Sciences of the United States of America* **2008**, *105*, 13403–13408.

(8) Liu, Z. X.; Reddy, G.; O'Brien, E. P.; Thirumalai, D. Collapse kinetics and chevron plots from simulations of denaturant-dependent folding of globular proteins. *Proceedings of the National Academy of Sciences of the United States of America* **2011**, *108*, 7787–7792.

(9) Liu, Z. X.; Reddy, G.; Thirumalai, D. Theory of the Molecular Transfer Model for Proteins with Applications to the Folding of the src-SH3 Domain. *Journal of Physical Chemistry B* **2012**, *116*, 6707–6716.

(10) Liu, Z.; Reddy, G.; Thirumalai, D. Folding PDZ2 domain using the molecular transfer model. *J. Phys. Chem. B.* **2016**, *120*, 8090–8101.

(11) Taketomi, H.; Udea, Y.; Go, N. Studies on protein folding, unfolding and fluctuations by computer simulation. I. The effect of specific amino acid sequence represented by specific inter-unit interactions. *Int. J. Pept. Res.* **1975**, *7*, 445–459.

(12) Go, Theoretical studies of protein folding. *Ann. Rev Biophys. Bioeng.* **1983**, *12*, 183–210.

(13) Camacho, C. J.; Thirumalai, D. Kinetics and Thermodynamics of Folding in Model Proteins. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 6369–6372.

(14) Dill, K. A.; Bromberg, S.; Yue, K. Z.; Fiebig, K. M.; Yee, D. P.; Thomas, P. D.; Chan, H. S. Principles of Protein-Folding - a Perspective from Simple Exact Models. *Prot. Sci.* **1995**, *4*, 561–602.

(15) Nymeyer, H.; Garcia, A. E.; Onuchic, J. N. Folding funnels and frustration in off-lattice minimalist protein landscapes. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 5921–5928.

(16) Klimov, D. K.; Thirumalai, D. Mechanisms and Kinetics of β-Hairpin Formation. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 2544–2549.

(17) Klimov, D. K.; Thirumalai, D. Multiple protein folding nuclei and the transition state ensemble in two-state proteins. *Proteins - Struct. Funct. Gene.* **2001**, *43*, 465–475.

(18) Best, R.; Hummer, G.; Eaton, W. Native contacts determine protein folding mechanisms in atomistic simulations. *Proc. Natl. Acad. Sci.* **2013**, *110*, 17874–17879.

(19) Bryngelson, J.; Onuchic, J.; Socci, N.; Wolynes, P. Funnels, pathways, and the energy landscape of protein-folding - A synthesis. *Proteins: Structure, Function and Genetics* **1995**, *21*, 167–195.

(20) Dill, K. A.; Chan, H. S. From Levinthal to Pathways to Funnels. *Nature Struct. Biol.* **1997**, *4*, 10–19.

(21) Onuchic, J. N.; Wolynes, P. G. Theory of protein folding. *Curr. Opin. Struct. Biol.* **2004**, *14*, 70–75.

(22) Nassar, R.; Dignon, G.; Razban, R.; Dill, K. The protein folding problem: The role of theory. *J. Mol. Biol.* **2021**, *x*, xxx–xxx.

(23) Dill, K. A.; Ozkan, S. B.; Shell, M. S.; Weikl, T. R. The protein folding problem. *Annu. Rev. Biophys.* **2008**, *37*, 289–316.

(24) Shakhnovich, E. Protein Folding Thermodynamics and Dynamics: Where Physics, Chemistry, and Biology Meet. *Chem. Rev.* **2006**, *106*, 1559–1588.

(25) Schuler, B.; Eaton, W. A. Protein folding studied by single-molecule FRET. *Curr. Opin. Struct. Biol.* **2008**, *18*, 16–26.

(26) Eaton, W. Modern Kinetics and Mechanism of Protein Folding: A Retrospective. *J. Phys. Chem. B.* **2021**, *125*, 3452–3467.

(27) Thirumalai, D.; O'Brien, E. P.; Morrison, G.; Hyeon, C. Theoretical Perspectives on Protein Folding. *Annu. Rev. Biophys.* **2010**, *39*, 159–183.

(28) Campos, L.; Sadqi, M.; Munoz, V. Lessons about Protein Folding and Binding from Archetypal Folds. *Acc. Chem. Res.* **2020**, *53*, 2180–2188.

(29) Thirumalai, D.; Klimov, D. K.; Dima, R. I. Insights into specific problems in protein folding using simple concepts. *Adv. Chem. Phys.* **2002**, *120*, 35–76.

(30) Thirumalai, D.; Liu, Z.; O'Brien, E.; Reddy, G. Protein folding: from theory to practice. 2013.

(31) Hyeon, C.; Dima, R. I.; Thirumalai, D. Pathways and kinetic barriers in mechanical unfolding and refolding of RNA and proteins. *Structure* **2006**, *14*, 1633–1645.

(32) Cornell, W.; Cieplak, P.; Bayly, C.; Gould, I.; Merz, K.; Ferguson, D.; David C. Spellmeyer, D.; Fox, J., T andCaldwell; Kollman, P. A. A Second Generation Force Field for the Simulation of Proteins, Nucleic Acids, and Organic Molecules. *J. Am. Chem. Soc.* **1995**, 5179–5197.

(33) Hornak, V.; Abel, R.; Okur, A.; Strockbine, B.; Roitberg, A.; Simmerling, C. Comparison of multiple Amber force fields and development of improved protein backbone parameters. *Proteins: Structure, Function, & genetics* **2006**, *65*, 712–725.

(34) Honeycutt, J. D.; Thirumalai, D. The nature of folded states of globular proteins. *Biopolymers* **1992**, *32*, 695–709.

(35) Reddy, G.; Thirumalai, D. Dissecting Ubiquitin Folding Using the Self-Organized Polymer Model. *Journal of Physical Chemistry B* **2015**, *119*, 11358–11370.

(36) Hayryan, S.; Hu, C.-K.; Skřivánek, J.; Hayryane, E.; Pokorný, I. A new analytical method for computing solvent-accessible surface area of macromolecules and its gradients. 2005.

(37) Reddy, G.; Liu, Z.; Thirumalai, D. Denaturant-dependent folding of GFP. *Proceedings of the National Academy of Sciences of the United States of America* **2012**, *109*, 17832–17838.

Figure 1: Folding trajectories at 310K for the 1PIN (2F21) domain on the left (right). The dynamics of folding is monitored using R_g as a function of time. Snapshots (a-e) and (a-f) are sampled conformations by 1PIN and 2F21, respectively. in the trajectory, respectively. In both the trajectories a conformation with a β-hairpin ((c) in 1PIN and (e) in F21) form upon collapse of the proteins.

Figure 2: Heat capacities (C_v) for 1PIN (black) and 2F21 (red) as a function of temperature. The folding temperatures correspond to the peaks in C_v. The experimental values are also shown.

Figure 3: Free energy difference between the folded and unfolded states as a function of temperature. Top and bottom panels are for 1PIN and 2F21, respectively. Black (red) lines are from simulations (experiments).

Figure 4: The dependence of C_v on temperature at various values of the GdmCl concentration. Top and bottom panels are for 1PIN and 2F21, respectively. The two-state transition at all values of denaturant concentration is reflected in a single peak in the heat capacity.

Figure 5: Predicted free energy difference between the folded and unfolded states as a function of GdmCl for 1PIN(black) and 2F21(red), respectively. The simulation temperatures are shown.