Mechanisms of genetic instability in a single S-phase following whole genome doubling

Simon Gemble[1,7], Sara Vanessa Bernhard[2], Nishit Srivastava[3], René Wardenaar[4] Maddalena Nano[1], Anne-Sophie Macé[5], Andréa E. Tijhuis[4], Kristina Keuper[2], Diana C.J. Spierings[4], Helfrid Hochegger[6], Matthieu Piel[3], Floris Foijer[4], Zuzana Storchová[2] and Renata Basto[1,7].

1- Institut Curie, PSL Research University, CNRS, UMR144, Biology of Centrosomes and Genetic Instability Laboratory, Paris, France.

2- Department of Molecular Genetics, TU Kaiserslautern, Kaiserslautern, Germany.

3- Institut Curie and Institut Pierre Gilles de Gennes, PSL Research University, CNRS, UMR 144, Systems Biology of Cell Polarity and Cell Division, Paris, France.

4- European Research Institute for the Biology of Ageing, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

5- Cell and Tissue Imaging Facility (PICT-IBiSA), Institut Curie, PSL Research University, Centre National de la Recherche Scientifique, Paris, France.

6- Genome Damage and Stability Centre, School of Life Sciences, University of Sussex, Brighton, UK.

7- Corresponding authors- simon.gemble@curie.fr and renata.basto@curie.fr

ABSTRACT

Doubling of the full chromosome content -whole genome duplications (WGDs)- is frequently found in human cancers and is responsible for the rapid evolution of genetically unstable karyotypes [1–3]. It has previously been established that WGDs fuel chromosome instability due to abnormal mitosis owing to the presence of extra centrosomes and extra chromosomes [4–8]. Tolerance to ploidy changes has been identified in different model organisms and cell types [5,6,9–12], revealing long term cellular adaptations that accommodate ploidy increase. Importantly, however, the immediate consequences of WGDs as cells become tetraploid are not known. It also remains unknown whether WGD triggers other events leading to genetic instability (GIN), independently of mitosis. In this study, we induced tetraploidy in diploid genetically stable RPE-1 cells and monitored the first interphase. We found that newly born tetraploids undergo high rates of DNA damage during DNA replication. Using DNA combing and single cell sequencing, we show that replication forks are unstable, perturbing DNA replication dynamics and generating under- and over-replicated regions at the end of S-phase. Mechanistically, we found that these defects result from lack of protein mass scaling up at the G1/S transition, which impairs the fidelity of DNA replication. This work shows that within a single interphase, unscheduled tetraploid cells can accumulate highly abnormal karyotypes. These findings provide an explanation for the GIN landscape that favors tumorigenesis after tetraploidization.

MAIN

Diploid and stable karyotypes are associated with health and fitness in animals. In contrast, whole genome duplications (WGDs) are linked to genetic instability (GIN) and cancer [1,2,13]. WGDs promote chromosomal instability (CIN) over time due to abnormal mitosis, contributing to the evolution of aneuploid karyotypes [3,6,8,14–16]. These represent cancer vulnerabilities with therapeutic potential [17,18]. Tolerance to ploidy changes has been identified in different model organisms and cell types [5,6,9–12], revealing long-term cellular adaptations that accommodate ploidy increase. Importantly, however, the immediate consequences of unscheduled WGDs are not known and their contribution to GIN remains to be identified. This is an essential question because single WGD events such as cytokinesis failure can promote tumorigenesis [19]. Identifying the initial defects derived from WGD and the underlying mechanisms establishing GIN in tetraploid cells is thus an important step, which requires understanding failure in maintaining genetic stability. This type of study has the potential to unravel the origins of GIN in one single cell cycle after WGDs.

High levels of DNA damage are generated in the first interphase following unscheduled Whole Genome Duplication (WGD)

To identify the immediate consequences of WGDs, we induced tetraploidization in the human diploid and genetically stable RPE-1 immortalized cell line, which contains an almost near-diploid chromosome content. Since WGDs can have different origins [15,20], we developed several approaches to induce tetraploidization through either cytokinesis failure (CF), endoreplication (EnR) or mitotic slippage (MS) (Fig. 1A and methods). While the large majority of cells resulting from CFs contained two nuclei, EnR or MS generated mainly mononucleated tetraploid cells (Fig. 1A-B and Extended data Fig. 1A-I). Importantly, taking together parameters such as cell size, nuclei number and size, and centrosome number, we were able to distinguish diploids from tetraploids in all the strategies used (Fig. 1B and Extended data Fig. 1A-I). For each strategy, a mix of diploid and tetraploid cells was obtained, allowing a comparison of an internal diploid control and the tetraploid population (Fig. 1C-H, and Extended data Fig. 1A-I). Importantly, from all the conditions used to induce WGDs, the large majority of tetraploid cells continued to cycle and enter and exit the first S-phase (see below). Thus, we have generated the conditions required to study the initial and immediate consequences of tetraploidy within the first cell cycle.

Using an early marker of DNA double strand breaks - γH2AX-, we characterized levels of DNA damage in the first interphase following tetraploidization and found high levels in tetraploid cells. Moreover, this was independent of the way tetraploid cells were generated. In contrast, diploid cells treated in the same conditions or untreated diploids showed low levels of DNA damage (Fig. 1C-H, and Extended data Fig. 1A-I, see Extended data Fig. 2A-C for additional methods of generating tetraploidy). While most of the diploid cells exhibited a low number of γH2AX foci, the percentage of tetraploid cells with more than 10 foci was high (Fig. 1C-H). We found a correlation between the number of γH2AX foci and the fluorescence intensity levels of this marker (Extended data Fig. 1J). For simplicity, we will include the information of the number of γH2AX foci per interphase nuclei throughout this study. Since MS generated the highest frequency of tetraploid cells (Extended data Fig. 1A), we chose to present data derived from MS throughout this study. To confirm some of our results, we also used CF or EnR and this will be mentioned in the figure legends and in the methods.

Since a tetraploid nucleus contains twice the amount of DNA than a diploid

nucleus, we excluded by normalization with nuclear area or nuclear fluorescence

intensity, an increase of γH2AX foci due to increased nuclear size (Extended data Fig.

1K-L). Additionally, we confirmed that the high levels of DNA damage found in the first

interphase after tetraploidization were not specific to RPE-1 cells, as the diploid BJ

fibroblast cell line as well as the pseudo diploid human colon carcinoma HCT116 cell

line also displayed high levels of DNA damage upon WGD (Extended data Fig. 2D-E).

We next compared the levels of DNA damage detected in tetraploid cells with

the levels of DNA damage in diploid cells generated by replication stress (RS). RS is

the slowing or stalling of replication forks, which can be induced by high doses of

Aphidicolin (APH) or Hydroxyurea (HU), among other challenges [21,22]. Interestingly,

APH or HU generated comparable levels of DNA damage in diploid cells, when

compared to untreated tetraploid cells (Extended data Fig. 1M).

Collectively, our results show that a transition from a diploid to tetraploid status

after unscheduled WGD, is accompanied by high levels of DNA damage within the first

cell cycle.

DNA damage and genetic instability in tetraploid cells is generated during S-

phase in a DNA replication-dependent manner

We then determined the cell cycle stage in which the DNA damage occurs. We

followed cell cycle progression right after WGD using the fluorescence ubiquitination

cell cycle indicator (FUCCI) system to map the timing of cell cycle progression in

tetraploid cells, allowing us to then monitored the number of γH2AX foci during the first

G1 and the first S-phase (Fig. 2A-B and Extended data Fig. 3A). During G1, the number

of γH2AX foci was quite low and comparable to controls. As cells enter S-phase

(t=10hrs), a slight increase in the number of foci in tetraploid nuclei could be observed,

which increased substantially at the end of S-phase (t=16hrs) (Fig. 2A-B and Extended

data Fig. 3A). These results were further confirmed by time lapse imaging using

tetraploid RPE-1 cells tagged with H2B-GFP to visualize DNA and 53BP1-RFP, which

is a double strand break repair factor [23] (Extended data Fig. 3B-C and Extended data

movies 1-2). To confirm that DNA damage in tetraploid cells was induced during S-

phase, we blocked cells at the G1/S transition using high doses of either CDK4/6 or

CDK2 inhibitors for 16 hrs (methods). We chose the 16hrs time period because it

corresponds to the end of S-phase in the cycling population (Fig. 2A-B). Afterwards,

CDKs inhibitors were washed out allowing cell cycle progression (Extended data Fig.
3D). G1-arrested tetraploid cells showed low levels of DNA damage, whereas cells
released in S-phase exhibited high levels of DNA damage (Extended data Fig. 3D-G).
Importantly, a certain proportion of γH2AX foci of S-phase tetraploid cells partially co-
localized with markers of active DNA replication sites visualized by Proliferating Cell
Nuclear Antigen (PCNA) and EdU incorporation (Extended data Fig. 3H).

To better characterize DNA damage in tetraploid cells during their first
interphase, we used other markers of the DNA damage signaling and repair pathways.
We found that the number of KU80 and XRCC1 foci, two proteins involved in Non-
Homologous End Joining (NHEJ) [24] remained low (Extended data Fig. 4A-B). In
contrast, the number of RAD51 foci, a protein involved in homologous recombination
(HR), was increased and co-localized with γH2AX in a fraction of tetraploid cells
(Extended data Fig. 4C-D). This was also the case for Replication protein A (RPA) and
FANCD2 foci, two markers of RS which also co-localized with γH2AX foci in tetraploid
cells (Extended data Fig. 4E-G). Together, these results demonstrate that tetraploid
cells experience high levels of DNA damage during S-phase progression, which are
recognized by bonafide DNA damage markers and by the HR repair pathway.

Based on these results, we hypothesized that DNA damage in tetraploid cells
arises from errors occurring during DNA replication. To test this possibility, we arrested
cells in G1 as described above (Extended data Fig. 3D). We then released them in the
presence of very low doses of APH or PHA, a Cdc7 inhibitor. We used low doses of
these compounds to inhibit DNA replication (detected by absence of EdU) without
generating DNA damage (methods). These conditions resulted in the inhibition of DNA
replication, albeit maintaining the biochemical activity typical of the S-phase nucleus.
Strikingly, the levels of DNA damage in tetraploid cells were dramatically decreased
when DNA replication was inhibited (Fig. 2C-D and Extended data Fig. 4H-J).
Importantly, in the few tetraploid cells that escaped DNA replication inhibition -
revealed by high EdU incorporation - a high number of γH2AX foci were noticed
(Extended data Fig. 4K-L), confirming the correlation between DNA replication and
DNA damage in tetraploid cells.

To characterize DNA replication, we established RPE-1 cell lines stably
expressing PCNA chromobodies. We showed that expression of PCNA chromobodies
does not affect cell cycle progression in RPE-1 cells, confirming its suitability to follow
DNA replication by live imaging (Extended data Fig. 4M). Using this cell line, we

performed quantitative 4D live imaging of endogenous DNA replication in diploid and tetraploid cells (methods). Surprisingly, the comparison between the total number of PCNA foci during S-phase in diploid and tetraploid RPE-1 cells, revealed a lack of scaling up with DNA content (Fig. 2E-G), also observed by quantifying the number of EdU foci (Extended data Fig. 4N). This result suggests that fewer replication sites were activated in tetraploid S-phase cells, when compared with diploid cells. The volume of PCNA foci was also lower in tetraploid cells (Fig. 2H). Moreover, timelapse analysis of PCNA dynamics revealed additional differences. As diploid cells enter S-phase, an exponential increase in the number of active replication sites was noticed, which was maintained before undergoing a steep, almost abrupt decreased (Fig. 2E-I, Extended data Fig. 5A and Extended data movie 3). In contrast, in tetraploid cells the increase in the number of active sites was more gradual, and the signals associated with DNA lingered for extended periods of time. Furthermore, the dissociation of PCNA from the DNA in tetraploid cells occurred much later and also in a progressive manner (Fig. 2E-I, Extended data Fig. 5A and Extended data movie 4). In line with this, by analyzing PCNA patterns as a readout of early and late S-phase [25], we showed that tetraploid cells spent more time in early S-phase compared to late S-phase (Extended data Fig. 4O). Surprisingly, even if S-phase was longer in tetraploid cells when compared to diploid cells (Extended data Fig. 4P), this was not sufficient to scale the number of active replication sites with DNA content. These results suggest that DNA replication in tetraploid cells is impaired due to both a lack of scaling up in the number of active replication sites and to a delayed DNA replication timing.

To ascertain if these defects impacted replication fork progression, we performed DNA combing, which allows the visualization of replication origins in single DNA fibers [26]. We failed to obtain fibers of the required quality in RPE-1 cells despite several attempts. To overcome this problem, we performed DNA combing in HCT116 cells since they also showed high levels of DNA damage within the first interphase (Extended data Fig. 2E). Inter-origin distances were not affected in tetraploid cells, however, and surprisingly, median fork speed was increased in tetraploid cells (Fig. 2J and Extended data Fig. 4Q). Further, a high increase in the percentage of unstable forks was also detected (Fig. 2J). These results show that the replication dynamics is perturbed in tetraploid cells when compared to diploid cells.

Since S-phase progression errors and inaccurate DNA replication are linked with a pleiotropy of DNA structural abnormalities, we assessed if unscheduled

tetraploidy was associated with abnormal karyotypes. We FACS sorted tetraploid from diploid cells (see below, Fig. 3C) in G1 and at the G2/M transition to perform single cell DNA sequencing (ssDNAseq) (methods). Normalization of the under and over replicating regions in G1 and G2/M diploid cells revealed already whole chromosome deviations in a certain number of cells. When present, they span along almost all chromosomes of a given cell (Extended data Fig. 5B). In G1 tetraploid cells, over replicated regions (5n) could also be identified, but these were restricted to a few chromosomes and might be explained by a caveat of the method (cells have initiated S-phase but were still selected as G1 by the FACS profile). Striking, however in G2/M tetraploid cells over duplicating chromosomes (> 10) could be identified in addition to frequent over and under replicated regions (9n, 7n and 4n) (Fig. 2K). In agreement with this variability and the extent of copy number deviations, both aneuploidy score and heterogeneity score were increased in G2/M tetraploid cells when compared to G2/M diploid cells (Aneuploidy score: 0.275 vs 0.102; Heterogeneity score 0.319 vs 0.158 respectively) (methods).

Together, our results show that unscheduled tetraploid cells cannot sustain normal DNA replication as they fail to scale proportionally the number of active replication sites and replication timing. Defects in S-phase result in the generation of highly aberrant karyotypes, demonstrating a causal relationship between tetraploidization and GIN within a single S-phase.

Lack of G1 lengthening in tetraploid cells leads to unprepared S-phase
The massive GIN described above, together with abnormal DNA replication dynamics, suggested that newly born tetraploid cells undergo the first S-phase in a non-optimal manner. We reasoned that cells might enter S-phase without the required protein levels to replicate a tetraploid genome. In principle, doubling the whole set of chromosomes should lead to an overall doubling of transcripts and protein translation, so that tetraploid cells should scale up by a factor of 2. To determine if cell mass was increased in tetraploid cells at the G1-S transition, we combined quantitative phase imaging with the cell cycle sensor FUCCI, which allows recording of mass measure trajectories through the cell cycle at the single cell level [27]. We found that the proportion of mass added during G1 was lower in tetraploid cells compared to diploid cells (Fig. 3A-B). These results establish that newly born tetraploid cells are not able to scale together protein and DNA content during the first G1 upon WGD. We next tested the

total levels of key S-phase components. To do so, we used a recently developed
protocol that enables to sort and isolate tetraploid from diploid cells based on FUCCI
and DNA content from a common population (Fig. 3C and Extended data Fig. 6A,
methods). Protein extracts from the two cell populations at the G1/S transition were
then probed by western blot. The same number of cells was loaded for diploid and
tetraploid conditions. Normalization of the chromatin associated H2B variant and the
cytoskeleton component Actin showed an increase in these two protein levels
consistent with DNA doubling (Fig. 3D-E and Extended data Fig. 6B). In stark contrast,
essential S-phase DNA replication factors such as the origin recognition complex 1
(ORC1) involved in the recognition of replication origins [28], the minichromosome
maintenance 2 (MCM2) helicase [29], CDC45, a member of the active helicase complex
[30], and PCNA did not scale up in tetraploid cells (Fig. 3D-F). Combined with quantitative
phase imaging data, these results suggest that tetraploid cells do not contain the
required protein levels to sustain timely and successful DNA replication during S-
phase.

In normal proliferative cell cycles, the growth phase occurring during G1 phase
prepares cells for DNA replication allowing the expression and accumulation of key S-
phase regulators [31,32]. We reasoned that a short G1 duration could account for
transition to S-phase in an unprepared manner, which is supported by the fact that
tetraploid cells did not scale up protein content with DNA doubling (Fig. 3A-F). Indeed,
time lapse analysis of tetraploid cells just after birth indicated only a slight increase in
G1 duration, which did not scale with DNA content when compared to diploid cells (Fig.
3G). Further, while we observed a significant correlation between cell mass and G1
duration in diploid cells, also described in other diploid conditions [33], this correlation
was absent in tetraploid cells suggesting that G1 duration is not dependent of cell mass
in tetraploid cells (Fig. 3H).

We next tested if imposing G1 lengthening was translated by increased protein
levels of S-phase factors and thus in principle enabled error-free DNA replication in
tetraploid cells. To test this model, we delayed S-phase entry using very low doses of
CDK4/6 or CDK2 inhibitors. These conditions were different from the ones described
above used to synchronize cells in G1. Indeed, while high doses of these inhibitors
result in a biochemical arrest, low inhibitor doses result in G1 lengthening [34,35]. The
different impact of high and low doses of CDK4/6 or CDK2 inhibitors could be noticed
by differences in the expression levels of DNA replication factors. Indeed, after G1

lengthening the levels of DNA replication factors scaled up with DNA content in
tetraploid cells, which was not the case when cells were arrested in G1 (Fig. 3I-K vs
3D-F). Consistent with these findings, after G1 lengthening the number and volume of
active replication sites in the subsequent S-phase visualized by monitoring PCNA or
EdU foci scaled up with DNA content in tetraploid cells (Extended data Fig. 6C-E).
Moreover, after G1 extension, PCNA dynamic behavior in tetraploid cells was
comparable to diploid cells (Fig. 3L and Extended data Fig. 6F and I and Extended
data movies 5-6). Even if the time spent in S-phase was not altered after G1
lengthening, we observed that increasing G1 duration restored the ratio between early
and late S-phase in tetraploid cells, suggesting that DNA replication timing was
reinstated (Extended data Fig. 6G-H). Strikingly, G1 lengthening was sufficient to
significantly reduce DNA damage in tetraploid S-phase cells (Fig. 3M and Extended
data Fig. 6J-L).

Altogether, our data show that tetraploid cells transition from G1 to S-phase
prematurely without undergoing scaling of global protein mass, and so they enter in S-
phase with insufficient amounts of DNA replication factors. This impacts the dynamics
and fidelity of DNA replication, generating DNA damage. Importantly, extension of G1
is sufficient to increase the levels of key DNA replication factors, which results in a
significant decrease in DNA damage in tetraploid cells.

G1 lengthening or increased E2F1 levels are sufficient to rescue GIN in tetraploid
cells and in polyploid cells *in vivo*
From yeast to mammals, the transition from G1 to S-phase is negatively regulated by
members of the retinoblastoma (Rb) protein family, which sequesters the transcription
factor E2F1 [31,32,36–38]. E2F1 targets several genes required for entry into S-phase and
DNA replication factors. Since a short G1 does not prepare tetraploid cells for S-phase,
we reasoned that increased levels of E2F1 might override the G1 lengthening defect.
We thus expressed E2F1 in diploid cells (Extended data Fig. 7A), allowing to increase
the expression of DNA replication proteins just before generating tetraploid cells.
Importantly, this was sufficient to rescue the levels of DNA damage in tetraploid cells
(Fig. 4A-B).

We recently characterized an *in vivo* model to study the consequences of
polyploidy in *Drosophila* neural stem cells also called neuroblasts (NBs) in the
developing brain [39]. These cells are normally diploid, but through repeated CF can

generate highly polyploid NBs (here referred to as unscheduled polyploidy) much beyond the tetraploidization status (Fig. 4C-D and Extended data Fig. 7B). A key prediction of our findings is that polyploid NBs should also accumulate high levels of DNA damage *in vivo*. To test this prediction, we determined the levels of DNA damage in unscheduled polyploid NBs during interphase using antibodies against γH2Av to determine the γH2Av index (methods). We compared it with diploid NBs and the programmed polyploid cells from the *Drosophila* salivary gland, which are extremely large and accumulate more than 2000 chromosomes [40]. This represents a 250- fold increase in DNA content, when compared with diploid *Drosophila* cells, which contain only 8 chromosomes. Interestingly, interphase polyploid NBs displayed high levels of DNA damage, which was not the case in diploid NBs or polyploid cells from the salivary gland (Fig. 4D-E and Extended data Fig. 7C). We next increased the levels of E2F1 and Rb by over-expressing (OE) in a tissue-specific manner using the UAS-Gal4 system. E2F1OE increases the expression of cell cycle regulators, while RbOE increases G1 lengthening [41,42]. Strikingly, this was sufficient to decrease substantially the levels of DNA damage in unscheduled polyploid NBs *in vivo* (Fig. 4F-G and Extended data Fig. 7D).

Taken together, these data show that *in vivo* unscheduled polyploidy is a source of DNA damage and GIN, which can be inhibited by increased E2F1 or Rb levels. These results put forward the idea that lack of cell cycle readjustment promotes GIN and the accumulation of highly complex karyotypes in cells that are not programmed to increase their DNA content (Fig. 4H).

Here, we analyzed the initial defects following WGD and identified a very early window of high GIN that could promote acquisitions of multiple mutations making it possible to bypass cell cycle controls while promoting tetraploid cell survival. Our results are consistent with a model where tetraploid cells transit through the first cell cycle without preparing the duplication of increased DNA content (Fig. 4H). We found defective fork progression rates in tetraploid cells, which surprisingly seem to progress faster than in diploid cells. Although the increased rates remain to be explained, these may contribute to RS and GIN as recently shown upon PARP inhibition [43] or in response to decreased levels of MCM proteins [44]. Strikingly, G1 extension or the increase in the expression levels of cell cycle proteins *in vivo* lowered considerably the high DNA damage levels of extreme polyploid cells such as the ones generated by

repeated CF. These results highlight the importance of keeping constant scaling up
between DNA and protein content to ensure genetic stability and cell homeostasis.

The most surprising finding of this study is the lack of scaling up between DNA
and protein content immediately after tetraploidization. In physiological conditions,
such as during animal development, WGDs and polyploidization lead to an overall
scaling up of cell mass and DNA content to favor increase in secretion and metabolic
activity for example [20,45,46]. Our work shows that unscheduled tetraploid or polyploid
cells do not increase cell mass as expected. Why certain key cell cycle and DNA
replication factors fail to be expressed at levels that allow optimal DNA replication
remains to be explained. Importantly, however these results show that an immediate
consequence of unscheduled genome doubling is loss of genetic integrity within a
single S-phase. Interestingly, studies performed on stable tetraploid cells have shown
a remarkable scaling up between protein and DNA content after long term adaptation
[47,48]. It is tempting to propose that in non-physiological conditions, as the ones studied
here, newly born tetraploids do not "feel" the increase in DNA content and so, cannot
adapt G1 duration or protein content in order to replicate a 4N genome. It will be
interesting to identify the molecular mechanisms that promote ploidy increase while
maintaining genetic stability and cell homeostasis.

Acknowledgments

The authors acknowledge the Cell and Tissue Imaging platform (PICT-IBiSA), member of the French National Research Infrastructure France-BioImaging (ANR10-INBS-04) and the Nikon Imaging center from Institut Curie for microscopy. We thank L. Guyonnet, A. Chipont and C. Guerrin from the Cytometry platform of Institut Curie. We thank V. Marthiens, S. Lambert, E. Schwob, J.S. Hoffmann, D. Fachinetti, M. Budzyk, F. Edwards, O. Goundiam, G. Fantozzi, R. Salamé, A. Goupil and C. Chen for helpful discussions and/or comments on the manuscript. This work was supported by FOR2800/STO918-7 to Z.S, ERC CoG (ChromoNumber-LS3, ERC-2016-COG) for R.B, Institut Curie and the CNRS. The Basto lab is a member of the Cell(n)Scale Labex.

Author contributions

S.G. and R.B. conceived the project and wrote the manuscript. S.G. did most of the experiments and data analysis presented here. M.N. did the initial observations of high levels of DNA damage in *Drosophila* polyploid NBs. S.V.B., K.K. and Z.S. did the DNA combing. R.W., A.E.T., D.C.J.S. and F.F. did the scSeq and BI analysis. A.S.M helped with image quantifications and analysis. N.S. and M.P. performed the quantitative phase imaging experiments and analysis and H.H. contributed with unpublished cell lines. All authors read and comment on the manuscript.

METHOD DETAILS:

Cell culture, generation of cell lines and treatments:

Cell culture:

Cells were maintained at 37°C in a 5% CO_2 atmosphere. hTERT RPE-1 cells (ATCC Cat# CRL-4000, RRID:CVCL_4388) and HEK293 cells (ATCC Cat# CRL-1573, RRID:CVCL_0045) were grown in Dulbecco's modified medium (DMEM) F12 (11320-033 from Gibco) containing 10% fetal bovine serum (GE Healthcare), 100 U/ml penicillin, 100 U/ml streptomycin (15140-122 from Gibco). BJ cells (ATCC Cat# CRL-4001, RRID:CVCL_6573) and HCT116 cells (ATCC Cat# CCL-247, RRID:CVCL_0291) were grown in Dulbecco's modified medium + GlutaMAX (61965-026 from Gibco) containing 10% fetal bovine serum (GE Healthcare), 100 U/ml penicillin, 100 U/ml streptomycin (15140-122 from Gibco).

All cells were routinely checked for mycoplasma infection.

Generation of RPE-1 PCNA^chromo stable cell line:

RPE-1 cells were transfected with 10µg Cell Cycle-Chromobody® plasmid (TagRFP) (From Chromotek, Planegg, Germany) using JET PRIME kit (Polyplus Transfection, 114-07) according to the manufacturer protocol. After 24 hours, 500µg/ml G418 (4727878001 from Sigma Aldrich) was added to the cell culture medium and then clones expressing PCNA chromobodies were selected.

Generation of a RPE-1 FUCCI or RPE-1 CCNB1^AID FUCCI stable cell line:

To produce lentiviral particles, HEK293 cells were transfected with 4µg pBOB-EF1-FastFUCCI-Puro (86849 from Addgene, RRID:Addgene_86849) + 4µg pMD2.G (12259 from Addgene, RRID:Addgene_12259) + 4µg psPAX2 (12260 from Addgene, RRID:Addgene_12260) using FuGENE HD Transfection Reagent (E2311 from Promega) in OptiMEM medium (51985034 from ThermoFisher). Cells were incubated at 37°C in a 5% CO2 atmosphere for 16 hours and then growth media were removed and replaced by 5 ml fresh OptiMEM. The following day, viral particles were isolated by filtering the medium containing the viral particles through a 0.45µm filter (16537 from Sartorius stedim biotech). Then, RPE-1 or RPE-1 CCNB1^AID [49] cells were incubated with viral particles in the presence of 8µg/ml polybrene (sc-134220 from Santa Cruz) at 37°C in a 5% CO_2 atmosphere for 24 hours. RPE-1 GFP and RFP-

positive cells were then collected using Sony SH800 FACS (BD FACSDiva Software

Version 8.0.1). RPE-1 or RPE-1 CCNB1[AID] clones expressing FUCCI were selected

and the cell lines were established from one single clone.

pBOB-EF1-FastFUCCI-Puro was a gift from Kevin Brindle & Duncan Jodrell (Addgene

plasmid # 86849 ; http://n2t.net/addgene:86849 ; RRID:Addgene_86849) [50].

Generation of RPE-1 GFP-53BP1 RFP-H2B stable cell line:

This cell line was obtained as described below. Briefly, to produce lentiviral particles,

HEK293 cells were transfected with 4µg pSMPUW-IRIS-Neo-H2BmRFP (Fachinetti

Lab) + 4µg pMD2.G (12259 from Addgene, RRID:Addgene_12259) + 4µg psPAX2

(12260 from Addgene, RRID:Addgene_12260). Then, RPE-1 cells were incubated with

viral particles and RPE-1 RFP-positive cells were collected using Sony SH800 FACS

(BD FACSDiva Software Version 8.0.1). RPE-1 clones expressing RFP-H2B were

selected, and the cell line was established from one single clone.

Then, new lentiviral particles were produced by transfecting HEK293 cells with 4µg

Apple-53BP1trunc (69531 from Addgene, RRID:Addgene_69531) + 4µg pMD2.G

(12259 from Addgene, RRID:Addgene_12259) + 4µg psPAX2 (12260 from Addgene,

RRID:Addgene_12260). RPE-1 RFP-H2B cells were incubated with viral particles and

RPE-1 clones expressing both RFP-H2B and GFP-53BP1 were selected using flow

cytometry (FACS SH800 from Sony) and the cell line was established from one single

clone.

Apple-53BP1trunc was a gift from Ralph Weissleder (Addgene plasmid # 69531 ;

http://n2t.net/addgene:69531 ; RRID:Addgene_69531) [51].

Generation of tetraploid cells:

Mitotic slippage using drugs: cells were incubated with DMSO (D8418 from Sigma

Aldrich) or with 50µM monastrol (S8439 from Selleckchem) + 1µM MPI-0479605

(S7488 from Selleckchem) for at least two hours. This approach was used in Fig. 1B-

D; Fig. 2A-I: Fig. 3L-M; Fig. 4A-B; Extended data Fig. 1A-C and I-M; Extended data

Fig. 2D-E; Extended data Fig.3B-E and G-H; Extended data Fig. 4A-H; K and N-P;

Extended data Fig. 5A; Extended data Fig. 6C-H and J-L; Extended data Fig. 7A.

Mitotic slippage using genetic tools: CCNB1 depletion in RPE CCNB1[AID] cells was

induced as described before[49]. Briefly, cells were treated with 2µg/ml doxycycline

(D3447 from Sigma Aldrich) + 3µM asunaprevir (S4935 from Selleckchem) for 2 hours.

Then, 500 µM auxin (I5148 from Sigma Aldrich) was added to the cell culture medium

for at least 4 hours. This approach was used in Fig1K; Fig. 3A-K; Extended data Fig.

2A; Extended data Fig. 6A-B.

Cytokinesis failure using drugs: cells were incubated with 10µM genistein (G6649

from Sigma Aldrich) for at least two hours. This approach was used in Fig. 1E-F;

Extended data Fig. 1DF; Extended data Fig. 2D-E; Extended data Fig. 3A and F;

Extended data Fig. 3I-J; Extended data Fig. 6I. Alternatively, cell were incubated with

0.75µM Dihydrocytochalasin D (D1641 from Sigma-ALdrich) for 1 hour. This approach

was used in Fig. 2J; Extended data Fig. 2B; Extended data Fig. 4Q.

Endoreplication using drugs: cells were incubated with 10µM SP600125 (S1460

from Selleckchem) for at least two hours. This approach was used in Fig. 1G-H and

Extended data Fig. 1G-I.

Endoreplication using genetic tools: CCNA2 depletion in RPE CCNA2^{AID} cells was

induced as described before[49]. Briefly, cells were treated with 2µg/ml doxycycline

(D3447 from Sigma Aldrich) for 2 hours. Then, 500 µM auxin (I5148 from Sigma

Aldrich) + 3µM asunaprevir (S4935 from Selleckchem) was added to the cell culture

medium for at least 4 hours. This approach was used in Extended data Fig. 2C.

Cell cycle synchronization and DNA replication inhibition:

Cells were treated with 1µM palbociclib (Cdk4/6 inhibitor, S1579 from Selleckchem),

with 0.5µM abemaciclib (Cdk4/6 inhibitor, S5716 from Selleckchem) or with 1µM

K03861(Cdk2 inhibitor, S8100 from Selleckchem) for 16 hours to synchronize cells at

G1/S transition and were collected (indicated by "G1 arrest" in the figures).

Alternatively, cells were then washed five times using PBS 1X and released in S-phase

for 10 hours before being collected (indicated by "Release in S-phase" in the figures).

To inhibit DNA replication, cells were released in S-phase in the presence of low doses

of Aphidicolin (APH, A0781 from Sigma-Aldrich), a DNA replication polymerase

inhibitor, or of PHA767491 (PZ0178 from Sigma-Aldrich), a Cdc7 inhibitor (indicated

by "Release in S-phase + APH" or "Release in S-phase + PHA", respectively, in the

figures). Doses were chosen to significantly decrease EdU incorporation without

affecting the levels of DNA damage.

Treatments:

Names:	Companies:	Targets:	References:	Concentrations:
Auxin	Sigma	AID system	I5148	500 µM
Doxycycline	Sigma Aldrich	AID system	D3447	2µg/ml
Asunaprevir	Selleckchem	AID system	S4935	3µM
Monastrol	Selleckchem	Eg5	S8439	50 µM
MPI-0479605	Selleckchem	MPS1	S7488	1µM
Genistein	Sigma Aldrich	MKLP1	G6649	10µM
SP600125	Selleckchem	JNK	S1460	10µM
Abemaciclib	Selleckchem	CDK4/6	S5716	50nM or 0.5µM
K03861	Selleckchem	CDK2	S8100	400nM or 1µM
Palbociclib	Selleckchem	CDK4/6	S1579	120nM or 1µM
Aphidicolin	Sigma Aldrich	DNA polymerase	A0781	0.4µM or 1µM
Hydroxyurea	Selleckchem	RNR	S1896	2mM
PHA 767491	Sigma Aldrich	Cdc7	PZ0178	1µM
RO3306	Calbiochem	CDK1	217699	10µM
Dihydrocytochalasin D	Sigma Aldrich	Actin	D1641	0.75µM
5'-Chloro-2'-deoxyuridine (CldU)	Sigma Aldrich	DNA	C6891	100µM
5'-Iodo-2'-deoxyuridine (IdU)	Sigma Aldrich	DNA	I7125	100µM

Fly husbandry and fly stocks:

Flies were raised on cornmeal medium (0.75% agar, 3.5% organic wheat flour, 5.0% yeast, 5.5% sugar, 2.5% nipagin, 1.0% penicillin-streptomycin and 0.4% propionic acid). Fly stocks were maintained at 18°C. Crosses were carried out in plastic vials and maintained at 25°C. Stocks were maintained using balancer inverted chromosomes to prevent recombination. Stocks used in this study: *sqh1* [52], *pavarotti* RNAi (BL#42573 from Bloomington Drosophila Stock Center, Indiana University, IN, USA) [39], UAS-E2F1 (F001065 from FlyORF, Zurich, Switzerland) and UAS-Rb (BL# 50746 from Bloomington Drosophila Stock Center, Indiana University, IN, USA).

In all experiments, larvae were staged to obtain comparable stages of development. Egg collection was performed at 25°C for 24 hours. After development at 25°C, third instar larvae were used for dissection.

Immunofluorescence microscopy and antibodies:

Preparation and imaging of human cells

Cells were plated on cover slips in 12-well plates and treated with the indicated drugs. To label cells, they were fixed using 4% of paraformaldehyde (15710 from Electron Microscopy Sciences) + Triton X-100 (2000-C from Euromedex) 0,1% in PBS (20 min at 4°C). Then, cells were washed three times using PBS-T (PBS 1X + 0,1% Triton X-100 + 0,02% Sodium Azide) and incubated with PBS-T + BSA (04-100-812-C from Euromedex) 1% for 30 min at RT. After three washes with PBS-T + BSA, primary and secondary antibodies were incubated in PBS-T + BSA 1% for 1 hr and 30 min at RT, respectively. After two washes with PBS, cells were incubated with 3 µg/ml DAPI (4',6-diamidino-2-phenylindole; D8417 from Sigma Aldrich) for 15 min at RT. After two washes with PBS slides were mounted using 1.25% n-propyl gallate (Sigma, P3130), 75% glycerol (bidistilled, 99.5%, VWR, 24388-295), 23.75% H2O.

Images were acquired on an upright widefield microscope (DM6B, Leica Systems, Germany) equipped with a motorized XY and a 40X objective (HCX PL APO 40X/1,40-0,70 Oil from Leica). Acquisitions were performed using Metamorph software (Molecular Devices, USA) and a sCMOS camera (Flash 4V2, Hamamatsu, Japan). Stacks of conventional fluorescence images were collected automatically at a Z-distance of 0.5 µm (Metamorph software; Molecular Devices, RRID:SCR_002368). Images are presented as maximum intensity projections generated with ImageJ software (RRID:SCR_002285).

Whole mount tissue preparation and imaging of Drosophila larval brains

Brains or Salivary glands from third instar larvae were dissected in PBS and fixed for 30 minutes in 4% paraformaldehyde in PBS. They were washed three times in PBST 0.3% (PBS, 0.3% Triton X-100 (T9284, Sigma), 10 minutes for each wash) and incubated for several hours in agitation at room temperature (RT) and O/N at 4°C with primary antibodies at the appropriate dilution in PBST 0.3%. Tissues were washed three times in PBST 0.3% (10 minutes for each wash) and incubated O/N at 4°C with secondary antibodies diluted in PBST 0.3%. Brains and salivary glands were then washed two times in PBST 0.3% (30 minutes for each wash), rinsed in PBS and incubated with 3 µg/ml DAPI (4',6-diamidino-2-phenylindole; D8417 from Sigma

528   Aldrich) at RT for 30min. Brains and salivary glands were then washed in PBST 0.3%
529   at RT for 30 minutes and mounted in mounting media. A standard mounting medium
530   was prepared with 1.25% n-propyl gallate (Sigma, P3130), 75% glycerol (bidistilled,
531   99.5%, VWR, 24388-295), 23.75% H2O.
532
Images were acquired on a spinning disk microscope (Gataca Systems, France).
Based on a CSU-W1 (Yokogawa, Japan), the spinning head was mounted on an
inverted Eclipse Ti2 microscope equipped with a motorized XY Stage (Nikon, Japan).
Images were acquired through a 40X NA 1.3 oil objective with a sCMOS camera
(Prime95B, Photometrics, USA). Optical sectioning was achieved using a piezo stage
(Nano-z series, Mad City Lab, USA). Gataca Systems' laser bench was equipped with
405, 491 and 561 nm laser diodes, delivering 150 mW each, coupled to the spinning
disk head through a single mode fibre. Multi-dimensional acquisitions were performed
using Metamorph 7.10.1 software (Molecular Devices, USA). Stacks of conventional
fluorescence images were collected automatically at a Z-distance of 1.5 μm
(Metamorph software; Molecular Devices, RRID:SCR_002368). Images are presented
as maximum intensity projections generated with ImageJ software
(RRID:SCR_002285).
546
Primary and secondary antibodies
Primary and secondary antibodies were used at the following concentrations: Guinea
pig anti CEP192 antibody (1/500; Basto lab)[53], rabbit anti beta catenin (1/250; C2206
from Sigma-Aldrich, RRID:AB_476831), mouse anti-gamma H2A.X phospho S139
(1/1000; ab22551 from Abcam, RRID:AB_447150), mouse anti-XRCC1 (1/500;
ab1838 from Abcam, RRID:AB_302636), rabbit anti-Rad51 (1/500; ab133534 from
Abcam, RRID:AB_2722613), mouse anti-KU80 (1/200; MA5-12933 from
ThermoFisher, RRID:AB_10983840), rabbit anti-FANCD2 (1/150; NB100-182SS from
Novusbio, RRID:AB_1108397), rabbit anti-γH2Av (1/500; 600-401-914 from Rockland;
RRID: AB_11183655), Alexa Fluor® 647 Phalloidin (1/250; A22287 from
ThermoFisher Scientific, RRID:AB_2620155), goat anti-Rabbit IgG (H+L) Highly
Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 (1/250; A21245 from
ThermoFisher, RRID:AB_2535813), Goat anti-Guinea Pig IgG (H+L) Highly Cross-
Adsorbed Secondary Antibody, Alexa Fluor 488 (1/250; A11073 from ThermoFisher,
RRID:AB_253411), Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody,

Alexa Fluor 546 (1/250, A11003 from ThermoFisher, RRID:AB_2534071), Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 546 (1/250; A-11035 from Thermo Fisher Scientific, RRID:AB_2534093).

Quantitative analysis of DNA damage:

Analysis of Drosophila NBs

Staged 3rd instar larval brains were dissected, stained and imaged using the procedures described above. We used the γH2Av primary antibody, which was preferentially detected using a secondary antibody conjugated Alexa Fluor 546. We used this secondary antibody because it was found to provide the best signal to noise ratio.

Quantitative analysis of DNA damage was carried out as previously described [39]. In brief, DNA damage was assessed using a γH2Av primary antibody detected with an Alexa Fluor secondary antibody. Confocal volumes were obtained with optical sections at 1.5µm intervals. Image analysis was performed using Fiji and a custom plugin developed by QUANTACELL. After manual segmentation of the nuclei, a thresholding operation was used to determine the percentage of γH2Av positive pixels (coverage) and their average intensity in a single z plane in the center of the nucleus. Coverage and intensity were multiplied to obtain the γH2Av index.

Analysis of human cell lines

For DNA damage quantification, the signals obtained in cultured cells were different from the signals found in *Drosophila* NBs. To asses DNA damage in human cells, we used an ImageJ software-based plugin developed by QUANTACELL, where γH2AX signals were measured using z-projection stacks after thresholding. Both FI and the percentage of nuclear coverage was obtained for each nucleus. γH2AX index was obtained multiplying FI by the coverage.

All data plotting and statistical analyses were performed using the GraphPad Prism software.

Time lapse microscopy:

Cells were plated on a dish (627870 from Dutscher) and treated with the indicated

drug. Images were acquired on a spinning disk microscope (Gataca Systems, France).

Based on a CSU-W1 (Yokogawa, Japan), the spinning head was mounted on an

inverted Eclipse Ti2 microscope equipped with a motorized XY Stage (Nikon, Japan).

Images were acquired through a 40X NA 1.3 oil objective with a sCMOS camera

(Prime95B, Photometrics, USA). Optical sectioning was achieved using a piezo stage

(Nano-z series, Mad City Lab, USA). Gataca Systems' laser bench was equipped with

405, 491 and 561 nm laser diodes, delivering 150 mW each, coupled to the spinning

disk head through a single mode fiber. Multi-dimensional acquisitions were performed

using Metamorph 7.10.1 software (Molecular Devices, USA). Stacks of conventional

fluorescence images were collected automatically at a Z-distance of 0.5 µm

(Metamorph software; Molecular Devices, RRID:SCR_002368). Images are presented

as maximum intensity projections generated with ImageJ software

(RRID:SCR_002285), from stacks deconvolved with an extension of Metamorph

software.

EdU staining:

EdU incorporation into DNA was visualized with the Click-it EdU imaging kit (C10338

from Life Technologies), according to the manufacturer's instructions. EdU was used

at a concentration of 1µM (Extended data Fig. 4N and 6E) or 10 µM (Extended data

Fig. 4A and I) for the indicated time. Cells were incubated with the Click-it reaction

cocktail for 15 minutes.

FACS sorting of diploid and tetraploid cells:

A mix of diploid and tetraploid cells (*see "generation of tetraploid cells" section*) were

incubated with 2µg/ml Hoescht (94403 from Sigma Aldrich) for 1 hour at 37°C, 5%

CO_2. Then, a single cell suspension was generated. Cells were washed using PBS 1X,

the supernatant was removed and cells were resuspended in cold cell culture medium

at $1x10^7$ cell per ml and kept at 4°C during all the experiment. FACS sorting was

performed using Sony SH800 FACS (BD FACSDiva Software Version 8.0.1).

Compensation was performed using the appropriate negative control samples.

Experimental samples were then recorded and sorted using gating tools to select the

populations of interest. RFP+ / GFP- negative cells (G1 cells) were first selected. Then,

in this population, DNA content was used to segregate diploid (2n) and tetraploid (4n)

G1 cells. Once gates have been determined, diploid and tetraploid G1 cells were

sorted into external collection tubes. Post-sort analysis was performed to determine

the purity of the sorted populations (*see Extended data Fig. 6A*).

E2F1 overexpression:

RPE-1 cells were transfected using 0.25µg pCMVHA E2F1 (24225 from Addgene,

RRID:Addgene_24225) using JET PRIME kit (Polyplus Transfection, 114-07)

according to the manufacturer's protocol. Five hours later, cells were incubated with

DMSO (D8418 from Sigma Aldrich) or with 50µM monastrol (S8439 from Selleckchem)

+ 1µM MPI-0479605 (S7488 from Selleckchem) to generate tetraploid cells. After 2

hours, DMSO or 1µM palbociclib (S1579 from Sellechem) were added to the cell

culture medium for 16 hours. Cells were then fixed in G1 (T0) or washed five times

using PBS and released in S-phase and fixed after 10 hours (T10). The

immunofluorescence protocol is described in the corresponding section.

pCMVHA E2F1 was a gift from Kristian Helin (Addgene plasmid # 24225 ;

http://n2t.net/addgene:24225 ; RRID:Addgene_24225) [54].

Western Blot analysis and antibodies:

Cells were lysed in 8 M urea, 50 mM Tris HCl, pH 7.5 and 150 mM β-mercaptoethanol

(161-0710 from Bio-Rad), sonicated and heated at 95°C for 10 minutes. Samples

(equivalent of 2×10^5 cells) were subjected to electrophoresis in NuPAGE Novex 4–

12% Bis-Tris pre-cast gels (NP0321 from Life Technologies). Protein fractions from the

gel were electrophoretically transferred to PVDF membranes (PVDF transfer

membrane; RPN303F from GE). After 1 hr saturation in PBS containing 5% dry non-

fat milk and 0.5% Tween 20, the membranes were incubated for 1 hr with a primary

antibody (*see below*) diluted in PBS containing 5% dry non-fat milk and 0.5% Tween

20. After three 10-min washes with PBS containing 0.5% Tween 20, the membranes

were incubated for 45 min with a 1/2 500 dilution of peroxidase-conjugated antibody

(*see below*). Membranes were then washed three times with PBS containing 0.5%

Tween 20, and the reaction was developed according to the manufacturer's

specifications using ECL reagent (SuperSignal West Pico Chemiluminescent

Substrate; 34080 from Thermo Scientific). Protein levels were normalized using H2B

signal and quantifications were done using Image Lab software version 6.0.1, Bio-Rad

Laboratories.

Primary and secondary antibodies were used at the following concentrations:

Mouse anti Tubulin (1/5000; T9026 from Sigma, RRID:AB_477593), mouse anti CDC45 (1/500; sc-55569 from Santa Cruz Biotechnology, RRID:AB_831146), rabbit anti PCNA (1/500; sc56 from Santa Cruz, RRID:AB_628110), rabbit anti Actin (1/2000; A5060 from Sigma-Aldrich, RRID:AB_476738), mouse anti-H2B (1/1000; sc-515808 from Santa Cruz Biotechnology), mouse anti ORC1 (1/500; sc-398734 from Santa Cruz Biotechnology), mouse anti MCM2 (1/500; 610701 from BD Biosciences, RRID:AB_398024), Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, HRP (1/2500; G21234 from ThermoFisher, RRID:AB_2536530), Peroxidase AffiniPure Goat Anti-Mouse IgG (H+L) (1/2500; 115-035-003 from Jackson ImmunoResearch, RRID:AB_10015289).

3D reconstruction and analysis on Imaris:

3D movies (see *time lapse microscopy* section) were imported into Imaris software v.9.6.0 (Bitplane, RRID:SCR_007370). For chosen cells, the module "Spot tracking" of Imaris was used to detect the foci, as spots of diameter 0.5 µm in the XY-direction and 1µm in Z-direction (modelling PSF elongation). Because the volume of the foci changes in time, the option "Enable growing regions" was used. In each movie, the threshold was chosen on the brightest frame (to detect a maximum of the correct spots) and then applied to the whole movie. For each cell, at each time point, the number of spots and volumes were recorded. For each condition, at least 10 cells were studied and the statistics from Imaris were averaged at each time point using a MATLAB script.

Molecular combing and antibodies:

Tetraploid HCT116 were generated by cytokinesis inhibition using 0.75 µM dihydrocytochalasin D (DCD, inhibitor of actin polymerization, D1641 from Sigma-Aldrich) for 18 h overnight. Afterwards, the cells were washed three times with PBS and cultured in DMEM supplemented with 10% FBS and 1% Pen/Strep for additional 10 h. Cells were pulse-labelled with 0.1 mM CIdU and 0.1 mM IdU for 30min and 100 000 cells per condition were collected for further analysis. The DNA was extracted from cells and prepped following the manufacturer's instructions using the FiberPrep® DNA Extraction Kit (Genomic Vision, Bagneux, France). Subsequently, the prepped DNA was stretched onto coated glass coverslips (CombiCoverslips™, Genomic Vision,

Bagneux, France) by using the FiberComb Molecular Combing System (Genomic
Vision, Bagneux, France). The Labelling was performed with antibodies against
ssDNA, IdU and CldU using the Replication Combing Assay (RCA) (Genomic Vision,
Bagneux, France). The imaging of the prepared cover slips was carried out by
Genomic Vision (Bagneux, France) and analysed using the FiberStudio® 2.0.1
Analysis Software by Genomic Vision.
702
Antibodies were used at the following concentrations:
Rabbit anti ssDNA (1/5; 18731 from IBL International, RRID:AB_494649), Rat anti
CldU (1/10; Ab6326 from Abcam, RRID:AB_2313786), Mouse anti IdU (1/10; 555627
from BD Biosciences, RRID:AB_10015222), mouse Alexa Fluor 647 Donkey (1/25;
JIM-715-605-151 from Biozol), Rat Alexa Fluor 594 Donkey (1/25; JIM-712-585-153
from Biozol), Rabbit Brilliant Violet 480 Donkey (1/25; 711-685-152 from Jakcson
Immuno Research, RRID:AB_2651109).
710
Quantitative phase imaging and measurements
Cells were plated on glass-bottom dishes coated with 50 µg/ml Fibronectin for 1 hour
and rinsed, and trypsinised cells were plated at a concentration of $1.5*10^6$ cells/ml. The
cells used for the experiments were seeded in T-25 dishes at a concentration of $0.7*10^6$
cells/ml 2 days before the actual experiment. On the day of the experiment, the cells
were detached with EDTA (versene), and plated at a concentration of $1.5*10^6$ cells/ml.
For inducing tetraploidy, cells were treated with 2µg/ml doxycycline (D3447 from Sigma
Aldrich) for 2 hours. Then, 500 µM auxin (I5148 from Sigma Aldrich) + 3µM asunaprevir
(S4935 from Selleckchem) was added to the cell culture medium for at least 4 hours.
The cells were then imaged for 35 hours every 20 minutes to track them throughout
their cell cycle.
The cell cycle state of the cells was indicated by the FUCCI system; G1 cells express
Cdt1-RFP while S/G2 cells express hGeminin-GFP and mitosis is indicated by the
NEBD with geminin being present through the cells [55]. To quantify the fluorescence of
geminin in the nucleus, first a background subtraction was performed on the images.
An ROI was used to define an area containing the background fluorescence in the
image. An average value of the ROI was then subtracted from all the frames.
Subsequently, a ROI was drawn as close as possible to the cell, and then the mean

gray value was measured across all the frames. This helped identify the frames of birth

and G1/S transition during cell cycle.

A detailed protocol for the mass measurement with phasics camera is available in [56,57].

Images were acquired by Phasics camera every 20 min for 35 hours for the duration

of the experiment. To obtain the reference image, 32 empty fields were acquired on

the dish and a median image was calculated. This reference image was subtracted

from the interferograms (images acquired by phasics) by custom written MATLAB

scripts to measure the optical path difference. They were then processed to calculate

the phase, intensity and phase cleaned images (the background set to 1000 and the

field cropped to remove edges). Background normalization was performed using a

gridfit method and a watershed algorithm was used to separate cells which came in

contact with each other. Mass was calculated by integrating the intensity of the whole

cell.

Sequencing and AneuFinder analysis:

A mixed population of diploid and tetraploid RPE-1 CCNB1[AID] FUCCI cells were

synchronized in G1 using 1µM palbociclib (S1579 from Selleckchem) for 16 hours or

released in S-phase for 20 hours in the presence of 10µM RO3306 (217699 from

Calbiochem) in order to block cells in the subsequent G2/M. G1 and G2/M diploid and

tetraploid cells were then isolated using cell sorting (see "*FACS sorting of diploid and*

tetraploid cells" section) and collected in a 96-well plate. Single-cell sequencing was

performed as described in detail in [58]. Briefly, cells were lysed to prepare a suspension

of nuclei and sorted as single nuclei in 96 or 384 well plates. Next, single cell

sequencing libraries were prepared using a semi-automated liquid handler platform

(Bravo, Agilent technologies). For library preparation, chromatin was fragmented by

micrococcal nuclease, end-repaired, and A-tailed, followed by Illumina adapter ligation.

Libraries were then cleaned up and PCR-amplified for 17 cycles that included the

addition of a library-specific barcode to uniquely label individual cell libraries. Up to 384

libraries were pooled and sequenced on a Nextseq 500 machine (Illumina; up to 77

cycles; single end). The generated data were subsequently demultiplexed using

sample-specific barcodes and changed into fastq files using bcl2fastq (Illumina;

version 1.8.4). Reads were afterwards aligned to the human reference genome

(GRCh38/hg38) using Bowtie2 (version 2.2.4)[59]. Duplicate reads were marked with

BamUtil (version 1.0.3)[60];. The aligned read data (bam files) were analyzed with a copy

number calling algarithm called AneuFinder (https://github.com/ataudt/aneufinder)[61]. Following GC correction and blacklisting of artefact-prone regions (extreme low or high coverage in control samples), libraries were analyzed using the dnacopy and edivisive copy number calling algorithms with variable width bins (average binsize = 1 Mb; step size = 500 kb). The G1 samples were used as reference for the analysis of the G2/M samples (G1 diploid for G2/M diploid and G1 polyploid for G2/M polyploid). The G1 samples were analyzed with an euploid reference[62]. Results were afterwards curated by requiring a minimum concordance of 90 % (4N and 8N samples) or 95% (2N samples) between the results of the two algorithms. Libraries with on average less than 10 reads per bin (~ 30,000 reads for a diploid genome) were discarded. The aneuploidy scores corresponds to the absolute difference from euploid genome and is the average from all bins and all libraries of one sample. Heterogeneity scores is calculated as the proportion of pairwise comparisons, between libraries, that shows different copy numbers. This is first calculated for each bin. To get to the final score a weighted average is applied.

QUANTIFICATION AND STATISTICAL ANALYSIS:

Quantifications:
Image analysis and quantifications were performed using Image J software V2.1.0/1.53c, https://imagej.net/software/fiji/downloads. To quantify the colocalizations between two signals (Extended data Fig. 3M and 4D) we used *JACOP* plugin with Image J software. 3D movies (Fig. 2F and Extended data Fig. 3B) were corrected using *3D correct drift* plugin with Image J software to keep the cell of interest at the centre of the region of interest. The nuclear area and DAPI intensity were measured using the *wand* tool with Image J software. For the figures, images were processed on Image J software, and mounted using Affinity Designer, https://affinity.serif.com/fr/designer/.

Statistical analysis:
At least three (n) independent experiments were carried out to generate each dataset, and the statistical significance of differences was calculated with Student's t-test. These tests were performed using GraphPad Prism (RRID:SCR_002798) version 7.00 for Mac, GraphPad Software, La Jolla California USA, www.graphpad.com.

REFERENCES:

1. Zack, T. I. *et al.* Pan-cancer patterns of somatic copy number alteration. *Nat. Genet.* **45**, 1134–1140 (2013).

2. Bielski, C. M. *et al.* Genome doubling shapes the evolution and prognosis of advanced cancers. *Nat. Genet.* **50**, 1189–1195 (2018).

3. López, S. *et al.* Interplay between whole-genome doubling and the accumulation of deleterious alterations in cancer evolution. *Nat. Genet.* **52**, 283–293 (2020).

4. Storchova, Z. & Kuffer, C. The consequences of tetraploidy and aneuploidy. *Journal of Cell Science* vol. 121 3859–3866 (2008).

5. Storchova, Z. & Pellman, D. From polyploidy to aneuploidy, genome instability and cancer. *Nature Reviews Molecular Cell Biology* vol. 5 45–54 (2004).

6. Dewhurst, S. M. *et al.* Tolerance of whole- genome doubling propagates chromosomal instability and accelerates cancer genome evolution. *Cancer Discov.* **4**, 175–185 (2014).

7. Fox, D. T., Gall, J. G. & Spradling, A. C. Error-prone polyploid mitosis during normal Drosophila development. *Genes Dev.* **24**, 2294–2302 (2010).

8. Goupil, A. *et al.* Chromosomes function as a barrier to mitotic spindle bipolarity in polyploid cells. *J. Cell Biol.* **219**, (2020).

9. Crockford, A. *et al.* Cyclin D mediates tolerance of genome-doubling in cancers with functional p53. *Ann. Oncol.* **28**, 149–156 (2017).

10. Potapova, T. A., Seidel, C. W., Box, A. C., Rancati, G. & Li, R. Transcriptome analysis of tetraploid cells identifes cyclin D2 as a facilitator of adaptation to genome doubling in the presence of p53. *Mol. Biol. Cell* **27**, 3065–3084 (2016).

11. Andreassen, P. R., Lohez, O. D., Lacroix, F. B. & Margolis, R. L. Tetraploid state induces p53-dependent arrest of nontransformed mammalian cells in G1. *Mol. Biol. Cell* **12**, 1315–1328 (2001).

12. Storchová, Z. *et al.* Genome-wide genetic analysis of polyploidy in yeast. *Nature* **443**, 541–547 (2006).

13. Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human cancer. *Nat. Biotechnol.* **30**, 413–421 (2012).

14. Ganem, N. J., Storchova, Z. & Pellman, D. Tetraploidy, aneuploidy and cancer. *Current Opinion in Genetics and Development* vol. 17 157–162 (2007).

832   15.   Gemble, S. & Basto, R. CHRONOCRISIS: When Cell Cycle Asynchrony
833   Generates DNA Damage in Polyploid Cells. *BioEssays* **42**, (2020).

834   16.   Kuznetsova, A. Y. *et al.* Chromosomal instability, tolerance of mitotic errors and
835   multidrug resistance are promoted by tetraploidization in human cells. *Cell*
836   *Cycle* **14**, 2810–2820 (2015).

837   17.   Jemaà, M. *et al.* Whole-genome duplication increases tumor cell sensitivity to
838   MPS1 inhibition. *Oncotarget* **7**, 885–901 (2016).

839   18.   Quinton, R. J. *et al.* Whole-genome doubling confers unique genetic
840   vulnerabilities on tumour cells. *Nature* **590**, 492–497 (2021).

841   19.   Fujiwara, T. *et al.* Cytokinesis failure generating tetraploids promotes
842   tumorigenesis in p53-null cells. *Nature* **437**, 1043–1047 (2005).

843   20.   Orr-Weaver, T. L. When bigger is better: The role of polyploidy in
844   organogenesis. *Trends in Genetics* vol. 31 307–315 (2015).

845   21.   Zeman, M. K. & Cimprich, K. A. Causes and Consequences of Replication
846   Stress. *Nat. Cell Biol.* **16**, 2 (2014).

847   22.   Koundrioukoff, S. *et al.* Stepwise activation of the ATR signaling pathway upon
848   increasing replication stress impacts fragile site integrity. *PLoS Genet.* **9**,
849   e1003643 (2013).

850   23.   Panier, S. & Boulton, S. J. Double-strand break repair: 53BP1 comes into
851   focus. *Nat. Rev. Mol. Cell Biol.* **15**, 7–18 (2014).

852   24.   Zhao, B., Rothenberg, E., Ramsden, D. A. & Lieber, M. R. The molecular basis
853   and disease relevance of non-homologous DNA end joining. *Nat. Rev. Mol.*
854   *Cell Biol.* **21**, 765–781 (2020).

855   25.   A, B., T, L. & A, C. Quantitative live imaging of endogenous DNA replication in
856   mammalian cells. *PLoS One* **7**, (2012).

857   26.   Michalet, X. *et al.* Dynamic molecular combing: Stretching the whole human
858   genome for high- resolution studies. *Science (80-.).* **277**, 1518–1523 (1997).

859   27.   Zlotek-Zlotkiewicz, E., Monnier, S., Cappello, G., Berre, M. Le & Piel, M.
860   Optical volume and mass measurements show that mammalian cells swell
861   during mitosis. *J. Cell Biol.* **211**, 765 (2015).

862   28.   Tatsumi, Y., Ohta, S., Kimura, H., Tsurimoto, T. & Obuse, C. The ORC1 cycle
863   in human cells: I. Cell cycle-regulated oscillation of human ORC1. *J. Biol.*
864   *Chem.* **278**, 41528–41534 (2003).

29.   Remus, D. *et al.* Concerted Loading of Mcm2-7 Double Hexamers around DNA

during DNA Replication Origin Licensing. *Cell* **139**, 719–730 (2009).

30.   Moyer, S. E., Lewis, P. W. & Botchan, M. R. Isolation of the Cdc45/Mcm2-

7/GINS (CMG) complex, a candidate for the eukaryotic DNA replication fork

helicase. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 10236–10241 (2006).

31.   Bertoli, C., Skotheim, J. M. & De Bruin, R. A. M. Control of cell cycle

transcription during G1 and S phases. *Nature Reviews Molecular Cell Biology*

vol. 14 518–528 (2013).

32.   Pardee, A. B. G1 events and regulation of cell proliferation. *Science (80-.).*

246, 603–608 (1989).

33.   Cadart, C. *et al.* Size control in mammalian cells involves modulation of both

growth rate and cell cycle duration. *Nat. Commun.* **9**, (2018).

34.   DW, F. *et al.* Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991

and associated antitumor activity in human tumor xenografts. *Mol. Cancer*

Ther. **3**, 1427–1437 (2004).

35.   Tan, C. *et al.* Cell size homeostasis is maintained by CDK4-dependent

activation of p38 MAPK. *Dev. Cell* **56**, 1756-1769.e7 (2021).

36.   Charvin, G., Oikonomou, C., Siggia, E. D. & Cross, F. R. Origin of irreversibility

of cell cycle start in budding yeast. *PLoS Biol.* **8**, (2010).

37.   Cross, F. R., Buchler, N. E. & Skotheim, J. M. Evolution of networks and

sequences in eukaryotic cell cycle control. *Philosophical Transactions of the*

Royal Society B: Biological Sciences vol. 366 3532–3544 (2011).

38.   Zatulovskiy, E., Zhang, S., Berenson, D. F., Topacio, B. R. & Skotheim, J. M.

Cell growth dilutes the cell cycle inhibitor Rb to trigger cell division. *Science*

(80-.). **369**, 466–471 (2020).

39.   Nano, M. *et al.* Cell-Cycle Asynchrony Generates DNA Damage at Mitotic

Entry in Polyploid Cells. *Curr. Biol.* **29**, (2019).

40.   Frawley, L. E. & Orr-Weaver, T. L. Polyploidy. *Current Biology* vol. 25 R353–

R358 (2015).

41.   Tseng, A.-S. K. & Hariharan, I. K. *An Overexpression Screen in Drosophila for*

Genes That Restrict Growth or Cell-Cycle Progression in the Developing Eye.

(2002).

42.   Duronio, R. J., Brook, A., Dyson, N. & O'Farrell, P. H. E2F-induced S phase

requires cyclin E. *Genes Dev.* **10**, 2505–2513 (1996).

43.   Maya-Mendoza, A. *et al.* High speed of fork progression induces DNA

replication stress and genomic instability. *Nature* **559**, 279–284 (2018).

44.   Sedlackova, H. *et al.* Equilibrium between nascent and parental MCM proteins

protects replicating genomes. *Nature* **587**, 297–302 (2020).

45.   M, T. *et al.* Functional reprogramming of polyploidization in megakaryocytes.

Dev. Cell **32**, 155–167 (2015).

46.   Klusza, S. & Deng, W.-M. At the crossroads of differentiation and proliferation:

Precise control of cell-cycle changes by multiple signaling pathways in

Drosophila follicle cells. *Bioessays* **33**, 124 (2011).

47.   Viganó, C. *et al.* Quantitative proteomic and phosphoproteomic comparison of

human colon cancer DLD-1 cells differing in ploidy and chromosome stability.

Mol. Biol. Cell **29**, 1031–1047 (2018).

48.   Wangsa, D. *et al.* Near-tetraploid cancer cells show chromosome instability

triggered by replication stress and exhibit enhanced invasiveness. *FASEB J.*

32, 3502–3517 (2018).

49.   Hégarat, N. *et al.* Cyclin A triggers Mitosis either via the Greatwall kinase

pathway or Cyclin B. *EMBO J.* **39**, (2020).

50.   SB, K. *et al.* A quantitative FastFUCCI assay defines cell cycle dynamics at a

single-cell level. *J. Cell Sci.* **130**, 512–520 (2017).

51.   KS, Y., RH, K., M, L., R, G. & R, W. Single cell resolution in vivo imaging of

DNA damage following PARP inhibition. *Sci. Rep.* **5**, (2015).

52.   Karess, R. E. *et al.* The regulatory light chain of nonmuscle myosin is encoded

by spaghetti-squash, a gene required for cytokinesis in Drosophila. *Cell* **65**,

1177–1189 (1991).

53.   Gemble, S. *et al.* Centromere Dysfunction Compromises Mitotic Spindle Pole

Integrity. *Curr. Biol.* **29**, (2019).

54.   J, L., BO, P., K, H., J, B. & K, H. Deregulated expression of E2F family

members induces S-phase entry and overcomes p16INK4A-mediated growth

suppression. *Mol. Cell. Biol.* **16**, 1047–1057 (1996).

55.   Sakaue-Sawano, A. *et al.* Visualizing Spatiotemporal Dynamics of Multicellular

Cell-Cycle Progression. *Cell* **132**, 487–498 (2008).

56.   Aknoun, S. *et al.* Living cell dry mass measurement using quantitative phase

imaging with quadriwave lateral shearing interferometry: an accuracy and

sensitivity discussion. *J. Biomed. Opt.* **20**, 126009 (2015).

57.   Bon, P., Maucort, G., Wattellier, B. & Monneret, S. Quadriwave lateral shearing

934  interferometry for quantitative phase microscopy of living cells. *Opt. Express*

935  **17**, 13080 (2009).

936 58. van den Bos, H. *et al.* Quantification of aneuploidy in mammalian systems. in

937  *Methods in Molecular Biology* vol. 1896 159–190 (Humana Press Inc., 2019).

938 59. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.

939  *Nat. Methods* **9**, 357–359 (2012).

940 60. Jun, G., Wing, M. K., Abecasis, G. R. & Kang, H. M. An efficient and scalable

941  analysis framework for variant extraction and refinement from population-scale

942  DNA sequence data. *Genome Res.* **25**, 918–925 (2015).

943 61. Bakker, B. *et al.* Single-cell sequencing reveals karyotype heterogeneity in

944  murine and human malignancies. *Genome Biol.* **17**, (2016).

945 62. van den Bos, H. *et al.* Single-cell whole genome sequencing reveals no

946  evidence for common aneuploidy in normal and Alzheimer's disease neurons.

947  *Genome Biol.* **17**, 1–9 (2016).

Figure 1

Figure 1: High levels of DNA damage are generated in the first interphase following unscheduled WGD.

(A) Schematic representation of the generation of tetraploid cells. **(B)** Representative immunofluorescence images of diploid and tetraploid RPE-1 cells generated by inhibiting Eg5 and MPS1 (MS), or MKLP1 (CF), or JNK (EnR). DNA was visualized using DAPI (in blue), centrosomes were stained using anti-CEP192 antibodies (in white) and membranes were stained using anti-β-Catenin antibodies (in red). The white squares correspond to higher magnifications presented in the lower panel and showing the centrosomes (in yellow). **(C, E and G)** Representative immunofluorescence images showing DNA damage in diploid and tetraploid RPE-1 cells were generated by inhibiting Eg5 and MPS1 (MS), or MPS1 (CF), or JNK (EnR). DNA was visualized using DAPI (in blue), DNA damage was visualized using anti-γH2AX antibodies (in red). >100 interphase cells were analyzed from at least three independent experiments. **(D, F and H)** Graph showing the number of γH2AX foci per interphase cells in diploid and tetraploid RPE-1 cells. The percentage of interphase cells with more than 10 γH2AX foci in diploid and tetraploid RPE-1 cells were indicated under the graph. >100 interphase cells were analyzed from at least three independent experiments. The dotted lines indicate the nuclear area. D: Diploid. T: Tetraploid. MS: mitotic slippage. CF: cytokinesis failure. EnR: endoreplication.

Figure 2

Figure 2: DNA damage and genetic instability in tetraploid cells is generated during S-phase in a DNA replication-dependent manner.
(A) Upper panel - Representative immunofluorescence images showing DNA damage in RPE-1 tetraploid cells over time. DNA was visualized using DAPI (in blue), DNA damage is visualized using anti-γH2AX antibodies (in red). Lower panel - Percentage of RPE-1 FUCCI tetraploid cells in G1 (red) or in S-G2 (green) over time. **(B)** Graph representing the number of γH2AX foci per interphase cells in diploid (gray) and tetraploid (blue) RPE-1 cells over time. The percentage of interphase cells with more than 10 γH2AX foci in diploid and tetraploid RPE-1 cells is indicated under the graph. >100 interphase cells were analyzed from at least three independent experiments. **(C)** Percentage of RPE-1 FUCCI tetraploid cells in G1 (red) or in S-G2 (green) and representative immunofluorescence images showing DNA damage in tetraploid cells synchronized in G1 using 1μM palbociclib or released in S-phase with or without 400nM aphidicolin (APH). DNA was visualized using DAPI (in blue), DNA damage was visualized using anti-γH2AX antibodies (in red). **(D)** Graph showing the number of γH2AX foci per interphase cells in diploid (gray) and tetraploid (blue) RPE-1 cells synchronized in G1 using 1μM palbociclib or released in S-phase with or without 400nM aphidicolin (APH). The percentage of interphase cells with more than 10 γH2AX foci in diploid and tetraploid RPE-1 cells are indicated under the graph. >100 interphase cells were analyzed from at least three independent experiments. **(E)** Schematic workflow showing the method used in this study to process and analyze DNA replication by live imaging. **(F)** Stills of time lapse of diploid and tetraploid RPE-1 PCNA^{chromo} cells. Active replication sites are visualized using PCNA chromobodies (in cyan) and reconstructed using Imaris in 3D (in red). **(G)** Graph showing the total number of active replication sites during S-phase in diploid (gray) and tetraploid (blue) RPE-1 cells. >20 S-phase cells were analyzed from three independent experiments. **(H)** Graph showing the volume of active replication sites in μm^3 in diploid (gray) and tetraploid (blue) RPE-1 PCNA^{chromo} cells. At least 1000 active replication sites were analyzed. **(I)** Graph showing the mean number of active replication sites over time in diploid (gray line) and tetraploid (blue line) RPE-1 cells. >20 S-phase cells were analyzed from three independent experiments. For other representative examples, see Figure S5A. **(J)** Left panel - Graph representing the replication fork speed in diploid (gray) and tetraploid (yellow) HCT116 cells. Right panel - Graph showing the percentage of unstable replication forks in diploid (gray) and tetraploid (yellow) HCT116 cells. More than 120 replication forks were analyzed. Lower panel - Representative immunofluorescence of DNA fibers obtained from diploid and tetraploid HCT116 cells. ssDNA was visualized using anti-ssDNA antibodies (in blue), CldU and IdU was visualized using anti-IdU and anti-CldU antibodies (in red and green), respectively. **(K)** Genome-wide copy number plots G2/M tetraploid RPE-1 cells were generated using a modified version of the Aneufinder algorithm and normalized using G1 tetraploid cells (see methods). Each row represents a cell and the copy number state (in 5-Mb bins) is indicated in colors (with aberrations contrasting from from dark green in G2/M (8n). The dotted lines indicate nuclear area. D: Diploid. T: Tetraploid. MS: mitotic slippage. CF: cytokinesis failure. EnR: endoreplication.

Figure 3

Figure 3: Lack of G1 lengthening in tetraploid cells leads to unprepared S-phase.
(A) Stills of time lapse of RPE CCNB1^{AID} FUCCI diploid and tetraploid cells. G1 cells are in red and S-G2 cells are in green. **(B)** Graph representing the amount of protein produced during G1 in diploid (gray) and tetraploid (blue) RPE-1 CCNB1^{AID} FUCCI cells. **(C)** Schematic representation of cell sorting of G1 diploid and tetraploid cells. At least 50 cells were analyzed. **(D)** H2B, actin, ORC1, MCM2, PCNA and Cdc45 levels assessed by western blot of cell lysates obtained from diploid (left) and tetraploid (right) RPE-1 CCNB1^{AID} FUCCI cells. The same number of cells was loaded for each condition. **(E)** Graph showing H2B levels normalized with diploid condition (fold change) in diploid (gray) and tetraploid (blue) cells. Mean +/- sd representing three independent experiments. **(F)** Graph representing the protein levels relative to H2B levels (fold change) in diploid (gray) and tetraploid (blue) cells. Mean +/- sd representing three independent experiments. **(G)** Left panel – Stills of time lapse of RPE FUCCI diploid and tetraploid cells. G1 cells are in red and S-G2 cells are in green. Right panel - Graph showing the time in G1 compared to the mass at birth in diploid (left panel, gray) and tetraploid (right panel, blue) RPE-1 FUCCI cells. At least 55 interphase cells from two independent experiments were analyzed. **(H)** Graph showing the correlation between the time in G1 and the mass at birth in diploid and tetraploid RPE-1 CCNB1^{AID} FUCCI cells. At least 50 interphase cells were analyzed. **(I)** H2B, ORC1, MCM2, PCNA and Cdc45 levels assessed by western blot of cell lysates obtained from diploid (left) and tetraploid (right) RPE-1 CCNB1^{AID} FUCCI cells with extended G1 duration. The same number of cells was loaded for each condition. **(J)** Graph showing H2B levels normalized with diploid condition (fold change) in diploid (gray) and tetraploid (blue) cells with extended G1 duration. Mean +/- sd representing three independent experiments. **(K)** Graph representing the protein levels relative to H2B levels (fold change) in diploid (gray) and tetraploid (blue) cells with extended G1 duration. Mean +/- sd representing three independent experiments. **(L)** Graph showing the average number of active replication sites over time in diploid (gray line) or tetraploid (blue line) RPE PCNA^{chromo} cells with extended G1 duration. For other representative examples, see Figure S6L. **(M)** Graph showing the number of γH2AX foci in interphase cells in diploid (gray) and tetraploid cells (blue) synchronized in G1 using 160nM (extended G1 duration) or 1µM (G1 arrest) palbociclib or released in S-phase. At least 100 interphase cells were analyzed from three independent experiments. D: Diploid. T: Tetraploid. MS: mitotic slippage.

Figure 4

Figure 4: G1 lengthening or increased E2F1 levels are sufficient to rescue GIN in tetraploid cells and in polyploid cells *in vivo*.
(A) Upper panel - Schematic workflow showing the method used to overexpress E2F1. Lower panel - Representative immunofluorescence images showing DNA damage in RPE-1 tetraploid cells overexpressing or not E2F1. DNA was visualized using DAPI (in blue), DNA damage was visualized using anti-γH2AX antibodies (in red). **(B)** Graph showing the number of γH2AX foci per interphase cells in diploid (gray) and tetraploid (blue) RPE-1 cells released in S-phase with or without E2F1 overexpression. The percentage of interphase cells with more than 10 γH2AX foci in diploid and tetraploid RPE-1 cells were indicated under the graph. >100 interphase cells were analyzed from at least three independent experiments. **(C)** Schematic representation of the brain of *drosophila* larvae. **(D)** Representative immunofluorescence images of *drosophila* brain lobe in control or *sqh* mutant (polyploid) and of salivary glands. DNA was visualized using DAPI (in blue), DNA damage was visualized using anti-γH2Av antibodies (in red), membranes were visualized using Phalloidin (in yellow). **(E)** γH2Av index in *drosophila* salivary glands (SG, gray) or in diploid (gray) and polyploid (yellow) neuroblasts (NB). At least 60 cells were analyzed per condition. **(F)** Representative immunofluorescence images of *drosophila* brain lobe in control or *sqh* mutant (polyploid) overexpressing or not E2F1. DNA was visualized using DAPI (in blue), DNA damage was visualized using anti-γH2Av antibodies (in red), membranes were visualized using Phalloidin (in yellow). **(G)** Graph showing the γH2Av index in diploid (gray) and polyploid (yellow) *drosophila* neuroblasts overexpressing or not E2F1. At least 30 cells were analyzed per condition. **(H)** Tetraploid cells are not able to sense an increase in DNA content and to adapt G1 duration. In consequence, G1 duration is not scaled up with DNA content and tetraploid cells enter S-phase with an insufficient amount of replication factors generating DNA replication-dependent DNA damage and abnormal karyotypes. The dotted lines indicate the nuclear (B) or cell area (D and F). The white squares correspond to higher magnifications presented in the lower panel. D: Diploid. T: Tetraploid. P: Polyploid. CF: cytokinesis failure. MS: mitotic slippage.