Epigenetic regulator genes direct lineage switching in *MLL-AF4* leukaemia

Ricky Tirtakusuma[1], Katarzyna Szoltysek[1,2,3], Paul Milne[4], Vasily V Grinev[5], Anetta Ptasinska[6],

Claus Meyer[7], Sirintra Nakjang[1], Jayne Y Hehir-Kwa[2], Daniel Williamson[1], Pierre Cauchy[6],

Salam A Assi[6], Maria R Imperato[6], Fotini Vogiatzi[9], Shan Lin[10], Mark Wunderlich[10], Janine

Stutterheim[2], Alexander Komkov[8], Elena Zerkalenkova[8], Paul Evans[11], Hesta McNeill[1], Alex

Elder[1], Natalia Martinez-Soria[1], Sarah E Fordham[1], Yuzhe Shi[1], Lisa J Russell[1], Deepali Pal[1],

Alex Smith[12], Zoya Kingsbury[13], Jennifer Becq[13], Cornelia Eckert[14], Oskar A Haas[15], Peter

Carey[16], Simon Bailey[1,16], Roderick Skinner[1,16], Natalia Miakova[8], Matthew Collin[4], Venetia

Bigley[4], Muzlifa Haniffa[17,18,19], Rolf Marschalek[7], Christine J Harrison[1], Catherine A Cargo[11],

Denis Schewe[9], Yulia Olshanskaya[8], Michael J Thirman[20], Peter N Cockerill[6], James C

Mulloy[10], Helen J Blair[1], Josef Vormoor[1,2], James M Allan[1], Constanze Bonifer[6*], Olaf

Heidenreich[1,2*†], Simon Bomken[1,16*†]

Author Affiliations

[1]Wolfson Childhood Cancer Research Centre, Centre for Cancer, Translational and Clinical
Research Institute, Newcastle University, Newcastle upon Tyne, UK

[2]Princess Maxima Center for Pediatric Oncology, Utrecht, The Netherlands

[3]Maria Sklodowska-Curie Institute - Oncology Center, Gliwice Branch, Gliwice, Poland

[4]Translational and Clinical Research Institute, Newcastle University, Framlington Place,
Newcastle upon Tyne, UK

[5]Department of Genetics, the Faculty of Biology, Belarusian State University, 220030 Minsk,
Republic of Belarus.

[6]Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK

[7]Institute of Pharmaceutical Biology/DCAL, Goethe-University, Frankfurt/Main, Germany

[8]Dmitry Rogachev National Research Center of Pediatric Hematology, Oncology, and
Immunology; Moscow, Russia

[9]Pediatric Hematology/Oncology, ALL-BFM Study Group, Christian Albrechts University Kiel
and University Hospital Schleswig-Holstein, Campus Kiel, Germany

[10]Experimental Hematology and Cancer Biology, Cancer and Blood Disease Institute,
Cincinnati Children's Hospital Medical Center, Cincinnati, USA

[11]Haematological Malignancy Diagnostic Service, St James's University Hospital, Leeds, UK

[12]Epidemiology and Cancer Statistics Group, University of York, York, United Kingdom

[13]Illumina Cambridge Ltd., Great Abington, UK

[14]Department of Pediatric Oncology/Hematology, Charité Universitätsmedizin Berlin, Berlin,
Germany

[15]Children's Cancer Research Institute, St. Anna Kinderkrebsforschung, Vienna, Austria

[16]Department of Paediatric Haematology and Oncology, The Great North Children's Hospital,
Newcastle upon Tyne, UK

[17]Biosciences Institute, Newcastle University, Framlington Place, Newcastle upon Tyne, UK

[18]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton UK

[19]Department of Dermatology and Newcastle NIHR Newcastle Biomedical Research Centre,
Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne

[20]Department of Medicine, Section of Hematology/Oncology, University of Chicago, Chicago,
USA

*Co-senior authors
†**Co-corresponding Authors.** Correspondance to: **s.n.bomken@ncl.ac.uk;**
O.T.Heidenreich@prinsesmaximacentrum.nl
48

Summary

The fusion gene *MLL-AF4* defines a high-risk subtype of pro-B acute lymphoblastic leukaemia.

However, relapse can be associated with a switch from acute lymphoblastic to acute myeloid

leukaemia. Here we show that these myeloid relapses share oncogene fusion breakpoints

with their matched lymphoid presentations and can originate in either early, multipotent

progenitors or committed B-cell precursors. Lineage switching is linked to substantial changes

in chromatin accessibility and rewiring of transcriptional programmes indicating that the

execution and maintenance of lymphoid lineage differentiation is impaired. We show that this

subversion is recurrently associated with the dysregulation of repressive chromatin modifiers,

notably the nucleosome remodelling and deacetylation complex, NuRD. In addition to

mutations, we show differential expression or alternative splicing of NuRD members and other

genes is able to reprogram the B lymphoid into a myeloid gene regulatory network. Lineage

switching in *MLL-AF4* leukaemia is therefore driven and maintained by defunct epigenetic

regulation.

Introduction

Translocation of Mixed Lineage Leukaemia (*MLL*) with one of over 130 alternative partner

genes is a recurrent cytogenetic finding in both acute myeloid and lymphoblastic leukaemias

and is generally associated with poor prognosis (Pieters *et al.*, 2007; Moorman *et al.*, 2010;

Meyer *et al.*, 2018). Amongst the most common translocations is t(4;11)(q21;q23), forming the

MLL-AF4 (also known as *KMT2A-AFF1*) fusion gene. Uniquely amongst *MLL* rearrangements

(*MLL*r), *MLL-AF4* is almost exclusively associated with pro-B cell acute lymphoblastic

leukaemia and is prototypical of infant acute lymphoblastic leukaemia (ALL) where it carries a

very poor prognosis (Meyer *et al.*, 2018). However, despite this general lymphoid presentation,

MLL-AF4 leukaemias have an intriguing characteristic - that of lineage switched relapses.

Lineage switch acute leukaemias (LSALs) lose their lymphoid specific features and gain

myeloid phenotype upon relapse (Jiang *et al.*, 2005; Germano *et al.*, 2006; Park *et al.*, 2011;

Carulli *et al.*, 2012; Rossi *et al.*, 2012; Ivanov *et al.*, 2013). Alternatively, *MLL-AF4* leukaemias

may harbour distinct lymphoid and myeloid populations at the same time, thus classifying as

mixed phenotype acute leukaemias (MPALs) of the bilineage subtype.

In order to understand the molecular basis of lineage promiscuity and switching, we examined

a unique cohort of *MLL-AF4*-positive LSAL presentation/relapse pairs and MPALs. We

demonstrate that disruption of the epigenetic machinery, including the nucleosome

remodelling and deacetylation complex (NuRD), is associated with the loss of lymphoid

restriction. Lineage switch is then enacted through redistribution of transcription factor binding

and chromatin reorganisation. Whilst identified here within this rare clinical context, our

findings bare relevance for our understanding of the transforming capacity of *MLL-AF4*, and

how this oncoprotein imposes lineage determination on haematopoietic precursor cells.

Furthermore, given the high-risk nature of this disease, we provide a novel insight into factors

which may prove critical to the effective implementation of lineage specific, epitope-directed

therapies such as chimeric antigen receptor T-cell (CAR-T) cell or bi-specific T-cell engaging

antibody (BiTE) approaches.

Results

Characterisation of *MLL-AF4* acute leukaemias with lineage switch

To characterize lineage promiscuity in *MLL-AF4* leukaemia and the underlying molecular

mechanisms, we collected a cohort of ten cases of *MLL-AF4* ALL comprising 6 infant, 2

paediatric and 2 adult patients who had relapsed with acute myeloid leukaemia (AML).

Amongst these, one infant patient (LS10) had relapsed following B-lineage directed

blinatumomab treatment (Table S1). The time to relapse ranged from 3 to 48 months. Seven

patients within the cohort subsequently died. Lineage switch was defined as loss of expression

of B lymphoid antigens (CD19, CD22, CD79A) with concomitant gain of expression of myeloid

antigens (CD33, CD117/KIT, CD64/FCGR1A) and/or an unequivocal change in morphology

to AML (Figure 1A and Table S1). In addition, we studied two *MLL-AF4* infant mixed phenotype

acute leukaemias (MPALs), comprising distinct lymphoid and myeloid populations (MPAL1,

MPAL2; Table S1).

All matched samples displayed identical oncogene fusion breakpoints at diagnosis and

relapse as shown by DNA (n=14) and/or RNA (n=10) sequencing, confirming a common clonal

origin and proving that the relapses are not *de novo* or therapy-associated AMLs (Figures 1B,

C, S1, Tables S1). Breakpoints of LSALs and MPALs show a similar distribution as *MLL-AF4*

ALL cases, clustering in *MLL* introns 9-11 and AF4 introns 3 and 4 (Meyer *et al.*, 2009; Jung

et al., 2010) (Figure 1D, Table S1) thus excluding that distinct, "non-canonical" chromosomal

breakpoints are causative for *MLL-AF4*-positive AML. These data raised the question of the

cellular origin of relapse and the nature of events secondary to *MLL-AF4* that affect lineage

commitment.

Cellular origin of lineage switched relapse

We hypothesised that myeloid leukaemias may not have undergone substantial B-cell

receptor (BCR) rearrangements. We used this feature to further interrogate the developmental

stage at which the relapse arose. To that end we examined BCR rearrangements within RNA-

seq and whole exome-seq (WES) derived data, using MiXCR software (Bolotin *et al.*, 2017).

All ALL cases showed classical oligoclonal rearrangements of BCR loci, supporting the

lymphoid lineage decision (Figure S2, Table S2). However, we observed three distinct patterns

for AML relapses (Figure 2A). Pattern 1 comprises AML relapse cells with no BCR

rearrangements implying presence of a relapse-initiating cell residing in a primitive precursor

population prior to early DJ recombination (Figure 2A, cases LS01, LS02, LS04). As a second

pattern, we found unrelated BCR rearrangements, which may indicate either aberrant

rearrangement in a myeloid cell or relapse initiating from either a B-lymphoid cell committed

to undergo rearrangement, or a transdifferentiated minor ALL clone with an alternative

rearrangement (Figure 2A, cases LS03, LS06, LS07, LS08, MPAL2). Interestingly, this pattern

is also found in a relapse after blinatumomab treatment (LS10). Pattern 3 comprises shared

BCR rearrangements between diagnostic and relapse material, which suggest a direct

transdifferentiated myeloid relapse from the ALL (Figure 2A, cases LS05, LS09, MPAL1).

These data demonstrate that AML relapses can originate from different stages of lymphoid

leukaemogenesis and suggest, at least for a subset, a common precursor preceding the pro-

B cell stage.

To functionally assess the plasticity of immunophenotypically defined diagnostic ALL and

relapsed AML, we transplanted NSG or MISTRG mice with either ALL or AML cells from

patient LS01 (Figure 2B). Because of the expression of several human myeloid growth factors,

the MISTRG strain more strongly supports AML engraftment than the NSG strain, which

shows a stronger lymphoid bias. Diagnostic ALL transplants rapidly produced representative

CD19+CD33- lymphoid leukaemias in both mouse strains. In contrast, transplantation of

relapsed AML engrafted only in the more myeloid-permissive MISTRG strain as a CD34+/-

CD19-CD33+ AML (Figure 2C). Thus, in contrast to the immunophenotypic plasticity seen in

MPAL leukaemias with wildtype *MLL* (Alexander *et al.*, 2018), transplantation of a relapsed,

fully switched AML was only capable of generating AML.

143   Lineage switch relapse can originate in HSPC compartments

To further investigate a potential origin of relapse in an early progenitor or stem cell, we purified

haematopoietic stem/progenitor cell (HSPC) populations from diagnostic ALL and relapsed

AML and tested purified populations for the presence of *MLL-AF4* targeted sequencing. The

MLL-AF4 translocation was found in the lymphoid-primed multipotent progenitor-like

population (LMPP, CD34+CD38-CD45RA+; lymphoid and myeloid potential, but not

megakaryocyte-erythroid potential), in the multipotential progenitor population (MPP,

CD34+CD38-CD45RA-CD90-; no lineage restriction) and for MPAL1 even in the

haematopoietic stem cell-like population (HSC, CD34+CD38-Lin-CD90+) (Figures 3A-C, S3;

Table S3). In line with these findings, serial xenotransplantation of LS01P identified a

persistent human CD34+CD38-CD45RA-CD90+ HSC compartment across four generations

of mice with maintenance of the *MLL-AF4* fusion gene within purified human CD34+ cells

(Figures 3D-F). These findings suggest maintenance of a (pre-)leukaemic clone with high

malignant self-renewal potential in HSPC populations and support our findings from BCR

analysis that, at least in a subgroup of cases, an early multipotent progenitor or HSC can act

as the cell of origin for the AML relapse.

In concordance with the translocation being present within the early HSPC compartment, we

sorted viable differentiated leukocytes and were able to detect the *MLL-AF4* fusion in myeloid

and lymphoid lineages including CD34-CD19/3-HLA-DR+CD14/11c+ monocytes, NK, B and

mature T cells (Figure 3C, S3A, B; Table S3). These findings imply the existence of a pre-

leukaemic progenitor cell, in which *MLL-AF4* does not impose a complete block on

haematopoietic differentiation but is compatible with myeloid and lymphoid differentiation.

These findings raise the question of which factors and molecular mechanisms affect the ALL

and AML lineage choice in *MLL-AF4* leukaemia.

Lineage switch leukaemia is associated with transcriptional reprogramming

We next investigated the underlying molecular events associated with a lineage switch. To this end we analysed differential gene expression across eight cases, including six LSALs for which RNA was available at both presentation and relapse, and sorted lymphoid and myeloid blast populations of two MPALs. This analysis identified in total 1374 up- (adj. p<0.01, Log Fold change >2) and 1323 down-regulated genes in the AML lineage switches and the myeloid populations of MPAL patients (Table S4). The most substantially down-regulated genes include lymphoid genes such as *PAX5* and *RAG2*, but also the polycomb PRC1 like complex component *AUTS2* and SWI/SNF complex component *BCL7A,* while up-regulated genes comprise several myeloid genes such as cathepsins, cystatins, *PRAM1* and *CSF3R* (Figure 4A). These fingings were consistent irrespective of which cellular origin of relapse the BCR rearrangement analysis supported (pattern 1, 2 or 3),

Both gene set enrichment analysis (GSEA) and non-negative matrix factorisation (NMF) showed that presentation and relapse cases with lineage switch have expression signatures similar to previously published *MLL-AF4* ALL and *MLL*r AML cases as well as normal lymphoid and myeloid cell types, respectively (Figures 4B, S4A, B) (Zangrando *et al.*, 2009; Novershtern *et al.*, 2011; Andersson *et al.*, 2015). More specifically, lineage switch included increased expression of factors controlling myeloid differentiation (e.g., *CSF3R, KIT)* and changes in haematopoietic surface marker expression (e.g., *CD19, CD22*, *CD33, CD14, FCGR1A/CD64*), loss of immunoglobulin recombination machinery genes (e.g., *RAG1*, *RAG2, DNTT*) and reduced expression of genes encoding heavy and light immunoglobulin chains (Figures 4C-E, S4C). Notably, GSEA also indicated impaired DNA repair and cell cycle progression of the AML relapse when compared with diagnostic ALL (Figures S4D). In particular, the reduced self-renewal potential might reflect the transition from ALL with a high incidence of leukaemic stem cells (LSCs) to AML with fewer LSCs.

MLL fusion proteins including MLL-AF4 have previously been shown to directly regulate multiple genes linked to haematopoietic and leukaemic stemness (Ayton and Cleary, 2003;

Somervaille *et al.*, 2009; Gessner *et al.*, 2010; Wilkinson *et al.*, 2013; Kerry *et al.*, 2017). For

instance, significant changes in gene expression across the *HOXA* cluster, notably with a 50-

fold reduction in *HOXA7* expression, represent a major additional disruption of *MLL*r

leukaemogenic transcriptional regulation (Figures S5A-C) (Ayton and Cleary, 2003;

Somervaille *et al.*, 2009; Gessner *et al.*, 2010). Furthermore, a very significant portion of target

genes including *PROM1*, *IKZF2* and chromatin modifying factors are highly enriched amongst

genes that show lower expression in myeloid lineages (Figures S5D, E, Table S4). In total,

996 out of 5208 bona fide direct target genes of *MLL-AF4* changed expression in the AML

relapse (Wilkinson *et al.*, 2013). These data suggest that the process of lineage switch is

associated with a major reorganisation of the *MLL-AF4* transcriptional network and pose the

question of which epigenetic regulators are involved in the lineage determination associated

with this fusion gene.

Reorganisation of chromatin accessibility and transcription factor binding upon

lineage switch

For case LS01 we had sufficient diagnostic material to perform DNase hypersensitivity site

(DHS) analysis and thus link transcriptional changes to altered genome-wide chromatin

accessibility. Many differentially expressed genes showed altered chromatin accessibility in

proximity to the transcriptional start site (TSS), including genes encoding key hematopoietic

surface markers CD33 and CD19, transcription factors and proteins related to differentiation

(Figures 5A-C, S6A).

Changes in chromatin accessibility were linked with an altered pattern of transcription factor

binding. High resolution DHS-seq (digital footprinting) of presentation ALL and relapse AML,

respectively, showed marked genome-wide alterations at sites distal to the TSS, from which

several AML and ALL specific *de novo* occupied transcription factor binding motifs were

identified (Figures 5C-E, S6B). Lineage switch from ALL to AML was associated with a loss of

occupancy of motifs binding lymphoid transcription factors such as EBF or PAX5 and a gain

of occupancy of motifs bound by C/EBP, IRF and NF-κB family members (Figure 5E-F). The

gain in C/EBP motif binding was associated with the expression of a C/EBPA regulated

myeloid transcriptional gene set (Figure 5G). We also observed a redistribution of occupancy

of transcription factors controlling both lymphoid and myeloid maturation such as RUNX, AP-

1 and ETS members to alternative cognate motifs (Figures 5E and S6B) (Hohaus *et al.*, 1995;

Nerlov and Graf, 1998; Leddin *et al.*, 2011). This finding is exemplified by decreased

accessibility of a region located 1 kb upstream of the *CD19* TSS with concomitant loss of EBF

binding at this element (Figure 5C). Differential motif enrichments were associated with

changes in RNA expression and chromatin accessibility at the genes encoding the

corresponding cognate transcription factors *EBF1*, *PAX5*, *LEF1* (B lymphoid determinants),

NFKB2 and *CEBPB/D/E* (myeloid determinants), particularly in regions flanking TSSs (Figures

5H, S6B).

In conclusion, the transition from lymphoid to myeloid immunophenotype is associated with

global lineage specific transcriptional reprogramming and genome-wide alteration in

chromatin accessibility and transcription factor binding.

Lineage switch changes alternative splicing patterns

Lineage fidelity and determination are not only linked to differential gene expression but may

also include co- or post-transcriptional mechanisms. It has been previously demonstrated that

lineage commitment during haematopoiesis leads to substantial changes in alternative mRNA

splicing patterns (Chen *et al.*, 2014). Furthermore, we recently showed that the *AML-ETO*

(RUNX1-RUNX1T1) fusion protein controls leukaemic self-renewal by both differential gene

transcription and alternative splicing (Grinev *et al.*, 2021). To complement the transcriptional

analysis we therefore sought to define the different composition of RNA isoforms in lymphoid

and myeloid populations from lineage switch and MPAL cases. Here we focussed on three

lineage switch patients and the two MPAL patients whose RNA-seq data provided sufficient

read depth for the analysis of exon-exon junctions, exon usage and intron retention (Figures

S7A-C). We detected in total 2630 retained introns (RIs) shared amongst the three lineage

switches with 653 and 343 RIs exclusively found in the diagnostic ALL or the AML relapses,

respectively. This was complemented by 97 exons (DEUs) and 193 exon-exon junctions

(DEEjs) differentially used between diagnostic ALL or the AML relapses. In contrast, this

analysis identified only 43 RIs present in both MPAL cases with 18 and 6 introns specifically

retained in either the lymphoid or myeloid subpopulation, respectively (Figure S7D, Table S5).

Intersection of the affected genes identified 21 shared genes out of 166 DEEjs and 74 DEUs.

MPALs had 420 DEUs and 155 DEEjs affecting 257 and 103 genes with 30 genes having both

DEUs and DEEjs (Figure 6A, Table S6). While more than 80% of the non-differential exon-

exon linkages were canonical, this was true for only 15% of the DEEjs. Here, non-canonical

exon skipping and complex splicing events contributed more than 30% each, most prominently

to differential alternative splicing (Figure 6B).

Pathway analysis revealed an enrichment of alternatively spliced genes in immune pathways

including antigen processing, membrane trafficking and FCGR-dependent phagocytosis

reflecting the change from a lymphoid to a myeloid state (Figures 6C, S7E). Furthermore, it

highlighted RNA processing and maturation including mRNA splicing, processing of capped

intron-containing pre-mRNAs and rRNA processing. Indeed, myeloid populations expressed

4-6-fold higher alternatively spliced *SF3B1* and *SRSF5* levels than their matched lymphoid

populations (Figure 6D, E). In addition, we noted a significant number of genes encoding

epigenetic modulators including *KDM5C, HDAC2* and several *CHD* members being

differentially spliced in AML relapse or myeloid subpopulations of MPALs (Figure 6D, E).

Analysing the occurrence of *MLL-AF4* isoforms showed that the fusion site of *MLL-AF4* itself

was subject to alternative splicing in lineage switch and MPAL. Three cases shared the

breakpoints in introns 10 and 4 of *MLL* and *AF4*, respectively (Figure 6F). By examining RNA-

seq and competitive RT-PCR data, we identified two co-occurring isoforms with either *MLL*

exon9 or exon10 joined to *AF4* exon4 (Figure 6F). Interestingly, lymphoid populations

expressed a higher ratio of exon10/exon4 over exon9/exon4 than the myeloid populations in

all three cases examined. Furthermore, the HSC and MPP-like populations of MPAL1 showed

mainly expression of the exon10/exon4 splice variant of *MLL-AF4*, thus resembling more the

lymphoid than the myeloid phenotype (Figure 6F). In conclusion, altered isoform expression

of *MLL-AF4* may contribute to lineage choice and the phenotypic switch.

The mutational landscape of lineage switch

Next, we examined the mutational landscape of lineage switched *MLL-AF4* leukaemias by

performing exome sequencing on the entire cohort. In keeping with published data on newly

presenting *MLL*r acute leukaemias (Dobbins *et al.*, 2013; Andersson *et al.*, 2015), exome

sequencing of presentation ALL samples confirmed a relatively quiet mutational landscape in

infant ALL cases, with median of 13 nonsynonymous somatic single nucleotide variants

(SNVs) or insertions/deletions (indels) predicted to be deleterious to protein function (Table

S7). Many of these were present in less than 30% of reads and considered sub-clonal. The

most commonly mutated genes at presentation were *NRAS* (3 cases) and *KRAS* (5 cases)

(Figure 7A) as described previously (Andersson *et al.,* 2015). In contrast, relapse AML

samples contained a median of 46 deleterious somatic SNVs and indels (Table S7). This

increase can be mainly attributed to three samples (LS03RAML, LS07RAML and LS08RAML)

that carried deleterious mutations in DNA polymerase genes in the respective major clones

linked to hypermutator phenotypes (Church *et al.*, 2013; Erson-Omay *et al.*, 2015; Mur *et al.*,

2020) (Figure 7A). However, we observed this phenotype only in three out of ten relapses,

arguing against this phenomenon being a general requirement for the lineage switch in

relapse. Similarly, many of the predominantly subclonal mutations identified in presentation

ALL samples, including half of *RAS* mutations, were subsequently lost at relapse, indicating

alternative subclones as the origin of relapse (Figure 7A). Finally, both MPALs harboured

several mutations that were exclusively found in either the lymphoid or myeloid subpopulation

indicating the presence of subclones with a lymphoid and myeloid bias (Figure 7A).

Next, we examined the global mutation patterns of patient LS01 in greater detail and focussed

on SNVs with a VAF of ≥0.3. ALL and AML contained 104 and 3196 SNVs above this

threshold, respectively, with only 22 shared SNVs between both samples. The most prevalent

type of SNV was the C to T transition in the DNA of both ALL and AML samples (Figure 7B).

However, the contribution of underlying single base substitution (SBS) signatures differed

between diagnosis and relapse. Three different signatures (SBS16, 5 and 1) explained about

60% of the SNPs found in ALL, while SBS1 seemed to explain more than 50% of all SNPs in

the AML (Figure 7B) suggesting a mutational clock as the main driver of the evolution of

relapse (Alexandrov et al., 2015; Blokzijl et al., 2016). Further inspection of the pattern

revealed a mutational signature mainly comprising C to T transitions and to a lesser degree C

to G transversions in NCG triplets, raising the possibility that thiopurine maintenance treatment

may have increased the mutational burden, resulting in lineage switched relapse in this patient

(Li *et al.*, 2020).

Twelve deleterious SNVs were identified as unique to the relapse sample of case LS01. The

availability of viable cellular material allowed us to investigate the order of acquisition of these

secondary mutations within the structure of the normal haematopoietic hierarchy. We sorted

this sample to isolate HSC-, MPP-, LMPP- and GMP-like and later populations. Using a

targeted deep sequencing approach we then examined these populations for the presence of

those 12 SNVs. This analysis showed an increasing number of mutations during the

differentiation from MPPs through LMPPs to GMPs. Amongst them, only *PHF3* and *CHD4*

mutations were present within the purified CD34+CD38-CD45RA-CD90- MPP-like fraction

with VAF≥0.3 (Figure 7C and Table S3). In contrast, LMPP- and GMP-like populations

contained all 12 SNVs at high VAF (Table S3). These findings identify the mutation of *CHD4*

and *PHF3* as the earliest genetic events during relapse evolution and suggest them as

potential drivers of an *MLL-AF4* positive, non-lineage committed, pre-leukaemic precursor

population. Subsequent accumulation of additional mutations likely establishes the fully

developed leukaemia with more mature haematopoietic/myeloid immunophenotypes.

Perturbation of *CHD4* and *PHF3* disrupts lymphoid development in *MLL-AF4*

expressing cells

The two earliest mutations within LS01 relapse were identified in the Nucleosome Remodelling

and Deacetylation complex (NuRD) member *CHD4* and the plant homeodomain finger

containing *PHF3*. PHF3 is a member of a family of transcriptional regulators that have been

suggested to link the deposition of histone marks to the regulation of transcription (Kinkelin *et*

al., 2013). PHF3 itself has been recently identified as an inhibitor of transcription elongation

by competing with TFIIS for binding to the C-terminal domain of RNA polymerase II (Appel *et*

al., 2020). NuRD is a multiprotein transcriptional co-repressor complex with both histone

deacetylase and ATP-dependent chromatin remodelling activity. It is a critical factor in the

lymphoid lineage determination in part directed by the transcription factor IKZF1 (Kim *et al.*,

1999; Yoshida *et al.*, 2006; Ng *et al.*, 2009; Zhang *et al.*, 2011; Arends *et al.*, 2019). Both the

CHD4 R1068H and PHF3 K1119I mutations affect highly conserved residues (Figures 8A,

S8A) that are predicted by the Condel classifier (González-Pérez and López-Bigas, 2011) to

disrupt protein function. Specifically, the CHD4 R1068H mutation has previously been linked

to defects in cardiac development (Sifrim *et al.*, 2016).

Whilst mRNA expression levels of the mutant CHD4 R1068H were unaffected in case LS01,

across the remaining cases analysed, expression of *CHD4* and additional NuRD complex

members *MBD3*, *MTA1* and *RBBP4* was reduced following lineage switched myeloid relapse

(Figures 8B). Furthermore, *CHD4* was affected in the MPAL myeloid populations by non-

canonical alternative splicing leading to premature termination of translation, indicating that

this particular pathway was severely disrupted across the cohort, irrespective of the putative

cell of origin of relapse.

The relevance of *CHD4* and *PHF3* in regulating lymphoid versus myeloid lineage choice was

further supported by ARACNE analysis using published ALL (Harvey *et al.*, 2010; Kang *et al.*,

2010) and AML (Gentles *et al.*, 2010) expression datasets (n=216) which we used to reverse-

engineer a mutual information network (Margolin *et al.*, 2006). This network was trimmed to

represent only genes significantly associated with the difference between AML and ALL thus

reflecting the likely influence of genes of interest upon genes associated with the difference

between AML and ALL. This analysis found *CHD4* and *PHF3* to be the mutated genes with

the highest number of edges within the network (*PHF3* – 21 edges, p=0.010; *CHD4* – 12

edges, p=0.0005, Figure S8B and Table S8), implicating them as causal to the

lymphoid/myeloid distinction identified within primary ALL/AML.

To establish a direct functional link from these mutations to the loss of lymphoid lineage

commitment in *MLL-AF4* ALL, we knocked down *CHD4* and *PHF3* in the *MLL-AF4* positive

cell line, SEM. Depletion of *CHD4* and *PHF3* alone or in combination resulted in increased

expression of the myeloid transcription factor *CEBPA* and reduced expression of lymphoid

transcription factors including *LEF1*, *PAX5, TCF3* and *TCF12* (Figure 8C). These changes

were accompanied by a more than twofold increase in CD33 expression in the two *MLL-AF4-*

expressing ALL cell lines SEM and RS4;11, while CD33 levels in two *MLL-AF4*-negative ALL

cell lines remained unaffected, supporting the importance of these epigenetic regulators

specifically within the *MLL-AF4* context (Figures 8D and S8C). GSEA following *PHF3* and

CHD4 knockdown indicated loss of HSC and B-lymphocyte progenitor gene expression

signatures (Figure 8E, left panel), similar to what was observed with the transcriptomes of the

lineage switch leukaemia cases (Figure 8E, right panel). In line with these cell line

experiments, knockdown of *CHD4* and *PHF3* in an ALL PDX generated from the first relapse

of patient LS03 also resulted in a more than twofold increase in CD19/CD33 double and CD33

single positive cells, demonstrating that perturbation of these two genes is also able to change

the immunophenotype of primary *MLL-AF4* ALL cells in a case susceptible to trans-

differentiated relapse (Figure S8D).

In order to examine the role of additional mutations of chromatin modifiers found in our cohort

and known to regulate lineage choice, we investigated the impact of the PRC1 members

PCGF6 and *AUTS2* on CD33 expression in SEM cells. *PCGF6* is mutated in LS07RAML and

LS08RAML and has known roles in B lymphoid malignancy (Ferreira *et al.*, 2008; Wang *et al.*,

2015). *AUTS2* is both mutated in LS08RAML and highly expressed in all lymphoid populations

examined (Figures 4A, 7A). STRING network (Szklarczyk *et al.*, 2019) analysis demonstrated

close functional associations within PRC1 and NuRD complexes and their shared associations

(Figure S8E). Mutual interactions include CBX2, a PRC1 complex member, which shows a

similar expression pattern between lineage switch and MPAL cases (Figure S8F). While

knockdown of *AUTS2* did not change CD33 levels, depletion of *PCGF6* also increased CD33

surface expression in SEM cells, further supporting the notion of epigenetic factors regulating

lineage determination in ALL (Figures S8G). Furthermore, GSEA also indicated impaired

function of PRC1 and PRC2 complexes in the AML relapse compared with the presentation

ALL with down-regulation of their respective target genes (Figure S8H).

Given that the relapse-initiating cell can arise within an uncommitted, *MLL-AF4* translocated

HSPC population, we went on to assess the impact of *CHD4* and *PHF3* function loss in a

human cord blood model, which harbours a chimeric *MLL-Af4* fusion and can be differentiated

both into myeloid and lymphoid lineages (Lin *et al.*, 2016). Knockdown of either *CHD4* or *PHF3*

under lymphoid culture conditions significantly impaired lymphoid differentiation potential,

whilst co-knockdown of *CHD4* and *PHF3* disrupted differentiation entirely (Figures 8F, G and

Table S9). Transcriptomic analysis of the sorted populations revealed that CD33 positive cells

exhibited metagene expression pattern similar to *MLL*r AML, while the pattern describing

CD19+ cells was most similar to *MLLr* ALL, thus confirming that changes in surface marker

expression were associated with the corresponding changes in the transcriptomic profiles

(Figure S8I).

Taken together, our data show the important role of the NuRD (CHD4), PHF3 and other

(PRC1*)* repressive complexes in the epigenetic control of lymphoid lineage choice. In

particular, dysregulation of *CHD4*/NuRD was mediated by mutation, down-regulation of

expression and differential splicing across the cohort, irrespective of the cellular/clonal origin

of relapse. These data support a role for these factors in the strong lineage determining

capacity of *MLL-AF4* whilst their loss undermines both the execution and the maintenance of

the lymphoid lineage fate.

Discussion

This study describes impaired epigenetic control as being central to the phenomenon of

lymphoid-myeloid lineage switch in *MLL-AF4*-positive leukaemia and identifies the cell of

origin of relapse into AML. We found that the origin of relapse was heterogeneous. Relapse

can directly evolve from pro-B-like ALL blast populations, which agrees with the general self-

renewal capacity of ALL cells (le Viseur *et al.*, 2008; Rehe *et al.*, 2013), but can also originate

within the HSPC compartment. Indeed, analysis of both patient and xenotransplanted cell

populations from diagnostic ALL identified *MLL-AF4* fusion transcripts in MPP- and HSC-like

cells. This finding agrees with recently published data pointing at MPP cells as the origin of

MLL-AF4 leukaemia (Malouf and Ottersbach, 2018) and is in line with transcriptomic

similarities between t(4;11) ALL and Lin-CD34+CD38-CD19- fetal liver cells, again suggesting

an HSPC as the cell of origin (Agraz-Doblas *et al.*, 2019).

Irrespective of the cellular origin of the relapse, lineage switching was associated with a major

rewiring of gene regulatory networks. At the level of transcriptional control, the decision for

lymphoid development relies not only on the activation of a lymphoid transcriptional program,

but also on the silencing of a default myeloid program (Nutt *et al.*, 1997, 1999). That decision

is enacted by lymphoid master regulators including EBF1, PAX5 and IKAROS, which

represent genes commonly mutated in precursor B-ALL (Mullighan *et al.*, 2007; Pongubala *et*

al., 2008; Boer *et al.*, 2016; Witkowski *et al.*, 2017). Pax5$^{-/-}$ pro-B cells which lack lymphoid

potential, whilst capable of differentiating down erythro-myeloid lineages *in vitro*, still maintain

expression of early B cell transcription factors *EBF1* and *E2A* (*TCF3*) (Nutt *et al.*, 1999). In

contrast, we show that lineage switching *MLL-AF4* pro-B leukaemic relapse is associated with

significant reduction in expression and binding of these earliest B lymphoid transcription

factors. Their loss is linked with changes in the *MLL-AF4* transcriptional programme, notably

within *HOXA* cluster genes (Ayton and Cleary, 2003; Somervaille *et al.*, 2009; Gessner *et al.*,

2010)., which likely results in a wider reorganisation of malignant haematopoietic

transcriptional networks, ultimately leading to a myeloid differentiation fate.

Similar to the *Pax5* knockout (Nutt *et al.*, 1999), loss of IKAROS DNA-binding activity prevents

lymphoid differentiation (Zhang *et al.*, 2011). NuRD co-operates directly with IKAROS to

repress HSC self-renewal and subsequent myeloid differentiation, permitting early lymphoid

development (O'Neill *et al.*, 2000; Williams *et al.*, 2004; Yoshida *et al.*, 2008; Zhang *et al.*,

2011; Hosokawa *et al.*, 2013; Lu *et al.*, 2019). We found that the abrogation of this pathway

through multiple mechanisms was central to the lineage switch from ALL to AML. Lineage

switch was either associated with mutation, reduced expression or, in the case of two MPALs,

alternative splicing of *CHD4* and other NuRD components. Long term knockdown of *CHD4*

was not tolerated in our cord blood culture. This is in line with reports showing that complete

loss of *CHD4* impairs leukaemic proliferation (Sperlazza *et al.*, 2015; Heshmati *et al.*, 2018),

both myeloid and lymphoid differentiation of HSPCs and causes exhaustion of HSC pools

(Yoshida *et al.*, 2008), indicating that basal CHD4 expression is required for maintaining AML.

Moreover, our observation that a 60% *CHD4* knockdown is associated with the activation of

pluripotency gene signatures is in line with the finding that a partial inhibition of *CHD4*

supported induction of pluripotency in iPSCs, while a complete deletion eliminated cell

proliferation (Mor *et al.*, 2018).

Whilst our study has investigated the rare clinical occurrence of lineage switching, recent

studies have identified core NuRD and PRC1 complex members as being direct targets of

MLL-AF4 binding (Kerry *et al.*, 2017; Harman *et al.*, 2021). We therefore hypothesise that

epigenetic regulator genes are co-opted during *MLL-AF4* leukemogenesis and mediate

fundamental lineage specific decision-making processes, in this case the suppression of the

myeloid lineage program. Multiple routes to their dysregulation may result in escape from this

lineage restriction. Our finding that frontline chemotherapy itself may contribute to relapse

highlights the urgent need to find alternative therapies for this high-risk leukaemia. Equally,

however, the associated loss of B cell surface markers (*e.g.*, CD19) provides an alternative

mechanism for relapse following CAR-T cell or blinatumomab therapy (Gardner *et al*., 2016;

Rayes, McMasters and O'Brien, 2016) in addition to mutations, alternative splicing (Sotillo *et*

al., 2015; Orlando *et al.*, 2018; Rabilloud *et al.*, 2021) and T cells trogocytosis (Hamieh *et al.*,

2019). Whilst these therapies target lineage specific surface markers, lineage-switched (pre-

)leukaemic progenitor populations escape epitope recognition and provide a potential clonal

source for the relapse. Given the increasing use of advanced immunological therapies, a

detailed understanding of the molecular processes underlying lineage determination and

switching will be critical for developing new strategies to avoid this route to clinical relapse.

Acknowledgements

We thank Jon Coxhead and Raf Hussain at the Newcastle University core genomics facility

as well as Marc van Tuil at the Princess Maxima Center Diagnostic department for

development of sequencing strategies. We acknowledge the Newcastle University Flow

Cytometry Core Facility and Tomasz Poplonski at the Princess Maxima Center Flow

Cytometry core facility for their assistance with the generation of flow cytometry data and cell

sorting strategies as well as the Newcastle University Bioinformatics Support Unit for helping

to develop the analysis approach for sequencing data. We thank Ruben van Boxtel and Eline

Bertrums at the Princess Maxima Center for the generation of single cell derived clones,

derived from early progenitors sorted from MPAL patient samples. We thank Monique den

Boer, Frank van Leeuwen and Ronald Stam for critically reading the manuscript. This study

was supported by a Cancer Research UK Centre Studentship (C27826/A17312) and

Newcastle University Overseas Research Scholarship to RT, a CRUK program grant to JV

and OH (C27943/A12788), grants from the North of England Children's Cancer Research

Fund to OH, JV and SB, by Bloodwise grants 12055 and 15005 to OH and by a grant from the

Kay Kendall Leukaemia Fund (KKL1142) to OH. SB was supported by an NIHR Academic

Clinical Lectureship (CL-2012-01-002) and the Sir Bobby Robson Foundation Clinical

Fellowship. Work in CB/PNC's lab was funded by a programme grant from Bloodwise (15001).

Work in JMA's lab was funded by a programme grant from Bloodwise (13044). EZ was

supported by an RFBR grant (№17-29-06052). Research in the VVG laboratory was

supported in part by the Ministry of Education of the Republic of Belarus, grant #3.04.3.

Research in the AK laboratory was supported by an RSF grant (20-75-10091).

This study makes use of data generated by the St. Jude Children's Research Hospital –

Washington University Pediatric Cancer Genome Project and the Therapeutically Applicable

Research to Generate Effective Treatments (TARGET) initiative, phs000218, managed by the

NCI (see supplementary methods).

Author contributions

Conceptualization, O.H., S.B., C.B.; Methodology, O.H., C.B., R.T., K.S., P.M., S.B., A.P.,

C.M., A.K., Z.K., J.B., V.B., R.M., J.V., J.M.A., S.L.; Software Programming, S.N., J.H.K.,

V.V.G., A.K., D.W., P.C.; Formal Analysis, S.N., J.H.K., V.V.G., A.K., D.W., P.C., C.B., O.H.;

Investigation, R.T., K.S., P.M., A.P., C.M., H.J.B., A.K., S.A., M.R.I., P.E., H.M., A.E., N.M.S.,

S.E.F., Y.S., D.P., P.C.; Resources, F.V., E.Z., A.S., J.C.M., L.J.R., C.E., O.A.H., S.Ba, R.S.,

N.M., M.C., V.B., R.M., M.W., C.J.H., C.A.C., D.S., Y.O., M.J.T., P.N.C., J.C.M., C.B., O.H.;

Data Curation, S.N., D.W., P.C.; Writing, S.B., O.H., C.B., R.T., K.S.; Supervision, O.H., S.B.,

J.M.A., J.V., C.B., ; Funding Acquisition, O.H., J.V., S.B., C.B., P.N.C., J.M.A., E.Z.

Declaration of interests

Z.K. and J.B. are employees of Illumina, a public company that develops and markets systems

for genetic analysis. The remaining authors declare no competing interests.

Figure titles and legends

Figure 1. Characterisation of the *MLL-AF4* lineage switch cases. (A) Morphological

change from lymphoblastic leukaemia (left panel) to acute monoblastic/monocytic leukaemia

(right panel). The scale bar represents 20 µm. (B) Sanger sequencing of *MLL-AF4* and

reciprocal *AF4/MLL* fusions in LS01 presentation ALL (upper panel) and relapse AML (lower

panel) identifies a common breakpoint with identical filler sequence in ALL and AML samples.

(C) Whole genome sequencing data of LS01 showing karyotype (outer circle), copy number

changes (log2 depth ratio in 1Mb windows, loss <2 green dots, gain >2 red dots) and structural

variants (translocations - red connecting lines, deletions – blue lines, inversions, purple lines).

(D) Schematic representation of identified fusion variants, located within the major breakpoint

MLL region, present in analysed t(4;11) cases (detailed breakpoint description presented at

Fig.S1 and Table S1). Red connecting line indicates MLL-AF4 translocation positions of each

gene partner.

Figure 2. Cellular origin of the *MLL-AF4* leukaemia. (A) Evaluation of B-cell receptor

repertoires on ALL (presentation) and AML (relapse) lineage switch, and MPAL cases

identified three different patterns. Pattern 1 - with clonotypes on the ALL only (LS01, LS02,

LS04). Pattern 2 - B-cell receptor-containing clones on ALL and AML, but distinct to each other

(LS03, LS06, LS07, LS08, LS10, MPAL2). Pattern 3 - B-cell receptor-containing clones shared

between ALL and AML (LS05, LS09, MPAL1). Graphs represent frequency of clones identified

with WES and/or RNAseq in LSAL and MPAL samples, respectively. (B) Presentation ALL

sample only produces ALL in all mouse strains transplanted. In contrast, Relapse AML sample

does not engraft in lymphoid supportive NSG mice and produces only AML in myeloid

supportive MISTRG mice. (C) Post-transplantation flow cytometric analysis of LS01PALL

produced in NSG mice (top panel) and LS01RAML produced in MISTRG mice (bottom panel).

Figure 3. Haematopoietic stem/progenitor populations carry *MLL-AF4*. (A) Flow

cytometric sorting strategy for haematopoietic stem/progenitor cell (HSPC) populations. (B)

PCR identification of the specific *MLL-AF4* fusion within sorted HSPC populations LS01 and

MPAL1 cases. (C) Summary of *MLL-AF4* positivity within different HSPC populations analysed

in patients LS01PALL and early progenitors and lymphoid fraction of MPAL1, presented as

red and pink circles, respectively. (D) Flow cytometric analysis showing sequential

transplantation of LS01 ALL presentation sample across four generations of NSG mouse

xenografts. The HSC population (CD34+CD38-CD45RA-CD90+) is maintained across four

mice generations. (E) Proportion of bone marrow human CD34+ cells in CD34+CD38-

CD45RA-CD90+ HSC gate in all analysed xenografts. (F) PCR identification of the specific

MLL-AF4 within the sorted HSPC xenograft sample.

Figure 4. Transcriptional reprogramming in lineage switch and MPAL cases.

(A) Heatmap showing the top 100 differentially expressed genes between ALL and AML from

six lineage switch (LS01, LS03, LS04, LS05, LS06, LS10) and two MPAL cases, ranked by

stat value. (B) Enrichment of myeloid growth and differentiation signature in relapsed samples

(left panel) identified by GSEA analyses, also pointing to downregulation of genes highly

correlated with acute lymphoblastic leukemia (middle and right panel). Gene set enrichment

analyses have been performed based on data derived from six lineage switch samples. FDR

– false discovery rate, NES – normalised enrichment score. (C-E) Differential expression of

(C) lineage specific, (D) immunoglobulin recombination machinery, and (E) genes encoding

immunoglobulin heavy and light chains in lineage switch and MPAL cases. Error bars show

standard error of the mean (SEM) for lineage switch cases and ranges for two MPAL cases.

Figure 5. Chromatin re-organisation and differential transcription factor binding

underpins lineage switching. (A) DNase hypersensitivity site sequencing identifies 13619

sites with a log2 fold reduction and 12203 sites with a log2 fold increase following lineage

switch to AML. (B) University of California, Santa Cruz (UCSC) genome browser screenshot

displaying differential expression at lineage specific loci (lower red tracks) accompanied by

altered DNase hypersensitivity (upper black tracks) proximal to the transcriptional start site

(TSS) of *CD33*. (C) UCSC genome browser screenshot for *CD19* zoomed in on an ALL-

associated DHS with EBF occupation as indicated by high resolution DHS-seq and Wellington

analysis. FP - footprint. (D) Heat maps showing distal DHS regions specific for AML relapse

on a genomic scale. Red and green indicate excess of positive and negative strand cuts,

respectively, per nucleotide position. Sites are sorted from top to bottom in order of decreasing

Footprint Occupancy Score. (E) De novo motif discovery in distal DHSs unique to AML relapse

as compared to ALL relapse as shown in (D). (F) EBF1 and C/EBP binding motifs demonstrate

differential motif density in presentation ALL and relapse AML. (G) Enrichment of a myeloid

C/EBPA network gene set in signatures associated with relapse AML and diagnostic ALL

samples as identified by GSEA (H) Differential expression of TFs cognately binding to

differentially accessible motifs shown in (F). TFs whose binding motifs show increased

accessibility in ALL are in blue whilst those showing increased accessibility in AML are in red.

RUNX1 in black reflects enriched accessibility of different RUNX1 binding sites in ALL and

AML. Error bars show SEM or ranges in LSAL and MPAL cases, respectively.

Figure 6. Alternative splicing in lineage switch and MPAL cases. (A) Volcano plots

demonstrating differential usage of exon-exon junctions in the transcriptome of AML/myeloid

versus ALL/lymphoid cells of lineage switch (LS01, LS03 & LS04) or MPAL patients. The

vertical dashed lines represent two-fold differences between the AML and ALL cells and the

horizontal dashed line shows the FDR-adjusted q-value threshold of 0.05 (upper panel). Venn

diagrams (lower panel) showing distribution of splice variants identified as significantly

changed in AML (or myeloid fraction of MPAL patients), including exon-exon junctions (DEEj),

differential exome usage (DEU) and retained introns (RI). (B) Pie charts showing the

classification of non-differential (non-DEEj) and differential (DEEj) exon-exon junctions.

Shown are the percentages of splicing events assigned to a particular mode of splicing.

Complex splicing event corresponds to several (two or more) alternative splicing incidents

occurring simultaneously in the same sample. (C) Enrichment analysis of affected signalling

pathways by the exon-exon junctions (DEEj) and differential exome usage (DEU) in the LSAL

AML relapse and myeloid compartment of MPAL patients. Pathway enrichment analysis has

been   performed   with   https://biit.cs.ut.ee/gprofiler/gost   under   the   highest   significance

threshold, with multiple testing correction (g:SCS algorithm). (D) Fold change expression levels

of total gene among genes identified to be affected by alternative splicing process (left panel)

and differentially spliced variants in lineage switched and myeloid compartments of MPAL

patients (right panel). (E) Schematic representation of the affected mRNA structure (and its

probable consequence depicted in red) within several selected genes (upper panel).

Corresponding normalized expression level (vs reference gene *TBP*) in two tested lineage

switch patients (LS03 and LS04) and one MPAL (lower panel). Shown is the ratio of analysed

splice variant expression level between AML (or myeloid) and ALL (or lymphoid) populations.

(F) Splice variants of *MLL-AF4* identified in MPAL patients and one lineage switch sample

(LS10). Left panel represents junction read counts of the fusion oncogene, identified by the

RNAseq analysis, with confirmation of the expression of both variants analysed by qRT-PCR

(MPAL1, right panel). Both splice variants, further confirmed by Sanger sequencing, showed

complete sequences of *MLL*ex9 or *MLL*e10 and either complete or truncated (for 3 nucleotides

at the 5'end) *AF*4ex4, respectively.

Figure 7. Molecular characterisation of lineage switch *MLL-AF4* leukaemias. (A) Whole

exome sequencing (WES) data showing genes recurrently mutated within the analysed

cohort, involved in the cell growth, communication and metabolism and genes mutated in

single cases belonging to the same function protein complexes (e.g. DNA polymerases,

epigenetic complexes). Data are presented according to the disease timepoint/cell lineage

and age of the patient. Depicted are major single nucleotide variants (SNVs) that were found

in >33% of reads and minor SNVs in <33% reads. (B) Clonal SNV patterns identified by whole

genome sequencing (WGS) in LS01 ALL and AML samples, constructed from counts of each

mutation-type at each mutation context, corrected for the frequency of each trinucleotide in

the reference genome. (C) Comparison of the whole exome sequencing and RNAseq data

obtained for LS01 patient identifies 6, 98, and 10 SNVs expressed only in ALL, AML, and

shared between ALL and AML, respectively. 12 SNVs exclusive for the AML relapse, predicted

(by Condel scoring) to have deleterious effects, were subjected to multiplex PCR followed by

next-generation sequencing analysed within each of the purified haematopoietic sub-

populations. Circles with solid colour indicate VAF >30%, light colour and dashed line indicates

VAF 5-30%. Remaining genes (yellow circle) represent the 10 other SNVs (out of 12 SNVs)

which showed the same pattern in the frequency of mutation acquisition.

Figure 8. Epigenetic modulatory genes influence lineage specific expression profiles.

(A) CHD4 scheme; the R1068H mutation is located in the critical helicase domain of CHD4 at

a highly conserved residue. An * (asterisk) indicates positions which have a single, fully

conserved residue, a : (colon) indicates conservation between groups of strongly similar

properties - scoring > 0.5 in the Gonnet PAM 250 matrix, a . (period) indicates conservation

between groups of weakly similar properties - scoring =< 0.5 in the Gonnet PAM 250 matrix.

(B) Fold change in expression of NuRD complex members (*CHD4*, *MTA1*, *RBBP4*, *MBD3*)

and *PHF3* following lineage switched relapse (left panel) and in MPAL cases (right panel). (C)

Expression of lineage specific genes following knockdown of *PHF3*, *CHD4* or the combination,

relative to non-targeting control construct in the *MLL-AF4* positive ALL cell line, SEM. (D) Flow

cytometric analysis of the surface CD33 expression following knockdown of *PHF3*, *CHD4* or

the combination in the SEM cell line. shNTC – non-targeting control. (E) Gene set enrichment

analysis of RNA sequencing data derived from: knockdown of *PHF3* and *CHD4* in the SEM

cell line (left panel) and lineage switch leukaemia cases (right panel). Shown is negative

correlation of shCHD4/shPHF3 and relapse samples with Jaatinen haematopoietic stem cell

signature (upper panel) and Haddad B-lymphocyte progenitor signature (lower panel). (F)

Expression of lineage specific cell surface markers CD19 (lymphoid) and CD33 (myeloid)

following culture of *MLL-Af4* transformed hCD34+ cord blood progenitor cells in lymphoid

permissive conditions. Knockdown of *PHF3*, *CHD4* or the combination disrupts the dominant

lymphoid differentiation pattern seen in non-targeting control (shNTC). (G) Assessment of

PHF3 knockdown influence on the surface marker expression after longer incubation period

(33 days); *CHD4* knockdown impaired cellular survival upon longer *in vitro* culture.

Figure S1. Sequencing of *MLL-AF4* fusion breakpoints from DNA (LS02, LS03, LS04,

LS07, LS08, LS09) or RNA (LS05, LS06, LS10, MPAL1, MPAL2).

Figure S2. Immunoglobulin rearrangement on ALL and AML LS01 according to the

standard protocol BIOMED-2. Clonal peaks are coloured in dark blue, as indicated in the

multiplex amplification FR1, FR2, and FR3 of the VH segments LS01PALL (left panel). The

multiplex results were confirmed by single amplification VH-JH primer sets, which further

showed two clonal VH3, one clonal VH1, and clonal VH4 rearrangement. No clonal peaks are

seen in LS01RAML (right panel).

Figure S3. Evaluation of *MLL-AF4* presence in HSPC populations. (A) Flow cytometry

plots of single sorted HLA-DR+CD14+CD11c+ monocytes LS01PALL. The two *MLL-AF4*

positive cells, cells 11 and 18, are highlighted in red throughout the sorting strategy (lower

panel). (B) Amplification of the *MLL-AF4* in cells 11, 18 and bulk LS01 sample shows the

expected 299 base pair band. (C) Flow cytometry plots showing sorting strategy for lymphoid

and myeloid populations of MPAL1 and MPAL2. (D) Bulk sort plots for HSC and MPP

populations (upper panel) and single cell plots for Lin-CD34+ populations (lower panel) sorted

from MPAL1 sample. (E) Amplification of the *MLL-AF4* in single- cell derived clones Lin-

CD34+, lymphoid, and myeloid fractions of MPAL1 patient.

Figure S4. Transcriptional profiles of ALL and AML lineage switch leukaemia.

(A) Non-negative matrix factorisation analysis of paired presentation ALL cases and relapse

AML cases, against normal haematopoietic precursors (Novershtern *et al*, 2011). Presentation

ALL cases demonstrate a high expression of B cell metagene signature, whilst relapse AML

cases continue to express this signature to a variable degree (upper panel). Relapse AML

cases demonstrate high expression of myeloid metagene signature, at a comparable level to

that of GMP (lower panel). (B) Non-negative matrix factorisation analysis of paired

presentation ALL cases and relapse AML cases, against ALL and AML cases, with and without

MLL rearrangement (Andersson *et al* 2015, Bolouri *et al* 2018). Heatmap shows clustering of

presentation cases (dark blue) with B-ALL *MLL-AF4* and relapse cases (dark green) with AML

with *MLL*r. (C) Differential promoter accessibility is associated with higher expression of *PAX5*,

LEF1, and *CD79A* in presentation ALL cases, and of *CSF3R*, *KIT* and *CSF2RA* in relapse

AML cases. TSS – transcriptional start site. (D) Gene set enrichment analysis indicates

impaired DNA repair and cell cycle progression in AML relapse. NES – normalised enrichment

score.

Figure S5. Impact of leukaemia lineage switch on *MLL-AF4*-regulated genes. (A) HOXA

cluster analysis includes a general reduction on HOXA3-10 in relapse AML cases. (B)

Differential chromatin accessibility and expression across the HOXA cluster at ALL

presentation and AML relapse of case LS01, demonstrating differential DNase hypersensitivity

of HOXA cluster. (C) Differential expression across the HOXA cluster in five cases of lineage

switch from ALL (upper panel for each case) and AML (lower panel). (D) The list of direct *MLL-*

AF4 target genes obtained by overlaying SEM cells and t(4;11) patient cells ChIPseq data

(Kerry *et al.*, 2017) and differentially expressed genes in our lineage switch and MPAL cases

(Venn diagram, upper panel). Highly enriched genes are further listed on the lower panel. (E)

GSEA showing loss of chromatin modifying enzymes signature in AML relapse.

Figure S6. High resolution DNase hypersensitivity sequencing. (A) UCSC genome

browser screenshot for *RUNX1* focused on an AML-associated DHS with C/EBP occupation

as indicated by high resolution DHS-seq and Wellington analysis. FP - footprint. (B) Heat maps

showing distal DHS regions specific for ALL presentation on a genomic scale. Red and green

indicate excess of positive and negative strand cuts, respectively, per nucleotide position.

Sites are sorted from top to bottom in order of decreasing Footprint Occupancy Score. *De*

novo motif discovery in distal DHSs unique to ALL as compared to AML relapse, as shown on

the table, right panel.

Figure S7. Alternative splicing analysis. (A) RNAseq read counts across all analysed

patient samples. (B) Qualitative analysis of exon-exon spanning reads from LS01, LS03 &

LS04 samples. Upper panel represents distribution of the unfiltered and unnormalized reads,

lower panel represents filtered and voom normalized reads. Expression (abundance) of the

exon-exon junctions is presented as log2-transformed counts per million (CPM). (C)

Abundance of non-differential versus differential exon-exon junctions in the transcriptome of

LSALs (left panel) and MPAL series of samples (right panel). According to limma/diffSplice

approach, junctions were divided into non-differential (upper panel) and differential (bottom

panel) splicing events. In each boxplot, the horizontal line represents the median of

distribution, box shows the interquartile range, and whiskers are the minimum and maximum.

p-values were calculated using the two-sided Mann-Whitney U test. (D) Venn diagrams

showing distribution of identified retained intron (RI) across both LSAL and MPAL patients with

corresponding GO terms describing cellular components affected in the AML relapse of

lineage switched patients. (E) Venn diagrams showing distribution of identified differential

exon-exon junctions (DEEj), significantly expressed in the AML relapse or myeloid

compartment of MPAL patients with corresponding GO terms describing affected Reactome

pathways and cellular components. GO terms analysis has been performed with

https://biit.cs.ut.ee/gprofiler/gost under the highest significance threshold, with multiple testing

correction (g:SCS algorithm).

Figure S8. The functional role of epigenetic modifiers on leukaemia lineage switching.

(A) PHF3 scheme; the K1119I mutation is located at a highly conserved residue. An *

(asterisk) indicates positions which have a single, fully conserved residue, a : (colon) indicates

conservation between groups of strongly similar properties - scoring > 0.5 in the Gonnet PAM

250 matrix, a . (period) indicates conservation between groups of weakly similar properties -

scoring < 0.5 in the Gonnet PAM 250 matrix. (B) Mutual information sub-networks illustrate

high centrality of mutated genes *PHF3* (left panel) and *CHD4* (right panel) within the AML/ALL

transcriptional network. Size and colour of nodes reflect differential expression between AML

and ALL. (C) Myeloid marker CD33 expression level upon knockdown of *CHD4*, *PHF3*, and

non-targeting control (NTC) in *MLL*r (t(4;11)) cell line RS4;11 and non-*MLL*r cell lines, 697 and

REH cells. (D) CD19 and CD33 surface expression level change upon knockdown of *CHD4*

and *PHF3* in the PDX sample generated from the first relapse of patient LS03 (ALL relapse).

(E) Protein network analysis of differentially expressed NuRD and PRC1 members, showing

interactions between proteins (and clusters, depicted as blue and red colour); interaction

strength is shown as increasing thickness of the joining line (https://string-db.org/). (F) Fold

change expression level changes of PRC1 members following lineage switched relapse and

in MPAL cases. (G) Expression of lineage specific surface markers following *PCGF6*

knockdown in SEM cells. Knockdown of *PCGF6* results in upregulation of *CD33*, *CSFR3* and

KIT mRNA (upper panel). Surface CD33 expression increases substantially with shPCGF6_1

and shPCGF6_3, compared with shNTC (lower panel). (H) GSEA showing increased

expression of PRC2 target genes and impaired function of PRC1 complex in the relapse AML

samples. (I) Boxplot of NMF projections of CD19+ cord blood (CB_19pos) and CD33+ cord

blood (CB_33pos) populations relative to the derived AML with *MLL*r metagene (red boxes)

and B precursor ALL with *MLL*r metagene (turquoise boxes).

Table titles and legends

Supplementary Table 1. Clinical characteristics of patient cases

Supplementary Table 2. B cell receptor repertoires in lymphoid and myeloid

populations. The frequency represents the stated specific sequence as a frequency of

the total reads of that clonotype.

Supplementary Table 3. Read counts of *MLL-AF4*, wild type (wt), and mutant (mut)

sequences from haematopoietic stem/progenitor cell populations sorted from

presentation and relapse samples of LS01. The MPP populations first showing clonal

mutations of candidates *CHD4* and *PHF3* are highlighted in yellow.

Supplementary Table 4. Differential gene expression across paired

presentation/lymphoid and relapse/myeloid samples.

Supplementary Table 5. Alternative splice variants - mRNAs with retained introns

(RI).

Supplementary Table 6. Alternative splice variants - genes with differential exon-

exon junctions (DEEj).

Supplementary Table 7.　Coding mutations identified in exome sequencing.

Genes in red lettering are major clonal variants with variant allele frequencies >0.33.

Genes in blue lettering are minor clones with variant allele frequencies 0.09-0.32.

Genes in black lettering show a variant allele frequency of <0.09. Shared genes

between presentation and relapse are indicated with red boxes.

**Supplementary Table 8.　**ARACNE analysis of coding mutations projected onto a

mutual information network constructed from published transcriptomic data from ALL

and AML cases.

757   STAR Methods

758

759   KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
CD3-APC-H7	BD Biosciences	Cat#641397; RRID:AB_1645731
CD3-FITC	BD Biosciences	Cat#345763
CD3-BV650	Biolegend	Cat#300468
CD10-BV650	BD Biosciences	Cat#563734; RRID:AB_2738393
CD11b-APC/Fire750	Biolegend	Cat#301351
CD14-FITC	BD Biosciences	Cat#555397; RRID:AB_395798
CD14-BV605	Biolegend	Cat#367125
CD16-FITC	BD Biosciences	Cat#335035
CD19-PE-CF594	BD Biosciences	Cat#562294; RRID:AB_11154408
CD19-APCCy7	Biolegend	Cat#363010; RRID:AB_2564193
CD19-BV421	Biolegend	Cat#302234; RRID:AB_11142678
CD19-PECy7	Biolegend	Cat#363011
CD20-PE	BD Biosciences	Cat#345793
CD20-FITC	Biolegend	Cat#302303
CD33-APC	BD Biosciences	Cat#345800
CD33-APCCy7	Biolegend	Cat#366614; RRID:AB_2566416
CD33-BV421	Biolegend	Cat#303416; RRID:AB_2561690
CD34-PerCPCy5.5	BD Biosciences	Cat#347222
CD34-APCCy7	Biolegend	Cat#343514; RRID:AB_1877168
CD34-APC	Biolegend	Cat#343607
CD38-PeCy7	BD Biosciences	Cat#335825
CD38-PE	Biolegend	Cat#303506
CD45RA-BV510	Biolegend	Cat#304142; RRID:AB_2561947
CD56-FITC	BD Biosciences	Cat#345811
CD64-BV711	Biolegend	Cat#305041
CD90-A700	Biolegend	Cat#328120; RRID:AB_2203302
CD90-PerCPCy5.5	Biolegend	Cat#328118; RRID:AB_2303335
CD90-BV421	Biolegend	Cat#328122

CD117-PE	BD Biosciences	Cat#332785
CD117-BV605	BD Biosciences	Cat#562687; RRID:AB_2737721
CD123-BV421	BD Biosciences	Cat#306018; RRID:AB_10962571
HLA-DR-BV786	Biolegend	Cat#307642; RRID:AB_2563461
HLA-DR-A700	Biolegend	Cat#560743; RRID:AB_1727526
HLA-DR-PerCPCy5.5	Biolegend	Cat#307629
Bacterial and Virus Strains		
One Shot STBL3 chemically competent *E. coli*	Invitrogen	Cat#C7373-03
Chemicals, Peptides, and Recombinant Proteins		
Cytox Green-FITC	Life Tech	Cat#S34860
Deoxyribonuclease I (DNase I)	Worthington-Biochem	Cat# LS006328
AmpliTaq Gold	Applied Biosystems	Cat#4311818
Phusion® High-Fidelity PCR Master Mix with HF Buffer	NEB	Cat#M0531L
IL-6	MACS Miltenyi Biotec	Cat#130-095-352
IL-3	R&D Systems	Cat#203-IL-010
IL-7	R&D Systems	Cat#207-IL-005
SCF	StemCell Technologies	Cat#78062.1
TPO	R&D Systems	Cat#288-TP
FLT-3 Ligand	R&D Systems	Cat#308-FK
Critical Commercial Assays		
AMPure XP	Beckman Coulter	Cat#A63880
QIAamp DNA FFPE Tissue Kit	Qiagen	Cat#56404
AllPrep DNA/RNA Mini Kit	Qiagen	Cat#80204
QIAamp DNA Mini Kit	Qiagen	Cat#51306
innuPREP DNA/RNA Mini Kit	Analytik Jena	Cat#845-KS-2080050
RNeasy Mini Kit	Qiagen	Cat#74106
NEBNext Ultra Directional RNA Kit	NEB	Cat#E7420S
REPLI-g Single Cell Kit	Qiagen	Cat#150345
EndoFree Plasmid Maxi Kit	Qiagen	Cat#12362
Fluidigm Access Array Barcode Library for Illumina Sequencers	Fluidigm	Cat#100-4876
RevertAid H Minus First Strand cDNA Synthesis Kit	Thermo Fisher Scientific	Cat#K1632

Platinum SYBR Green qPCR SuperMix-UDG with ROX, 2X	Invitrogen	Cat#11744500
Deposited Data		
Exome-seq data	This paper	PRJNA547947
Genome-seq data	This paper	PRJNA547815
Ig rearrangement sequencing data	This paper	PRJNA511413
RNA-seq data	This paper	GSE132396
DNase hypersensitivity sequencing data	This paper	GSE130142
Experimental Models: Cell Lines		
SEM	DSMZ	Cat#ACC546; RRID:CVCL_0095
REH	DSMZ	Cat#ACC22
697	DSMZ	Cat#ACC42
RS4;11	DSMZ	Cat#ACC508
MS-5	DSMZ	Cat#ACC441; RRID:CVCL_2128
Human cord blood CD34$^+$ MLL-Af4	(Lin *et al.*, 2016)	N/A
Experimental Models: Organisms/Strains		
NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ (NSG) mice	Charles River Laboratories	N/A
M-CSF^h; IL-3/GM-CSF^h; hSIRPA^{tg};TPO^h;Rag2$^-$;γc$^-$ (MISTRG) mice	Jackson Laboratory	N/A
Oligonucleotides		
shPHF3-1	GPP, Broad Institute	TRCN0000019118
shPHF3-2	GPP, Broad Institute	TRCN0000019114
shPHF3-3	GPP, Broad Institute	TRCN0000274376
shCHD4-1	GPP, Broad Institute	TRCN0000380981
shCHD4-2	GPP, Broad Institute	TRCN0000021363
shCHD4-3	GPP, Broad Institute	TRCN0000021360
shPCGF6-1	GPP, Broad Institute	TRCN0000229804
shPCGF6-2	GPP, Broad Institute	TRCN0000073109
shPCGF6-3	GPP, Broad Institute	TRCN0000073109
shAUTS2-1	GPP, Broad Institute	TRCN0000119058
shAUTS2-2	GPP, Broad Institute	TRCN0000304019
shAUTS2-3	GPP, Broad Institute	TRCN0000304081
shNTC (ATCTCGCTTGGGCGAGAGTAAG)	(Schwarzer *et al.*, 2017)	N/A
PHF3 Fw (ATGGACCTGGGCTTGAACTG)	This paper	N/A

PHF3 Rev (TGGTGGTGCACTTTCAGGAG)	This paper	N/A
CHD4 Fw (TGCTGACACAGTTATTATCTATGACTCTGA)	This paper	N/A
CHD4 Rev (ACGCACGGGTCACAAACC)	This paper	N/A
PCGF6 Fw (GGGAAATCCGACGTGCAAT)	This paper	N/A
PCGF6 Rev (GGAGAAACCACAAGACCATAATGA)	This paper	N/A
AUTS2 Fw (AAAAGGACCCGAGGTTGACA)	This paper	N/A
AUTS2 Rev (GCGATGTGAACATGCATAGCA)	This paper	N/A
GAPDH Fw (GAAGGTGAAGGTCGGAGTC)	This paper	N/A
GAPDH Rev (GAAGATGGTGATGGGATTTC)	This paper	N/A
CD19 Fw (TGACCCCACCAGGAGATTCTT)	This paper	N/A
CD19 Rev (CACGTTCCCGTACTGGTTCTG)	This paper	N/A
CD33 Fw (CTCGTGCCCTGCACTTTCTT)	This paper	N/A
CD33 Rev (CCCGGAACCAGTAACCATGA)	This paper	N/A
CSF3R Fw (CCCAGGCGATCTGCATACTT)	This paper	N/A
CSF3R Rev (AACAAGCACAAAAGGCCATTG)	This paper	N/A
KIT Fw (GGACCAGGAGGGCAAGTCA)	This paper	N/A
KIT Rev (GATAGCTTGCTTTGGACACAGACA)	This paper	N/A
Recombinant DNA		
pCMVD8.91 packaging vector	Life Science Market	Cat# PVT2323
pMD2.G envelope vector	Addgene	Cat#12259
pLKO5d.SFFV.miRNA30n	(Schwarzer *et al.*, 2017)	N/A
Software and Algorithms		
Bowtie2	(Langmead *et al.*, 2009)	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
RUVSeq	(Risso *et al.*, 2014)	http://bioconductor.org/packages/RUVSeq
DESeq2	(Love, Huber and Anders, 2014)	http://bioconductor.org/packages/DESeq2
GSEA	(Subramanian *et al.*, 2005)	http://software.broadinstitute.org/gsea
ARACNe2	(Margolin *et al.*, 2006)	N/A
MiXCR	(Bolotin *et al.*, 2015)	http://mixcr.milaboratory.com/
MuTect	(Cibulskis *et al.*, 2013)	http://www.broadinstitute.org/cancer/cga/MuTect

MuTect2	(Cibulskis *et al.*, 2013)	http://www.broadinstitute.org/cancer/cga/MuTect
VEP, version 90	(Cibulskis *et al.*, 2013)	https://github.com/Ensembl/ensembl-vep
Tximport	(Soneson, Love and Robinson, 2016)	http://bioconductor.org/packages/tximport/
ISAAC	(Raczy *et al.*, 2013)	https://github.com/sequencing
Strelka	(Saunders *et al.*, 2012)	ftp://strelka@ftp.illumina.com/
Manta	(Chen *et al.*, 2016)	https://github.com/Illumina/manta
Wellington	(Piper *et al.*, 2015)	http://jpiper.github.com/pyDNase
Java Treeview	(Saldanha, 2004)	http://jtreeview.sourceforge.net
Cytoscape	(Shannon *et al.*, 2003)	http://www.cytoscape.org/

CONTACT FOR REAGENT AND RESOURCE SHARING

As Lead Contact, Olaf Heidenreich (O.T.Heidenreich@prinsesmaximacentrum.nl) is responsible for all reagent and resource requests.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Patient samples and data

Patients were diagnosed by local haematology specialists according to contemporary clinical diagnostic criteria based on morphology and immunophenotypic analysis. All patient samples were collected at the point of diagnosis, remission following treatment or relapse and stored with written informed consent for research in one of six centres (Newcastle Haematology Biobank, Newcastle, UK; University Hospital Schleswig-Holstein, Kiel, Germany; Dmitry Rogachev National Medical Research Center of Pediatric Hematology, Oncology and Immunology, Moscow, Russia; Haematological Malignancy Diagnostic Service, Leeds, UK; Princess Maxima Center for Pediatric Oncology, Utrecht, The Netherlands; Cincinnati Children's Hospital Medical Center, Cincinnati, USA). Mononuclear cells were isolated from

bone marrow or peripheral blood by density centrifugation followed by immediate extraction of

DNA or RNA, or cryopreservation in the presence of 10% v/v DMSO.

Samples were requested and used in accordance with the ethical approvals granted to each

of the local/institutional ethical review boards (NRES Committee North East - Newcastle &

North Tyneside 1, UK, reference 07/H0906/109+5; Medical Faculty Christian-Albrechts

University, Kiel, reference A 103/08; Dmitry Rogachev National Medical Research Center,

Moscow, references MB2008: 22.01.2008, MB2015: 22.01.2015, ALL-REZ-2014: 28.01.2014;

Haematological Malignancy Research Network, Yorkshire, UK, reference 04/Q1205/69;

Haematological Malignancy Diagnostic Service, Leeds, UK, reference 14/WS/0098; Erasmus

MC METC, Netherlands, reference MEC-2016-739; IRB of Cincinnati Children's Hospital,

USA, reference 2010-0658) and in accordance with the Declaration of Helsinki. Each

patient/sample was allocated an anonymised reference and no identifiable information was

shared.

Cell Lines

SEM, 697, REH and RS4;11 cells were grown in RPMI-1640 with 10% fetal bovine serum

(FBS, Sigma)] at 37°C in a humidified 5% CO2 incubator. All cell lines contain no PHF3,

CHD4 (with exception of the REH cell line - missense mutation at p.R61Q), PCGF6, AUTS2

mutation according to the CCLE (https://portals.broadinstitute.org/ccle) database.

MLL-Af4 transduced cord blood cells (Lin *et al.*, 2016) were cultured in IMDM with 10% FBS,

2 mM Glutamine supplemented with recombinant human SCF, IL-3, IL-6, FLT-3L, and TPO

(10 ng/ml each) at 37°C in a humidified 5% CO_2 incubator. To prime towards lymphoid

differentiation, the cells were co-cultured with MS-5 (RRID:CVCL_2128, murine) stroma cells

in α-MEM with 10% FBS, 2 mM glutamine supplemented with SCF, FLT-3L, and IL-7 (10 ng/ml

each). The cells were demi-populated weekly.

In Vivo **Mouse Studies**

In vivo studies were conducted in accordance with the UK Animals (Scientific Procedures) Act

1986 under project licences PPL60/4552 and PPL60/4222 following institutional ethical

review.

NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wjl}*/SzJ (NSG, Charles River Labs and bred in-house) mice (both

sexes) aged 8 -10 weeks old were transplanted intra-femorally under isoflurane anaesthesia

with 10^6 cells from sample LS01PALL. Sample LS01RAML was transplanted into 5 NSG mice

without evidence of engraftment.

MISTRG   mice,   expressing   four   human   cytokines   (M-CSF^h;   IL-3/GM-CSF^h;

hSIRPA^{tg};TPO^h;Rag2$^-$;γc$^-$, Jax Lab (Rongvaux *et al.*, 2014), bred in house) were transplanted

intra-hepatically. Three day old mice (n = 2) were sub-lethally irradiated (1.5 Gy) 24 h before

the transplantation. Each mouse received 10^6 mononuclear cells from sample LS01RAML.

Mice were humanely killed before or when they displayed end points specified by the licenses

(tumours reached 1.5 cm in diameter, lost >10% weight in 3 consecutive days or 20% at any

time, or displayed signs of ill). PDX cells were harvested from engrafted spleen (NSG) or

abdominal masses (MISTRG).

METHOD DETAILS

Long-distance inverse PCR experiments

MLL rearrangement sequences were identified using long-distance inverse PCR (LDI-PCR)

(Meyer *et al.*, 2005, 2013). DNA (1 μg) from each patient was digested with restriction enzyme

*Bam*HI and purified by phenol extraction and ethanol precipitations to remove the residual

enzyme. The digested product was re-ligated by incubation at 16°C overnight in the presence

of 5 units T4 DNA ligase, 50 µl total volume, to form DNA circles. Five microliters (100 ng) re-

ligated genomic DNA was used for the LDI-PCR.

Five *MLL*-specific primers (primers A-E (Meyer *et al.*, 2005)) were designed and used in four

different combinations (A–B, A–C, A–D, and A–E). The LDI-PCR reactions were performed

using TripleMaster PCR System (Eppendorf) according to the manufacturer's instructions. The

amplicons were separated on 0.8% agarose gels, extracted, and sequenced. Sequencing

results   were   annotated   by   blasting   the   human   genome   database

www.ncbi.nlm.nih.gov/genome/seq/Blast.

PCR-based clonality immunoglobulin gene targets

Clonality analysis based on immunoglobulin gene rearrangements was performed according

to the BIOMED-2 Concerted Action BMH-CT98-3936 using standardised protocols and primer

sets (van Dongen *et al.*, 2003). DNA was amplified by multiplex PCR assays IGH (VH–JH and

DH–JH) and IGK (Vκ–Jκ and Kde primers) using AmpliTaq Gold (Applied Biosystems) in

duplicate reactions. The PCR reaction consists of 100 ng DNA, 1 X ABI Gold Buffer, 10 pmol

of each primer, 200 µM final concentration dNTP, 1.5 mM final concentration $MgCl_2$, and 1 U

Taq enzyme in 50 µl volume. The PCR amplification parameters were: one cycle at 95°C for

10 min, thirty five cycles of 95°C for 45 s, 60°C for 45 s, and 72°C for 1.5 min, followed by one

cycle at 72°C for 10 min. The duplicate samples were analysed by GeneScan analysis on a

Life Technologies 3100 platform.

Exome sequencing

Germline DNA from cases LS08 and LS09 were extracted from formalin fixed paraffin

embedded remission bone marrow using QIAamp DNA FFPE Tissue Kit (Qiagen,

Cat#56404). Other DNA samples were extracted from either bone marrow or peripheral blood

using AllPrep DNA/RNA Mini Kit (Qiagen, Cat#80204), QIAamp DNA Mini Kit (Qiagen,

Cat#51306), or innuPREP DNA/RNA Mini Kit (Analytik Jena, Cat#845-KS-2080050),

according to manufacturers' instructions. The exons were captured using SureSelect XT2

Human All Exon V6 (Agilent), and sequenced by paired-end 75 bp sequencing on HiSeq4000

(Illumina), resulting in roughly 45 million reads per sample. DNA from the myeloid and

lymphoid cellular compartments derived from MPAL patients samples, were pre-processed

with KAPA HyperPlus Kit (Roche) followed by exons enrichment with KAPA HyperCapture Kit

(Roche), and sequenced by paired-end 300 bp sequencing on NovaSeq6000 (Illumina),

resulting in roughly 25 million reads per sample.

Raw reads were aligned to human reference genome (hg19 or hg38 for lineage switch or

MPAL patients, respectively) using Burrows-Wheeler Aligner (BWA) 0.7.12 (Li and Durbin,

2009) and were processed using the Genome Analysis Toolkit (GATK, v3.8 or 4.1) best

practices recommended workflow for variant discovery analysis (Mckenna *et al.*, 2010;

DePristo *et al.*, 2011; van der Auwera *et al.*, 2013). MuTect (v1.1.7) and MuTect2 (4.1) were

used to identify somatic variants for each matched sample pair (Cibulskis *et al.*, 2013).

Variants were annotated using Ensembl Variant Effect Predictor (VEP, version 90) (Cibulskis

et al., 2013).

RNA sequencing

Total RNA was extracted with AllPrep DNA/RNA Mini Kit (Qiagen, Cat#80204), innuPREP

DNA/RNA Mini Kit (Analytik Jena, Cat#845-KS-2080050), or TRIzol (Thermo Fisher Scientific,

Cat# 15596026) followed by RNeasy Mini Kit (Qiagen, Cat#74106) from either bone marrow

or peripheral blood, according to manufacturers' instructions. Messenger RNA was captured

using NEBNext Ultra Directional RNA Kit in combination with NEBNext poly(A) mRNA

Magnetic Isolation Module or KAPA RNA HyperPrep Kit with RiboErase (HMR) in case of

lineage switch or MPAL patients respectively, and submitted for paired-end 150 bp sequencing

on HiSeq4000 (Illumina) or paired-end 300 bp sequencing on NovaSeq6000 (Illumina)

depending on the analysed patients group. For each sample, transcript abundance was

quantified from raw reads with Salmon (version 0.8.2) (Patro *et al.*, 2017) using the reference

human transcriptome (hg38) defined by GENCODE release 27. An R package Tximport

(version 1.4.0) (Soneson, Love and Robinson, 2016) was used to estimate gene-level

abundance from Salmon's transcript-level counts. Gene-level differential expression analysis

was performed using DESeq2 (version 1.16.1) (Love, Huber and Anders, 2014).

Alternative splicing analysis

The differential splicing events were detected in RNAseq data obtained from all analysed

patient samples. Detailed analyses were restricted to samples with more than 15 million reads

per sample, which included four lineage switched (LS01, LS03, LS04 and LS10) and both

MPAL patients. The differential splicing events, including exon-exon junctions (DEEj),

differential exon usage (DEU) and retained introns (RI) were identified in both

presentation/relapse pairs or lymphoid/myeloid cellular fractions, using pipeline described

previously (Grinev *et al.*, 2021). Validation of identified different splice variants was performed

by real-time qPCR experiments.

Whole genome sequencing

Presentation, remission and relapse DNA samples from case LS01 were sequenced by

Illumina UK and analysed using the remission sample as the matching normal. Sequencing

reads were aligned to the human GRCh37.1 reference genome using ISAAC (Raczy *et al.*,

2013). Identification of somatic SNVs and small somatic indels (<50 bp) was performed by

Strelka (Saunders *et al.*, 2012). Large structural variants (including deletions, inversions,

duplications and insertions all >50bp and translocations) were called by Manta (Chen *et al.*,

2016).

Non-negative matrix factorization analysis

Non-negative matrix factorisation (NMF) was performed using data sets obtained with

permission. Data from Bolouri *et al.* (Bolouri *et al.*, 2018) were generated by the

Therapeutically Applicable Research to Generate Effective Treatments (TARGET) initiative,

phs000218, managed by the NCI. The data used for this analysis are available

[https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000218.v21.p7].

Information about TARGET can be found at http://ocg.cancer.gov/programs/target. Data from

Andersson *et al.* (Andersson *et al.*, 2015) was obtained with permission of the Pediatric Cancer

Genome Project from the European Genome-Phenome Archive (EGA study accession

EGAS00001000246). Data from Novoshtern *et al.* (Novershtern *et al.*, 2011) was obtained

openly from Gene Expression Omnibus (GEO dataset GSE24759). NMF was used to extract

metagenes from three model gene expression datasets including only leukaemic samples for

which information on recurrent genetic abnormalities was known (Novershtern *et al.*, 2011;

Andersson *et al.*, 2015; Bolouri *et al.*, 2018). We classified samples into the following 7

subgroups of interest: ALL without *MLL*r, ALL with *MLL-AF4* fusion, ALL with other types of

*MLL*r, AML with *MLL*r, AML with inv(16), AML with *RUNX1-ETO* fusion, AML with normal

karyotype. A matrix of gene expression data combining per gene read counts of samples from

each dataset was created, normalized and variance-stabilized transformed (vst) using

DESeq2 (Love, Huber and Anders, 2014). Batch effect between leukaemia derived datasets

(Andersson *et al.*, 2015; Bolouri *et al.*, 2018) was minimized using the remove unwanted

variation method from RUVSeq (Risso *et al.*, 2014) on upper-quartile normalized counts whilst

retaining variation associated with biological covariates of interest. To remove genes that did

not vary sufficiently across the model dataset, for RNAseq data, we capped expression values

at 20 and 10,000 read counts and only included genes with > 10-fold and >1000 read counts

between minimum and maximum. For the Affymetrix U133A microarray, we capped

expression values at 0.01 and 10,000 units and only probes with > 3-fold and >5 units

difference between minimum and maximum were considered in the NMF analysis. To identify

the most robust combination of metagenes and subgroups/clusters in the model datasets,

consensus bootstrapped NMF clustering was performed on filtered gene count matrix as

previously described (Schwalbe *et al.*, 2013). Briefly, we performed NMF and K-means

clustering, testing all combinations of 2-15 metagenes and clusters with bootstrapped

resampling method (n=100) to test for reproducibility. Cluster stability measures (Cohen's

kappa, average silhouette scores) were assessed to determine optimal combinations of

metagenes and clusters. Samples assigned to the same cluster fewer than 90% of replicates

were removed from the dataset. All genes in the refined model dataset were then column rank

normalised and metagenes scores were recalculated.

Metagenes were projected onto our RNAseq datasets (both lineage switch cases and *MLL-*

Af4 cord blood samples) by the pseudoinverse method as outlined in (Brunet *et al.*, 2004;

Tamayo *et al.*, 2007). All analysis and figures were generated using R/Bioconductor and by

using a modified version of the NMF scripts provided by (Brunet *et al.*, 2004).

DHS library generation, sequencing, and mapping

DHS analysis was performed as described previously (Ptasinska *et al.*, 2014; Cauchy *et al.*,

2015). DNase I (Worthington, Cat# LS006328) digestion was performed using ~5 million

patient sample cells using 8 units (presentation) or 14 units (relapse) for 3 min at 22°C in a 1

mM $CaCl_2$ supplemented buffer. Nuclear proteins were digested with 1 mg/ml Proteinase K

overnight at 37°C. DNase I digestion products were size-selected on an agarose gel, cutting

below 150 bp. High-throughput sequencing libraries were prepared from 10 ng of size-selected

material, using the Kapa Hyperprep kit as per manufacturer's instruction. Libraries were

sequenced with 50 bp single-end reads on an Illumina HiSeq 2500 sequencer according to

manufacturer's instructions.

Fastq files were generated using bcl2fastq (1.8.4) and subsequently aligned to the hg19

assembly (NCBI Build 37) with the use of bowtie2 (2.1.0) (Langmead *et al.*, 2009), with –very-

sensitive-local as a parameter. Read coverage generation and peak detection were carried

out using MACS 1.4.1 (Zhang *et al.*, 2008) using --keep-dup=all –g hs -w -S. Pairwise

comparisons were performed as previously described (Cauchy *et al.*, 2015). Digital

footprinting was carried out using the Wellington package using default parameters (Piper *et*

al., 2013). Differential footprinting analysis was carried out on footprints using the Wellington-

bootstrap (Piper *et al.*, 2015) package with default parameters. Average profiles and heatmaps

were obtained using the functions dnase_average_profile and dnase_to_javatreeview from

the Wellington package. Heatmaps were plotted using Java TreeView (Saldanha, 2004).

Flow cytometry and cell sorting

One million cells (of all tested cell lines, cord blood or LS03ALL PDX cells) were used for flow

cytometry analysis of CD19 (BV421) and CD33 (APCCy7). The antibody details are listed in

the Key Resources Table. Cells were collected and washed in a 3 ml wash buffer containing

0.2% BSA in PBS, re-suspended in 100 µl wash buffer and 5 µl each antibody was added,

followed by incubation at RT for 20 min in the dark. Cells were washed and re-suspended in

a 0.5 ml wash buffer. Flow cytometry was performed on FACSCanto II from Becton Dickinson

(BD) and data analysed with FlowJo (Treestar).

Haematopoietic hierarchy analysis was performed from 10^7 cells of primary and PDX samples.

The cells were collected and incubated for 30 min in a wash buffer containing 0.5% FBS, 2

mM EDTA in PBS. Following the incubation, the cells were washed and resuspended in a

wash buffer. Fluorescence activated cell sorting (FACS) was performed using a FACS Aria

Fusion (BD) or ASTRIOS EQ (Beckman Coulter). In order to separate early progenitors or

lymphoid and myeloid fractions, MPAL patient samples were sorted from 10^7 or $3*10^6$ cells for

MPAL1 and MPAL2 respectively. Sorted cells were collected into 1.5 ml microfuge tubes

containing 500 µl RPMI with 10% fetal calf serum or SFEM II media in case of cell lines or

primary samples, respectively. In order to analyse single cell derived clones, sorted early

progenitor cells derived from MPAL samples were deposited onto 384-well plates containing

75ul of SFEM II media (supplemented with SCF, FLT3L, IL6, IL3 and TPO). Single cells were

grown into 60-90% confluency colonies followed by RNA/DNA extraction and MLL-AF4

detection by PCR.

Single cell sorting and whole genome amplification (WGA)

Single cells were labelled (see flow cytometry and cell sorting section above) and sorted into

96 well plate (Eppendorf twin.tec PCR Plate 96 full skirt) using FACSAria Fusion (BD). The

plates were filled with 3.5 µl PBS/well prior to the sorting. Each plate included forty-five single

cell samples, two 0 cells as negative control, and one bulk (300 – 1,000 cells) as positive

control sample. The cells were prepared from as many as 10^5 cells in 500 µl buffer (0.5% FBS,

2 mM EDTA in PBS).

Cell sorting was performed using the "single cell" setting on FACSAria Fusion. Following the

sorting, the plate was centrifuged at 900 g for 1 min, snap-frozen on dry ice, and then kept at

-20°C until the whole genome amplification (WGA) procedure.

WGA was performed using REPLI-g Single Cell Kit (Qiagen, Cat#150345). The cells were

lysed by adding 1.5 µl Buffer D2 and incubation at 65°C for 10 min in a thermal cycler (HYBAID

PCRExpress). Subsequently, the reaction was terminated by adding 1.5 µl Stop Solution and

stored on the ice. The mixture of 20 µl REPLI-g sc Reaction Buffer and REPLI-g sc DNA

Polymerase was added and incubated at 30°C for 8 h. The reaction was terminated by heat

inactivation at 65°C for 3 min. The products were diluted 100 fold with TE Buffer ahead of PCR

amplification.

Nested multiplex PCR and targeted sequencing

The 12 mutation candidate driver genes and *MLL-AF4* LS01RAML were amplified using gDNA

as the template by nested multiplex PCR method. The primers were designed using Primer

Express (Applied Biosystems) software.

PCR amplifications were carried out using Phusion® High-Fidelity PCR Master Mix with HF

Buffer (NEB, Cat#M0531L) according to the manufacturer's instructions (25 µl reaction), plus

80 – 200 nM of each primer set. The first multiplex reaction parameters were: one cycle at

98°C for 2 min, thirty cycles of 98°C for 10 s, 63°C for 30 s, and 72°C for 30 s, followed by one

cycle at 72°C for 10 min. The products were diluted 500-fold, and 1 µl used as the template

for the second PCR reactions (25 µl reaction). The second/nested multiplex reaction

parameters were: one cycle at 98°C for 2 min, twenty cycles of 98°C for 10 s, 65°C for 30 s,

and 72°C for 30 s, followed by one cycle at 72°C for 10 min. The amplicons that were taken

forward   for   next-generation   targeted   sequencing   had   additional   CS1

(ACACTGACGACATGGTTCTACA) and CS2 (TACGGTAGCAGAGACTTGGTCT) Fluidigm

tag sequences on the nested PCR primer forward and reverse, respectively. The amplicons

were barcoded using Fluidigm Access Array Barcode Library for Illumina Sequencers

(Cat#100-4876) by taking 0.8 µl multiplex PCR products (multiplex group A-C), 4 µl Fluidigm

barcode primer (400 nM final concentration), 10 µl of 2X Phusion Master Mix (NEB,

Cat#M0531L), and 3.6 µl H_2O (20 µl reaction). The barcoding PCR reaction parameters were:

one cycle at 98°C for 2 min, six cycles of 98°C for 10 s, 60°C for 30 s, and 72°C for 1 min,

followed by one cycle at 72°C for 10 min. The products were run on 2% agarose gel and

extracted using the QIAquick Gel Extraction Kit (Qiagen, Cat#28706). The purified products

were submitted for paired-end 300 bp sequencing on MiSeq (Illumina), resulting in >1,000

coverage per gene.

***MLL-Af4* stem cell expression analysis**

Following myeloid or lymphoid culture of *MLL/Af4* transduced CD34+ cord blood cells,

CD19+CD33-, CD19-CD33+ and CD19-CD33- populations were flow sorted, lysed and RNA

extracted using RNeasy Micro Kit (Qiagen), according to the manufacturer's instructions. Input

RNA was equilibrated to a starting input cell number of 300 cells per population before cDNA

and sequencing library production were performed using SMARTSeqv4 (Clontech) and

NexteraXT (Illumina) kits, according to manufacturer's instructions.   The resultant libraries

were submitted for paired-end 150 bp sequencing on a NEXTSeq500 (Illumina). For each

sample, transcript abundance was quantified from raw reads with Salmon (version 0.8.2)

(Patro *et al.*, 2017) using the reference human transcriptome (hg38) defined by GENCODE

release 27. An R package Tximport (version 1.4.0) (Soneson, Love and Robinson, 2015) was

used to estimate gene-level abundance from Salmon's transcript-level counts. Gene-level

differential expression analysis was performed using DESeq2 (version 1.16.1) (Love, Huber

and Anders, 2014) prior NMF analysis as described above.

***PHF3*, *CHD4*, *PCGF6*, and *AUTS2* shRNAs**

shRNAs against PHF3 (TRCN0000019118, TRCN0000019114, TRCN0000274376), CHD4

(TRCN0000380981, TRCN0000021363, TRCN0000021360), PCGF6 (TRCN0000229804,

TRCN0000073109, TRCN0000073109), AUTS2 (TRCN0000119058, TRCN0000304019,

TRCN0000304081), or non-targeting control (ATCTCGCTTGGGCGAGAGTAAG) were

cloned into pLKO5d.SFFV.miR30n (Schwarzer *et al.*, 2017). Each target gene was linked with

different fluorescent protein, including dTomato (PHF3 and AUTS2), eGFP (CHD4 and

PCGF6), and RFP657 (non-targeting control).

The vector contained BsmBI at the cloning sites. The oligonucleotides were designed with the

appropriate complementary overhang sequences of BsmBI-cleaved vector. They were ligated

and transformed into STBL3 chemically competent cells (Invitrogen) according to

manufacturer's instructions and plated on agar plates containing 100 µg/ml ampicillin

incubated for 16 h at 37°C. Single colonies were inoculated for DNA preparation, and extracted

using EndoFree Plasmid Maxi Kit (Qiagen, Cat#12362). All clones were verified by sequencing

prior to lentivirus production.

Lentivirus Production and Transduction

Viral particles were produced using calcium phosphate precipitation method on 293T cells by

co-transfecting an envelope plasmid pMD2.G, packaging plasmid pCMVΔR8.91, and the

shRNA vector pLKO5d.SFFV.miR30n. The cells were grown in 100 mm tissue culture dishes

at a concentration of 1-2 x 10^6 cells in 10 ml medium the day prior to co-transfection. On the

following day when the cells had reached 30-50% confluence, 20 µg of shRNA vector, 15 µg

of pCMVΔR8.91, and 5 µg of pMD2.G were mixed. The volume of the mixture was adjusted

with HEPES buffer solution (2.5 mM HEPES containing deionized water at pH7.3) to 250 μl.

A volume of 250 μl of 0.5 M $CaCl_2$ was added to the mixture. This solution was added to 500

μl of 2X HeBS (0.28 M NaCl, 0.05 M HEPES and 1.5 mM Na_2HPO_4 in deionized water at pH

7.00) and mixed by vortexing. It was incubated at RT for 30-40 min to allow the formation of

the calcium phosphate precipitate, before adding dropwise on the 293T cells. After 16 h, the

cells were washed with 10 ml PBS and added with 10 ml culture media. They were incubated

at 37°C in a humidified atmosphere with 5% CO_2 for the next two days. Lentivirus particles

were collected by centrifuging the supernatant at 400 g for 10 min at 4°C and filtered through

Acrodisc Syringe 0.45 filters. They were stored in aliquots at -80°C. Cell lines and PDX

samples were transduced with lentivirus as previously described (Bomken *et al.*, 2013).

Lymphoid differentiation of transduced *MLL-Af4* cord blood cells

MLL-Af4 cord blood cells (5 x 10^5 cells per transduction, Lin *et al.*, 2016) were transduced

with short hairpin constructs as described and co-cultured with MS-5 stroma cells in

lymphoid culture conditions described above. Single and triple transduced populations were

identified using the construct specific fluorophores and lineage specific surface markers

assessed as a proportion of the total transduced leukocyte population.

qRT-PCR

One million cells were collected and RNA was extracted using RNeasy Mini Kit (Qiagen,

Cat#74106). cDNA was synthesised from 1 μg RNA using RevertAid H Minus First Strand

cDNA Synthesis Kit (Thermo Fisher Scientific, Cat#K1632). The product was diluted by

adding 80 μl H_2O (Livak and Schmittgen, 2001). Sequences of primers used in this study were

designed using Primer Express (Applied Biosystems) software.

QUANTIFICATION AND STATISTICAL ANALYSIS

Reverse Engineering of Transcriptional Networks

A mutual information network was developed such that the connections between genes within

the network represented an inferred likelihood of a causal relationship within ALL/AML. The

network was further refined by retaining only those nodes (genes) and edges (causal

connections) which are significantly associated, in expression terms, to the distinction

between normal myeloid and lymphoid cells. Thus, the centrality of the mutations within the

network reflects the estimated extent of their causal influence upon a lymphoid/myeloid

distinction within the framework of primary ALL/AML.

Mutual Information networks were reverse engineered from 216 published AML/ALL

Affymetrix HGU133p2 expression profiles (GSE11877 & GSE24006) processed using Rma

(affy package R/Bioconductor) using the ARACNe2 algorithm (Margolin *et al.*, 2006) set to

adaptive partitioning with a DPI tolerance of 0.1 and a p-value threshold of 0.01 and 1000

bootstraps. The probes/genes were filtered to remove those that were insufficiently variable

as per the Aracne method. Probes had to be greater than 3 fold expression and 300 delta

between the min and max excluding the 5% most extreme values to be included. Probes

without >20 intensity in greater than 20% of samples were also removed. All probes which

fulfilled these requirements were included in the construction of this network (n=9780).

Nodes within the network were annotated with a metagene value calculated using Non-

Negative Matrix Factorisation (k=2) and reflecting the differences between myeloid and

lymphoid cells (GSE24759) and the other. A metagene cutoff of 0.13 (based on a calculated

alpha of 0.1) was applied to trim the network. Cytoscape (Shannon *et al.*, 2003) was used to

merge nodes where probe sets represented common genes and to calculate centrality

statistics for each of the mutated genes of interest. Nodes of interest (i.e. mutated nodes) were

ranked according to their centrality (e.g. degree). Nodes were sized and coloured according

to their differential expression between AML and ALL types; size = significance, colour = log2

fold change.

DATA AND SOFTWARE AVAILABILITY

Exome sequencing data and genome sequencing data presented in this manuscript have

been deposited in the NCBI Sequence Read Archive (SRA) under project numbers

PRJNA547947 and PRJNA547815 respectively. Immunoglobulin and TCR sequencing data

have been deposited in NCBI SRA under project number PRJNA511413. RNA sequencing

data and DNase hypersensitivity sequencing data were deposited in Gene Expression

Omnibus under accession numbers GSE132396 and GSE130142 respectively. All deposited

data will be publically available following publication of the manuscript.

1111

References

Agraz-Doblas, A., Bueno, C., Bashford-Rogers, R., Roy, A., Schneider, P., Bardini, M., Ballerini, P., Cazzaniga, G., Moreno, T., Revilla, C., Agraz-Doblas, A., *et al.* (2019). Unravelling the cellular origin and clinical prognostic markers of infant B-cell acute lymphoblastic leukemia using genome-wide analysis. Haematologica, 104, 1176-1188. 10.3324/haematol.2018.206375.

Alexander, T.B., Gu, Z., Iacobucci, I., Dickerson, K., Choi, J.K., Xu, B., Payne-Turner, D., Yoshihara, H., Loh, M.L., Horan, J., *et al*. (2018) The genetic basis and cell of origin of mixed phenotype acute leukaemia. Nature 562, 373-379. 10.1038/s41586-018-0436-0.

Alexandrov, L. B., Jones, P.H., Wedge, D.C., Sale, J.E., Campbell, P.J., Nik-Zainal, S., Stratton, M.R. (2015) Clock-like mutational processes in human somatic cells. Nature Genetics 47, 1402–1407. 10.1038/ng.3441.

Andersson, A.K., Ma, J., Wang, J., Chen, X., Gedman, A.L., Dang, J., Nakitandwe, J., Holmfeldt, L., Parker, M., Easton, J., *et al*. (2015). The landscape of somatic mutations in infant *MLL*-rearranged acute lymphoblastic leukemias. Nature Genetics 47, 330-337. 10.1038/ng.3230.

Appel, L-M., Franke, V., Bruno, M., Grishkovskaya, I., Kasiliauskaite, A., Schoeberl, U.E., Puchinger, M.G., Kostrhon, S., Beltzung, E., Mechtler, K., *et al*. (2020) PHF3 regulates neuronal gene expression through the new Pol II CTD reader domain SPOC. bioRxiv, p. 2020.02.11.943159. 10.1101/2020.02.11.943159.

Arends, T., Dege, C., Bortnick, A., Danhorn, T., Knapp, J.R., Jia, H., Harmacek, L., Fleenor, C.J., Straign, D., Walton, K., *et al*. (2019). CHD4 is essential for transcriptional repression and lineage progression in B lymphopoiesis. Proc Natl Acad Sci U S A 116, 10927-10936. 10.1073/pnas.1821301116.

Ayton, P. M., and Cleary, M. L. (2003). Transformation of myeloid progenitors by MLL oncoproteins is dependent on Hoxa7 and Hoxa9. Genes Dev 17, 2298–2307. 10.1101/gad.1111603.

Ferreira, B.I., García, J.F., Suela, J., Mollejo, M., Camacho, F.I., Carro, A., Montes, S., Piris, M.A., Cigudosa, J.C. (2008). Comparative genome profiling across subtypes of low-grade B-cell lymphoma identifies type-specific and common aberrations that target genes with a role in B-cell neoplasia. Haematologica 93, 670–679. 10.3324/haematol.12221.

Blokzijl, F. de Ligt, J., Jager, M., Sasselli, V., Roerink, S., Sasaki, N., Huch, M., Boymans, S., Kuijk, E., Prins, P. (2016) Tissue-specific mutation accumulation in human adult stem cells during life. Nature 538, 260–264. doi: 10.1038/nature19768.

Boer, J.M., van der Veer, A., Rizopoulos, D., Fiocco, M., Sonneveld, E., de Groot-Kruseman, H.A., Kuiper, R.P., Hoogerbrugge, P., Horstmann, M., Zaliova, M., *et al*. (2016). Prognostic value of rare IKZF1 deletion in childhood B-cell precursor acute lymphoblastic leukemia: an international collaborative study. Leukemia, 30, 32–38. 10.1038/leu.2015.199.

Bolotin, D.A., Poslavsky, S., Davydov, A.N., Frenkel, F.E., Fanchi, L., Zolotareva, O.I., Hemmers, S., Putintseva, E.V., Obraztsova, A.S., Shugay, M., *et al*. (2017). Antigen receptor repertoire profiling from RNA-seq data. Nat Biotechnol, 35, 908–911. 10.1038/nbt.3979.

Bolouri, H., Farrar, J.E., Triche, T.Jr., Ries, R.E., Lim, E.L., Alonzo, T.A., Ma, Y., Moore, R., Mungall, A.J., Marra, M.A., *et al*. (2018). The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational interactions. Nat Med, 24, 103–112.

10.1038/nm.4439 PMID - 29227476.

Bomken, S., Buechler, L., Rehe, K., Ponthan, F., Elder, A., Blair, H., Bacon, C.M., Vormoor, J.,
Heidenreich, O. (2013). Lentiviral marking of patient-derived acute lymphoblastic leukaemic cells
allows in vivo tracking of disease progression. Leukemia, 27, 718-721. 10.1038/leu.2012.206

Brunet, J.P., Tamayo, P., Golub, T.R., Mesirov, J.P. (2004). Metagenes and molecular pattern discovery
using matrix factorization. Proc Natl Acad Sci U S A 101, 4164–4169. 10.1073/pnas.0308531101.

Carulli, G., Marini, A., Ferreri, M.I., Azzarà, A., Ottaviano, V., Lari, T., Rocco, M., Giuntini, S., Petrini, M.
(2012). B-cell acute lymphoblastic leukemia with t(4;11)(q21;q23) in a young woman: evolution into
mixed phenotype acute leukemia with additional chromosomal aberrations in the course of therapy.
Hematol Rep 4, e15. 10.4081/hr.2012.e15.

Cauchy, P., James, S.R., Zacarias-Cabeza, J., Ptasinska, A., Imperato, M.R., Assi, S.A., Piper, J.,
Canestraro, M., Hoogenkamp, M., Raghavan, M., *et al*. (2015). Chronic FLT3-ITD Signaling in Acute
Myeloid Leukemia Is Connected to a Specific Chromatin Signature. Cell Rep 12, 821–836.
10.1016/j.celrep.2015.06.069.

Chen, L., Kostadima, M., Martens, J.H.A., Canu, G., Garcia, S.P., Turro, E., Downes, K., Macaulay, I.C.,
Bielczyk-Maczynska, E., Coe, S., *et al*. (2014). Transcriptional diversity during lineage commitment of
human blood progenitors. Science 345, 1251033. 10.1126/science.1251033.

Chen, X., Schulz-Trieglaff, O., Shaw, R., Barnes, B., Schlesinger, F., Källberg, M., Cox, A.J., Kruglyak, S.,
Saunders, C.T. (2016). Manta: rapid detection of structural variants and indels for germline and cancer
sequencing applications. Bioinformatics 32, 1220–1222. 10.1093/bioinformatics/btv710.

Church, D.N., Briggs, S.E., Palles, C., Domingo, E., Kearsey, S.J., Grimes, J.M., Gorman, M., Martin, L.,
Howarth, K.M., Hodgson, S.V., *et al*. (2013). DNA polymerase ε and δ exonuclease domain mutations
in endometrial cancer. Hum Mol Genet 22, 2820–2828. 10.1093/hmg/ddt131.

Cibulskis, K., Lawrence, M.S., Carter, S.L., Sivachenko, A., Jaffe, D., Sougnez, C., Gabriel, S., Meyerson,
M., Lander, E.S., Getz, G. (2013). Sensitive detection of somatic point mutations in impure and
heterogeneous cancer samples. Nat Biotechnol 31, 213–219. 10.1038/nbt.2514.

DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel,
G., Rivas, M.A., Hanna, M., *et al*. (2011). A framework for variation discovery and genotyping using
next-generation DNA sequencing data. Nat Genet 43, 491–498. 10.1038/ng.806.

Dobbins, S.E., Sherborne, A.L., Ma, Y.P., Bardini, M., Biondi, A., Cazzaniga, G., Lloyd, A., Chubb, D.,
Greaves, M.F., Houlston, R.S. (2013). The silent mutational landscape of infant MLL-AF4 pro-B acute
lymphoblastic leukemia. Genes Chromosomes Cancer 52, 954–960. 10.1002/gcc.22090.

Erson-Omay, E.Z., Çağlayan, A.O., Schultz, N., Weinhold, N., Omay, S.B., Özduman, K., Köksal, Y., Li, J.,
Serin Harmancı, A., Clark, V., *et al*. (2015). Somatic POLE mutations cause an ultramutated giant cell
high-grade glioma subtype with better prognosis. Neuro Oncol 17, 1356–1364.
10.1093/neuonc/nov027.

Gardner, R., Wu, D., Cherian, S., Fang, M., Hanafi, L.A., Finney, O., Smithers, H., Jensen, M.C., Riddell,
S.R., Maloney, D.G., *et al*. (2016). Acquisition of a CD19-negative myeloid phenotype allows immune
escape of *MLL*-rearranged B-ALL from CD19 CAR-T-cell therapy. Blood 127, 2406-2410.
10.1182/blood-2015-08-665547.

Germano, G., Pigazzi, M., del Giudice, L., Campo Dell'Orto, M., Spinelli, M., Zangrando, A., Paolucci, P.,
Ladogana, S., Basso, G. (2006). Two consecutive immunophenotypic switches in a child with MLL-
rearranged acute lymphoblastic leukemia. Haematologica 91, ECR09.

Gentles, A.J., Plevritis, S.K., Majeti, R., Alizadeh, A.A. (2010). Association of a leukemic stem cell gene
expression signature with clinical outcomes in acute myeloid leukemia. JAMA 304, 2706–2715.
10.1001/jama.2010.1862.

Gessner, A., Thomas, M., Castro, P.G., Büchler, L., Scholz, A., Brümmendorf, T.H., Soria, N.M., Vormoor,
J., Greil, J., Heidenreich, O. (2010). Leukemic fusion genes MLL/AF4 and AML1/MTG8 support leukemic
self-renewal by controlling expression of the telomerase subunit TERT. Leukemia 24, 1751–1759.
10.1038/leu.2010.155.

González-Pérez, A., and López-Bigas, N. (2011). Improving the assessment of the outcome of
nonsynonymous SNVs with a consensus deleteriousness score, Condel', Am J Hum Genet 88, 440–449.
10.1016/j.ajhg.2011.03.004.

Grinev, V.V., Barneh, F., Ilyushonak, I.M., Nakjang, S., Smink, J., van Oort, A., Clough, R., Seyani, M.,
McNeill, H., Reza, M., *et al*. (2021). RUNX1/RUNX1T1 mediates alternative splicing and reorganises the
transcriptional landscape in leukemia. Nat Commun 12, 520. 10.1038/s41467-020-20848-z.

Hamieh, M., Dobrin, A., Cabriolu, A., van der Stegen, S.J.C., Giavridis, T., Mansilla-Soto, J., Eyquem, J.,
Zhao, Z., Whitlock, B.M., Miele, M.M., *et al*. (2019). CAR T cell trogocytosis and cooperative killing
regulate tumour antigen escape. Nature 568, 112–116. 10.1038/s41586-019-1054-1.

Harman, J.R., Thorne, R., Jamilly, M., Tapia, M., Crump, N.T., Rice, S., Beveridge, R., Morrissey, E., de
Bruijn, M.F.T.R., Roberts, I., *et al*. (2021) A KMT2A-AFF1 gene regulatory network highlights the role
of core transcription factors and reveals the regulatory logic of key downstream target genes. Genome
Res, Jun 4 online ahead of print. doi: 10.1101/gr.268490.120.

Harvey, R.C., Mulligan, C.G., Wang, X., Dobbin, K.K., Davidson, G.S., Bedrick, E.J., Chen, I.M., Atlas,
S.R., Kang, H., Ar, K. *et al*. (2010). Identification of novel cluster groups in pediatric high-risk B-
precursor acute lymphoblastic leukemia with gene expression profiling: correlation with genome-wide
DNA copy number alterations, clinical characteristics, and outcome. Blood. 116, 4874–4884.
10.1182/blood-2009-08-239681.

Heshmati, Y., Türköz, G., Harisankar, A., Kharazi, S., Boström, J., Dolatabadi, E.K., Krstic, A., Chang, D.,
Månsson, R., Altun, M., *et al*. (2018). The chromatin-remodeling factor CHD4 is required for
maintenance of childhood acute myeloid leukemia. Haematologica 103, 1169-1181.
10.3324/haematol.2017.183970

Hohaus, S., Petrovick, M.S., Voso, M.T., Sun, Z., Zhang, D.E., Tenen, D.G. (1995). PU.1 (Spi-1) and C/EBP
alpha regulate expression of the granulocyte-macrophage colony-stimulating factor receptor alpha
gene. Mol Cell Biol 15, 5830–5845. 10.1128/mcb.15.10.5830.

Hosokawa, H., Tanaka, T., Suzuki, Y., Iwamura, C., Ohkubo, S., Endoh, K., Kato, M., Endo, Y., Onodera,
A., Tumes, D.J., *et al*. (2013). Functionally distinct Gata3/Chd4 complexes coordinately establish T
helper 2 (Th2) cell identity. Proc Natl Acad Sci U S A 110, 4691–4696. 10.1073/pnas.1220865110.

Ivanov, I.C., Jitam, D., Grigore, G.E., Zlei, M., Ivanoc, A.V., Dumitras, S., Carasevici, E., Miron, I.C. (2013)
Infant acute leukemia with lineage switch at relapse expressing a novel t(4;11)(q21;q23) MLL-AF4
fusion transcript. Romanian Review of Laboratory Medicine 21, 47–58. 10.2478/rrlm-2013-0017.

Jiang, J.G., Roman, E., Nandula, S.V., Murty, V.V., Bhagat, G., Alobeid, B. (2005). Congenital MLL-
positive B-cell acute lymphoblastic leukemia (B-ALL) switched lineage at relapse to acute myelocytic
leukemia (AML) with persistent t(4;11) and t(1;6) translocations and JH gene rearrangement. Leuk
Lymphoma 46, 1223-1227. 10.1080/10428190500086055.

Jung, R., Jacobs, U., Krumbholz, M., Langer, T., Keller, T., De Lorenzo, P., Valsecchi, M.G., van der
Velden, V.H., Moericke, A., Stanulla, M., *et al*. (2010). Bimodal distribution of genomic MLL
breakpoints in infant acute lymphoblastic leukemia treatment. Leukemia 24, 903–907.
10.1038/leu.2010.14.

Kang, H., Chen, I.M., Wilson, C.S., Bedrick, E.J., Harvey, R.C., Atlas, S.R., Devidas, M., Mullighan, C.G.,
Wang, X., Murphy, M., *et al*. (2010). Gene expression classifiers for relapse-free survival and minimal
residual disease improve risk classification and outcome prediction in pediatric B-precursor acute
lymphoblastic leukemia. Blood 115, 1394–1405. 10.1182/blood-2009-05-218560.

Kerry, J., Godfrey, L., Repapi, E., Tapia, M., Blackledge, N.P., Ma, H., Ballabio, E., O'Byrne, S., Ponthan,
F., Heidenreich, O., *et al*. (2017). MLL-AF4 Spreading Identifies Binding Sites that Are Distinct from
Super-Enhancers and that Govern Sensitivity to DOT1L Inhibition in Leukemia. Cell Rep 18, 482–495.
10.1016/j.celrep.2016.12.054.

Kim, J., Sif, S., Jones, B., Jackson, A., Koipally, J., Heller, E., Winandy, S., Viel, A., Sawyer, A., Ikeda, T.,
et al. (1999). Ikaros DNA-Binding Proteins Direct Formation of Chromatin Remodeling Complexes in
Lymphocytes. Immunity 10, 345-355. 10.1016/s1074-7613(00)80034-5.

Kinkelin, K., Wozniak, G.G., Rothbart, S.B., Lidschreiber, M., Strahl, B.D., Cramer, P. (2013). Structures
of RNA polymerase II complexes with Bye1, a chromatin-binding PHF3/DIDO homologue. Proc Natl
Acad Sci U S A 110, 15277–15282. 10.1073/pnas.1311010110.

Langmead, B., Trapnell, C., Pop, M., Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biol 10, R25. 10.1186/gb-2009-10-3-r25.

Leddin, M., Perrod, C., Hoogenkamp, M., Ghani, S., Assi, S., Heinz, S., Wilson, N.K., Follows, G.,
Schönheit, J., Vockentanz, L., *et al*. (2011). Two distinct auto-regulatory loops operate at the PU.1 locus
in B cells and myeloid cells. Blood 117, 2827–2838. 10.1182/blood-2010-08-302976.

le Viseur, C., Hotfilder, M., Bomken, S., Wilson, K., Röttgers, S., Schrauder, A., Rosemann, A., Irving, J.,
Stam, R.W., Shultz, L.D., *et al*. (2008) In childhood acute lymphoblastic leukemia, blasts at different
stages of immunophenotypic maturation have stem cell properties. Cancer Cell, 14, 47–58.
10.1016/j.ccr.2008.05.015

Li, B., Brady, S.W., Ma, X., Shen, S., Zhang, Y., Li, Y., Szlachta, K., Dong, L., Liu, Y., Yang, F., *et al*. (2020).
Therapy-induced mutations drive the genomic landscape of relapsed acute lymphoblastic leukemia.
Blood 135, 41–55. 10.1182/blood.2019002220.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics 25, 1754–1760. 10.1093/bioinformatics/btp324.

Lin, S., Luo, R.T., Ptasinska, A., Kerry, J., Assi, S.A., Wunderlich, M., Imamura, T., Kaberlein, J.J., Rayes,
A., Althoff, M.J., *et al*. (2016). Instructive Role of MLL-Fusion Proteins Revealed by a Model of t(4;11)
Pro-B Acute Lymphoblastic Leukemia. Cancer Cell 30, 737–749. 10.1016/j.ccell.2016.10.008.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. Methods 25, 402–408. 10.1006/meth.2001.1262.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15, 550. 10.1186/s13059-014-0550-8.

Lu, X., Chu, C.S., Fang, T., Rayon-Estrada, V., Fang, F., Patke, A., Qian, Y., Clarke, S.H., Melnick, A.M.,
Zhang, Y., *et al*. (2019). MTA2/NuRD Regulates B Cell Development and Cooperates with OCA-B in
Controlling the Pre-B to Immature B Cell Transition. Cell Rep 28, 472-485.
10.1016/j.celrep.2019.06.029.

Malouf, C., and Ottersbach, K. (2018). The fetal liver lymphoid-primed multipotent progenitor
provides the prerequisites for the initiation of t(4;11) MLL-AF4 infant leukemia. Haematologica, 103,
e571–e574. 10.3324/haematol.2018.191718.

Margolin, A.A., Nemenman, I., Basso, K., Wiggins, C., Stolovitzky, G., Dalla Favera, R., Califano, A.
(2006). ARACNE: An Algorithm for the Reconstruction of Gene Regulatory Networks in a Mammalian
Cellular Context. BMC Bioinformatics 7, S7. doi: 10.1186/1471-2105-7-S1-S7.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler,
D., Gabriel, S., Daly, M., DePristo, M.A. (2010). The genome analysis toolkit: A MapReduce framework
for analyzing next-generation DNA sequencing data. Genome Res 20, 1297–1303.
10.1101/gr.107524.110.

Meyer, C., Schneider B., Dingermann, T., Klingebiel, T., Marschalek, R. (2005). A LDI-PCR based method
allows the identification of any MLL rearrangement. Blood 106, 2848.
10.1182/blood.V106.11.2848.2848.

Meyer, C., Kowarz, E., Hofmann, J., Renneville, A., Zuna, J., Trka, J., Ben Abdelali, R., Macintyre, E., De
Braekeleer, E., De Braekeleer, M., *et al*. (2009). New insights to the MLL recombinome of acute
leukemias. Leukemia 23, 1490–1499. 10.1038/leu.2009.33.

Meyer, C., Hofmann, J., Burmeister, T., Gröger, D., Park, T.S., Emerenciano, M., Pombo de Oliveira, M.,
Renneville, A., Villarese, P., Macintyre, E., *et al*. (2013). The MLL recombinome of acute leukemias in
2013. Leukemia 27, 2165–2176. 10.1038/leu.2013.135.

Meyer, C., Burmeister, T., Gröger, D., Tsaur, G., Fechina, L., Renneville, A., Sutton, R., Venn, N.C.,
Emerenciano, M., Pombo-de-Oliveira, M.S. (2018). The MLL recombinome of acute leukemias in 2017.
Leukemia 32, 273–284. 10.1038/leu.2017.213.

Moorman, A.V., Ensor, H.M., Richards, S.M., Chilton, L., Schwab, C., Kinsey, S.E., Vora, A., Mitchell,
C.D., Harrison, C.J. (2010). Prognostic effect of chromosomal abnormalities in childhood B-cell
precursor acute lymphoblastic leukaemia: results from the UK Medical Research Council ALL97/99
randomised trial. Lancet Oncol 11, 429–438. 10.1016/S1470-2045(10)70066-8.

Mor, N., Rais, Y., Sheban, D., Peles, S., Aguilera-Castrejon, A., Zviran, A., Elinger, D., Viukov, S., Geula,
S., Krupalnik, V., *et al*. (2018). Neutralizing Gatad2a-Chd4-Mbd3/NuRD Complex Facilitates
Deterministic Induction of Naive Pluripotency. Cell Stem Cell 23, 412-425.
10.1016/j.stem.2018.07.004.

Mullighan, C.G., Goorha, S., Radtke, I., Miller, C.B., Coustan-Smith, E., Dalton, J.D., Girtman, K.,
Mathew, S., Ma, J., Pounds, S.B., *et al*. (2007). Genome-wide analysis of genetic alterations in acute
lymphoblastic leukaemia. Nature 446, 758–764. 10.1038/nature05690.

Mur, P., García-Mulero, S., Del Valle, J., Magraner-Pardo, L., Vidal, A., Pineda, M., Cinnirella, G., Martín-
Ramos, E., Pons, T., López-Doriga, A., *et al*. (2020). Role of POLE and POLD1 in familial cancer. Genet

Med 22, 2089–2100. 10.1038/s41436-020-0922-2.

Nerlov, C., and Graf, T. (1998). PU.1 induces myeloid lineage commitment in multipotent
hematopoietic progenitors. Genes Dev 12, 2403–2412. 10.1101/gad.12.15.2403.

Ng, S.Y., Yoshida, T., Zhang, J., Georgopoulos, K. (2009). Genome-wide lineage-specific transcriptional
networks underscore Ikaros-dependent lymphoid priming in hematopoietic stem cells. Immunity 30,
493–507. 10.1016/j.immuni.2009.01.014.

Novershtern, N., Subramanian, A., Lawton, L.N., Mak, R.H., Haining, W.N., McConkey, M.E., Habib, N.,
Yosef, N., Chang, C.Y., Shay, T., *et al*. (2011). Densely Interconnected Transcriptional Circuits Control
Cell States in Human Hematopoiesis. Cell 144, 296–309. 10.1016/j.cell.2011.01.004.

Nutt, S.L., Urbánek, P., Rolink, A., Busslinger, M. (1997). Essential functions of Pax5 (BSAP) in pro-B cell
development: difference between fetal and adult B lymphopoiesis and reduced V-to-DJ recombination
at the IgH locus. Genes Dev 11, 476–491. 10.1101/gad.11.4.476.

Nutt, S.L., Heavey, B., Rolink, A.G., Busslinger, M. (1999). Commitment to the B-lymphoid lineage
depends on the transcription factor Pax5. Nature 402, 14–20. 10.1038/35005514.

O'Neill, D. W., Schoetz, S.S., Lopez, R.A., Castle, M., Rabinowitz, L., Shor, E., Krawchuk, D., Goll, M.G.,
Renz, M., Seelig, H.P., Han, S., *et al*. (2000). An ikaros-containing chromatin-remodeling complex in
adult-type erythroid cells. Mol Cell Biol 20, 7572–7582. 10.1128/mcb.20.20.7572-7582.2000.

Orlando, E.J., Han, X., Tribouley, C., Wood, P.A., Leary, R.J., Riester, M., Levine, J.E., Qayed, M., Grupp,
S.A., Boyer, M., *et al*. (2018). Genetic mechanisms of target antigen loss in CAR19 therapy of acute
lymphoblastic leukemia. Nat Med 24, 1504–1506. 10.1038/s41591-018-0146-z.

Park, M., Koh, K.N., Kim, B.E., Im, H.J., Jang, S., Park, C.J., Chi, H.S., Seo, J.J. (2011). Lineage Switch at
Relapse of Childhood Acute Leukemia: A Report of Four Cases. J Korean Med Sci 26, 829–831.
10.3346/jkms.2011.26.6.829.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., Kingsford, C. (2017). Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods 14, 417–419. 10.1038/nmeth.4197.

Pieters, R., Schrappe, M., De Lorenzo, P., Hann, I., De Rossi, G., Felice, M., Hovi, L., LeBlanc, T.,
Szczepanski, T., Ferster, A., *et al*. (2007). A treatment protocol for infants younger than 1 year with
acute lymphoblastic leukaemia (Interfant-99): an observational study and a multicentre randomised
trial. Lancet 370, 240–250. 10.1016/S0140-6736(07)61126-X.

Piper, J., Elze, M.C., Cauchy, P., Cockerill, P.N., Bonifer, C., Ott, S. (2013). Wellington: a novel method
for the accurate identification of digital genomic footprints from DNase-seq data. Nucleic Acids
Research 41, e201–e201. 10.1093/nar/gkt850.

Piper, J., Assi, S.A., Cauchy, P., Ladroue, C., Cockerill, P.N., Bonifer, C., Ott, S. (2015). Wellington-
bootstrap: differential DNase-seq footprinting identifies cell-type determining transcription factors.
BMC Genomics 16, 1000. 10.1186/s12864-015-2081-4.

Pongubala, J.M., Northrup, D.L., Lancki, D.W., Medina, K.L., Treiber, T., Bertolino, E., Thomas, M.,
Grosschedl, R., Allman, D., Singh, H. (2008). Transcription factor EBF restricts alternative lineage
options and promotes B cell fate commitment independently of Pax5. Nat Immunol 9, 203–215.
10.1038/ni1555.

Ptasinska, A., Assi, S.A., Martinez-Soria, N., Imperato, M.R., Piper, J., Cauchy, P., Pickin, A., James, S.R., Hoogenkamp, M., Willimason, D. *et al*. (2014). Identification of a Dynamic Core Transcriptional Network in t(8;21) AML that Regulates Differentiation Block and Self-Renewal. Cell Rep 8, 1974–1988. 10.1016/j.celrep.2014.08.024 .

Rabilloud, T.,Potier, D., Pankaew, S., Nozais, M., Loosveld, M., Payet-Bornet, D. (2021). Single-cell profiling identifies pre-existing CD19-negative subclones in a B-ALL patient with CD19-negative relapse after CAR-T therapy. Nature Commun 12, 865. 10.1038/s41467-021-21168-6.

Raczy, C., Petrowski, R., Saunders, C.T., Chorny, I., Kruglyak, S., Margulies, E.H., Chuang, H.-Y., Källberg, M., Kumar, S.A., Liao, A . (2013). Isaac: Ultra-fast whole-genome secondary analysis on Illumina sequencing platforms. Bioinformatics 29, 2041–2043. 10.1093/bioinformatics/btt314.

Rayes, A., McMasters, R.L., O'Brien, M.M. (2016). Lineage switch in MLL-rearranged infant leukemia following CD19-directed therapy. Pediatr Blood Cancer 63, 1113-510. 10.1002/pbc.25953

Rehe, K. Wilson, K., Bomken, S., Williamson, D., Irving, J., den Boer, M.L., Stanulla, M., Schrappe, M., Hall, A.G., Heidenreich, O., Vormoor, J.(2013). Acute B lymphoblastic leukaemia-propagating cells are present at high frequency in diverse lymphoblast populations. EMBO Mol Med 5, 38–51. 10.1002/emmm.201201703.

Risso, D. Ngai, J., Speed, T.P., Dudoit, S. (2014). Normalization of RNA-seq data using factor analysis of control genes or samples.Nature Biotechnol 32, 896–902. 10.1038/nbt.2931.

Rongvaux, A. Willinger, T., Martinek, J., Strowig, T., Gearty, S.V., Teichmann, L.L., Saito, Y., Marches, F., Halene, S., Palucka, A.K. *et al*. (2014). Development and function of human innate immune cells in a humanized mouse model.Nature Biotechnol 32, 364–372. 10.1038/nbt.2858.

Rossi, J.G., Andrea R Bernasconi, Cristina N Alonso, Patricia L Rubio, Marta S Gallego, Carolina A Carrara, Myriam R Guitter, Silvia Eandi Eberle, Mariela Cocce, Pedro A Zubizarreta, *et al*. (2012). Lineage switch in childhood acute leukemia: An unusual event with poor outcome. Am J Hematol 87, 890–897. 10.1002/ajh.23266.

Saldanha, A. J. (2004). Java Treeview - Extensible visualization of microarray data. Bioinformatics 20, 3246–3248.10.1093/bioinformatics/bth349.

Saunders, C.T. Wong, W.S.W., Swamy, S., Becq, J., Murray, L.J., Cheetham, R.K. (2012). Strelka: Accurate somatic small-variant calling from sequenced tumor-normal sample pairs. Bioinformatics 28, 1811–1817. 10.1093/bioinformatics/bts271.

Schwalbe, E.C., Williamson, D., Lindsey, J.C., Hamilton, D., Ryan, S.L., Megahed, H., Garami, M., Hauser, P., Dembowska-Baginska, B., Perek, D. *et al*. (2013). DNA methylation profiling of medulloblastoma allows robust subclassification and improved outcome prediction using formalin-fixed biopsies. Acta Neuropathol 125, 359–371. 10.1007/s00401-012-1077-2.

Schwarzer, A. Emmrich, S., Schmidt, F., Beck, D., Ng, M., Reimer, C., Adams, F.F., Grasedieck, S., Witte, D., Käbler, S. *et al*. (2017). The non-coding RNA landscape of human hematopoiesis and leukemia. Nature Commun 8, 218.10.1038/s41467-017-00212-4.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Research 13, 2498–2504. 10.1101/gr.1239303.

Sifrim, A. Hitz, M.-P., Wilsdon, A., Breckpot, J., Al Turki, S.H., Thienpont, B., McRae, J., Fitzgerald,
T.W.F., Singh, T., Swaminathan, G.J. *et al*. (2016). Distinct genetic architectures for syndromic and
nonsyndromic congenital heart defects identified by exome sequencing. Nature Gen 48, 1060–1065.
10.1038/ng.3627.

Somervaille, T.C.P., Christina J Matheny, C.J., Gary J Spencer, G.J., Masayuki Iwasaki, M., John L Rinn,
J.L., Daniela M Witten, D.M., Howard Y Chang, H.Y., Sheila A Shurtleff, S.A., James R Downing, J.R.,
Michael L Cleary, M.L. (2009). Hierarchical maintenance of MLL myeloid leukemia stem cells employs
a transcriptional program shared with embryonic rather than adult stem cells. Cell Stem Cell 4, 129–
140. 10.1016/j.stem.2008.11.015.

Soneson, C., Love, M.I., Robinson, M.D. (2015). Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000 Research 4, 1521. 10.12688/f1000research.7563.

Sotillo, E., Barrett, D.M., Black, K.L., Bagashev, A., Oldridge, D., Wu, G., Sussman, R., Lanauze, C., Ruella,
M., Gazzara, M.R., *et al*. (2015). Convergence of Acquired Mutations and Alternative Splicing of CD19
Enables Resistance to CART-19 Immunotherapy. Cancer Discovery 5, 1282–1295. 10.1158/2159-
8290.CD-15-1020.

Sperlazza, J., Rahmani, M., Beckta, J., Aust, M., Hawkins, E., Wang, S., Zhu, S., Podder, S., Dumur, C.,
Archer, K., *et al*. (2015). Depletion of the chromatin remodeler CHD4 sensitizes AML blasts to
genotoxic agents and reduces tumor formation. Blood 126, 1462–1472. 10.1182/blood-2015-03-
631606.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,
Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: A knowledge-
based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A 102,
15545–15550. 10.1073/pnas.0506580102

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva,
N.T., Morris, J.H., Bork, P., *et al*. (2019). STRING v11: protein-protein association networks with
increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic
Acids Research 47, D607–D613. 10.1093/nar/gky1131.

Tamayo, P., Scanfeld, D., Ebert, B.L., Gillette, M.A., Roberts, C.W.M., Mesirov, J.P. (2007). Metagene
projection for cross-platform, cross-species characterization of global transcriptional states. . Proc Natl
Acad Sci U S A 104, 5959 LP – 5964. 10.1073/pnas.0701068104.

van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., Jordan,
T., Shakir, K., Roazen, D., Thibault, J., *et al*. (2013). From FastQ data to high-confidence variant calls:
the Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 43, 11.10.1-11.10.33.
10.1002/0471250953.bi1110s43.

van Dongen, J.J.M., Langerak, A.W., Brüggemann, M., Evans, P.A.S., Hummel, M., Lavender, F.L.,
Delabesse, E., Davi, F., Schuuring, E., García-Sanz, R. *et al*. (2003). Design and standardization of PCR
primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene
recombinations in suspect lymphoproliferations: Report of the BIOMED-2 Concerted Action BMH4-
CT98-3936. Leukemia 17, 2257–2317. 10.1038/sj.leu.2403202.

Wang, J., Mi, J.-Q., Debernardi, A., Vitte, A.-L., Emadali, A., Meyer, J.A., Charmpi, K., Ycart, B., Callanan,
M.B., Carroll, W.L., *et al*. (2015). A six gene expression signature defines aggressive subtypes and
predicts outcome in childhood and adult acute lymphoblastic leukemia. Oncotarget 6, 16527–16542.
10.18632/oncotarget.4113.

Wilkinson, A.C., Ballabio, E., Geng, H., North, P., Tapia, M., Kerry, J., Biswas, D., Roeder, R.G., Allis, C.D.,
Melnick, A., *et al*. (2013). RUNX1is a key target in t(4;11) leukemias that contributes to gene activation
through an AF4-MLL complex interaction. Cell Reports 3, 116–127. 10.1016/j.celrep.2012.12.016.

Williams, C.J., Naito, T., Arco, P.G., Seavitt, J.R., Cashman, S.M., De Souza, B., Qi, X., Keables, P., Von
Andrian, U.H., Georgopoulos, K. (2004). The chromatin remodeler Mi-2β is required for CD4
expression and T cell development. Immunity 20, 719–733. 10.1016/j.immuni.2004.05.005.

Witkowski, M.T., Hu, Y., Roberts, K.G., Boer, J.M., McKenzie, M.D., Liu, G.J., Le Grice, O.D., Tremblay,
C.S., Ghisi, M., Willson, T.A., *et al*. (2017). Conserved IKAROS-regulated genes associated with B-
progenitor acute lymphoblastic leukemia outcome. J Exp Med 214, 773–791. 10.1084/jem.20160048.

Yoshida, T., Ng, S.Y.-M., Zuniga-Pflucker, J.C., and Georgopoulos, K. (2006). Early hematopoietic
lineage restrictions directed by Ikaros. Nature Immunology 7, 382–391. 10.1038/ni1314.

Yoshida, T., Hazan, I., Zhang, J., Ng, S.Y., Naito, T., Snippert, H.J., Heller, E.J., Qi, X., Lawton, L.N.,
Williams, C.J., *et al*. (2008).The role of the chromatin remodeler Mi-2beta in hematopoietic stem cell
self-renewal and multilineage differentiation. Genes Dev 22, 1174–1189. 10.1101/gad.1642808.

Zangrando, A., Dell'orto, M.C., Te Kronnie, G., Basso, G. (2009). MLL rearrangements in pediatric acute
lymphoblastic and myeloblastic leukemias: MLL specific and lineage specific signatures. BMC Med
Genomics 236, 10.1186/1755-8794-2-36.

Zhang, J., Jackson, A.F., Naito, T., Dose, M., Seavitt, J., Liu, F., Heller, E.J., Kashiwagi, M., Yoshida, T.,
Gounari, F., et al. (2011). Harnessing of the nucleosome-remodeling-deacetylase complex controls
lymphocyte development and prevents leukemogenesis. Nature Immunology 13, 86–94.
10.1038/ni.2150.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M.,
Brown, M., Li, W., et al. (2008). Model-based Analysis of ChIP-Seq (MACS). Genome Biology 9, R137.
10.1186/gb-2008-9-9-r137.

1461
1462

Figure 1

A

Presentation ALL   **Relapse AML**

B

MLL intron 9   Filler   AF4 intron 4   MLL intron 9   Filler   AF4 intron 4

ALL

AML

der11; *MLL-AF4*   **der4; *AF4-MLL***

C

Presentation ALL

Relapse AML

D

MLL
Ex10   Ex11
Ex9   Ex12
Ex8   Ex13
Ex7
Ex3
Ex7
Ex6
Ex4
Ex5
AF4

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

A

B

motif	match	p-value	% FP
AACAGGAAGT	ETS	1e-516	13.6
AAACCACA	RUNX	1e-378	10.8
GTCCCAGGCAACTC	EBF	1e-984	10.30
CAGCTG	Ebox	1e-113	9.9
ATGAGTCATT	AP1	1e-640	6.7
CACGTGGC	Ebox	1e-71	4.9
ATGCATGACTGATT	PAX	1e-146	1.6

Figure S7

Figure S8