Epigenetic regulator genes direct lineage switching in *MLL-AF4*

leukaemia

Ricky Tirtakusuma[1], Katarzyna Szoltysek[1,2,3], Paul Milne[4], Vasily V Grinev[5], Anetta

Ptasinska[6], Claus Meyer[7], Sirintra Nakjang[1], Jayne Y Hehir-Kwa[2], Daniel Williamson[1], Pierre

Cauchy[6], Salam A Assi[6], Maria R Imperato[6], Fotini Vogiatzi[9], Shan Lin[10], Mark Wunderlich[10],

Janine Stutterheim[2], Alexander Komkov[8], Elena Zerkalenkova[8], Paul Evans[11], Hesta

McNeill[1], Alex Elder[1], Natalia Martinez-Soria[1], Sarah E Fordham[1], Yuzhe Shi[1], Lisa J

Russell[1], Deepali Pal[1], Alex Smith[12], Zoya Kingsbury[13], Jennifer Becq[13], Cornelia Eckert[14],

Oskar A Haas[15], Peter Carey[16], Simon Bailey[1,16], Roderick Skinner[1,16], Natalia Miakova[8],

Matthew Collin[4], Venetia Bigley[4], Muzlifah Haniffa[17,18,19], Rolf Marschalek[7], Christine J

Harrison[1], Catherine A Cargo[11], Denis Schewe[9], Yulia Olshanskaya[8], Michael J Thirman[20],

Peter N Cockerill[6], James C Mulloy[10], Helen J Blair[1], Josef Vormoor[1,2], James M Allan[1],

Constanze Bonifer[6*], Olaf Heidenreich[1,2*†], Simon Bomken[1,16*†]

Author Affiliations

[1]Wolfson Childhood Cancer Research Centre, Centre for Cancer, Translational and Clinical
Research Institute, Newcastle University, Newcastle upon Tyne, UK

[2]Princess Maxima Center for Pediatric Oncology, Utrecht, The Netherlands

[3]Maria Sklodowska-Curie Institute - Oncology Center, Gliwice Branch, Gliwice, Poland

[4]Translational and Clinical Research Institute, Newcastle University, Framlington Place,
Newcastle upon Tyne, UK

[5]Department of Genetics, the Faculty of Biology, Belarusian State University, 220030 Minsk,
Republic of Belarus.

[6]Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK

[7]Institute of Pharmaceutical Biology/DCAL, Goethe-University, Frankfurt/Main, Germany

[8]Dmitry Rogachev National Research Center of Pediatric Hematology, Oncology, and
Immunology; Moscow, Russia

[9]Pediatric Hematology/Oncology, ALL-BFM Study Group, Christian Albrechts University Kiel
and University Hospital Schleswig-Holstein, Campus Kiel, Germany

[10]Experimental Hematology and Cancer Biology, Cancer and Blood Disease Institute,
Cincinnati Children's Hospital Medical Center, Cincinnati, USA

[11]Haematological Malignancy Diagnostic Service, St James's University Hospital, Leeds, UK

[12]Epidemiology and Cancer Statistics Group, University of York, York, United Kingdom

[13]Illumina Cambridge Ltd., Great Abington, UK

[14]Department of Pediatric Oncology/Hematology, Charité Universitätsmedizin Berlin, Berlin,
Germany

[15]Children's Cancer Research Institute, St. Anna Kinderkrebsforschung, Vienna, Austria

[16]Department of Paediatric Haematology and Oncology, The Great North Children's
Hospital, Newcastle upon Tyne, UK

[17]Biosciences Institute, Newcastle University, Framlington Place, Newcastle upon Tyne, UK

[18]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton UK

[19]Department of Dermatology and Newcastle NIHR Newcastle Biomedical Research Centre,
Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne

[20]Department of Medicine, Section of Hematology/Oncology, University of Chicago, Chicago,
USA

*Co-senior authors
†**Co-corresponding Authors.**
47

Running title: Lineage switching in *MLL-AF4* leukaemias

Keywords: *MLL-AF4*; *KMT2A-AFF1*; acute lymphoblastic leukaemia; nucleosome

remodelling and deactylation complex (NuRD); chromatin remodelling

Corresponding authors

Dr Simon Bomken
Wolfson Childhood Cancer Research Centre
Translational and Clinical Research Institute
Level 6 Herschel Building
Brewery Lane
Newcastle University
Newcastle upon Tyne
NE1 7RU, UK
60
61   Tel: +44 (0)191 2082231
62   E mail: s.n.bomken@ncl.ac.uk
63
64   Professor Olaf Heidenreich
65   Princess Maxima Center for Pediatric Oncology
66   Heidelberglaan 25
67   3584 CS Utrecht
68   The Netherlands
69
70   Tel: +31 (0)88 972 7272
71   E mail: O.T.Heidenreich@prinsesmaximacentrum.nl
72

Conflict of interest statement

Z.K. and J.B. are employees of Illumina, a public company that develops and markets

systems for genetic analysis. The remaining authors declare no potential conflicts of interest.

Financial support

This study was supported by a Cancer Research UK Centre Studentship (C27826/A17312)

and Newcastle University Overseas Research Scholarship to RT, a CRUK program grant to

JV and OH (C27943/A12788), grants from the North of England Children's Cancer Research

Fund to OH, JV and SB, by Bloodwise grants 12055 and 15005 to OH and by a grant from

the Kay Kendall Leukaemia Fund (KKL1142) to OH. SB was supported by an NIHR

Academic Clinical Lectureship (CL-2012-01-002), the Sir Bobby Robson Foundation Clinical

Fellowship and a Medical Research Council Clinician Scientist Fellowship (MR/S021590/1).

Work in CB/PNC's lab was funded by a programme grant from Bloodwise (15001). Work in

JMA's lab was funded by a programme grant from Bloodwise (13044). EZ was supported by

an RFBR grant (№17-29-06052). Research in the VVG laboratory was supported in part by

the Ministry of Education of the Republic of Belarus, grant #3.04.3. Research in the AK

laboratory was supported by an RSF grant (20-75-10091).

Abstract word count - 150
Statement of significance word count - 47
Manuscript word count - 6207
Figures - 7
Tables – 0
References - 60

Abstract

The fusion gene *MLL-AF4* defines a high-risk subtype of pro-B acute lymphoblastic leukaemia. However, relapse can be associated with a switch from acute lymphoblastic to acute myeloid leukaemia. Here we show that these myeloid relapses share oncogene fusion breakpoints with their matched lymphoid presentations and can originate in either early, multipotent progenitors or committed B-cell precursors. Lineage switching is linked to substantial changes in chromatin accessibility and rewiring of transcriptional programmes indicating that the execution and maintenance of lymphoid lineage differentiation is impaired. We show that this subversion is recurrently associated with the dysregulation of repressive chromatin modifiers, notably the nucleosome remodelling and deacetylation complex, NuRD. In addition to mutations, we show differential expression or alternative splicing of NuRD members and other genes is able to reprogram the B lymphoid into a myeloid gene regulatory network. Lineage switching in *MLL-AF4* leukaemia is therefore driven and maintained by defunct epigenetic regulation.

Statement of Significance

We demonstrate diverse cellular origins of lineage switched relapse within *MLL-AF4* pro-B acute leukaemia. Irrespective of the developmental origin of relapse, dysregulation of NuRD and/or other epigenetic machinery underpins fundamental lineage reprogramming with profound implications for the increasing use of epitope directed therapies in this high-risk leukaemia.

Introduction

Translocation of Mixed Lineage Leukaemia (*MLL*) with one of over 130 alternative partner

genes is a recurrent cytogenetic finding in both acute myeloid and lymphoblastic leukaemias

and is generally associated with poor prognosis (1, 2). Amongst the most common

translocations is t(4;11)(q21;q23), forming the *MLL-AF4* (also known as *KMT2A-AFF1*)

fusion gene. Uniquely amongst *MLL* rearrangements (*MLL*r), *MLL-AF4* is almost exclusively

associated with pro-B cell acute lymphoblastic leukaemia and is prototypical of infant acute

lymphoblastic leukaemia (ALL) where it carries a very poor prognosis (2). However, despite

this general lymphoid presentation, *MLL-AF4* leukaemias have an intriguing characteristic -

that of lineage switched relapses. Lineage switch acute leukaemias (LSALs) lose their

lymphoid specific features and gain myeloid phenotype upon relapse (3-5). Alternatively,

MLL-AF4 leukaemias may harbour distinct lymphoid and myeloid populations at the same

time, thus classifying as mixed phenotype acute leukaemias (MPALs) of the bilineage

subtype.

In order to understand the molecular basis of lineage promiscuity and switching, we

examined a unique cohort of *MLL-AF4*-positive LSAL presentation/relapse pairs and MPALs.

We demonstrate that disruption of the epigenetic machinery, including the nucleosome

remodelling and deacetylation complex (NuRD), is associated with the loss of lymphoid

restriction. Lineage switch is then enacted through redistribution of transcription factor

binding and chromatin reorganisation. Whilst identified here within this rare clinical context,

our findings bare relevance for our understanding of the transforming capacity of *MLL-AF4*,

and how this oncoprotein imposes lineage determination on haematopoietic precursor cells.

Furthermore, given the high-risk nature of this disease, we provide a novel insight into

factors which may prove critical to the effective implementation of lineage specific, epitope-

directed therapies such as chimeric antigen receptor T-cell (CAR-T) cell or bi-specific T-cell

engaging antibody (BiTE) approaches.

Results

Characterisation of MLL-AF4 acute leukaemias with lineage switch

To characterize lineage promiscuity in *MLL-AF4* leukaemia and the underlying molecular mechanisms, we collected a cohort of ten cases of *MLL-AF4* ALL comprising 6 infant, 2 paediatric and 2 adult patients who had relapsed with acute myeloid leukaemia (AML). Amongst these, one infant patient (LS10) had relapsed following B-lineage directed blinatumomab treatment (Table S1). The time to relapse ranged from 3 to 48 months. Seven patients within the cohort subsequently died. Lineage switch was defined as loss of expression of B lymphoid antigens (CD19, CD22, CD79A) with concomitant gain of expression of myeloid antigens (CD33, CD117/KIT, CD64/FCGR1A) and/or an unequivocal change in morphology to AML (Figure 1A and Table S1). In addition, we studied two *MLL-AF4* infant mixed phenotype acute leukaemias (MPALs), comprising distinct lymphoid and myeloid populations (MPAL1, MPAL2; Table S1).

All matched samples displayed identical oncogene fusion breakpoints at diagnosis and relapse as shown by DNA (n=14) and/or RNA (n=10) sequencing, confirming a common clonal origin and proving that the relapses are not *de novo* or therapy-associated AMLs (Figures 1B, S1A, S1B, Table S1). Breakpoints of LSALs and MPALs show a similar distribution as *MLL-AF4* ALL cases, clustering in *MLL* introns 9-11 and AF4 introns 3 and 4 (6, 7) (Figure S1C, Table S1) thus excluding that distinct, "non-canonical" chromosomal breakpoints are causative for *MLL-AF4*-positive AML. These data raised the question of the cellular origin of relapse and the nature of events secondary to *MLL-AF4* that affect lineage commitment.

Cellular origin of lineage switched relapse

We hypothesised that myeloid leukaemias may not have undergone substantial B-cell receptor (BCR) rearrangements. We used this feature to further interrogate the developmental stage at which the relapse arose. To that end we examined BCR

rearrangements within RNA-seq and whole exome-seq (WES) derived data, using MiXCR

software (8). All ALL cases showed classical oligoclonal rearrangements of BCR loci,

supporting the lymphoid lineage decision (Figure S2A, Table S2). However, we observed

three distinct patterns for AML relapses (Figure 1C). Pattern 1 comprises AML relapse cells

with no BCR rearrangements implying presence of a relapse-initiating cell residing in a

primitive precursor population prior to early DJ recombination (Figure 1C, cases LS01, LS02,

LS04). As a second pattern, we found unrelated BCR rearrangements, which may indicate

either aberrant rearrangement in a myeloid cell or relapse initiating from either a B-lymphoid

cell committed to undergo rearrangement, or a transdifferentiated minor ALL clone with an

alternative rearrangement (Figure 1C, cases LS03, LS06, LS07, LS08, MPAL2).

Interestingly, this pattern is also found in a relapse after blinatumomab treatment (LS10).

Pattern 3 comprises shared BCR rearrangements between diagnostic and relapse material,

which suggest a direct transdifferentiated myeloid relapse from the ALL (Figure 1C, cases

LS05, LS09, MPAL1). These data demonstrate that AML relapses can originate from

different stages of lymphoid leukaemogenesis and suggest, at least for a subset, a common

precursor preceding the pro-B cell stage.

To functionally assess the plasticity of immunophenotypically defined diagnostic ALL and

relapsed AML, we transplanted NSG or MISTRG mice with either ALL or AML cells from

patient LS01 (Figure S2B). Because of the expression of several human myeloid growth

factors, the MISTRG strain more strongly supports AML engraftment than the NSG strain,

which shows a stronger lymphoid bias. Diagnostic ALL transplants rapidly produced

representative CD19+CD33- lymphoid leukaemias in both mouse strains. In contrast,

transplantation of relapsed AML engrafted only in the more myeloid-permissive MISTRG

strain as a CD34+/-CD19-CD33+ AML (Figure S2C). Thus, in contrast to the

immunophenotypic plasticity seen in MPAL leukaemias with wildtype *MLL* (9),

transplantation of a relapsed, fully switched AML was only capable of generating AML.

Lineage switch relapse can originate in HSPC compartments

To further investigate a potential origin of relapse in an early progenitor or stem cell, we

purified haematopoietic stem/progenitor cell (HSPC) populations from diagnostic ALL and

relapsed AML and tested purified populations for the presence of *MLL-AF4* targeted

sequencing. The *MLL-AF4* translocation was found in the lymphoid-primed multipotent

progenitor-like population (LMPP, CD34+CD38-CD45RA+; lymphoid and myeloid potential,

but not megakaryocyte-erythroid potential), in the multipotential progenitor population (MPP,

CD34+CD38-CD45RA-CD90-; no lineage restriction) and for MPAL1 even in the

haematopoietic stem cell-like population (HSC, CD34+CD38-Lin-CD90+) (Figures 2A-C, S3;

Table S3). In line with these findings, serial xenotransplantation of LS01P identified a

persistent human CD34+CD38-CD45RA-CD90+ HSC compartment across four generations

of mice with maintenance of the *MLL-AF4* fusion gene within purified human CD34+ cells

(Figures 2D-F). These findings suggest maintenance of a (pre-)leukaemic clone with high

malignant self-renewal potential in HSPC populations and support our findings from BCR

analysis that, at least in a subgroup of cases, an early multipotent progenitor or HSC can act

as the cell of origin for the AML relapse.

In concordance with the translocation being present within the early HSPC compartment, we

sorted viable differentiated leukocytes and were able to detect the *MLL-AF4* fusion in

myeloid and lymphoid lineages including CD34-CD19/3-HLA-DR+CD14/11c+ monocytes,

NK, B and mature T cells (Figure 2C, S3A, B; Table S3). These findings imply the existence

of a pre-leukaemic progenitor cell, in which *MLL-AF4* does not impose a complete block on

haematopoietic differentiation but is compatible with myeloid and lymphoid differentiation.

These findings raise the question of which factors and molecular mechanisms affect the ALL

and AML lineage choice in *MLL-AF4* leukaemia.

Lineage switch leukaemia is associated with transcriptional reprogramming

We next investigated the underlying molecular events associated with a lineage switch. To

this end we analysed differential gene expression across eight cases, including six LSALs

for which RNA was available at both presentation and relapse, and sorted lymphoid and

myeloid blast populations of two MPALs. This analysis identified in total 1374 up- (adj.

p<0.01, Log Fold change >2) and 1323 down-regulated genes in the AML lineage switches

and the myeloid populations of MPAL patients (Table S4). The most substantially down-

regulated genes include lymphoid genes such as *PAX5* and *RAG2*, but also the polycomb

PRC1 like complex component *AUTS2* and SWI/SNF complex component *BCL7A,* while up-

regulated genes comprise several myeloid genes such as cathepsins, cystatins, *PRAM1* and

CSF3R (Figure 3A). These fingings were consistent, irrespective of which cellular origin of

relapse the BCR rearrangement analysis supported (pattern 1, 2 or 3),

Both gene set enrichment analysis (GSEA) and non-negative matrix factorisation (NMF)

showed that presentation and relapse cases with lineage switch have expression signatures

similar to previously published *MLL-AF4* ALL and *MLL*r AML cases as well as normal

lymphoid and myeloid cell types, respectively (Figures 3B, S4A, B) (10-12). More

specifically, lineage switch included increased expression of factors controlling myeloid

differentiation (e.g., *CSF3R, KIT)* and changes in haematopoietic surface marker expression

(e.g., *CD19, CD22*, *CD33, CD14, FCGR1A/CD64*), loss of immunoglobulin recombination

machinery genes (e.g., *RAG1*, *RAG2, DNTT*) and reduced expression of genes encoding

heavy and light immunoglobulin chains (Figures 3C-E, S4C). Notably, GSEA also indicated

impaired DNA repair and cell cycle progression of the AML relapse when compared with

diagnostic ALL (Figures S4D). In particular, the reduced self-renewal potential might reflect

the transition from ALL with a high incidence of leukaemic stem cells (LSCs) to AML with

fewer LSCs.

MLL fusion proteins including MLL-AF4 have previously been shown to directly regulate

multiple genes linked to haematopoietic and leukaemic stemness (13-16). For instance,

significant changes in gene expression across the *HOXA* cluster, notably with a 50-fold

reduction in *HOXA7* expression, represent a major additional disruption of *MLL*r

leukaemogenic transcriptional regulation (Figures S5A-C) (13, 14). Furthermore, a very

significant portion of target genes including *PROM1*, *IKZF2* and chromatin modifying factors

are highly enriched amongst genes that show lower expression in myeloid lineages (Figures

S5D, E, Table S4). In total, 996 out of 5208 bona fide direct target genes of *MLL-AF4*

changed expression in the AML relapse (15). These data suggest that the process of lineage

switch is associated with a major reorganisation of the *MLL-AF4* transcriptional network and

pose the question of which epigenetic regulators are involved in the lineage determination

associated with this fusion gene.

Reorganisation of chromatin accessibility and transcription factor binding upon

lineage switch

For case LS01 we had sufficient diagnostic material to perform DNase hypersensitivity site

(DHS) analysis and thus link transcriptional changes to altered genome-wide chromatin

accessibility. Many differentially expressed genes showed altered chromatin accessibility in

proximity to the transcriptional start site (TSS), including genes encoding key hematopoietic

surface markers CD33 and CD19, transcription factors and proteins related to differentiation

(Figures 4A-C, S6A).

Changes in chromatin accessibility were linked with an altered pattern of transcription factor

binding. High resolution DHS-seq (digital footprinting) of presentation ALL and relapse AML,

respectively, showed marked genome-wide alterations at sites distal to the TSS, from which

several AML and ALL specific *de novo* occupied transcription factor binding motifs were

identified (Figures 4C-E, S6B). Lineage switch from ALL to AML was associated with a loss

of occupancy of motifs binding lymphoid transcription factors such as EBF or PAX5 and a

gain of occupancy of motifs bound by C/EBP, IRF and NF-κB family members (Figure 4E-F).

The gain in C/EBP motif binding was associated with the expression of a C/EBPA regulated

myeloid transcriptional gene set (Figure 5G). We also observed a redistribution of occupancy

of transcription factors controlling both lymphoid and myeloid maturation such as RUNX, AP-

1 and ETS members to alternative cognate motifs (Figures 4E and S6B) (17, 18). This

finding is exemplified by decreased accessibility of a region located 1 kb upstream of the

CD19 TSS with concomitant loss of EBF binding at this element (Figure 4C). Differential

motif enrichments were associated with changes in RNA expression and chromatin

accessibility at the genes encoding the corresponding cognate transcription factors *EBF1*,

PAX5, *LEF1* (B lymphoid determinants), *NFKB2* and *CEBPB/D/E* (myeloid determinants),

particularly in regions flanking TSSs (Figures 4H, S6B).

In conclusion, the transition from lymphoid to myeloid immunophenotype is associated with

global lineage specific transcriptional reprogramming and genome-wide alteration in

chromatin accessibility and transcription factor binding.

Lineage switch changes alternative splicing patterns

Lineage fidelity and determination are not only linked to differential gene expression but may

also include co- or post-transcriptional mechanisms. It has been previously demonstrated

that lineage commitment during haematopoiesis leads to substantial changes in alternative

mRNA splicing patterns (19). Furthermore, we recently showed that the *AML-ETO* (RUNX1-

RUNX1T1) fusion protein controls leukaemic self-renewal by both differential gene

transcription and alternative splicing (20). To complement the transcriptional analysis we

therefore sought to define the different composition of RNA isoforms in lymphoid and

myeloid populations from lineage switch and MPAL cases. Here we focussed on three

lineage switch patients and the two MPAL patients whose RNA-seq data provided sufficient

read depth for the analysis of exon-exon junctions, exon usage and intron retention (Figures

S7A-C). We detected in total 2630 retained introns (RIs) shared amongst the three lineage

switches with 653 and 343 RIs exclusively found in the diagnostic ALL or the AML relapses,

respectively. This was complemented by 97 exons (DEUs) and 193 exon-exon junctions

(DEEjs) differentially used between diagnostic ALL or the AML relapses. In contrast, this

analysis identified only 43 RIs present in both MPAL cases with 18 and 6 introns specifically

retained in either the lymphoid or myeloid subpopulation, respectively (Figure S7D, Table

S5). Intersection of the affected genes identified 21 shared genes out of 166 DEEjs and 74

DEUs. MPALs had 420 DEUs and 155 DEEjs affecting 257 and 103 genes with 30 genes

having both DEUs and DEEjs (Figure 5A, Table S6). While more than 80% of the non-

differential exon-exon linkages were canonical, this was true for only 15% of the DEEjs.

Here, non-canonical exon skipping and complex splicing events contributed more than 30%

each, most prominently to differential alternative splicing (Figure 5B).

Pathway analysis revealed an enrichment of alternatively spliced genes in immune pathways

including antigen processing, membrane trafficking and FCGR-dependent phagocytosis

reflecting the change from a lymphoid to a myeloid state (Figures 5C, S7E). Furthermore, it

highlighted RNA processing and maturation including mRNA splicing, processing of capped

intron-containing pre-mRNAs and rRNA processing. Indeed, myeloid populations expressed

4-6-fold higher alternatively spliced *SF3B1* and *SRSF5* levels than their matched lymphoid

populations (Figure 5D, E). In addition, we noted a significant number of genes encoding

epigenetic modulators including *KDM5C, HDAC2* and several *CHD* members being

differentially spliced in AML relapse or myeloid subpopulations of MPALs (Figure 5D, E).

Analysing the occurrence of *MLL-AF4* isoforms showed that the fusion site of *MLL-AF4* itself

was subject to alternative splicing in lineage switch and MPAL. Three cases shared the

breakpoints in introns 10 and 4 of *MLL* and *AF4*, respectively. By examining RNA-seq and

competitive RT-PCR data, we identified two co-occurring isoforms with either *MLL* exon9 or

exon10 joined to *AF4* exon4 (Figure 5F). Interestingly, lymphoid populations expressed a

higher ratio of exon10/exon4 over exon9/exon4 than the myeloid populations in all three

cases examined. Furthermore, the HSC and MPP-like populations of MPAL1 showed mainly

expression of the exon10/exon4 splice variant of *MLL-AF4*, thus resembling more the

lymphoid than the myeloid phenotype (Figure 5F). In conclusion, altered isoform expression

of *MLL-AF4* may contribute to lineage choice and the phenotypic switch.

The mutational landscape of lineage switch

Next, we examined the mutational landscape of lineage switched *MLL-AF4* leukaemias by

performing exome sequencing on the entire cohort. In keeping with published data on newly

presenting *MLL*r acute leukaemias (12, 21), exome sequencing of presentation ALL samples

confirmed a relatively quiet mutational landscape in infant ALL cases, with median of 13

nonsynonymous somatic single nucleotide variants (SNVs) or insertions/deletions (indels)

predicted to be deleterious to protein function (Table S7). Many of these were present in less

than 30% of reads and considered sub-clonal. The most commonly mutated genes at

presentation were *NRAS* (3 cases) and *KRAS* (5 cases) (Figure 6A) as described previously

(12). In contrast, relapse AML samples contained a median of 46 deleterious somatic SNVs

and indels (Table S7). This increase can be mainly attributed to three samples (LS03RAML,

LS07RAML and LS08RAML) that carried deleterious mutations in DNA polymerase genes in

the respective major clones linked to hypermutator phenotypes (22, 23) (Figure 6A).

However, we observed this phenotype only in three out of ten relapses, arguing against this

phenomenon being a general requirement for the lineage switch in relapse. Similarly, many

of the predominantly subclonal mutations identified in presentation ALL samples, including

half of *RAS* mutations, were subsequently lost at relapse, indicating alternative subclones as

the origin of relapse (Figure 6A). Finally, both MPALs harboured several mutations that were

exclusively found in either the lymphoid or myeloid subpopulation indicating the presence of

subclones with a lymphoid and myeloid bias (Figure 6A).

Next, we examined the mutation patterns of patient LS01 in greater detail. ALL and AML

contained 104 and 3196 SNVs with a variant allele frequency (VAF) ≥0.3 respectively, with

only 22 shared SNVs between both samples. Globally the most prevalent type of SNV was

the C to T transition in the DNA of both ALL and AML samples (Figure 6B). However, the

contribution of underlying single base substitution (SBS) signatures differed between

diagnosis and relapse. Three different signatures (SBS16, 5 and 1) explained about 60% of

the SNPs found in ALL, while SBS1 seemed to explain more than 50% of all SNPs in the

AML (Figure 6B) suggesting a mutational clock as the main driver of the evolution of relapse

(24). Further inspection of the pattern revealed a mutational signature mainly comprising C

to T transitions and to a lesser degree C to G transversions in NCG triplets, raising the

possibility that thiopurine maintenance treatment may have increased the mutational burden,

resulting in lineage switched relapse in this patient (25).

Twelve deleterious SNVs were identified as unique to the relapse sample of case LS01. The

availability of viable cellular material allowed us to investigate the order of acquisition of

these secondary mutations within the structure of the normal haematopoietic hierarchy. We

sorted this sample to isolate HSC-, MPP-, LMPP- and GMP-like and later populations. Using

a targeted deep sequencing approach we then examined these populations for the presence

of those 12 SNVs. This analysis showed an increasing number of mutations during the

differentiation from MPPs through LMPPs to GMPs. Amongst them, only *PHF3* and *CHD4*

mutations were present within the purified CD34+CD38-CD45RA-CD90- MPP-like fraction

with VAF≥0.3 (Figure 6C and Table S3). In contrast, LMPP- and GMP-like populations

contained all 12 SNVs at high VAF (Table S3). These findings identify the mutation of *CHD4*

and *PHF3* as the earliest genetic events during relapse evolution and suggest them as

potential drivers of an *MLL-AF4* positive, non-lineage committed, pre-leukaemic precursor

population. Subsequent accumulation of additional mutations likely establishes the fully

developed leukaemia with more mature haematopoietic/myeloid immunophenotypes.

Perturbation of CHD4 and PHF3 disrupts lymphoid development in MLL-AF4

expressing cells

The two earliest mutations within LS01 relapse were identified in the Nucleosome

Remodelling and Deacetylation complex (NuRD) member *CHD4* and the plant

homeodomain finger containing *PHF3*. PHF3 is a member of a family of transcriptional

regulators that have been suggested to link the deposition of histone marks to the regulation

of transcription (26). PHF3 itself has been recently identified as an inhibitor of transcription

elongation by competing with TFIIS for binding to the C-terminal domain of RNA polymerase

II (27). NuRD is a multiprotein transcriptional co-repressor complex with both histone

deacetylase and ATP-dependent chromatin remodelling activity. It is a critical factor in the

lymphoid lineage determination in part directed by the transcription factor IKZF1 (28-30).

Both the CHD4 R1068H and PHF3 K1119I mutations affect highly conserved residues

(Figures 7A, S8A) that are predicted by the Condel classifier (31) to disrupt protein function.

Specifically, the CHD4 R1068H mutation has previously been linked to defects in cardiac

development (32).

Whilst mRNA expression levels of the mutant CHD4 R1068H were unaffected in case LS01,

across the remaining cases analysed, expression of *CHD4* and additional NuRD complex

members *MBD3*, *MTA1* and *RBBP4* was reduced following lineage switched myeloid relapse

(Figures 7B). Furthermore, *CHD4* was affected in the MPAL myeloid populations by non-

canonical alternative splicing leading to premature termination of translation, indicating that

this particular pathway was severely disrupted across the cohort, irrespective of the putative

cell of origin of relapse.

The relevance of *CHD4* and *PHF3* in regulating lymphoid versus myeloid lineage choice was

further supported by ARACNE analysis using published ALL (33) and AML (34) expression

datasets (n=216) which we used to reverse-engineer a mutual information network. This

network was trimmed to represent only genes significantly associated with the difference

between AML and ALL thus reflecting the likely influence of genes of interest upon genes

associated with the difference between AML and ALL. This analysis found *CHD4* and *PHF3*

to be the mutated genes with the highest number of edges within the network (*PHF3* – 21

edges, p=0.010; *CHD4* – 12 edges, p=0.0005, Figure S8B and Table S8), implicating them

as causal to the lymphoid/myeloid distinction identified within primary ALL/AML.

To establish a direct functional link from these mutations to the loss of lymphoid lineage

commitment in *MLL-AF4* ALL, we knocked down *CHD4* and *PHF3* in the *MLL-AF4* positive

cell line, SEM. Depletion of *CHD4* and *PHF3* alone or in combination resulted in increased

expression of the myeloid transcription factor *CEBPA* and reduced expression of lymphoid

transcription factors including *LEF1*, *PAX5, TCF3* and *TCF12* (Figure 7C). These changes

were accompanied by a more than twofold increase in CD33 expression in the two *MLL-*

AF4-expressing ALL cell lines SEM and RS4;11, while CD33 levels in two *MLL-AF4-*

negative ALL cell lines remained unaffected, supporting the importance of these epigenetic

regulators specifically within the *MLL-AF4* context (Figures 7D and S8C). GSEA following

PHF3 and *CHD4* knockdown indicated loss of HSC and B-lymphocyte progenitor gene

expression signatures (Figure 7E, left panel), similar to what was observed with the

transcriptomes of the lineage switch leukaemia cases (Figure 7E, right panel). In line with

these cell line experiments, knockdown of *CHD4* and *PHF3* in an ALL PDX generated from

the first relapse of patient LS03 also resulted in a more than twofold increase in CD19/CD33

double and CD33 single positive cells, demonstrating that perturbation of these two genes is

also able to change the immunophenotype of primary *MLL-AF4* ALL cells in a case

susceptible to trans-differentiated relapse (Figure S8D).

In order to examine the role of additional mutations of chromatin modifiers found in our

cohort and known to regulate lineage choice, we investigated the impact of the PRC1

members *PCGF6* and *AUTS2* on CD33 expression in SEM cells. *PCGF6* is mutated in

LS07RAML and LS08RAML and has known roles in B lymphoid malignancy (35). *AUTS2* is

both mutated in LS08RAML and highly expressed in all lymphoid populations examined

(Figures 3A, 6A). STRING network analysis demonstrated close functional associations

within PRC1 and NuRD complexes and their shared associations (Figure S8E). Mutual

interactions include CBX2, a PRC1 complex member, which shows a similar expression

pattern between lineage switch and MPAL cases (Figure S8F). While knockdown of *AUTS2*

did not change CD33 levels, depletion of *PCGF6* also increased CD33 surface expression in

SEM cells, further supporting the notion of epigenetic factors regulating lineage

determination in ALL (Figures S8G). Furthermore, GSEA also indicated impaired function of

PRC1 and PRC2 complexes in the AML relapse compared with the presentation ALL with

down-regulation of their respective target genes (Figure S8H).

Given that the relapse-initiating cell can arise within an uncommitted, *MLL-AF4* translocated

HSPC population, we went on to assess the impact of *CHD4* and *PHF3* function loss in a

human cord blood model, which harbours a chimeric *MLL-Af4* fusion and can be

differentiated both into myeloid and lymphoid lineages (36). Knockdown of either *CHD4* or

PHF3 under lymphoid culture conditions significantly impaired lymphoid differentiation

potential, whilst co-knockdown of *CHD4* and *PHF3* disrupted differentiation entirely (Figures

7F, G and Table S9). Transcriptomic analysis of the sorted populations revealed that CD33

positive cells exhibited metagene expression pattern similar to *MLL*r AML, while the pattern

describing CD19+ cells was most similar to *MLLr* ALL, thus confirming that changes in

surface marker expression were associated with the corresponding changes in the

transcriptomic profiles (Figure S8I).

Taken together, our data show the important role of the NuRD (CHD4), PHF3 and other

(PRC1*)* repressive complexes in the epigenetic control of lymphoid lineage choice. In

particular, dysregulation of *CHD4*/NuRD was mediated by mutation, down-regulation of

expression and differential splicing across the cohort, irrespective of the cellular/clonal origin

of relapse. These data support a role for these factors in the strong lineage determining

capacity of *MLL-AF4* whilst their loss undermines both the execution and the maintenance of

the lymphoid lineage fate.

Discussion

This study describes impaired epigenetic control as being central to the phenomenon of

lymphoid-myeloid lineage switch in *MLL-AF4*-positive leukaemia and identifies the cell of

origin of relapse into AML. We found that the origin of relapse was heterogeneous. Relapse

can directly evolve from pro-B-like ALL blast populations, which agrees with the general self-

renewal capacity of ALL cells (37), but can also originate within the HSPC compartment.

Indeed, analysis of both patient and xenotransplanted cell populations from diagnostic ALL

identified *MLL-AF4* fusion transcripts in MPP- and HSC-like cells. This finding agrees with

recently published data pointing at MPP cells as the origin of *MLL-AF4* leukaemia (38) and is

in line with transcriptomic similarities between t(4;11) ALL and Lin-CD34+CD38-CD19- fetal

liver cells, again suggesting an HSPC as the cell of origin (39).

Irrespective of the cellular origin of the relapse, lineage switching was associated with a

major rewiring of gene regulatory networks. At the level of transcriptional control, the

decision for lymphoid development relies not only on the activation of a lymphoid

transcriptional program, but also on the silencing of a default myeloid program (40). That

decision is enacted by lymphoid master regulators including EBF1, PAX5 and IKAROS,

which represent genes commonly mutated in precursor B-ALL (41-43). Pax5$^{-/-}$ pro-B cells

which lack lymphoid potential, whilst capable of differentiating down erythro-myeloid lineages

in vitro, still maintain expression of early B cell transcription factors *EBF1* and *E2A* (*TCF3*)

(40). In contrast, we show that lineage switching *MLL-AF4* pro-B leukaemic relapse is

associated with significant reduction in expression and binding of these earliest B lymphoid

transcription factors. Their loss is linked with changes in the *MLL-AF4* transcriptional

programme, notably within *HOXA* cluster genes (13, 14) which likely results in a wider

reorganisation of malignant haematopoietic transcriptional networks, ultimately leading to a

myeloid differentiation fate.

Similar to the *Pax5* knockout (40), loss of IKAROS DNA-binding activity prevents lymphoid

differentiation (29). NuRD co-operates directly with IKAROS to repress HSC self-renewal

and subsequent myeloid differentiation, permitting early lymphoid development (29, 44, 45).

We found that the abrogation of this pathway through multiple mechanisms was central to

the lineage switch from ALL to AML. Lineage switch was either associated with mutation,

reduced expression or, in the case of two MPALs, alternative splicing of *CHD4* and other

NuRD components. Long term knockdown of *CHD4* was not tolerated in our cord blood

culture. This is in line with reports showing that complete loss of *CHD4* impairs leukaemic

proliferation (46, 47), both myeloid and lymphoid differentiation of HSPCs and causes

exhaustion of HSC pools (44), indicating that basal CHD4 expression is required for

maintaining AML. Moreover, our observation that a 60% *CHD4* knockdown is associated

with the activation of pluripotency gene signatures is in line with the finding that a partial

inhibition of *CHD4* supported induction of pluripotency in iPSCs, while a complete deletion

eliminated cell proliferation (48).

Whilst our study has investigated the rare clinical occurrence of lineage switching, recent

studies have identified core NuRD and PRC1 complex members as being direct targets of

MLL-AF4 binding (16, 49). We therefore hypothesise that epigenetic regulator genes are co-

opted during *MLL-AF4* leukemogenesis and mediate fundamental lineage specific decision-

making processes, in this case the suppression of the myeloid lineage program. Multiple

routes to their dysregulation may result in escape from this lineage restriction. Our finding

that frontline chemotherapy itself may contribute to relapse highlights the urgent need to find

alternative therapies for this high-risk leukaemia. Equally, however, the associated loss of B

cell surface markers (*e.g.*, CD19) provides an alternative mechanism for relapse following

CAR-T cell or blinatumomab therapy (50, 51) in addition to mutations, alternative splicing

(52, 53) and T cells trogocytosis (54). Whilst these therapies target lineage specific surface

markers, lineage-switched (pre-)leukaemic progenitor populations escape epitope

recognition and provide a potential clonal source for the relapse. Given the increasing use of

advanced immunological therapies, a detailed understanding of the molecular processes

underlying lineage determination and switching will be critical for developing new strategies

to avoid this route to clinical relapse.

Methods

Patient samples and data

Patients were diagnosed by local haematology specialists according to contemporary clinical

diagnostic criteria based on morphology and immunophenotypic analysis. All patient

samples were collected at the point of diagnosis, remission following treatment or relapse

and stored with written informed consent for research in one of six centres (Newcastle

Haematology Biobank, Newcastle, UK; University Hospital Schleswig-Holstein, Kiel,

Germany; Dmitry Rogachev National Medical Research Center of Pediatric Hematology,

Oncology and Immunology, Moscow, Russia; Haematological Malignancy Diagnostic

Service, Leeds, UK; Princess Maxima Center for Pediatric Oncology, Utrecht, The

Netherlands; Cincinnati Children's Hospital Medical Center, Cincinnati, USA). Mononuclear

cells were isolated from bone marrow or peripheral blood by density centrifugation followed

by immediate extraction of DNA or RNA, or cryopreservation in the presence of 10% v/v

DMSO.

Samples were requested and used in accordance with the ethical approvals granted to each

of the local/institutional ethical review boards (NRES Committee North East - Newcastle &

North Tyneside 1, UK, reference 07/H0906/109+5; Medical Faculty Christian-Albrechts

University, Kiel, reference A 103/08; Dmitry Rogachev National Medical Research Center,

Moscow, references MB2008: 22.01.2008, MB2015: 22.01.2015, ALL-REZ-2014:

28.01.2014; Haematological Malignancy Research Network, Yorkshire, UK, reference

04/Q1205/69; Haematological Malignancy Diagnostic Service, Leeds, UK, reference

14/WS/0098; Erasmus MC METC, Netherlands, reference MEC-2016-739; IRB of Cincinnati

Children's Hospital, USA, reference 2010-0658) and in accordance with the Declaration of

Helsinki. Each patient/sample was allocated an anonymised reference and no identifiable

information was shared.

DHS library generation, sequencing, and mapping

DHS analysis was performed as described previously (55). DNase I (Worthington, Cat#

LS006328) digestion was performed using ~5 million patient sample cells using 8 units

(presentation) or 14 units (relapse) for 3 min at 22°C in a 1 mM $CaCl_2$ supplemented buffer.

Nuclear proteins were digested with 1 mg/ml Proteinase K overnight at 37°C. DNase I

digestion products were size-selected on an agarose gel, cutting below 150 bp. High-

throughput sequencing libraries were prepared from 10 ng of size-selected material, using

the Kapa Hyperprep kit as per manufacturer's instruction. Libraries were sequenced with 50

bp single-end reads on an Illumina HiSeq 2500 sequencer according to manufacturer's

instructions.

Fastq files were generated using bcl2fastq (1.8.4) and subsequently aligned to the hg19

assembly (NCBI Build 37) with the use of bowtie2 (2.1.0), with –very-sensitive-local as a

parameter. Read coverage generation and peak detection were carried out using MACS

1.4.1 using --keep-dup=all –g hs -w -S. Pairwise comparisons were performed as previously

described (55). Digital footprinting was carried out using the Wellington package using

default parameters (56). Differential footprinting analysis was carried out on footprints using

the Wellington-bootstrap (57) package with default parameters. Average profiles and

heatmaps were obtained using the functions dnase_average_profile and

dnase_to_javatreeview from the Wellington package. Heatmaps were plotted using Java

TreeView.

Exome sequencing

Germline DNA from cases LS08 and LS09 were extracted from formalin fixed paraffin

embedded remission bone marrow using QIAamp DNA FFPE Tissue Kit (Qiagen,

Cat#56404). Other DNA samples were extracted from either bone marrow or peripheral

blood using AllPrep DNA/RNA Mini Kit (Qiagen, Cat#80204), QIAamp DNA Mini Kit (Qiagen,

Cat#51306), or innuPREP DNA/RNA Mini Kit (Analytik Jena, Cat#845-KS-2080050),

according to manufacturers' instructions. The exons were captured using SureSelect XT2

Human All Exon V6 (Agilent), and sequenced by paired-end 75 bp sequencing on

HiSeq4000 (Illumina), resulting in roughly 45 million reads per sample. DNA from the

myeloid and lymphoid cellular compartments derived from MPAL patients samples, were

pre-processed with KAPA HyperPlus Kit (Roche) followed by exons enrichment with KAPA

HyperCapture Kit (Roche), and sequenced by paired-end 300 bp sequencing on

NovaSeq6000 (Illumina), resulting in roughly 25 million reads per sample.

Raw reads were aligned to human reference genome (hg19 or hg38 for lineage switch or

MPAL patients, respectively) using Burrows-Wheeler Aligner (BWA) 0.7.12 and were

processed using the Genome Analysis Toolkit (GATK, v3.8 or 4.1). MuTect (v1.1.7) and

MuTect2 (4.1) were used to identify somatic variants for each matched sample pair. Variants

were annotated using Ensembl Variant Effect Predictor (VEP, version 90).

RNA sequencing

Total RNA was extracted with AllPrep DNA/RNA Mini Kit (Qiagen, Cat#80204), innuPREP

DNA/RNA Mini Kit (Analytik Jena, Cat#845-KS-2080050), or TRIzol (Thermo Fisher

Scientific, Cat# 15596026) followed by RNeasy Mini Kit (Qiagen, Cat#74106) from either

bone marrow or peripheral blood, according to manufacturers' instructions. Messenger RNA

was captured using NEBNext Ultra Directional RNA Kit in combination with NEBNext poly(A)

mRNA Magnetic Isolation Module or KAPA RNA HyperPrep Kit with RiboErase (HMR) in

case of lineage switch or MPAL patients respectively, and submitted for paired-end 150 bp

sequencing on HiSeq4000 (Illumina) or paired-end 300 bp sequencing on NovaSeq6000

(Illumina) depending on the analysed patients group. For each sample, transcript abundance

was quantified from raw reads with Salmon (version 0.8.2) using the reference human

transcriptome (hg38) defined by GENCODE release 27. An R package Tximport (version

1.4.0) was used to estimate gene-level abundance from Salmon's transcript-level counts.

Gene-level differential expression analysis was performed using DESeq2 (version 1.16.1).

Differential splicing events were identified in both presentation/relapse pairs or

lymphoid/myeloid cellular fractions, using pipeline described previously (20).

Whole genome sequencing

Presentation, remission and relapse DNA samples from case LS01 were sequenced by

Illumina UK and analysed using the remission sample as the matching normal. Sequencing

reads were aligned to the human GRCh37.1 reference genome using ISAAC (58).

Identification of somatic SNVs and small somatic indels (<50 bp) was performed by Strelka

(59). Large structural variants (including deletions, inversions, duplications and insertions all

>50bp and translocations) were called by Manta (60).

Nested multiplex PCR and targeted sequencing

The 12 mutation candidate driver genes and *MLL-AF4* LS01RAML were amplified using

gDNA as the template by nested multiplex PCR method. The primers were designed using

Primer Express (Applied Biosystems) software.

PCR amplifications were carried out using Phusion® High-Fidelity PCR Master Mix with HF

Buffer (NEB, Cat#M0531L) according to the manufacturer's instructions (25 µl reaction), plus

80 – 200 nM of each primer set. The first multiplex reaction parameters were: one cycle at

98°C for 2 min, thirty cycles of 98°C for 10 s, 63°C for 30 s, and 72°C for 30 s, followed by

one cycle at 72°C for 10 min. The products were diluted 500-fold, and 1 µl used as the

template for the second PCR reactions (25 µl reaction). The second/nested multiplex

reaction parameters were: one cycle at 98°C for 2 min, twenty cycles of 98°C for 10 s, 65°C

for 30 s, and 72°C for 30 s, followed by one cycle at 72°C for 10 min. The amplicons that

were taken forward for next-generation targeted sequencing had additional CS1

(ACACTGACGACATGGTTCTACA) and CS2 (TACGGTAGCAGAGACTTGGTCT) Fluidigm

tag sequences on the nested PCR primer forward and reverse, respectively. The amplicons

were barcoded using Fluidigm Access Array Barcode Library for Illumina Sequencers

(Cat#100-4876) by taking 0.8 µl multiplex PCR products (multiplex group A-C), 4 µl Fluidigm

barcode primer (400 nM final concentration), 10 µl of 2X Phusion Master Mix (NEB,

Cat#M0531L), and 3.6 µl H_2O (20 µl reaction). The barcoding PCR reaction parameters

were: one cycle at 98°C for 2 min, six cycles of 98°C for 10 s, 60°C for 30 s, and 72°C for 1

min, followed by one cycle at 72°C for 10 min. The products were run on 2% agarose gel

and extracted using the QIAquick Gel Extraction Kit (Qiagen, Cat#28706). The purified

products were submitted for paired-end 300 bp sequencing on MiSeq (Illumina), resulting in

>1,000 coverage per gene.

Lymphoid differentiation of transduced MLL-Af4 cord blood cells

MLL-Af4 cord blood cells (36) were transduced with short hairpin constructs targeting CHD4,

PHF3 or NTC control (as described in supplementary methods) and co-cultured with MS-5

stroma cells in lymphoid culture conditions. Single and triple transduced populations were

identified using the construct specific fluorophores and lineage specific surface markers

assessed as a proportion of the total transduced leukocyte population.

MLL-Af4 stem cell expression analysis

Following myeloid or lymphoid culture of *MLL/Af4* transduced CD34+ cord blood cells,

CD19+CD33-, CD19-CD33+ and CD19-CD33- populations were flow sorted, lysed and RNA

extracted using RNeasy Micro Kit (Qiagen), according to the manufacturer's instructions.

Input RNA was equilibrated to a starting input cell number of 300 cells per population before

cDNA and sequencing library production were performed using SMARTSeqv4 (Clontech)

and NexteraXT (Illumina) kits, according to manufacturer's instructions.   The resultant

libraries were submitted for paired-end 150 bp sequencing on a NEXTSeq500 (Illumina). For

each sample, transcript abundance was quantified from raw reads with Salmon (version

0.8.2) using the reference human transcriptome (hg38) defined by GENCODE release 27.

An R package Tximport (version 1.4.0) was used to estimate gene-level abundance from

Salmon's transcript-level counts. Gene-level differential expression analysis was performed

using DESeq2 (version 1.16.1) prior NMF analysis as described above.

Data availability

Exome sequencing data and genome sequencing data presented in this manuscript have

been deposited in the NCBI Sequence Read Archive (SRA) under project numbers

PRJNA547947 and PRJNA547815 respectively. Immunoglobulin and TCR sequencing data

have been deposited in NCBI SRA under project number PRJNA511413. RNA sequencing

data and DNase hypersensitivity sequencing data were deposited in Gene Expression

Omnibus under accession numbers GSE132396 and GSE130142 respectively. All deposited

data will be publically available following publication of the manuscript. Requests for

additional specific data/materials should be made to Olaf Heidenreich

(O.T.Heidenreich@prinsesmaximacentrum.nl).

Acknowledgements

We thank Jon Coxhead and Raf Hussain at the Newcastle University core genomics facility

as well as Marc van Tuil at the Princess Maxima Center Diagnostic department for

development of sequencing strategies. We acknowledge the Newcastle University Flow

Cytometry Core Facility and Tomasz Poplonski at the Princess Maxima Center Flow

Cytometry core facility for their assistance with the generation of flow cytometry data and cell

sorting strategies as well as the Newcastle University Bioinformatics Support Unit for helping

to develop the analysis approach for sequencing data. We thank Ruben van Boxtel and

Eline Bertrums at the Princess Maxima Center for the generation of single cell derived

clones, derived from early progenitors sorted from MPAL patient samples. We thank

Monique den Boer, Frank van Leeuwen and Ronald Stam for critically reading the

manuscript.

This study makes use of data generated by the St. Jude Children's Research Hospital –

Washington University Pediatric Cancer Genome Project and the Therapeutically Applicable

Research to Generate Effective Treatments (TARGET) initiative, phs000218, managed by

the NCI (see supplementary methods).

Author contributions

Conceptualization, O.H., S.B., C.B.;

Methodology, O.H., C.B., R.T., K.S., P.M., S.B., A.P., C.M., A.K., Z.K., J.B., V.B., M.H.,

R.M., J.V., J.M.A., S.L.;

Software Programming, S.N., J.H.K., V.V.G., A.K., D.W., Pi.C.;

Formal Analysis, S.N., J.H.K., V.V.G., A.K., Z.K., J.B., D.W., Pi.C., C.B., O.H.;

Investigation, R.T., K.S., P.M., A.P., C.M., H.J.B., A.K., S.A., M.R.I., P.E., H.M., A.E.,

N.M.S., S.E.F., Y.S., D.P., Pi.C.;

Resources, F.V., E.Z., S.L., J.S., A.S., J.C.M., L.J.R., C.E., O.A.H., S.Ba., R.S., N.M., M.C.,

V.B., R.M., M.W., C.J.H., C.A.C., Pe.C., M.H., D.S., Y.O., M.J.T., P.N.C., J.C.M., C.B., O.H.;

Data Curation, S.N., D.W., Pi.C.;

Writing, S.B., O.H., C.B., R.T., K.S.;

Supervision, O.H., S.B., J.M.A., J.V., C.B., ;

Funding Acquisition, O.H., J.V., S.B., C.B., P.N.C., J.M.A., E.Z.**References**

1.   Moorman AV, Ensor HM, Richards SM, Chilton L, Schwab C, Kinsey SE, *et al.*
Prognostic effect of chromosomal abnormalities in childhood B-cell precursor acute
lymphoblastic leukaemia: results from the UK Medical Research Council ALL97/99
randomised trial. *Lancet Oncol* **2010**; 11:429–438.
2.   Meyer C, Burmeister T, Gröger D, Tsaur G, Fechina L, Renneville A, *et al.* The MLL
recombinome of acute leukemias in 2017. *Leukemia* **2018**; 32:273–284.
3.   Jiang JG, Roman E, Nandula SV, Murty VV, Bhagat G, Alobeid B. Congenital MLL-
positive B-cell acute lymphoblastic leukemia (B-ALL) switched lineage at relapse to
acute myelocytic leukemia (AML) with persistent t(4;11) and t(1;6) translocations and JH
gene rearrangement. *Leuk Lymphoma* **2005**; 46:1223-1227.
4.   Germano G, Pigazzi M, del Giudice L, Campo Dell'Orto M, Spinelli M, *et al.* Two
consecutive immunophenotypic switches in a child with MLL-rearranged acute
lymphoblastic leukemia. *Haematologica* **2006**; 91:ECR09.
5.   Rossi JG, Bernasconi AR, Alonso CN, Rubio PL, Gallego MS, Carrara CA, *et al.*
Lineage switch in childhood acute leukemia: An unusual event with poor outcome. *Am J*
Hematol **2012**; 87:890–897.
6.   Meyer C, Kowarz E, Hofmann J, Renneville A, Zuna J, Trka J, *et al.* New insights to the
MLL recombinome of acute leukemias. *Leukemia* **2009**; 23:1490–1499.
7.   Jung R, Jacobs U, Krumbholz M, Langer T, Keller T, De Lorenzo P, *et al.* Bimodal
distribution of genomic MLL breakpoints in infant acute lymphoblastic leukemia
treatment. *Leukemia* **2010**; 24:903–907.

8. Bolotin DA, Poslavsky S, Davydov AN, Frenkel FE, Fanchi L, Zolotareva OI, *et al.* Antigen receptor repertoire profiling from RNA-seq data. *Nat Biotechnol* **2017**; 35:908–911.

9. Alexander TB, Gu Z, Iacobucci I, Dickerson K, Choi JK, Xu B, *et al.* The genetic basis and cell of origin of mixed phenotype acute leukaemia. *Nature* **2018**; 562;373-379.

10. Zangrando A, Dell'orto MC, Te Kronnie G, Basso G. MLL rearrangements in pediatric acute lymphoblastic and myeloblastic leukemias: MLL specific and lineage specific signatures. *BMC Med Genomics* **2009**; 2-36

11. Novershtern N, Subramanian A, Lawton LN, Mak RH, Haining WN, McConkey ME, *et al.* Densely Interconnected Transcriptional Circuits Control Cell States in Human Hematopoiesis. *Cell* **2011**; 144:296–309.

12. Andersson AK, Ma J, Wang J, Chen X, Gedman AL, Dang J, *et al.* The landscape of somatic mutations in infant MLL-rearranged acute lymphoblastic leukemias. *Nat Genetics* **2015**; 47:330-337.

13. Somervaille TCP, Matheny CJ, Spencer GJ, Iwasaki M, Rinn JL, Witten DM, *et al.* Hierarchical maintenance of MLL myeloid leukemia stem cells employs a transcriptional program shared with embryonic rather than adult stem cells. *Cell Stem Cell* **2009**; 4:129–140.

14. Gessner A, Thomas M, Castro PG, Büchler L, Scholz A, Brümmendorf TH, *et al.* Leukemic fusion genes MLL/AF4 and AML1/MTG8 support leukemic self-renewal by controlling expression of the telomerase subunit TERT. *Leukemia* **2010**; 24:1751–1759.

15. Wilkinson AC, Ballabio E, Geng H, North P, Tapia M, Kerry J, *et al.* RUNX1 is a key target in t(4;11) leukemias that contributes to gene activation through an AF4-MLL complex interaction. *Cell Rep* **2013**; 3:116–127.

16. Kerry J, Godfrey L, Repapi E, Tapia M, Blackledge NP, Ma H, *et al.* MLL-AF4 Spreading Identifies Binding Sites that Are Distinct from Super-Enhancers and that Govern Sensitivity to DOT1L Inhibition in Leukemia. *Cell Rep* **2017**; 18:482–495.

17. Hohaus S, Petrovick MS, Voso MT, Sun Z, Zhang DE, Tenen DG. PU.1 (Spi-1) and C/EBP alpha regulate expression of the granulocyte-macrophage colony-stimulating factor receptor alpha gene. *Mol Cell Biol* **1995**; 15:5830–5845.

18. Leddin M, Perrod C, Hoogenkamp M, Ghani S, Assi S, Heinz S, *et al.* Two distinct auto-regulatory loops operate at the PU.1 locus in B cells and myeloid cells. *Blood* **2011**; 117:2827–2838.

19. Chen L, Kostadima M, Martens JHA, Canu G, Garcia SP, Turro E, *et al.* Transcriptional diversity during lineage commitment of human blood progenitors. *Science* **2014**; 345:1251033.

20. Grinev VV, Barneh F, Ilyushonak IM, Nakjang S, Smink J, van Oort A, *et al.* RUNX1/RUNX1T1 mediates alternative splicing and reorganises the transcriptional landscape in leukemia. *Nat Commun* **2021**; 12:520.

21. Dobbins SE, Sherborne AL, Ma YP, Bardini M, Biondi A, Cazzaniga G, *et al.* The silent mutational landscape of infant MLL-AF4 pro-B acute lymphoblastic leukemia. *Genes Chromosomes Cancer* **2013**; 52:954–960.

22. Erson-Omay EZ, Çağlayan AO, Schultz N, Weinhold N, Omay SB, Özduman K, *et al.* Somatic POLE mutations cause an ultramutated giant cell high-grade glioma subtype with better prognosis. *Neuro Oncol* **2015**; 17:1356–1364.

23. Mur P, García-Mulero S, Del Valle J, Magraner-Pardo L, Vidal A, Pineda M, *et al.* Role of POLE and POLD1 in familial cancer. *Genet Med* **2020**; 22:2089–2100.

24. Alexandrov LB, Jones PH, Wedge DC, Sale JE, Campbell PJ, Nik-Zainal S, Stratton, *et al.* Clock-like mutational processes in human somatic cells. *Nat Genetics* **2015**; 47:1402–1407.

25. Li B, Brady SW, Ma X, Shen S, Zhang Y, Li Y, *et al.* Therapy-induced mutations drive
the genomic landscape of relapsed acute lymphoblastic leukemia. *Blood* **2020**; 135:41–
55.
26. Kinkelin K, Wozniak GG, Rothbart SB, Lidschreiber M, Strahl BD, Cramer P. Structures
of RNA polymerase II complexes with Bye1, a chromatin-binding PHF3/DIDO
homologue. *Proc Natl Acad Sci U S A* **2013**; 110:15277–15282.
27. Appel LM, Franke V, Bruno M, Grishkovskaya I, Kasiliauskaite A, Schoeberl UE, *et al.*
PHF3 regulates neuronal gene expression through the new Pol II CTD reader domain
SPOC. bioRxiv **2020**; p. 2020.02.11.943159.
28. Ng SY, Yoshida T, Zhang J, Georgopoulos K. Genome-wide lineage-specific
transcriptional networks underscore Ikaros-dependent lymphoid priming in
hematopoietic stem cells. *Immunity* **2009**; 30:493–507.
29. Zhang J, Jackson AF, Naito T, Dose M, Seavitt J, Liu F, *et al.* Harnessing of the
nucleosome-remodeling-deacetylase complex controls lymphocyte development and
prevents leukemogenesis. *Nat Immunology* **2011**; 13:86–94.
30. Arends T, Dege C, Bortnick A, Danhorn T, Knapp JR, Jia H, *et al.* CHD4 is essential for
transcriptional repression and lineage progression in B lymphopoiesis. *Proc Natl Acad
Sci U S A* **2019**; 116:10927-10936.
31. González-Pérez A, López-Bigas N. Improving the assessment of the outcome of
nonsynonymous SNVs with a consensus deleteriousness score, Condel. *Am J Hum
Genet* **2011**; 88:440–449.
32. Sifrim A, Hitz MP, Wilsdon A, Breckpot J, Al Turki SH, Thienpont B, *et al.* Distinct
genetic architectures for syndromic and nonsyndromic congenital heart defects
identified by exome sequencing. *Nat Gen* **2016**; 48:1060–1065.
33. Kang H, Chen IM, Wilson CS, Bedrick EJ, Harvey RC, Atlas SR, *et al*. Gene expression
classifiers for relapse-free survival and minimal residual disease improve risk
classification and outcome prediction in pediatric B-precursor acute lymphoblastic
leukemia. *Blood* **2010**; 115:1394–1405.
34. Gentles AJ, Plevritis SK, Majeti R, Alizadeh AA. Association of a leukemic stem cell
gene expression signature with clinical outcomes in acute myeloid leukemia. *JAMA*
2010; 304:2706–2715.
35. Ferreira BI, García JF, Suela J, Mollejo M, Camacho FI, Carro A, *et al.* Comparative
genome profiling across subtypes of low-grade B-cell lymphoma identifies type-specific
and common aberrations that target genes with a role in B-cell neoplasia.
Haematologica **2008**; 93:670–679.
36. Lin S, Luo RT, Ptasinska A, Kerry J, Assi SA, Wunderlich M, *et al.* Instructive Role of
MLL-Fusion Proteins Revealed by a Model of t(4;11) Pro-B Acute Lymphoblastic
Leukemia. *Cancer Cell* **2016**; 30:737–749.
37. le Viseur C, Hotfilder M, Bomken S, Wilson K, Röttgers S, Schrauder A, *et al*. In
childhood acute lymphoblastic leukemia, blasts at different stages of immunophenotypic
maturation have stem cell properties. *Cancer Cell* **2008**; 14:47–58.
38. Malouf C, Ottersbach K. The fetal liver lymphoid-primed multipotent progenitor provides
the prerequisites for the initiation of t(4;11) MLL-AF4 infant leukemia. *Haematologica*
2018; 103:e571–e574.
39. Agraz-Doblas A, Bueno C, Bashford-Rogers R, Roy A, Schneider P, Bardini M, *et al*.
Unravelling the cellular origin and clinical prognostic markers of infant B-cell acute
lymphoblastic leukemia using genome-wide analysis. *Haematologica* **2019**; 104:1176-
1188.
40. Nutt SL, Heavey B, Rolink AG, Busslinger M. Commitment to the B-lymphoid lineage
depends on the transcription factor Pax5. *Nature* **1999**; 402:14–20.

41. Mullighan CG, Goorha S, Radtke I, Miller CB, Coustan-Smith E, Dalton JD, *et al*.
Genome-wide analysis of genetic alterations in acute lymphoblastic leukaemia. *Nature*
2007; 446:758–764.
42. Boer JM, van der Veer A, Rizopoulos D, Fiocco M, Sonneveld E, de Groot-Kruseman
HA, *et al*. Prognostic value of rare IKZF1 deletion in childhood B-cell precursor acute
lymphoblastic leukemia: an international collaborative study. *Leukemia* **2016**; 30:32–38.
43. Witkowski MT, Hu Y, Roberts KG, Boer JM, McKenzie MD, Liu GJ, *et al.* Conserved
IKAROS-regulated genes associated with B-progenitor acute lymphoblastic leukemia
outcome. *J Exp Med* **2017**; 214:773–791.
44. Yoshida T, Hazan I, Zhang J, Ng SY, Naito T, Snippert HJ, *et al.* The role of the
chromatin remodeler Mi-2beta in hematopoietic stem cell self-renewal and multilineage
differentiation. *Genes Dev* **2008**; 22:1174–1189.
45. Lu X, Chu CS, Fang T, Rayon-Estrada V, Fang F, Patke A, *et al.* MTA2/NuRD
Regulates B Cell Development and Cooperates with OCA-B in Controlling the Pre-B to
Immature B Cell Transition. *Cell Rep* **2019**; 28:472-485.
46. Sperlazza J, Rahmani M, Beckta J, Aust M, Hawkins E, Wang S, *et al*. Depletion of the
chromatin remodeler CHD4 sensitizes AML blasts to genotoxic agents and reduces
tumor formation. *Blood* **2015**; 126:1462–1472.
47. Heshmati Y, Türköz G, Harisankar A, Kharazi S, Boström J, Dolatabadi EK, *et al.* The
chromatin-remodeling factor CHD4 is required for maintenance of childhood acute
myeloid leukemia. *Haematologica* **2018**; 103:1169-1181
48. Mor N, Rais Y, Sheban D, Peles S, Aguilera-Castrejon A, Zviran A, *et al.* Neutralizing
Gatad2a-Chd4-Mbd3/NuRD Complex Facilitates Deterministic Induction of Naive
Pluripotency. *Cell Stem Cell* **2018**; 23:412-425.
49. Harman JR, Thorne R, Jamilly M, Tapia M, Crump NT, Rice S, *et al*. A KMT2A-AFF1
gene regulatory network highlights the role of core transcription factors and reveals the
regulatory logic of key downstream target genes. *Genome Res* **2021**; online ahead of
print.
50. Gardner R, Wu D, Cherian S, Fang M, Hanafi LA, Finney O, *et al.* Acquisition of a
CD19-negative myeloid phenotype allows immune escape of MLL-rearranged B-ALL
from CD19 CAR-T-cell therapy. *Blood* **2016**; 127:2406-2410.
51. Rayes A, McMasters RL, O'Brien MM. Lineage switch in MLL-rearranged infant
leukemia following CD19-directed therapy. *Pediatr Blood Cancer* **2016**; 63:1113-510.
52. Sotillo E, Barrett DM, Black KL, Bagashev A, Oldridge D, Wu G, *et al*. Convergence of
Acquired Mutations and Alternative Splicing of CD19 Enables Resistance to CART-19
Immunotherapy. *Cancer Discovery* **2015**; 5:1282–1295.
53. Rabilloud T, Potier D, Pankaew S, Nozais M, Loosveld M, Payet-Bornet D. Single-cell
profiling identifies pre-existing CD19-negative subclones in a B-ALL patient with CD19-
negative relapse after CAR-T therapy. *Nat Commun* **2021**; 12:865.
54. Hamieh M, Dobrin A, Cabriolu A, van der Stegen SJC, Giavridis T, Mansilla-Soto J, *et*
al. CAR T cell trogocytosis and cooperative killing regulate tumour antigen escape.
Nature **2019**; 568:112–116.
55. Cauchy P, James SR, Zacarias-Cabeza J, Ptasinska A, Imperato MR, Assi SA, *et al*.
Chronic FLT3-ITD Signaling in Acute Myeloid Leukemia Is Connected to a Specific
Chromatin Signature. *Cell Rep* **2015**; 12:821–836.
56. Piper J, Elze MC, Cauchy P, Cockerill PN, Bonifer C, Ott S. Wellington: a novel method
for the accurate identification of digital genomic footprints from DNase-seq data. *Nucleic*
Acids Research **2013**; 41:e201–e201.
57. Piper J, Assi SA, Cauchy P, Ladroue C, Cockerill PN, Bonifer C, Ott S. Wellington-
bootstrap: differential DNase-seq footprinting identifies cell-type determining
transcription factors. *BMC Genomics* **2015**; 16:1000.

58. Raczy C, Petrowski R, Saunders CT, Chorny I, Kruglyak S, Margulies EH, *et al*. Isaac:
850   Ultra-fast whole-genome secondary analysis on Illumina sequencing platforms.
851   *Bioinformatics* **2013**; 29:2041–2043.
852   59. Saunders CT, Wong WSW, Swamy S, Becq J, Murray LJ, Cheetham RK. Strelka:
853   Accurate somatic small-variant calling from sequenced tumor-normal sample pairs.
854   *Bioinformatics* **2012**; 28:1811–1817.
855   60. Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Källberg M, *et al*. Manta:
856   rapid detection of structural variants and indels for germline and cancer sequencing
857   applications. *Bioinformatics* **2016**; 32:1220–1222.

859   Figure 1

Figure 1. Characterisation of the *MLL-AF4* lineage switch cases. (A) Morphological change from

lymphoblastic leukaemia (left panel) to acute monoblastic/monocytic leukaemia (right panel). The

scale bar represents 20 μm. (B) Sanger sequencing of *MLL-AF4* and reciprocal *AF4/MLL* fusions in

LS01 presentation ALL (upper panel) and relapse AML (lower panel) identifies a common breakpoint

with identical filler sequence in ALL and AML samples. (C) Evaluation of B-cell receptor repertoires in

ALL (presentation) and AML (relapse) lineage switch, and MPAL cases identified three distinct

patterns. Pattern 1 - with clonotypes identified only in the ALL (LS01, LS02, LS04). Pattern 2 – distinct

clonotypes identified in ALL and AML (LS03, LS06, LS07, LS08, LS10, MPAL2). Pattern 3 – identical

clonotypes shared between ALL and AML (LS05, LS09, MPAL1). Graphs represent frequency of

clones identified with WES and/or RNAseq in LSAL and MPAL samples, respectively.

Figure 2

Figure 2. Haematopoietic stem/progenitor populations carry *MLL-AF4.* (A) Flow cytometric

sorting strategy for haematopoietic stem/progenitor cell (HSPC) populations. (B) PCR identification of

the specific *MLL-AF4* fusion within sorted HSPC populations LS01 and MPAL1 cases. (C) Summary

of *MLL-AF4* positivity within different HSPC populations analysed in patients LS01PALL and early

progenitors and lymphoid fraction of MPAL1, presented as red and pink circles, respectively. (D) Flow

cytometric analysis showing sequential transplantation of LS01 ALL presentation sample across four

generations of NSG mouse xenografts. The HSC population (CD34+CD38-CD45RA-CD90+) is

maintained across four mice generations. (E) Proportion of bone marrow human CD34+ cells in

CD34+CD38-CD45RA-CD90+ HSC gate in all analysed xenografts. (F) PCR identification of the

specific *MLL-AF4* fusion within the sorted HSPC xenograft sample.

Figure 3

Figure 3. Transcriptional reprogramming in lineage switch and MPAL cases.
(A) Heatmap showing the top 100 differentially expressed genes between ALL and AML from six lineage switch (LS01, LS03, LS04, LS05, LS06, LS10) and two MPAL cases, ranked by stat value. (B) Enrichment of myeloid growth and differentiation signature in relapsed samples (left panel) identified by GSEA analyses, also pointing to downregulation of genes highly correlated with acute lymphoblastic leukemia (middle and right panel). Gene set enrichment analyses have been performed

based on data derived from six lineage switch samples. FDR – false discovery rate, NES –

normalised enrichment score. (C-E) Differential expression of (C) lineage specific, (D) immunoglobulin

recombination machinery, and (E) genes encoding immunoglobulin heavy and light chains in lineage

switch and MPAL cases. Error bars show standard error of the mean (SEM) for lineage switch cases

and ranges for two MPAL cases.

Figure 4

Figure 4. Chromatin re-organisation and differential transcription factor binding underpins

lineage switching. (A) DNase hypersensitivity site sequencing identifies 13619 sites with a log2 fold

reduction and 12203 sites with a log2 fold increase following lineage switch to AML. (B) University of

California, Santa Cruz (UCSC) genome browser screenshot displaying differential expression at

lineage specific loci (lower red tracks) accompanied by altered DNase hypersensitivity (upper black

tracks) proximal to the transcriptional start site (TSS) of *CD33*. (C) UCSC genome browser

screenshot for *CD19* zoomed in on an ALL-associated DHS with EBF occupation as indicated by high

resolution DHS-seq and Wellington analysis. FP - footprint. (D) Heat maps showing distal DHS

regions specific for AML relapse on a genomic scale. Red and green indicate excess of positive and

negative strand cuts, respectively, per nucleotide position. Sites are sorted from top to bottom in order

of decreasing Footprint Occupancy Score. (E) De novo motif discovery in distal DHSs unique to AML

relapse as compared to ALL relapse as shown in (D). (F) EBF1 and C/EBP binding motifs

demonstrate differential motif density in presentation ALL and relapse AML. (G) Enrichment of a

myeloid C/EBPA network gene set in signatures associated with relapse AML and diagnostic ALL

samples as identified by GSEA (H) Differential expression of TFs cognately binding to differentially

accessible motifs shown in (F). TFs whose binding motifs show increased accessibility in ALL are in

blue whilst those showing increased accessibility in AML are in red. RUNX1 in black reflects enriched

accessibility of different RUNX1 binding sites in ALL and AML. Error bars show SEM or ranges in

LSAL and MPAL cases, respectively.

Figure 5

Figure 5. Alternative splicing in lineage switch and MPAL cases. (A) Volcano plots demonstrating

differential usage of exon-exon junctions in the transcriptome of AML/myeloid versus ALL/lymphoid

cells of lineage switch (LS01, LS03 & LS04) or MPAL patients. The vertical dashed lines represent

two-fold differences between the AML and ALL cells and the horizontal dashed line shows the FDR-

adjusted q-value threshold of 0.05 (upper panel). Venn diagrams (lower panel) showing distribution of

splice variants identified as significantly changed in AML (or myeloid fraction of MPAL patients),

including exon-exon junctions (DEEj), differential exome usage (DEU) and retained introns (RI). (B)

Pie charts showing the classification of non-differential (non-DEEj) and differential (DEEj) exon-exon

junctions. Shown are the percentages of splicing events assigned to a particular mode of splicing.

Complex splicing event corresponds to several (two or more) alternative splicing incidents occurring

simultaneously in the same sample. (C) Enrichment analysis of affected signalling pathways by the

exon-exon junctions (DEEj) and differential exome usage (DEU) in the LSAL AML relapse and

myeloid compartment of MPAL patients. Pathway enrichment analysis has been performed with

https://biit.cs.ut.ee/gprofiler/gost under the highest significance threshold, with multiple testing

correction (g:SCS algorithm). (D) Fold change expression levels of total gene among genes identified

to be affected by alternative splicing process (left panel) and differentially spliced variants in lineage

switched and myeloid compartments of MPAL patients (right panel). (E) Schematic representation of

the affected mRNA structure (and its probable consequence depicted in red) within several selected

genes (upper panel). Corresponding normalized expression level (vs reference gene *TBP*) in two

tested lineage switch patients (LS03 and LS04) and one MPAL (lower panel). Shown is the ratio of

analysed splice variant expression level between AML (or myeloid) and ALL (or lymphoid)

populations. (F) Splice variants of *MLL-AF4* identified in MPAL patients and one lineage switch

sample (LS10). Left panel represents junction read counts of the fusion oncogene, identified by the

RNAseq analysis, with confirmation of the expression of both variants analysed by qRT-PCR (MPAL1,

right panel). Both splice variants, further confirmed by Sanger sequencing, showed complete

sequences of *MLL*ex9 or *MLL*e10 and either complete or truncated (for 3 nucleotides at the 5'end)

*AF4*ex4, respectively.

Figure 6

Figure 6. Molecular characterisation of lineage switch *MLL-AF4* leukaemias. (A) Whole exome

sequencing (WES) data showing genes recurrently mutated within the analysed cohort, involved in

the cell growth, communication and metabolism and genes mutated in single cases belonging to the

same function protein complexes (e.g. DNA polymerases, epigenetic complexes). Data are presented

according to the disease timepoint/cell lineage and age of the patient. Depicted are major single

nucleotide variants (SNVs) that were found in >33% of reads and minor SNVs in <33% reads. (B)

Clonal SNV patterns identified by whole genome sequencing (WGS) in LS01 ALL and AML samples,

constructed from counts of each mutation-type at each mutation context, corrected for the frequency

of each trinucleotide in the reference genome. (C) Comparison of the whole exome sequencing and

RNAseq data obtained for LS01 patient identifies 6, 98, and 10 SNVs expressed only in ALL, AML,

and shared between ALL and AML, respectively. 12 SNVs exclusive for the AML relapse, predicted

(by Condel scoring) to have deleterious effects, were subjected to multiplex PCR followed by next-

generation sequencing analysed within each of the purified haematopoietic sub-populations. Circles

with solid colour indicate VAF >30%, light colour and dashed line indicates VAF 5-30%. Remaining

genes (yellow circle) represent the 10 other SNVs (out of 12 SNVs) which showed the same pattern in

the frequency of mutation acquisition.

Figure 7

Figure 7. Epigenetic modulatory genes influence lineage specific expression profiles. (A) CHD4 scheme; the R1068H mutation is located in the critical helicase domain of CHD4 at a highly conserved residue. An * (asterisk) indicates positions which have a single, fully conserved residue, a : (colon) indicates conservation between groups of strongly similar properties - scoring > 0.5 in the Gonnet PAM 250 matrix, a . (period) indicates conservation between groups of weakly similar properties - scoring =< 0.5 in the Gonnet PAM 250 matrix. (B) Fold change in expression of NuRD complex members (*CHD4*, *MTA1*, *RBBP4*, *MBD3*) and *PHF3* following lineage switched relapse (left

panel) and in MPAL cases (right panel). (C) Expression of lineage specific genes following

knockdown of *PHF3*, *CHD4* or the combination, relative to non-targeting control construct in the *MLL-*

AF4 positive ALL cell line, SEM. (D) Flow cytometric analysis of the surface CD33 expression

following knockdown of *PHF3*, *CHD4* or the combination in the SEM cell line. shNTC – non-targeting

control. (E) Gene set enrichment analysis of RNA sequencing data derived from: knockdown of *PHF3*

and *CHD4* in the SEM cell line (left panel) and lineage switch leukaemia cases (right panel). Shown is

negative correlation of shCHD4/shPHF3 and relapse samples with Jaatinen haematopoietic stem cell

signature (upper panel) and Haddad B-lymphocyte progenitor signature (lower panel). (F) Expression

of lineage specific cell surface markers CD19 (lymphoid) and CD33 (myeloid) following culture of

MLL-Af4 transformed hCD34+ cord blood progenitor cells in lymphoid permissive conditions.

Knockdown of *PHF3*, *CHD4* or the combination disrupts the dominant lymphoid differentiation pattern

seen in non-targeting control (shNTC). (G) Assessment of *PHF3* knockdown influence on the surface

marker expression after longer incubation period (33 days); *CHD4* knockdown impaired cellular

survival upon longer *in vitro* culture.