Social niche construction in captive Gouldian Finch (*Erythrura gouldiae*) Flocks

Gregory M. Kohn[1,2], M. Ryan Nugent[2], Xzavier Dail[2], Taylor R. Orlandi[2]

[1]University of North Florida
[2]Jacksonville Zoo and Gardens

Abstract

The organization of animal groups is both a cause of, and consequence of, patterns of interactions among individuals. The stability of animal social organization reflects how individuals construct and maintain resilient patterns of interactions across changes in group size and membership. In this study we describe patterns of social interactions in captive flocks of Gouldian Finches (*Erythrura gouldiae*) that were exposed to changing social conditions. A flock of adult Gouldian Finches was exposed to two changing conditions, an introduction condition where juveniles were introduced to the flock, and a fission-fusion condition where the flock was split into two smaller flocks and a new group of juveniles introduced to each. We show that the social organization of captive Gouldian finch flocks is characterized by stable homophilic communities of adult females and juveniles. Females showed higher rates of approaches to other adult females, while juveniles maintained higher rates of approaches towards other juveniles. These findings highlight how the stable interactive decisions made by individuals contribute to higher-order patterns of organization in bird flocks. Furthermore, this study is one of the few to demonstrate self-assortment of juveniles in passerine flocks and suggests that peer and sibling interactions are an unexplored area in avian social development.

INTRODUCTION

Describing the factors that shape social connections, and how those connections create enduring patterns of organization at the group level is core feature of research on animal social systems (Lott 1991). Most social species inhabit dynamic and loosely cohesive fission-fusion societies where group size and composition fluctuate over short timescales. The stability and organization of groups reflects the ability of individuals to reconstruct patterns of connections across fission-fusion changes (Kohn 2019, Mason 1978). If changing conditions force individuals to "rewire" their connections, the social organization will change, but if individuals respond by reconstructing similar social connections than the social organization will remain resilient. Thus, an important aspect of understanding animal social systems is describing how behavioral responses to period of social change allow the social organization to be continually reconstructed across changing contexts. This study aims to describes the processes maintaining patterns of social organization in an passerine, the Gouldian Finch (*Erythrura gouldiae*).

Homophily reflects preferential associations with like individuals and is a core feature of social systems from economies to small group dynamics (McPherson, Smith-Lovin, & Cook 2001). In animals, variation in sex, age, familiarity, morphology and personality play a large role in shaping who associates with whom (Ebenau, 2019; Kohn, King, Scherschel, & West, 2011). In some species, homophily is caused by morphological and physiological differences resulting in divergent patterns of activity synchronization between individuals (Conradt, Ruckstuhl, & Neuhaus, 2005). During parts of the year, this can result complete segregation, with individuals of different sex and ages classes inhabiting separate groups (Ruckstuhl & Neuhaus, P. 2000). More frequently, incomplete social assortment is observed in groups. While a diverse range of individuals inhabit the same group, distinct subgroups and communities emerge as individuals have stronger social connections with like conspecifics and other group members. While such subgroups are prevalent throughout the vertebrates, little is known about the behavioral processes underlying the emergence and stability of these subgroups across group changes.

Assortment by sex has been documented extensively in social mammals and increasingly in birds and invertebrates. Across taxa there has been a tendency for females to form strong connections with other females, whereas males disperse to neighboring groups (Ward & Webster, 2016). In mammals (Stanley, Liddiard Williams, & Preziosi, 2018), birds (Kohn, King, Scherschel, & West, 2011), and insects (Stanley, Williams, Preziosi 2017) females have been observed to form cohesive female subgroups within the larger group. Strong connections among familiar females often have beneficial consequences for females such as reduced risk from male harassment (Cameron, Setsaas & Linklater,. 2009), higher offspring survival (Silk et al. 2009), lowering stress responses, and facilitating access to social information (Gros-Louis, White, King & West 2003). In horses (*Equus ferus*) and yellow baboons (*Papio cynocephalus*), strong and stable connections between females are associated with higher offspring survival and lower mortality (Carter, Lee, Marshall, Ticó, & Cowlishaw, 2015; Silk et al. 2010, 2009). In Brown-headed Cowbirds (*Molothrus ater*) females form stable subgroups with familiar females across group changes, and females with the strongest preference for familiar females during the fall have higher rates of reproductive output during the spring (Kohn, 2017).

Age is a significant determinant of social connections in many animal societies. While
extensively documented in primates, the existence of juvenile groups and subgroups has been
observed in animals ranging from fish to equids to invertebrates. Many factors can facilitate the
emergence of juvenile assortment independent of mutual attraction between juveniles. Juveniles
often show distinct morphological and physiological differences from adults, and this can cause
juveniles to cluster based on shared physiological needs and habitat preferences. Some studies
have shown that juvenile form "gangs" around contested resources to better compete with more
experienced adults (Dall & Wright, 2009; Franks et al. 2020) Adults may also corral their
offspring into all-juvenile groups for protection from predation and preferentially avoid
interacting with the more active and disruptive juveniles (Munro, & Bédard,1977; Wanker,
Bernate & Franck,1996). In summary, while juvenile groups are commonly observed across taxa
the behavioral processes underlying their formation and stability is largely unknown.
149
Juveniles can self-assort by preferring to interact with similar-aged peers who are more
behaviorally compatible than adults (Lindgren & Shapiro, 1990). Goat kids (*Capra aegagrus*
hircus) will form subgroups with peers two weeks after birth by preferring interactions with
peers over parents (Lickliter, 1987). This pattern is observed across many equids (Boyd, 1988)
and other social mammals. In birds little is known about the social relationships, experiences,
and interaction preferences of juveniles upon becoming independent. While some have suggested
that birds associate with siblings and peers for an extended time after independence (Hötker,
1982; Loretto, Fraser, & Bugnyar, 2012; Nicolai, 1956), it's unknown if juvenile subgroups exist
in many species, and if mutual interaction preferences play a role in their formation. One aim of
the current study is to describe patterns of age-based assortment in a Gouldian finch flocks and
investigate if shared interaction preferences among juveniles lead to the formation of juvenile
subgroups.
162
Peer interactions among juveniles are often necessary for the development of species-typical
behaviors and competencies. When isolated from peers, juveniles often develop distinct
behavioral repertoires and have long-lasting behavioral deficiencies when compared to typically
reared individuals (Suomi & Harlow, 1972). Juveniles are much more active, playful, and social
than adults, and juvenile subgroups may allow for young individuals to explore their behavioral
potentials and refine their behavioral abilities. Ultimately behaviors acquired through peer
interactions can allow juveniles to become socialized into the wider group (Pellis & Pellis, 2007;
Poirier & Smith, 1974). For example, in Brown-headed cowbirds, interactions within juvenile
subgroups predicts the development of later courtship competence used to secure pair bonds as
adults (Kohn, King, Dohme, Meredith & West, 2013), and rhesus macaques (*Macaca mulatta)*
infants raised in social isolation were able to acquire species-normative social behavior after an
extensive period of social interaction with peers (Suomi & Harlow, 1972).
175
This study aims to describe the social organization in flocks of captive Gouldian finches.
Gouldian finches are an endangered estrildid finch that inhabits the subtropical grasslands of
northern Australia. While little is known about the social organization or behavior of Gouldian
finches, in the wild, large fission-fusion flocks composed of hundreds of individuals of mixed
age and sex composition are commonly observed. The fluidity and diversity of estrildid flocks
has led some to speculate that they lack significant patterns of within-group structure (Bolton, et
al. 2016; Dostine, Johnson, Franklin, Zhang, Hempel, 2001).

183
Gouldian finches also exhibit a startling color polymorphism. Adults show discrete differences in
head coloration with some having either red, black, or yellow heads, while juveniles have grey
plumage with olive backs. Differences in head coloration correspond with differences in
behavior and personality. Red headed individuals are more aggressive, with fights between red-
headed individuals being more intense than others, and red-headed females displacing black-
headed females more often (Pryke, 2007). In the wild, black-headed individuals are the first to
descend to waterholes (O'Reilly, Hofmann & Mettke-Hofmann, 2019). In captivity, some studies
have shown that black-headed individuals were more likely to approach and interact with a novel
object faster and return to a food bowl sooner after the presentation of a simulated predator
(Williams, King, & Hoffman 2012, Mettke-Hofmann 2012), while others have found that red-
headed individuals are more neophilic in other contexts (Fragueira & Beaulieu, 2019).
195
It is currently unknown how the different color morphs, sexes, and age classes associate with
each other in a semi-naturalistic context. Due to the behavioral differences across individuals of
various sex, age, and head-color status, it is hypothesized that Gouldian finch flocks will show
significant assortment by both age, sex, and color morph status. Research on animal social
networks often focuses on associations, where social connections are measured when two or
more individuals inhabit the same place in close proximity. While associations can be powerful
measures of social organization, the social choices that build connections can be difficult to
measure without observing directed interactions between known individuals. By recording
interaction preferences, we can discover how these social choices construct higher-order patterns
of organization (Hinde, 1976). Who individuals prefer and avoid engaging will often directly
contribute to the communities, subgroups, hierarchies at the group and population level. By
describing the stable social preferences creating patterns of assortment in changing Gouldian
Finch flocks this study will highlight the behavioral mechanisms generating the social network
organization by discovering *stable* social preferences consistently expressed across different
contexts which correspond to higher order structural attributes of group organization.
211
212 **METHODS**
213
Subjects: All individuals in this study were Gouldian finches. All finches were acquired from
professional breeders in Arizona and Florida U.S.A. Each bird was provided with a uniquely
colored leg band for individual recognition. Males, females, and juveniles were distinguished
from each other based on plumage and behavior differences, with adult females being much
more subdued in their chest coloration in contrast to males.
All juveniles were hatched and raised in separate cages prior to being introduced to
experimental flocks. At around 40 days after fledging juveniles become independent of their
parents and can feed and drink on their own without any sign of parental provisioning. Juveniles
were distinguished from adults based on their grey head's olive back and lack of typical adult
plumate. In this study all juveniles were under one year of age since fledging the nest and adults
were older than a year and exhibited adult plumage. Head color was easily observed in adults
and assessed in juveniles at the time of their first molt after the end of the data collection period.
We used a total of 5 adult males and 6 females and 10 juveniles for this study.

All birds were provided with a daily diet consisting of finch seed mix (Kaytee) and
soaked Perle Morbide pellets, with eggshell, boiled eggs, and dandelion greens provided twice a
week. Fresh water was continually available.

Aviaries: We used a series of two aviaries in this study. Each aviary consisted of identical
dimensions (23 x 72 x 41 in) and similar environmental conditions. The aviaries were located
inside next to a large window allowing for sunlight and ventilation. Each aviary contained both
natural eucalyptus and plastic privacy perches, small shrubs, a feeding station, and a bowl for
water baths.

Data Collection: We used focal sampling during data collection. During observation periods all
individuals in the flock were observed for a 5-minute focal block. We recorded all approaches,
occurred within a 2-inch radius around the focal individual being observed. All observations
were done using voice recognition technology allowing the observer to maintain continual visual
contact with the focal bird throughout the sampling block. An approach was scored when another
individual approached another within a radius of 2 inch. Before data collection, video of marked
perches was made to establish that 2 inches was the typical social distance observed between two
resting adult finches. All focal samples were collected by GMK.

Procedure

Baseline Session: From 8/7/2020 to 8/17/2020 focal samples were taken measuring approach
behavior. A total of 352 focal samples were taken. Each individual was observed 32 blocks for a
total of 160 minutes each.

Fusion session: On 10/5/2020 to four novel juveniles (40 days post fledging) were introduced
into the aviary. From 10/06/2020 to 12/22/2020 an additional 503 focal samples were taken. All
individuals observed 34 blocks for a total of 170 minutes except one individual who was
removed from the flock temporarily for health reasons and was observed for 26 blocks.

Fission-Fusion Session: On 2/10/2021 the fusion flock was split up into two separate flocks.
Three males, two females, and two juveniles were moved into flock 1, and four females, two
males, and one juvenile were moved to flock 2. In each flock, three new juveniles (40 days post
fledging) were randomly introduced to both flocks. From 2/11/2021 to 4/20/2021 a total of 84
blocks were recorded in each aviary. Each individual was observed for 8 blocks for a total of 40
minutes each.

ANALYSIS

Due to the small non-independent sample sizes, permutation statistics was used. Permutation
tests are designed to deal with the unique challenges of small non-independent and sample sizes
while maintaining an appropriate type I error rate.

Modeling social network structure: Generative network modeling and clustering algorithms were
used to model social network structure. We used exponential random graph models (hereafter
ERGMs) for valued and directed networks (R package ergm.count in the statnet suite). ERGMs

are like generalized linear models, except the dependent variable is the distribution of
connections in the social network (the network topology) and the explanatory variables are
statistical terms that assess the influence social processes on the distribution of connections in the
network (Silk, M. J., & Fisher, D. N. 2017). We conducted an ERGM for each seperate flock in
each condition, and for all models the terms included as explanatory factors were the influence of
age, sex, and head-color based homophily.
279
When a model identified any significant homophilous terms a post hoc clustering algorithm was
used to look at the subgroup structure of the network. A subgroup is a set of individuals who
share more ties within the community then they do outside the community. A spinglass cluster
algorithm was used to identify subgroups. This algorithm is suited to smaller networks with
weighted and directed connections between individuals and seeks to rewards ties between
community members while minimizing edges for within community members and places an
emphasis on missing edges between individuals (Pasta & Zaidi, 2017).
287
Permutation tests for social structure: After conducting ERGMs and cluster analysis,
permutation tests were used to assess the significance and direction of social preferences. A null
distribution was computed for each flock by doing row-wise permutations. For each permutation,
the number and distribution of approaches directed towards others was preserved but the
individual they were directed towards was randomized. A total of 100,000,000 permuted
networks were generated. For each permuted network, the approach rate per-individual toward
the different age, sex, and head color categories was calculated by taking the total number of
approaches directed towards those categories and dividing it by the number of group members in
those categories. The null distributions reflected the approach rates across all permutations.
Observed interaction rates were then compared to the null distribution and empirical p-values
were calculated to assess if specific approach rates occurred above or below expected values.
These p-values were calculated by taking the number of observations either above or below the
observed value and dividing them by the total number of permutations. All permutation tests
used a two-tailed significance with an alpha of 0.05 (above or below 0.025 in each tail)
Pairwise differences in approach rates were used to see if individuals maintained higher
rates of approaches toward like individuals in contrast to other group members. For each
individual we subtracted the approach rate directed toward homophilous conspecifics from
approaches directed towards other conspecifics and compared the mean of those differences to a
permuted null distribution. For example, to see if females preferred females over males we
subtracted the approaches to males from the approaches to females. This observed mean
difference was compared to null distributions of mean differences across the permuted networks.
Empirical p-values were calculated to show if approach rates toward homophilous conspecifics
were significantly higher than expected with an alpha of 0.05. All permutation tests were also
conducted using data stream permutations with equivalent results (Farine, 2017), and multiple
comparisons were corrected using the False Discovery Rate (FDR) correction.
313
314 *Repeatability of adult network structure:* To assess if social networks remained resilient to group
315 changes, we specifically looked at the adult network from baseline to the introduction period. To
316 see if the adult network was significantly robust to the introduction of juveniles a uniform graph
317 correlation (gscor in statnet) was performed with a conditioned uniform graph test (cugtest in
318 statnet) being used to calculate empirical-p values.

319
Results:

Basic: During the baseline period a total of 5,655 approaches were recorded. After the baseline
period a group of four juveniles were added to the group and a total number of 8840 approaches
were recorded. After the this a fission-fusion was performed where the flock was split up into
two separate flocks and three juveniles added to each one, here a total of 2184 approaches were
observed in flock 1 and 2345 approaches were observed in flock two.

Social network structure: Our Gouldian finch flock was organized around sex and age-based
preferences. During both the baseline and introduction period the models found significant
patterns of social assortment. During the baseline, fusion, and fission-fusion periods the ERGM
model found significant assortment by both sex, age, and head color, with assortment by sex
showing the strongest positive coefficient in all conditions (Table 1).

Community Structure: During the baseline period the clustering algorithm identified two
communities, one consisting of the six females and the other consisting of the five males (Fig 1).
During the fusion condition, three different communities were identified, one community
containing the six females, another contained all five males, and the other contained all four
juveniles. During the fission-fusion period, distinct communities were observed in both flocks. In
the first flock, three communities were observed, one containing three adult individuals including
both adult females and their paired male, another community contained the three juveniles that
were introduced to the flock during this period, and the final community contained two of the
juveniles introduced during the fusion period and two adult males. The second flock during the
fission-fusion condition showed two distinct communities, one containing only adults including
four adult females and two adult males, and another community containing all four juveniles.

Permutation tests for social preferences: In all conditions, social preferences differed
significantly from the null expectations. Females had significantly higher rates of approaches
towards other females, and juveniles had significantly higher rates of approaches towards other
juveniles in all flocks. Adult males showed significant preference to interact with females in all
conditions (Table 2). In all conditions, juveniles showed a tendency to avoid interacting with
other adult males, but no preference towards approaching or avoiding adult females.
In both the baseline and fusion conditions, females were significantly more likely to
interact with females over males and juveniles (Fig 1, Table 3). During the fission-fusion period,
the sex ratio became skewed towards females (Flock 2) or males (Flock1) and females showed a
tendency to prefer females over male interactions when the ratio was skewed towards males in
flock 1. Males showed no consistent pairwise social preferences except with preferring to
interact with females over males in flock 1 during the fission-fusion condition. Across all
conditions juveniles showed a significant preference to interact with juveniles over adults (Table
3, Figure 1). Across all condition we found no consistent patterns in the social preferences for
head colors (Sup Materials 1).

Repeatability of network structure: Adults were able to maintain the connections they had with
others between the baseline and introduction period. Structural graph correlation showed high

positive correlations between the baseline and introduction adult networks (gscor = 0.79, CUG
test: p = 0.00001).
366
367 **DISCUSSION**
368
This study described the fine-scale structure of social approaches across changing Gouldian finch
flocks. Our results showed that the social organization of Gouldian finch flocks is characterized
by sex and age-based assortment patterns. These patterns were robust to changes in group size
and composition, with higher levels of female and juvenile homophily occurring in almost every
flock we observed. While social assortment can be influenced by a wide range of social and non-
social factors, our results suggests that these patterns are driven by the social decisions of
juveniles and females. Females maintained higher levels of female-to-female approaches than
expected, and significantly preferred to approach females over others in most flocks. Across all
conditions, juvenile-to-juvenile approach rates were higher in comparison null expectations, and
juveniles preferred to approach other juveniles over adults.
379
The ability to navigate the risks and benefits of interactions is essential for success in the group
(Farine, et al. 2015). There can be significant risks when interacting with novel or aggressive
conspecifics, and individuals may buffer against these interactions by becoming selective in
whom they approach. Some interactions can have significant benefits, such as access to
beneficial social information or experiences necessary for the development of competent
behavioral abilities (West & King, 2002). In such cases, individuals may be selective by
preferentially interacting with individuals due to the beneficial consequences of those
interactions (Hausberger, 1997; Mann & Slater, 1995). To understand the potential risks and
benefits of interactions, we must first have a detailed description of how group members allocate
their social decisions in the group to see which individuals are preferred or avoided.
390
Our study found that Gouldian finch females preferentially formed subgroups with other
females. In many mammals, females form stable groups with other females whereas males
disperse to new groups or all male groups (Ruckstuhl, 2007). Increasingly similar patterns are
also being discovered in birds. In red junglefowl, pheasants, brown-headed cowbirds, and others
female birds often form cohesive groups and subgroups while males do not (Kohn, King,
Scherschel & West, 2011; McDonald, Spurgin, Fairfield, Richardson & Pizzari, 2019;
Whiteside, van Horik, Langley, Beardsworth, Capstick & Madden, 2019).
398
Sex assortment can occur through various behavioral mechanisms. Some have suggested that
females prefer to interact with other females to buffer against destabilizing interactions with
males. Studies in baboons, horses, and others have shown that the presence of strong stable
female relationships are associated with increased survival and fitness (Cameron, Setsaas &
Linklater, 2009; Carter, Lee, Marshall, Ticó & Cowlishaw, 2015; Silk, et al. 2009, 2010) and
studies in the cockroach *Diploptera punctata* and guppies have suggested that female
aggregations limit intrusions by males (Stanley, Liddiard Williams & Preziosi, 2018). If male
interactions have a destabilizing effect, we should also observe that they are preferentially
avoided. However, in this study we found that while females interacted more often with other
females than expected, male interactions did not consistently occur less likely than expected,
suggesting that female preferences may carry other motivations for avoiding males.

410
Females will also preferentially associate with other females to access valuable social
information. Cues displayed by females can communicate mate and song preferences and
exposure to these cues can shape reproductive, competitive, and social decisions in females
(Freed-Brown & White, 2009; West, King, White, Gros-Louis & Freed-Brown, 2006). In brown-
headed cowbirds, females acquire mate preferences through interactions with other females, and
females who were more familiar with each other during the fall are more competitive with each
other during the breeding season (Kohn, in prep). Cowbird flocks also segregate according to age
and sex in response to specific stimuli, such as playback of specific songs (Gros-Louis, White,
King & West, 2003). While little is known about the mate preferences of Gouldian finches,
relationships between females may allow females to monitor the preferences of others. In our
flock two of the strongest connected females (FYER and FB2G) were both paired with the same
male (MBNB). While further research should investigate the broader consequences of female-to-
female interactions, our findings suggests that female interactions may be preferred beyond
simply avoiding males.
425
Our study documented a consistent pattern of age assortment in Gouldian finch flocks, with
juveniles preferring interactions with other juveniles across flock changes. Most studies of avian
development focus on adult-juvenile interactions to understand the horizontal transmission of
social information from adults to juveniles. Nonetheless, it is increasingly recognized that peer
and sibling interactions are essential for species-typical development, and isolation from peers
causes behavioral deficiencies that negatively impact performance in a wide range of social
contexts, from integration into groups to reproductive success (Garnetzke-Stollmann & Franck,
D. 1991; Hudson, Bautista, Reyes-Meza, Montor & Rödel, 2011; Liévin-Bazin, Pineaux, Le
Covec, Gahr, Bovet & von Bayern 2019). While studies of peer interactions have been largely
restricted to social mammals, especially primates, this study suggests that peer interactions are an
unexplored area in avian social development. Little is known about sibling and peer interactions
in juvenile birds, especially after individuals become independent of their parents. Nonetheless,
when followed beyond independence, studies have shown that birds often sustain connections
with siblings and peers until sexual maturity .
440
A diverse range of factors may cause the formation of sibling and peer groups in juvenile birds.
Some have suggested that competition around patchy food resources may drive juvenile
subgroups, in which juveniles form "gangs" with other juveniles that allow them to compete with
others for access to limited food sites. Studies have found that in juvenile hihi (*Notiomystis
cincta*) and ravens (*Corvus corax*), individuals preferentially associate with juveniles around
contested spaces (Franks, Ewen, McCready, Rowcliffe, Smith, D & Thorogood, 2020; Marzluff,
Heinrich & Marzluff, 1996). Others have suggested that social groups formed to protect
individuals from predation. In some birds such as spectacled parottlets, parents bring their
offspring to areas with unrelated juveniles that serve as crèches, suggesting adults play the causal
role in the formation of subgroups (Garnetzke-Stollmann & Franck, 1991, Velando, 2001,
Wanker, Bernate, & Franck, 1996). In other cases, adults avoid or are aggressive towards
juveniles, forcing juveniles to preferentially associate with peers.
453
Our results suggest that juvenile Gouldian finches self-assort by preferentially choosing to
interact with siblings and peers after becoming independent of their parents. This is among the

few examples of self-assortment of juveniles within passerine flocks (Hötker, 1982; Kohn, King,
Scherschel, & West, 2011; Nicolai, 1956; Templeton, Reed, Campbell & Beecher, 2012). While
adults did tend to avoid interactions with juveniles, juveniles significantly preferred to interact
with each other over adults. Both male and female juveniles spend considerable time engaging in
peering displays and exchanging courtship-like hopping and head shaking displays with other
juvenile siblings in a play-like manner (Kohn, personal observation). Play behavior is expressed
preferentially in juveniles and allows individuals to acquire the skills to dynamically adjust
behavior to ambiguous social cues (Pellis, Pellis & Bell, 2010). Thus, juvenile subgroups may
facilitate the horizontal co-construction of social skills necessary to navigate interactions during
later periods (Kohn, 2019). While play has been documented in birds (Diamond & Bond, 2003;
Pellis, 1981) its extent and influence in juvenile passerines is not fully known. Current studies
are looking at the specific details of juvenile-to-juvenile interactions and comparing it to the
interactions between juveniles and adults.
469
Head color did not seem to be an important factor in shaping the social organization of Gouldian
finch flocks. While significant in the ERGM model, further analysis in each condition showed no
clear pattern in either approach or avoidance between individual of differing head colors. This
result held both when juveniles were included with their eventual adult plumage, and when
juveniles were removed form the analysis. However, in this study, head colors were biased
according to the sex of the individual, with black being more prominent in females and red being
more prominent in males. Further studies will likely be needed to conclusively determine the
extent that head color shapes social networks in this species.
478
From guiding the ontogeny of species-typical abilities to constraining patterns of cultural
evolution across populations, the social organization of groups is the pivot upon which many
social phenomenon orbit. Describing the social decisions individuals make in fluid naturalistic
conditions is essential to understanding the factors underlying the emergence of group
organization. This study is among the first to show that the behavioral decisions made across
age-sex classes reconstruct patterns of social organization in fission-fusion Gouldian finch
groups. As birds are heavily dependent on socially transmitted information this study suggests
that the emergence of an organized network of social connections provides the structured
environment necessary the development of avian behavioral competencies.
488

489 **ACKNOWLEDGMENTS**
490
491 This study was supported with a grant from the Van Vleck Award. All research was done in
492 accordance with ASAB/ABS Guidelines for the Use of Animals in Research and was approved
493 by the University of North Florida's Institutional Animal Care and Use Committee protocol
494 number 20-006.

495
496
497   REFERENCE
498

Bildstein, K. L., Frederick, P. C., & Spalding, M. G. (1991). Feeding patterns and aggressive
behavior in juvenile and adult American flamingos. *The Condor*, *93*(4), 916-925.

Bolton, P. E., West, A. J., Cardilini, A. P., Clark, J. A., Maute, K. L., Legge, S., ... & Rollins, L.
A. (2016). Three molecular markers show no evidence of population genetic structure in the
Gouldian finch (Erythrura gouldiae). *PloS one*, *11*(12), e0167723.

Boyd, L. E. (1988). Ontogeny of behavior in Przewalski horses. *Applied Animal Behaviour*
Science, *21*(1-2), 41-69.

Carter, A. J., Lee, A. E., Marshall, H. H., Ticó, M. T., & Cowlishaw, G. (2015). Phenotypic
assortment in wild primate networks: implications for the dissemination of information. *Royal*
Society Open Science, *2*(5), 140444.

Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between unrelated
females increase reproductive success in feral horses. *Proceedings of the National Academy of*
Sciences, *106*(33), 13850-13853.

Chambers, M. S., Sidhu, L. A., O'Neill, B., & Sibanda, N. (2017). Evidence of separate
subgroups of juvenile southern bluefin tuna. *Ecology and evolution*, *7*(22), 9818-9844.

Clutton-Brock, T., & Janson, C. (2012). Primate socioecology at the crossroads: past, present,
and future. *Evolutionary Anthropology: Issues, News, and Reviews*, *21*(4), 136-150.

Conradt, L., Ruckstuhl, K. E., & Neuhaus, P. (2005). Definitions, hypotheses, models and
measures in the study of animal segregation. *Sexual segregation in vertebrates*, 11-32.

Dall, S. R., & Wright, J. (2009). Rich pickings near large communal roosts favor 'gang'foraging
by juvenile common ravens, Corvus corax. *PloS one*, *4*(2), e4530.

Darden SK, Croft DP (2008). Male harassment drives females to alter habitat use and leads to
segregation of the sexes. Biology letters, 4(5), 449–451.

Diamond, J., & Bond, A. B. (2003). A comparative analysis of social play in birds. *Behaviour*,
1091-1115.

Dostine, P. L., Johnson, G. C., Franklin, D. C., Zhang, Y., & Hempel, C. (2001). Seasonal use of
savanna landscapes by the Gouldian finch, Erythrura gouldiae, in the Yinberrie Hills area,
Northern Territory. *Wildlife Research*, *28*(4), 445-458.

Ebenau, A., von Borell, C., Penke, L., Ostner, J., & Schülke, O. (2019). Personality homophily
affects male social bonding in wild Assamese macaques, Macaca assamensis. *Animal*
Behaviour, *155*, 21-35.

Edgar, J., Held, S., Paul, E., Pettersson, I., Price, R. I. A., & Nicol, C. (2015). Social buffering in
a bird. *Animal behaviour*, *105*, 11-19.

Farine, D. R. (2017). A guide to null models for animal social network analysis. *Methods in*
Ecology and Evolution, *8*(10), 1309-1320.

Farine, D. R., Firth, J. A., Aplin, L. M., Crates, R. A., Culina, A., Garroway, C. J., ... & Sheldon,
B. C. (2015). The role of social and ecological processes in structuring animal populations: a
case study from automated tracking of wild birds. *Royal Society Open Science*, *2*(4), 150057.

Fragueira, R., & Beaulieu, M. (2019). Red does not always outperform black: morph-specific
behavioural variation in response to environmental changes. *Animal Behaviour*, *148*, 81-91.

Franks, V. R., Ewen, J. G., McCready, M., Rowcliffe, J. M., Smith, D., & Thorogood, R. (2020).
Analysing age structure, residency and relatedness uncovers social network structure in
aggregations of young birds. *Animal Behaviour*, *166*, 73-84.

Freed-Brown, G., & White, D. J. (2009). Acoustic mate copying: female cowbirds attend to other
females' vocalizations to modify their song preferences. *Proceedings of the Royal Society B:
Biological Sciences*, *276*(1671), 3319-3325.

Garnetzke-Stollmann, K., & Franck, D. (1991). Socialisation tactics of the spectacled parrotlet
(*Forpus conspicillatus*). *Behaviour*, *119*(1-2), 1-29.

Gros-Louis, J., White, D. J., King, A. P., & West, M. J. (2003). Female brown-headed
cowbirds'(Molothrus ater) social assortment changes in response to male song: a potential source
of public information. *Behavioral Ecology and Sociobiology*, *53*(3), 163-173.

Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, 1-17.

Hausberger, M. (1997). Social influences on song acquisition and sharing in the European
starling (Sturnus vulgaris).

Hötker, H. (1982). Studies of the behaviour of young meadow pipits (Anthus pratensis) after
fledging. *Vogelwelt*, *103*, 1-16.

Hudson, R., Bautista, A., Reyes-Meza, V., Montor, J. M., & Rödel, H. G. (2011). The effect of
siblings on early development: a potential contributor to personality differences in
mammals. *Developmental psychobiology*, *53*(6), 564-574.

Kohn, G. M. (2017). Friends give benefits: autumn social familiarity preferences predict
reproductive output. *Animal behaviour*, *132*, 201-208.

Kohn, G. M., King, A. P., Dohme, R., Meredith, G. R., & West, M. J. (2013). Robust autumn
social attributes predict spring courtship skills in juvenile female brown-headed cowbirds,
Molothrus ater. *Animal Behaviour*, *85*(4), 727-732.

Kohn, G. M., King, A. P., Scherschel, L. L., & West, M. J. (2011). Social niches and sex
assortment: uncovering the developmental ecology of brown-headed cowbirds, Molothrus
ater. *Animal Behaviour*, *82*(5), 1015-1022.

Liévin-Bazin, A., Pineaux, M., Le Covec, M., Gahr, M., Bovet, D., & von Bayern, A. M. (2019). Food sharing and affiliation: an experimental and longitudinal study in cockatiels (Nymphicus hollandicus). *Ethology*, *125*(5), 276-288.

Lickliter, R. E. (1987). Activity patterns and companion preferences of domestic goat kids. *Applied Animal Behaviour Science*, *19*(1-2), 137-145.

Lindgren, C. J., & Shapiro, L. J. (1990). Preferences of mallard ducklings for strange broods of different ages. *Bird Behavior*, *8*(2), 79-86.

Loretto, M. C., Fraser, O. N., & Bugnyar, T. (2012). Ontogeny of social relations and coalition formation in common ravens (Corvus corax). *International journal of comparative psychology/ISCP; sponsored by the International Society for Comparative Psychology and the University of Calabria*, *25*(3), 180.

Lott, D. F. (1991). *Intraspecific variation in the social systems of wild vertebrates* (Vol. 2). Cambridge University Press.

McDonald, G. C., Spurgin, L. G., Fairfield, E. A., Richardson, D. S., & Pizzari, T. (2019). Differential female sociality is linked with the fine-scale structure of sexual interactions in replicate groups of red junglefowl, Gallus gallus. *Proceedings of the Royal Society B*, *286*(1913), 20191734.

Mann, N. I., & Slater, P. J. B. (1995). Song tutor choice by zebra finches in aviaries. *Animal Behaviour*, *49*(3), 811-820.

Marzluff, J. M., Heinrich, B., & Marzluff, C. S. (1996). Raven roosts are mobile information centres. *Animal Behaviour*, *51*(1), 89-103.

Mason, W. A. (1978). Ontogeny of social systems. *Recent advances in primatology*, *1*, 5-14.

McPherson, M., Smith-Lovin, L., & Cook, J. M. (2001). Birds of a feather: Homophily in social networks. *Annual review of sociology*, *27*(1), 415-444.

Mettke-Hofmann, C. (2012). Head colour and age relate to personality traits in Gouldian finches. *Ethology*, *118*(9), 906-916.

Mettke-Hofmann, C., Eccles, G., Greggor, A., & Bethell, E. J. (2020). Cognition in a changing world: Red-headed Gouldian finches enter spatially unfamiliar habitats more readily than do black-headed birds. *Frontiers in Ecology and Evolution*, *8*.

Munro, J., & Bédard, J. (1977). Crèche formation in the common eider. *The Auk*, *94*(4), 759-771.

Nicolai, J. (1956). Zur Biologie und Ethologie des Gimpels (*Pyrrhula pyrrhula L.)* D. 77. *Zeitschrift für Tierpsychologie*, *13*(1), 93-132.

640 O'Reilly, A. O., Hofmann, G., & Mettke-Hofmann, C. (2019). Gouldian finches are followers
641 with black-headed females taking the lead. *Plos one*, *14*(4), e0214531.
642
643 Pasta, M. Q., & Zaidi, F. (2017). Topology of complex networks and performance limitations of
644 community detection algorithms. *IEEE Access*, *5*, 10901-10914.
645
646 Pellis, S. M. (1981). A description of social play by the Australian magpie Gymnorhina tibicen
647 based on Eshkol-Wachman notation. *Bird Behavior*, *3*(3), 61-79.
648
649 Pellis, S. M., Pellis, V. C., & Bell, H. C. (2010). The function of play in the development of the
650 social brain. *American Journal of Play*, *2*(3), 278-296.
651
652 Pellis, S. M., & Pellis, V. C. (2007). Rough-and-tumble play and the development of the social
653 brain. *Current directions in psychological science*, *16*(2), 95-98.
654
655 Poirier, F. E., & Smith, E. O. (1974). Socializing functions of primate play. *American
656 Zoologist*, *14*(1), 275-287.
657
658 Pryke, S. R. (2007). Fiery red heads: female dominance among head color morphs in the
659 Gouldian finch. *Behavioral Ecology*, *18*(3), 621-627.
660
661 Romey WL, Wallace AC (2007) Sex and the selfish herd: sexual segregation within nonmating
662 whirligig groups. Behav Ecol 18(5):910–915. doi: 10.1093/beheco/arm057
663
664 Ruckstuhl, K. E. (2007). Sexual segregation in vertebrates: proximate and ultimate
665 causes. *Integrative and Comparative Biology*, *47*(2), 245-257.
666
667 Ruckstuhl, K. E., & Neuhaus, P. (2000). Sexual segregation in ungulates: a new
668 approach. *Behaviour*, 361-377.
669
670 Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., ... &
671 Cheney, D. L. (2009). The benefits of social capital: close social bonds among female baboons
672 enhance offspring survival. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1670),
673 3099-3104.
674
675 Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., ... &
676 Cheney, D. L. (2010). Strong and consistent social bonds enhance the longevity of female
677 baboons. *Current biology*, *20*(15), 1359-1361.
678
679 Silk, M. J., & Fisher, D. N. (2017). Understanding animal social structure: exponential random
680 graph models in animal behaviour research. *Animal Behaviour*, *132*, 137-146.
681
682 Stanley, C. R., Liddiard Williams, H., & Preziosi, R. F. (2018). Female clustering in cockroach
683 aggregations—A case of social niche construction?. *Ethology*, *124*(10), 706-718.
684

685 Stanley, C. R., Mettke-Hofmann, C., Hager, R., & Shultz, S. (2018). Social stability in semiferal
686 ponies: networks show interannual stability alongside seasonal flexibility. *Animal*
687 *Behaviour*, *136*, 175-184.
688
689 Suomi, S. J., & Harlow, H. F. (1972). Social rehabilitation of isolate-reared
690 monkeys. *Developmental Psychology*, *6*(3), 487.
691
692 Templeton, C. N., Reed, V. A., Campbell, S. E., & Beecher, M. D. (2012). Spatial movements
693 and social networks in juvenile male song sparrows. *Behavioral Ecology*, *23*(1), 141-152.
694
695 Velando, A. (2001). Postfledging creche behavior in the European shag. *Journal of*
696 *Ethology*, *19*(2), 121-127.
697
698 Wanker, R. (1999). Socialization in spectacled parrotlets (Forpus conspicillatus): how juveniles
699 compensate for the lack of siblings. *acta ethologica*, *2*(1), 23-28.
700
701 Wanker, R., Bernate, L. C., & Franck, D. (1996). Socialization of spectacled parrotlets Forpus
702 conspicillatus: the role of parents, creches and sibling groups in nature. *Journal für*
703 *Ornithologie*, *137*(4), 447-461.
704
705 Ward, A., & Webster, M. (2016). Sociality: the behaviour of group-living animals.
706
707 West, M. J., & King, A. P. (2002). The ontogeny of competence. In *Conceptions of*
708 *development* (pp. 77-105). Psychology Press.
709
710 West, M. J., King, A. P., White, D. J., Gros-Louis, J., & Freed-Brown, G. (2006). The
711 development of local song preferences in female cowbirds (Molothrus ater): Flock living
712 stimulates learning. *Ethology*, *112*(11), 1095-1107.
713
714 Whiteside, M. A., van Horik, J. O., Langley, E. J., Beardsworth, C. E., Capstick, L. A., &
715 Madden, J. R. (2019). Patterns of association at feeder stations for Common Pheasants released
716 into the wild: sexual segregation by space and time. *Ibis*, *161*(2), 325-336.
717
718 Williams, L. J., King, A. J., & Mettke-Hofmann, C. (2012). Colourful characters: head colour
719 reflects personality in a social bird, the Gouldian finch, Erythrura gouldiae. *Animal*
720 *Behaviour*, *84*(1), 159-165.

Table 1: Results from the Exponential Random Graph Models

Condition	Homophilly	Coefficient	s-Error	z-value
Baseline	Sex	3.58345	0.035	102.38*
	Age	0.62427	0.03701	16.87*
	Head Color	0.44437	0.03884	11.44*
Fusion	Sex	3.26371	0.02299	141.99*
	Age	0.71065	0.02992	23.75*
	Head Color	0.87981	0.03046	28.89*
Fission-Fusion Flock 1	Sex	2.18036	0.05905	36.92*
	Age	1.42661	0.05497	25.95*
	Head Color	0.9629	0.05097	18.89*
Fission-Fusion Flock 2	Sex	2.17984	0.06002	36.32*
	Age	1.42576	0.05448	26.17*
	Head Color	0.96428	0.05127	18.81*

Results from the ERGM model for the baseline, fusion and fission-fusion flocks. The * symbol indicates a significant effect below 0.0001.

758   Table 2: Permutation tests for Interaction Types
759

Condition	Interaction	Mean	97.5 Int	p-value Up	p-value Down
Baseline	F-F	66	39 -53	< 0.0001*	0.99
	F-M	18	34-55	0.65	0.35
	M-F	58	37-61	0.06	0.95
	M-M	38	35 - 63	0.92	0.07
Introduction	F-F	80.97	33-57	< 0.0001*	1.00
	F-M	13.46	30-59	1.00	< 0.0001*
	F-J	28.62	29-61	0.97	0.02*
	M-F	57.36	25-51	< 0.0001*	0.99
	M-M	28.24	24-54	0.89	0.11
	M-J	21.15	23-58	0.01*	0.99
	J-F	27.54	24-43	0.86	0.13
	J-M	18.65	22-44	0.99	< 0.0001*
	J-J	59.37	20-47	< 0.0001*	0.99
Flock 1	F-F	60	4-63.75	0.04	0.96
	F-M	15.5	5.83-49	0.55	0.45
	F-J	10.3	8.2-35.4	0.91	0.10
	M-F	47.33	5.33-39.33	< 0.0001*	1.00
	M-M	12	6.67-32.11	0.70	0.30
	M-J	8.07	8.6-25.6	0.99	0.01
	J-F	18	12.1-39.3	0.81	0.20
	J-M	15.01	14.67-35.53	0.97	0.03
	J-J	32.96	17.8-31.52	0.01*	0.99
Flock 2	F-F	37.7	13-30	< 0.0001*	1.00
	F-M	33.12	9.25-36.4	0.06	0.94
	F-J	10.375	14-30	0.99	< 0.0001*
	M-F	31.5	6.75-29.5	0.01*	0.99
	M-M	11.75	4.5-38.25	0.70	0.30
	M-J	7	6.75-29.625	0.97	0.03
	J-F	18.65	17.65-38.875	0.96	0.04
	J-M	11	12.375- 46.50	0.99	0.01*
	J-J	46	17.625-38.9375	< 0.0001*	0.99

760
This table shows if specific interaction types occurred above and below null expectations. A
significant p-value are indicated by the * symbol. Mean represents the observed mean approach
rate and the 97.5 column represent the 95.6 is the confidence interval for the permuted mean
approach rate. P-value up above represents mean approach rates that occurred above null
expectations, whereas the p-value down below represent mean approach rates that occurred
below null expectations.
767

768   Table 3 Permutation tests for Interaction Preferences
769

Condition	Comparison	Mean Difference	97.5 Int	p-value
Baseline	FF-FM	23.1	-37.52	0.007*
	MM-MF	-20.14	-24.7-25.03	0.94
Introduction	FF-FM	67.51	-2.37-2.31	< 0.00001*
	FF-FJ	52.35	-2.55-2.43	< 0.00001*
	MM-MF	-29.12	-24.1-25.8	0.99
	MM-MJ	7.09	29.02-27.1	0.34
	JJ-JF	31.83	-19.42-20.44	< 0.00001*
	JJ-JM	40.72	-20.3-21.0	< 0.00001*
Flock 1	FF-FM	44.5	-40-52.4	0.04
	FF-FJ	49.7	-40-52.4	0.03
	MM-MF	-35.33	-28.6-23.6	1.00
	MM-MJ	3.933333	-17.6-22.13	0.36
	JJ-JF	14.46	-19.08-13.6	0.01*
	JJ-JM	17.89333	-16.21-15.4	0.001*
Flock 2	FF-FM	4.5625	-19.7-17.8	0.33
	FF-FJ	27.3125	-15.2-15.2	< 0.00001*
	MM-MF	-19.75	-22.12-27.1	0.95
	MM-MJ	4.75	-22-27.25	0.35
	JJ-JF	35	-19.7-17.8	< 0.00001*
	JJ-JM	27.375	-19.7-17.8	< 0.00001*

770
771
This table shows significant pairwise interaction preferences. The letter F stands for adult
females, M for adult males and J for juveniles. The type of interaction is indicated by two letters
such that the direction of approach is indicated by the first letter ad initiator and the second letter
as the receiver. For example, FF is female-to-female and MJ is male-to-juvenile approaches.
Comparison column represents which differences between mean approach rates are investigated.
For example, FF-FJ represents the mean difference in approach rates from female-to-female
approaches subtracted by female-to-juvenile approach rates. Significant p-values are indicated by
the * symbol. Mean difference represents the observed mean difference.The 97.5 column
represent the 95.6 is the confidence interval for the permuted mean differences.

Changes in social network subgroup structure across the baseline, fusion, and fission-fusion conditions. Shapes represent individuals and arrows represent the approaches between individuals. Communities identified via the spinglass algorithm are outlined in each social network as a cluster of individuals. Rectangles represent juveniles, circles represent adult males and squares represent adult females. Bar graphs represent the mean approach rates of adult females and juveniles in the three conditions. Error bars represent 95% confidence intervals.

808
809
810

811
812
813 Picture of juvenile subgroup. Associations between juveniles often resulted in visually clustered
814 associations within the larger flock.