1   # Juvenile Gouldian Finches (*Erythrura gouldiae*) form

2   # sibling sub-groups during social integration

5   Gregory M. Kohn[1,2], M. Ryan Nugent[2], Xzavier Dail[2],

7   [1]University of North Florida, Department of Psychology

8   [2]Jacksonville Zoo and Gardens, Department of Wildlife Wellness

ABSTRACT

The formation of social relationships in complex groups is critical in shaping patterns of social organization and behavioral development. In many birds, young individuals remain dependent on their parents for extended periods but must abruptly transition to navigating interactions in the wider group after independence. While lack of social relationships during this period is detrimental in the development of later social skills, little is known about the social relationship's juveniles form after independence in many bird species. In this study we describe patterns of social interactions in juvenile Gouldian Finches (*Erythrura gouldiae*) during transitions from family groups into flocks of unfamiliar individuals. A group of 20 juveniles from 4 families were introduced into two flocks. After introductions juveniles showed a gradient of approach rates with most approaches directed towards siblings, followed by juvenile peers, adult females, and lastly adult males. Significant preferences for siblings resulted in the emergence of sibling subgroups within the larger social network. This active self-assortment of siblings suggests that sibling sub-groups are an important bridge linking social connections within the family and the wider group. Such findings suggests that sibling relationships have a critical role in the socialization after independence, as well as structuring the social organization of Gouldian finch flocks.

INTRODUCTION

Social integration involves establishing relationships in new social contexts. Across the vertebrates, from fish, to birds, to mammals, young individuals remain dependent on their parents for extended periods after birth and switch to interacting with unfamiliar conspecifics during adolescence. In many bird species this switch is abrupt, with juveniles suddenly being cut off from parental support (Collias, 1952; Stamps, Kus, Clark, & Arrowood, 1990). This sudden transition to independence is an important developmental period as it's associated with high levels of mortality (Lack, 1954) and interactions during this time are necessary for learning to navigate later social challenges (M. West, King, & Kohn, 2011).

Integrating into a new group carries considerable risks and benefits. Being a new group member is associated with having low social status and higher frequencies of aggression from established group members (Marler, 1976; Senar, Camerino, & Metcalfe, 1990). Such risks may be magnified in juveniles who don't yet possess the necessary skills or experiences to interact with unfamiliar conspecifics. Nonetheless, social integration is a critical component of behavioral development, as juveniles isolated from interactions with others, especially during adolescence, exhibit behavioral and biological deficiencies that impair their ability to thrive in the group (Preiss & Franck, 1974). Therefore, understanding the types of social connections juveniles form as they enter new groups may reveal how they cope with the challenges of social integration.

Many factors influence the formation of juvenile social connections upon entering a group for the first time. Some of these can be passive, caused by constraints in the social or physical environment that force juveniles into close proximity. Passive processes suggest that social connections do not reflect the ability to recognize and prefer specific individuals. For example, in some bird species adults corral their offspring into all-juvenile groups for protection from predation (Munro & Bédard, 1977; Ralf Wanker, Bernate, & Franck, 1996). Such crèches are commonly seen in colonially nesting seabirds and other species, where the risk of predation is high, and juveniles derive substantial protection through the amalgamation of offspring from numerous families. In other cases, juveniles show distinct morphological and physiological differences from adults, which causes them to passively cluster based on shared needs and habitat preferences. For Instance, due to their reduced flying ability fledglings converge on similar spaces as they are unable to synchronize their activities with older individuals. For example, in Piñon Jays (*Gymnorhinus cyanocephalus*), fledglings initially cluster on the ground right after fledging as their lack of flying abilities makes it difficult for them to follow their parents (Balda & Balda, 1978).

Alongside these passive factors, juveniles can form social connections through an active process. In contrast to passive processes active ones involve the recognition and preference for interacting with specific individuals. Upon entering the group, juveniles are exposed to a wide range of potential interaction partners. In groups with a mixed

age and sex composition, the choice between interacting with unfamiliar adults,
juveniles or siblings may be the first active social decision juveniles make. Numerous
studies have shown that birds who inhabit flocks not characterized by strong family ties
still possess the ability to recognize siblings and similar-aged peers (Nakagawa &
Waas, 2004). Studies on both adults and dependent young (fledglings still being cared
for by parents) have shown that birds can differentiate between siblings and unrelated
individuals. Adult zebra finch (*Taeniopygia guttata)* males prefer to perch closer to male
siblings (Burley, Minor, & Strachan, 1990), and juvenile Greylag Geese (*Anser*
anser) can be trained to peck at a specific geometric symbol when in close proximity to
particular siblings (Scheiber, Hohnstein, Kotrschal, & Weiß, 2011).
133
While such studies show that birds are capable of sibling recognition, we know
remarkably little about the social connections juveniles make as they integrate into new
groups. The evidence we have suggests that sibling relationships provide a bridge
between the family group and integration into the wider flock. In Spectacled Parrotlets
(*Forpus conspicillatus*), juveniles are herded into crèches by their parents right after
fledgling. Here they form strong relationships with siblings that are used as a stable
social position to explore interactions with unrelated group memebers (Ralf Wanker et
al., 1996). In house sparrows (*Passer domesticus*), individuals follow siblings over non-
siblings (Tóth et al., 2009), and in budgerigars (*Melosittacus undulatus*) fledglings
attract more interaction from their siblings than expected, and often form an exclusive
bond with one specific sibling (Stamps et al., 1990). Cockatiels (*Nymphicus hollandicus*)
are more likely to share food with siblings than non-siblings (Liévin-Bazin et al., 2019),
and movement and foraging patterns in meadow pipits (*Anthus partensis),* Ospreys
(Pandion haliaetus) and lilac-crowned amazons (*Amazona finschi*), suggest siblings
stay in contact well into independence (Edwards, 1989; Hötker, 1982; Salinas-Melgoza
& Renton, 2007).
150
This study aims to describe the social relationships of newly independent juvenile
Gouldian finches during their first transition to a novel flock. Gouldian finches are an
endangered estrildid finch that inhabits the subtropical grasslands of northern Australia
and exhibit a color polymorphism. Adults show discrete differences in head coloration
with some having either red, black, or yellow heads, while juveniles have grey plumage
with olive backs. Differences in head coloration correspond with differences in behavior
and personality (O'Reilly, Hofmann & Mettke-Hofmann, 2019; Williams, King, & Hoffman
2012; Mettke-Hofmann 2012; Fragueira & Beaulieu, 2019; Pryke, 2007). While little is
known about the social structure of Gouldian finches, large fission-fusion flocks
composed of hundreds of individuals in mixed age and sex composition are commonly
observed in the wild. The fluidity and diversity of estrildid flocks has led some to
speculate that they lack significant patterns of within-group structure (Bolton et al.,
2016; Dostine et al., 2001).
164
165
166

The social relationships individuals form will directly shape the organization and
experience of living in a group (Aureli & Schino, 2019). By describing how juvenile
Gouldian finches integrate into novel flocks, this study aims to uncover significant
aspects of this species developmental ecology. By identifying which social relationships
are preferred and avoided during integration, we can uncover if juveniles are
preferentially exposed to influences from specific group members during this critical
period. In the wild, small aggregations of juvenile Gouldian finches are observed around
waterholes, with adults descending to the waterhole first, followed by groups of juveniles
(O'Reilly, Hofmann, & Mettke-Hofmann, 2019). However, it's unknown if juveniles
maintain connections with peers and siblings outside the waterhole, or if associations
between juveniles at these locations reflect sibling or peer relationships. Due to the
presence of sibling recognition abilities in related species (Nakagawa & Waas, 2004) we
hypothesize that newly independent juvenile Gouldian finches will maintain strong
relationships with siblings upon entering the flock for the first time.

181
182

METHODS

Subjects: All individuals in this study were Gouldian finches. All finches were acquired
from professional breeders in Arizona and Florida, U.S.A. Each bird was provided with a
uniquely colored leg band for individual recognition. Males, females, and juveniles were
distinguished from each other based on plumage and behavioral differences, with adult
females being much more subdued in their chest coloration in contrast to males.
All juveniles were hatched and raised in separate cages prior to being introduced
to experimental flocks. At around 40 days after fledging, juveniles become independent
of their parents and can feed and drink on their own without any sign of parental
provisioning. Juveniles were distinguished from adults based on their grey heads, olive
backs, and lack of typical adult plumage. In this study, all juveniles were under one year
of age since fledging the nest and adults were older than a year and exhibited adult
plumage. Head color was easily observed in adults and assessed in juveniles at the
time of their first molt after the end of the data collection period. We used a total of 11
adults and 20 juveniles for this study.
All birds were provided with a daily diet consisting of finch seed mix (Kaytee),
with soaked Perle Morbide pellets, eggshell, boiled eggs, and dandelion greens
provided twice a week. Fresh water was continually available.

Aviaries: We used a series of two aviaries in this study. Each aviary consisted of
identical dimensions (23 x 72 x 41 in) and similar environmental conditions. The aviaries
were located indoors next to a large window allowing for natural sunlight and ventilation.
Each aviary contained both natural eucalyptus and plastic privacy perches, small
shrubs, a feeding station, and a bowl for water baths.

Data Collection: We used focal sampling during data collection. During observation
periods, all individuals in the flock were observed for a 5-minute focal block. We

recorded all approaches that occurred within a 2-inch radius around the focal individual being observed. All observations were done using voice recognition technology allowing the observer to maintain continual visual contact with the focal bird throughout the sampling block. An approach was scored when another individual approached another within a radius of 2 inches. Before data collection, video of marked perches was made to establish that 2 inches was the typical social distance observed between two resting adult finches. All focal samples were collected by GMK.

Procedure

First introduction: On 10/5/2020, four novel juvenile siblings were introduced into an aviary consisting of 6 adult female and 5 adult males. From 10/06/2020 to 12/22/2020, 503 focal samples were taken. All individuals observed for 34 blocks for a total of 170 minutes except one individual who was removed from the flock temporarily for health reasons and was observed for 26 blocks.

Second Introduction: On 2/10/2021, the aviary from the first introduction was split up into two separate flocks: three adult males, two females, and two juveniles were moved into Aviary 1, while four females, two males, and one juvenile were moved to Aviary 2. In each flock, three new juvenile siblings were randomly introduced. From 2/11/2021 to 4/20/2021, a total of 84 blocks were recorded in each aviary. Everyone was observed for a total of 8 blocks measuring 40 minutes each.

Third Introduction: On 05/25/2021, a group of 5 juveniles were transferred from their family group and introduced into Aviary 1. On 06/11/2021, another group of 5 siblings were transferred to Aviary 2. From 05/26/2021 to 06/27/2021, all individuals in the flock were focal sampled, collecting a total of 240 blocks in each aviary. Each individual was observed for 16 blocks for a total of 80 minutes each.

ANALYSIS

We used generalized linear mixed models to investigate the social preferences of juveniles when integrating into the flock. Three models were constructed for this study: the dyad-model, age-model, and sibling-model. The dyad-model was a multi-membership model performed using the MCMCglmm package and was used to predict the dyad weights (how often two individuals in the flock interacted) for all dyads involving juveniles. The MCMCglmm package is a Bayesian mixed model package that allows for multi-membership and random effects (Hadfield, 2010), and has been used to investigate the factors shaping social networks and relationships in animals (Hart, Weiss, Brent, & Franks, 2021; Raulo et al., 2021). Trace plots were examined to assess if the dyad-model was well-mixed, and Heidelberger and Welch's diagnostics were used to test if the Markov chain was from a stationary distribution. The dyad weights represented the number of times each dyad in the flock approached each other. For each dyad we classified if it was between members of the same sex, age, family, or

255 head color as a fixed effect. The random effects included the effect of round, aviary, and
256 a multi-membership term that contained the initiator and receiver of each dyad.
257
258   Post-hoc tests included a permutation test where raw approach data was
259 permuted 100,000 times to create a null distribution of approaches for each aviary. The
260 observed approach rates were then compared to the permuted approach rates to see if
261 they occurred above or below null expectations with empirical p-values calculated at an
262 alpha of 0.025 for each tail of the distribution.
263
The second model, the age-model investigated if juveniles showed preferences
towards adults or juveniles in the aviaries. The third model, the "sibling-model",
investigated if juveniles showed preferences towards siblings and unrelated individuals.
Both models were generalized linear mixed models conducted using the lmerTest
package (Kuznetsova, Brockhoff, & Christensen, 2017). We first calculated the rate of
approaches towards juvenile, adult male, adult female, sibling and non-sibling
conspecifics. This was done by dividing the sum of the approaches toward members of
a specific category, such as juveniles, by the number of individuals in that category in
the flock. Both models included the approach rate as the dependent factor. The age-
model included approach rates towards adult males, adult females, and juveniles, while
the sibling-model included the approach rates for siblings and unrelated flock members.
Each model included the direction of approach's (age-model: adult male, adult females,
or juveniles; sibling-model: siblings or non-siblings), the individual's sex, and head color
as fixed effects with random effects of individual identity, round, and aviary. Post hoc
tests used Wilcoxon tests to run pairwise comparisons between interaction preferences
while controlling for multiple comparisons using the False Discovery Index.
280
281 **RESULTS**
282
During the first introduction a group of four juveniles was added to the aviary and
a total number of 8,840 approaches were recorded. In the second introduction the flock
was split up into two separate aviaries and three juveniles added to each aviary. Here a
total of 2,184 approaches were recorded in Aviary 1 and 2,345 were recorded in Aviary
2. During the third introduction where 5 juveniles were introduced to each aviary a total
of 4,425 approaches were recorded in Aviary 1 while 3,990 approaches were recorded
in Aviary 2.
290
The dyad-model showed a significant influence of family, age, and head color on
the frequency of dyads. The strongest dyads involving juveniles were more likely to
occur between members of the same family and between individuals of the same age.
Our model also uncovered a significant effect of head color with the coefficient
indicating stronger dyads for individuals of differing head colors (Table 1). Post-hoc
permutation analysis showed that, across all the flocks, the juvenile-to-juvenile dyads
occurred at a higher rate than expected under null conditions, while in many

introductions the juvenile-to-male dyads occurred at lower-than-expected rates (Table
2).
300
Both the age and sibling model showed a significant main effect of approach
direction. The age-model showed a significant main effect of approach direction toward
juveniles or adults (Table 3) and the sibling-model showed a significant main effect of
approach direction towards siblings and non-siblings (Table 3). Post hoc tests indicated
juveniles had higher approach rates towards other juveniles than towards adult males
and females (N = 20, Mean juvenile approach rate = 40.65, Mean female approach rate
= 18.01, Mean male approach rate = 11.83, Wilcoxon Rank Sum test, juvenile-adult
female, V = 0, $p < 0.0001$, Wilcoxon Rank Sum test, juvenile-adult male, V = 0, $p <$
0.0001, Fig 1). When interacting with adults, juveniles showed higher rates of
approaches towards females in contrast to adult males (N = 20, Wilcoxon Rank Sum
test, adult male-adult female, V = 26, $p = 0.005$). Approach rates towards siblings where
higher than approach rates to unrelated conspecifics (N = 20, Mean sibling approach
rate = 23.6, Mean non-sibling approach rate = 14.15, Wilcoxon Rank Sum sibling-non-
sibling, V = 163, $p = 0.03172$, Fig 1).
315
To see if juveniles preferred related siblings over unrelated juveniles, we did an
additional analysis restricted to our third introduction. During the third introduction, the
flock size was large enough to calculate approach rates towards both siblings and
unrelated juveniles for the 10 juveniles introduced during this period. An additional
GLMM was performed using the lmerTest package that looked at the fixed effect of
direction (sibling versus non-sibling juveniles), sex, and head color with random effects
of aviary and individual. Our model showed a significant influence of direction (Table 3)
with the juveniles preferring to interact with siblings over unrelated peers (N = 10, Mean
Sibling = 39.02, Mean unrelated juvenile = 16.7, , Wilcoxon Rank Sum test, siblings-
unrelated juveniles, V = 55, $p = 0.002$, Fig 1 C). Permutating approaches during the
third period indicated that while sibling directed approaches occurred above null
expectations, approaches to unrelated juveniles occurred below null expectations
(Table 2).
329
330
331 **DISCUSSION**
332
This study described the social preferences and relationships of newly independent
Gouldian finches during their transition from the family group to the larger flock. The
period between fledging and adulthood is one of the most critical stages for survival and
development in birds. Our results show that juvenile Gouldian finches preferentially
choose to interact with siblings after becoming independent of their parents. This is
among the few examples of sibling-assortment of independent juveniles in passerine
flocks (Hötker, 1982; Kohn, King, Scherschel, & West, 2011; Nicolai, 1956; Templeton,
Reed, Campbell, & Beecher, 2012; Tóth et al., 2009). Upon entering the group for the
first time, juveniles showed a gradient of approach rates with most approaches directed

towards siblings, followed by juvenile peers, adult females, and lastly towards adult
males (Fig 2). Juveniles showed significantly higher approach rates towards siblings
over both unrelated juveniles and adults. This resulted in the emergence of sub-groups
that remained consistent across changes in the age-sex composition of the flock.
346
A diverse range of factors can cause the formation of sibling and peer groups in juvenile
birds. In zebra finches, fledglings initially follow their parents around the colony but
quickly cluster around areas with same aged-peers (Goodwin, 1982; Ogino, Maldonado-
Chaparro, & Farine, 2021; Zann, 1996). In these contexts, it is unclear if the close
association between juveniles is caused by preferences to interact with other juveniles,
collective herding of juveniles by parents, or a juvenile's inability to synchronize their
behavior with parents during foraging trips. Therefore, understanding the role that the
environment, experiences, morphology, and active social preferences play in the
formation of juvenile subgroups is essential to understanding their function in the
evolution and development of avian behavior.
357
Our results suggest that the formation of juvenile subgroups in Gouldian flocks is an
active process, reflecting the ability of juveniles to recognize their siblings and mutually
prefer interacting with them. Studies in birds have shown that individuals can recognize
and preferentially attend to siblings (Nakagawa & Waas, 2004). In a cross-fostering
study with juvenile Bengalese finches (*Lonchura striata domestica*), preening bouts
between genetic siblings were longer and more frequent that preening bouts between
foster siblings (Ju & Lee, 2016). In house sparrows (*Passer domesticus*), juveniles
preferred to follow siblings (Tóth et al., 2009) over unrelated conspecifics. Our analysis
showed that sibling-to-sibling approaches were significantly higher than approaches to
all others, and that juvenile subgroups were more likely to contain siblings than
unrelated juveniles (Fig 2). In captive contexts, competition over patchy resources is
minimized and observers did not find that either juvenile or adults consistently occupied
different areas of the enclosure (Kohn, personal observations). Outside of plumage
differences, juvenile Gouldian finches do not show substantial morphological or
physiological differences from adults (Gelis, 2003). This suggests that the higher
approach rates towards siblings observed in juvenile Gouldian finches reflects decisions
made in accordance with underlying social preferences.
375
In the wild, the period from fledging to independence is characterized by high mortality
and unique developmental challenges (Cox, Thompson III, Cox, & Faaborg, 2014).
Specifically, one of the largest challenges juveniles face is integrating into novel groups
for the first time with little experiences outside the family (Collias, 1952). Our study
highlights that siblings play an important role in social integration as they are the
strongest relationships juveniles have after becoming independent of their parents. The
existence of sibling sub-groups assures some social connections will remain stable as
juveniles move from the family to the larger flock. This provides a predictable social
context that can buffer against aggressive interactions with novel individuals.
Interactions with novel conspecifics carries increased risk of aggression since adults are

often more aggressive toward unrelated juveniles in birds (Senar et al., 1990). Juveniles
may respond to adult aggression by quickly becoming selective in their interactions and
preferentially associating with familiar siblings. Sub-groups of familiar siblings will soften
the impact of stressful interactions with novel individuals while assuring individuals get
the predictable social experiences needed for development.
391
Interactions with siblings can have significant benefits, such as access to beneficial
social information or experiences necessary for the development of competent
behavioral abilities (Alberts, 2008). Among mammals and birds, early interactions are
critical for the development of species-typical abilities (King & West, 2002; M. J. West &
King, 1987). When isolated from peers, juveniles often develop distinct behavioral
repertoires or long-lasting behavioral deficiencies when compared to typically reared
individuals (Suomi & Harlow, 1972;). For instance, Rosy-faced lovebird (*Agapornis
roseicollis*) chicks raised in isolation were more aggressive, showed increased
avoidance behavior, were unable to form a pair bond, or become integrated into the
flock (Preiss & Franck, 1974). In mallard ducklings (*Anas platyrhynchos*) chicks require
direct interactions with siblings in semi-naturalistic environments to properly imprint on
maternal cues (R. Lickliter & Gottlieb, 1985). Ducklings will even switch their imprinting
preferences to be in line with their siblings if they show a preference for different objects
(R. Lickliter & Gottlieb, 1986). In cross fostering studies on zebra finches, it was shown
that the ability to imprint on foster Bengalese finch parents is shaped by the presence of
siblings. Zebra finches raised by Bengalese finch parents, but with 4 to 5 zebra finch
siblings showed species-typical sexual preferences whereas those with fewer siblings
developed sexual preferences for Bengalese finches (Kruijt, Ten Cate, & Meeuwissen,
1983). While such results show that siblings are necessary resources in development of
species-typical abilities, less is known about the ontogenetic role that siblings play after
independence.
413
Upon independence, behaviors acquired through sibling and peer interactions can
facilitate integration into the group (Pellis & Pellis, 2007). Interactions with siblings
allows individuals to explore their behavioral potentials and refine their behavioral
abilities without the risks associated with adult interactions. In Spectacled parrotlets
(*Forpus conspicillatus*), interactions within siblings serve as a springboard to the
formation of their first pair bonds (Garnetzke-Stollmann & Franck, 1991; Ralf Wanker et
al., 1996), and singletons who hatch without siblings compensate by prolonging their
relationship with parents and forming wider social networks at independence (R.
Wanker, 1999). In Brown-headed cowbirds (*Molothrus ater*), interactions in juvenile
subgroups predict the development of competent courtship behavior during an
individual's first breeding season (Kohn, King, Dohme, Meredith, & West, 2013).
Rhesus macaque (*Macaca mulatta)* infants raised in isolation were able to acquire
species-normative social behavior after an extensive period of social interaction with
peers (Suomi & Harlow, 1972). Some studies have even shown that juveniles cluster to
form "gangs" around contested resources to better compete with more experienced
adults (Dall & Wright, 2009; Franks et al. 2020). These studies suggest that juvenile and

sibling subgroups may facilitate the horizontal co-construction of social skills necessary
to navigate interactions during later periods (Kohn, 2019). Current studies are looking at
the details of juvenile-to-juvenile interactions in Gouldian finches to explore how they
might guide the development of competence social and reproductive skills in adulthood.
434
When interacting with adults, juvenile Gouldian finches showed a significant preference
to interact with females over males. In many species, females are less aggressive
towards juveniles in contrast to males (Ward & Webster, 2016). When exploring
connections outside sibling subgroups, juveniles experience less displacement and
aggression from females in contrast to males (Kohn, in prep). Across the different
flocks, juvenile approaches towards males were more likely to occur below null
expectations, suggesting that females may be the first unrelated adults juveniles have
consistent contact with. While little is currently known about the shifting patterns of
social relationships across development in birds this study suggests that adult females
may be an important factor in social integration.
445
The stability of a group's social organization reflects the ability of juveniles to
reconstruct patterns of relationships across generations (Deputte, 2000; Kohn, 2019).
Consequently, the social relationships made by juvenile finches as they transition into
the flock for the first time will influence how they experience their social environment
and the social organization of the group itself. By constructing sibling sub-groups,
juvenile Gouldian finches will disproportionately have access to information from
siblings. This study and others suggest that sibling relationships are likely an important
component in the social development of passerines. As passerines are heavily
dependent on socially transmitted information, the emergence of a stable network of
siblings may create the predictable social contexts necessary for the development of
competent social skills.
457
458 **ACKNOWLEDGMENTS**
459

460 This study was supported with a grant from the Van Vleck Award. All research was
461 done in accordance with ASAB/ABS Guidelines for the Use of Animals in Research and
462 was approved by the University of North Florida's Institutional Animal Care and Use
463 Committee protocol number 20-006.

464
465   REFERENCE
466

Alberts, J. R. (2008). The Nature of Nurturant Niches in Ontogeny. *Philosophical*

Psychology, *21*(3), 295–303. https://doi.org/10.1080/09515080802169814

Aureli, F., & Schino, G. (2019). Social complexity from within: How individuals

experience the structure and organization of their groups. *Behavioral Ecology*

and Sociobiology, *73*(1), 6. https://doi.org/10.1007/s00265-018-2604-5

Balda, R. P., & Balda, J. H. (1978). The care of young Piñon Jays (Gymnorhinus

cyanocephalus) and their integration into the flock. *Journal Für Ornithologie*,

119(2), 146–171. https://doi.org/10.1007/BF01644586

Burley, N., Minor, C., & Strachan, C. (1990). Social preference of zebra finches for

siblings, cousins and non-kin. *Animal Behaviour*, *39*(4), 775–784.

https://doi.org/10.1016/S0003-3472(05)80389-6

Collias, N. E. (1952). The development of social behavior in birds. *The Auk*, *69*, 127–

159. https://doi.org/10.2307/4081265

Cox, W. A., Thompson III, F. R., Cox, A. S., & Faaborg, J. (2014). Post-fledging survival

in passerine birds and the value of post-fledging studies to conservation. *The*

Journal of Wildlife Management, *78*(2), 183–193.

https://doi.org/10.1002/jwmg.670

Deputte, B. L. (2000). Primate Socialization Revisited: Theoretical and Practical Issues

in Social Ontogeny. In P. J. B. Slater, J. S. Rosenblatt, C. T. Snowdon, & T. J.

Roper (Eds.), *Advances in the Study of Behavior* (Vol. 29, pp. 99–157).

Academic Press. https://doi.org/10.1016/S0065-3454(08)60104-7

Edwards, T. C. (1989). The Ontogeny of Diet Selection in Fledgling Ospreys. *Ecology*,

70(4), 881–896. https://doi.org/10.2307/1941356

Garnetzke-Stollmann, K., & Franck, D. (1991). Socialisation Tactics of the Spectacled

Parrotlet (Forpus conspicillatus). *Behaviour*, *119*(1/2), 1–29.

Gelis, S. (2003). The gouldian finch (erythrura gouldiae) in health and disease.

Seminars in Avian and Exotic Pet Medicine, *12*(4), 215–227.

https://doi.org/10.1053/S1055-937X(03)00037-9

Goodwin, D. (1982). *Estrildid Finches of the World*. British Museum (Natural History).

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. *Journal of Statistical Software*, *33*(2).

https://doi.org/10.18637/jss.v033.i02

Hart, J. D. A., Weiss, M. N., Brent, L. J. N., & Franks, D. W. (2021). *Common*

Permutation Methods in Animal Social Network Analysis Do Not Control for Non-

independence [Preprint]. Animal Behavior and Cognition.

https://doi.org/10.1101/2021.06.04.447124

Hötker, H. (1982). Studies of the behaviour of young meadow pipits (Anthus pratensis)

after fledging. *Vogelwelt*, (103), 1–16.

Ju, S., & Lee, S. (2016). Effect of kinship on the allopreening among juvenile Bengalese

finches. *Animal Cells and Systems*, *20*(4), 213–217.

https://doi.org/10.1080/19768354.2016.1194318

King, A. P., & West, M. J. (2002). The Ontogeny of Competence. In *Conceptions of*

Development. Psychology Press.

Kohn, G. M. (2019). How social systems persist: Learning to build a social network in an

uncertain world. *Animal Behaviour*, *154*, 1–6.

https://doi.org/10.1016/j.anbehav.2019.06.006

Kohn, G. M., King, A. P., Dohme, R., Meredith, G. R., & West, M. J. (2013). Robust

autumn social attributes predict spring courtship skills in juvenile female brown-

headed cowbirds, Molothrus ater. *Animal Behaviour*, *85*(4), 727–732.

https://doi.org/10.1016/j.anbehav.2013.01.009

Kohn, G. M., King, A. P., Scherschel, L. L., & West, M. J. (2011). Social niches and sex

assortment: Uncovering the developmental ecology of brown-headed cowbirds,

Molothrus ater. *Animal Behaviour*, *82*(5), 1015–1022.

https://doi.org/10.1016/j.anbehav.2011.07.035

Kruijt, J. P., Ten Cate, C. J., & Meeuwissen, G. B. (1983). The influence of siblings on

the development of sexual preferences of male zebra finches. *Developmental*

Psychobiology, *16*(3), 233–239. https://doi.org/10.1002/dev.420160309

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package:

Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Lack, D. (1954). The Natural Regulation of Animal Numbers. *The Natural Regulation of*

Animal Numbers. Retrieved from

https://www.cabdirect.org/cabdirect/abstract/19552902187

Lickliter, R., & Gottlieb, G. (1985). Social interaction with siblings is necessary for visual

imprinting of species-specific maternal preferences in ducklings (Anas

platyrhynchos). *Journal of Comparative Psychology*, *99*(4), 371–379.

https://doi.org/10.1037/0735-7036.99.4.371

Lickliter, R., & Gottlieb, G. (1986). Visually imprinted maternal preference in ducklings is

redirected by social interaction with siblings. *Developmental Psychobiology*,

19(3), 265–277. https://doi.org/10.1002/dev.420190312

Liévin-Bazin, A., Pineaux, M., Covec, M. L., Gahr, M., Bovet, D., & Bayern, A. M. P.

von. (2019). Food sharing and affiliation: An experimental and longitudinal study

in cockatiels (Nymphicus hollandicus). *Ethology*, *125*(5), 276–288.

https://doi.org/10.1111/eth.12850

Marler, P. (1976). On animal aggression: The roles of strangeness and familiarity.

American Psychologist, *31*(3), 239–246. (1977-00540-001).

https://doi.org/10.1037/0003-066X.31.3.239

Munro, J., & Bédard, J. (1977). Crèche Formation in the Common Eider. *The Auk*,

94(4), 759–771. https://doi.org/10.2307/4085272

Nakagawa, S., & Waas, J. R. (2004). O sibling, where art thou? A review of avian

sibling recognition with respect to the mammalian literature. *Biological Reviews*,

79(1), 101–119. https://doi.org/10.1017/S1464793103006249

Nicolai, J. (1956). Zur Biologie und Ethologie des Gimpels (Pyrrhula pyrrhula L.) D. 77.

Zeitschrift Für Tierpsychologie, *13*(1), 93–132. https://doi.org/10.1111/j.1439-

0310.1956.tb01549.x

Ogino, M., Maldonado-Chaparro, A. A., & Farine, D. R. (2021). Drivers of alloparental

provisioning of fledglings in a colonially breeding bird. *Behavioral Ecology*, *32*(2),

316–326. https://doi.org/10.1093/beheco/araa137

O'Reilly, A. O., Hofmann, G., & Mettke-Hofmann, C. (2019). Gouldian finches are

followers with black-headed females taking the lead. *PLOS ONE*, *14*(4),

e0214531. https://doi.org/10.1371/journal.pone.0214531

Pellis, S. M., & Pellis, V. C. (2007). Rough-and-Tumble Play and the Development of

the Social Brain. *Current Directions in Psychological Science*, *16*(2), 95–98.

https://doi.org/10.1111/j.1467-8721.2007.00483.x

Preiss, H.-J., & Franck, D. (1974). Verhaltensentwicklung isoliert handaufgezogener

Rosenköpfchen (Agapornis roseicollis [Vieillot]). *Zeitschrift für Tierpsychologie*,

34(5), 459–463. https://doi.org/10.1111/j.1439-0310.1974.tb01815.x

Raulo, A., Allen, B. E., Troitsky, T., Husby, A., Firth, J. A., Coulson, T., & Knowles, S. C.

L. (2021). Social networks strongly predict the gut microbiota of wild mice. *The*

ISME Journal, *15*(9), 2601–2613. https://doi.org/10.1038/s41396-021-00949-3

Salinas-Melgoza, A., & Renton, K. (2007). Postfledging Survival and Development of

Juvenile Lilac-Crowned Parrots. *Journal of Wildlife Management*, *71*(1), 43–50.

https://doi.org/10.2193/2005-646

Scheiber, I. B. R., Hohnstein, A., Kotrschal, K., & Weiß, B. M. (2011). Juvenile Greylag

Geese (Anser anser) Discriminate between Individual Siblings. *PLOS ONE*, *6*(8),

e22853. https://doi.org/10.1371/journal.pone.0022853

Senar, J. C., Camerino, M., & Metcalfe, N. B. (1990). Familiarity Breeds Tolerance: The

Development of Social Stability in Flocking Siskins (Carduelis spinus). *Ethology*,

85(1), 13–24. https://doi.org/10.1111/j.1439-0310.1990.tb00381.x

Stamps, J., Kus, B., Clark, A., & Arrowood, P. (1990). Social relationships of fledgling

budgerigars, Melopsitticus undulatus. *Animal Behaviour*, *40*(4), 688–700.

https://doi.org/10.1016/S0003-3472(05)80698-0

Suomi, S. J., & Harlow, H. F. (1972). Social rehabilitation of isolate-reared monkeys.

Developmental Psychology, *6*(3), 487–496. (1972-26700-001).

https://doi.org/10.1037/h0032545

Templeton, C. N., Reed, V. A., Campbell, S. E., & Beecher, M. D. (2012). Spatial

movements and social networks in juvenile male song sparrows. *Behavioral*

Ecology, *23*(1), 141–152. https://doi.org/10.1093/beheco/arr167

Tóth, Z., Bokony, V., Lendvai, A., Szabó, K., Pénzes, Z., & Liker, A. (2009). Whom do

the sparrows follow? The effect of kinship on social preference in House Sparrow

flocks. *Behavioural Processes*, *82*, 173–177.

https://doi.org/10.1016/j.beproc.2009.06.003

Wanker, R. (1999). Socialization in spectacled parrotlets (Forpus conspicillatus): How

juveniles compensate for the lack of siblings. *Acta Ethologica*, *2*(1), 23–28.

https://doi.org/10.1007/PL00012228

Wanker, Ralf, Bernate, L. C., & Franck, D. (1996). Socialization of Spectacled

ParrotletsForpus conspicillatus: The role of parents, crèches and sibling groups

594   in nature. *Journal Für Ornithologie, 137*(4), 447–461.

595   https://doi.org/10.1007/BF01661101

596 Ward, A., & Webster, M. (2016). *Sociality: The Behaviour of Group-Living Animals.*

597   Springer International Publishing. https://doi.org/10.1007/978-3-319-28585-6

598 West, M. J., & King, A. P. (1987). Settling nature and nurture into an ontogenetic niche.

599   *Developmental Psychobiology, 20*(5), 549–562.

600   https://doi.org/10.1002/dev.420200508

601 West, M., King, A., & Kohn, G. (2011). Developmental ecology: Platform for designing a

602   communication system. *Interaction Studies, 12*(2), 351–371.

603 Zann, R. (1996). *The Zebra Finch: A Synthesis of Field and Laboratory Studies.*

604   https://doi.org/10.5860/choice.34-5095

641 Table 1: Results from the Dyad Model
642

Fixed Effects	Posterior means	Lower 95% CI	Upper 95% CI	Bayesian P value
Sex	2.76	-0.8377	6.3081	0.64
Family	30.39	25.5280	34.9899	<0.001
Age	10.88	5.8812	16.1538	<0.001
Head Color	-7.17	-11.4511	-3.1861	<0.001

643
644 Results from the Monte-Carlo Markov Chain Generalized Linear mixed model. The
645 effect of same sex, age, and family was included in the model as fixed effect. The

Table 2: Permutation Tests for Approaches by Age and Sex

Condition	Interaction	Mean	95% CI	p-value Up	p-value Down
First	J-F	27	24-43	0.86	0.13
	J-M	18	22-44	0.99	< 0.0001*
	J-J	59	20-47	< 0.0001*	0.99
Second - Aviary 1	J-F	18	12-39	0.81	0.20
	J-M	15	15-35	0.97	0.03
	J-J	33	17-31	0.001*	0.99
Second - Flock 2	J-F	18	18-38	0.96	0.04
	J-M	11	12-46	0.99	0.01*
	J-J	46	17-39	< 0.0001*	0.99
Third-Flock 1	J-F	13.3	13-31	0.95	0.05
	J-M	10.5	15 – 29	0.99	< 0.0001*
	J-J	27.0	19 -24	< 0.0001*	0.99
	J-Sib	49.24	16 - 32	< 0.0001*	0.99
	J-non-Sib J	12	16 - 32	0.99	< 0.0001*
Third-Flock 2	J-F	10.14	14 - 22	0.99	< 0.0001*
	J-M	10.0	11 - 24	0.99	0.003
	J-J	23	15 - 20	< 0.0001*	0.99
	J-Sib	29	13 - 23	< 0.0001*	0.99
	J-non-Sib J	12	13 - 23	0.94	0.06

Table 3: Results from the age, sibling, and sibling-juvenile model

Model	Fixed-Effect	Coefficient	Std. Error	t value	P value
Age	Direction	22.556	2.559	8.813	<0.001*
	Sex	-5.817	4.862	-1.196	0.2455
	Head Color	2.641	2.314	1.141	0.2673
Sibling	Direction	9.4500	2.8992	3.260	0.0023*
	Sex	6.1024	3.7654	1.621	0.12
	Head Color	2.3855	5.2048	0.458	0.68
Sibling-Juvenile	Direction	6.3000	1.9893	1.146	0.005*
	Sex	2.7647	2.4123	1.146	0.27
	Head Color	7.0147	4.0366	1.543	0.35

731   Figure 1: Social Networks and Social Preferences

732
733
Fig 1. A) Social Networks formed during the different introductions. Shapes
represent individuals, with triangles as juveniles, circles as adult males and
squares as adult females. Arrows represent approaches between individuals.
The spinglass communities are shown in the grouping of different shapes. B) Bar

graph of the median approaches directed towards juveniles, adult females and
adult males. The error bars represent 95% confidence intervals in all graphs. C)
Bar graphs represent median approaches directed towards siblings and non-
siblings. D) Bar graphs represent median approaches siblings and non-sibling
juveniles during the third introduction.
752   Figure 2: Picture of Juvenile Subgroup
753
754

755
756
757 Picture of juvenile subgroup. Associations between juveniles often resulted in visually
758 clustered associations within the larger flock.
759
760