Rescue of stalled clathrin-mediated endocytosis by asymmetric Arp2/3-mediated actin

assembly

Meiyan Jin[1*], Cyna Shirazinejad[1, 2*], Bowen Wang[3], Amy Yan[1], Johannes Schöneberg[1, 4],

Srigokul Upadhyayula[1], Ke Xu[3], David G. Drubin[1#]

[1] Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720

[2] Biophysics Graduate Group, University of California Berkeley; Berkeley, CA, 94720

[3] Department of Chemistry, University of California, Berkeley, CA 94720

[4] Current address: Department of Pharmacology, and Department of Chemistry and

Biochemistry, University of California, San Diego, CA 92093

[*]These authors contributed equally.

[#]Correspondence to DGD (drubin@berkeley.edu)

Author contributions

MJ, CS and DGD conceived the study and experiments. MJ and AY generated the genome-

edited cell lines. MJ performed live cell data acquisition and sample preparation for super-

resolution microscopy. BW and KX performed super-resolution microscopy and super-resolution

data reconstruction. CS developed computational analysis tools and SU, JS, and MJ supported

the data analysis. MJ and CS prepared the figures and MJ and DGD wrote the manuscript with

feedback from the other authors.

Abstract

Actin assembly facilitates vesicle formation in several trafficking pathways. Clathrin-mediated endocytosis (CME) shows elevated actin assembly dependence under high membrane tension. Why actin assembly at CME sites occurs heterogeneously even within the same cell, and how assembly forces are harnessed, are not fully understood. Here, endocytic dynamics, actin presence, and geometry of CME proteins from three different functional modules, were analyzed using three-dimensional (3D) super-resolution microscopy, live-cell imaging, and machine-learning-based computation. When hundreds of CME events were compared, sites with actin assembly showed a distinct signature, a delay between completion of coat expansion and vesicle scission, indicating that actin assembly occurs preferentially at stalled CME sites. N-WASP is recruited to one side of CME sites where it is positioned to stimulate asymmetric actin assembly. We propose that asymmetric actin assembly rescues stalled CME sites by pulling vesicles into the cell much like a bottle opener pulls off a bottle cap.

Introduction

Formation of clathrin-coated vesicles requires forces to first bend the membrane into a sphere or tube, and to then break the thin neck that connects the vesicle to the plasma membrane. These forces are generated through the combined actions of actin filament assembly and proteins that directly bend the membrane[1–5] (Supplementary Fig. 1a). Several studies have demonstrated that dependence of CME on actin assembly increases under elevated membrane tension[6–9]. Interestingly, actin does not assemble at all CME sites in mammalian cells, suggesting highly localized differences in requirement for actin assembly, that nature of which are obscure[10–12]. A detailed understanding of how actin forces are harnessed to aid vesicle formation and scission,

and whether and how actin assembly might mediate an adaptive response to the opposing forces

such as membrane tension and turgor pressure, depends on understanding which CME sites

assembly actin, where filament assembly occurs around the endocytic membrane and when. In

yeast cells, where turgor pressure is particularly high, super-resolution data suggest that actin

assembles symmetrically around CME sites and indicate that actin regulators including Las17,

which is yeast WASP, are present in a ring surrounding the base of the clathrin coat

symmetrically[13]. On the other hand, studies on fixed mammalian cells raised the possibility that

actin assembly may at least in some cases be initiated asymmetrically at clathrin coats[14,15].

However, methods used for these studies prevented analysis of large numbers of sites, and

suffered from possible loss of actin filaments during unroofing and extraction of the cells. Which

CME sites assemble actin, and how actin networks are organized with respect to CME sites, has

not been determined systematically, in a large-scale, unbiased manner, particularly in live

mammalian cells. This information is essential to understanding how and why actin assembly

forces are harnessed for CME.

Here, by combining fixed and live-cell imaging of triple-genome-edited, human induced

pluripotent stem cells (iPSCs), and newly developed machine-learning-based computational

analysis tools, we report that N-WASP and Arp2/3 complex localize at one side of the coat and

neck of invaginating endocytic sites until the scission, similar to what was proposed

previously[14,15]. Most importantly, by comparing recruitment dynamics of proteins from three

distinct endocytic modules for over one thousand endocytic events, we found that branched actin

assembly occurs predominantly at sites that have stalled between coat expansion and scission.

We propose that these branched actin networks rescue stalled CME.

Results

Super-resolution imaging reveals asymmetric actin distribution around endocytic sites

To investigate the physiological roles and spatiotemporal regulation of actin assembly at CME sites in mammalian cells, we applied genome-editing techniques to generate a human iPSC line (hereafter referred to as ADA cells) that co-expresses a TagRFP-T fusion of the mu1 subunit of the AP2 adaptor complex (AP2M1), a TagGFP2 fusion of dynamin2 (DNM2), and a HaloTag fusion of the ARPC3 subunit of the Arp2/3 complex as representatives of the CME coat, scission and actin modules respectively[2,16,17] (Supplementary Fig. 1 and Supplementary Video 1, 2). Previous studies showed that endogenously tagged AP2M1, DNM2 and ARPC3 can serve as reliable markers of these CME functional modules that avoid disruption of physiological spatiotemporal organization of the process as might be caused by overexpression of fluorescently labeled proteins[12,18–21]. We observed dynamic CME events on the basal plasma membrane of the genome-edited cells using Total Internal Reflection Fluorescence (TIRF) microscopy (Supplementary Fig. 1c). Consistent with previous studies, AP2 is recruited at early CME stages while DNM2 is recruited in two phases[10,16,20,22]. At the early stage of CME, a relatively small amount of DNM2 is recruited to CME sites. Shortly before the end of a CME event, the DNM2 recruitment rate increases rapidly with DNM2 levels reaching a peak concomitant with vesicle scission[16,20,23] (Supplementary Fig. 1c). This later rapid-recruitment phase represents the assembly of the dynamin helix on the highly curved neck of the budding vesicle after the U to Ω shape transition of the endocytic membrane[20,23–27].

Super-resolution imaging of fixed human skin melanoma SKMEL cells observed asymmetry of actin arrangement around CME sites[7], consistent with observations from previous studies[14,15]. To analyze how actin networks are organized at CME sites in iPS cells, we first performed two-

color 3D Stochastic Optical Reconstruction Microscopy (STORM) imaging[28] on fixed ADA

cells, localizing either AF647 phalloidin-labeled actin filaments[29] or HaloTag-fused ARPC3 at

CME sites. Due to the dense phalloidin labelling of cortical actin filaments under the plasma

membrane, it was often challenging to unambiguously identify the CME-specific actin structures

in iPSCs. However, in regions with thinner cortical actin layers, we observed that actin was

typically distributed asymmetrically around CME sites (Fig. 1a, b), consistent with what has

been observed in different mammalian cell lines by STORM or EM imaging approaches[7,15].

Antibody labeling of ARPC3-Halotag in the ADA cells had the advantage of a less complex

staining pattern. Besides being highly concentrated in lamellipodia, ARPC3 was associated with

CME sites asymmetrically, like actin (Fig 1c, d). These data suggest an asymmetric Arp2/3-

mediated actin network arrangement around CME sites.

Asymmetric branched actin networks assembled at CME sites persist through scission

We next used ADA cells to investigate actin assembly at CME sites in live cells, which has

several advantages over studies in fixed cells. During the fixation and subsequent sample

preparation, actin structures may not be faithfully preserved. In addition, in fixed cells it is very

difficult to identify the stage of the CME, so the timing, geometry and dynamics of actin

assembly cannot be related to the endocytic stage. More importantly, only by using live cells is it

possible to trace a single CME event from start to finish, and to therefore identify those CME

events wherein no detectable actin is ever assembled so key parameters can be compared

between events with and without associated actin assembly.

By visualizing endogenously tagged AP2M1 to mark the coat and CME initiation, and

DNM2 to mark the neck and scission, together with ARPC3 to specifically label Arp2/3-

nucleated, branched actin filaments (Supplementary Fig. 1a), we were able to precisely study the

spatial and temporal regulation of actin assembly during CME. Three-color labeling and analysis

of the displacement between markers for the three modules allowed us to distinguish *bona fide*

asymmetric actin assembly from events that artificially might appear asymmetric because the

invaginations were elongated and tilted (Fig2 a). Using TIRF live-cell imaging, we observed

ARPC3-labeled branched actin networks at lamellipodia and a subpopulation of CME sites (Fig.

2b, c). Dynamic actin assembly and disassembly occurred at CME sites with different spatio-

temporal characteristics, including discrete CME sites, clathrin plaques and at clathrin coat

splitting sites, as previously reported[14] (Fig. 2d and Supplementary Fig. 2a, b). In the analysis

described below, we focus on the discrete CME events and not the more complex ones (plaques

and splitting events). Analysis of these events with 1s/frame temporal resolution revealed that

ARPC3 is most robustly recruited during the late stages of CME shortly before scission[12] (Fig.

2c, d). Interestingly, we observed clear spatial displacement between ARPC3 (actin module) and

AP2 (coat module) as well as between ARPC3 and DNM2 (neck) before vesicle scission (Fig.

2d). This observation supports the conclusion that asymmetric branched actin networks provide

forces at endocytic sites through the time of scission. Imaging fluorescent beads using the same

settings indicates that the displacement is not an artifact caused by misalignment between

different imaging channels (Supplementary Video 3 and Supplementary Fig. 2c).

To analyze the intrinsic recruitment order and timing for up to three endocytic proteins at

CME sites quantitatively and systematically, we developed an automated, high-throughput

method that avoids bias because it does not involve manual selection of CME sites (see Materials

and Methods). Briefly, AP2 tracks were identified using standard particle-tracking algorithms[30].

Novel filtering methods then extracted DNM2-positive events marked by one or more DNM2

burst. The AP2 and DNM2 tracks were decomposed into dynamic features describing the events'

position and brightness. These features were used for clustering via unsupervised machine

learning, which enabled grouping of similarly-behaved tracks (Supplementary Fig. 3a and b).

DNM2-positive events were refined by a detection scheme that determined the number of DNM2

peaks using various characteristics of a single DNM2-peak: the peak height, width, and

minimum peak-to-peak distance (Supplementary Fig. 3c). Events with a single DNM2 peak were

analyzed as described below. The method detects low signals from endogenously tagged CME

proteins, such as the low-level recruitment of DNM2 at the early stages of CME, and accurately

reveals the different CME stages (Extended Data Fig. 3d).

Next, the timing of actin network assembly at CME sites was determined using ARPC3 as a

branched actin filament marker by analyzing over one thousand CME events. Although actin

appearance early in CME has been reported[14], determining the actin assembly timing is

challenging because it is difficult to distinguish newly assembled branched actin at CME sites

from the nearby cortical actin filaments or actin filaments attached to other vesicles or

organelles. Also, whether actin functions during the early stage of CME has not yet been shown

conclusively due to the potential side effects such as changes in membrane tension caused by

actin inhibitors. Our endogenous ARPC3 tagging and large-scale computational analysis

approach sidesteps these problems. We classified ARPC3 positive CME events into two groups:

one group with ARPC3 appearance early in CME, and the other with late appearance in CME

(Fig. 3a). We observed that in most of the events (N=1,385, 67.8%) a sharply increasing ARPC3

signal appears with similar timing to the rapid-recruitment phase of DNM2 concomitant with the

U to Ω membrane shape transition. This timing is consistent with previously proposed role for

actin in membrane invagination, as studies showed that actin inhibitors block the U to Ω

endocytic membrane shape transition[6,14]. In some cases (N=657, 32.2%) we detected ARPC3

signals at early CME stages. To test whether random overlap between nearby actin structures and

CME sites might be responsible for the apparent early actin recruitment, we generated a

randomized data set by pairing ARPC3 images with AP2 and DNM2 images from an unrelated

movie (Fig. 3b). In this data set, we detected early "assembly" of actin in the majority of ARPC3

positive CME events (N=17,282, 72.9%), and the intensity profiles of these events resembled the

early-actin CME events we observed in the real data set (Fig. 3a, b). Therefore, we conclude that

the presence of actin early in CME is very likely due to unrelated nearby actin structures

overlapping with CME sites.

Our live-cell analysis allowed the timing of branched actin network assembly to be

compared to the scission timing, and the spatial offset between the clathrin coat and the

associated actin network to be determined. Super-resolution imaging of yeast CME sites

suggested that actin and actin nucleators localize symmetrically in a ring around CME sites, and

computational modeling suggested that an asymmetric actin arrangement would not provide

sufficient force for the membrane invagination during yeast CME[13]. In contrast, in mammalian

cells, which require less actin force production during CME, imaging of fixed cells suggested

that actin structures associate adjacent to apparent flat clathrin coats. However, these studies

proposed that at the later CME stages the actin structures become larger and more symmetric to

provide sufficient force for membrane deformation and scission[14,15]. Surprisingly, in our live cell

studies designed to highlight sites of new actin assembly, we observed off-centered branched

actin networks at CME sites throughout even the latest CME stages (Fig. 2d). Furthermore, most

ARPC3-positive CME sites accomplish scission within 30s from the initiation of ARPC3

recruitment (Fig. 3c). The actin networks were off center from the coat and neck signals by

approximately 150nm at the time of vesicle scission (Fig. 3d). Given the temporal separation

between channel acquisition and the movement of AP2 spots, the separation between channels

can be attributed in part to an imaging artifact. Therefore, when we measured the average

movement of AP2 spots leading up to scission as a control, we found that over 95% of the events

had AP2-ARPC3 separations that exceed the frame-to-frame motility of AP2 (Supplementary

Fig 4). Also, the uncertainties measured by a standard deviation, when measuring the fitted

position of AP2, range up to 40 nm. Therefore, we include the AP2-DNM2 separation as a basis

for comparison to the AP2-ARPC3 and DNM2-ARPC3 separations (Fig 3d). These results

further support our conclusion that branched actin networks assemble asymmetrically at CME

sites through the time of scission (Fig 2d). This observation is consistent with the observation

that ring-shaped actin structures at clathrin coats were rarely observed in the high-resolution,

live-cell imaging in a previous study[31]. In total, these live-cell data suggest that in mammalian

cells, asymmetric actin network assembly can provide enough force to assist membrane

deformation and scission during the late stages of CME.

Asymmetric branched actin networks facilitate CME at stalled sites

To gain additional insights into the function of this asymmetric actin network assembly, we

quantitatively compared kinetics of CME events with or without ARPC3 recruitment. We

observed that about 30% of CME events are completed in the absence of detectable actin

assembly, which is consistent with the hypothesis that in mammalian cells actin assembly is

required for CME only under relatively high membrane tension, which can vary regionally

within cells[6,7,9]. Consistent with the possibility that increased membrane tension stalls membrane

deformation during CME[4,8,9,32–34], CME lifetimes were markedly longer for ARPC3 positive

events compared to the ARPC3 negative events (Fig. 4a). In addition, when the AP2M1 intensity

vs time profiles were compared between ARPC3 positive and negative CME sites, a plateau,

which lasts for approximately 10 seconds, was observed for the ARPC3 positive events (Fig. 4b).

Based on these observations and previous experimental and computational modeling data[4,6,7], we

propose that this plateau in branched actin-positive CME events represents stalled membrane

bending due to an unfavorable local membrane environment, such as higher membrane

tension[4,32,33].

We next tested the hypothesis that the asymmetric actin network might affect the lateral

movements of endocytic coats on the plasma membrane. Interestingly, the ARPC3 positive CME

sites showed significantly slower, but more directional lateral movement before the scission

compared to the ARPC3 negative CME sites (Fig. 4c, d). After scission both ARPC3 positive

and negative vesicles showed fast, apparently random movements (Fig. 4c, d). These data

suggest that the asymmetric actin can stabilize the forming endocytic coat while pushing it in the

plane of the plasma membrane with a lateral directional force.

N-WASP is recruited asymmetrically to the stalled CME sites

To further explore how the asymmetrical assembly of actin networks at CME sites is

regulated, we endogenously tagged N-WASP, an actin nucleation promoting factor (NPF) that

plays roles in CME, in AP2M1-tagRFP-T/ DNM2-tagGFP2 genome-edited iPSCs (hereafter

referred to as ADW cells, Fig. 5a and Supplementary Fig. 5a). Quantitative imaging of budding

yeasts demonstrated that initiation of productive actin assembly at CME sites requires the

accumulation of yeast WASP or WIP to a certain amount[35]. In our genome-edited iPS cells, we

observed that N-WASP is recruited asymmetrically to CME sites mostly at the late stage of CME

(Fig. 5b, c and Supplementary Fig. 5b, c). Longer lifetimes and a plateau in the AP2 intensity vs

time plot were observed specifically in the N-WASP positive CME events (Fig. 5d, e), similar to

the ARPC3 positive events (Fig 4 a, b). These data indicate that asymmetric NPF recruitment

underlies the asymmetric architecture of branched actin networks at CME sites.

Discussion

Using large-scale, comprehensive analysis of thousands of CME sites in unperturbed live

cells, our study demonstrates that in mammalian cells coat assembly dynamics predict which

sites will assemble actin, and show that at apparently stalled sites, actin assembles

asymmetrically to facilitate successful vesicle formation.

Based on the data presented here, we propose an updated model for actin assembly at

mammalian CME sites in which, beyond global tension-dependent changes in requirement for

actin assembly, highly localized differences give rise to heterogeneity even within the same

patch of plasma membrane in the same cell (Fig. 6): (1) Where the local membrane tension is

lower (Fig. 6 upper scenario), the membrane can undergo flat-U-Ω shape transitions without

actin assembly in a relatively short time. When the coat grows large enough to form a Ω-shaped

bud, sufficient dynamin can be recruited to perform scission, and there is little delay between

coat expansion and scission; (2) Where the local conditions are not favorable, presumably under

high membrane tension and possibly other impediments, the coat protein-membrane interaction

does not generate sufficient force to curve the membrane (Fig. 6 lower scenario). Here, extra

force generation from actin assembly is required[4,6]. Asymmetric N-WASP recruitment activates

actin nucleation mostly at one side of the clathrin coat, generating an asymmetric force that pulls

the membrane into the cell with a similar action to a bottle cap opener. We speculate that this

asymmetrical force contributes to asymmetric membrane deformation at endocytic sites observed

by high-speed atomic force microscopy[36] and may act with dynamin[37] to twist the clathrin pit to

promote scission at the neck. CME events with associated actin assembly have longer lifetimes,

likely due to a delay between coat expansion and scission, requiring adaptive recruitment of actin

regulators followed by actin network assembly and membrane remodeling. This result reinforces

the conclusion from previous studies[6–9] that increased membrane tension enhances the

requirement for actin assembly during CME, but also establishes that site-to-site heterogeneity in

actin dependence and involvement can be observed without manipulating membrane tension.

Future computational modeling studies of how asymmetric actin network assembly provides

forces for vesicle formation and membrane remodeling will deepen our understanding of actin's

functions in a host of actin-mediated processes.

Our model provides further insights into the basis for inconsistent effects of actin drugs on

CME[6,14,18,38–43]. Actin plays crucial roles in membrane shaping, cell adhesion, and membrane

tension. Global disruption of actin dynamics is expected to dramatically change membrane

tension and the available pool of actin and associated proteins and therefore to have both direct

and indirect effects on CME. Here, we focused on in-depth analysis of the unperturbed process

and detected preference for actin assembly at stalled CME events.

The results presented here may prove relevant to the constant coat area vs constant coat

curvature debate for how the clathrin coat assembles and develops curvature[32,44–48]. In the

constant area model, flat clathrin coats grow close to their final size before curvature develops as

a vesicle forms. In the constant curvature model, clathrin coats grow with a fixed curvature. Our

observations suggest that coat expansion and curvature generation may be regulated via distinct

mechanisms, with different actin requirements. At actin-positive CME sites, actin assembles

primarily at the late stage of CME when coat assembly is mostly complete (Fig. 3 and

Supplementary Fig. 6a). If the constant curvature model holds, actin should only be associated

with clathrin coats with highly curved dome and spherical shapes (Supplementary Fig. 6b).

However, actin associated with flat or shallow clathrin coats has been observed in multiple

studies[7,15], which supports the constant area model at actin-positive CME sites (Supplementary

Fig. 6c). On the other hand, mathematical modeling predicts that at actin-negative CME sites,

where coat and other curvature-promoting proteins provide sufficient force to bend the

membrane[23], the membrane is smoothly and continuously shaped by these proteins into a budded

morphology as the coat area increases[4], which follows the constant curvature model. Perhaps the

constant area model applies primarily for actin-negative sites and the constant area model applies

primarily for actin-positive sites, which we have shown here are mostly stalled CME sites. We

suggest that in future studies the constant coat area and constant coat curvature models be tested

at individual CME events to test the possibility that both mechanisms operate in the same cell.

Acknowledgments: MJ was funded by American Heart Association Postdoctoral Fellowship

(18POST34000029). DGD was funded by NIH MIRA grant R35GM118149. KX is a

Chan Zuckerberg Biohub investigator and acknowledges support from NIH (DP2GM132681).

SU was funded by Philomathia Foundation and the Chan Zuckerberg Initiative Imaging Scientist

program. The authors would like to thank Dr. Yidi Sun and Dr. Matthew Akamatsu for insightful

comments on the manuscript; the Conklin Lab at UCSF for providing WTC10 human iPSC line;

the Lavis Lab at Janelia Research Campus for providing JF635 HaloTag ligand; Dr. Sun Hae

Hong for generating the AP2-tagRFP-T iPSC cell line; the UC Berkeley QB3 MacroLab for

purified S. pyogenes NLS-Cas9; the UC Berkeley Cancer Research Laboratory Flow Cytometry

Facility for iPSC sorting.

Data availability

The raw live-cell imaging data (TIRF) can be found at

https://github.com/DrubinBarnes/Jin_Shirazinejad_et_al_branched_actin_manuscript. All other

raw data are available from the corresponding author upon request.

Code availability

The Jupyter Notebooks used for live-cell imaging analysis can be found at

https://github.com/DrubinBarnes/Jin_Shirazinejad_et_al_branched_actin_manuscript.

METHODS

Cell culture

The WTC10 hiPSC line was obtained from the Bruce Conklin Lab at UCSF. hiPSCs were cultured on Matrigel (hESC-Qualified Matrix, Corning) in StemFlex medium (Thermo Fisher) with Penicillin/ Streptomycin in 37°C, 5% CO_2. Cultures were passaged with Gentle Cell Dissociation reagent (StemCell Technologies, Cat#: 100-0485) twice every week.

Genome-editing

The AP2M1 gene was edited in WTC10 hiPSCs as previously described using TALENs targeting exon 7 of the AP2M1 gene[49]. Both alleles of AP2M1 were tagged with tagRFP-T. The Cas9-crRNAtracrRNA complex electroporation method was used sequentially to edit DNM2 and ARPC3 gene in AP2M1-tagRFP-T genome-edited hiPSCs, as previously described[12,18]. The same method was used to edit the WASL gene in AP2M1-tagRFP-T/DNM2-tagGFP2 genome edited hiPSCs. *S. pyogenes* NLS-Cas9 was purified in the University of California Berkeley QB3 MacroLab. TracrRNA and crRNA that target CCTGCTCGACTAGGCCTCGA (DNM2), CCTGGACAGTGAAGGGAGCC (ARPC3) and AGCTCATGGTTTCGCCGGCG (WASL), were purchased from IDT. Gibson assembly (New England Biolabs) was used to construct donor plasmids containing DNM2 5' homology-ggtaccagtggcggaagc-tagGFP2-DNM2 3' homology, ARPC3 5' homology-ggatccggtaccagcgatccaccggtcgccacc-HaloTag-ARPC3 3' homology, and WASL 5' homology-HaloTag-agcgatccaccggtcgccaccggatcc-WASL 3' homology sequences, respectively. Three days after electroporation (Lonza, Cat#: VPH-5012) of the Cas9-crRNA-tracrRNA complex and donor plasmid, the tagGFP2 or HaloTag positive cells were single cell

sorted using a BD Bioscience Influx sorter (BD Bioscience) into Matrigel-coated 96-well plates.

Clones were confirmed by PCR and Sanger sequencing of the genomic DNA locus around the

insertion site. Both alleles of DNM2 and ARPC3 were tagged with tagGFP2 and HaloTag,

respectively, and one allele of WASL was tagged with HaloTag in the hiPSC lines used in this

study.

Western blotting

Cells were dissociated from the well using Gentle Cell Dissociation reagent (StemCell

Technologies, Cat#: 100-0485). Total proteins were extracted by adding 1ml of cold 10% TCA

to the cell pellets, incubated on ice for 30min, and spun down by centrifuging at 4 °C, 12000rpm

for 10min. Protein pellets were dissolved in loading buffer (50 mM HEPES, pH 7.4, 150 mM

NaCl, 1 mM MgCl2, 5% BME, 5mM DTT and protease inhibitor) and loaded onto an

acrylamide gel for SDS-PAGE and transferred to nitrocellulose membranes for immunoblotting.

Blots were incubated overnight at 4°C with primary antibodies targeting Tag(CGY)FP (1:2000

dilution in 1% milk, Evrogen, Cat#: AB121), HaloTag (1:1000 dilution in 0.5% milk, Promega,

Cat#: G9211), GAPDH (1:100,000 dilution in 0.5% milk, Proteintech, Cat#: 10494-1-AP),

respectively, and subsequently incubated in the dark at room temperature for 1hr with secondary

antibodies.

TIRF live-cell imaging

Two days before imaging, hiPSCs were seeded onto Matrigel-coated 4-well chambered cover

glasses (Cellvis). Halotag was labeled by JF635- HaloTag ligand[50]. Cells were incubated in

StemFlex medium with 100 mM JF635-HaloTag for 45min and the unbound ligands were

washed away by three washes with 5 min incubation in prewarmed StemFlex medium. Cells

were imaged on a Nikon Ti-2 inverted microscope fitted with TIRF optics and a sCMOS camera

(Hamamatsu). Cells were maintained at 37 °C with a stage top incubator (OKO Lab) in

StemFlex medium with 10mM HEPES. Images were acquired with Nikon Elements. Channels

were acquired sequentially at a 1 sec interval and 300ms exposure time over 4 minutes.

TIRF image processing

Four generalized processing steps were applied identify of clathrin-coated pits with single

DNM2 peaks: track feature abstraction, feature dimensionality reduction, event clustering, and

DNM2-peak detection. First, tracks that are defined by fitted positions and intensities for single

events were generated using cmeAnalysis[30]. Then, AP2 and DNM2 tracks were decomposed into

dynamic features describing the dynamics of the events' position and brightness. The mapping of

each track to discrete features was done to generalize the dynamics of tracked events into a set of

interpretable coordinates. These features were clustered after feature scaling to normal

distributions, dimensionality reduction with principal component analysis, and Gaussian mixture

modeling. DNM2-positive events represented a distinct cluster of tracks that had detectable

DNM2 throughout the event, were long lived, and were below the threshold of motility expected

for transient, non-CME-derived clathrin-coated vesicle "visitors" at the TIRF field. Single

DNM2-peak events were found by searching over a range of values set for the minimum DNM2

peak height, width, and peak-to-peak temporal distance. After finding single-peaked events in a

fixed peak-parameter combination, the lifetime distribution of single peak events' lifetimes were

fit to the expected underlying distribution, a Rayleigh distribution[51], where the best-fitting

parameter combination was chosen to identify single-peaked events. Single DNM2-peaked

events were kept as CME sites for the remainder of the analysis. All code associated with this

analysis, generating Figures 3-5, and a detailed step-by-step protocol, are available at

https://github.com/DrubinBarnes/Jin_Shirazinejad_et_al_branched_actin_manuscript.

Two-color 3D STORM imaging

12 mm round coverslips were sonicated in distilled water and sterilized for 20 min in 70%

ethanol, air-dried and coated with Matrigel in 24-well plates. Cells were seeded onto Matrigel-

coated coverslips two days before fixation. For clathrin and actin two-color imaging, cells were

fixed first for 1 min in 0.3% (v/v) glutaraldehyde (GA) solution containing 0.25% (v/v) Triton in

cytoskeleton buffer (CB: 10mM MES, 150mM NaCl, 5mM EGTA, 5mM Glucose, 5mM $MgCl_2$,

0.005% NaN_3, pH 6.1) and then immediately fixed for 20 min in 2% (v/v) GA solution in CB.

Both solutions were prepared fresh from a 10% GA stock (Electron Microscopy Science, cat

#16120). After fixation, samples were incubated twice for 5 min in freshly prepared 0.1% (w/v)

$NaBH_4$ in PBS. For clathrin and ARPC3-HaloTag imaging, cells were fixed for 20 min in 4%

(v/v) PFA (Electron Microscopy Sciences, Cat#: 15710) in CB. Subsequently, both types of

samples were washed 3 times for 10 min in PBS. Samples were then blocked for 20 min in

blocking buffer [3% (w/v) BSA and 0.1% (w/v) Saponin in PBS]. Clathrin light chain

(Invitrogen, Cat#: MA5-11860, 1:200 dilution) and Halotag (Promega, Cat#: G9281, 1:200

dilution) antibodies were used in blocking solution. Primary antibody immunostaining was

performed overnight at 4°C. On the next day, samples were washed three times in washing

buffer (0.1x blocking buffer in PBS) for 10 min. Samples were incubated with secondary

antibody in blocking buffer for 30 min at room temperature in the dark and were washed three

times for 10 min in washing buffer, and then three times for 10 min in PBS. Homemade mouse

secondary antibody-CF680 (1:50) was used to stain clathrin and actin samples. Commercial

mouse secondary antibody-AF647 (ThermoFisher, cat#A32787; 1:400) and homemade rabbit

secondary antibody-CF680 (1:50) were used to stain the clathrin and ARPC3-HaloTag. Clathrin

and actin samples were then stained with 0.5μM Phalloidin-AF647 (Fisher Scientific, Cat#:

A22287) in PBS and kept at room temperature in the dark for 2 hours. Samples were washed

three times with PBS before STORM imaging.

STORM imaging was performed as previously described on a homebuilt STORM setup[7,52].

Samples labeled by AF647 and CF680 were excited by an 647nm laser. The emission of both

AF647 and CF680 was then split into two light paths as two channels using a dichroic mirror

(Chroma, cat#T685lpxr), and each channel was projected onto one-half of an EMCCD camera

(Andor iXon Ultra 897). Color assignment of each localization was based on its intensity in the

two channels. A cylindrical lens was inserted into the transmitted channel to acquire 3D

localization[28]. 3D position of each localization was determined from the ellipticity of each point

spread function.

Fig. 1: Two-color, 3D stochastic optical reconstruction microscopy (STORM) shows that actin structures are off-centered with respect to clathrin coats.

Fig. 1: Two-color, 3D stochastic optical reconstruction microscopy (STORM) shows that

actin structures are off-centered with respect to clathrin coats. a, b, Two-color 3D STORM

image of bottom membrane of ADA cells immunolabeled with clathrin light chain antibody

(clathrin, CF-680, magenta) and phalloidin (actin, AF647, rainbow). **c, d,** Two color 3D STORM

423   image of the bottom membrane of ADA cells immunolabeled with clathrin light chain antibody

424   (clathrin, AF647, magenta) and HaloTag antibody (ARPC3-HaloTag, CF-680, rainbow). Dotted

425   lines label lamellipodia. **b, d,** The highlighted CME sites, which are labeled by white arrows in

426   (**a**) and (**c**), are rotated and shown in magnified top and side view projections. Color bar shows

427   the z position of ARPC3-HaloTag. Scale bars: 2µm, 100nm.

Fig. 2: Triple-genome-edited iPS cells reveal dynamic actin organization at CME sites.

Fig. 2: Triple-genome-edited iPS cells reveal dynamic actin organization at CME sites. a,
Models of branched actin assembly at invaginating CME sites. Model 1: Asymmetric actin
assembly at CME sites results in separated actin-coat and actin-neck signals. Model 2:
Symmetric actin assembly at tilted CME sites results in separated actin-coat signals but

overlapped actin-neck signals. Model 3: Symmetric actin assembly at perpendicularly

invaginating CME sites will result overlapped actin, coat and neck signals. **b,** A representative

single time frame image of a TIRF movie (Supplementary Video 2) of AP2M1-tagRFP-T

(magenta), DNM2-tagGFP2 (green) and JF635 ligand[50]-conjugated ARPC3-HaloTag (cyan) in

ADA cells. The highlighted region is boxed by a dashed line. Scale bar: 5µm. **c,** A representative

kymograph of AP2M1-tagRFP-T (magenta), DNM2-tagGFP2 (green) and JF635 ligand-

conjugated ARPC3-HaloTag (cyan) at CME sites in ADA cells. Scale bar: 5µm. **d,** Montage of a

representative ARPC3 positive CME site in ADA cells. Individual channels and pair-wise

merges are shown. *: Images from one frame before scission (maximum DNM2 intensity) are

marked to show the displacement between the CME coat (AP2)-ARPC3 and CME neck

(DNM2)-ARPC3. Size of field of view: 2µm x 2µm. Intervals: 1sec.

Fig. 3: Computational analysis of ARPC3 positive CME sites reveals asymmetric actin network assembly at the late stage of CME.

Fig. 3: Computational analysis of ARPC3 positive CME sites reveals asymmetric actin network assembly at the late stage of CME. a, b, Averaged intensity vs time plots of cohorts of ARPC3 positive CME sites in ADA cells (**a**) and in the randomized data set (**b**). Events are grouped by the timing of ARPC3-labeled branched actin network recruitment (early: top, late: bottom), and then grouped into cohorts by the lifetimes of AP2 and aligned to the frames showing the maximum DNM2 intensity (time = 0s). Total number of CME sites in each group is shown in parentheses. Percentage of the number of the CME sites in each cohort is shown next to the plot. **c,** Histogram of ARPC3-mediated actin network assembly duration. The assembly duration is measured from the first frame of the ARPC3 signal to the presumed scission time (the peak of DNM2 signal). **d,** Averaged intensity (solid lines) and distance (dashed lines) vs time plots of ARPC3 positive CME sites in ADA cells. Events are aligned to the frames showing the maximum DNM2 intensity (time = 0s). Distance between centers of two signals are shown from -10s to 3s when DNM2 and ARPC3 signals are relatively high. N=1,385. **a, b, d,** Error bar: ¼ standard deviation.

Fig. 4: Actin positive CME sites show distinct dynamics.

Fig. 4: Actin positive CME sites show distinct dynamics. a, Histograms of ARPC3 negative

(blue) and positive (orange) CME lifetimes. CME lifetime is measured from the first frame of the

AP2 signal to the presumed scission time (the peak of DNM2 signal). ARPC3 positive CME

events have longer lifetimes. **b,** Averaged intensity vs time plots of ARPC3 negative (top) and

positive (bottom) CME sites in ADA cells. Events were aligned to the frames showing the

maximum DNM2 intensity. Error bar: ¼ standard deviation. **c,** Lateral motility of ARPC3

negative (blue) and positive (yellow) CME sites before (solid line) and after (dashed line) vesicle

scission. ARPC3 positive CME sites move slower than ARPC3 negative ones. **d,** Straightness-

index of ARPC3 negative (blue) and positive (yellow) CME sites before (solid line) and after

(dashed line) scission. The straightness-index is defined by the ratio between the sum of frame-

to-frame distances to the end-to-end distance of a single event's trajectory, where a perfectly

straight-lined trajectory would have an index of 1. APRC3 positive CME sites move with a

straighter trajectory. **a-d,** ARPC3 -: N=840, ARPC3 +: N=1,385.

Fig. 5: Asymmetric N-WASP recruitment to stalled CME sites.

Fig. 5: Asymmetric N-WASP recruitment to stalled CME sites. a, A representative single

time frame image of a TIRF movie (Supplementary Video 4) of AP2M1-tagRFP-T (magenta),

DNM2-tagGFP2 (green) and JF635 ligand-conjugated HaloTag-N-WASP (cyan) in ADW cells.

The highlighted region is boxed by a dashed line. Scale bar: 5μm. **b,** A representative

kymograph of CME sites in ADW cells over a 4 min movie. Scale bar: 5μm. **c,** Averaged

intensity (solid line) and distance (dashed line) vs time plots of N-WASP positive CME sites in

ADW cells. Events are aligned to the frames showing the maximum DNM2 intensity. Intensity is

scaled to 1 at peaks for each channel. Error bar: ¼ standard deviation. **d,** N-WASP positive CME

483　　events have longer lifetimes. **e,** Intensity vs time plots of averaged N-WASP negative (top) and

484　　positive (bottom) CME sites in ADW cells. Events were aligned to the frames showing the

485　　maximum DNM2 intensity. Error bar: ¼ standard deviation. **c-d,** N-WASP negative CME sites:

486　　N=385, N-WASP positive CME sites: N=1,381

Fig. 6: An updated schematic model of actin negative and actin positive CCPs in human cells.

Fig. 6: An updated schematic model of actin-negative and actin-positive clathrin-coated pits in human cells. Actin assembly is induced at stalled CME sites, where asymmetric forces pull, bend and possibly twist the plasma membrane against membrane tension to drive membrane invagination and vesicle scission.

Supplementary Fig. 1: Genome-edited iPSCs show dynamic CME sites.

Supplementary Fig 1: Genome-edited iPSCs show dynamic CME sites. a, Schematic model

of CME. Mammalian CME proteins can be grouped into several modules, including the coat,

WASP and Myosin / actin nucleation promoting factor (NPF), actin and scission modules[5]. Actin

networks provide pulling forces to invaginate the membrane against membrane tension[4,6,9,12]. **b,**

Immunoblot analysis of cell extracts from the WT (WTC) and genome-edited (AP2M1-tagRFP-

T/DNM2-tagGFP2/ARPC3-HaloTag; ADA) human iPSCs. The labeled proteins were detected

with tag(CGY)FP, HaloTag, and GAPDH (loading control) antisera respectively. **c,** Kymograph

of representative CME sites of double-edited (AP2M1-tagRFP-T/DNM2-tagGFP2; AD) and

triple-edited (AP2M1-tagRFP-T/DNM2-tagGFP2/ARPC3-HaloTag; ADA) cells.

Supplementary Fig 2: Actin assembles at different types of CME sites.

Supplementary Fig 2. Actin assembles at different types of CME sites. a, Montage of a

representative ARPC3 positive CME plaque from a TIRF movie of triple-edited (AP2M1-

tagRFP-T/DNM2-tagGFP2/ARPC3-HaloTag; ADA) human iPSCs (Supplementary Video 2). **b,**

Montage of a representative ARPC3 positive splitting CME site from a TIRF movie of triple-

edited (AP2M1-tagRFP-T/DNM2-tagGFP2/ARPC3-HaloTag; ADA) human iPSCs

510   (Supplementary Video 2). **c,** Montage from a TIRF movie of a multi-fluorescence bead

511   (Supplementary Video 3). Size of field of view: 2μm x 2μm. Intervals: 1sec.

Supplementary Fig 3: Filtering methods for selection of CME sites.

Supplementary Fig 3. Filtering methods for selection of CME sites. a, 2-D histogram of the first two principal components (PCs) of AP2 and DNM2 dynamic features. The shaded underlay represents simulated data points in principal component space and their individual probabilities of belonging to the nearest cluster center. Cluster 0 shows data points in the DNM2-positive cluster. **b,** Cohort plots of the shortest AP2 events (<40 seconds) from each cluster. Cluster 0 represents DNM2-positive events where a strong DNM2 signal is detected. **c,** DNM2-positive events are sorted by the number of DNM2 peaks using a peak-detection scheme. Representative intensity vs time plots of a single-peaked event (left) and a multi-peaked event (right). **d,** Single-peaked DNM2 events, hereon named CME sites, are grouped into lifetime cohorts and aligned to the peak of the DNM2 channel.

Supplementary Fig 4: AP2-ARPC3 separation is not due to imaging artifacts.

Supplementary Fig 4: AP2-ARPC3 separation is not due to imaging artifacts. A heat map

graph of distance between AP2 and ARPC3 signals before scission, and average AP2 frame to

frame displacement within 6 seconds before scission. Over 95% of the CME events present

larger AP2-ARPC3 separation than AP2 displacement. N= 1,385.

Supplementary Fig 5. Dynamics of N-WASP at CME sites.

Supplementary Fig 5: Dynamics of N-WASP at CME sites. a, Immunoblot analysis of cell extracts from the control and genome-edited (AP2M1-tagRFP-T/DNM2-tagGFP2/HaloTag-WASL; ADW) human iPSCs. The labeled proteins were detected with HaloTag and GAPDH (loading control) antisera respectively. **b,** Histogram of N-WASP lifetime at CME sites. The lifetime is measured from the first frame of the N-WASP signal to the presumed scission time (the peak of DNM2 signal). **c,** Intensity vs time plots of cohorts of N-WASP positive CME sites in ADW cells. Events are grouped into cohorts by the lifetimes of AP2 and aligned to the frames showing the maximum DNM2 intensity. N=1,381. Error bar: 1/4 standard deviation.

540 **Supplementary Fig 6: Constant curvature vs constant area models for how clathrin coats**

541 **assemble at actin-positive CME sites. a,** A sketch showing amounts of CME coat and actin

542 module proteins at actin-positive CME sites as a function of time based on the data in Fig. 4b.

543 The CME coat is assembled to its maximum area around the time of actin assembly initiation. **b,**

544 Schematic representation of constant curvature model for CME. CME coat assembles during

545 invagination and actin assembles only at deep invaginations. **c,** Schematic representation of

546 constant area model for CME. The CME coat expands to its maximum area first and bends

547 during membrane invagination. In these two different scenarios, actin assembles at CME sites

548 with different curvatures.

REFERENCES

1. Rottner, K., Faix, J., Bogdan, S., Linder, S. & Kerkhoff, E. Actin assembly mechanisms at a glance. *J. Cell Sci.* **130**, 3427–3435 (2017).

2. Lu, R., Drubin, D. G. & Sun, Y. Clathrin-mediated endocytosis in budding yeast at a glance. *J. Cell Sci.* **129**, 1531–6 (2016).

3. Lacy, M. M., Ma, R., Ravindra, N. G. & Berro, J. Molecular mechanisms of force production in clathrin-mediated endocytosis. *FEBS Letters* (2018). doi:10.1002/1873-3468.13192

4. Hassinger, J. E., Oster, G., Drubin, D. G. & Rangamani, P. Design principles for robust vesiculation in clathrin-mediated endocytosis. *Proc. Natl. Acad. Sci. U. S. A.* (2017). doi:10.1073/pnas.1617705114

5. Lanzetti, L. Actin in membrane trafficking. *Current Opinion in Cell Biology* (2007). doi:10.1016/j.ceb.2007.04.017

6. Boulant, S., Kural, C., Zeeh, J. C., Ubelmann, F. & Kirchhausen, T. Actin dynamics counteract membrane tension during clathrin-mediated endocytosis. *Nat. Cell Biol.* **13**, 1124–1132 (2011).

7. Kaplan, C. *et al.* Adaptive actin organization buffers endocytosis against changes in membrane tension. *bioRxiv* (2021).

8. Batchelder, E. M. & Yarar, D. Differential requirements for clathrin-dependent endocytosis at sites of cell-substrate adhesion. *Mol. Biol. Cell* **21**, 3070–3079 (2010).

9. Kaur, S., Fielding, A. B., Gassner, G., Carter, N. J. & Royle, S. J. An unmet actin requirement explains the mitotic inhibition of clathrin-mediated endocytosis. *Elife* **3**, 1–17

(2014).

10.   Grassart, A. *et al.* Actin and dynamin2 dynamics and interplay during clathrin-mediated

endocytosis. *J. Cell Biol.* (2014). doi:10.1083/jcb.201403041

11.   Boulant, S., Kural, C., Zeeh, J. C., Ubelmann, F. & Kirchhausen, T. Actin dynamics

counteract membrane tension during clathrin-mediated endocytosis. *Nat. Cell Biol.*

(2011). doi:10.1038/ncb2307

12.   Akamatsu, M. *et al.* Principles of self-organization and load adaptation by the actin

cytoskeleton during clathrin-mediated endocytosis. *Elife* (2020). doi:10.7554/eLife.49840

13.   Mund, M. *et al.* Systematic Nanoscale Analysis of Endocytosis Links Efficient Vesicle

Formation to Patterned Actin Nucleation. *Cell* (2018). doi:10.1016/j.cell.2018.06.032

14.   Yarar, D., Waterman-Storer, C. M. & Schmid, S. L. A dynamic actin cytoskeleton

functions at multiple stages of clathrin-mediated endocytosis. *Mol. Biol. Cell* **16**, 964–975

(2005).

15.   Collins, A., Warrington, A., Taylor, K. A. & Svitkina, T. Structural organization of the

actin cytoskeleton at sites of clathrin-mediated endocytosis. *Curr. Biol.* (2011).

doi:10.1016/j.cub.2011.05.048

16.   Taylor, M. J., Perrais, D. & Merrifield, C. J. A high precision survey of the molecular

dynamics of mammalian clathrin-mediated endocytosis. *PLoS Biol.* **9**, e1000604 (2011).

17.   Kaksonen, M., Toret, C. P. & Drubin, D. G. A modular design for the clathrin- and actin-

mediated endocytosis machinery. *Cell* **123**, 305–20 (2005).

18.   Dambournet, D. *et al.* Genome-edited human stem cells expressing fluorescently labeled

endocytic markers allow quantitative analysis of clathrin-mediated endocytosis during

differentiation. *J. Cell Biol.* **217**, 3301–3311 (2018).

19.   Doyon, J. B. *et al.* Rapid and efficient clathrin-mediated endocytosis revealed in genome-
edited mammalian cells. *Nat. Cell Biol.* **13**, 331–7 (2011).

20.   Cocucci, E., Gaudin, R. & Kirchhausen, T. Dynamin recruitment and membrane scission
at the neck of a clathrin-coated pit. *Mol. Biol. Cell* **25**, 3595–3609 (2014).

21.   Gibson, T. J., Seiler, M. & Veitia, R. A. The transience of transient overexpression. *Nat.*
Methods **10**, 715–21 (2013).

22.   Taylor, M. J., Lampe, M. & Merrifield, C. J. A feedback loop between dynamin and actin
recruitment during clathrin-mediated endocytosis. *PLoS Biol.* **10**, e1001302 (2012).

23.   Merrifield, C. J., Feldman, M. E., Wan, L. & Almers, W. Imaging actin and dynamin
recruitment during invagination of single clathrin-coated pits. *Nat. Cell Biol.* (2002).
doi:10.1038/ncb837

24.   María Cabeza, J., Acosta, J. & Alés, E. Dynamics and Regulation of Endocytotic Fission
Pores: Role of Calcium and Dynamin. *Traffic* (2010). doi:10.1111/j.1600-
0854.2010.01120.x

25.   Ramachandran, R. & Schmid, S. L. Real-time detection reveals that effectors couple
dynamin's GTP-dependent conformational changes to the membrane. *EMBO J.* (2008).
doi:10.1038/sj.emboj.7601961

26.   Roux, A. *et al.* Membrane curvature controls dynamin polymerization. *Proc. Natl. Acad.*
Sci. U. S. A. (2010). doi:10.1073/pnas.0913734107

27.   Iversen, T. G., Skretting, G., Van Deurs, B. & Sandvig, K. Clathrin-coated pits with long,
dynamin-wrapped necks upon expression of a clathrin antisense RNA. *Proc. Natl. Acad.*
Sci. U. S. A. (2003). doi:10.1073/pnas.0534231100

28.   Huang, B., Wang, W., Bates, M. & Zhuang, X. Three-Dimensional Super-Resolution

619   Reconstruction Microscopy. *Science.* **319**, 810–813 (2008).

620 29. Xu, K., Babcock, H. P. & Zhuang, X. Dual-objective STORM reveals three-dimensional

621   filament organization in the actin cytoskeleton. *Nat. Methods* **9**, 185–188 (2012).

622 30. Aguet, F., Antonescu, C. N., Mettlen, M., Schmid, S. L. & Danuser, G. Advances in

623   analysis of low signal-to-noise images link dynamin and AP2 to the functions of an

624   endocytic checkpoint. *Dev. Cell* (2013). doi:10.1016/j.devcel.2013.06.019

625 31. Li, D. *et al.* Extended-resolution structured illumination imaging of endocytic and

626   cytoskeletal dynamics. *Science.* (2015). doi:10.1126/science.aab3500

627 32. Bucher, D. *et al.* Clathrin-Adaptor ratio and membrane tension regulate the flat-To-curved

628   transition of the clathrin coat during endocytosis. *Nat. Commun.* (2018).

629   doi:10.1038/s41467-018-03533-0

630 33. Saleem, M. *et al.* A balance between membrane elasticity and polymerization energy sets

631   the shape of spherical clathrin coats. *Nat. Commun.* (2015). doi:10.1038/ncomms7249

632 34. Liu, A. P., Loerke, D., Schmid, S. L. & Danuser, G. Global and local regulation of

633   clathrin-coated pit dynamics detected on patterned substrates. *Biophys. J.* **97**, 1038–1047

634   (2009).

635 35. Sun, Y. *et al.* Switch-like Arp2/3 activation upon WASP and WIP recruitment to an

636   apparent threshold level by multivalent linker proteins in vivo. *Elife* **6**, 1–25 (2017).

637 36. Yoshida, A. *et al.* Morphological changes of plasma membrane and protein assembly

638   during clathrin-mediated endocytosis. *PLoS Biol.* **16**, (2018).

639 37. Cheng, X. *et al.* Dynamin-dependent vesicle twist at the final stage of clathrin-mediated

640   endocytosis. *Nat. Cell Biol.* **23**, 859–869 (2021).

641 38. Kaksonen, M., Sun, Y. & Drubin, D. G. A Pathway for Association of Receptors,

642   Adaptors, and Actin during Endocytic Internalization. *Cell* **115**, 475–87 (2003).

643 39. Aghamohammadzadeh, S. & Ayscough, K. R. Differential requirements for actin during

644   yeast and mammalian endocytosis. *Nat. Cell Biol.* **11**, 1039–42 (2009).

645 40. Ferguson, S. *et al.* Coordinated Actions of Actin and BAR Proteins Upstream of Dynamin

646   at Endocytic Clathrin-Coated Pits. *Dev. Cell* **17**, 811–22 (2009).

647 41. Durrbach, A., Louvard, D. & Coudrier, E. Actin filaments facilitate two steps of

648   endocytosis. *J. Cell Sci.* **109**, 457–65 (1996).

649 42. Lamaze, C., Fujimoto, L. M., Yin, H. L. & Schmid, S. L. The actin cytoskeleton is

650   required for receptor-mediated endocytosis in mammalian cells. *J. Biol. Chem.* **272**,

651   20332–5 (1997).

652 43. Miya Fujimoto, L., Roth, R., Heuser, J. E. & Schmid, S. L. Actin assembly plays a

653   variable, but not obligatory role in receptor-mediated endocytosis in mammalian cells.

654   *Traffic* (2000). doi:10.1034/j.1600-0854.2000.010208.x

655 44. Lampe, M., Vassilopoulos, S. & Merrifield, C. Clathrin coated pits, plaques and adhesion.

656   *J. Struct. Biol.* (2016). doi:10.1016/j.jsb.2016.07.009

657 45. Avinoam, O., Schorb, M., Beese, C. J., Briggs, J. A. G. & Kaksonen, M. Endocytic sites

658   mature by continuous bending and remodeling of the clathrin coat. *Science.* (2015).

659   doi:10.1126/science.aaa9555

660 46. Scott, B. L. *et al.* Membrane bending occurs at all stages of clathrincoat assembly and

661   defines endocytic dynamics. *Nat. Commun.* (2018). doi:10.1038/s41467-018-02818-8

662 47. Sochacki, K. A. *et al.* The structure and spontaneous curvature of clathrin lattices at the

663   plasma membrane. *bioRxiv* (2020).

664 48. Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nature Reviews*

Molecular Cell Biology (2018). doi:10.1038/nrm.2017.132

49.   Hong, S. H., Cortesio, C. L. & Drubin, D. G. Machine-Learning-Based Analysis in

Genome-Edited Cells Reveals the Efficiency of Clathrin-Mediated Endocytosis. *Cell Rep.*

12, 2121–2130 (2015).

50.   Grimm, J. B. *et al.* A general method to fine-tune fluorophores for live-cell and in vivo

imaging. *Nat. Methods* **14**, 987–994 (2017).

51.   Loerke, D. *et al.* Cargo and dynamin regulate clathrin-coated pit maturation. *PLoS Biol.* **7**,

0628–0639 (2009).

52.   Wojcik, M., Hauser, M., Li, W., Moon, S. & Xu, K. Graphene-enabled electron

microscopy and correlated super-resolution microscopy of wet cells. *Nat. Commun.* **6**, 2–7

(2015).