Autophagosomes fuse to phagosomes and facilitate the degradation of apoptotic cells in *Caenorhabditis elegans*

Omar Peña-Ramos[1], Lucia Chiao[1], Xianghua Liu[1,2], Xiaomeng Yu[1], Tianyou Yao[1], Henry He[1,3], and Zheng Zhou[1,4]

[1] Verna and Marrs McLean Department of Biochemistry and Molecular Biology, Baylor College of Medicine, Houston, TX 77030

[2] Current address: Caris Life Sciences, Inc., Tempe, AZ 85281

[3] Current address: Department of Neurology Residency Program, University of Texas Southwestern Medical Center, Dallas, TX 75390

[4] Corresponding author, zhengz@bcm.tmc.edu, 713-798-6489

Abstract

Autophagosomes are double-membrane intracellular vesicles that degrade protein aggregates, intracellular organelles, and other cellular components. During the development of the nematode *Caenorhabditis elegans*, many somatic and germ cells undergo apoptosis. These cells are engulfed and degraded by their neighboring cells. We discovered a novel role of autophagosomes in facilitating the degradation of apoptotic cells using a real-time imaging technique. Specifically, the double-membrane autophagosomes in engulfing cells are recruited to the surfaces of phagosomes containing apoptotic cells and subsequently fuse to phagosomes, allowing the inner vesicle to enter the phagosomal lumen. Mutants defective in the production of autophagosomes display significant defects in the degradation of apoptotic cells, demonstrating the importance of autophagosomes to this process. The signaling pathway led by the phagocytic receptor CED-1, the adaptor protein CED-6, and the large GTPase dynamin (DYN-1) promotes the recruitment of autophagosomes to phagosomes. Moreover, the subsequent fusion of autophagosomes with phagosomes requires the functions of the small GTPase RAB-7 and the HOPS complex components. Further observations suggest that autophagosomes provide apoptotic cell-degradation activities in addition to and in parallel of lysosomes. Our findings reveal that, unlike the single-membrane, LC3-associated phagocytosis (LAP) vesicles reported for mammalian phagocytes, the canonical double-membrane autophagosomes facilitate the clearance of *C. elegans* apoptotic cells. These findings add autophagosomes to the collection of intracellular organelles that contribute to

30   phagosome maturation, identify novel crosstalk between the autophagy and phagosome maturation

31   pathways, and discover the upstream signaling molecules that initiate this crosstalk.

 Introduction

During metazoan development and adulthood, a large number of cells undergo apoptosis or cell

suicide; these dying cells are engulfed by phagocytes and degraded inside phagosomes, vacuoles

composed of the lipid bilayers originated from the plasma membrane [1,2]. Swift engulfment and

degradation of apoptotic cells are critical for tissue remodeling, the resolution of the wound area, and the

prevention and suppression of harmful inflammatory and autoimmune responses induced by the content

of the dying cells [2]. Critical to the degradation of phagosomal contents is the fusion of intracellular

organelles, including lysosomes and early endosomes, to phagosomes, which results in the delivery of

the content of these organelles to the phagosomal lumen [3]. Lysosomes, which contribute many kinds

of hydrolytic enzymes, including proteases, nucleases, lipases, and hydrolyzing enzymes for

polysaccharides to the lumen of phagosomes, are the most pivotal organelles that support phagosomal

degradation [3]. The fusion of lysosomes to phagosomes also helps acidify the phagosomal lumen,

creating a low pH condition in which the digestive enzymes are active [3]. Besides lysosomes and

endosomes, whether other kinds of intracellular organelles fuse to phagosomes and contribute to the

degradation of the apoptotic cells inside remains elusive.

Mammalian microtubule-associated protein 1 light chain 3 (MAP1-LC3, or LC3) protein is a

member of the ATG8 protein family [4]. LC3 molecules conjugated to the lipid

phosphatidylethanolamine (PE) are most often observed on the surfaces of autophagosomes, double-

membrane organelles that are a key structure of autophagy [4]. In fact, LC3 is a well-established marker

for autophagosomes [4]. In mammalian cells, lipidated LC3 molecules were also reported to label a

novel kind of vesicles referred to as LC3-associated phagocytosis (LAP) vesicles, which are single-

membrane vesicles [5]. LAP vesicles were reported to be the third kind of intracellular vesicles that fuse

to phagosomes and facilitate the degradation of apoptotic cells in mice [6-8].

Autophagy is an evolutionarily conserved cellular event that plays an essential role in

maintaining cellular homeostasis by enveloping harmful protein aggregates and damaged cellular

organelles in double-membrane autophagosomes and subsequently degrading them via fusion with

lysosomes [9]. Autophagy also supports cell survival during nutrient starvation by capturing

intracellular organelles into autophagosomes and converting them to nutrients and energy sources [9].

Autophagosome formation requires the organized action of a set of proteins known as autophagic

related (ATG) proteins. It is a process of three sequential steps: initiation, nucleation, and expansion,

until an autophagosome fully forms and closes [10]. After formation, autophagosomes undergo a

maturation process through fusion with lysosomes, which provide digestive enzymes to degrade

autophagosomal contents, generating a fusion product referred to as "autolysosomes" [10]. In the nematode *C. elegans*, several autophagy genes have been reported to facilitate the clearance of apoptotic cells [11-13]. However, it is unknown whether autophagosome, as a particular type of cellular organelle, is involved in the clearance of apoptotic cells or whether these *atg* genes have additional functions, such as forming LAP vesicles.

Although both are labeled with LC3, mammalian LAP vesicles and autophagosomes are different in several key aspects. First, LAP vesicles are single-membrane vesicles, whereas autophagosomes are double-membrane vesicles [5,10]. In addition, although the formation of LAP vesicles relies on many autophagy genes, *ulk1*, *atg13*, and *atg14*, three genes whose products act in the initiation complexes for autophagosomes, are dispensable for the generation of LAP vesicles [5,7,14]. During the initiation of autophagosomes formation, which starts with the appearance of a membrane structure known as a phagophore, ULK1, a serine-threonine kinase, forms a protein complex with ATG13 and two other proteins and phosphorylates the class-III phosphoinositide 3-kinase (PI3K) VPS34 as well as the rest of the VPS34 complex (ATG6, ATG14, and VPS15), triggering the production of PtdIns(3)P on the phagosphore [10]. These distinct features of autophagosomes and LAP vesicles are critical for distinguishing whether an LC3-labeled vesicle is an autophagosome or a LAP vesicle.

During *C. elegans* embryonic development, 131 somatic cells undergo apoptosis and are swiftly engulfed and degraded by neighboring cells [15,16]. Apoptotic cells display a "button-like" structure under the Differential Contrast Interference (DIC) microscopy and are referred to as cell corpses [15,16]. Mutants defective in the clearance of cell corpses exhibit an increased number of persistent cell corpses, a phenotype known as **ce**ll **d**eath abnormal (Ced) [17]. In the *C. elegans* hermaphrodite gonad, 300-500 germ cells undergo apoptosis and are cleared by the neighboring gonadal sheath cells [18]. Previous genetic studies revealed two parallel, partially redundant pathways that primarily drive the clearance of *C. elegans* cell corpses. These include a signaling pathway led by CED-1, a phagocytic receptor for apoptotic cells, CED-6, an adaptor protein for CED-1, and DYN-1, a large GTPase playing many roles in membrane trafficking, and the other pathway led by the small Rac1 GTPase CED-10, and CED-5 and CED-12, the bipartite nucleotide exchange factor for CED-10 [19]. Unlike the CED-10 pathway, which primarily regulates cell corpse engulfment, the CED-1 pathway has two separate functions, regulating both the engulfment and degradation of cell corpses [20]. CED-1 on neighboring engulfing cells recognizes the "eat me" signal on the surfaces of cell corpses and is enriched to the region of the plasma membrane facing a cell corpse [21]. This enrichment initiates the extension of pseudopods along the

cell corpse and the subsequent closure of the phagocytic cup to form a nascent phagosome [22,23].

Moreover, CED-1 plays a distinct role in initiating the degradation of phagosomal contents [20]. CED-1

remains transiently enriched on the surface of nascent phagosomes, where it facilitates the CED-6 and

DYN-1-mediated sequential recruitment of the Class II phosphatidylinositol 3-bisphosphate (PtdIns(3)P)

kinase PIKI-1, the Class III PtdIns(3)P kinase VPS-34, and the small GTPases RAB-5 and RAB-7 to

phagosomal surfaces [20,24]. The robustly produced PtdIns(3)P and the RAB-5 and RAB-7 GTPases

further recruit effectors for PtdIns(3)P and the RAB proteins, respectively, which drive the recruitment

and fusion of early endosomes and lysosomes to a phagosome, leading to the degradation of the cell

corpse inside [24-26]. RAB-7, in particular, is essential for the fusion of lysosomes to phagosomes [20].

Autophagy and phagocytosis are two distinct lysosomal-mediated cellular degradation pathways

designated to eliminate intracellular and extracellular components, respectively. Previously, whether

canonical autophagosomes were involved in the degradation of phagosomal contents was unknown. We

report here that, in *C. elegans*, during the maturation of phagosomes that contain apoptotic cells, LC3-

labeled, double-membrane canonical autophagosomes are recruited to phagosomal surfaces and

subsequently fuse to these phagosomes. We have further discovered that this event facilitates the

degradation of apoptotic cells and is driven by the signaling pathway led by CED-1. This

autophagosome-phagosome fusion represents a novel mechanism that contributes to the degradation of

phagosomal contents and reveals a new function of autophagosomes.

Results

Vesicles labeled with GFP-tagged LC3 are recruited to the surfaces of phagosomes

The ATG8 protein family is composed of two subfamilies, the LC3 subfamily and the

GABARAP subfamily, which are very close to each other in sequence; the ATG8 family is also referred

to as the LC3/GABARAP family [4]. *C. elegans* has two LC3/GABARAP family members, LGG-1 and

LGG-2, which belong to the LC3 and GABARAP subfamilies, respectively (**Fig 1A**) [27]. Both LGG-1

and LGG-2 are attached to autophagosomes, except they each label autophagosomes of different

maturity [27].

To determine whether autophagosomes interact with phagosomes that contain apoptotic cells in

C. elegans embryos, we constructed the GFP-tagged LGG-1 and LGG-2 reporters that were expressed

under the control of the *ced-1* promoter (P*ced-1*), a well-documented engulfing cell-specific promoter

[21,28]. In embryos, we observed numerous GFP::LGG-1[+] and GFP::LGG-2[+] puncta (**Fig 1 D,G**).

Using our previously established time-lapse recording protocol [28], we observed the enrichment of

GFP::LGG-1[+] and GFP-LGG-2[+] puncta to the surfaces of the phagosomes, including the phagosomes

containing apoptotic cells C1, C2, and C3 (**Fig 1 B, D, G**). C1, C2, and C3 are localized to the ventral

surface of an embryo and undergo apoptosis at approximately 330-min post-1[st] cleavage [28]. Apoptotic

C1, C2, and C3 are each engulfed and degraded by a particular ventral hypodermal cell (**Fig 1 B**).

Inside the engulfing cell that expresses GFP::LGG-1 in the cytoplasm, the phagosomes containing C1,

C2, and C3 (also referred to as phagosomes C1, C2, and C3 for convenience) appear like dark discs (**Fig

1 D, G**). The GFP-labeled puncta were not observed inside the phagosomal lumen (**Fig 1 D**, **G**, top

panels).

C. elegans LGG-1 and LGG-2 were both reported to specifically attach to autophagosomes

through their lipid tails [27,29]. To verify that the observed GFP puncta are LGG-labeled lipid vesicles

and not artifacts of protein aggregation, we tested two mutant constructs, GFP::LGG-1(G116A) and

GFP::LGG-2(G130A), which bear mutations in the lipidation sites of the LGG proteins and are deficient

for membrane targeting [27]. We found that both GFP::LGG-1(G116A) and GFP::LGG-2(G130A)

display a diffuse cytosolic localization pattern (**Fig 1 D**, **G**, bottom panels), in stark contrast to the

punctate pattern presented by GFP::LGG-1 and GFP::LGG-2. This result indicates that GFP::LGG-1

and ::LGG-2 are membrane attached and thus label lipid vesicles.

The LGG-tagged puncta that fuse to phagosomes are double-membrane vesicles

The fluorophore within GFP is sensitive to acidic pH (pKa=6.0) [30], thus its signal diminishes

when GFP is inside the acidic lumen of lysosomes and phagosomes. To further monitor the fate of the

LGG[+] puncta after they are recruited to the surfaces of phagosomes, we replaced GFP with mCherry

(pKa<4.5) or mNeonGreen (mNG) (pKa=5.1), both of which are more resistant than GFP to the acidic

pH environment [31,32], allowing fluorescence signal inside the phagosomal lumen to be detected. Co-

expressed with the mCherry::LGG or mNG::LGG reporter are the CED-1::GFP or PH(hPLCγ)::mRFP

reporters, which were used in the time-lapse experiments as markers for the extending pseudopods,

allowing us to determine the moment when the pseudopods sealed up and a nascent phagosome was

born (**Fig 1 E**, **F**, **H**, **I,** top panels) [21,23]. In time-lapse image series of the clearance process of C1,

C2, and C3, we found that like the GFP::LGG reporters, the mCherry::LGG and mNG::LGG reporters

were enriched on the surfaces of phagosomes; moreover, unlike the GFP::LGG reporters, the mCherry

and mNG tagged reporters subsequently entered phagosomal lumen (**Fig 1 E**, **F**, **H**, **I**). The fluorescence

signal intensity increases over time with the continuous recruitment of the LGG[+] puncta on phagosomal

surfaces (**Fig 1 E**, **F**, **H**, **I, Figure 1-videos 1, 2, 3**). We measured the intensity of the mCherry::LGG-1

and mCherry::LGG-2 signal in the center of the C3 phagosomes over time from the start of the

phagosome (**Materials and Methods**) (**Fig 1C**) and observed over 10-fold increases of the signal

intensity within 60 min (**Fig 1J**). The appearance of the membrane-attached mCherry::LGG and

mNG::LGG signal in the phagosomal lumen indicates that the LGG$^+$ vesicles fuse with the phagosomal

membrane (**Fig 1K**). They further suggest that these vesicles are composed of double membranes

labeled with the LGG reporter molecules on both the outer and inner membranes (**Fig 1K**). If these

LGG$^+$ vesicles were single-membrane, no membrane-attached LGG reporter would end up inside the

phagosomal lumen because, as a result of fusion, the reporter molecules will be retained on the

phagosomal membrane (**Fig 1K**). Indeed, when CTNS-1, a lysosomal transmembrane protein, is tagged

with mRFP, which is acid-resistant (pKa=4.5 [31]), on its C-terminus, the lysosome-phagosome fusion

event resulted in the incorporation of the mRFP signal into the phagosomal membrane; as a result, the

CTNS-1::mRFP signal is only observed on the phagosomal surface, not in the phagosomal lumen (**Fig 1**

– figure supplement 1) [20].

174
Figure 1 with 2 supplements . The vesicles labeled with LGG-1 or LGG-2 are recruited to the surface of phagosomes
and subsequently fuse to phagosomes.

(A) Domain structures of mammalian LC3 and *C. elegans* LGG-1 and LGG-2. The green box indicates the conserved
ubiquitin-like domain. Residues in white are conserved among the three proteins. The glycine residue in red is the site where
the lipid tail is attached to.
(B) A diagram illustrating the three phagosomes that contain cell corpses C1, C2, and C3, with which we monitor the dynamic
recruitment and fusion of autophagosomes, at ~330 min post-the 1st embryonic division. Both the positions of C1, C2, and C3
(brown dots) and the identities of their engulfing cells are shown.
(C) A diagram illustrating that the relative mNG signal in the center of a phagosome is measured over time to create sub-figure
(J). At time point T (time after engulfment), the Relative signal intensity $_T$ = (Unit Intensity(phagosome center)$_T$ – Unit Intensity
(background)$_T$) / (Unit Intensity(phagosome center)$_{T0}$ – Unit Intensity(background)$_{T0}$). (Figure 1-source data 1)
(D-I) Time-lapse images of indicated reporters starting when a nascent phagosome (white arrowheads) just formed (time point
"0 min"). All reporters were expressed under the control of P$_{ced-1}$. Scale bars are 2 μm. Solid white arrowheads label nascent
phagosomes. Yellow arrows mark a few LGG-labeled puncta on the surface of phagosomes. Open white arrows in (E, F, H, I)
label the phagosomal lumen where the LGG signal is observed. "*" is the time point when the LGG signal is first seen inside
the phagosomal lumen. CED-1::GFP (E, H) and PH(PLCγ)::mRFP (F, I) are co-expressed markers that label the surfaces of
nascent phagosomes.
(D) GFP::LGG-1-labeled puncta are observed on the surface of a C3 phagosome, but the GFP signal is not seen inside the
phagosomal lumen. No GFP::LGG-1(G116A)-labeled puncta are seen on the surface of phagosomes.
(E-F) The mCherry::LGG-1 (E) and mNG::LGG-1 (F) puncta are observed on the surface of a C3 phagosome and
subsequently accumulate inside the phagosome lumen.
(G) GFP::LGG-2-labeled puncta are observed to attach on the surface of a C2 phagosome, but the GFP signal does not enter
the phagosomal lumen, whereas no GFP::LGG-1(G130A)-labeled puncta are seen on the surface of phagosomes.
(H-I) The mCherry::LGG-2 (H) and mNG::LGG-2 (I) puncta are observed on the surface of a C2 (H) or C1 (I) phagosome,
respectively, and subsequently accumulate inside the phagosome lumen.
(J) The relative mCherry::LGG-1 or -2 signal intensity in the center of a phagosome (Y-axis) over time (in the 2-min interval)
(X-axis). "0 min" indicates the moment when a nascent phagosome just formed. One blue horizontal line indicates value "1",
where no signal enrichment above background level is observed.
(K) A diagram illustrating that those double membrane-vesicles labeled with mCherry::LGG on their outer and inner
membranes are recruited to phagosomal surfaces and fused to the phagosomal membrane. After the fusion between the
outer membrane of these vesicles and the phagosomal membrane, the mCherry::LGG-tagged inner membrane is released
into the phagosomal lumen. The continuing incorporation of these double-membrane vesicles to phagosomes increases the
mCherry signal level in the phagosomal lumen over time. If the LGG-1 or LGG-2-labeled vesicles are of a single membrane,
no fluorescence signal is expected to enter the phagosomal lumen.
209

The LGG$^+$ vesicles that are incorporated into phagosomes are canonical autophagosomes

The observation that the LGG$^+$ vesicles incorporated into phagosomes are likely double-

membrane vesicles reminded us of canonical autophagosomes rather than LAP vesicles. In support of

this model, previously, using electron microscopy and immune-gold staining, Segalen *et al.* have shown

that in *C. elegans* embryos, LGG-1 or LGG-2 labeled vesicles are double-membrane autophagosomes

[27]. To further determine whether these vesicles represent canonical autophagosomes, we examined

whether loss-of-function mutations of *atg-7*, *atg-13*, and *epg-8*, which are defective for the biogenesis of

autophagosomes [33,34], impaired the production of these vesicles. *C. elegans atg-13* and *epg-8* encode

homologs of mammalian ATG13 and ATG14, respectively, which are essential for the biogenesis of

autophagosomes but dispensable for that of LAP vesicles (**Introduction**) [35]. *atg-7* encodes a

homolog of mammalian ATG7, a protein essential for conjugating a phospholipid tail onto the LC3

family proteins and thus for the biogenesis of both autophagosomes and LAP vesicles [10,35]. We first

scored whether the mCherry::LGG-1/-2 reporters were observed in the center of 15 phagosomes during

the phagosome maturation process (**Figs 1C** and **2**), which is indicative of the fusion of double-

membrane mCherry::LGG $^+$ vesicles to phagosomes. In wild-type embryos, the steady entry of mCherry

into phagosomes over time (**Fig 1 E**, **H**) results in the increase of the average mCherry::LGG-1 and -
LGG-2 intensities to 9.2- and 8.4-fold of that at 0 min time point, respectively, at 60 min after the
formation of a phagosome (**Fig 2 H**, **J**). However, in *atg-7(bp411)* [36] mutant embryos, hardly any
mCherry signal was observed inside phagosomes **(Fig 2 A, D, G-J)**. At 60 min after the phagosome
formation, the average mCherry signal intensities were merely 1.2 and 1.4-fold at 0 min time point **(Fig
2 H, J).** These observations indicate a lack of LGG$^+$ vesicles that fuse to phagosomes. In *atg-13(bp414)*
and *epg-8(bp251)* mutant embryos, similar observations were made except that the defects were slightly
weaker (**Fig 2 B, C, E, F, G-J**).

We next examined whether the mCherry::LGG-1/-2$^+$ vesicles are produced in cells of the *atg-7*, *atg-9*,
atg-13, and *epg-8* mutant embryos. In wild-type embryos at mid-embryonic developmental stages
(~330, ~350, ~400 min post-1st embryonic cell division), numerous mCherry$^+$ puncta were observed
(**Fig 2 – figure supplements 1A** and **2A**). In the *atg-7* mutant embryos, rarely any such puncta existed
(**Fig 2 – figure supplements 1B** and **2B**), consistent with a previous report [33]. *atg-9* encodes ATG-9,
the only transmembrane protein in the core autophagy machinery [33,35]. ATG9 plays an essential role
in the expansion of phagophore and the biogenesis of autophagosomes [10]. In *atg-9(bp564)* mutant
embryos that expressed mNG::LGG-1 or ::LGG-2, much fewer mNG$^+$ puncta that might represent
autophagosomes were observed, and the mNG reporters label large aggregates (**Fig 3 – figure
supplement 1**), consistent with a previous report [37]. In *atg-13* and *epg-8* mutant embryos, the
numbers of mCherry-LGG$^+$ puncta were also significantly reduced (**Fig 2 – figure supplements 1 C-D
and 2 C-D**), suggesting that the *atg-13(bp414)* and *epn-8(bp251)* mutations severely impaired the
biogenesis of autophagosomes, a phenotype that is consistent with previous reports [33,34]. The drastic
reduction in the number of LGG-1/-2-labeled puncta in *atg-7*, *atg-9, atg-13,* and *epg-8* loss-of-function
mutants strongly indicates that these puncta belong to canonical autophagosomes.

248
Figure 2 with 2 supplements. In autophagosome-formation mutants, the enrichment of the LGG⁺ vesicles on the
phagosomal surface and the entry of the LGG signal into the phagosomal lumen are severely defective.
(A-F) Time-lapse image series monitoring the enrichment of the puncta labeled with mCherry-tagged LGG-1 (A-C) or LGG-2
(D-F) on phagosomes (white arrowheads) and the subsequent entry of mCherry signal into the phagosomal lumen in *atg-7*,
atg-13, and *epg-8* mutant embryos. "0 min" is when a phagosome is just sealed (determined by CED-1::GFP). Open white
arrow denotes the time point that the mCherry signal starts to appear inside the phagosomal lumen. Scale bars are 2µm.
Yellow arrows in (F) mark mCherry::LGG-2 puncta on the surface of a phagosome.
(G and I) The relative mCherry::LGG-1 or -2 signal intensity in the center of a phagosome (Y-axis) over time (in the 2-min
interval) (X-axis). "0 min" indicates the moment when a phagosome is just sealed. (G) The data for the wild-type, *atg-7(bp411)*,
atg-13(bp414), and *epg-8(bp251)* mutant embryos are from Figures 1E and 2(A-C), respectively. (I) The data for the wild-type,
atg-7(bp411), *atg-13(bp414)*, and *epg-8(bp251)* mutant embryos are from Figures 1I and 2(D-F), respectively. (Figure 2-
source data 1)
(H and J) Box-and-Whiskers plots of the relative mCherry signal intensity measured in the center of phagosomes 60 min-post
the formation of nascent C3 phagosomes from 15 each of wild-type, *atg-7(bp411)*, *atg-13(bp414)*, and *epg-8(bp251)* mutant
embryos. Red dashed lines indicate the position of value 1, which represents no signal enrichment relative to the background
signal. (Figure 2-source data 2)
***, p<0.001, **, <0.001p<0.01, Student *t*-test of each mutant comparing to the wild-type value.
266

To further confirm that autophagosomes fuse to phagosomes, we examined the subcellular

localization of a mCherry::ATG-9 reporter expressed in engulfing cells. Besides the LGG proteins,

ATG-9 is also an established autophagosomes marker as it is the only transmembrane protein in

autophagosomes [38]. In embryonic hypodermal cells that co-expressed mCherry::ATG-9 and either

mNG::LGG-1 or::LGG-2, puncta labeled with mCherry are recruited to the surfaces of phagosomes (**Fig**

3). Furthermore, the mCherry signal gradually accumulates in the phagosomal lumen like the LGG

reporters (**Fig 3**). During the phagosome maturation process, the co-localization between

mCherry::ATG-9 and each of the two LGG reporters on the puncta on phagosomal surfaces and inside

phagosomal lumen was nearly perfect (**Fig 3**). Together, the above observations verified that the ATG-9

and LGG double-positive autophagosomes are recruited to phagosomal surfaces and subsequently fuse

to phagosomes containing apoptotic cells.

Figure 3 with 1 supplement. LGG-1⁺ and LGG-2⁺ puncta colocalize with ATG-9, a component of autophagosomes.
The mNG- and mCherry- tagged reporters are expressed in wild-type embryos under the control of P*ced-1*. White arrowheads
mark nascent phagosomes. Small white arrows mark the regions where LGG⁺ and ATG-9⁺ puncta colocalize. Open white
arrows indicate when the fluorescent signal is first detected inside the phagosomal lumen. Scale bars are 5 µm.
(A) Time-lapse microscopy showing the localization of mNG::LGG-1 and mCherry::ATG-9. Images are from ABplaapppp,
which engulfs C3.
(B) Time-lapse microscopy showing the localization of mNG::LGG-2 and mCherry::ATG-9. Images are from ABplaapppa,
which engulfs C1.
287

The autophagosome-phagosome interaction is a general phenomenon observed in embryos and the

adult gonad

Besides the C1, C2, and C3 phagosomes, in embryos co-expressing mCherry::LGG-1 or -2 and

CED-1::GFP reporters, we also observed the recruitment of the mCherry signal to other phagosomes and

the subsequent entry of the mCherry signal to the phagosomal lumen (**Fig 1 – figure supplement 2 A-**

E). Time-lapse image series (**Fig 1 – figure supplement 2 A-E**) demonstrated the recruitment and

fusion of LGG⁺ puncta to apoptotic cells C4 and C5 in the tail [23]. In mid-stage (1.5-fold) embryos,

multiple phagosomes are observed to either have LGG$^+$ puncta attached to the surface (**Fig 1 – figure**

supplement 2 F(b, d), yellow arrows) or LGG$^+$ signal inside the lumen (**Fig 1 – figure supplement 2**

F(b,d), white open arrows). None of these phagosomes are C1, C2, C3, C4, or C5, judging by their

positions in the embryo. These observations indicate that the fusion between autophagosomes and

phagosomes is a general phenomenon during embryogenesis.

300   In each *C. elegans* adult hermaphrodite gonad, germ cells that die of apoptosis undergo

cellularization and are engulfed by surrounding gonadal sheath cells [18]. We used transmission

electron microscopy (TEM) (**Materials and Methods**) to examine whether there were any double-

membrane vesicles attaching to phagosomes containing germ cells. In wild-type adults, due to the swift

engulfment and degradation activities, on average only 2-4 germ cell corpses are observed in each

gonadal arm in an adult hermaphrodite 48-hrs post mid-L4 larval stage [20], making it very difficult to

find any phagosomes in the 50-nm thin sections in TEM. We thus chose to analyze phagosomal

surfaces in the *rab-7* mutants, in which many germ cell corpses persist in the gonad due to the blockage

of phagosome degradation [20]. In TEM images, germ cell corpses are identified by their higher density

than live germ cells and complete cellularization [20]. The *rab-7(ok511)* mutants are maternal-effect

embryonic lethal. *rab-7(ok511)* adult hermaphrodites are viable and produce dead embryos [20]. In

rab-7(ok511) mutant adult gonads, many undegraded phagosomes containing germ cell corpses are

observed inside the sheath cell in thin (50 nm) TEM sections (**Fig 4 B, C**), as reported previously [20].

Furthermore, in the *rab-7* mutant gonad, mCherry::LGG-1 expressed in sheath cells (under P$_{ced-1}$) is

observed to accumulate on the surfaces of phagosomes (**Fig 4 A**), like in *rab-7* mutant embryos (see

Results section "The small GTPase RAB-7 and the HOPS complex are essential for the fusion between

autophagosomes and phagosomes" below). This result is consistent with the observation of LGG$^+$

puncta on phagosomal surfaces in embryos. In *rab-7* mutant gonad, we did identify multiple double-

membrane vesicles in close contact with the phagosomal surfaces in TEM thin sections (**Fig 4 D-H**).

Besides being composed of double membranes, these vesicles also resemble autophagosomes identified

in C. elegans cells [39,40]. These vesicles vary in diameter from 200 nm to 800 nm, consistent with that

of autophagosomes reported in the literature [39,40]. The above observations support our conclusion

that the LGG$^+$ puncta that are recruited to phagosomal surfaces are indeed canonical autophagosomes.

They further suggest that the autophagosome-phagosome interaction is general rather than a cell-specific

phenomenon.

Figure 4. Double-membrane vesicles were observed to attach on the surfaces of phagosomes bearing germ cell corpses

All samples are from the distal gonadal arms of *rab-7(ok511)* adult hermaphrodites.

(A) In the gonadal arm of a *rab-7(ok511)* hermaphrodite expressing P*ced-1 mCherry::lgg-1* in gonadal sheath cells, mCherry[+] puncta (arrows) are found on the surfaces of phagosomes carrying germ cell corpses (white (a) and black (b) asterisks). Scale bars are 10 μm.

(B) A thin cross-section (50 nm in thickness) TEM image of half of a distal gonad. The scale bar is 1 μm.

(C) Traces of membranes corresponding to (B). All three germ cell corpses are inside the gonadal sheath cell. Two asterisks mark the remnants of two engulfed germ cell corpses.

(D-H) Scale bars are 500 μm. Examples of five double-membrane vesicles (yellow asterisks) were observed on phagosomal surfaces. White arrows mark each layer of the double-layer membranes of the vesicles of interest. Open arrows mark the phagosomal (white) and germ cell corpse (red) membranes. (E) is an enlarged image of the region framed by the red box in (D). (G) The luminal content of the double-membrane vesicle (*) is missing due to the damage in sample preparation. Due to the same damage, the phagosomal membrane in (G) is unclear.

Autophagosomes facilitate the degradation of apoptotic cells inside phagosomes

To examine whether the incorporation of autophagosomes into phagosomes affects the clearance

of the engulfed apoptotic cells, we first quantified whether, in mutants of genes essential for the

biogenesis of autophagosomes, apoptotic cells were un-degraded and thus persisted in embryos. In

addition to the *atg-7*, *atg-9*, *atg-13*, and *epg-8* mutants characterized above, we also characterized loss-

of-function mutants of *lgg-1* and *lgg-2*, and of *atg-3*, whose gene product is essential for the conjugation

of a phosphatidylethanolamine (PE) tail to the LC3 family proteins [33,41], of *atg-2* and *atg-18*, whose

gene products function together with ATG-9 in the expansion of phagophore [38], and of *unc-51*, which

encodes a *C. elegans* homolog of ULK1, an autophagic protein kinase [38]. In 2-fold stage wild-type

embryos, which are ~460 min post the first embryonic cell division, an average of 11.3 cell corpses were

scored (**Fig 5 A**). The mutant 2-fold embryos examined bore 46.9% - 91.2% more cell corpses (**Fig 5**),

indicating that the clearance of cell corpses is defective. Together, the findings reported in **Figs 2** to **5**

indicate that autophagosomes made a substantial contribution to the clearance of cell corpses.

Figure 5 with 1 supplement. Mutations in autophagy genes impair the clearance of apoptotic cells.
(A-B) Bar graph displaying the average numbers of somatic cell corpses in 2-fold stage wild-type and various mutant embryos.
Bars and error bars represent mean and standard deviation (sd), respectively, the actual values of which are displayed inside
the bars. Numbers in parentheses indicate the number of embryos scored. ***, p<0.001, Student *t*-test of each mutant
comparing to the wild-type value. (Figure 5-source data 1)
(C) DIC images of cell corpses in 2-fold stage embryos of various genotypes. White arrows indicate button-like structures
characteristic of cell corpses. Scale bars are 5 μm.

To determine whether the lack of autophagosomes impairs the engulfment or degradation of cell

corpses, we monitored the formation and degradation of phagosomes containing apoptotic cells C1, C2,

and C3 (**Fig 1 B**) in wild-type, *atg-7*, and *lgg* mutant embryos in real-time using established protocol

(**Materials and Methods**) [25,28]. The CED-1::GFP expressed in engulfing cells labels the extending

pseudopods and enables us to monitor the process of phagosome formation, starting from the budding

and ending at the sealing of the pseudopods [21,22]. 2xFYVE::mRFP (also expressed under P$_{ced-1}$), a

reporter for phagosomal surface PtdIns(3)P, enables us to monitor the shrinking of a phagosome, an

indication of phagosome degradation [20,24,25]. In *atg-7*, *lgg-1,* and *lgg-2* mutant strains that co-

expressed CED-1::GFP and 2xFYVE::mRFP, we found that engulfment was completed in 4-8 min, just

like in wild-type embryos (**Fig 6 A-D, F**), indicating that defects in autophagosomes biogenesis do not

affect the engulfment of cell corpses. However, the lifespans of phagosomes (**Materials & Methods**)

were much longer in *atg-7*, *lgg-1*, and *lgg-2* mutants than in wild-type embryos (**Fig 6 A-D, G**). All

wild-type phagosomes have a lifespan between 40-60 min (**Fig 6 G**). In *atg-7*, *lgg-1,* and *lgg-2* mutants,

the lifespan varied in a much more extensive range. Remarkably, 43.8%, 82.4%, and 53.3% of

phagosomes in *atg-7*, *lgg-1,* and *lgg-2* mutant embryos, respectively, have lifespans longer than 60 min

(**Fig 6 G**). These observations indicate that autophagosomes made a specific contribution to the

efficient degradation of phagosomal contents.

Figure 6. Mutations in *rab-7, atg-7, lgg-1,* and *lgg-2* impair the degradation of cell corpses to different degrees.
(A-E) Time-lapse recording conducted in wild-type and different mutant embryos monitoring the dynamics of the pseudopod marker CED-1::GFP and the phagosome marker 2xFYVE::mRFP (both expressed in engulfing cells) during the engulfment and degradation processes of cell corpse C3 by ABplaapppp. "0 min" is the first time point when a nascent phagosome (white arrowheads) is formed, as indicated by the closure of a green GFP::CED-1 ring. 2xFYVE::mRFP labels the surface of a phagosome (yellow arrows) until it is degraded. Scale bars are 2 µm.
(F) Histogram depicting the distribution of the time it takes to engulf 15 C3 cell corpses in wild-type, *atg-7*, *lgg-1*, and *lgg-2* and 7 cell corpses in *rab-7*(m⁻z⁻) homozygous embryos. The engulfment time is defined as the period between the first time when pseudopods (labeled with CED-1::GFP) are observed and when a full circle is observed forms around C3. (Figure 6-source data 1)
(G) Histogram depicting the lifespan distribution of 15 C3 phagosomes in wild-type, *atg-7*,*lgg-1*, *lgg-2,* and 7 phagosomes in *rab-7*(m⁻z⁻) homozygous embryos. Phagosome lifespan is measured as the time interval between the "0 min" time point when a nascent phagosome is just sealed and when the phagosome shrinks to one-half of its original diameter. (Figure 6-source data 2)

LGG-1 and LGG-2 act in engulfing cells, and together they define three subpopulations of autophagosomes that are incorporated into phagosomes

LGG-1 and LGG-2 have distinct structural features and were observed to attach to different sub-populations of autophagosomes, which represent autophagosomes at different maturation stages [27,29]. The sub-populations of autophagosomes labeled by LGG-1 or LGG-2 are incorporated into phagosomes (**Fig 1**). *lgg-1* and *lgg-2* single mutants are both inefficient in clearing cell corpses (**Fig 5B**). Furthermore, in *lgg-1; lgg-2* double mutant embryos at 2-fold, the Ced phenotype is further enhanced significantly (**Fig 5B**) -- the number of cell corpses is 22.0% and 27.1% larger than in the *lgg-1* and *lgg-2* single mutants, respectively. This enhanced phenotype indicates an additive effect and suggests that the autophagosomes labeled with LGG-1 or LGG-2 play parallel and partially redundant roles in promoting phagosome degradation. In embryos co-expressing GFP::LGG-1 and mCherry::LGG-2, we observed that both in the cytoplasm of the engulfing cells and on the surfaces of phagosomes, puncta that were either labeled with GFP or mCherry alone or with both GFP and mCherry (**Fig 7 A, B**). In hypodermal cells, the average distribution of LGG-1$^+$ LGG-2$^-$, LGG-1$^-$ LGG-2$^+$, and LGG-1$^+$ LGG-2$^+$ puncta is 35.8%, 40.7%, and 23.5%, respectively (**Fig 7C**). These observations indicate that in addition to the LGG-1$^+$-only and LGG-2$^+$-only sub-populations, a third, LGG-1$^+$ LGG-2$^+$ double-positive sub-population of autophagosomes exists. This sub-population likely corresponds to an intermediate stage in the maturation path of autophagosomes.

To determine whether LGG-1 and LGG-2 act in engulfing cells to facilitate phagosome degradation, we examined whether the specific expression of each gene in engulfing cells, under the control of P$_{ced-1}$, would rescue the Ced phenotype of the corresponding mutant embryos. We tested the *lgg* cDNAs that are tagged with either *gfp* or *mCherry* for the rescuing activity by counting the number of cell corpses in 1.5-fold stage transgenic embryos. In the *lgg-1* and *lgg-2* null mutants, both the *gfp*- and *mCherry*-tagged corresponding *lgg* cDNA efficiently rescued the Ced phenotype (**Fig 5 – figure supplement 1**). The *gfp::lgg-1* and *mCherry::lgg-1* transgenes lowered the number of cell corpses from on-average 152% of wild-type level observed in the *lgg-1(tm3489)* mutants to 118% and 123% of the wild-type level, respectively. Similarly, the *gfp::lgg-2* and *mCherry::lgg-2* transgenes lowered the number of cell corpses from on-average 143% of wild-type level observed in the *lgg-2(tm5755)* mutants to 109% and 112% of the wild-type level, respectively (**Fig 5 – figure supplement 1**). These results indicate that *lgg-1* and *lgg-2* primarily act in engulfing cells to facilitate the clearance of apoptotic cells.

Figure 7. The puncta labeled with LGG-1 and/or LGG-2 define three distinct populations of vesicles
The GFP- and mCherry- tagged reporters are expressed in wild-type embryos under the control of P*ced-1*.
 (A) Time-lapse image series of a C2 phagosome (big white arrowhead) monitoring the localization of LGG-1[+], LGG-2[+,] and
LGG-1[+] - LGG-2[+] double-positive puncta on the surface of the phagosome. "0 min" is the moment when a nascent phagosome
just seals. The scale bar is 2 µm. White arrows mark GFP[+] mCherry[-] puncta, white open arrows mark GFP[-] mCherry[+] puncta ,
and yellow arrows mark GFP[+] mCherry[+] puncta.
(B) DIC and fluorescence images of an embryo exhibiting LGG[+] puncta outside phagosomes in multiple cells. The scale bar is
10 µm. The white arrows, white open arrows, and yellow arrows mark GFP[+] mCherry[-], GFP[-] mCherry[+], and GFP[+] mCherry[+]
puncta on the surface of the phagosome, respectively. The big white arrowheads in (c) and (d) indicate the mCherry signal
internalization to the phagosomal lumen.
(C) Bar graph indicating the distribution of single and double-labeled puncta in the LGG-labeled population in the engulfing
cells for C1, C2, and C3, scored immediately prior to the point when the LGG signal was observed inside the phagosomal
lumen. Eight engulfing cells were scored. Bars represent the mean, the error bars indicate standard deviation, and each dot
represents one sample. (Figure 7-source data 1)

The small GTPase RAB-7 is enriched on the surfaces of autophagosomes

The small GTPase Rab7 is well known to specifically label late endosomes and lysosomes [42].

In addition, Rab7 proteins in mammalian and *C. elegans* are also recruited from the cytoplasm to the

phagosomal membrane shortly after the formation of a phagosome and mediate the fusion between the

maturing phagosome and lysosomes and do that through their effector, the HOPs complex [3,26].

Furthermore, in yeast, *Drosophila*, and mammalian cells, Rab7 is directly recruited to the surfaces of

autophagosomes and plays an important role in the fusion between autophagosomes and lysosomes [43-

46]. *C. elegans* RAB-7 also plays an essential role in the fusion between autophagosomes and

lysosomes [27]. To examine whether RAB-7 is localized to LGG-1[+] and LGG-2[+] autophagosomes that

fuse to *C. elegans* phagosomes, we generated two transgenic *C. elegans* strains that co-expressed the

mCherry::LGG-1/ GFP::RAB-7 or mCherry::LGG-2/ GFP::RAB-7 pairs of reporters (**Materials and**

Methods). We observed that GFP::RAB-7 was localized to some but not all of the LGG-1[+] or LGG-2[+]

puncta (**Fig 8 B-D**). The LGG$^+$ RAB-7$^+$ double-positive autophagosomes were observed both freely

distributed in the cytoplasm of ventral hypodermal cells (**Fig 8B(a-c)** and **C**) and on phagosomal

surfaces (**Fig 8B(d-f)**, **D**). In addition, **Fig 8B(d-f)** depicts that GFP::RAB-7 is both evenly distributed

to the surface of a phagosome (d) as previously reported [20], and highly enriched on LGG-1$^+$

autophagosomes that are recruited to the phagosomal surface (**d-f**, white arrows). **Fig 8D** shows a time-

lapse series of a dynamic fusion event of an LGG-2$^+$/RAB-7$^+$ punctum (marked by the bottom arrow in

the "0 min" time point) to the phagosome: this punctum is first seen attached to the phagosome

membrane and subsequently becoming part of the phagosomal surface at the "+6 min" time point.

Quantitative analysis of the green and red puncta distribution reveals that on average, 66.2 % and 63.5 %

of LGG-1$^+$ and LGG-2$^+$ puncta observed in the cytoplasm of the engulfing cells are RAB-7$^+$,

respectively (**Fig 8 E**). The LGG$^+$ but RAB-7$^-$ puncta (**Fig 8 B, C**, white arrowheads) are likely

immature autophagosomes not acquired RAB-7 yet. We also observed puncta that were LGG$^-$ but RAB-

7$^+$ (Fig 8 **B, C**, yellow arrows). These puncta represent intracellular organelles such as late endosomes

or lysosomes. Different from the observation made in the cytoplasm, 100% of the autophagosomes that

were observed on phagosomal surfaces were RAB-7$^+$ (**Fig 8 B, D, F**). This observation suggests that

RAB-7 might play an essential role in the interaction between autophagosomes and phagosomes.

Figure 8. RAB-7 is enriched on a portion of autophagosomes, and RAB-7 and VPS-18 are essential for the fusion between autophagosomes and phagosomes.

(A) Diagram of the ventral surface of an embryo at ~330 min post 1st embryonic division.

(B-D) Images of part of the ventral surface of an embryo co-expressing P$_{ced-1}$gfp::rab-7 and P$_{ced-1}$mCherry::lgg-1 (B) or P$_{ced-1}$mCherry::lgg-2 (C-D). B(a-c) depicts the region framed by the blue box in (A). B(d-f) depicts a C1 phagosome (a yellow arrowhead). C(a-c) depicts the region framed by the red box in (A). (D) A time-lapse image series of a C1 phagosome (a yellow arrowhead) indicates the dynamic recruitment and fusion of GFP and mCherry double-positive puncta to the phagosomal membrane. "0 min" is when the first puncta are observed on the phagosomal surface. White arrows mark several puncta that are both GFP[+] and mCherry[+]. Yellow arrows mark puncta that are GFP[+] but mCherry[-]. White arrowheads mark puncta that are GFP[-] but mCherry[+]. Scale bars for B(a-c) and (C) are 5 μm, and for B(d-f) and (D) are 2 μm.

(E-F) Bar graphs depicting the percentage of LGG-1+ or LGG-2+ puncta that are also RAB-7+ in the cytoplasm of the engulfing
cells for C1, C2, and C3 (E) or on the surfaces of the phagosomes 2-min before the autophagosome-phagosome fusion
occurs (F). Nine engulfing cells and the phagosomes inside were scored for each of the LGG-1+ and LGG-2+ categories. Bars
and error bars represent the mean and standard deviation values. Each dot represents one sample. ns, not significant.
(Figure 8-source data 1)
(G-H) Time-lapse images monitoring the recruitment and fusion of puncta labeled with mCherry::LGG-1 (E) or::LGG-2 (F) to
the C1, C2, and C3 phagosomes in *rab-7(ok511)* and *vps-18(tm1125)* mutant embryos. "0 min" represents the moment when
a phagosome just seals (white arrowheads). Yellow arrows mark the mCherry+ puncta on the phagosomal surfaces. Open
white arrow marks the mCherry signal inside the phagosomal lumen. Scale bars are 2 µm.
(I-J) Histograms depicting the distribution of the time it takes for LGG-1+ or LGG-2+ puncta to fuse to phagosomes, measured
from the "0 min" point to the time when mCherry was detected in the center of a phagosome. C1, C2, and C3 phagosomes
were recorded. *n*, the number of phagosomes scored. "No fusion": no mCherry signal entry was observed even after 72-114
min post-nascent phagosome formation. NA: not applicable. (Figure 8-source data 2)
(K) A Diagram depicts the mechanism driving autophagosome-phagosome fusion. RAB-7 is enriched on the surfaces of both
phagosomes and autophagosomes. RAB-7 and VPS-18, a subunit of the HOPs complex, are proven essential for
autophagosome-phagosome fusion. Other factors are proposed to play roles in this event based on the general knowledge of
intracellular membrane fusion.
498

RAB-7 and the HOPS complex are essential for the fusion between autophagosomes and

phagosomes

We examined whether the recruitment and fusion of mCherry::LGG-1+- or LGG-2+-

autophagosomes to the C1, C2, and C3 phagosomes were normal in *rab-7(ok511)* null mutant embryos

[20]. The recruitment event can be evaluated by the level of enrichment of mCherry puncta on the

surfaces of phagosomes, whereas the subsequent fusion event can be measured by the level of

accumulation of the mCherry signal inside the phagosomal lumen. In *rab-7(ok511)*(m$^-$z$^-$) mutant

embryos produced by *rab-7(ok511)*(m$^+$z$^-$) mothers, robust enrichment of both the mCherry::LGG-1$^+$

and ::LGG-2$^+$ puncta to phagosomal surfaces are prominent (**Fig 8 G-H**), indicating that the recruitment

of autophagosomes was normal. However, no mCherry signal was observed entering the phagosomal

lumen (**Fig 8 G-H Figure 8-videos 1, 2**). We quantified the time between the moments a phagosome

was just born and that when an obvious mCherry signal was observed inside the phagosomal lumen. In

the wild-type embryos, this time is <30 min in 100% of the samples analyzed, and the median value is

14 and 18 min for LGG-1$^+$ and LGG-2$^+$ autophagosomes, respectively (**Figs 1**(**E** and **H**) and **8(I-J)**). In

stark contrast, in *rab-7* mutants, the mCherry signal was not detected in the lumen inside any of the

phagosomes for LGG-1 or LGG-2 reporters over a time span of 0-60 min after phagosome formation

(**Fig 8 I-J**). For over half of the samples, the observation period was extended beyond 90 min after

phagosome formation, and still, no mCherry signal was observed in the phagosomal lumen within this

period. These results strongly indicate that the function of RAB-7 is essential for autophagosome-

phagosome fusion but not required for the recruitment of autophagosomes to phagosomal surfaces.

The HOPS complex acts as an effector for Rab7 [47]. In *C. elegans*, a null mutation in *vps-18*,

which encodes a subunit of the HOPS complex, impairs phagosome maturation [48]. The lack of

autophagosome-phagosome fusion observed in *rab-7* mutants led us to subsequently examine the *vps-18*

null mutant embryos. We found that, like in *rab-7* mutants, the recruitment of autophagosomes to

phagosomes appeared normal in *vps-18* mutants (**Fig 8 G-H**). Also, like in *rab-7* mutants, in *vps-18*

mutants, the fusion of LGG-1$^+$ and LGG-2$^+$ autophagosomes to phagosomes was severely defective (**Fig**

8 G-J). However, whereas in *rab-7* mutants, the accumulation of the mCherry signal in the phagosomal

lumen was blocked completely in all samples, in some *vps-18* mutant embryos, the entry of mCherry

signal still occurred, *albeit* severely delayed (**Fig 8 I-J**). In contrast, in other samples, the entry was

blocked **(Fig 8 I-J)**, indicating a fusion defect that is less severe than that caused by the *rab-7* null

mutation and suggesting the existence of the residual HOPS function in *vps-18* mutants. Together, our

observations indicate that RAB-7 and the HOPS complex play a critical and specific role in driving the

fusion between autophagosomes and phagosomes.

To evaluate the defect in phagosome degradation caused by the *rab-7(ok511)* null mutation,

which abolishes the fusion of both lysosomes and autophagosomes to phagosomes, we measured the

phagosome lifespan in *rab-7(ok511)(m⁻z⁻)* embryos using the phagosome lifespan analysis used in **Fig 6**

A-D). Compared to that observed in the *atg-7*, *lgg-1*, and *lgg-2* mutant embryos, the phagosome

lifespan is significantly longer in *rab-7* mutants. In *rab-7* mutants, 100% of phagosomes last longer

than 60 min, among which 71.4% last longer than 80 min and 57% last longer than 90 min (**Fig 6 E, G**).

The difference in severity in the phagosome degradation defects displayed by *rab-7* and autophagy

mutants suggest that autophagosomes and lysosomes both contribute to phagosome degradation, and

that autophagosomes provide a phagosome degradation activity in an additive and possibly independent

manner to lysosomes.

542
The CED-1 pathway drives the recruitment of autophagosomes to phagosomes

The signaling pathway led by the phagocytic receptor CED-1 plays essential roles in initiating

the maturation of phagosomes containing apoptotic cells [20]. The CED-1 pathway is known to drive

the incorporation of early endosomes and lysosomes to phagosomes [20,22]. Here we further examined

whether the loss-of-function mutations in members of the CED-1 pathway affect the incorporation of

autophagosomes to phagosomes. CED-1 promotes the recognition, engulfment, and degradation of

apoptotic cells [20]. In *ced-1(e1735)* null mutant embryos, although the recognition and engulfment are

delayed or blocked due to the loss of CED-1's engulfment activity, the majority of the cell corpses are

eventually engulfed inside phagosomes [20]. This is why in *ced-1* mutants, there are phagosomes that

contain C1, C2, or C3 available for analysis of phagosome degradation. The partially penetrant

recognition and engulfment defects observed in the *ced-1* mutants are due to the compensation of the

activities by the *ced-5/-10/-12* and the *rab-35* pathways [1,22,49]. In *ced-1(e1735)* mutants expressing

the mNG::LGG-1 or -2 reporters, we observed severe defects in the incorporation of autophagosomes

into phagosomes. First of all, only a very dim mNG signal was observed inside the phagosomal lumen

50 min post phagosome formation, a time point well past the observed initiation time for

autophagosomes/phagosome fusion in the wild-type condition (**Fig 9 A**, **C**, **F**, and **G, Figure 9-video 1**).

Whereas in wild-type embryos, the median relative LGG-1 and -2 signal intensities are 6.3 and 5.3 at 50

min-post phagosome formation, respectively, in *ced-1* mutant embryos, the median values are merely

2.1 and 1.6, respectively (**Fig 9 K-L**). Secondly, unlike in *rab-7* mutants, where autophagosomes were

observed accumulating on phagosomal surfaces (**Fig 8 G-J**), in *ced-1* mutants, very few LGG-1- or

LGG-2-labeled puncta were observed on phagosomal surfaces (**Fig 9 A**, **C**).

Further quantitative measurement of the samples presented in (**Fig 9 A**, **C**) and 14 additional

samples for each of the reporters confirmed that the mNG signal was not enriched on the surfaces of

phagosomes, in contrast to wild-type embryos (**Fig 9 H-L**). These results indicate a severe defect in the

recruitment of autophagosomes to phagosomes. Unlike in the *atg* mutants that we have examined (**Fig**

2 – figure supplements 1-2), in *ced-1* mutant embryos, normal numbers of LGG-1$^+$ or LGG-2$^+$ puncta

were observed (**Fig 9 – figure supplement 1**), indicating that the biogenesis of autophagosomes is

normal. Thus the recruitment defect observed in *ced-1* mutants is not a consequence of the lack of

autophagosomes; rather, it is likely a result of a defect in signaling between phagosomes and

autophagosomes.

We further examined whether CED-6 and DYN-1, two other members of the CED-1 pathway,

were also needed for the incorporation of autophagosomes into phagosomes. In the *ced-6(n2095)* and

dyn-1(n4039) loss-of-function mutant embryos, the median relative LGG-1 and LGG-2 signal intensities

in the center of phagosomes are much lower than that in wild-type samples, respectively, at 50 min-post

phagosome formation mutants (**Figs 9(B, D, F, G, K-L)** and **10(A, D E, F, I, J)**), although the defects

are not as severe as in *ced-1* mutants (**Fig 9 A, C, F, G, K, L**). Further observation discovered that the

LGG-1 or LGG-2-labeled puncta were rarely observed on the surfaces of phagosomes in these mutants

(**Figs 9(B, D, I, J)** and **10(A, F)**), demonstrating severe defects in the recruitment of autophagosomes to

phagosomal surfaces.

In *ced-1*, *ced-6*, and *dyn-1* mutants, the fusion between autophagosomes and phagosomes might

also be defective. However, the severe recruitment defects resulted in the lack of LGG-labeled puncta

on phagosomal surfaces, making it difficult to evaluate whether there are additional fusion defects and

how severe the fusion defects are.

Figure 9 with 1 supplement. CED-1 and DYN-1 are essential for the incorporation of autophagosomes into phagosomes

(A-D) Time-lapse image series monitoring the presence or absence of puncta (white arrows) labeled with mNG::LGG-1 (A-B) or -2 (C-D) on C3 phagosomes (white arrowheads) and the subsequent entry of the mNG signal into the lumen in *ced-1* and *dyn-1* mutant embryos. DIC images mark the position of the cell corpse. "0 min" is the moment when phagosomes are just sealed. Scale bars are 2μm.

(E) A diagram illustrating that the relative mNG signal in the center of a phagosome is measured over time to create sub-figures (F) and (G). At time point t (time after "0 min"), the Relative Signal Intensity $_T$ = (Unit Intensity(phagosome center)$_T$ − Unit Intensity(background)$_T$) / (Unit Intensity(phagosome center)$_{T0}$- Unit Intensity (background)$_{T0}$).

(F-G) The relative mNG::LGG-1 (F) or -2 (G) signal intensity in the center of a phagosome (Y-axis) over time in the 2-min interval (X-axis). "0 min" is the moment when pseudopods are sealed and a nascent phagosome forms. (F) The data for the wild-type, *ced-1(e1735),* and *dyn-1(n4039)* mutant embryos are from Figs 1F and 8(A-B), respectively. (G) The data for the wild-type, *ced-1(e1735),* and *dyn-1(n4039)* mutant embryos are from Figures 1I and 8(C-D), respectively. (Figure 9-source data 1)

(H) A diagram illustrating that the relative mNG signal on the surface of a phagosome is measured over time to create sub-figures (I) and (J). At time point T (time after "0" min), the Relative signal intensity $_T$ = (Unit Intensity(phagosome surface (the green ring))$_T$ −Unit Intensity(background)$_T$) / (Unit Intensity(phagosome surface)$_{T0}$ - Unit Intensity(background)$_{T0}$).

(I-J) The relative mNG::LGG-1 or -2 signal intensity on the surface of a phagosome (Y-axis) over time in the 2-min interval (X-axis). "0 min" indicates the moment when pseudopods are sealed and nascent phagosome forms. (I) The data for the wild-

type, *ced-1(e1735),* and *dyn-1(n4039)* mutant embryos are from Figures 1F and 8(A-B), respectively. (J) The data for the wild-
type, *ced-1(e1735),* and *dyn-1(n4039)* mutant embryos are from Figs 1I and 8(C-D), respectively. (Figure 9-source data 2)
(K-L) Box-and-Whiskers plots of the relative mNG signal intensity measured in the center of phagosomes 50 min-post the
formation of nascent C3 phagosomes from 15 each of wild-type, *ced-1(e1735),* and *dyn-1(n4039)* mutant embryos. Red
dashed lines indicate the position of value 1, which represents no signal enrichment relative to the background signal. ***,
p<0.001, Student *t*-test of each mutant compared to the wild-type value. (Figure 9-source data 3)
612

We also examined whether the pathway composed of the small GTPase CED-10 and its bipartite

Guanine Nucleotide Exchange Factor (GEF) CED-5 and CED-12, which acts parallel to the CED-1

pathway in the engulfment of cell corpses, plays any role in promoting the incorporation of

autophagosomes into phagosomes. In the *ced-5(n1812)* null mutant and *ced-10(n1993)* loss-of-function

mutant embryos, due to the presence of the parallel engulfment pathways (the CED-1 and RAB-35

pathways), some C1, C2, or C3 phagosomes eventually form, although they usually suffer a severe delay,

allowing the analysis of phagosome degradation [20,22,49]. In these two mutants, the accumulation of

the LGG-1 and LGG-2 signals on the surfaces of the phagosomes and the subsequent accumulation of

signals inside phagosomal lumen were normal both in the time course and in the levels of signal

enrichment (**Fig 10 B, C-E, G, H-J**), indicating that both the recruitment and fusion of autophagosomes

to phagosomes are normal. We thus conclude that, unlike the CED-1/-6/DYN-1 pathway, the CED-5/-

10 pathway is not involved in regulating the incorporation of autophagosomes to phagosomes (**Fig 10**

K).

626
Figure 10. *ced-6*, but not *ced-5* or *ced-10*, is required for the incorporation of autophagosomes into phagosomes
(A-C and F-H) Time-lapse image series monitoring the presence or absence of puncta (white arrows) labeled with
mCherry::LGG-1 (A-C) or -2 (F-H) on C3 phagosomes (white arrowheads) and the subsequent entry of the mCherry signal into
the phagosomal lumen in *ced-6, ced-5,* and *ced-10* mutant embryos. DIC images mark the position of the cell corpse. "0 min"
is the moment when a nascent phagosome just seals. Scale bars are 2μm.
(D and I) The relative mCherry::LGG-1 (D) or -2 (I) signal intensity in the center of a phagosome (Y-axis) over time (in the 2-
min interval) (X-axis). "0 min" indicates the moment when a nascent phagosome just seals. (D) The data for the wild-type,
ced-6(n2095), ced-5(n1812), and *ced-10(n1993)* mutant embryos are from Figs 1E and 9(A-C), respectively. (I) The data for
the wild-type, *ced-6(n2095), ced-5(n1812),* and *ced-10(n1993)* mutant embryos are from Figs 1H and 9(F-H), respectively.
(Figure 10-source data1)

The incorporation of lysosomes into phagosomes is not significantly affected by the lack of autophagosome biogenesis

The incorporation of lysosomes into phagosomes is an essential force that drives the degradation of phagosomal contents [3]. As a portion of the autophagosomes would fuse with lysosomes and become autolysosomes that retain lysosomal features [9], it is possible that among all LGG$^+$ puncta, the autolysosomes are the sole portion that contributes to phagosome degradation by depositing lysosomal luminal proteins to the phagosomal lumen. To test whether this is the case, we examined whether mutations that specifically impair autophagosomes biogenesis would reduce the amount or speed of the incorporation of organelles with lysosomal features into phagosomes. *C. elegans* NUC-1 is an endonuclease belonging to the DNase II family and resides in the lysosomal lumen [50,51]. Using a NUC-1::mCherry reporter expressed in engulfing cells as a lysosomal luminal marker, we quantified the level and rate of lysosome-phagosome fusion over time. In wild-type embryos, in phagosomes containing cell corpses C1, C2, and C3, we observed first the attachment of NUC-1::mCherry puncta on the surfaces of phagosomes and subsequently the accumulation of the mCherry signal inside the phagosomal lumen (**Fig 11A**). This dynamic process represents the recruitment and the subsequent fusion of lysosomal particles to phagosomes, which result in the delivery of NUC-1::mCherry into the phagosomal lumen. We next monitored NUC-1::mCherry inside the phagosomal lumen in *lgg-1(tm3489)*, *lgg-2(tm5755)*, and *atg-7(bp411)* embryos, which are defective in the biogenesis of autophagosomes. In these mutants, numerous mCherry::NUC-1 puncta are enriched on the surfaces of phagosomes C1, C2, and C3 and are subsequently fused to phagosomes (**Fig 11 B-D**). As a result, the accumulation of the mCherry signal is observed in the phagosomal lumen in all samples like in wild-type embryos (**Fig 11 B-E**). Quantitative analysis of 15 phagosomes for each genotype found that 60 min after phagosome formation, the average folds of increase of the luminal mCherry signal over the "0 min" time point in *lgg-1*, *lgg-2*, and *atg-7* mutants are not significantly different from that observed from wild-type samples (**Fig 11F**). In addition, the median value of the first time point when NUC-1::mCherry signal was detected inside the phagosomal lumen was not significantly different in all four genotypes (**Fig 11G**). These data strongly suggest that the incorporation of lysosomes into phagosomes in *lgg-1*, *lgg-2*, and *atg-7* mutants is as efficient as in wild-type embryos, at least within the detection

673 range of our assay. Therefore, defects in the biogenesis of autophagosomes, which potentially would

674 indirectly result in the lack of autolysosomes, do not appear to significantly affect the incorporation of

675 lysosomes into phagosomes.

Figure 11. Defects in the formation of autophagosomes do not significantly affect the timing or efficiency of lysosomes incorporation into phagosomes

The time-lapse recording was conducted on phagosomes containing C1, C2, and C3 in wild-type and named mutant embryos carrying P*ced-1nuc-1::mCherry*, the lysosomal lumen reporter, and P*ced-1PH(PLCγ)::gfp*, the marker for extending pseudopods and nascent phagosomes. (A-D) Fluorescence time-lapse images of a C3 phagosome in each strain with the indicated genotype. "0 min" is the moment when a nascent phagosome (white arrowhead) just seals. Yellow arrows mark the lysosomal particles that are located on phagosomal surfaces. White open arrows mark the phagosomes with mCherry signals in the lumen. Scale bars are 2.5 µm.

(E) The relative NUC-1::mCherry signal intensity in the center of a phagosome (Y-axis) over time (in the 2-min interval) (X-axis). "0 min" indicates the moment when a nascent phagosome is just sealed. Data are from Fig 10 (A-D). (Figure 11-source data 1)

(F) Bar graphs of the average fold-of-increase of the mCherry signal intensity at the center of phagosomal lumen 60 min-post the formation of nascent C3 phagosomes. Bars represent the mean, the error bars indicate standard deviation, and each dot represents a sample. 15 phagosomes of the indicated genotype were scored. Student *t*-test of each mutant compared to the wild-type value. ns, not significant. (Figure 11-source data 2)

(G) Bar graphs of the average time when the NUC-1::mCherry signal is first detected inside the lumen of 15 C3 phagosomes in the indicated genotypes. Bars represent the mean, the error bars indicate standard deviation, and each dot represents a sample. Student *t*-test of each mutant compared to the wild-type value. ns, not significant. (Figure 11-source data 2)

In the wild-type embryos that co-express the mNG::LGG-1 or -2 reporters with the NUC-

1::mCherry reporter in engulfing cells, we observed that, in addition to the lysosomal particles that were

mNG[-] but mCherry[+], there were two kinds of mNG[+] puncta that were recruited to phagosomal surfaces

and subsequently fused to phagosomes. The puncta are not labeled with mCherry represent

autophagosomes that are not fused to lysosomes; on the other hand, the mNG$^+$ mCherry$^+$ puncta

represent autolysosomes (**Fig 12 A-B**). Quantification of these two kinds of puncta on the surfaces of

the C1, C2, and C3 phagosomes reveals that 40.6% and 36.5% of LGG-1$^+$ and -2$^+$ puncta are

autolysosomes, respectively (**Fig 12 C**). Whereas the LGG$^+$ NUC-1$^+$ autolysosomes are capable of

contributing materials that are autophagosome-specific and lysosome-specific to phagosomes after

fusion, the LGG$^+$ NUC-1$^-$ population of autophagosomes are likely to deliver certain autophagosome-

specific material to phagosomes.

Figure 12. Visualizing the enrichment of lysosomes, autophagosomes, and autolysosomes on phagosomal surfaces.
(A-B) DIC and fluorescence images of C2 and C3 phagosomes in wild-type embryos co-expressing P$_{ced-1}$ *nuc-1::mCherry* with
P$_{ced-1}$ *mNG::lgg-1 or* P$_{ced-1}$ *mNG::lgg-2.* Images were captured at the indicated time point after phagosome formation ("0 min").
DIC images mark the position of the cell corpse (white arrows). Yellow arrows, open white arrows, and white arrowheads label
the GFP$^+$ mCherry$^-$ puncta (autophagosomes), GFP$^-$ mCherry$^+$ puncta (lysosomes), and GFP$^+$ mCherry$^+$ double-positive
puncta (autolysosomes), respectively, on phagosomal surfaces. Scale bars are 2μm.
(C) Bar graph depicting the percentage of autolysosomes (GFP$^+$ mCherry$^+$ puncta) among LGG-1$^+$ or -2$^+$ puncta on the
surface of phagosomes. Each sample (dot) represents the distribution of the aforementioned puncta collected from the C1, C2,
and C3 phagosomes of one embryo. Seven embryos were scored. Bars and error bars represent mean and standard deviation
values. (Figure 12-source data 1)

Impairing autophagosomes biogenesis results in moderate defects in phagosome acidification and

the digestion of DNA from apoptotic cells

Acidification of the phagosomal lumen is a critical event for the degradation of phagosomal

content. To examine whether defects in autophagosome biogenesis would affect the acidification of

phagosomes, we developed an acidification reporter for the phagosomal lumen. This reporter, P$_{his-72}$ *his-*

72::gfp::mCherry, expresses a HIS-72 (histone H3.3)::GFP::mCherry fusion protein in all cells,

including cells that undergo programmed cell death [52]. We quantified the GFP/mCherry signal ratio

in the phagosomal lumen over time and normalized it over the GFP/mCherry signal ratio at the "0 min"

time point when a nascent phagosome just formed. We refer to this normalized value as an

"acidification index". In a typical phagosome in a wild-type embryo, the acidification index reduces

continuously from 1.0 (t=0 min post engulfment) to 0.4 (t=51 min) (**Fig 13 A, D**). In addition, in 100%

of the phagosomes (n=6), the acidification index values at the 51 min time point are <0.6, among which

in 35% of the samples, the index values are <0.4 (**Fig 13 E**). On the contrary, in *cup-5(n3264)* mutant

embryos, the acidification of the phagosomal lumen is severely defective (**Fig 13 B, D, E**). The

acidification index values of all the phagosomes (n=13) at t=51 min are >0.6, and in 53.8% of the

phagosomes, the index values are >0.9, indicating a minimal reduction of the GFP signal. *cup-5*

encodes a lysosomal TRP channel homologous to human Mucolipin IV [53,54]. CUP-5 and its

mammalian homologs play conserved and essential roles in supporting lysosomal biogenesis and

functions [53,54]. The severe acidification defect observed here emphasizes the essential role of

lysosomes for phagosome degradation. Compared to the *cup-5* mutant embryos, in *atg-7(bp411)* null

mutant embryos, we observed a moderate phagosome acidification defect (**Fig 13 C, D, E**). The average

acidification index values for wild-type, *atg-7(bp411)*, and *cup-5(n3264)* mutants are 0.443, 0.679, and

0.912, respectively, at t=51 min. In *atg-7(bp411)* mutants, the acidification index values in 69.2% of the

phagosomes at t=51 min are >0.6, yet the values of only 15.3% phagosomes are >0.9. This moderate

defect is distinct from that observed from the *cup-5* mutants.

Figure 13. Inactivation of autophagy results in a modest phagosomal acidification defect.

(A-C) Time-lapse imaging series of phagosomes (white arrowheads in DIC images) of wild-type, *cup-5*, and *atg-7* mutant
embryos expressing P*his-72* *his-72::gfp::mCherry*. Open white arrows depict the nuclei of engulfed cell corpses, labeled with
both the GFP and mCherry markers. Reduction of the GFP signal intensity over time is indicative of phagosome acidification.
"0 min" is when a phagosome is just sealed. Scale bars are 2μm.
(D) The acidification index curves of three phagosomes (Y-axis) over time (in the 3-min interval) (X-axis) in embryos with the
labeled genotypes. 0 min" indicates the moment when a phagosome is just sealed. The data of the wild-type, *cup-5 (n3265),*
and *atg-7 (bp411)* are from A-C, respectively. (Figure 13-source data 1)
(E) Histogram depicting the distribution of the acidification index measured at 51-min-post the formation of nascent
phagosomes. In wild-type, *cup-5 (n3265)*, and *atg-7 (bp411)* mutant embryos, 6, 12, and 12 phagosomes were scored.
(Figure 13-source data 2)

The overall phagosome degradation efficiency in wild-type and autophagy mutant embryos has

been measured by the reduction of phagosomal size over time (**Fig 6**). To further evaluate the

hydrolytic activities inside the phagosomal lumen in the autophagy mutant embryos, we measured the

rate of the degradation of chromatin DNA in engulfed apoptotic cells. We found that, unlike a previous

report that concluded that the NUC-1 endonuclease acts in cells undergoing programmed cell death to

generate chromatin DNA fragments [50], NUC-1 actually acts in engulfing cells, in the phagosomal

lumen to degrade chromatin DNA of engulfed apoptotic cells (Pickett J., Auld N., Lucas, L., Pena-

Ramos O., and Zhou, Z., unpublished results). During the above study, we established HIS-

72::mCherry as a reporter for chromatin DNA. Histone b H3.3, together with other histones and the

DNA wrapping around them, form the nucleosome, the fundamental subunit of the chromatin. In Z-

section images, inside the phagosomal lumen, the HIS-72::mCherry signal appears as a condensed red

disc, outlining the nucleus of the engulfed apoptotic cell. We observed that in the wild-type embryos,

the degradation of the apoptotic cell-chromatin DNA inside the phagosomal lumen allowed the

reduction of the size of the mCherry$^+$ disc over time (**Fig 14 A, D**). Within 60 min after the formation of

a phagosome, the average diameter of the mCherry$^+$ disc reduces to 49% of the 0 min value (**Fig 14 E**).

On the contrary, in the *nuc-1* null mutant embryos, the diameter of the mCherry$^+$ disc remains the same

over the entire period of time-lapse observation (>80 min) (Pickett J., Auld N., Lucas, L., Pena-Ramos

O., and Zhou, Z., unpublished observation), indicating the lack of degradation of chromatin DNA when

NUC-1 activity is absent.

We observed that, as predicted for a mutant severely defective in lysosomal biogenesis and

functions, in *cup-5(n3264)* mutant embryos, within 60 min post phagosome formation, the average

diameter of the mCherry+ disc remains at 89% of the 0 min-value (**Fig 14 B, E**). In comparison, in *atg-*

7(bp411) mutant embryos, that value is reduced to 66% of the 0 min-value; in addition, the diameters of

the mCherry$^+$ discs in 25% of the phagosomes are > 80% of the 0 min-value. These results indicate that

in *atg-7* mutants, there is a defect in the digestion of chromatin DNA, yet this defect is relatively

moderate.

Figure 14: Inactivation of autophagy results in a modest delay of the degradation of apoptotic cell DNA.
(A-C) Time-lapse imaging series monitoring the shrinkage of the apoptotic cell nucleus inside a C1, C2, or C3 phagosome (white arrowhead in DIC images) in three different strains expressing P*his-72 his-72::mCherry*. Apoptotic cell nuclei are labeled with HIS-72::mCherry (open white arrows). "0 min" is when a phagosome is just sealed. Scale bars are 2μm.
(D) The ratio of the nuclear diameter of a phagosome (Y-axis) at labeled time points compared to that of the "0 min" diameter (in the 3-min interval) (X-axis). "0 min" indicates the moment when a phagosome is just sealed. The data of the wild-type, *cup-5 (n3265),* and *atg-7 (bp411)* are from figures 1(A-C), respectively. (Figure 14-source data 1)
(E) Histogram depicting the distribution of the ratio of nuclear diameters measured at 60min-post phagosome formation. In wild-type, *cup-5 (n3265)*, and *atg-7 (bp411)* mutant strains, 9, 9, and 12 engulfed apoptotic cells were scored. (Figure 14-source data 2)

Discussion

Autophagosomes play essential roles in cellular homeostasis by eliminating harmful protein aggregates and damaged organelles, and in stress response by converting intracellular organelles to nutrients during starvation [9]. We have identified a novel function of autophagosomes in facilitating the degradation of apoptotic cells through fusing to phagosomes. In *C. elegans* cells that engulf apoptotic cells, autophagosomes are recruited to the surfaces of phagosomes and subsequently fuse to the phagosomal membrane. Through this previously unknown interaction, autophagosomes facilitate the degradation of apoptotic cells inside phagosomes, presumably through providing certain materials to phagosomes. We have further identified a signaling pathway that promotes the recruitment and subsequent fusion of autophagosomes to phagosomes. Together, these findings reveal a novel mechanism through which the pathways that control autophagy and phagocytosis converge, underlining the importance of this mechanism in the degradation of apoptotic cells. We propose that this mechanism might be conserved in the metazoans and has a significant impact in the regulation of the immunological consequences of dying cells inside human bodies.

Canonical autophagosomes play an essential role in the degradation of apoptotic cells

Previously, *C. elegans* autophagy genes have been implicated in facilitating the clearance of
apoptotic cells. Autophagy occurring in apoptotic cells was proposed to facilitate the exposure of
phosphatidylserine, which is the "eat me" signal that attracts engulfing cells to the surfaces of apoptotic
cells in mice embryonic bodies and *C. elegans* [55,56]. Multiple autophagy genes were reported to
participate in the clearance of apoptotic cells in *C. elegans* engulfing cells [11-13,56]. They were
reported to work together with Class II PtdIns3 kinase PIKI-1 to facilitate the production of PtdIns(3)P
on phagosomal surfaces [11], to promote the recruitment of RAB-5 and RAB-7 to the phagosomal
surfaces [13], and to promote phagosome degradation [56]. All of the above findings suggest that the
autophagy genes examined might have separate functions in the degradation of phagosomal contents
unrelated to their canonical roles in autophagosome biogenesis. Our work reported here, on the other
hand, presents a different discovery regarding how the autophagy machinery regulates phagosome
degradation.

We found that the canonical double-membrane autophagosomes actively participate in the
degradation of apoptotic cells inside phagosomes. In the time-lapse recording series, we first observed
that vesicles labeled with LGG-1/or -2 reporters are recruited to phagosomal surfaces; subsequently, the
LGG-tagged reporters enter the phagosomal lumen. We propose that the entry of the LGG reporters is a
result of the fusion of the outer membrane of a double-membrane vesicle with the membrane of a
phagosome based on two lines of evidence. First, the fusion between these two membranes would allow
the release of the inner vesicle into the phagosomal lumen. Moreover, the RAB-7 GTPase and its
effector -- the HOPS complex, are well-known membrane tethering factors that promote the fusion
between two intracellular membranes [42,47]. Their essential roles in supporting the entry of the LGG
reporters into the phagosomal lumen that we have discovered strongly indicate that this event is a result
of the fusion between the outer membrane of autophagosomes and phagosomes. Formally, other
possible explanations of the entry of the LGG reporters exist, such as that phagosomes engulf the LGG-
labeled vesicles. However, currently, there are no reported findings that phagosomes would engulf
intracellular vesicles.

Using the immuno-electron microscopy technique, Segalen et al previously found that the LGG-
1 and -2 are specifically enriched on double-membrane autophagosomes in *C. elegans* embryos [27].
Five critical sets of evidence we report here further indicate that the LGG-1- and -2-labeled vesicles that
are incorporated into phagosomes are double-membrane autophagosomes, not the single-membrane

LAP vesicles. First, only when the vesicles are composed of double membranes and when the LGG-1/
or -2 reporter molecules label both the inside and outside membranes, the entry of the reporters into
phagosomal lumen is possible (**Fig 1K**). Conversely, if the LGG-1 or -2 labeled vesicles are of a single
membrane, after vesicle-phagosome fusion, the reporter signal would remain on the phagosomal
membrane, just like the membrane-bound lysosomal marker CTNS-1::mRFP (**Fig 1 – figure**
supplement 1). Second, these vesicles are also labeled with GFP-tagged ATG-9, an integral membrane
protein inserted into the autophagosomal membranes. Third, we found that besides in embryos, in the
gonad of adult hermaphrodites, many LGG-1 labeled vesicles are also attached to the surfaces of
phagosomes that contain germ cell corpses in the *rab-7* mutant background in which membrane fusion is
presumably blocked. Also, in the *rab-7* mutant adult hermaphrodites, thin-section transmission electron
micrographs reveal multiple double-membrane vesicles that are in close contact with the phagosomal
membranes. These vesicles resemble autophagosomes in morphology [39,40]. Fourth, the production
of these vesicles relies on genes pivotal for autophagosome biogenesis, such as *atg-3*, *atg-7*, and *atg-9*.
Last but not least, genes encoding the *C. elegans* homologs of mammalian ULK1, ATG13, and ATG14,
which are dispensable for the generation of LAP vesicles in mammalian cells [6], are required for the
formation of the LGG-labeled vesicles discussed here and for the efficient clearance of apoptotic cells,
further verifying that these vesicles are distinct from the single-membrane LAP vesicles. Together,
these five sets of findings demonstrate that double-membrane autophagosomes, not single-membrane
LAP vesicles, are being incorporated into phagosomes.
In *C. elegans*, LC3-dependent phagocytosis was reported to function in the clearance of the
midbody, a structure that is essential for the completion of cytokinesis, and the second polar body
generated during female meiosis [57,58]. The clearance of midbodies and second polar body is
considered independent of autophagosomes based on the observation that neither *unc-51* nor *epg-8* is
required for the above processes [57,58]. However, whether the LGG markers attached to phagosomes
carrying midbodies or polar bodies represent double-membrane or single-membrane vesicles have not
been investigated [57,58]. Whether LAP vesicles exist in *C. elegans* cells still needs to be demonstrated.
On the other hand, we found that in *atg-13* mutant embryos, in which the biogenesis of the LAP vesicles
is not supposed to be affected, the LGG-labeled puncta largely disappeared, indicating that these puncta
are primarily autophagosomes instead of LAP vesicles (**Fig 2 – figure supplements 1 -2**). This
observation does not support the existence of a substantial population of the LAP vesicles in *C. elegans*
embryos, at least not in the cells in which the LGG reporters are expressed.

The observation of the LGG-1 or -2 signals inside the phagosomal lumen is made using the

mCherry reporter. The mCherry protein is resistant to acidic pH [31]. Phagosome luminal pH value

reduces from 5.5-6.0 to 4.5-5.5 after incorporating lysosomes [59], leading to the inactivation of the

fluorophore in the commonly used GFP reporter [30]. In fact, in *C. elegans* engulfing cells, GFP-tagged

LGG-1 or -2, unlike the mCherry-tagged reporters, are only observed on the surfaces but not in the

lumen of phagosomes, presumably due to the acidic environment of the phagosomal lumen (**Fig 1**). It is

thus conceivable that in other experimental systems, a mCherry-tagged LC3 marker might disclose a

previously overlooked fusion between autophagosomes and phagosomes.

Based on the evolutionary conservation of almost all molecular mechanisms, in particular the

strong conservation of the known mechanisms that control phagocytosis and autophagy in metazoans

[26,35], we propose that in mammalian cells, besides LAP vesicles, canonical autophagosomes might

also contribute to the degradation of apoptotic cells in phagosomes. The demonstration of an

autophagosome-phagosome fusion event in *C. elegans*, a well-established model organism for studying

cell death-related events, opens a path to investigate whether the event and mechanism of

autophagosome-phagosome fusion is an evolutionarily conserved event. In addition, the

autophagosome-phagosome fusion might contribute to the degradation of other kinds of phagosomal

cargos besides apoptotic cells.

LGG-1 and LGG-2 define three subpopulations of autophagosomes that are incorporated into

phagosomes

LGG-1 and -2 are close homologs that each labels autophagosomes. We have observed three

separate classes of autophagosomes that are incorporated into phagosomes: LGG-1$^+$ LGG-2$^-$, LGG-1$^-$

LGG-2$^+$, or LGG-1$^+$ LGG-2$^+$. These subpopulations might represent autophagosomes at different stages

of maturation [27]. Remarkably, we have found that the *lgg-1*; *lgg-2* double mutants display a further

enhanced Ced phenotype than that displayed by each of the *lgg-1* or *lgg-2* single mutants. Together,

these results indicate that autophagosomes at different stages of maturity all contribute to the

degradation of apoptotic cells. These results are also consistent with our finding that the

autophagosome-phagosome fusion is primarily independent of lysosome-phagosome fusion (**Fig 11**)

(Also see the next section).

What do autophagosomes contribute to the degradation of apoptotic cells

We have found that the blockage of the biogenesis of autophagosomes or the recruitment or
fusion of autophagosomes to phagosomes results in a significant delay in the degradation of phagosomal
content and, consequently, the persistent appearance of engulfed cell corpses. Among other possible
mechanisms, the incorporation of autophagosomes into phagosomes presumably delivers certain
substances to the lumen and/or membrane of phagosomes, substances that are important for the
degradation of phagosomal contents. Here we discuss several possible candidates and mechanisms
supporting the roles of autophagosomes in phagosome degradation.

We have observed that the LGG$^+$ puncta inside engulfing cells are a collection of two
subpopulations: autophagosomes that are already fused to lysosomes and become autolysosomes (LGG$^+$
NUC-1$^+$) and autophagosomes that are not fused to lysosomes (LGG$^+$ NUC-1$^-$) (**Fig 12**). The
autolysosome populations, which occupied 40.7% and 36.5% of the LGG-1$^+$ and LGG-2$^+$ puncta
observed on phagosomal surfaces, respectively, might contribute both autosomal and lysosomal
materials to the phagosomes. In *atg-7* mutants, in which the biogenesis of autophagosomes is virtually
blocked, the acidification of the phagosomal lumen and the degradation of the nuclear DNA of engulfed
apoptotic cells are modestly defective (**Figs 13** and **14**). These defects might be partially attribute to the
lack of autolysosomes that are incorporated into phagosomes. However, in cells that maintain a stable
lysosome population, the lack of autophagosomes to fuse with lysosomes is not expected to influence
the overall activities of lysosomes. In *atg-7*, *lgg-1*, and *lgg-2* mutant embryos, we found that the lack of
autophagosomes does not significantly affect the efficiency of lysosome-phagosome fusion (**Fig 11**).
This result could be due to that in an engulfing cell, the population of autolysosomes, comparing to that
of lysosomes, is a rather minor population. It supports the notion that the autophagy pathway primarily
regulates activities that are additive to the lysosomal activities for promoting phagosome degradation,
although the assay might not be sensitive enough to detect a modest decrease. Moreover, in *rab-7(m$^-$z$^-$)*
mutant embryos in which the fusion of both the lysosomes and the LGG$^+$ particles to phagosomes are
blocked, the phagosome degradation defect is much more severe than in autophagosome biogenesis
mutants. These results suggest that the collected population of LGG$^+$ vesicles contribute to phagosome
degradation in a manner in parallel to lysosomes. In addition to the autolysosomal subpopulation, the
LGG-1$^+$ NUC-1$^-$ (59.3% of the LGG$^+$ puncta) or LGG-2$^+$ NUC-1$^-$ (63.5% of the LGG2$^+$ puncta)
subpopulations that fuse to phagosomes are likely to contribute certain unique, lysosome-independent
activity(ies) to the phagosomes.

What could be the unique activities or mechanisms provided by the autophagosomes? Through
fusion to phagosomes and releasing the inner vesicles to the phagosomal lumen, autophagosomes might

deliver certain protein or lipid molecules to facilitate the degradation of phagosomal cargos. For

example, the outer membrane of the autophagosomes might provide important signaling molecules to

the phagosomal membrane. One such candidate is PtdIns(3)P, a membrane signaling molecule essential

for many membrane trafficking events. On the phagosomal surfaces, PtdIns(3)P recruits PtdIns(3)P-

binding proteins, including the sorting nexins SNX-1, SNX-6, and LST-4/SNX-9 and the HOPs complex,

which subsequently drive multiple membrane remodeling events that promote phagosome maturation

[25,26]. In *C. elegans*, PtdIns(3)P is presented on phagosomal membrane over time in a two-peak

pattern [24]. Autophagosomes are coated with PtdIns(3)P [10]. Judging by the timing of

autophagosome-phagosome fusion, autophagosomal membranes are likely to contribute PtdIns(3)P for

the rise of the second peak of PtdIns(3)P.

Alternatively, the autophagosome-phagosome fusion incorporates membrane materials to the

phagosomal membrane and thus should increase the size of phagosomal membrane. Increasing the

overall amount of phagosomal membrane might aid the degradation of the apoptotic cell by facilitating

the recruitment of lysosomes to the phagosomal surfaces. We previously reported that phagosomes

extend transient lipid tubules to capture lysosomal particles in the cytoplasm and bring these particles

back to the phagosomes for fusion [20,25]. The increased phagosomal membrane material might

support the extension of these lipid tubules. A substantial body of future investigation is required to

discover the mechanisms employed by autophagosomes that facilitate phagosome degradation.

The CED-1 signaling pathway drives the incorporation of autophagosomes to phagosomes

Autophagosomes are incorporated into phagosomes in two sequential steps: (1) they are recruited

to the surfaces of phagosomes, detected by the enrichment of punctated LGG-1 or -2 fluorescence

reporters on phagosomal surfaces, and (2) they subsequently fuse to phagosomes, detected by the

enrichment of the LGG-1 or -2 reporter signals inside the phagosomal lumen. RAB-7 and its effector,

the HOPS complex, are known to act as tethering factors that facilitate the fusion of various intracellular

organelles, including autophagosomes, to lysosomes [27,60]. Closely related to this study, RAB-7 and

VPS-18 are pivotal for the fusion between lysosomes and phagosomes and for the degradation of

apoptotic cells inside phagosomes in *C. elegans* [20,48,61]. Here we observed that 100% of the LGG[+]

puncta attached to phagosomal surfaces were also labeled with RAB-7. We further discovered that

RAB-7 and VPS-18 played essential and specific roles in the fusion but not for the recruitment of

autophagosomes to phagosomes. Our finding adds a new pair of organelles that depend on the RAB-

7/HOPS complex for fusion to each other. As RAB-7 is enriched on the surfaces of both phagosomes

and autophagosomes ([20] and (**Fig 8 (B-D)**)), and as the HOPS complexes in mammalian cells and

Drosophila are known to interact with the SNARE complex, the membrane fusion machinery [62,63],

we propose that the RAB-7/HOPS complex acts on the surfaces of phagosomes and autophagosomes to

facilitate autophagosome-phagosome fusion via promoting the interaction between the SNARE

complexes on phagosomes and autophagosomes (**Fig 8 K**). Currently, the specific SNARE proteins that

catalyze the autophagosome-phagosome fusion have not been identified.

The CED-1 signaling pathway, which initiates the maturation of phagosomes that bear apoptotic

cells, is essential for the enrichment of GTP-bound RAB-7 to the surfaces of phagosomes [20]. Here we

have found that CED-1, CED-6, and DYN-1 drive the incorporation of autophagosomes to phagosomes,

in addition to driving the incorporation of early endosomes and lysosomes previously discovered [20,22].

In *ced-1* mutants, for example, the recruitment of autophagosomes to the surfaces of phagosomes is

almost completely blocked. Due to the severe recruitment defect, whether the *ced-1* null mutation

further impairs the fusion between autophagosomes and phagosomes cannot be readily visualized.

However, since a *ced-1* null mutation impairs the recruitment of RAB-7 to [20] and the production of

PtdIns(3)P on phagosomal surfaces [20,24], and since the recruitment of the HOPS complex to the

surfaces of intracellular organelles requires both RAB7 and PtdIns(3)P [47,64,65], we predict that the

CED-1 pathway would also control the RAB-7/HOPS complex-mediated autophagosome-phagosome

fusion (**Fig 9 K**). Identifying the CED-1 signaling pathway as the driving force for the incorporation of

autophagosomes to phagosomes helps to reveal the molecular mechanisms behind the crosstalk between

autophagy and phagocytosis.

Key Resources Table

Reagent type	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*E. coli*)	OP50	CGC	OP50	
Strain, strain background (*C. elegans*)	N2	CGC	Wild-type Bristol N2	
Strain, strain background (*C. elegans*)	VC308	CGC	*rab-7(ok511) /mIn1 II*	
Strain, strain background (*C. elegans*)	ZH0989	[20]	*unc-76(e911) enIs36 [punc-76(+), Pced-1 ced-1::gfp, Pced-1 2xFYVE::mRFP] V*	Figure 6 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2059	This study	*unc-76(e911)V; enEx979 [punc-76(+), Phis-72 HIS-72::GFP::mCherry]*	Figure 13 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2105	This study	*cup-5(n3264) III; unc-76(e911) V; enEx979*	Figure 13 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2573	This study	*lgg-2(tm5755) IV; unc-76(e911) V; enEx1223 [Pced-1mCherry::lgg-2, punc-76(+)]*	Figure 5-figure supplement 1 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2632	This study	*lgg-2(tm5755) IV; unc-76(e911) V; enEx1267 [Punc-76(+), Pced-1::gfp::lgg2]*	Figure 5-figure supplement 1 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2715	This study	*rab-7(ok511) II / mIn1 II; unc-76(e911) V; enEx1320 [punc-76(+), Pced-1mCherry::lgg-2]*	Figure 8 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2782	This study	*rab-7 (ok511) II / mIn1 II; unc-76(e911) V; enEx1376 [punc-76(+), Pced-1 mCherry::lgg-1]*	Figure 8 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2831	This study	*lgg-1(tm3489) II / mIn1 I ; lgg-2(tm5755) IV*	Figure 5 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2835	This study	*lgg-1(tm3489) II / mIn1 I ; unc-76(e911) enIs36 V*	Figure 6 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2838	This study	*lgg-1(tm3489) II / mIn1 II; unc-76(e911) V; enEx1428 [punc-76(+), Pced-1 gfp::lgg-1]*	Figure 5-figure supplement 1 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2841	This study	*lgg-1(tm3489) II / mIn1 II; unc-76(e911) V; enEx1431 [punc-76(+), Pced-1 mCherry ::lgg-1]*	Figure 5-figure supplement 1 Available from the Zhou Lab
Strain, strain background (*C. elegans*)	ZH2875	This study	*lgg-2(tm5755) IV; unc-76(e911) enIs36 V*	Figure 6 Available from the Zhou Lab

Strain, strain background (C. elegans)	ZH2889	This study	unc-76(e911) V; enEx1459 [punc-76, Pced-1 PH::mrfp),Pced-1 mNG::lgg-2]	Figure 1 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2903	This study	atg-9(bp564) him-5(e1490)V; lin-15AB(n765ts) X; enEx1468 [Plin-15(+), Pced-1 PH::mrfp, Pced-1 mNG::lgg-1]	Figure 3-figure supplement 1, Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2907	This study	vps-18(tm1125) II; enIs80 [punc-76(+), Pced-1 ced-1::gfp, Pced-1mCherry::lgg-1] IV; unc-76(e911) V	Figure 8 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2916	This study	ced-1(e1735) I; unc-76(e911) V; enEx1470 [Pced-1 mNG::lgg-1, punc-76(+), Pced-1 PH::mrfp]	Figure 9 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2919	This study	enIs82 [unc-76(+), Pced-1 ced-1::gfp, Pced-1mCherry::lgg-1] II; unc-76(e911) V	Figure 1 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2921	This study	unc-76(e911) enIs85 [punc-76(+), Pced-1 PH::mrfp, Pced-1mNG::lgg-2] V.	Figure 1 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2922	This study	atg-9(bp564) him-5(e1490) V; lin-15AB(n765ts) X ; enEx1472 [Plin-15(+), Pced-1 PH::mrfp, Pced-1 mNG::lgg-2]	Figure 3-figure supplement 1, Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2929	This study	vps-18(tm1125) II; unc-76(e911) V; enIs83 [punc-76(+), Pced-1 ced-1::gfp, Pced-1mCherry::lgg-2] X	Figure 8 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2934	This study	ced-1(e1735) I; unc-76(e911) enIs85 V	Figure 9 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2950	This study	enIs82 II; atg-7(bp411) IV; unc-76(e911) V	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2951	This study	atg-7(bp411) IV; enIs83 X	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2952	This study	enIs82 II; atg-13(bp414) III	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2953	This study	atg-13(bp414) III; enIs83 X	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2954	This study	epg-8(bp251) I; enIs82 II; unc-76(e911) V	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2955	This study	epg-8(bp251) I; unc-76(e911) him-5(1490) V; enIs83 X	Figure 2 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2980	This study	atg-7(bp411) IV; unc-76(e911) enIs36 V	Figure 6 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2992	This study	enIs87 [punc-76, Pced-1 PH::mrfp, Pced-1 mNG::lgg-1] I; unc-76(e911) V	Figure 1 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH2994	This study	enIs82 II; ced-6(n2095) III; unc-76(e911) V	Figure 10 Available from the Zhou Lab

Strain, strain background (C. elegans)	ZH2995	This study	ced-6(n2095) III; unc-76(e911) V; enIs83 X	Figure 10 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3009	This study	ced-5(n1812) IV; unc-76(e911) V; enIs83 X	Figure 10 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3010	This study	enIs82 II; ced-5(n1812) IV; unc-76(e911) V	Figure 10 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3011	This study	enIs82 II; ced-10(n1993) IV; unc-76(e911) V	Figure 10 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3012	This study	ced-10(n1993) IV; unc-76(e911) V; enIs83 X	Figure 10 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3014	This study	enIs87 I; unc-76(e911) V; dyn-1(n4039) X; enEx21[Pdyn-1 dyn-1]	Figure 9 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3015	This study	unc-76(e911) enIs85 V; dyn-1(n4039) X; enEx21	Figure 9 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3485	This study	unc-76(e911)V; enEx1791 [punc-76(+), Pced-1 mNG::LGG-1, Pced nuc-1::mCherry]	Figure 12 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3489	This study	rab-7 (ok511) II / mIn1 II; unc-76(e911) V; enEx1705 [punc-76(+), Pced-1 CED-1::GFP, Pced-1 2xFYVE::mRFP]	Figure 6 Available from the Zhou Lab
Strain, strain background (C. elegans)	ZH3492	This study	atg-7(bp411) IV; unc-76(e911) V, enEx979	Figure 13 Available from the Zhou Lab

991
992

Mutations, strains, and transgenic arrays

C. elegans strains were grown at 20ºC as previously described [66] unless indicated otherwise. The N2

Bristol strain was used as the wild-type control strain. Mutations are described in [67] and by the

Wormbase (http://www.wormbase.org) unless noted otherwise (**Key Resources Table**): LG1: *ced-*

1(e1735), epg-8(bp251); LGII: *lgg-1(tm3489), rab-7(ok511), vps-18(tm1126)*; LGIII: *atg-13(bp414),*

ced-6(n2095), cup-5(n3264); LGIV: *atg-3(bp412), atg-7(bp411), ced-5(n1812), ced-10(n1993), lgg-*

2(tm5755 and *tm6474)*; LGV: *atg-9(bp564), atg-18(gk378), unc-76(e911), unc-51(e369)*; LGX: *atg-*

2(bp576), dyn-1(n4039). *dyn-1(n4039)* homozygous mutants, which are zygotic embryonic lethal, were

maintained by an extrachromosomal array carrying a wild-type *dyn-1* gene and a co-expressed P$_{egl-13}$*gfp*

marker (**Key Resources Table**) [22]. *dyn-1(n4039)* homozygous embryos losing the rescuing transgene

were identified as the embryos not carrying P$_{egl-13}$*gfp*. The *rab-7(ok511) and lgg-1(tm3489)*

homozygous strains are both maternal-effect embryonic lethal, and the mIn1 balancer maintained each

allele with an integrated pharyngeal GFP marker (**Key Resources Table**) [68]. To obtain *rab-7(ok511)*

m⁻z⁻ homozygous embryos, GFP⁻ *rab-7(ok511)* homozygous hermaphrodites were isolated among the

progeny of the strain VC308, and their progeny were collected as embryos. The same protocol was used

to collect *lgg-1(tm3489)* m⁻z⁻ embryos from the strain GK738 (**Key Resources Table**). Double mutants

between *lgg-1(tm3489)/mIn1* and *lgg-2(tm6474)* were generated by standard genetic crosses.

Extrachromosomal arrays were generated by the microinjection of plasmids with the co-injection

marker p76-18B [*punc-76(+)*] into *unc-76(e911)* mutants [69,70]. Non-Unc mutants were identified as

transgenic animals. Integrated transgenic arrays were generated by gamma irradiation [70]. Integrated

arrays generated in this study are as follows (**Key Resources Table**): LGI: *enIs87[P$_{ced-}$*

$_1$PH(hPLCγ)::mrfp and P$_{ced-1}$mNG::lgg-1]; LGII: enIs82[P$_{ced-1}$ ced-1::gfp and P$_{ced-1}$ mCherry::lgg-1];

LGV: *enIs85[P$_{ced-1}$ PH(hPLCγ)::mrfp and P$_{ced-1}$ mNG::lgg-2]; LGX: enIs83[P$_{ced-1}$ ced-1::gfp and P$_{ced-1}$*

mCherry::lgg-2].

1017

Plasmid construction

lgg-1 and *lgg-2* cDNAs were PCR amplified from a mixed-stage *C. elegans* cDNA library [49].

To generate the P$_{ced-1}$ *gfp::lgg*-1 and P$_{ced-1}$ *gfp::lgg-2* plasmids, *lgg-1* and *lgg-2* cDNAs were cloned into

the XmaI and KpnI sites of plasmid pZZ956 (P$_{ced-1}$ *5'gfp*) [49]. P$_{ced-1}$ *mCherry::lgg-1 or -2* were

constructed by replacing the cDNA of *gfp* with that of mCherry [31]. P$_{ced-1}$ *mNG (mNeonGreen)::lgg-1*

or -2 were constructed by replacing the cDNA of *gfp* with that of mNeonGreen [71]. To generate P$_{ced-}$

$_1$gfp::lgg-1(G116A) and P$_{ced-1}$*gfp::lgg-2(G130A),* the QuickChange Site-directed Mutagenesis Kit

(Stratagene, La Jolla, CA) was used to introduce the above mutations into the constructs. To construct

P$_{ced-1}$ *atg-9::mCherry, the atg-9a* open reading frame was PCR-amplified from a *C. elegans* mixed-stage

cDNA library and inserted between P$_{ced-1}$ and *gfp* in pZZ829 (P$_{ced-1}$ *3'gfp*) [49]. The *gfp* cDNA was then

replaced by the *mCherry* cDNA [31]. All plasmids contain an *unc-54* 3' UTR.

1029

Transmission electron microscopy.

In a mixed population of the strain VC308, isolate mid-L4 stage hermaphrodite larvae that were rab-

7(ok511) and place them in a new plate. Forty-eight hrs later, these worms, which are adults, were

processed for transmission electron microscopic studies as previously described [20,22]. Cross-sections

of the gonadal region were generated. In 50-nm thin sections, germ cell corpses and the neighboring

gonadal sheath cells were identified as previously described. A phagosome containing a germ cell

corpses was identified in serial 50-nm thin sections that cover the entire length of each germ cell corpse

were analyzed by determining whether a germ-cell corpse is entirely inside the vacuole inside the

gonadal sheath cells. Phagosomal surfaces were carefully monitored in a thin section electron

micrograph for the presence of double-membrane vesicles.

1040

Quantification of the number of cell corpses using Nomarski DIC microscopy

Cell corpses display a highly refractive button-like morphology under Differential Interference

Contrast (DIC) microscopy. An Axionplan 2 compound microscope (Carl Zeiss, Thornwood, NY)

equipped with Nomarski DIC optics, an AxioCam digital camera, and AxioVision imaging software was

used for DIC microscopy. Using a previously established protocol [28], we quantified the number of

cell corpses in the head region at the 1.5-fold and 2-fold stage embryos, which are ~420 and ~460 min

post-first cleavage.

1048

Fluorescence microscopy and time-lapse imaging

A DeltaVision Elite Deconvolution Imaging System (GE Healthcare, Inc.) equipped with a DIC

imaging apparatus and a Photometrics Coolsnap 2 digital camera was used to capture fluorescence, and

DIC images Applied Precision SoftWoRx 5.5 software was utilized for deconvolving and analyzing the

images [28]. To observe the amount of autophagosomes in ventral hypodermal cells that express P_{ced-1},

fourteen serial Z-sections in 0.5 μm interval between adjacent optical sections, starting at the ventral

surface of embryos at mid-embryonic stages, were collected. The 2D projection image of each Z-stack

was generated and compared among different genetic backgrounds. To track fluorescence markers on

pseudopods, on the surfaces of phagosomes, or inside phagosomal lumen during the clearance process of

cell corpses C1, C2, and C3, embryos were monitored on their ventral surface starting at ~310 min post-

first cleavage using an established time-lapse imaging protocol [28]. Twelve to 16 serial Z-sections (at

0.5-μm intervals) were captured every 2 min, with recordings typically lasting between 60 to 180 min.

Embryos that exhibited normal elongation and movement were considered developing properly. The

moment engulfment starts is defined as when the extension of pseudopods around C1, C2, or C3 is first

observed. The moment a nascent phagosome form is defined as when the pseudopods around a cell

corpse join and make a full closure. The life span of a phagosome is defined as the time interval

between the moments when the nascent phagosome forms and when the phagosome shrinks to one-half

of its original diameter.

The time spans of the engulfment and degradation processes of cell corpses C1, C2, or C3 were

measured as previously established [49]. Briefly, a pseudopod marker, either CED-1::GFP or

PH(PLCγ)::GFP, was monitored over time. The moment engulfment starts is defined as when the

budding pseudopods around C1, C2, or C3 is first observed. The moment a nascent phagosome form is

defined as the moment when the pseudopods around a cell corpse join and make a full closure. The

period between the budding and the sealing of the pseudopods is the time span of engulfment. To

measure phagosome duration, a co-expressed phagosomal surface marker mCherry::2xFYVE was used

to track the diameter of the phagosome over time. The life span of a phagosome is defined as the time

interval between the time points when a nascent phagosome was born and when the phagosome shrank

to one-half of its original diameter.

To monitor the degradation process of the chromatin DNA of apoptotic cells inside phagosomes,

we measured the size of the nucleus of engulfed C1, C2, or C3 over time by following the ubiquitously

expressed P_{his-72} *his-72::mCherry* reporter. Briefly, the "0 min" timepoint of phagosome formation was

determined by the appearance of the "bottom-like" structure under DIC optics. The nuclear diameter of

each phagosome was measured as the diameter of the $mCherry^+$ disc inside the phagosome. The

measurement is performed in regular time intervals until 60-min after the phagosome formation.

1083

Quantitative measurement of signal intensity

Measuring the signal intensity inside the phagosomal lumen

In embryos expressing mCherry- or mNG-tagged LGG-1 or LGG-2, to measure the fluorescence

signal intensity inside the phagosomal lumen over time, we identified the boundary of a phagosome and

the "0 min" time point when a nascent phagosome was formed by observing the co-expressed marker for

a nascent phagosome such as CED-1::GFP or PH(PLCγ)::GFP. At each time point, the total LGG-1 or -

2 image intensity of a fixed area (4x4 pixels) at the center of a phagosome ($Int_{phagosome}$) was recorded

(**Fig 1C**), so was the intensity of an area of the same size (4x4 pixels) outside the embryo as the

background image intensity ($Int_{background}$). The Relative image intensity (RInt) at a particular time point

(Tn) comparing to the start point (T0) is calculated as $RInt_{Tn} = (Int_{phagosome}-Int_{background})_{Tn} / (Int_{phagosome}-$

$Int_{background})_{T0}$. The $RInt_{Tn}$ value of 1.0 indicates no entry of LGG-1- or LGG-2-labeled autophagosomes

into the phagosomal lumen.

1096

Measuring the signal intensity on the surface of a phagosome

To measure the efficiency of recruitment of autophagosomes to phagosomes, we quantified the

intensity of mCherry- or mNG-labeled LGG-1 or LGG-2 on the surfaces of phagosomes. First, we

identified the boundary of a phagosome and the "0 min" time point when a nascent phagosome just

formed by observing co-expressed CED-1::GFP or PH(PLCγ)::GFP. At a particular time point, Tn or

T0, the surface of a phagosome was outlined by two closed polygons (**Fig 8E**). The total signal

intensities, as well as the areas of the polygons, were recorded. The unit signal intensity of the "donut-

shape" area between the two polygons was calculated as follows:

Unit Intensity ($UI_{phagosome}$) = ($Intensity_{external\ polygon}$ − $Intensity_{internal\ polygon}$)/($Area_{external\ polygon}$ − $Area_{internal}$

$_{polygon}$). The Unit Background Intensity ($UI_{background}$) was measured from a polygon outside the embryo

was calculated as follow: $UI_{background}$ = $Intensity_{background}$/$Area_{background}$.

At the time point Tn, the relative signal intensity ($RInt_{Tn}$) = ($UI_{phagosome}$ − $UI_{background}$)$_{Tn}$ / ($UI_{phagosome}$ −

$UI_{background}$)$_{T0}$. The $RInt_{Tn}$ value 1.0 indicates no enrichment of LGG-1- or LGG-2-labeled

autophagosomes on phagosomal surfaces comparing to the "0 min" time point.

1111

Measuring the acidification index of a phagosome

In embryos expressing P$_{his-72}$ *his-72::gfp::mCherry,* we quantified the fluorescence intensity of

the mCherry and GFP signals at the center of phagosomes in regular time intervals. The "0 min"

timepoint of phagosome formation was determined by the appearance of the "bottom-like" structure

under DIC optics. At each time point, the total mCherry and GFP signal intensity of a fixed area (3x3

pixels) at the center of a phagosome ($Int_{mCherry}$, Int_{GFP}) was recorded. The acidification index (A_{idx}) at a

particular time point (Tn) is defined as: A_{idx} = (Int_{GFP}/$Int_{mCherry}$)$_{Tn}$ / (Int_{GFP}/$Int_{mCherry}$)$_{T0}$. The A_{idx} value

1.0 indicates no phagosome acidification compared to the "0 min" value.

1120

Quantifying the percentage of autolysosomes among the LGG-1/or-2-labeled vesicles

In wild-type embryos co-expressing either mNG::LGG-1 or mNG:: LGG-2 together with NUC-

1::mCherry (mCh), we counted the numbers of mNG$^+$ mCherry$^-$ puncta and the mNG$^+$ mCherry$^+$ puncta

observed on the surfaces of phagosomes. For each set of reporters, C1, C2, and C3 phagosomes were

quantified in each embryo, and the numbers of puncta were added. Seven embryos were scored (n=7),

each representing one sample. The percentage of autolysosomes (%$_{AL}$) for each embryo was calculated

as follow: %$_{AL}$= 100*((# of mNG$^+$ mCh$^+$ puncta)/ [(# of mNG$^+$ mCh$^+$) + (# of mNG$^+$ mCh$^-$)]. The

average value and standard deviation were calculated and reported.

1129

Quantifying the percentage of LGG-labeled puncta that are also labeled with RAB-7

In wild-type embryos co-expressing P$_{ced-1}$ *mCherry-lgg-1* or *-lgg-2* together with P$_{ced-1}$ *gfp::rab-7*,

we counted the numbers of mCh$^+$ GFP$^-$ and mCh$^+$ GFP$^+$ puncta in the engulfing cells for C1, C2, and C3

in two different locations: (1) on the surfaces of the phagosomes, and (2) in the cytoplasm. For each set

of co-expressed reporters, 9 engulfing cells were scored. N_{mCh+} (the total number of mCh$^+$ puncta) =

# of mCh$^+$ GFP$^-$ puncta + # of mCherry$^+$ GFP$^+$ puncta. % mCh$^+$ GFP$^+$ = 100 * (# mCh+ GFP+ puncta)/

N_{mCh+}. The average values and standard deviation values were calculated and reported.

1137

1138

1139

Acknowledgment

We thank Ying Wang and Lidan Gao for technical support and R. Haley for helpful comments.

We thank H. Robert Horvitz for advice and the use of the electron microscope. We thank Erika

Hartwieg for the technical support for the TEM study. We thank the C. elegans Genetics Center (CGC),

funded by the NIH Office of Research Infrastructure Programs (P40 OD010440), for providing some

strains. We also thank the National BioResource Project of Japan and Dr. Shohei Mitani for providing

some strains.

1147

1148 Funding Source

1149 NIH R01GM067848 supports this work.

1150

1151 Competing Interests

1152   The authors declare no competing interests

1153

1154

References

1. Reddien PW, Horvitz HR (2004) The engulfment process of programmed cell death in
caenorhabditis elegans. Annu Rev Cell Dev Biol 20: 193-221.
2. Nagata S (2018) Apoptosis and Clearance of Apoptotic Cells. Annu Rev Immunol.
3. Levin R, Grinstein S, Canton J (2016) The life cycle of phagosomes: formation, maturation, and
resolution. Immunol Rev 273: 156-179.
4. Schaaf MB, Keulers TG, Vooijs MA, Rouschop KM (2016) LC3/GABARAP family proteins:
autophagy-(un)related functions. FASEB J 30: 3961-3978.
5. Sanjuan MA, Dillon CP, Tait SW, Moshiach S, Dorsey F, et al. (2007) Toll-like receptor signalling
in macrophages links the autophagy pathway to phagocytosis. Nature 450: 1253-1257.
6. Green DR, Oguin TH, Martinez J (2016) The clearance of dying cells: table for two. Cell Death
Differ 23: 915-926.
7. Martinez J, Almendinger J, Oberst A, Ness R, Dillon CP, et al. (2011) Microtubule-associated
protein 1 light chain 3 alpha (LC3)-associated phagocytosis is required for the efficient
clearance of dead cells. Proc Natl Acad Sci U S A 108: 17396-17401.
8. Martinez J, Cunha LD, Park S, Yang M, Lu Q, et al. (2016) Noncanonical autophagy inhibits the
autoinflammatory, lupus-like response to dying cells. Nature 533: 115-119.
9. Morishita H, Mizushima N (2019) Diverse Cellular Roles of Autophagy. Annu Rev Cell Dev Biol
35: 453-475.
10. Nakatogawa H (2020) Mechanisms governing autophagosome biogenesis. Nat Rev Mol Cell
Biol 21: 439-458.
11. Cheng S, Wu Y, Lu Q, Yan J, Zhang H, et al. (2013) Autophagy genes coordinate with the class II
PI/PtdIns 3-kinase PIKI-1 to regulate apoptotic cell clearance in C. elegans. Autophagy 9:
2022-2032.
12. Huang S, Jia K, Wang Y, Zhou Z, Levine B (2013) Autophagy genes function in apoptotic cell
corpse clearance during C. elegans embryonic development. Autophagy 9: 138-149.
13. Li W, Zou W, Yang Y, Chai Y, Chen B, et al. (2012) Autophagy genes function sequentially to
promote apoptotic cell corpse degradation in the engulfing cell. J Cell Biol 197: 27-35.
14. Martinez J, Malireddi RK, Lu Q, Cunha LD, Pelletier S, et al. (2015) Molecular characterization
of LC3-associated phagocytosis reveals distinct roles for Rubicon, NOX2 and autophagy
proteins. Nat Cell Biol 17: 893-906.
15. Sulston JE, Horvitz HR (1977) Post-embryonic cell lineages of the nematode, Caenorhabditis
elegans. Dev Biol 56: 110-156.
16. Sulston JE, Albertson DG, Thomson JN (1980) The Caenorhabditis elegans male:
postembryonic development of nongonadal structures. Dev Biol 78: 542-576.
17. Ellis RE, Jacobson DM, Horvitz HR (1991a) Genes required for the engulfment of cell corpses
during programmed cell death in Caenorhabditis elegans. Genetics 129: 79-94.
18. Gumienny TL, Lambie E, Hartwieg E, Horvitz HR, Hengartner MO (1999) Genetic control of
programmed cell death in the *Caenorhabditis elegans* hermaphrodite germline.
Development 126: 1011-1022.
19. Mangahas PM, Zhou Z (2005) Clearance of apoptotic cells in Caenorhabditis elegans. Semin Cell
Dev Biol 16: 295-306.
20. Yu X, Lu N, Zhou Z (2008) Phagocytic receptor CED-1 initiates a signaling pathway for
degrading engulfed apoptotic cells. PLoS Biol 6: e61.

21. Zhou Z, Hartwieg E, Horvitz HR (2001b) CED-1 is a transmembrane receptor that mediates cell
corpse engulfment in C. elegans. Cell 104: 43-56.
22. Yu X, Odera S, Chuang CH, Lu N, Zhou Z (2006) C. elegans Dynamin mediates the signaling of
phagocytic receptor CED-1 for the engulfment and degradation of apoptotic cells. Dev Cell
10: 743-757.
23. Shen Q, He B, Lu N, Conradt B, Grant BD, et al. (2013) Phagocytic receptor signaling regulates
clathrin and epsin-mediated cytoskeletal remodeling during apoptotic cell engulfment in C.
elegans. Development 140: 3230-3243.
24. Lu N, Shen Q, Mahoney TR, Neukomm LJ, Wang Y, et al. (2012) Two PI 3-kinases and one PI 3-
phosphatase together establish the cyclic waves of phagosomal PtdIns(3)P critical for the
degradation of apoptotic cells. PLoS Biol 10: e1001245.
25. Lu N, Shen Q, Mahoney TR, Liu X, Zhou Z (2011) Three sorting nexins drive the degradation of
apoptotic cells in response to PtdIns(3)P signaling. Mol Biol Cell 22: 354-374.
26. Lu N, Zhou Z (2012) Membrane trafficking and phagosome maturation during the clearance of
apoptotic cells. Int Rev Cell Mol Biol 293: 269-309.
27. Manil-Segalen M, Lefebvre C, Jenzer C, Trichet M, Boulogne C, et al. (2014) The C. elegans LC3
acts downstream of GABARAP to degrade autophagosomes by interacting with the HOPS
subunit VPS39. Dev Cell 28: 43-55.
28. Lu N, Yu X, He X, Zhou Z (2009) Detecting apoptotic cells and monitoring their clearance in the
nematode Caenorhabditis elegans. Methods Mol Biol 559: 357-370.
29. Wu F, Watanabe Y, Guo XY, Qi X, Wang P, et al. (2015) Structural Basis of the Differential
Function of the Two C. elegans Atg8 Homologs, LGG-1 and LGG-2, in Autophagy. Mol Cell 60:
914-929.
30. Tsien RY (1998) The green fluorescent protein. Annu Rev Biochem 67: 509-544.
31. Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, et al. (2004) Improved
monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red
fluorescent protein. Nat Biotechnol 22: 1567-1572.
32. Shinoda H, Ma Y, Nakashima R, Sakurai K, Matsuda T, et al. (2018) Acid-Tolerant Monomeric
GFP from Olindias formosa. Cell Chem Biol 25: 330-338 e337.
33. Tian Y, Li Z, Hu W, Ren H, Tian E, et al. (2010) C. elegans screen identifies autophagy genes
specific to multicellular organisms. Cell 141: 1042-1055.
34. Yang P, Zhang H (2011) The coiled-coil domain protein EPG-8 plays an essential role in the
autophagy pathway in C. elegans. Autophagy 7: 159-165.
35. Zhang H, Baehrecke EH (2015) Eaten alive: novel insights into autophagy from multicellular
model systems. Trends Cell Biol.
36. Gomes LC, Odedra D, Dikic I, Pohl C (2016) Autophagy and modular restructuring of
metabolism control germline tumor differentiation and proliferation in C. elegans.
Autophagy 12: 529-546.
37. Lin L, Yang P, Huang X, Zhang H, Lu Q, et al. (2013) The scaffold protein EPG-7 links cargo-
receptor complexes with the autophagic assembly machinery. J Cell Biol 201: 113-129.
38. Lu Q, Yang P, Huang X, Hu W, Guo B, et al. (2011) The WD40 repeat PtdIns(3)P-binding protein
EPG-6 regulates progression of omegasomes to autophagosomes. Dev Cell 21: 343-357.
39. Kovacs AL, Vellai T, Muller F (2013) Autophagy in Caenorhabditis elegans. In: Klionsky DJ,
editor. Autophagy: CRC Press.
40. Zhang H, Chang JT, Guo B, Hansen M, Jia K, et al. (2015) Guidelines for monitoring autophagy in
Caenorhabditis elegans. Autophagy 11: 9-27.

41. Wu F, Wang P, Shen Y, Noda NN, Zhang H (2016) Small differences make a big impact:
Structural insights into the differential function of the 2 Atg8 homologs in C. elegans.
Autophagy 12: 606-607.
42. Stenmark H (2009) Rab GTPases as coordinators of vesicle traffic. Nat Rev Mol Cell Biol 10:
513-525.
43. Szatmari Z, Sass M (2014) The autophagic roles of Rab small GTPases and their upstream
regulators: a review. Autophagy 10: 1154-1166.
44. Gao J, Langemeyer L, Kümmel D, Reggiori F, Ungermann C (2018) Molecular mechanism to
target the endosomal Mon1-Ccz1 GEF complex to the pre-autophagosomal structure. Elife 7.
45. Hegedűs K, Takáts S, Boda A, Jipa A, Nagy P, et al. (2016) The Ccz1-Mon1-Rab7 module and
Rab5 control distinct steps of autophagy. Mol Biol Cell 27: 3132-3142.
46. Vaites LP, Paulo JA, Huttlin EL, Harper JW (2018) Systematic Analysis of Human Cells Lacking
ATG8 Proteins Uncovers Roles for GABARAPs and the CCZ1/MON1 Regulator
C18orf8/RMC1 in Macroautophagic and Selective Autophagic Flux. Mol Cell Biol 38.
47. Balderhaar HJ, Ungermann C (2013) CORVET and HOPS tethering complexes - coordinators of
endosome and lysosome fusion. J Cell Sci 126: 1307-1316.
48. Xiao H, Chen D, Fang Z, Xu J, Sun X, et al. (2009) Lysosome biogenesis mediated by vps-18
affects apoptotic cell degradation in Caenorhabditis elegans. Mol Biol Cell 20: 21-32.
49. Haley R, Wang Y, Zhou Z (2018) The small GTPase RAB-35 defines a third pathway that is
required for the recognition and degradation of apoptotic cells. PLoS Genet 14: e1007558.
50. Wu YC, Stanfield GM, Horvitz HR (2000) NUC-1, a caenorhabditis elegans DNase II homolog,
functions in an intermediate step of DNA degradation during apoptosis. Genes Dev 14: 536-
548.
51. Guo P, Hu T, Zhang J, Jiang S, Wang X (2010) Sequential action of Caenorhabditis elegans Rab
GTPases regulates phagolysosome formation during apoptotic cell degradation. Proc Natl
Acad Sci U S A 107: 18016-18021.
52. Ooi SL, Priess JR, Henikoff S (2006) Histone H3.3 variant dynamics in the germline of
Caenorhabditis elegans. PLoS Genet 2: e97.
53. Treusch S, Knuth S, Slaugenhaupt SA, Goldin E, Grant BD, et al. (2004) Caenorhabditis elegans
functional orthologue of human protein h-mucolipin-1 is required for lysosome biogenesis.
Proc Natl Acad Sci U S A 101: 4483-4488.
54. Campbell EM, Fares H (2010) Roles of CUP-5, the Caenorhabditis elegans orthologue of human
TRPML1, in lysosome and gut granule biogenesis. BMC Cell Biol 11: 40.
55. Qu X, Zou Z, Sun Q, Luby-Phelps K, Cheng P, et al. (2007) Autophagy gene-dependent clearance
of apoptotic cells during embryonic development. Cell 128: 931-946.
56. Jenzer C, Simionato E, Largeau C, Scarcelli V, Lefebvre C, et al. (2019) Autophagy mediates
phosphatidylserine exposure and phagosome degradation during apoptosis through
specific functions of GABARAP/LGG-1 and LC3/LGG-2. Autophagy 15: 228-241.
57. Fazeli G, Stetter M, Lisack JN, Wehman AM (2018) C. elegans Blastomeres Clear the Corpse of
the Second Polar Body by LC3-Associated Phagocytosis. Cell Rep 23: 2070-2082.
58. Fazeli G, Trinkwalder M, Irmisch L, Wehman AM (2016) C. elegans midbodies are released,
phagocytosed and undergo LC3-dependent degradation independent of macroautophagy. J
Cell Sci 129: 3721-3731.
59. Vieira OV, Botelho RJ, Grinstein S (2002) Phagosome maturation: aging gracefully. Biochem J
366: 689-704.
60. Spang A (2016) Membrane Tethering Complexes in the Endosomal System. Front Cell Dev Biol
4: 35.

61. Kinchen JM, Doukoumetzidis K, Almendinger J, Stergiou L, Tosello-Trampont A, et al. (2008) A
pathway for phagosome maturation during engulfment of apoptotic cells. Nat Cell Biol 10:
556-566.
62. Jiang P, Nishimura T, Sakamaki Y, Itakura E, Hatta T, et al. (2014) The HOPS complex mediates
autophagosome-lysosome fusion through interaction with syntaxin 17. Mol Biol Cell 25:
1327-1337.
63. Takáts S, Pircs K, Nagy P, Varga Á, Kárpáti M, et al. (2014) Interaction of the HOPS complex
with Syntaxin 17 mediates autophagosome clearance in Drosophila. Mol Biol Cell 25: 1338-
1354.
64. Stroupe C, Collins KM, Fratti RA, Wickner W (2006) Purification of active HOPS complex
reveals its affinities for phosphoinositides and the SNARE Vam7p. EMBO J 25: 1579-1589.
65. Jeschke A, Haas A (2018) Sequential actions of phosphatidylinositol phosphates regulate
phagosome-lysosome fusion. Mol Biol Cell 29: 452-465.
66. Wood WB, Researchers tCo*Ce* (1988) The Nematode *Caenorhabditis elegans*. Cold Spring
Harbor, NY: Cold Spring Harbor Laboratory.
67. Riddle DL, Blumenthal T, Meyer BJ, Priess JR, editors (1997) C. elegans II. Plainview, NY: Cold
Spring harbor Laboratory Press.
68. Edgley ML, Baillie DL, Riddle DL, Rose AM (2006) Genetic balancers. WormBook: 1-32.
69. Bloom L, Horvitz HR (1997) The Caenorhabditis elegans gene unc-76 and its human homologs
define a new gene family involved in axonal outgrowth and fasciculation. Proc Natl Acad Sci
U S A 94: 3414-3419.
70. Jin Y (1999) Transformation. In: Hope IA, editor. C elegans, a practical approach. Oxford:
Oxford University Press. pp. 69-96.
71. Shaner NC, Lambert GG, Chammas A, Ni Y, Cranfill PJ, et al. (2013) A bright monomeric green
fluorescent protein derived from Branchiostoma lanceolatum. Nat Methods 10: 407-409.
1317
1318

1320

1321
Figure 1 – figure supplement 1. The fusion of lysosomal particles to phagosomes results in the incorporation of
lysosomal membrane protein CTNS-1 to the phagosomal membrane but not the lumen
(A) Time-lapse images of a C2 phagosome following the lysosomal recruitment and fusion. CTNS-1::mRFP is expressed in
engulfing cells under P_{ced-1}. "0 in" is the time point when a C2 phagosome (white arrowheads) just forms. Yellow arrows mark
CTNS-1::mRFP-labeled lysosomal particles. One open white arrow marks the phagosomal surface that is evenly colored by
mRFP. Scale bars are 2μm.
(B) A diagram illustrating that the single membrane-vesicles labeled with CTNS-1::mRFP on their membranes are recruited to
phagosomal surfaces and fused to the phagosomal membrane. After the fusion between the membranes of these vesicles
and the phagosome, the mRFP signal is evenly distributed to the phagosomal surface. However, no mRFP signal enters the
phagosomal lumen.

1332

1333
Figure 1 – figure supplement 2. Besides the phagosomes containing C1, C2, and C3, the attachment and fusion of
LGG+-vesicles to other phagosomes are also observed during embryonic development
Wild-type embryos co-expressing P*ced-1* *ced-1::gfp* and P*ced-1* *mCherry::lgg-1* or P*ced-1* *mCherry::lgg-2* reporters were analyzed.
(A-B) Time-lapse images following the indicated reporters in C4 (A) or C5 (B) phagosomes
starting when the nascent phagosome (white arrowheads) just formed (time point "0 min"). Scale bars are 2 μm. Yellow
arrows mark the LGG+ puncta on phagosome surfaces. Open white arrows mark when mCherry signal is detected in the
phagosomal lumen, indicating autophagosome/phagosome fusion events.
(C) A diagram illustrating the two phagosomes that contain cell corpses C4 and C5, with which we monitor the dynamic
recruitment and fusion of autophagosomes, at ~310 min (A) and ~280 min (B) post-the 1st embryonic division. Both the
positions of C4 and C5 (brown dots) and the identities of their engulfing cells are shown.
(D-E) Images of wild-type embryos illustrate the positions of the C4 and C5 phagosomes shown in (A) and (B), respectively.
Scale bars are 5 μm.
(F) Images of two wild-type embryos at a 1.5-fold developmental stage. White arrowheads (a and c) indicate phagosomes in
which the mCherry+ puncta are observed to be recruited to their surfaces (yellow arrows in (b and d)), and that mCherry signal
is detected in the phagosomal lumen (open white arrows in (b and d)). Scale bars are 5 μm.
1349

1350

1351
Figure 2 – figure supplement 1 The *atg-7*, *atg-13*, and *epg-8* mutants are severely defective in the production of LGG-
1-labeled autophagosomes
Images of wild-type (A), *atg-7(bp411)* (B), *atg-13(bp414)* (C), and *epg-8(bp251)* mutant (D) embryos expressing P$_{ced-1}$
mCherry::lgg-1 are presented. "0 min" labels embryos that are ~330 min post-1st cleavage. The mCherry images are 2-D
projections of 14 Z-sections at 0.5 μm intervals. Yellow arrows mark LGG-1 autophagosomes. White arrows in (A(d)) mark
three phagosomes, on the surface of which mCherry signal is recruited in (a-b), and 20 min later, enters the phagosome,
demonstrating the autophagosomes/phagosome fusion event. Scale bars are 5 μm.
1359

1360
**Figure 2 – figure supplement 2. The *atg-7*, *atg-13*, and *epg-8* mutants are severely defective in the production of LGG-
2-labeled autophagosomes**
Images of wild-type (A), *atg-7(bp411)* (B), *atg-13(bp414)* (C), and *epg-8(bp251)* mutant (D) embryos expressing P$_{ced-1}$
mCherry::lgg-2 are presented. "0 min" labels embryos that are ~330 min post-1st cleavage. The mCherry images are 2-D
projections of 14 Z-sections at 0.5 µm intervals. Yellow arrows mark LGG-2 autophagosomes. White arrows in (A(b, d, and f))
mark phagosomes in which mCherry signal was detected, indicating the existence of autophagosomes/phagosome fusion
events. Scale bars are 5 µm.
1368

1369
1370 **Figure 3 – figure supplement 1.** ***atg-9*** **is another gene essential for the production of autophagosomes and the**
1371 **incorporation of autophagosomes into phagosomes**
1372 Images of *atg-9(bp564)* mutant embryos expressing P*ced-1 mNG::lgg-1* (A) or *::lgg-2* (B) are presented. "0 min" embryos are
1373 embryos at ~330 min post-1st cleavage. The mNG images are 2-D projections of 14 Z-sections at 0.5 µm intervals. Green
1374 puncta (yellow arrows) represent LGG-1 or LGG-2 autophagosomes. Scale bars are 5 µm.
1375

1376
1377
Figure 5 – figure supplement 1. The expression of *lgg-1* and *lgg-2* cDNA in engulfing cells suppresses the Ced
phenotype of *lgg-1* and *lgg-2* null mutants, respectively.
GFP- and mCherry-tagged reporters for LGG-1 and LGG-2 are expressed in engulfing cells under the control of the P$_{ced-1}$
promoter in *lgg-1(tm3489)* and *lgg-2(tm5755)* mutant embryos, respectively. The bar graph displays the average number of
somatic cell corpses in *lgg-1(tm3489)* and *lgg-2(tm5755)* mutant embryos at the 1.5-fold stage, both with and without
expression rescuing construct. The number of embryos scored for each strain is in parentheses. Bars represent the mean,
and the error bars indicate standard deviation. Brackets above the bars indicate the samples compared by the Student *t*-test.
p-values are summarized as such: *, 0.001 < p < 0.05; **, 0.0001 < p <0.001; ***, p <0.0001; ns, no significant difference.
(Figure 5 – figure supplement 1 – source data 1)
1387

1388
Figure 9 – figure supplement 1. The generation of autophagosomes is normal in *ced-1* mutants.
Images of wild-type and *ced-1(e1735)* mutant embryos expressing P*ced-1* *mNG::lgg-1* (A) or P*ced-1lgg-2::mNG* (B) are presented.
(a-b and e-f) Embryos are at ~370 min post-1st cleavage; (c-d and g-h) Embryos 20 min later. (a, c, e, g) DIC images. (b, d, f,
h) 2-D projection of 14 consecutive Z-sections at 0.5 µm interval each. Green puncta represent LGG autophagosomes (yellow
arrows). White arrows in (A(a-d)) marks one C3 phagosome, onto which LGG-1::mNG puncta are recruited in (a-b); 20 min
later (c-d), LGG-1::mNG enters this phagosome, showing an autophagosomes/phagosome fusion event. Yellow arrowheads
in (A(e-h)) marks one C3 phagosome on the surface, of which only one mNG punctum was found (f), and no mNG signal
enters the phagosome (h), indicating the lack of recruitment and fusion. Yellow arrowhead in (B(g-h)) marks one C2
phagosome on the surface, of which one mNG[+] punctum is found, and no fusion is visible at 20 min. Scale bars are 5 µm.
1398

1399

Figure 1-video 1. mCherry::LGG-1-labeled vesicles are recruited to the surface of a phagosome

and subsequently fuse to the phagosome

Related to **Fig 1E**. This movie shows time-lapse recording images of a C3 phagosome (white

arrowheads) in a wild-type embryo, starting at ~330 min post-1st cleavage. CED-1::GFP labels

pseudopods, allowing the visualization of the phagosome formation process. "0 min" is the moment

when pseudopods just seal. Yellow arrowheads indicate the mCherry::LGG-1^{+} puncta, and open white

arrows point to the phagosome lumen containing the mCherry signal.

1408

Figure 1-video 2. mCherry::LGG-2-labeled vesicles are recruited to the surface of a phagosome

and subsequently fuse to the phagosome

Related to **Fig 1H**. This movie shows time-lapse recording images of a C3 phagosome (white

arrowheads) in a wild-type embryo, starting at ~330 min post-1st cleavage. CED-1::GFP labels

pseudopods, allowing the visualization of the phagosome formation process. "0 min" is the moment

when pseudopods just seal. Yellow arrowheads indicate the mCherry::LGG-2^{+} puncta, and open white

arrows point to the phagosome lumen containing the mCherry signal.

1416

Figure 1-video 3. mNG::LGG-1-labeled vesicles are recruited to the surface of a phagosome and

subsequently fuse to the phagosome.

Related to **Fig 1F**. This movie shows time-lapse recording images of a C3 phagosome in a wild-type

embryo, starting at ~330 min post-1st cleavage. PH(hPLCγ)::mRFP labels pseudopods, allowing the

visualization of the phagosome formation process. "0 min" is the moment when pseudopods just seal.

Yellow arrowheads indicate mNG::LGG-1^{+} puncta, and open white arrows point to the phagosome with

mNG signal in its lumen.

1424

Figure 8-video 1. In *rab-7 (ok511)*, mCherry::LGG-1-labeled vesicles fail to fuse with

phagosomes.

Related to **Fig 8G**. This movie shows time-lapse recording images of a C3 phagosome in a *rab-*

7(ok511) m^{-}z^{-} homozygous embryo, starting at ~330 min post-1st cleavage. "0 min" is the moment when

a nascent phagosome is just formed. White arrowheads mark the phagosome. Yellow arrows mark

mCherry::LGG-1^{+} puncta.

1431

Figure 8-video 2. In *rab-7(ok511)* mutants, mCherry::LGG-2-labeled vesicles fail to fuse with

phagosomes.

Related to **Fig 8H**. This movie shows time-lapse recording images of a C3 phagosome in a *rab-*
7(ok511) m⁻z⁻ homozygous embryos, starting at ~330 min post-1st cleavage. "0 min" is the moment
when a nascent phagosome is just formed. White arrowheads mark the phagosome. Yellow arrows
mark mCherry::LGG-2⁺ puncta.

1438

Figure 9-video 1. In *ced-1(e1735)* mutants*,* mNG::LGG-1-labeled vesicles fail to be recruited to
the phagosomal surface.
Related to **Fig 9A**. This movie shows time-lapse recording images of a C3 phagosome in a *ced-*
1(e1735) embryo, starting at ~300 min post-1st cleavage. "0 min" is the moment when a nascent
phagosome is just formed. White arrowheads mark the phagosome; yellow arrows label mNG::LGG-1⁺
puncta.

1445