The M2 Gene Is a Determinant of Reovirus-Induced Myocarditis

Marcelle Dina Zita[1*], Matthew B. Phillips[1*], Johnasha D. Stuart[1*], Asangi R. Kumarapeli[2],

Vijayalakshmi Sridharan [3], Marjan Boerma[3,4], Anthony J. Snyder[5], Pranav Danthi[5], and

Karl W. Boehme[1,5,6‡]

[1]Department of Microbiology and Immunology, [2]Department of Pathology, University of

Arkansas for Medical Sciences, [3]Department of Pharmaceutical Sciences, Division of

Radiation Health, [4]Winthrop P. Rockefeller Cancer Institute, [5]Department of Biology,

Indiana University-Bloomington, and [6]Center for Microbial Pathogenesis and Host

Inflammatory Responses, University of Arkansas for Medical Sciences

* These authors contributed equally to this work.

[‡] Correspondence to Karl W. Boehme

Mailing address: 4301 W. Markham Dr. #511, Little Rock, AR 72205

Phone: (501) 686-5189

Email: kwboehme@uams.edu

23   Running Title: M2 is a determinant of reovirus myocarditis

ABSTRACT

Although a broad range of viruses cause myocarditis, the mechanisms that underlie viral myocarditis are poorly understood. Here, we report that the M2 gene is a determinant of reovirus myocarditis. The reovirus M2 gene encodes outer capsid protein µ1, which influences both cell entry and cell death. We infected newborn C57BL/6 mice with reovirus strain type 1 Lang (T1L) or a reassortant reovirus in which the M2 gene from strain type 3 Dearing (T3D) was substituted into the T1L genetic background (T1L/T3DM2). T1L was non-lethal in wild-type mice, whereas greater than 90% of mice succumbed to T1L/T3DM2 infection. T1L/T3DM2 produced higher viral loads than T1L at the site of inoculation. In secondary organs, T1L/T3DM2 was detected with more rapid kinetics and reached higher peak titers than T1L. We found that hearts from T1L/T3DM2-infected mice were grossly abnormal, with large lesions corresponding to substantial cardiac injury with inflammatory infiltrates. Lesions in T1L/T3DM2-infected mice contained aggregates of necrotic cardiomyocytes with pyknotic debris, and prominent lymphocyte and histiocyte infiltration. In contrast, T1L induced the formation of smaller lesions in a subset of animals, consistent with T1L being mildly myocarditic. Finally, more activated caspase-3-positive cells were observed in hearts from animals infected with T1L/T3DM2 compared to T1L. Together, our findings indicate that substitution of the T3D M2 allele into an otherwise T1L genetic background is sufficient to change a non-lethal infection into a lethal infection. Our results further indicate that T3D M2 enhances T1L replication and dissemination *in vivo*, which potentiates the capacity of reovirus to cause myocarditis.

IMPORTANCE

Reovirus is a non-enveloped virus with a segmented double-stranded RNA genome that serves as a model for studying virus-induced injury to the heart. However, the mechanisms by which reovirus affects heart disease are not fully elucidated. We found that substituting a single gene from reovirus strain type 3 Dearing (T3D) into an otherwise type 1 Lang (T1L) genetic background (T1L/T3DM2) was sufficient to convert the non-lethal T1L strain into a lethal infection in neonatal mice. In comparison to T1L, T1L/T3DM2 reached distal organs such as the heart more efficiently. Furthermore, it replicated to higher levels at those sites. Hearts from mice infected with T1L resulted in limited recruitment of immune cells. In contrast, hearts from mice infected with T1L/T3DM2 contained substantially more immune cell invasion and cell death. Together, our findings identify the reovirus M2 gene as a new determinant of reovirus induced heart disease.

INTRODUCTION

Numerous viruses, such as cytomegalovirus (CMV), coxsackievirus B3 (CVB3),

parvovirus B19, and severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2)

can cause myocarditis, which is characterized by inflammation of the heart muscle.

Myocarditis can lead to cardiac damage, electrophysiological abnormalities, heart

failure, or sudden death (1). Specifically, viral myocarditis is driven by a combination of

virus-induced cytopathic effects in cardiac cells and the host inflammatory response to

infection (2-5). Although a significant public health risk, many open questions remain

concerning the viral factors and host mechanisms that underlie viral myocarditis.

Mammalian orthoreovirus (reovirus) is a non-enveloped, double-stranded RNA

(dsRNA) virus with a segmented genome that serves as a model for studying viral

myocarditis (6). Some, but not all, reovirus strains cause myocarditis in neonatal mice

(7, 8). The segmented nature of the reovirus genome has enabled the identification of

specific genes that associate with myocarditis induction. Such studies have identified

proteins that comprise the viral inner shell (core) as determinants of myocarditis (7, 9,

10). Yet, amino acid changes in core proteins of myocarditic and non-myocarditic

strains that define differences in heart disease have remained unidentified. Interestingly,

many myocarditic strains are reassortants . Clone 8B, a highly myocarditic reovirus

strain, was generated via reassortment between the mildly myocarditic reovirus strain

type 1 Lang (T1L) and the non-myocarditic strain type 3 Dearing (T3D) (11). Clone 8B

contains 8 gene segments from T1L in combination with the S1 and M2 genes from T3D

(11). The reovirus S1 gene encodes viral attachment protein σ1 and non-structural

protein σ1s, while M2 encodes outer capsid protein μ1 (6). T1L/T3Dμ1σ3 contains 8

gene segments from T1L with the M2 and S4 genes from T3D (12). The S4 gene

encodes outer capsid protein σ3 that complexes with μ1 on the reovirus virion (6). The

KC19 strain, which contains 7 genes from T1L and the S1, S3 (encoding nonstructural

protein σNS), and M2 genes from T3D also has higher myocarditic potential than T1L

(7). The underlying commonality among clone 8B, T1L/T3Dμ1σ3, and KC19 is that they

contain the M2 gene derived from T3D. This observation raises the possibility that the

T3D-derived M2 gene impacts the myocarditic potential of reovirus strains. Yet, this idea

has not been formally tested.

Reovirus infection is initiated by binding of viral attachment protein σ1 to cell

surface receptors, followed by internalization of the particle by receptor-mediated

endocytosis (13, 14). In the endosome, heterohexameric complexes of μ1 and σ3 that

form the viral outer capsid are cleaved by the activity of endosome-resident cathepsin

proteases to facilitate virus entry (15-17). The σ3 protein is degraded, and μ1 is cleaved

to enable conformational changes that are required for endosomal membrane

penetration and the release of viral cores into the cytoplasm for transcription and

subsequent replication (6, 15-17). In addition to its role in entry, the μ1 protein also

contributes to the induction of apoptosis in reovirus-infected cells (6, 18, 19). T3 and T1

reoviruses differ in many aspects of virus replication and host response. Among them,

several phenotypes are associated with T3-derived M2. A single-gene reassortant

containing the M2 gene from T3D strain in an otherwise T1L genetic background

(T1L/T3DM2) has increased attachment to host cells compared to T1L (20). Enhanced

binding of T1L/T3DM2 correlated with greater infectivity and cell death induction in

cultured cells. These observations suggest that the reovirus M2 gene, and by extension

the µ1 protein, can influence reovirus tropism and disease *in vivo*. The presence of T3D

M2 in the virion also allows more efficient conformational changes that occur during cell

entry. Consequently, membrane penetration and core delivery also are more efficient for

viruses containing a T3D-derived M2 gene (21-23). The higher apoptotic potential of T3

strains in comparison to T1 strains also maps to the µ1-encoding M2 gene segment (18,

19). The precise function of µ1 in apoptosis induction is unclear since seemingly

contradictory data exist to support the role of both incoming µ1 from capsids and newly

synthesized µ1 produced during virus replication (18, 24-26). Nonetheless, mutations in

the ɸ domain of T3D µ1 diminish the apoptotic capacity of reovirus-induced apoptosis *in*

vivo, a phenotype associated with attenuated virulence (27).

Here, we assessed the function of M2 in reovirus pathogenesis. In contrast to

T1L, T1L/T3DM2 was lethal in neonatal mice. We found that T1L/T3DM2 replicated to

higher titers than T1L in all organs tested. Hearts from mice infected with T1L/T3DM2

displayed signs of overt myocarditis, including extensive lesions with marked

inflammatory infiltrates. In contrast, hearts from T1L-infected animals appeared normal

or had small lesions consistent with T1L being mildly myocarditic. Together, these

findings indicate that the T3D M2 gene is a determinant of reovirus-induced myocarditis.

RESULTS

T1L/T3DM2 is more lethal than T1L in neonatal mice. To determine whether

the T3D M2 allele modulates reovirus virulence, we assessed survival of neonatal

C57BL/6 mice infected orally with T1L or T1L/T3DM2 viruses (Fig 1A). Consistent with

previous findings (6), all mice inoculated with T1L survived. In contrast, 9 of 10 mice

(90%) succumbed to infection with T1L/T3DM2. Similar results were obtained following

intracranial inoculation (Fig 1B). Following T1L infection, 11 of 12 mice (91.7%)

survived, whereas 1 of 14 mice (7.1%) survived T1L/T3DM2 infection. These data

indicate that replacing the T1L M2 allele with the T3D M2 allele converts the non-lethal

T1L strain into a lethal infection in neonatal mice. Further because infection with

T1L/T3DM2 is lethal following both peroral and intracranial inoculation, our work

indicates that the effect of T3D M2 on reovirus virulence is independent of the

inoculation route.

T1L/T3DM2 replicates and disseminates more efficiently than T1L *in vivo*.

Next, we measured viral burdens following oral inoculation with T1L or T1L/T3DM2

viruses (Fig 2). At days 1 and 2 post-infection, T1L/T3DM2 produced higher viral titers

in the intestine than T1L. However, T1L and T1L/T3DM2 intestinal viral titers

equilibrated by day 4. In secondary organs (liver, spleen, heart, and brain), T1L/T3DM2

produced higher viral loads than T1L at most time points and achieved higher peak viral

titers than T1L in each organ tested. The largest difference in peak viral titers between

T1L/T3DM2 and T1L was observed in the heart, where T1L/T3DM2 viral loads were

approximately 50-fold higher than T1L at 8 days post-infection.

We also assessed viral loads following intracranial (IC) inoculation. Similar to oral

inoculation, T1 reoviruses also disseminate systemically following IC inoculation (28,

29). As observed following oral inoculations T1L/T3DM2 produced higher viral titers at

earlier time points and had higher peak viral loads compared to T1L in each organ when

virus was introduced IC (Fig 3). Note that only a single animal inoculated with

T1L/T3DM2 survived until day 8. Again, the largest difference in peak viral titers

between T1L/T3DM2 and T1L was found in the heart. T1L/T3DM2 viral loads were 10-

fold higher than T1L at 4 days post-infection. Together, these data indicate that the M2

gene influences reovirus replication and dissemination *in vivo*.

Systemic dissemination is required for T3D M2-mediated lethality. Both type

1 and type 3 reoviruses require junctional adhesion molecule-A (JAM-A) for systemic

dissemination (30). To determine whether systemic spread is required for the virulence

differences between T1L and T1L/T3DM2, we measured survival of neonatal JAM-A$^{-/-}$

mice infected with T1L or T1L/T3DM2 viruses (Fig 4). Following oral inoculation, none of

the JAM-A$^{-/-}$ mice succumbed to T1L and 10 of 11 (>90%) mice inoculated with

T1L/T3DM2 survived (Fig 4A). T1L/T3DM2 produced higher viral titers in the intestine

than T1L at days 1 and 2 post-infection, but T1L/T3DM2 and T1L viral loads were equal

by day 8 (Fig 4A). Intestinal viral loads in JAM-A$^{-/-}$ mice were comparable to those

observed in wild-type mice (Fig 2 vs Fig 4A). This observation is consistent with

published results demonstrating that JAM-A is dispensable for reovirus replication in the

intestine (30). A small number of JAM-A$^{-/-}$ mice had detectable virus in the heart and

brain but the majority of animals had no virus. Although most animals had measurable

viral titers of T1L and T1L/T3DM2 in the liver and spleen, viral progeny produced in

each organ in JAM-A$^{-/-}$ mice were substantially lower than in wild-type mice (Fig 2 vs Fig

4A).

Similar results were obtained following intracranial inoculation. All JAM-A$^{-/-}$ mice

infected with T1L or T1L/T3DM2 survived (Fig 4B). These data indicate that JAM-A

expression is required for T1L/T3DM2 lethality *in vivo*. T1L and T1L/T3DM2 replicated

comparably the brain (Fig 4B), which is consistent with JAM-A being dispensable for

reovirus replication in the central nervous system (30). In secondary organs, T1L and

T1L/T3DM2 viral titers were below the level of detection at 1 and 2 days post-infection,

but some virus was detected in the heart, liver, and spleen at days 4 and 8 (Fig 4B).

However, the viral titers produced in JAM-A$^{-/-}$ mice were substantially lower than the

viral loads measured in wild-type animals (Fig 3 vs Fig 4B). T1L and T1L/T3DM2 viral

titers in the intestine increased over time, and T1L/T3DM2 produced higher intestinal

viral loads than T1L at day 8 (Fig 4B). These data suggest that systemic dissemination

is required for the enhanced virulence of T1L/T3DM2 compared to T1L.

T1L/T3DM2 is more myocarditic than T1L. Regardless of the inoculation route,

T1L/T3DM2 was substantially more virulent than T1L in wild-type mice (Fig 1). During

necropsy, we observed that hearts from T1L infected mice were normal, with a few mice

having small cardiac lesions (Fig 5A and B). This result is consistent with published

reports that T1L is mildly myocarditic (7, 31, 32). In contrast, hearts from T1L/T3DM2-

infected mice were grossly abnormal. In some animals the entire heart appeared

opaque, which is suggestive of severe myocarditis. These data indicate that substitution

of reovirus T3D M2 allele enhances the capacity of T1L to cause cardiac pathology. In

JAM-A$^{-/-}$ mice, hearts from T1L or T1L/T3DM2-infected animals lacked any signs of

inflammation (data not shown), which is consistent with the residual viral titer detected

in the heart following oral and IC inoculation (Fig 4). This finding indicates that reovirus

dissemination to the heart is required for the development of cardiac lesions upon

reovirus infection.

We performed histological analysis of hearts from wild-type mice infected with

T1L or T1L/T3DM2 (Fig 6A). In contrast to mock-infected hearts, T1L-infected hearts

had small patches of cardiomyocyte necrosis admixed with pyknotic nuclear debris and

histiocytes. T1L lesions typically contained 2-3 cardiomyocytes exhibiting myolysis,

which is an indicator of destruction of cardiac muscle tissue. In addition, scant

lymphocyte and histiocyte infiltration were observed. These data are consistent with

previous observations of T1L inducing mild myocarditis in neonatal wild-type mice. On

the other hand, hearts from mice infected with T1L/T3DM2 displayed more frequent

cardiomyocyte damage and had increased lymphocytic and histiocytic infiltration

scattered in the myocardium as well as prominently in the epicardium compared to T1L-

infected hearts (Fig 6A). Hearts infected with T1L/T3DM2 showed markedly more viral

antigen compared to T1L (Fig 6B). This observation is consistent with the higher peak

titers produced by T1L/T3DM2 compared to T1L in the heart at day 8 post-infection (Fig

2 and Fig 3). Additionally, regions of viral antigen staining in T1L and T1L/T3DM2-

infected hearts overlapped with myocardial lesions (Fig 6B). This finding indicates that

reovirus replication in cardiomyocytes is associated with lesion formation (11). Lesions

from T1L/T3DM2-infected hearts contained cells that also stained positive for activated

caspase-3, which is a marker of apoptosis (Fig 6C). Together, these data indicate that

T1L- and T1L/T3DM2-induced cardiac lesions contain reovirus-infected cells, infiltrating

immune cells, and apoptotic and necrotic cells, with T1L/T3DM2 causing more

extensive lesion formation than T1L.

DISCUSSION

In this study, we found that the M2 gene is a determinant of reovirus myocarditis.

Reovirus strain T1L is non-lethal in neonatal mice although it induces mild heart

inflammation in a subset of animals. However, replacement of the T1L M2 allele with the

T3D M2 allele (T1L/T3DM2) was sufficient to confer a lethal phenotype that is

characterized by marked inflammation in approximately 90% of infected animals.

T1L/T3DM2 replicated to significantly higher titers at initial site of inoculation (Fig 2 and

Fig 3), disseminated to secondary organs with more rapid kinetics and produced higher

peak organ titers than T1L (Fig 2 and Fig 3). T1L/T3DM2 binds and infects L929 cells

more efficiently than T1L, leading to a higher percentage of cells becoming infected and

ultimately producing higher progeny yields than T1L (20). It is possible that T1L/T3DM2

initially infects more cells than T1L, leading to higher progeny yields at the primary site

of replication that subsequently disseminate systemically and infect target organs more

efficiently, including the heart. The greatest difference in peak viral titers between T1L

and T1L/T3DM2 was observed in the heart, where T1L/T3DM2 viral progeny were

approximately 50-fold and 10-fold higher than T1L following oral and intracranial

inoculation, respectively (Fig 2, Fig 3). Previous studies indicate that reovirus replication

levels in the heart do not correlate with myocarditis induction (10). It is possible that

delivery of T1L/T3DM2 virus to the heart more rapidly than T1L allows T1L/T3DM2 to

reach a threshold that contributes to the myocarditic capacity of T1L/T3DM2.

Apoptosis is a key pathogenic mechanism that contributes to reovirus

myocarditis (6, 33). Clone 8B induces caspase-3 activation within the myocardium of

infected hearts, enhancing myocardial injury (34). Moreover, myocarditis development is

dramatically reduced in caspase-3-deficient mice and mice treated with caspase

inhibitors. (34-36). Reoviruses display serotype-specific differences in apoptosis

induction, with T3 reoviruses inducing substantially more apoptosis than T1 strains (18).

One determinant of differences in the capacity of T1 and T3 reoviruses to induce

apoptosis is the M2 gene (6, 18, 27). We found that hearts from mice infected with

T1L/T3DM2 had more prominent activated caspase-3 staining than T1L-infected hearts

(Fig 6). Therefore, we hypothesize that the T3D M2 allele enhances apoptosis in

infected cells, which promotes myocardial injury and inflammation, the hallmark of

myocarditis.

The inflammatory response also may contribute to the enhanced myocarditic

capacity of the T1L/T3DM2 virus. The adaptive immune response is dispensable for

reovirus-induced myocarditis (6, 37). However, innate immune cells may contribute to

the heart inflammation and cell death observed following infection with the T1L/T3DM2

virus. Similar to what we observed for T1L/T3DM2, clone 8B induced significant

mononuclear cardiac infiltration (11). The myocarditic reovirus strain T1L/T3DM2S4,

which contains the M2 and S4 genes from T3D, elicits substantial pro-inflammatory

cytokine production in the heart that promotes cardiomyocyte damage (12).

T1L is mildly myocarditic, whereas T3D is non-myocarditic (6, 7, 31). However,

reassortant viruses generated from cells coinfected with T1L and T3D can cause severe

myocarditis (11, 38). Several highly myocarditic reassortants contain the T3D M2 gene.

Clone 8B contains the S1 and M2 alleles from T3D, while clone KC19 contains the T3D

S1, M2, and S3 alleles (7, 11). Clone EW60 is genetically similar to the T1L/T3DM2

used in this study as it is a single-gene reassortant containing the T3D M2 allele in a

T1L genetic background (31). Here, we used T1L/T3DM2 produced using reverse

genetics to clarify and characterize how M2 contributes to reovirus myocarditis.

However, the M2 gene is not the sole determinant of reovirus myocarditis. The M1

gene, which encodes inner core protein µ2 that functions as a co-factor for the reovirus

RNA-dependent RNA polymerase, also is a determinant of reovirus myocarditis (6, 8,

10, 32). M1 enhances reovirus myocarditis by potentiating replication in cardiac cells

and modulating the sensitivity to type-1 interferon (8, 39). It is likely that a combination

of mechanisms contribute to increased myocarditis caused by T1L/T3DM2. It is possible

that enhanced dissemination to and replication in the heart coupled with amplified

apoptotic capacity increase the level of cardiac damage induced by T1L/T3DM2. In turn,

the recruitment of immune cells to fight the infection may further exacerbate cardiac

damage by the production of pro-inflammatory cytokines and directly or indirectly killing

cells.

Other models of virus-induced myocarditis share similarities with reovirus (40-

43). Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) replication in the

myocardium is hypothesized to drive the pathological changes observed in the heart

tissue (44). Coxsackievirus B3 and murine cytomegalovirus (CMV) myocarditis require

viral replication and apoptosis induction (40-43). Studies of how reovirus causes

pathology in cardiac muscle cells can provide valuable information about how the heart

combats viral infection.

MATERIALS AND METHODS

Cells. Spinner-adapted murine L929 fibroblasts were maintained in Joklik's

modified minimum essential medium (JMEM, Sigma) supplemented with 5% heat-

inactivated fetal bovine serum (FBS, Invitrogen), 1% 200 mM L-glutamine (Invitrogen),

1% 10,000 U/ml penicillin/ 10,000 µg/ml streptomycin (Invitrogen), and 0.1% 250 µg/ml

amphotericin B (Sigma).

Viruses. Recombinant reoviruses (T1L and T1L/T3DM2) were generated using

plasmid-based reverse genetics (45, 46). Reovirus virions were purified from second- or

third-passage L929 cell lysates infected with twice-plaque-purified reovirus (47).

Reovirus particles were Vertrel (TMC industries) extracted from infected cell lysates,

layered onto 1.2- to 1.4-g/cm^3 CsCl gradients, and centrifuged at 25,000 rpm for 18h or

29,000 rpm for 5h. Virions were collected and dialyzed against dialysis buffer (150 mM

NaCl, 15 mM MgCl, and 10 mM Tris-HCl [pH 7.4]). Viral titers were determined by

plaque assay using L929 fibroblasts (48). To confirm the viral gene segments, viral

dsRNA was purified from virions and separated by SDS-PAGE. Gene segments were

visualized by ethidium bromide staining (45).

Mouse experiments. C57BL/6 mice were obtained from Jackson Laboratory.

C57BL/6-JAM-A$^{-/-}$ mice were obtained from the Dermody Laboratory at the University of

Pittsburgh. Animal husbandry and housing were performed following the guidelines of

the Division of Laboratory Animal Medicine (DLAM) at the University of Arkansas for

Medical Sciences (UAMS). Neonatal mice (3-5 days old) were infected orally with 10^4

PFU or intracranially with 100 PFU of rsT1L or rsT1L/T3DM2. For survival experiments,

infected mice were monitored for 20 or 25 days and sacrificed if moribund. Brain, heart,

spleen, liver, and intestine organs were resected at the times indicated in the figure

legends, homogenized, and viral titers were determined by plaque assay on L929 cells.

Resected hearts were imaged at the times indicated in the figure legends using a hand-

held camera.

Histology and immunohistochemistry. Neonatal C57BL/6 mice (3-5 days old)

were mock-infected (PBS) or infected orally with 10^4 PFU of rsT1L or rsT1L/T3DM2. At

8 days post-infection, mice were euthanized and the hearts were resected. Vertical

longitudinal cuts of the infected hearts were prepared and fixed in methanol Carnoy's for

24h followed by 70% ethanol until embedding. The fixed hearts were submitted to the

UAMS Experimental Pathology Core Laboratory for paraffin embedding and sectioning

(4 μm thickness). The cardiac sections were stained with hematoxylin and eosin (H&E)

and a pathologist blinded to the identity of the inoculum examined them under a light

microscope. Consecutive sections were processed for immunohistochemical staining

with reovirus-specific rabbit polyclonal antiserum (1:1000) or activated caspase-3

(Abcam, 1:100). Images were taken on an EVOS FL Auto 2 (Invitrogen) using the 10x

and 20x objectives (133x and 266x total magnification, respectively).

Statistics. Statistical analysis was performed using Prism software (GraphPad

Software Inc.). Differences in survival were determined by Log-rank test. Differences in

mean viral titers were determined using Mann-Whitney non-parametric *t*-test. *P* values

less than 0.05 were considered statistically significant.

ACKNOWLEDGMENTS

We thank the Dermody Lab (University of Pittsburgh) for providing the

JAM-A$^{-/-}$ mice. We are grateful to Jennifer James at the UAMS Experimental

Pathology Core (UAMS) for assistance with histological sample preparations.

This research was supported by Public Health Award R01 AI118801 (K.W.B.)

 FIGURE LEGENDS

Figure 1. T1L/T3DM2 is more lethal than T1L in neonatal mice. (A) Neonatal

C57BL/6 mice were inoculated orally with 10^4 PFU of T1L or T1L/T3DM2 viruses. Mice

were monitored for 25 days. (B) Neonatal C57BL/6 mice were inoculated intracranially

with 100 PFU of T1L or T1L/T3DM2 viruses. Mice were monitored for 20 days. ****, $P <$

0.0001 as determined by log-rank test in comparison to T1L.

Figure 2. T1L/T3DM2 disseminates more efficiently than T1L following oral

inoculation. Neonatal C57BL/6 mice were inoculated orally with 10^4 PFU of T1L or

T1L/T3DM2 viruses. On days 1, 2, 4, and 8, mice were euthanized and the indicated

organs were resected. Viral titers in each organ were determined by plaque assay. Error

bars represent SD. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ as determined by Mann-

Whitney test.

Figure 3. T1L/T3DM2 disseminates more efficiently than T1L following intracranial

inoculation. Neonatal C57BL/6 mice were inoculated intracranially with 100 PFU of

T1L or T1L/T3DM2 viruses. On days 1, 2, 4, and 8, mice were euthanized and the

indicated organs were resected. Viral titers in each organ were determined by plaque

assay. Error bars represent SD. *, $P < 0.05$; ***, $P < 0.001$ as determined by Mann-

Whitney test.

Figure 4. Systemic dissemination is required for T1L/T3DM2 lethality. Neonatal

JAM-A$^{-/-}$ mice were inoculated (A) orally with 10^4 PFU or (B) intracranially with 100 PFU

of T1L or T1L/T3DM2 viruses. For survival experiments, mice were monitored for 25

days. For titers experiments, on days 1, 2, 4, and 8, mice were euthanized, and the

indicated organs were resected. Viral titers in each organ were determined by plaque

assay. Error bars represent SD. *, $P < 0.05$; **, $P < 0.01$ as determined by Mann-

Whitney test.

Figure 5. T1L/T3DM2 causes more cardiac pathology than T1L. Neonatal C57BL/6

mice were inoculated (A) orally with 10^4 PFU or (B) intracranially with 100 PFU of T1L or

T1L/T3DM2. At 8 days, mice were euthanized and hearts were resected and imaged.

Figure 6. T1L/T3DM2 is more myocarditic than T1L. Neonatal C57BL/6 mice were

mock-infected with PBS or inoculated orally with 10^4 PFU of T1L or T1L/T3DM2 viruses.

At 8 days, the mice were euthanized, the hearts were resected, paraffin embedded,

sectioned, and stained with (A) hematoxylin and eosin (H&E), (B) polyclonal reovirus

antiserum (Reo), or (C) cleaved-caspase-3 antibodies. Shown are sections imaged at

10x magnification (top panels) or 20x (bottom panels).

REFERENCES

1. Fung G, Luo H, Qiu Y, Yang D, McManus B. 2016. Myocarditis. Circ Res 118:496-514. https://doi.org/10.1161/CIRCRESAHA.115.306573.
2. Opavsky MA, Penninger J, Aitken K, Wen WH, Dawood F, Mak T, Liu P. 1999. Susceptibility to myocarditis is dependent on the response of alphabeta T lymphocytes to coxsackieviral infection. Circ Res 85:551-8. https://doi.org/10.1161/01.res.85.6.551.
3. Bock CT, Klingel K, Kandolf R. 2010. Human parvovirus B19-associated myocarditis. N Engl J Med 362:1248-9. https://doi.org/10.1056/NEJMc0911362.
4. Duechting A, Tschope C, Kaiser H, Lamkemeyer T, Tanaka N, Aberle S, Lang F, Torresi J, Kandolf R, Bock CT. 2008. Human parvovirus B19 NS1 protein modulates inflammatory signaling by activation of STAT3/PIAS3 in human endothelial cells. J Virol 82:7942-52. https://doi.org/10.1128/JVI.00891-08.
5. Perez-Bermejo JA, Kang S, Rockwood SJ, Simoneau CR, Joy DA, Silva AC, Ramadoss GN, Flanigan WR, Fozouni P, Li H, Chen PY, Nakamura K, Whitman JD, Hanson PJ, McManus BM, Ott M, Conklin BR, McDevitt TC. 2021. SARS-CoV-2 infection of human iPSC-derived cardiac cells reflects cytopathic features in hearts of patients with COVID-19. Sci Transl Med 13. https://doi.org/10.1126/scitranslmed.abf7872.
6. Dermody TS, Parker JSL, Sherry B. 2013. Orthoreovirus, p 1304-1346. *In* Knipe DM, Howley PM (ed), Fields Virology, 6th ed, vol 2. Lippincott, Williams, & Wilkins, Philadelphia, PA.
7. Sherry B, Blum MA. 1994. Multiple viral core proteins are determinants of reovirus-induced acute myocarditis. J Virol 68:8461-5.
8. Sherry B, Torres J, Blum MA. 1998. Reovirus induction of and sensitivity to beta interferon in cardiac myocyte cultures correlate with induction of myocarditis and are determined by viral core proteins. J Virol 72:1314-23.
9. Matoba Y, Sherry B, Fields BN, Smith TW. 1991. Identification of the viral genes responsible for growth of strains of reovirus in cultured mouse heart cells. J Clin Investig 87:1628-33. https://doi.org/10.1172/JCI115177.
10. Sherry B, Fields BN. 1989. The reovirus M1 gene, encoding a viral core protein, is associated with the myocarditic phenotype of a reovirus variant. J Virol 63:4850-6.
11. Sherry B, Schoen FJ, Wenske E, Fields BN. 1989. Derivation and characterization of an efficiently myocarditic reovirus variant. J Virol 63:4840-9.
12. Doyle JD, Stencel-Baerenwald JE, Copeland CA, Rhoads JP, Brown JJ, Boyd KL, Atkinson JB, Dermody TS. 2015. Diminished reovirus capsid stability alters disease pathogenesis and littermate transmission. PLoS Pathog 11:e1004693. https://doi.org/10.1371/journal.ppat.1004693.
13. Barton ES, Chappell JD, Connolly JL, Forrest JC, Dermody TS. 2001. Reovirus receptors and apoptosis. Virology 290:173-80. https://doi.org/10.1006/viro.2001.1160.
14. Campbell JA, Schelling P, Wetzel JD, Johnson EM, Forrest JC, Wilson GA, Aurrand-Lions M, Imhof BA, Stehle T, Dermody TS. 2005. Junctional adhesion molecule a serves as a receptor for prototype and field-isolate strains of

420 mammalian reovirus. J Virol 79:7967-78. https://doi.org/10.1128/JVI.79.13.7967-
421 7978.2005.
15. Liemann S, Chandran K, Baker TS, Nibert ML, Harrison SC. 2002. Structure of
     the reovirus membrane-penetration protein, Mu1, in a complex with is protector
     protein, Sigma3. Cell 108:283-95.
16. Snyder AJ, Danthi P. 2018. Cleavage of the C-Terminal Fragment of Reovirus
     mu1 Is Required for Optimal Infectivity. J Virol 92.
     https://doi.org/10.1128/JVI.01848-17.
17. Snyder AJ, Danthi P. 2018. Infectious Subviral Particle to Membrane Penetration
     Active Particle (ISVP-to-ISVP*) Conversion Assay for Mammalian Orthoreovirus.
     Bio Protoc 8. https://doi.org/10.21769/BioProtoc.2700.
18. Tyler KL, Squier MK, Brown AL, Pike B, Willis D, Oberhaus SM, Dermody TS,
     Cohen JJ. 1996. Linkage between reovirus-induced apoptosis and inhibition of
     cellular DNA synthesis: role of the S1 and M2 genes. J Virol 70:7984-91.
19. Oberhaus SM, Dermody TS, Tyler KL. 1998. Apoptosis and the cytopathic effects
     of reovirus. Curr Top Microbiol Immunol 233:23-49.
20. Thete D, Snyder AJ, Mainou BA, Danthi P. 2016. Reovirus mu1 Protein Affects
     Infectivity by Altering Virus-Receptor Interactions. J Virol 90:10951-10962.
     https://doi.org/10.1128/JVI.01843-16.
21. Lucia-Jandris P, Hooper JW, Fields BN. 1993. Reovirus M2 gene is associated
     with chromium release from mouse L cells. J Virol 67:5339-45.
22. Chandran K, Walker SB, Chen Y, Contreras CM, Schiff LA, Baker TS, Nibert ML.
     1999. In vitro recoating of reovirus cores with baculovirus-expressed outer-capsid
     proteins mu1 and sigma3. J Virol 73:3941-50.
23. Sarkar P, Danthi P. 2010. Determinants of strain-specific differences in efficiency
     of reovirus entry. J Virol 84:12723-32. https://doi.org/10.1128/JVI.01385-10.
24. Danthi P, Coffey CM, Parker JS, Abel TW, Dermody TS. 2008. Independent
     regulation of reovirus membrane penetration and apoptosis by the mu1 phi
     domain. PLoS Path 4:e1000248. https://doi.org/10.1371/journal.ppat.1000248.
25. Danthi P, Hansberger MW, Campbell JA, Forrest JC, Dermody TS. 2006. JAM-A-
     independent, antibody-mediated uptake of reovirus into cells leads to apoptosis.
     J Virol 80:1261-70. https://doi.org/10.1128/JVI.80.3.1261-1270.2006.
26. Roebke KE, Danthi P. 2019. Cell Entry-Independent Role for the Reovirus mu1
     Protein in Regulating Necroptosis and the Accumulation of Viral Gene Products.
     J Virol 93. https://doi.org/10.1128/JVI.00199-19.
27. Oberhaus SM, Smith RL, Clayton GH, Dermody TS, Tyler KL. 1997. Reovirus
     infection and tissue injury in the mouse central nervous system are associated
     with apoptosis. J Virol 71:2100-6.
28. Boehme KW, Guglielmi KM, Dermody TS. 2009. Reovirus nonstructural protein
     sigma1s is required for establishment of viremia and systemic dissemination.
     Proc Natl Acad Sci U S A 106:19986-91.
     https://doi.org/10.1073/pnas.0907412106.
29. Phillips MB, Dina Zita M, Howells MA, Weinkopff T, Boehme KW. 2021.
     Lymphatic Type 1 Interferon Responses Are Critical for Control of Systemic
     Reovirus Dissemination. J Virol 95. https://doi.org/10.1128/JVI.02167-20.

30. Antar AA, Konopka JL, Campbell JA, Henry RA, Perdigoto AL, Carter BD, Pozzi A, Abel TW, Dermody TS. 2009. Junctional adhesion molecule-A is required for hematogenous dissemination of reovirus. Cell Host Microbe 5:59-71. https://doi.org/10.1016/j.chom.2008.12.001.

31. Sherry B, Baty CJ, Blum MA. 1996. Reovirus-induced acute myocarditis in mice correlates with viral RNA synthesis rather than generation of infectious virus in cardiac myocytes. J Virol 70:6709-15.

32. Irvin SC, Zurney J, Ooms LS, Chappell JD, Dermody TS, Sherry B. 2012. A single-amino-acid polymorphism in reovirus protein mu2 determines repression of interferon signaling and modulates myocarditis. J Virol 86:2302-11. https://doi.org/10.1128/JVI.06236-11.

33. DeBiasi RL, Robinson BA, Leser JS, Brown RD, Long CS, Clarke P. 2010. Critical role for death-receptor mediated apoptotic signaling in viral myocarditis. J Card Fail 16:901-10. https://doi.org/10.1016/j.cardfail.2010.05.030.

34. DeBiasi RL, Robinson BA, Sherry B, Bouchard R, Brown RD, Rizeq M, Long C, Tyler KL. 2004. Caspase inhibition protects against reovirus-induced myocardial injury in vitro and in vivo. J Virol 78:11040-50. https://doi.org/10.1128/JVI.78.20.11040-11050.2004.

35. DeBiasi RL, Edelstein CL, Sherry B, Tyler KL. 2001. Calpain inhibition protects against virus-induced apoptotic myocardial injury. J Virol 75:351-61. https://doi.org/10.1128/JVI.75.1.351-361.2001.

36. Kim M, Hansen KK, Davis L, van Marle G, Gill MJ, Fox JD, Hollenberg MD, Rancourt DE, Lee PW, Yun CO, Johnston RN. 2010. Z-FA-FMK as a novel potent inhibitor of reovirus pathogenesis and oncolysis in vivo. Antivir Ther 15:897-905. https://doi.org/10.3851/IMP1646.

37. Sherry B, Li XY, Tyler KL, Cullen JM, Virgin HWt. 1993. Lymphocytes protect against and are not required for reovirus-induced myocarditis. J Virol 67:6119-24.

38. Wenske EA, Chanock SJ, Krata L, Fields BN. 1985. Genetic reassortment of mammalian reoviruses in mice. J Virol 56:613-6.

39. Stewart MJ, Smoak K, Blum MA, Sherry B. 2005. Basal and reovirus-induced beta interferon (IFN-beta) and IFN-beta-stimulated gene expression are cell type specific in the cardiac protective response. J Virol 79:2979-87. https://doi.org/10.1128/JVI.79.5.2979-2987.2005.

40. Lenzo JC, Fairweather D, Cull V, Shellam GR, James Lawson CM. 2002. Characterisation of murine cytomegalovirus myocarditis: cellular infiltration of the heart and virus persistence. J Mol Cell Cardiol 34:629-40. https://doi.org/10.1006/jmcc.2002.2003.

41. Henke A, Launhardt H, Klement K, Stelzner A, Zell R, Munder T. 2000. Apoptosis in coxsackievirus B3-caused diseases: interaction between the capsid protein VP2 and the proapoptotic protein siva. J Virol 74:4284-90. https://doi.org/10.1128/jvi.74.9.4284-4290.2000.

42. Olivetti G, Abbi R, Quaini F, Kajstura J, Cheng W, Nitahara JA, Quaini E, Di Loreto C, Beltrami CA, Krajewski S, Reed JC, Anversa P. 1997. Apoptosis in the failing human heart. N Engl J Med 336:1131-41. https://doi.org/10.1056/NEJM199704173361603.

43.   Colston JT, Chandrasekar B, Freeman GL. 1998. Expression of apoptosis-
related proteins in experimental coxsackievirus myocarditis. Cardiovasc Res
38:158-68. https://doi.org/10.1016/s0008-6363(97)00323-4.
44.   Zeng JH, Liu YX, Yuan J, Wang FX, Wu WB, Li JX, Wang LF, Gao H, Wang Y,
Dong CF, Li YJ, Xie XJ, Feng C, Liu L. 2020. First case of COVID-19
complicated with fulminant myocarditis: a case report and insights. Infection
48:773-777. https://doi.org/10.1007/s15010-020-01424-5.
45.   Kobayashi T, Antar AA, Boehme KW, Danthi P, Eby EA, Guglielmi KM, Holm
GH, Johnson EM, Maginnis MS, Naik S, Skelton WB, Wetzel JD, Wilson GJ,
Chappell JD, Dermody TS. 2007. A plasmid-based reverse genetics system for
animal double-stranded RNA viruses. Cell Host Microbe 1:147-57.
https://doi.org/10.1016/j.chom.2007.03.003.
46.   Kobayashi T, Ooms LS, Ikizler M, Chappell JD, Dermody TS. 2010. An improved
reverse genetics system for mammalian orthoreoviruses. Virology 398:194-200.
https://doi.org/10.1016/j.virol.2009.11.037.
47.   Furlong DB, Nibert ML, Fields BN. 1988. Sigma 1 protein of mammalian
reoviruses extends from the surfaces of viral particles. J Virol 62:246-56.
https://doi.org/10.1128/JVI.62.1.246-256.1988.
48.   Tyler KL, Bronson RT, Byers KB, Fields B. 1985. Molecular basis of viral
neurotropism: experimental reovirus infection. Neurology 35:88-92.
https://doi.org/10.1212/wnl.35.1.88.

Figure 1. T1L/T3DM2 is more lethal than T1L in neonatal mice. (A) Neonatal

C57BL/6 mice were inoculated orally with 10^4 PFU of T1L or T1L/T3DM2 viruses. Mice

were monitored for 25 days. (B) Neonatal C57BL/6 mice were inoculated intracranially

with 100 PFU of T1L or T1L/T3DM2 viruses. Mice were monitored for 20 days. ****, *P* <

0.0001 as determined by log-rank test in comparison to T1L.

Figure 2. T1L/T3DM2 disseminates more efficiently than T1L following oral inoculation. Neonatal C57BL/6 mice were inoculated orally with 10^4 PFU of T1L or T1L/T3DM2 viruses. On days 1, 2, 4, and 8, mice were euthanized and the indicated organs were resected. Viral titers in each organ were determined by plaque assay. Error bars represent SD. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ as determined by Mann-Whitney test.

Figure 3. T1L/T3DM2 disseminates more efficiently than T1L following intracranial inoculation. Neonatal C57BL/6 mice were inoculated intracranially with 100 PFU of T1L or T1L/T3DM2 viruses. On days 1, 2, 4, and 8, mice were euthanized and the indicated organs were resected. Viral titers in each organ were determined by plaque assay. Error bars represent SD. *, $P < 0.05$; ***, $P < 0.001$ as determined by Mann-Whitney test.

Figure 4. Systemic dissemination is required for T1L/T3DM2 lethality. Neonatal JAM-A$^{-/-}$ mice were inoculated (A) orally with 10^4 PFU or (B) intracranially with 100 PFU of T1L or T1L/T3DM2 viruses. For survival experiments, mice were monitored for 25 days. For titers experiments, on days 1, 2, 4, and 8, mice were euthanized, and the indicated organs were resected. Viral titers in each organ were determined by plaque assay. Error bars represent SD. *, $P < 0.05$; **, $P < 0.01$ as determined by Mann-Whitney test.

Figure 5. T1L/T3DM2 causes more cardiac pathology than T1L. Neonatal C57BL/6

mice were inoculated (A) orally with 10^4 PFU or (B) intracranially with 100 PFU of T1L or

T1L/T3DM2. At 8 days, mice were euthanized and hearts were resected and imaged.

Figure 6. T1L/T3DM2 is more myocarditic than T1L. Neonatal C57BL/6 mice were mock-infected with PBS or inoculated orally with 10^4 PFU of T1L or T1L/T3DM2 viruses. At 8 days, the mice were euthanized, the hearts were resected, paraffin embedded, sectioned, and stained with (A) hematoxylin and eosin (H&E), (B) polyclonal reovirus antiserum (Reo), or (C) cleaved-caspase-3 antibodies. Shown are sections imaged at 10x magnification (top panels) or 20x (bottom panels).

A

T1L

T1L/T3DM2

B

T1L

T1L/T3DM2

	Mock	T1L	T1L/T3DM2

A