`

A vast evolutionarily transient translatome contributes to phenotype and fitness

Aaron Wacholder[12], Saurin Bipin Parikh[123], Nelson Castilho Coelho[12], Omer Acar[124], Carly
Houghton[124], Lin Chou[123], and Anne-Ruxandra Carvunis[125]*

1. Department of Computational and Systems Biology, School of Medicine, University of
Pittsburgh, Pittsburgh, PA, 15213, United States
2. Pittsburgh Center for Evolutionary Biology and Medicine, School of Medicine, University of
Pittsburgh, Pittsburgh, PA, 15213, United States
3. Integrative Systems Biology Program, School of Medicine, University of Pittsburgh,
Pittsburgh, PA, 15213, United States
4. Joint CMU-Pitt Ph.D. Program in Computational Biology, University of Pittsburgh, Pittsburgh,
PA, 15213, United States
5. Lead contact

*Correspondence: **anc201@pitt.edu**

`

15 **Summary**

Translation is the process by which ribosomes synthesize proteins. Ribosome profiling recently revealed

that many short sequences previously thought to be noncoding are pervasively translated. To identify

protein-coding genes in this noncanonical translatome, we combine an integrative framework for

extremely sensitive ribosome profiling analysis, iRibo, with high-powered selection inferences tailored

for short sequences. We construct a reference translatome for *Saccharomyces cerevisiae* comprising

5,400 canonical and almost 19,000 noncanonical translated elements. Only 14 noncanonical elements

were evolving under detectable purifying selection. Surprisingly, a representative subset of translated

elements lacking signatures of selection demonstrated involvement in processes including DNA repair,

stress response and post-transcriptional regulation. Our results suggest that most translated elements

are not conserved protein-coding genes and contribute to genotype-phenotype relationships through

fast-evolving molecular mechanisms.

Keywords:

Noncanonical translation, ribosome profiling, de novo gene birth, protein evolution, evolutionary

genomics, microproteins, smORFs, genome annotation

`

Introduction

The central role played by protein-coding genes in biological processes has made their identification and characterization an essential project for understanding organismal biology. Over the past decade, the scope of this project has expanded as ribosome profiling (ribo-seq) studies have revealed pervasive translation of eukaryotic genomes.[1–4] These experiments demonstrate that genomes encode not only the "canonical translatome", consisting of the open reading frames (ORFs) identified as protein-coding genes in genome databases like RefSeq[5], but also a large "noncanonical translatome" consisting of ORFs that are not annotated as genes. Despite lack of annotation, large-scale studies find that many noncanonical ORFs are translated to express stable proteins and show evidence of association with cellular phenotypes.[6–10] Additionally, a handful of previously unannotated coding sequences, identified by RNA-seq or ribo-seq experiments, have now been characterized in depth, revealing that they play key roles in biological pathways and are important to organism fitness.[11–15] Yet, these well-studied examples represent only a small fraction of the noncanonical translatome. Most noncanonical translation could simply be biologically insignificant "translational noise" resulting from the imperfect specificity of translation processes.[16–19] Alternatively, thousands of missing protein-coding genes that contribute to phenotype and fitness could be hidden in the noncanonical translatome.

A common and powerful approach to identifying biologically significant genomic sequences is to look for evidence of selection.[20–22] Many canonical genes were annotated on the basis of such evidence[23,24], and this approach has also been applied to noncanonical ORFs detected by ribo-seq.[25–28] However, in the case of noncanonical translation, evolutionary analysis is often limited by a lack of sufficient statistical power to confidently detect selection. Most noncanonical ORFs are much shorter than canonical genes[7,12,29], thus having fewer genetic variants that can be analyzed for evolutionary inference. As a result, short coding sequences are sometimes missed by genome-wide evolutionary analyses despite long-term evolutionary conservation.[13,30] It is especially challenging to detect selection among noncanonical ORFs that are evolutionarily novel, as a short evolutionary history also provides less time for enough genetic variants to accumulate the signatures that allow for statistically distinguishing selective from neutral evolution.[31] Several young genes of recent *de novo* origin (i.e., coding genes that evolved from previously nongenic sequences) have been discovered from within the noncanonical translatome.[3,32,33]

In addition to the challenges short ORF length poses for detection of selection, it also poses challenges for unequivocal detection of translation in the first place. Microproteins are often missed by most

proteomics techniques, though specialized methods have had some success.[9,10,34–36] In ribo-seq data, the

most robust evidence of translation comes from a pattern of triplet periodicity in reads corresponding to

the progression of the ribosome across codons.[6,37,38] Ribo-seq analysis methods are less capable of

detecting translation of short ORFs, as they contain fewer positions to use to establish periodicity.[39] The

low expression levels of some noncanonical ORFs further increases the difficulties in identification.[3,27]

Perhaps as a result of these limitations, less than half of the noncanonical ORFs detected as translated in

humans are reproducible across studies.[31]

Here, we designed an approach to increase sensitivity in detection of both translation and selection

among noncanonical ORFs. We address the challenges in detecting translation through the development

of a ribo-seq analysis framework (iRibo) that identifies signatures of translation with high sensitivity and

high specificity by integrating data across hundreds of experiments from many published studies. This

facilitates detection of sequences that are short or poorly expressed. We address the challenges in

detecting selection through a comparative genomics framework that analyzes translated sequences

collectively across evolutionary scales within- and between-species.

We applied our approach to define a "reference translatome" for the model organism *Saccharomyces*

cerevisiae and to characterize the biological significance of noncanonical ORFs. Using iRibo, we identified

~19,000 noncanonical ORFs translated at high confidence and established the dependence of

noncanonical translation on both genomic context and environmental condition. Using genomic data

both within strains of *S. cerevisiae* and across budding yeast species[40,41], we identified a handful of

undiscovered conserved genes within the yeast noncanonical translatome. However, we find that most

of the yeast noncanonical translatome is evolutionarily young and of *de novo* origin, having emerged

recently from noncoding sequence. These young ORFs differ greatly from conserved genes in their

length, amino acid composition, and expression level, and show no signs of purifying selection.

Nevertheless, we report experimental evidence based on fluorescent protein tagging and conditional

loss-of-function fitness measurements showing that translation of evolutionarily young noncanonical

ORFs can generate stable protein products and affect cellular phenotypes. We thus propose that much

of the noncanonical translatome is composed of neither translational noise nor conserved genes, but

rather of a distinct class of evolutionarily short-lived coding sequences with important biological

implications. This "transient translatome" is larger than, and categorically distinct from, the conserved

translatome made mostly of canonical protein-coding genes that have been studied for decades.

Results

`

An integrative approach to defining the translatome

We designed iRibo to detect translation events with high sensitivity and high specificity. High sensitivity

is achieved through integration of ribo-seq data across hundreds of diverse experiments, which provides

sufficient read depth for detection of translated ORFs that are short or weakly expressed. High

specificity is achieved through the use of three nucleotide periodicity as the sole basis for translation

inference. Three nucleotide periodicity corresponds to the progression of the ribosome codon by codon

across a transcript, a dynamic unique to translation. Three nucleotide periodicity is therefore robust

against false inference of translation from other sources of ribo-seq reads.[37,38,42] High specificity is

further achieved by controlling confidence levels using an empirical false discovery rate approach that

relies on minimal modeling assumptions. iRibo consists of four components (**Figure 1A**). First, a set of

"candidate" ORFs that could potentially be translated are identified in the genome. Second, reads from

multiple ribo-seq experiments are pooled and mapped to these ORFs. Third, the translation status of

each candidate ORF is assessed based on whether the reads mapping to the ORF exhibit a pattern of

triplet nucleotide periodicity according to a binomial test. Finally, a list of translated ORFs is constructed

with a specified false discovery rate, derived from applying the same translation calling method on a

negative control set constructed to exhibit no genuine signatures of translation.

`

Figure 1: The iRibo framework enables detection of thousands of noncanonical translated sequences.
A) The iRibo framework. 1) Candidate ORFs, both canonical (cORFs; red) and noncanonical (nORFs; blue),
are identified in the genome. 2) Reads aggregated from published datasets are then mapped to these

`

ORFs. 3) Translation is inferred from triplet periodicity of reads. 4) The false discovery rate is estimated
by scrambling the ribo-seq reads of each ORF and then assessing periodicity in this scrambled set. **B)**
iRibo identifies translated ORFs that are undetectable in any single experiment. Mapped ribo-seq reads
(y-axis) across an example nORF located on chromosome II, 604674-604748 (x-axis). The top five graphs
correspond to five individual experiments with reads mapping to the ORF while the bottom graph
includes all reads integrated across all experiments. Reads are colored according to their position on the
codon. **C)** iRibo identifies 18,953 translated nORFs at 5% false discovery rate. The number of nORFs
found to be translated using iRibo (y-axis) at a range of p-value thresholds (x-axis) is shown as a solid
blue line. Translation calls for a negative control set, constructed by scrambling the actual ribo-seq reads
for each nORF, is also plotted (dashed blue line). The dashed vertical line indicates false discovery rate of
5% among nORFs. **D)** iRibo identifies 5,364 cORFs**.** The number of cORFs found to be translated using
iRibo at a range of p-value thresholds, contrasted with negative controls constructed by scrambling the
ribo-seq reads of each cORF.

iRibo can be applied to a set of ribo-seq experiments conducted under a single environmental condition

to identify ORFs that are translated under that condition. Alternatively, iRibo can be deployed on a

broader set of ribo-seq experiments conducted in many different contexts to construct a "reference

translatome" consisting of all elements within a genome with sufficient evidence of translation.

We used iRibo to identify translated ORFs across the *S. cerevisiae* genome (**Supplementary Figure 1**)*.*

First, we constructed the set of candidate ORFs by collecting all genomic sequences at least three

codons in length that start with ATG and end with a stop codon in the same frame. For ORFs overlapping

in the same frame, only the longest ORF was kept. Each candidate ORF was classified either as canonical

(cORF), if it was annotated as "verified," "uncharacterized," or "transposable element" in the

Saccharomyces Genome Database (SGD)[43] or as noncanonical (nORF), if it was annotated as "dubious,"

"pseudogene," or was unannotated. We excluded nORFs that overlap cORFs on the same strand. This

process generated a list of 179,441 candidate ORFs: 173,868 nORFs and 5,573 cORFs. We assessed

translation status for candidate ORFs using data from 412 ribo-seq experiments across 42 studies

(**Supplementary Table 1, Supplementary Table 2**).

As expected, integrating data from many experiments allowed for identification of translated ORFs that

would otherwise have too few reads in any individual experiment (**Figure 1B**). Setting a confidence

threshold to ensure a 5% false discovery rate (FDR) among nORFs, we identified 18,953 nORFs (**Figure**

1C) as translated along with 5,364 cORFs (**Figure 1D**), for a total of 24,317 ORFs making up the yeast

reference translatome. This corresponds to an identification rate of 99% for "verified" cORFs, 77% for

"uncharacterized" cORFs, 37% for "dubious" nORFs and only 11% for unannotated nORFs (**Figure 2A**).

Despite the low rate of identified translation, unannotated nORFs make up a large majority of translated

`

148   sequences (**Figure 2B**). In general, translated cORFs are much longer (**Figure 2C)** and translated at much

149   higher rates (**Figure 2D**) than translated nORFs.

`

Figure 2: The noncanonical yeast translatome is larger than the canonical. A) A majority of cORFs, and
a minority of nORFs, are translated. The percent of ORFs (y-axis) in each Saccharomyces Genome

Database annotation class that are detected as translated by iRibo, with canonical classes indicated in
red and noncanonical in blue. **B)** Unannotated nORFs make up a large majority of translated sequences.
The number of ORFs (y-axis) of each annotation class (x-axis) that are detected using iRibo. **C)** nORFs are
shorter than cORFs. ORF length distributions for translated cORFs and nORFs. **D)** nORFs are translated at
lower rates than cORFs. Distribution of translation rate (in-frame ribo-seq reads per base) for translated
cORFs and nORFs. **E)** Translation calls are highly reproducible. For six large studies (y-axis), the
proportion of nORFs identified using reads from that study that are replicated using reads from the
largest study, Gerashchenko et al. 2014[44] (x-axis). Random expectation is the proportion that would be
expected to replicate by chance. **F)** CHX facilitates detection of translated nORFs. Ratio of total ribo-seq
read counts mapping to cORFs or nORFs in experiments with vs. without CHX treatment. Note that the
same number of total reads are sampled from each condition. **G)** nORFs identified as translated only
with CHX nevertheless show preference for the first codon position in its absence. Among nORFs
identified as translated by iRibo only in the CHX condition, all codons among these nORFs are classed
based on which of the three positions in the codon have the most reads from experiments without CHX.

To assess replicability in translation calls for nORFs, we applied iRibo separately to each of the largest

individual studies by read count. We then counted, among the nORFs that could be inferred to be

translated using only the reads in each study, how many were also found in the largest study,

Gerashchenko and Gladyshev, 2014.[44] For all studies, at least 75% of detected ORFs were also detected

in the largest study (**Figure 2E**). In general, translation rates among ORFs were highly correlated among

independent studies (**Supplementary Figure 2**). These observations demonstrate that noncanonical

translation patterns are highly reproducible, suggesting that they are driven by regulated biological

processes rather than technical artifacts or stochastic ribosome errors.

A large fraction of ribo-seq experiments use the translation elongation inhibitor cycloheximide (CHX).

This drug is known to influence ribo-seq results in several ways.[44–46] We therefore wished to specifically

examine whether the size of the noncanonical translatome we identified could have been artificially

inflated by CHX usage. To this aim, we compared translation signatures from experiments with (N=139)

and without (N=170) CHX, randomly sampling the same number of reads from both groups of

experiments. We observed a large enrichment in ribo-seq read counts among nORFs with CHX

treatment ($p < 10^{-10}$, Fisher's exact test, **Figure 2F**), resulting in 56% more nORFs identified as translated

($p < 10^{-10}$, Fisher's exact test). The nORFs identified as translated only with CHX treatment nevertheless

displayed a strong collective signal of triplet periodicity (i.e., preferential mapping to the first position in

the codon) in experiments without CHX treatment when reads were aggregated across all such nORFs

(**Figure 2G**). These results indicate that CHX treatment aids detection of translation events that also

occur but are more difficult to detect without CHX.

Noncanonical translation patterns depend on genomic and environmental context

`

We examined to what extent translation of nORFs depends on genomic context. We classified nORFs as:

upstream nORFs (uORFs) located on the 5' untranslated regions of transcripts containing cORFs;

downstream nORFs (dORFs) located on the 3' untranslated regions of transcripts containing cORFs;

intergenic nORFs that do not share transcripts with cORFs (independent); nORFs antisense to a cORF

and located entirely within the bounds of that cORF (antisense full overlap); nORFs overlapping the

boundaries of a cORF on the opposite strand (antisense partial overlap) (**Figure 3A**). Additionally, for

nORFs sharing a transcript with an RNA gene, the nORF was classified based on the type of RNA gene.

The transcripts used for these classifications were derived from the TIF-seq data collected by Pelechano

et al. 2014[47], which provide transcript start and end sites.

`

Figure 3: Noncanonical translation patterns depend on both genomic and environmental context. A) Potential genomic contexts for nORFs in relation to nearby canonical genes. Transcripts are defined from published TIF-seq data[47]. **B)** Counts of translated nORFs identified by iRibo (y-axis) in each considered

`

genomic context (x-axis), determined by which elements share a transcript with the nORF and its
position within the transcript. For nORFs that share a transcript with RNA genes, the annotation of the
RNA gene is specified. **C)** Proportion of nORFs detected as translated by iRibo (y-axis) in each genomic
context considered, among nORFs completely covered by a TIF-seq transcript (x-axis). **D)** Amino acid
composition of translated nORFs differs from that of translated cORFs and depends on genomic context.
Amino acid frequencies among predicted protein products of translated nORFs in each genomic context
and of cORFs. The start codon methionine is excluded from frequency estimates. **E)** Amino acid
composition of translated nORFs is similar to that of context-matched controls. For each genomic
context, the amino acid frequency of translated nORFs relative to that of length-matched untranslated
nORFs in that same context. The start codon methionine is excluded from frequency estimates. **F)** More
nORFs are identified as translated in minimal than rich media. Number of translated nORFs identified (y-
axis) for experiments on yeast grown in either minimal (SD, solid line) or rich media (YPD, dashed line) at
a range of read depths (x-axis). For each read depth, reads are sampled at random from experiments in
each condition. **G)** For each nORF called translated by iRibo in minimal media (SD), rich media (YPD), or
both, the log reads per base in each condition is indicated. Total read count in each condition was held
constant by randomly sampling reads from YPD experiments until the read count in SD experiments was
matched. nORFs with significantly more reads in one condition than the other are colored, green for SD
and brown for YPD. Lists of nORFs with significantly different translation rates were obtained as follows:
p-values for differential translation of each nORF were calculated from Fisher's exact test on in-frame
ribo-seq reads mapping to the ORF in each condition and a 5% FDR was set using the Benjamini-
Hochberg approach.[48] An nORF had to be detected as translated in a condition by iRibo to be identified
as more highly translated in that condition.

Most nonoverlapping translated nORFs were independent (6,373, 52%) and around 47% shared a

transcript with a cORF, including 3,512 uORFs and 2,278 dORFs, while 1.5% (186) shared a transcript

with an annotated RNA gene (**Figure 3B**). Among antisense nORFs, 73% (4,844) overlapped fully with the

opposite-strand gene while 27% (1,760) overlapped partially.

We next calculated the frequency at which candidate nORFs were identified as translated for each

genomic context (**Figure 3C**); for purposes of comparison, we considered only those nORFs fully

contained within a TIF-seq transcript. Consistent with prior research[49], uORFs were translated at

significantly higher rates than other classes, with 30% of considered uORFs found to be translated

compared to only 17% of dORFs (p < 10^{-10}, Fisher's Exact Test) and 20% of independent nORFs (p < 10^{-10},

Fisher's Exact Test). nORFs antisense to cORFs and only partially overlapping them were translated at

the lowest rate of any context, with a rate of 10% compared to 26% for fully overlapping antisense

nORFs (p < 10^{-10}, Fisher's Exact Test).

The amino acid frequencies of the proteins expressed from translated nORFs differ greatly from those of

cORFs and depend on genomic context (p<10^{-10} for any comparison between cORF amino acid

`

frequencies and nORF frequencies in a given context, chi-square test; **Figure 3D**). The translation

products of nORFs present a large excess of cysteine, phenylalanine, isoleucine, arginine, and tyrosine

and deficiency in alanine, asparagine, glutamic acid, and glycine relative to cORFs. Notably, aside from

arginine, the amino acids with large excess in nORFs relative to cORFs are all hydrophobic. Amino acid

frequencies of nORFs appear to largely reflect underlying DNA sequence composition biases that differ

between the distinct genomic contexts. Indeed, within each genomic context, amino acid frequencies of

translated nORF are generally similar (with less than 15% difference in frequency) to that of length- and

context- matched nORFs that lack evidence of translation, though they do show significant differences

(p < 10^{-10} for all contexts, chi-square test; **Figure 3E**). The most striking differences include a large excess

of methionine residues and a deficiency in tryptophan and glycine residues among translated nORFs

compared to the untranslated control group.

In addition to genomic context, we assessed how environmental context affects noncanonical

translation. To this aim, we leveraged the power of iRibo to construct separate datasets of nORFs found

translated in rich media (YPD) or in nutrient-limited minimal media (SD) (**Supplementary Table 3**).

Previous research has reported an increase in detected noncanonical translation events relative to

canonical translation events in response to starvation.[1,3] Consistent with these results, more nORFs were

identified as translated in minimal than in rich media at equal read counts (**Figure 3F**). Furthermore,

2968 nORFs were supported by a significantly higher number of in-frame reads in minimal media than

rich media while the converse was true for only 1265 nORFs (5% FDR, Fisher's exact test with Benjamini-

Hochberg procedure[48]; **Figure 3G**). These results suggest that starvation conditions may increase

noncanonical translation, or alternatively that noncanonical translation is less affected by the general

translation inhibition that occurs in starvation conditions.[50] Either way, these results support the

hypothesis that nORF translation is regulated in response to changing environments.

Two translatomes, transient and conserved

Given the large numbers of nORFs translated in the yeast genome, we next sought to assess the

biological significance of this translation by determining the extent to which these nORFs are evolving

under selection. We assessed selection acting on nORFs, as well as on cORFs for purpose of comparison,

across three evolutionary scales. At the population level, we analyzed 1011 distinct *S. cerevisiae* isolates

sequenced by Peter et al. 2018.[40] At the species level, we compared *S. cerevisiae* ORFs to their orthologs

in the *Saccharomyces* genus, a taxon consisting of *S. cerevisiae* and its close relatives.[51] To detect long

term evolutionary conservation, we looked for homologs of *S. cerevisiae* ORFs among 332 budding yeast

`

genomes (excluding *Saccharomyces*) in the subphylum *Saccharomycotina* collected by Shen et al. 2018.[41]

The power to detect selection on an ORF depends on the amount of genetic variation in the ORF

available for evolutionary inference, which in turn depends on its length, the density of genetic variants

across its length, and the number of genomes available for comparison. Given that many translated

nORFs are very short (**Figure 2C**), we employed a two-stage strategy to increase power for detecting

signatures of selection. First, we investigated selection in a set of "high information" ORFs for which we

have sufficient statistical power to potentially detect selection. Second, we investigated the remaining

"low information" ORFs in groups to quantify collective evidence of selection (**Figure 4A**). Group level

analysis increases power to detect the presence of selection but does not enable identification of the

individual ORFs under selection. The "high information" set consisted of the ORFs that 1) have

homologous DNA sequence in at least four other *Saccharomyces* species and 2) have a median count of

nucleotide differences between the *S. cerevisiae* ORF and its orthologs of at least 20. We found these

criteria are sufficient to distinguish ORFs evolving under strong purifying selection (**Supplementary**

Figure 3). Under this definition, 9,440 translated ORFs that do not overlap a different cORF (henceforth

"nonoverlapping ORFs", including 4,248 nORFs, and 5,192 cORFs) and 3,022 ORFs that overlap a cORF on

the opposite strand ("antisense ORFs", including 2,962 nORFs and 60 cORFs) were placed in the "high

information" set.

`

`

Figure 4: Two distinct translatomes: transient and conserved. A) Selection inference analyses
conducted on low-information and high-information ORFs to classify them as evolutionarily conserved,
transient, or unclassified. **B)** A bimodal distribution of reading frame conservation (RFC) among high
information translated ORFs. The distribution of RFC (x-axis), indicating how well reading frame of the
ORF is conserved in the Saccharomyces genus, is shown for all translated high information ORFs (top),
only cORFs (middle) and only nORFs (bottom). See Methods for details. Dashed lines separate RFC < 0.6
and RFC > 0.8, the thresholds used to distinguish ORFs preserved or not preserved by selection. **C)** No
evidence of purifying selection acting on low information nORFs. pN/pS and dN/dS ratios are shown for
each group of ORFs. Low information nonoverlapping nORFs that lack a conserved homolog are divided
into deciles of translation rate (in-frame ribo-seq reads per base), coding score, or ORF length, and into
three genomic contexts. Untranslated nORFs are the set of all nORFs in the genome not called as
translated by iRibo. Low information nonoverlapping cORFs are assembled into a single group, with the
set of all nonoverlapping cORFs shown for comparison. Low information antisense nORFs were also
assembled into a single group, with the set of all antisense cORFs shown for comparison. pN/pS is
calculated from variation at each ORF codon among *S. cerevisiae* isolates.[40] dN/dS is calculated among
all codons that share the same frame between *S. cerevisiae* ORFs and aligned orthologous ORFs in *S.*
paradoxus. Note that the displayed pN/pS and dN/dS values are not averages of these ratios among
ORFs. Rather, synonymous and nonsynonymous variants among all ORFs in each class are counted, and
a single ratio is calculated from the summed counts. Error bars indicate standard errors estimated from
bootstrapping. The dashed blue line indicates a ratio of one, the expected ratio under neutral evolution.
D) No evidence of distant homology for low information nORFs. The frequency of nORFs with weak
TBLASTN matches (10^{-4} < e-value < .05) in each group of nORFs (dark bars) and negative controls (light
bars) consisting of the sequences of the nORFs of each group randomly scrambled. Error bars indicate
standard errors estimated from bootstrapping. **E)** ORFs that are translated yet evolutionarily transient
make up 72% of the yeast reference translatome. The components of the translatome (transient,
conserved, unclassified) are represented with area proportional to frequency. Each box represents sets
of 15 ORFs.

We attempted to detect purifying selection in the high information set within the *Saccharomyces* genus

and within the *Saccharomycotina* subphylum. For the *Saccharomyces* analysis, we adapted reading

frame conservation (RFC), a sensitive approach developed by Kellis et al. 2003[20] to distinguish ORFs

evolving under selection from other ORFs in the yeast genome. RFC is an index ranging from 0 to 1 that

indicates how well reading frame is conserved between an ORF in a given species (here, *S. cerevisiae*)

and its orthologous sequences in related species (other species in the *Saccharomyces* genus). An RFC

value of 1 indicates perfect agreement of reading frame, such that all bases that make up the first

nucleotide in a codon in the *S. cerevisiae* ORF also make up the first nucleotide in a codon in each

orthologous ORF. An RFC value of 0 indicates that all bases in the *S. cerevisiae* ORF align to bases with a

different within-codon position in orthologous ORFs, or that the aligned bases exist outside of any ORF.

We found a bimodal distribution of RFC among nonoverlapping ORFs in the yeast translatome,

considering cORFs and nORFs together: 53.3% have RFC above 0.8 and 45.5% have RFC less than 0.6,

`

with only 1.2% of ORFs intermediate between these values (**Figure 4B)**. The bimodal distribution of RFC

among translated ORFs is similar to the bimodal distribution observed among all candidate ORFs,

regardless of translation status (**Supplementary Figure 4A**), as observed previously by Kellis et al. 2003.[20]

The modes of the distribution largely correspond to annotation status, with 96.7% of cORFs having RFC >

0.8 and 98.5% of nORFs having RFC < 0.6. This pattern holds when evaluated only in the last 100 bp of

ORFs, suggesting that it is not affected by potential incorrect inference of nORF start positions

(**Supplementary Figure 4B**). The clean separation between well-conserved and poorly-conserved ORFs

indicate that most high-information ORFs can be straightforwardly classified into one of the two groups,

and thus nearly all high-information nonoverlapping nORFs can be placed in the poorly-conserved class.

High RFC among antisense ORFs does not demonstrate selection on the ORF itself, as it might be caused

by selective constraints on the opposite-strand gene, but low RFC still indicates lack of purifying

selection. A majority of antisense translated nORFs (64.1%) have RFC <0.6, indicating that most are not

preserved by selection across the genus (**Supplementary Figure 4C**). Overall, we find no evidence for

purifying selection acting on nORFs on a large scale.

In light of the general correspondence between annotation and conservation, the exceptions are of

interest: 110 cORFs had RFC < 0.6, and 13 nonoverlapping unannotated nORFs had RFC > 0.8. To further

assess conservation among these two sets of ORFs, we performed a BLAST analysis (using both BLASTP

and TBLASTN with default parameters) to search for homologs of each ORF among the budding yeast

genomes assembled by Shen et al. 2018.[41] Among the 110 cORFs with low RFC, 101 also had no detected

homology to other *S. cerevisiae* genes or any budding yeast genome outside of *Saccharomyces*,

indicating that these are likely annotated ORFs of recent *de novo* origin. For the 13 nORFs with high RFC,

several additional lines of evidence suggest that these are indeed evolving under purifying selection

(**Table 1)**. For nine of the thirteen, we identified a homolog among budding yeast genomes outside of

the *Saccharomyces* genus by either a BLASTP or TBLASTN search. The existence of a homolog in a

distantly related species indicates that the ORF existed in the common ancestor of *S. cerevisiae* and that

distant species, implying long-term preservation of the ORF by purifying selection in both lineages. We

also performed pN/pS analysis for each ORF on *S. cerevisiae* isolates and dN/dS analysis for each ORF

among the *Saccharomyces* genus species (**Table 1**). A pN/pS or dN/dS ratio significantly below 1

indicates purifying selection on the ORF amino acid sequence among *S. cerevisiae* strains or among

Saccharomyces genus species, respectively, while a ratio above 1 indicates positive selection. By these

measures, two ORFs showed significant evidence of purifying selection by pN/pS and three by dN/dS

`

(**Table 1**). Thus, a small number of nORFs appear to be evolving under selection, indicating significant

biological roles.

We next assessed selection among the full set of nORFs (both high and low information) at the

subphylum scale, searching for addition nORFs that exhibited long term conservation and thus purifying

selection. Towards this end, we searched for distant homologs of all translated nonoverlapping *S.*

cerevisiae nORFs using TBLASTN against budding yeast genomes in the *Saccharomycotina* subphylum,

excluding species in the *Saccharomyces* genus. After excluding matches that appeared non-genic or

pseudo-genic (**Supplementary Figure 5**) we identified a single additional nORF with both distant

TBLASTN matches and recent signatures of purifying selection (dN/dS = 0.5, p=.039 for test of difference

from 1.0): YBR012C, annotated as "dubious" on SGD. Thus, combining the 13 nORFs that appeared

conserved by RFC analysis and the single additional nORF found using TBLASTN, we identified 14

translated nORFs that show evidence of preservation by purifying selection (**Table 1**).

To analyze collective evidence of selection among "low information" ORFs, we first divided low

information nonoverlapping nORFs (7,855 nORFs, after excluding those with homology to conserved *S.*

cerevisiae cORFs) according to properties that we expected to be potentially associated with selection:

rate of translation (as measured by ribo-seq reads mapped to the first position within codons divided by

the length of the ORF), coding score[28,52] (a measure of sequence similarity to annotated coding

sequences), ORF length, and genomic context. For each group, we calculated the pN/pS ratio among

1,011 *S. cerevisiae* isolates[40] and the dN/dS ratio based on alignments of the *S. cerevisiae* ORFs with their

orthologous DNA sequence in *S. paradoxus*. We also analyzed low information nonoverlapping cORFs

(22 cORFs) in the same manner. For low information antisense nORFs (3642 nORFs; only 2 cORFS fell in

this category and were not analyzed), we calculated the pN/pS and dN/dS ratios restricted to

substitutions that were synonymous on the opposite-strand cORF.[53,54] Unlike the RFC, dN/dS and pN/pS

analyses conducted above on individual high information ORFs, these analyses were conducted by

aggregating substitutions among all low information ORFs in each group to assess evidence for selection

(i.e., a ratio significantly different from 1) within the group as a whole. We expected that, if low

information nORFs were evolving under selection, then more highly translated ORFs, longer ORFs, and

ORFs with coding scores more similar to conserved genes, would be enriched in biologically relevant

nORFs and thus show stronger signatures of selection. Low information nonoverlapping cORFs did show

collective pN/pS and dN/dS ratios significantly below 1, indicating that some ORFs in this group are

evolving under purifying selection (**Supplementary Table 4**, **Figure 4C**). In contrast, for all groups of low

`

information nORFs examined, we observed no significant difference in the pN/pS or dN/dS ratio from 1,

providing no evidence for either purifying or positive selection (**Supplementary Table 4, Figure 4C**).

Finally, we assessed collective evidence of long-term evolutionary conservation in each group. To do

this, we calculated the frequency of weak TBLASTN matches (e-values between 10^{-4} and .05, above our

threshold for homology detection at the individual level) of ORFs in each group to the *Saccharomycotina*

subphylum genomes outside of *Saccharomyces* as compared to a negative control set consisting of

scrambled sequences of the ORFs in each group. Applying this strategy to the full set of 362

nonoverlapping cORFs that lacked TBLASTN matches outside *Saccharomyces* at the e-value < 10^{-4} level,

we found a large excess of weak matches relative to controls (p=.0001, Fisher's exact test;

Supplementary Figure 6), demonstrating the ability of this approach to detect faint signals of homology

within a group of ORFs. However, we identified no significant difference in the frequency of weak

TBLASTN hits between any nonoverlapping nORF group and scrambled controls (**Figure 4D**), nor among

nonoverlapping nORFs overall (p>.05, Fisher's exact test). The lack of a significant result does not

exclude the possibility that a small subset of short conserved nORFs could be lost in the noise of a much

larger set of nORFs without distant homology. However, our TBLASTN, dN/dS and pN/pS analyses

altogether indicate that ORFs evolving under strong purifying selection are not a major component of

the yeast noncanonical translatome.

Overall, our analyses distinguish two distinct yeast translatomes: a conserved, mostly canonical

translatome with intact ORFs preserved by selection; and a mostly noncanonical translatome where

ORFs are not preserved over evolutionary time. This distinction is rooted in evolutionary evidence rather

than annotation history. We thus propose to group the translated ORFs that showed neither evidence of

selection nor homology to conserved ORFs in our high-information and low-information sets as the

"transient translatome." The "transient translatome" designation indicates membership in a set of ORFs

that are expected to exist in the genome for only a short time on an evolutionary scale, though we

cannot be certain that any particular translated ORF that currently exists in the yeast genome will be

rapidly lost. The transient translatome includes 4,051 nonoverlapping and 1,923 antisense nORFs

identified as not preserved by selection using RFC analyses and having no conserved homologs, along

with 86 nonoverlapping and 15 antisense cORFs (total 101) matching the same criteria. Also included are

7,855 nonoverlapping and 3,644 antisense nORFs that lack sufficient information to analyze at the

individual level but were found to show no selective signal in group-level analyses. Together, this set of

`

17,574 ORFs that are translated yet likely evolutionarily transient makes up 72% of the yeast reference

translatome (**Figure 4E**).

Transient cORFs are representative of the transient translatome overall

By general theory and practice in evolutionary genomics, the lack of selective signal suggests that the

transient translatome does not meaningfully contribute to fitness.[55] Surprisingly, however, 101 cORFs

belong to the transient set, suggesting that some transient ORFs have phenotypes. To assess whether

these cORFs are representative of the transient translatome overall, we compared their evolutionary

and sequence properties with those of transient "dubious" nORFs (annotated but presumed

nonfunctional) and transient unannotated nORFs. We found transient cORFs, transient dubious nORFs

and transient unannotated nORF to all have pN/pS ratios indistinguishable from 1.0 (**Figure 5A**),

providing no evidence for purifying selection. Similarly, the average nucleotide diversity (mean number

of nucleotide differences per site between pairs of isolates) of transient cORFs was indistinguishable

from that of transient nORFs or untranslated controls, and much higher than that of conserved cORFs

(**Figure 5B**). In addition, no class of transient ORFs showed differences from each other in RFC between

S. cerevisiae and *S. paradoxus* (**Figure 5C**), rate of translation (**Figure 5D**) or coding score (**Figure 5E)**.

`

Figure 5: Canonical and noncanonical transient ORFs are similar except for their length. Properties of transient cORFs (n=86), transient dubious nORFs (n=25), transient unannotated nORFs (n=12,160), untranslated controls (n=100) and conserved cORFs (n=5130). Untranslated controls consist of nonoverlapping ORFs that would be grouped in the transient class (RFC <.6) but are not inferred to be translated based on ribo-seq evidence. Conserved cORFs are nonoverlapping cORFs with RFC >.8. All groups are restricted to nonoverlapping ORFs. Error bars represent standard error. Stars indicate significant differences from untranslated controls by permutation test: P-value <.001: ***. **A)** pN/pS values for each group among *S. cerevisiae* strains. **B)** Average nucleotide diversity (π) among each group. **C)** Average reading frame conservation between *S. cerevisiae* and *S. paradoxus* ORFs. **D)** Average ribo-seq reads per base (logged), considering only in-frame reads. Unannotated nORFs and untranslated controls are sampled to match the length distribution of transient cORFs. **E)** Coding scores for each group. **F)** ORF lengths in nucleotides for each group.

`

The only distinguishing property between classes of transient ORFs was their length: annotated

transient cORFs and transient "dubious" nORFs are much longer on average than unannotated transient

nORFs (**Figure 5F**). This is a consequence of the history of annotation of the *S. cerevisiae* genome, where

a length threshold of 300 nt was set for annotation of ORFs.[56,57] The sharp 300 nt threshold is still clearly

reflected in annotations. For example, genome annotations include 96% of nonoverlapping transient

ORFs in the 300-400 nt range (55/57), but only 4% in the 252-297 nt range (4/101). Given that transient

nORFs resemble transient cORFs in all respects besides length, we hypothesized that numerous never-

studied transient nORFs are just as likely to have phenotypes as transient cORFs.

Transient ORFs are detected in the cell and mediate diverse phenotypes

To gain further insights into the potential biological roles of transient ORFs, we examined published

reports about annotated ORFs (transient cORFs and transient dubious nORFs) in the *S. cerevisiae*

experimental literature and performed additional experiments to investigate transient unannotated

nORFs. We examined whether transient ORF products could be detected experimentally, whether they

affect phenotypes, and whether they interact with specific biological pathways.

We first assessed whether the proteins encoded by transient ORFs can be detected in the cell. We

examined the CYCLoPs database[58,59], the C-SWAT tagging library[60], and the YeastRGB database[61], which

contain collections of fluorescently tagged proteins expressed from their native promoters and

terminators, including both cORFs and dubious nORFs. Together these studies detected expression of a

fluorescent protein product for 90 of 93 (97%) transient cORFs tested, along with 37 of 41 (90%)

transient dubious nORFs tested (**Figure 6A**). For comparison, we C-terminally tagged 21 highly expressed

unannotated transient nORFs with mNeonGreen at their endogenous locus and examined their

expression using microscopy. We detected 8 of 21 tagged nORF proteins (38%) (**Figure 6A-B,**

Supplementary Figure 7). Thus, translation of tagged proteins can be detected for both annotated and

unannotated transient ORFs.

`

Figure 6: Transient nORFs and cORFs can be detected in the cell and exhibit phenotypes. A) Transient
ORFs are detected by fluorescent microscopy. For cORFs or dubious nORFs, the proportion of proteins
expressed by transient ORFs detected in the C-SWAT[60], CYCLoPs[59], or YeastRGB[61] microscopy datasets
out of those tested. For unannotated transient nORFs, the proportion detected by mNeonGreen tagging

`

in this study. Error bars indicate standard error of the proportion. **B)** Tagged unannotated transient
nORFs show varied sub-cellular localizations. Microscopy images of unannotated transient nORFs taken
at 100X. Left panel shows the expression of the nORFs tagged with mNeonGreen, middle panels the
dyes CMAC Blue and MitoTracker Red for mitochondria and vacuoles identification, respectively, and
the right panel the merge all the above channels with DIC. Top panel show the nORF (orf45629) with a
cytosolic expression and the bottom panel the nORF (orf231865) with expression localizing to the
mitochondria. **C)** Loss of transient nORFs can affect phenotype despite lack of evolutionary conservation
The proportion of deletion mutants with reported loss-of-function phenotypes in two groups: transient
cORFs in published deletion mutant screens, and transient nORFs assayed in this study. Reported
phenotypes in published data was taken from literature associated with each ORF on SGD. In this study,
deleterious deletion mutant phenotypes were identified from a high-throughput colony fitness screen in
six stress conditions using a 5% FDR threshold. **D)** Transient ORFs engage in epistatic relationships. The
percent of transient ORFs and nonessential genes with at least one genetic interaction at given
threshold are shown. Differences between groups were tested using Fisher's exact test. **E)** Genetic
interactions of the transient ORF YER175W-A. Five interactors are related to exosome (striped circles). **F)**
Presence of phenotypes among annotated transient ORFs. "Protein detected" indicates that the ORF
product was found in either the C-SWAT or CYCLoPs database. Phenotypes of deletion collection,
deletion and overexpression screens were taken from reported findings in the yeast experimental
literature (**Supplementary Table 5**). "Genetic interaction" indicates a statistically significant genetic
interaction with Ɛ< -0.2, and "GO-associated interactors" indicates a GO enrichment was found among
significant interactors at 5% FDR.

We next examined the evidence that transient ORFs affect phenotype. Five transient cORFs have been

studied in depth. Two of these, *MDF1*[62] and YBR196C-A[63], have been previously described as having

emerged *de novo* from non-genic sequences. *MDF1* inhibits the mating pathway in favor of vegetative

growth[62,64] and YBR196C-A is an ER-located transmembrane protein whose expression is beneficial

under nutrient limitations.[65] The remaining three have been experimentally characterized, although

their evolutionary properties were not analyzed in the corresponding studies: *HUR1* plays an important

role in non-homologous end-joining DNA repair[66]; *YPR096C* regulates translation of *PGM2*[67]; *ICS3* is

involved in copper homeostasis.[68] These cases demonstrate that some transient ORFs do affect

phenotypes and have the potential to play important biological roles.

To determine whether transient cORFs that are not well described also affect phenotypes, we examined

all literature listed as associated with the ORF on SGD. Many of these transient cORFs have direct

evidence of phenotype (**Supplementary Table 5**). Of 101 transient cORFs, 45 were reported to have

deletion mutant phenotypes (i.e., a phenotype observed when the ORF is deleted) and 12 to have

overexpression phenotypes. Overall, we found phenotypes reported in the literature for 50 of 101

transient cORFs (50%).

`

As unannotated transient nORFs have not been systematically investigated for phenotype, we sought to

experimentally determine whether these ORFs too might have deletion mutant phenotypes. We thus

conducted a deletion mutant screen of 49 unannotated transient nORFs selected for high translation

rate and to avoid intersecting cORFs, annotated ncRNAs, or promoters (200 bp upstream of canonical

genes). We fully deleted the nORF using homologous recombination and each strain was assayed for

colony growth in seven conditions. Eight nORF deletion mutant strains showed deleterious phenotypes

in at least one condition at a 5% FDR (**Figure 6C**, **Supplementary Table 6**). Thus, loss of transient nORFs,

as with cORFs, can affect phenotype despite lack of evolutionary conservation.

To begin to understand the specific biological processes in which transient ORFs might be involved, we

leveraged the large yeast genetic interaction network assembled in Costanzo et al. 2016.[69] This dataset

includes 75 non-overlapping transient cORFs and 9 non-overlapping dubious transient nORFs. Genetic

interaction strength, Ɛ, measures the difference between the observed fitness of a strain in which two

genes are deleted and the expected fitness given the fitness of the two single gene deletion strains; a

negative value of high magnitude suggests that the two mutated genes are involved in related

processes. Of the 84 transient ORFs in the dataset, 79 (94%) have at least one negative genetic

interaction at the high-stringency cut-off defined by Costanzo et al.[69] ($\varepsilon < -0.2$ and p-value<0.05) and 51

(61%) have synthetic lethal interactions ($\varepsilon < -0.35$ and p-value<0.05) as defined in that study (**Figure 6D**).

This was only a slightly lower rate than for conserved non-essential ORFs, 98% of which had negative

interactions at the high stringency cut-off and 76% of which had synthetic lethal interactions. At the high

stringency threshold, 27 transient ORFs were found to interact with groups of related genes enriched in

specific gene ontology (GO) terms (5% FDR; **Supplementary Table 7**). For example, the interactors of

YER175W-A are associated with the GO category "cryptic unstable transcript (CUT) metabolic processes"

with high confidence, and five of its eleven interactors are components or co-factors of the exosome

(**Figure 6E**), indicating likely involvement in CUT degradation or a closely related post-transcriptional

regulation pathway. Other enrichments included diverse processes such as "mating projection tip" or

"Golgi sub-compartment". In contrast, when we applied GO enrichment analysis to the full set of genes

that interact with any transient ORF, no significant enrichment was observed. These results suggest that

transient ORFs in general do not participate in one shared biological process, but rather are involved in a

wide variety of cellular processes.

Overall, we uncovered evidence that 131 of 250 (53%) annotated transient ORFs have at least one

indicator of biological significance (detection of a protein product, a reported phenotype in a screen, or

a genetic interaction in the Costanzo et al. 2016[69] network) (**Figure 6F**). Additionally, we demonstrate

that unannotated transient ORFs encode proteins that can be detected in the cell (38% of tested in this

study) and influence cellular fitness when deleted (17% of tested in this study). Given that this class has

received almost no study compared to the great number of experiments that have been conducted on

cORFs, the number of transient ORFs with biological relevance may be substantially larger than that

which has been annotated.

A limitation on much of the experimental evidence available on deletion mutant phenotypes is that

most deletion mutant and genetic interaction screens are based on a full gene replacement strategy in

which the entire ORF is lost, leaving the possibility that some deletion phenotypes could be caused by

loss of a ncRNA or a DNA regulatory element located at the same position as the ORF rather than loss of

the ORF translation (**Figure 7A**). To examine this possibility, we constructed a set of strains where the

ORF start codon ATG was replaced with an AAG codon while keeping the rest of the ORF intact. This set

included three transient cORFs that have previously been characterized on the basis of overexpression

or full deletion mutants, *ICS3*[68], YPR096C[67], and YBR196C-A[65], along with four transient nORFs that

showed strong deleterious phenotypes in our full ORF deletion screen (**Supplementary Table 8**; *HUR1*

and *MDF1* were not tested because they overlap other cORFs). Each deletion strain was tested in seven

environmental conditions. The single nucleotide ATG→AAG mutation caused significantly reduced

colony size for all three transient cORFs tested and for three of four transient nORFs tested in at least

one condition (**Figure 7B**). We gave these three nORFs systematic names YDL204W-A, YGR016C-A, and

YNL040C-A. The remaining nORF, YDR073C-A, showed a weak beneficial phenotype from the ATG→AAG

mutation in some conditions, as did two other nORFs, YGR016C-A and YNL040C-A. The largest growth

reductions were observed from disabling translation in YDL204W-A: this strain reached only 64% of

wildtype growth in hydroxyurea and 63% in high salt concentration, with a smaller reduction to 94%

growth in rich media (YPDA). These growth defects were also observed in a liquid growth setting (**Figure

7C-E**). To confirm that these phenotypes were caused by loss of the YDL204W-A protein rather than *cis*

effects at the locus, we expressed the intact YDL204W-A ORF from a plasmid in the ATG→AAG mutant

strain. Plasmid expression of the ORF fully restored the wildtype phenotype in the mutant strains (**Figure

7F-H**), providing further evidence that blocking YDL204W-A translation causes a loss of function

phenotype mediated by loss of the encoded protein.

In our translation dataset, YDL204W-A has a translation rate at the top percentile among transient ORFs

(**Figure 7I**), higher than 10% of cORFs. Comparing its sequence to the homologous region of other

`

Saccharomyces genus species, only *S. paradoxus* and *S. mikatae* have a homologous start codon, but a

2bp insertion in *S. cerevisiae* results in a frameshift such that little of the ORF is shared in any other

species (**Figure 7I**); thus, this ORF has a reading frame conservation score of only 0.2 (**Table 2**). The other

transient ORFs with phenotypes induced by an ATG→AAG mutation also showed no signs of selection

(Table 2). Thus, our results exemplify the potential for unannotated coding sequences with no evident

evolutionary conservation to affect cellular phenotypes and fitness.

`

Figure 7: Translation inhibition of transient ORFs causes strong phenotypes. A) A two-step strategy for inhibiting nORF translation. An ORF may overlap a DNA regulatory element or an RNA with a noncoding function (Wildtype ORF), both of which are disrupted in a gene replacement strategy in addition to the loss of translation (Full gene replacement). This creates ambiguity in interpreting comparisons between deletion mutants and wildtype strains. Following a deletion screen using gene replacement, we used a second round of homologous recombination to restore either the full ORF (Restored ORF) or an ORF

`

with its start codon mutated from ATG to AAG (Translation inhibition). As these mutants differ only by
this single base, the specific effects of translation inhibition can be inferred. **B)** Inhibiting translation of
transient ORFs triggers colony growth phenotypes. The fitness of AAG mutants (translation inhibition) is
shown for seven transient ORFs under stress conditions (colors). Fitness is assessed by comparing colony
size between AAG mutants and ATG controls (restored ORFs). A cross symbol after the ORF names
indicates unannotated nORFs assigned systematic names in this study. Relative fitness values
significantly different from one are indicated as follows: *p<.05 **p<.01 ***p<.001. **C-E)** Deleterious
impact of inhibiting translation of transient nORF YDL204W-A in a liquid growth assay. Liquid growth
curve of a strain in which YDL204W-A translation is inhibited by mutating its start codon (AAG) and a
strain with the initial codon as ATG in: 1M NaCl (C), 100mM hydroxyurea (D), and YPDA (E), with three
technical replicates for each strain. **F-H)** Expression from plasmid restores wildtype growth to YDL204W-
A start codon mutants. Liquid growth curves of an attempted rescue of the YDL204W-A AAG mutant by
expressing intact YDL204W-A from a plasmid. The AAG start codon mutants were transformed with
either an empty plasmid or a plasmid expressing the intact ORF; the ATG controls were transformed
with an empty plasmid. All strains were then assayed for growth in liquid media in either 1M NaCl (F),
100 mM hydroxyurea (G) or YPDA (H) with three technical replicates each. The shaded area covers 1 SD
from the mean OD value among replicates. **I)** YDL204W-A is translated and not conserved. Top:
ribosome profiling reads mapped by iRibo to YDL204W-A show triplet periodicity. Bottom: alignment of
the YDL204W-A ORF against homologous DNA in the *Saccharomyces* genus.

Discussion

Since the advent of ribosome profiling, it has been evident that large parts of eukaryotic genomes are

translated outside of canonical protein-coding genes[1], but the nature and full significance of this

translation has remained elusive. To facilitate study of this noncanonical translatome, we developed

iRibo, a framework for integrating ribosome profiling data to sensitively detect ORF translation across a

variety of environmental conditions. The iRibo framework can be applied to any species and set of

candidate ORFs of interest. Here, we deployed iRibo to map a high confidence yeast reference

translatome almost five times larger than the canonical translatome. This resource can serve as the basis

for further investigations into the yeast noncanonical translatome, including the prioritization of nORFs

for experimental study.

We designed iRibo to be highly sensitive at detecting patterns of triplet periodicity through the genome,

but there are some limitations to our strategy. We focused exclusively on ORFs with AUG start codons

and therefore missed the non-AUG codons that are sometimes used as starts.[70] Similarly, we did not

consider ORFs overlapping canonical genes in a different frame on the same strand, though some such

nORFs are known to be translated.[71,72] Finally, candidate ORFs were selected as the longest ORF in any

reading frame, which means the true boundaries of identified ORFs could be shorter than described. We

`

expect these limitations to cause underestimation of the number of translated nORFs, suggesting that

the true count is even larger than identified here.

We used the iRibo yeast reference translatome to address a fundamental question: to what extent does

the noncanonical translatome consist of conserved coding sequences that were missed in prior

annotation attempts? In a thorough evolutionary investigation, we identified 14 translated nORFs that

show evidence of being conserved under purifying selection. Only one of these ORFs, YJR107C-A,

appears to have been previously described[34], though it was not annotated on Saccharomyces Genome

Database at the time of our analysis. Thus, even a genome as well-studied as *S. cerevisiae*'s contains

undiscovered conserved genes, likely missed in prior analyses due to difficulties in analyzing ORFs of

short length. These 14 nORFs are, however, the exception: the great majority of translated nORF show

no signatures of selection, comprising a large pool of evolutionarily transient translated sequences.

The yeast genome thus encodes two translatomes, one conserved, one transient. The conserved

translatome consists of coding sequences that are preserved by strong purifying selection and usually

have a long evolutionary history. They tend to be relatively long, well expressed, and with sequence

properties highly distinct from noncoding sequences. The transient translatome, by contrast, is

evolutionarily young, of recent *de novo* origin from previously noncoding sequence and still similar to

noncoding sequences in nucleotide composition. Evolving in the absence of strong purifying selection,

transient translated ORFs appear to be frequently lost to disrupting mutations, only to be replaced by

other transient translated ORFs upon translation-enabling mutations. Despite these profound

differences, transient translated ORFs, like conserved ones, can affect the phenotype and fitness of the

organism. Several well-characterized coding sequences unique to *S. cerevisiae*, such as *HUR1*[66] and

MDF1[62], play key roles in biological processes through encoding lineage-specific proteins that physically

interact with conserved proteins. Additionally, around 100 transient ORFs are annotated as coding genes

and have therefore been extensively screened; a majority express stable proteins and many have known

loss-of-function phenotypes. Their genetic interaction patterns suggest involvement in a wide array of

specialized cellular processes. Our experiments revealed that disabling the start codons of unannotated

transient translated ORFs can cause large fitness reductions in stress conditions. The strength of the

fitness reduction observed was highly dependent on the stressor applied in the environment, suggesting

again specialized cellular roles. In some cases, disabling the start codon resulted in growth increases,

perhaps indicating that disabling translation saved the cell energy.

`

Our work adds to the growing research on the roles noncanonical coding play across many species,

including humans.[7,73] We note that "noncanonical" is not a coherent biological category, as it simply

indicates the class of sequences that have not been annotated in genome databases. We demonstrate

that the division between "canonical" and noncanonical" translation in *S. cerevisiae* corresponds largely,

but not perfectly, to a biological division between transient and conserved. It is this biological division

that is fundamental: the 101 yeast canonical ORFs classified as transient have sequence and

evolutionary properties nearly identical to noncanonical transient ORFs, except for sequence length, and

should be placed in the same category. We can thus reclassify the translatome according to biology

rather than annotation history.

It is perhaps surprising that a coding sequence can affect organism phenotype despite showing no

evidence of selection. However, this result is consistent with evidence from the field of *de novo* gene

birth. Species-specific coding sequences have been characterized in numerous species.[32] For example,

Xie et al. 2019[74] identified a mouse protein contributing to reproductive success that experienced no

evident period of adaptive evolution. Sequences that contribute to phenotype without conservation

have also been described outside of coding sequences. Regulatory sequences, such as transcription

factor binding sites, are a mix of relatively well-conserved elements and elements that are not preserved

even between close species[75]; it is plausible that translated sequences also show such a division. There

are several explanations for why translated ORFs may lack detectable signatures of selection. Most

transient ORFs are expressed at much lower levels than canonical genes, and therefore may have

minimal effects on phenotype. For those that do have large and beneficial effects in some

environmental conditions, these may be balanced by deleterious effects in other conditions. Moreover,

selection may occur, and be biologically important, below the limits of detectability for the genomic

approaches we used. Our findings do not imply an absence of selective forces in shaping the patterns of

noncanonical translation. Rather, the particular selective environment favoring expression of these

sequences may be too short-lived to detect selection using traditional comparative genomics

approaches. Previous research, such as the proto-gene model of *de novo* gene birth[3], have proposed

that recently emerged translated ORFs serve as an intermediary between noncoding sequences and

mature genes. Our results add to the evidence that these ORFs provide many potential phenotypes from

which selection could preserve beneficial ones for the long term.[65] Still, the observation that even ORFs

with phenotypes lack evidence of conservation at the population level suggests that there are important

filters that prevent the vast majority of recently emerged translated ORFs, even those with beneficial

phenotypes, from evolving into mature genes that are preserved over long evolutionary time. The

`

primary significance of the great majority of transient translated ORFs is in their biological activity over

their short lifespans.

The yeast reference translatome resource we constructed with iRibo is meant to facilitate community

efforts to decipher the specific physiological implications of transient translated ORFs. Our proof-of

concept analyses of subcellular localization, genetic interactions and ATG->AAG mutants suggest

involvement in diverse cellular processes and pathways. It is important to note that some transient

translatome phenotypes may be mediated by a protein product, by the process of translation itself, or

both. Translation of both uORFs[76] and dORFs[77] can affect expression of nearby genes. Translation also

plays a major role in the regulation of RNA metabolism through the nonsense-mediated decay pathway.

[78,79] Dissection of the molecular mechanisms mediating transient translatome phenotypes is an exciting

area for future research.

Our results indicate that the yeast noncanonical translatome is neither a major reservoir of conserved

genes missed by annotation, nor mere "translational noise." Instead, many translated nORFs are

evolutionarily novel and likely affect the biology, fitness, and phenotype of the organism through

species-specific molecular mechanisms. As transient ORFs differ greatly in their evolutionary and

sequence properties from conserved ORFs, they should be understood as representing a distinct class of

coding element from most canonical genes. Nevertheless, as with conserved genes, understanding the

biology of transient ORFs is necessary for understanding the relationship between genotype and

phenotypes.

Acknowledgments

We thank Dr. Emmanuel Doram Levy's at the Weizmann Institute of Science for sharing the fluorescence

intensity data displayed in YeastRGB. We thank Dr. Benjamin Dubreuil for the helpful discussion over

YeastRGB data. We thank Dr. Allyson O'Donnell for her help in microscopy image acquisition. We thank

Drs. Craig Kaplan and Nikolaos Vakirlis for helpful discussions of an earlier preprinted version of this

manuscript. This work was supported by funds provided by the Searle Scholars Program to A.-R.C., the

National Science Foundation grant MCB-2144349 to A.-R.C., and the National Institute of General

Medical Sciences of the National Institutes of Health grants R00GM108865 and DP2GM137422

(awarded to A.-R.C.).

Author contributions

Conceptualization, A.W. and A.-R.C. Methodology, A.W., A.-R.C., S.B.P., N.C.C., O.A. Investigation, A.W.,

N.C.C., S.B.P., O.A., C.H., L.C. Writing – Original Draft, A.W., S.B.P., O.A., N.C.C. Writing – Review &

Editing, A.W., A.-R.C., S.B.P., N.C.C., O.A., C.H., L.C. Supervision, A.-R.C.

Declaration of interests

A.-R.C. is a member of the scientific advisory board for Flagship Labs 69, Inc (ProFound Therapeutics).

Tables

Table 1: Properties of well-conserved nORFs. Systematic name refers to either the systematic name

annotated on SGD, or the name assigned here according to SGD conventions. BLASTP and TBLASTN e-

values are the minimum BLASTP or TBLASTN e-value observed in a search of the ORF against the yeast

genomes assembled by Shen et al.[41], excluding those in the *Saccharomyces* genus. BLAST coverage is the

length of the segment that aligns to the best identified homolog (lowest e-value) in the BLAST search.

RFC is reading frame conservation of the ORF among species in the Saccharomyces genus. Length is the

length of the ORF in nucleotides. The pN/pS ratio is obtained from nucleotide variation in the ORF

among the 1011 S. cerevisiae strains assembled by Peter et al.[40]; significant values below 1 indicate

purifying selection. The dN/dS ratio was obtained from nucleotide variation in the ORF among

Saccharomyces genus species; significant values below 1 indicate purifying selection. Translation

percentile indicates the percentage of nORFs with a lower ribo-seq read count per codon than the

indicated ORF.

Systematic Name	Coordinates	BLASTP e-value	TBLASTN e-value	RFC	BLAST coverage (nt)	Length (nt)	pN/pS (p-value)	dN/dS (p-value)	Translation percentile
YBL029W-B[a]	chrII:164192-164368	6.5×10^{-4}	8.0×10^{-3}	0.82	107	177	1.65 (.33)	0.88 (.68)	67
YBL014W-A[a]	chrII:196737-196889	4.1×10^{-5}	1.0×10^{-4}	1	116	153	0.47 (.11)	$0.14 (3.46 \times 10^{-12})$	86
YBR085W-B[a]	chrII:417494-417556	1	1	0.86	0	63	0.72 (.48)	1.26 (.62)	58
YBR268W-A[a]	chrII:741844-742005	1	1	0.99	0	162	0.61 (.15)	$0.35 (3.18 \times 10^{-7})$	97
YBR292W-A[a]	chrII:786745-786903	1.9×10^{-7}	5.0×10^{-3}	0.96	146	159	0.72 (.43)	0.57 (.0026)	83
YER186W-A[a]	chrV:565603-565800	6.0×10^{-6}	1	0.92	143	198	0.55 (.02)	1.0 (1)	97
YGL262W-A[a]	chrVII:4663-4872	1	1.0×10^{-3}	0.88	113	210	0.96 (.86)	1.0 (1)	86
YGR238W-A[a]	chrVII:969015-969089	1	1	0.87	0	75	0.20 (.01)	1.18 (.74)	94
YBL049C-A[a]	chrII:126330-126461	8.3×10^{-5}	6.0×10^{-4}	0.84	92	132	1.36 (.79)	1.5 (.22)	75
YBL026C-A[a]	chrII:169634-169870	6.8×10^{-12}	9.0×10^{-10}	0.88	116	237	1.30 (.6)	0.87 (.42)	99.96
YJR107C-A[a]	chrX:628457-628693	3.8×10^{-8}	3.0×10^{-18}	0.99	161	237	0.39 (.005)	1.42 (.13)	99.91

YLR349C-A[a]	chrXII:828276-828338	1	1	0.81	0	63	0.30 (.02)	0.73 (.24)	73
YNR062C-A[a]	chrXIV:745640-745792	5.0×10^{-14}	5.0×10^{-13}	0.89	110	153	0.65 (.44)	1.49 (.15)	44
YBR012C	chrII:259147-259566	6.51×10^{-59}	1×10^{-16}	0.70	120	420	.62 (.1)	.50 (.039)	92

735 [a]We assigned this unannotated ORF a systematic name based on SGD conventions.

736 **Table 2: Evolutionary properties of transient ORFs with phenotypes induced by inhibiting translation.**

737 The pN/pS ratio is obtained from nucleotide variation in the ORF among the 1011 S. cerevisiae strains

738 assembled by Peter et al.[40] TBLASTN was run for each ORF against genomes in the subphylum

739 *Saccharomycotina*, excluding the genus *Saccharomyces*, with an e-value threshold of 10^{-4}.

ORF Name	Reading frame conservation	pN/pS (p-value)	TBLASTN matches
YBR196C-A	.29	1.34 (0.65)	0
YDL204W-A[a]	.20	1.25 (0.83)	0
YGR016C-A[a]	.29	0.66 (0.36)	0
YJL077C	.21	0.74 (0.19)	0
YNL040C-A[a]	.38	0.97 (1.00)	0
YPR096C	.20	1.39 (0.47)	0

740 [a]We assigned this unannotated ORF a systematic name based on SGD conventions.

741 **Supplementary figure legends**

`

Supplementary Figure 1: Workflow to identify translated ORFs in the *S. cerevisiae* genome using published datasets.

`

Supplementary Figure 2: Translation patterns in candidate ORFs show high replicability between studies. A) Pairwise correlation between ribo-seq coverage of all candidate ORFs between studies included in the dataset. B) For each candidate ORF, the reads per base (considering only in-frame reads) are plotted for the two largest studies in the dataset.

Supplementary Figure 3: Nucleotide variation determines ability to distinguish conserved ORFs. Reading frame conservation for each nonoverlapping ORF is plotted against the median count of

differences between the *S. cerevisiae* ORF and the aligned homologous sequence in each *Saccharomyces*

relative. Colors indicate SGD annotation categories. To the right of the vertical line, there are two

distinct populations separable by reading frame conservation; the intermediate region contains few

ORFs. For ORFs to the left of the vertical line, with few differences in the ORF between species, there is

no clear separation between high-RFC and low-RFC ORFs.

Supplementary Figure 4: Distribution of frame conservation among classes of ORFs. A) The distribution

of frame conservation among candidate ORFs in the genome, including both translated and untranslated

ORFs. B) For all ORFs in the high information set at least 100 nt in length, RFC was calculated considering

only the final 100 nt of the ORF. RFC was then plotted for both cORFs and nORFs. This was done to test

whether low RFC in nORFs could be caused by inferring start codons upstream of the actual start codons

for conserved nORF, which would lead to false inference of a low RFC value. However, the pattern

considering only the final 100 nt is similar to the pattern observed for the full ORFs in Figure 4B, with a

clear bimodal distribution, indicating that false start codon inference is likely not driving the pattern. C)

The distribution of frame conservation is plotted for translated cORFs and nORFs that are antisense to

canonical genes. In contrast to frame conservation among nonoverlapping ORFs, the distribution does

not appear bimodal.

Supplementary Figure 5: Identification of conserved genes in the noncanonical translatome using

TBLASTN. A) Process for identification of conserved nORFs evolving under purifying selection. To be

identified as conserved, an nORF could not overlap any annotated feature on the *S. cerevisiae* genome

or have any homology to an *S. cerevisiae* cORF at a 10^{-4} BLASTP e-value threshold (as this makes BLAST

results ambiguous) and have at least two identified homologs in a TBLASTN search at a 10^{-4} e-value

threshold. Then, an additional indicator of selection was required (RFC > .8, or p-value < .05 in a test of

neutrality using dN/dS or pN/pS). B) Among translated *S. cerevisiae* ORFs with a single TBLASTN hit (at a

10^{-4} e-value threshold) among budding yeasts outside the *Saccharomyces* genus, the distribution of

sequence identities with that match is plotted. The existence of only a single match together with the

prevalence of high sequence identities (>80%) suggests that the matches may be the result of genomic

contamination rather than genuine homology, so at least two matches are required to accept homology

as valid.

`

786 **Supplementary Figure 6: cORFs lacking high-confidence homologs are enriched in weak TBLASTN**

787 **matches.** The frequency of weak TBLASTN matches (10^{-4} < e-value < .05) among budding yeast genomes

788 for cORFs that lack any strong matches, and controls consisting of the same sequences randomly

789 scrambled. Error bars indicate standard errors estimated from bootstrapping.

`

791   **Supplementary Figure 7: Microscopy of detected transient nORFs.** Microscopy images of unannotated

792   transient nORFs taken at 40X. Left panel show the expression of the nORFs tagged with mNeonGreen,

793   middle panels the dyes CMAC Blue and MitoTracker Red for mitochondria and vacuoles identification,

794   respectively, and the right panel the merge all the above channels with DIC. Two representative images

`

are shown per strain; expression of orf55981 was not uniformly detected, with some cells showing

expression and some not.

Supplementary tables

Supplementary Table 1: Ribosome profiling experiments used for translation inference.

Supplementary Table 2: Ribosome profiling studies used for translation inference.

Supplementary Table 3: The yeast translatome.

Supplementary Table 4: Selection analysis of ORF groups in S. cerevisiae strains.

Supplementary Table 5: Phenotypes of canonical evolutionarily transient ORFs reported in literature.

Supplementary Table 6: Results of deletion mutant screen on transient nORFs using a gene replacement
strategy.

Supplementary Table 7: Gene ontology analysis of genetic interactors of annotated transient ORFs.

Supplementary Table 8: Information on ORFs tested in translation inhibition experiment.

Supplementary Table 9: Strains used in this study.

STAR Methods

Key resources table

Reagent or Resource	Source	Identifier
Deposited Data		
Deletion screen colony growth images	This paper	https://figshare.com/articles/dataset/A_vast_evolutionarily_transient_translatome_contributes_to_phenotype_and_fitness_-_Deletion_screen_data/21741434
C-SWAT collection	Meurer et al.[60]	Supplementary Table
YeastRGB collection	Dubreuil et al.[61]	Yeastrgb.org
CYCLoPs collection	Ko et al.[59]	https://thecellvision.org/cyclops/
Saccharomyces cerevisiae S288C reference genome	Saccharomyces genome database	S288C reference sequence R64.2.1
S. paradoxus genome	Liti et al. 2009[80]	http://www.saccharomycessensustricto.org/
S. arboricolus genome	Liti et al. 2013[81]	GCF_000292725.1
S. jurei genome	Naseeb et al. 2018[82]	GCA_900290405.1
S. mikatae, *S. bayanus var. uvarum*, *S. bayanus var. bayanus,* and *S. kudriavzevii* genome	Scannell et al. 2011[51]	http://www.saccharomycessensustricto.org/
TIF-seq data	Pelechano et al. 2014[47]	GSE39128
S. cerevisiae strain genomes	Peter et al. 2018[40]	http://1002genomes.u-strasbg.fr/files/
Budding yeast genomes	Shen et al. 2018[41]	https://y1000plus.wei.wisc.edu/data
Reagents		
Yeast Extract	BD Difco	DF0127179
Peptone	BD Difco	DF0118170
G-418	RPI	G64000-1.0
D(+) Glucose	Thermo Fisher	AAA168280E
Hygromycin B	RPI	H75020-1.0
CellTracker Blue CMAC Dye	Invitrogen	C2110
MitoTracker Red CMXRos	Invitrogen	M7512

`

Tunicamycin	Sigma	SML1287-1ML
Fluconazole	Sigma	PHR1160-1G
Sodium Chloride	Spectrum	S1240-1KG
Hydroxyurea	Thermo Scientific	A10831.14
Hydrogen Peroxide	Fisher Scientific	H323-500
DMSO	Amresco	0231-500ML
Poly(ethylene-glycol) 3350	Sigma	P4338-500G
ssDNA	Life Technologies	15632011
Lithium Acetate dihydrate	Sigma	L4158-100G
Experimental Models: Organisms, Strains		
Saccharomyces cerevisiae: BY4741	Dharmacon	YSC1048
Saccharomyces cerevisiae: BY4741, deletion collection	Dharmacon	YSC1053
Saccharomyces cerevisiae: BY4741, ORF::KanMx (mini collection with the 49 nORFs and 3 cORFs deleted)	This study	
Saccharomyces cerevisiae: BY4741, ORF-mNG:HYG (mini collection with the selected ORFs tagged with mNeonGreen)	This study	
BY4741, YDL204W-A(wt):HYG	This study	
BY4741, YDL204W-A(ATG->AAG):HYG	This study	
BY4741, YBR196C-A(wt):HYG	This study	
BY4741, YBR196C-A(ATG->AAG):HYG	This study	
BY4741, YDR073C-A(wt):HYG	This study	
BY4741, YDR073C-A(ATG->AAG):HYG	This study	

`

BY4741, YGR016C-A(wt):HYG	This study	
BY4741, YGR016C-A(ATG->AAG):HYG	This study	
BY4741, YJL077C(wt):HYG	This study	
BY4741, YJL077C(ATG->AAG):HYG	This study	
BY4741, YNL040C-A(wt):HYG	This study	
BY4741, YNL040C-A(ATG->AAG):HYG	This study	
BY4741, YPR096C(wt):HYG	This study	
BY4741, YPR096C(ATG->AAG):HYG	This study	
BY4741, YDL204W-A(wt):HYG, pAG-GPD-ccdB1-KanMx	This study	
BY4741, YDL204W-A(ATG->AAG):HYG, pAG-GPD-ccdB1-KanMx	This study	
BY4741, YDL204W-A(wt):HYG, pAG-GPD-YDL204W-A-KanMx	This study	
BY4741, YDL204W-A(ATG->AAG):HYG, pAG-GPD-YDL204W-A-KanMx	This study	
Plasmids		
pAG-GPD-ccdB1-KanMx	This study	
pAG-GPD-YDL204W-A-KanMx	This study	
Software and algorithms		
Code for analyses conducted	This paper	https://zenodo.org/badge/latestdoi/446910374
R	R	R version 4.1.2

`

BLAST	National Library of Medicine	BLAST 2.9.0+
Ontologizer 2.0	Bauer et al. 2008[83]	http://ontologizer.de/
water	EMBOSS	https://www.ebi.ac.uk/Tools/psa/emboss_water/
MUSCLE 3.8.31	Edgar 2004[84]	https://www.drive5.com/muscle/

Resources availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled
by the lead contact, Anne-Ruxandra Carvunis (anc201@pitt.edu).

Materials availability

All materials will be made available on request.

Data and code availability

- All original code has been deposited on GitHub and is publicly available as of the date of publication. DOIs are listed in the key resources table.
- Plate images for colony growth assays are available at Figshare and are publicly available as of the data of publication. DOIs are listed in the key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Experimental Model and Subject Details

Yeast strains

All strains used in this study are derived from BY4741 (Dharmacon, YSC1048). The parental strain and all
derivatives produced in this study are listed in Supplementary Table 9. The lithium acetate method[85] was
used to create new strains and selection was performed on appropriate selection plates. For genomic
integration, the inserts were PCR amplified from plasmids or GBlocks.

Method Details

Defining candidate ORFs

To identify a set of translated ORFs, a set of candidate ORFs was constructed for which translation status
could be inferred using ribo-seq data. ORFs were identified on the R64.2.1 *Saccharomyces cerevisiae*
genome assembly downloaded from SGD.[43] The initial set of candidates consisted of all possible single-

exon reading frames starting with an ATG, ending with a canonical stop codon, and having at least one

additional codon between the start and stop. Among all ORFs that shared a stop codon, all but the

longest were discarded. An ORF was considered canonical if it shared a stop codon with an ORF

annotated as "verified", "uncharacterized", or "transposable element gene" on SGD. All other ORFs that

overlapped a canonical ORF on the same strand were removed (including pairs of overlapping canonical

genes) while ORFs that overlapped cORFs on the opposite strand were classified as antisense ORFs.

Yeast ribo-seq dataset collection and read mapping

A list of *S. cerevisiae* ribosome profiling (ribo-seq) studies was identified by conducting a broad literature

search. For each study, all ribo-seq experiments were added to the dataset except those conducted on

mutants designed to alter wildtype translation patterns. The full list of experiments and studies included

is given in **Supplementary Tables 1 and 2**, respectively. The fastq files associated with each experiment

were downloaded from Sequence Read Archive[86] or the European Nucleotide Archive.[87] If adaptors were

present in the fastq file, they were trimmed. Reads were filtered to exclude reads in which any base had

a Phred score below 20. For each remaining read, the number of perfect matches in the *S. cerevisiae*

genome were identified, and only unique perfect matches were kept.

In initial mapping, reads were assigned to the genomic position aligning with the first base of the read. It

was necessary to remap the reads such that the position assigned to the read instead corresponded to

the first amino acid in the P-site of the translating ribosome, as in previous ribo-seq analyses[37], so that

the triplet periodic signal indicative of active translation overlaps precisely the bounds of translated

ORFs. This was done by shifting all reads by the same number of positions, with the number determined

separately for each read length and each experiment. To determine this number, a metagene profile

was constructed: the number of reads in each of the -20 to +20 positions relative to the start codon was

counted, accumulated over all annotated genes on Saccharomyces Genome Database (SGD)[43]. As there

should be many more reads on the start codon of annotated genes than the sequence immediately

upstream of these genes, the first attempt was to remap the first position with read count above a

threshold to the first amino acid on the start codons, which then requires all other reads to shift by the

same amount. The threshold selected was 5% of the total reads within 20 bases of the annotated start

codons. The attempted shift was accepted if the expected triplet periodic pattern was obtained; i.e.,

there were more remapped reads on the first base of the codons of annotated genes than on the

second or third base. Otherwise, a second shift was attempted from the next position exceeding the

read count threshold, and so on until both criteria were met.

For quality control, presence of triplet periodicity was then tested for each read length in each

experiment. The number of reads mapping (after remapping) to the first, second, and third position of

each codon was counted among annotated genes, requiring at least twice as many reads in the first

position than each of the second and third. If a read length failed this test for a given experiment it was

excluded from further analysis, and if all read lengths for an experiment failed the experiment itself was

excluded. All read lengths from 25 to 35 nucleotides were tested.

Translation calling

The iRibo program can be applied to any set of ribo-seq experiments to identify a set of ORFs with

evidence of translation among those experiments. To construct a reference translatome, translation was

inferred using ribo-seq data from the full set of experiments we collected that passed quality control

(**Supplementary Table 3**). Separately, iRibo was also run on specific subsets of the full collection,

including: experiments with or without the drug cycloheximide, experiments only on cells grown in YPD;

only on cells grown on SD; and only on cells grown in YPD without cycloheximide (**Supplementary Table**

3). iRibo was also run separately for each individual study, generating lists of translated ORFs within

each study.

Translation was assessed as follows: for each codon in each candidate ORF, the position within the

codon with the most reads was noted, if any. The number of times each codon position had the highest

read count across the ORF was then counted. The binomial test was then used to calculate a p-value for

the null hypothesis that all positions were equally likely, against the alternative that the first position

was favored. This p-value is an indicator of the strength of evidence for triplet periodicity favoring the

first codon position.

To estimate the false discovery rate (FDR), a set of ORFs corresponding to the null hypothesis was

constructed. For each ORF, the ribo-seq reads were scrambled randomly position by position (not read

by read); e.g., if 10 reads mapped to the first base on the actual ORF, a random position in the

scrambled ORF was assigned 10 reads, and so on. In this way the read distribution across positions was

maintained but the spatial structure was eliminated. The same binomial test as used for the actual reads

was then used on all scrambled-read ORFs. For every p-value threshold, the FDR can then be calculated

as the number of scrambled ORFs with p-value below the threshold divided by the number of actual

ORFs with p-values below the threshold. For each list of translated ORFs, the p-value threshold was set

`

896 to give a 5% FDR among noncanonical ORFs; all ORFs with p-values below this threshold were then

897 included in the translated set, whether canonical or noncanonical.

898 Estimating translation rates across different genomic contexts

All nORFs were partitioned into genomic contexts, with nonoverlapping nORFs classified by the relation

between the nORF and any cORF located on the same transcript and antisense nORFs classified by

partial or complete overlap of the opposite strand gene. The transcripts reported in Pelechano et al.

2014[47] based on TIF-seq data were used for this analysis. An nORF was considered antisense if it

overlapped an ORF annotated as "verified", "uncharacterized", "transposable element" or "blocked" on

SGD on the opposite strand and nonoverlapping otherwise (ORFs overlapping annotated genes on the

same strand were excluded from analysis, as described above). A nonoverlapping nORF was considered

to share a transcript with a cORF or annotated non-coding RNA if any transcript fully contained both the

nORF and the cORF or annotated RNA sequence; the ORF was then further classified as being either a

uORF or dORF based on whether it was upstream or downstream of the cORF or RNA. If an nORF shared

a transcript with both its upstream and downstream neighboring cORFs, it was classified according to

the cORF that was closer.

911 Identifying homologous sequences of the *S. cerevisiae* ORF in other Saccharomyces genus species

Genomes were obtained from seven relatives of *S. cerevisiae* within the *Saccharomyces* genus: *S.*

paradoxus from Liti et al. 2009[80], *S. arboricolus* from Liti et al. 2013[81], *S. jurei* from Naseeb et al. 2018[82],

and *S. mikatae*, *S. bayanus var. uvarum*, *S. bayanus var. bayanus,* and *S. kudriavzevii* from Scannell et al.

2011.[51] Alignments were constructed between each *S. cerevisiae* ORF and its homologs in each

Saccharomyces relative using synteny information. To identify anchor genes for syntenic blocks, BLASTP

was run for each annotated ORF in *S. cerevisiae* against each ORF in the comparison species. Identified

homolog pairs with e-value < 10^{-7} were selected as potential anchors. For each ORF in the *S. cerevisiae*

genome, the upstream anchor G_0 and downstream anchor G_1 were selected that minimized the sum of

the distance between the anchors in *S. cerevisiae* and the distance between the anchors in the

comparison species; this sum was required to be less than 60 kb. The sequence between and including

G_0 and G_1 were then extracted from both the *S. cerevisiae* genome and the comparison species and a

pairwise alignment of the syntenic region was generated using MUSCLE 3.8.31.[84]

To confirm that the ORF was matched to genuinely homologous DNA, the alignment of the *S. cerevisiae*

ORF along with its 50 bp flanking regions was extracted from the full syntenic alignment. The extracted

`

region was then realigned using the Smith-Waterman algorithm[88] with a match bonus of 5, a mismatch

penalty of 4, and a gap penalty of 4. To test homology, 1000 alignments were constructed using the

same score system in which the sequence of the comparison species was shuffled at random, reflecting

a null hypothesis that the region was not homologous. The proportion of times the alignment of the real

sequence scored better than the shuffled ones is a p-value indicating the strength of the null hypothesis

against the alternative that the region is homologous. Homology was accepted as confirmed if the p-

value was less than 1%, and alignments were excluded from analysis if homology was not confirmed.

If a syntenic alignment could not be constructed for a particular *S. cerevisiae* ORF and comparison

species (because homology failed or there were no appropriate anchors), BLAST was attempted as an

alternative method of finding the homologous DNA sequence. For these ORF sequences, BLASTn was

run against the genome of the comparison species. For each reciprocal best matching pair with e-value <

10^{-4}, the matched sequences in both species were extracted, together with a 1000 bp flanking region in

both ends, and aligned using MUSCLE.[84] DNA homology was then tested using Smith-Waterman

alignment as described above.

Division of ORFs into high information and low information sets

Evolutionary analysis of ORFs was done separately for those ORFs for which there existed substantial

information to test selection ("high information ORFs") and those for which less information was

available ("low information ORFs"). To be placed in the high information set, the ORF had to meet a

homology criterion and a diversity criterion. The homology criterion required that DNA homology was

confirmed in either a synteny or BLAST-based pairwise alignment with at least four other species in the

Saccharomyces genus. For the diversity criterion, the number of single nucleotide differences (excluding

gaps) was counted between the *S. cerevisiae* ORF and all its aligned sequence with confirmed homology

among *Saccharomyces* genomes. The diversity criterion was satisfied if the median count of differences

exceeded 20.

Reading frame conservation

Reading frame conservation is a measure of conservation of codon structure developed by Kellis et al.

2003[20] and used here with some modifications. Calculation of reading frame conservation was done on a

pairwise alignment of a genomic region containing the *S. cerevisiae* ORF (either a syntenic block

between conserved genes or the 1000 bp flanking region around a BLAST hit). All single-exon ORFs (ATG

to stop codon) in the comparison species were identified across this region. For each ORF in the

`

comparison species, the reading frame conservation was calculated by summing up all points in the

alignment where the pair of aligned bases are in the same position within the codon (i.e., both are in

either the first, second, or third position) and dividing by the length of the *S. cerevisiae* ORF in

nucleotides (including start and stop codons). Positions that align to gaps or are outside the range of the

S. cerevisiae ORF are always considered to be not in the same codon position and do not add to the

numerator. The ORF in the comparison species with the highest reading frame conservation is

considered the best match, and the reading frame conservation of the *S. cerevisiae* ORF in relation to

each other *Saccharomyces* species is defined as its reading frame conservation with its best match. In

addition to the pairwise reading frame conservation of each *S. cerevisiae* ORF in relation to its homologs

in all other species, an index of reading frame conservation (RFC) was defined equal to the average

reading frame conservation of the *S. cerevisiae* ORF against all species in the *Saccharomyces* genus for

which homologous DNA could be identified.

Detecting distant homology among *S. cerevisiae* ORFs

The genomes of 332 budding yeasts were taken from Shen et al. 2018.[41] We applied TBLASTN and

BLASTP for each *S. cerevisiae* translated ORF against each genome in this dataset (excluding the

Saccharomyces genus). Default settings were used except for setting an e-value threshold of 0.1; results

were then filtered by a stricter e-value threshold as described in each analysis. The BLASTP analysis was

run against the list of protein coding genes used in Shen et al. 2018[41] while the TBLASTN analysis was

run against each entire genome. In the TBLASTN analysis, scrambled sequences of each *S. cerevisiae* ORF

were also included as queries to serve as a negative control.

Tests of selection using the dN/dS and pN/pS ratios

Variant call file data for 1011 *S. cerevisiae* isolates was taken from Peter et al. 2018.[40] For each ORF,

nucleotide diversity was estimated from the full set of isolates. Nucleotide diversity was estimated as

the mean number of differences per site in the ORF between any pair of isolates. To calculate dN/dS, the

consensus sequence among all isolates was determined. At each position in the consensus, the three

possible nucleotide variations were recorded as possible polymorphisms and distinguished by

polymorphism type (12 possible combinations of consensus and variant nucleotide) and whether they

would result in a synonymous or nonsynonymous difference from the consensus. If at least one isolate

had the polymorphism, the polymorphism was also recorded as observed. All possible and observed

polymorphisms were counted among all considered ORFs.

`

986 The pN/pS ratio was calculated in a similar manner to Ruiz-Orera et al. 2018[28] and could be applied to

987 either a single ORF or a group of ORFs. For each ORF under consideration, the consensus sequence

988 among all isolates was determined. At each position in the consensus, the three possible nucleotide

989 variations were recorded as possible polymorphisms and distinguished by polymorphism type (12

990 possible combinations of consensus and variant nucleotide) and whether they would result in a

991 synonymous or nonsynonymous difference from the consensus. If at least one isolate had the

992 polymorphism, the polymorphism was also recorded as observed. All possible and observed

993 polymorphisms were counted among all considered ORFs.

994 Consider a variant $X \rightarrow Y$ where X is the consensus at a site and Y is a possible variant. The probability of

995 observing variant Y at a position with consensus X, $p_{x \rightarrow y}$ was estimated as the observed count of $X \rightarrow Y$

996 variant sites divided by the possible count of $X \rightarrow Y$ variant sites. Under neutrality, the expected count of

997 either synonymous or nonsynonymous $X \rightarrow Y$ variant sites is then the product of $p_{x \rightarrow y}$ and the number of

998 possible synonymous or nonsynonymous $X \rightarrow Y$ variant sites. In this manner the expected and observed

999 counts of synonymous and nonsynonymous variants were calculated. The pN/pS ratio is then estimated

1000 as:

1001 $$\omega = \frac{nonsyn_{obs}/nonsyn_{exp}}{syn_{obs}/syn_{exp}}$$

1002 Under neutrality, then, the expected count of $X \rightarrow Y$ nonsynonymous variant sites is the number of

1003 possible such variant sites times the expected probability of this variant. In this manner the expected

1004 and observed counts of all synonymous variant types were calculated. To test for deviation from

1005 neutrality, we used a chi-squared test with one degree of freedom to compare observed vs. expected

1006 counts of synonymous and nonsynonymous variants. Standard errors for the pN/pS ratio in group

1007 analyses were estimated by bootstrapping: the ORFs in the group were resampled with replacement

1008 1000 times and the pN/pS ratio was calculated each time. The standard error was then estimated as the

1009 sample standard deviation among the 1000 pN/pS ratios.

1010 The dN/dS ratio was calculated based on differences in the pairwise ORF alignments *S. cerevisiae* and its

1011 closest relative *S. paradoxus*. Each S. cerevisiae ORF was associated with an *S. paradoxus* ORF for which

1012 the pair had the highest reading frame conservation (or none if homology with *S. paradoxus* was not

1013 confirmed or the highest reading frame conservation was 0). Counts of differences were made only for

1014 codons that shared the same frame between these ORFs and with at most one nucleotide difference

1015 between the codons. For every eligible position in the *S. cerevisiae* ORF, each possible *S. paradoxus*

difference was counted and distinguished by whether the difference was synonymous or

nonsynonymous and by type (four *S. cerevisiae* nucleotides, each with three possible *S. paradoxus*

differences). These observed and possible differences were then used to estimate the dN/dS ratio in the

same way as described above for the pN/pS ratio.

Among nORFs with high RFC, the strong conservation in *Saccharomyces* permitted calculation of dN/dS

over the entire *Saccharomyces* tree, and so this was done as an additional test of selection (as reported

in Table 1). For this analysis, ancestral reconstruction of the *Saccharomyces* phylogeny was conducted

using PRANK[89] with parameters -showanc -showevents -once -prunetree -keep. Ancestral reconstruction

included all species in which DNA homology was confirmed. Codons were only used for counting

substitutions if they shared frame conservation among all species. Observed and possible substitutions

were counted across each branch and distinguished by substitution type and whether the substitutions

were synonymous or nonsynonymous. Then, dN/dS was estimated in the same way as described for

pN/pS above.

Classification of ORFs into transient and conserved sets

All high-information nonoverlapping translated ORFs with RFC > 0.8 were classified as conserved (**Figure**

4A). An nORF was also classified as conserved if it overlapped no annotated feature on SGD, had

TBLASTN matches with e-value < 10^{-4} with at least two species outside the *Saccharomyces* genus and

showed at least one additional signature of purifying selection (RFC > 0.8 or a p-value < 0.05 in a test of

neutrality using dN/dS or pN/pS) (**Supplementary Figure 5A**).

Nonoverlapping ORFs were excluded from classification in the transient set if they showed homology to

an ORF classified as conserved in *S. cerevisiae* (e-value < 10^{-4} using BLASTP) or to any sequence among

budding yeasts outside *Saccharomyces*[41] (e-value < 10^{-4} using TBLASTN). Among remaining translated

ORFs, all high-information ORFs with RFC < 0.6 were classified as transient. Low information ORFs were

divided into groups and classified as transient if no group they belonged to showed evidence of

selection in dN/dS analysis, pN/pS analysis, or weak homology matching analysis. Two low-information

groups were cORFs and antisense nORFs. Low information nonoverlapping nORFs were each assigned to

three groups corresponding to deciles of translation rate, coding score and ORF length. Analyses of

dN/dS and pN/pS are described above. For weak homology detection, the number of ORFs with at least

two weak TBLASTN matches (e-value < 0.05) to budding yeast genomes collected by Shen et al. 2018[41]

(excluding *Saccharomyces* species) was counted for both actual and scrambled ORF sequences. Selection

`

was inferred if actual matches significantly (p < 0.05) exceeded scrambled matches using Fisher's exact

test. Only ORFs that did not overlap any annotated feature on SGD were included in weak homology

detection analysis.

Coding score calculation

The coding score, described by Ruiz-Orera et al. 2014[90], is a measure of how close the hexamer (i.e., the

nucleotide sequence of a pair of adjacent codons) frequency of an ORF is to the hexamer of coding vs.

noncoding sequences. Higher scores indicate a more gene-like hexamer distribution. Coding hexamer

frequencies were calculated among all ORFs annotated as "verified" or "uncharacterized" by

Saccharomyces Genome Database.[43] Noncoding hexamer frequencies were calculated for all intergenic

sequences (sequences in between verified or uncharacterized ORFs) in the *S. cerevisiae* genome. As

intergenic sequence has no codon structure, hexamer frequencies for intergenic sequence were counted

as if read in each possible coding frame. The score was then calculated as described in Ruiz-Orera et al.

2014.[90]

Identification to transient ORFs with detectable translation products in published microscopy studies

Published results were examined from fluorescent tagging experiments where the expression of ORFs

was driven by native promoters and terminators. A list of ORFs detected in 15 GFP-tagged screens on

wildtype strains in either normal conditions or with chemical treatment (hydroxyurea or rapamycin)

were retrieved from the CYCLoPs database.[58,59] Lists of ORFs detected in the C-SWAT tagging library

were taken from Meurer et al. 2018[60] and from YeastRGB[61]. ORFs with fluorescent intensity below the

reported detection threshold in each screen were filtered out. Transient ORFs that showed detectable

translation products in at least one screen were considered as detected.

Literature analysis of transient translatome cORFs

For each transient translatome cORF, we examined all publications listed on SGD as "primary" or

"additional" literature for the ORF. If the ORF had a phenotype in any listed publication, we noted the

evidence for the phenotype (**Supplementary Table 5**).

Genetic interaction analysis

Single mutant fitness and genetic interaction data were downloaded from TheCellMap.org.[91] In this

dataset, mutants of nonessential genes are full deletions and mutants of essential genes are

temperature-sensitive alleles. Transient ORFs were all nonessential. Different temperature-sensitive

alleles for the same essential gene were treated separately. We removed all genes or transient ORFs

with a genomic overlap to another genetic element from our analyses as it is not possible to assign the

observed phenotypes to either of the overlapping pairs.

We counted the number of transient ORF and nonessential genes that showed at least one genetic

interaction with Ɛ<-.2 and p-value < 0.05 (a negative genetic interaction) or Ɛ<-.35 with a p-value<0.05 (a

synthetic lethal interaction). We then divided this number by the total number of transient ORFs or

nonessential genes in the Costanzo et al. 2016[69] genetic interaction dataset to calculate the percentage

showing at least one genetic interaction. We used Fisher's exact test to assess the significance of

differences between percentages of nonessential genes and transient ORFs.

Gene ontology analysis of the interactors of each ORF was conducted with Ontologizer[83], using

Benjamini-Hochberg multiple testing correction and the term-for-term calculation method. The gene

association file was downloaded from SGD. Gene ontology evidence codes relating to genetic

interactions (IGI and HGI) were not used.

Creation of yeast strains

Deletion mutant strains for 49 transient nORFs and 3 transient cORFs were created by using homologous

recombination to replace the ORFs with a KanMX cassette. Transformations were done using the

LiAc/PEG protocol[85] in the background BY4741 strain, and selected in media containing G-418. After an

initial screen of these strains, a subset of the deletion strains that showed strong deleterious effects

were transformed a second time, also using the LiAc/PEG protocol[85], to replace the KanMx cassette with

either an intact copy of the original ORF, or a mutant copy of the ORF with the start codon ATG and (in

some cases) additional in-frame ATG codons mutated to AAG to prevent translation. This was

accomplished by using homologous recombination to replace the KanMx cassette with a construct

containing the intact or mutant ORF followed by a hygromycin resistance cassette. These constructs

were synthesized by IDT (Integrated DNA Technologies). The resulting transformants were selected in

agar plates containing hygromycin. All positive clones were sequenced to confirm presence of either the

restored wildtype ORF or the ORF with a mutated start codon.

Strains containing an mNeonGreen tag for microscopy purposes were also made by homologous

recombination using the LiAc/PEG protocol[85] in the BY4741 background. The mNeonGreen and

hygromycin cassette sequences were amplified from a plasmid using primers containing homology to

the 3' of each ORF. The primers were designed to remove the STOP codon of each ORF and place the

`

mNeonGreen in frame with the ORF, to be expressed under its native promoter. Positive clones were

selected on agar plates containing hygromycin.

All strains were kept in glycerol stocks at −80 °C in 96 and 384-well format until used for screening.

Strain genotypes are listed in **Supplementary Table 9**.

Screening strategy for fitness estimation

Both rounds of deletion screening were conducted at 1536 colony density, with 1 in 4 colonies on the

plate being reference strains used to correct for spatial biases as described in Parikh et al. 2021.[92] In the

initial deletion screen, each mutant strain was tested using 12 replicates; 72 replicates were tested per

strain in the start codon mutant screen. Conditions tested were YPDA and YPDA+DMSO as unstressed

conditions and five stress conditions: YPDA supplemented with 1M NaCl, 100mM Hydroxyurea, 0.6μM

Tunicamycin, 25μg/ml Fluconazole, or 30mM Hydrogen peroxide (H_2O_2). Agar plates were incubated and

imaged periodically until the colonies reached saturation. The plate handler Singer ROTOR (Singer

Instrument Co. Ltd) was used to prepare all plates starting from glycerol stocks. Serial imaging of the

plates was conducted using the spImager Automated Imaging System (S & P Robotics Inc., Ontario,

Canada). The images were analyzed in bulk using a custom script made using functions from the

MATLAB Colony Analyzer Toolkit[92] to provide colony size estimations

(https://github.com/sauriiiin/lid_personal/blob/master/justanalyze.m). The output files containing

colony size information along with the images is available at https://bit.ly/3xtzHJO. The LI Detector

analytical pipeline[92] was used to correct for spatial biases in colony size and obtain colony fitness

estimates. Strain fitness was estimated as the median of bias-corrected colony size among replicates of

the strain at 40 hours in the initial screen and 90 hours in the start codon mutant screen. In the LI

Detector pipeline[92], sets of reference colonies are treated as if they were replicates of a mutant strain,

with their median fitness calculated in order to construct an empirical null distribution of median fitness

values to compare with estimated strain fitness. Strains were called as beneficial or deleterious using a

5% false discovery rate threshold based on this empirical null distribution. For any selected fitness

threshold used to infer deleterious strains, the false discovery rate can be calculated as the proportion

of null distribution fitness values below that threshold divided by the proportion of mutant strain fitness

values below the threshold. Thus, fitness thresholds were selected such that a 5% FDR was obtained and

strains with fitness below that threshold were inferred to be deleterious. In the same manner, a list of

beneficial strains at 5% FDR was also selected.

`

Liquid growth assay

For liquid growth assays, cells were first grown in liquid YPDA media overnight at 30°C in a 96-density

microplate. These were then used to inoculate a new 96-density microplate with 150μl YPDA+ stress

conditions (1M NaCl, 100mM Hydroxyurea)) using the Singer ROTOR (Singer Instrument Co. Ltd). This

microplate was incubated at 30°C with constant double orbital shaking for a period of 72h on microplate

reader Biotek Synergy H1 (Aligent Technology Inc.). Optical density readings at 600nm (OD$_{600}$) were

taken every 15 minutes.

Microscopy

The strains containing the ORFs tagged with mNeonGreen were imaged on a Nikon TiE2 inverted A1R

confocal microscope. A first screening was done at high density in 96-well plates with a 40x water

objective, to assess the success of the transformations. Plates were incubated with CellTracker Blue

CMAC Dye (Invitrogen) and MitoTracker Red CMXRos Dye (Invitrogen) at least 10 min prior to imaging.

Plates were then imaged in 4 channels (405, 488, 561, and DIC), and 3 fields of view were taken for each

strain that contained many cells. Strains that demonstrated visibly higher signal in the green channel

(488nm) compared to a non-transformed background strain were selected to examine in single dishes

under a 100X oil objective to more accurately evaluate sub-cellular localization. All strains were imaged

in triplicate at high density and triplicate in dishes (once without CMAC and MitoTracker and two times

with the dyes).

Quantification and statistical analysis

Statistical analyses were performed in R version 4.1.2. Details for each statistical test and analysis can be

found in the results section and figure legends.

1156

1157

`

1158

References

1. Ingolia, N.T., Brar, G.A., Stern-Ginossar, N., Harris, M.S., Talhouarne, G.J.S., Jackson, S.E., Wills, M.R.,
and Weissman, J.S. (2014). Ribosome Profiling Reveals Pervasive Translation Outside of Annotated
Protein-Coding Genes. Cell Rep. *8*, 1365–1379. 10.1016/j.celrep.2014.07.045.

2. Ingolia, N.T. (2014). Ribosome profiling: new views of translation, from single codons to genome
scale. Nat. Rev. Genet. *15*, 205–213. 10.1038/nrg3645.

3. Carvunis, A.-R., Rolland, T., Wapinski, I., Calderwood, M.A., Yildirim, M.A., Simonis, N., Charloteaux,
B., Hidalgo, C.A., Barbette, J., Santhanam, B., et al. (2012). Proto-genes and *de novo* gene birth.
Nature *487*, 370–374. 10.1038/nature11184.

4. Wilson, B.A., and Masel, J. (2011). Putatively Noncoding Transcripts Show Extensive Association with
Ribosomes. Genome Biol. Evol. *3*, 1245–1252. 10.1093/gbe/evr099.

5. Pruitt, K.D., and Maglott, D.R. (2001). RefSeq and LocusLink: NCBI gene-centered resources. Nucleic
Acids Res. *29*, 137–140. 10.1093/nar/29.1.137.

6. Erhard, F., Halenius, A., Zimmermann, C., L'Hernault, A., Kowalewski, D.J., Weekes, M.P., Stevanovic,
S., Zimmer, R., and Dölken, L. (2018). Improved Ribo-seq enables identification of cryptic translation
events. Nat. Methods *15*, 363–366. 10.1038/nmeth.4631.

7. Chen, J., Brunner, A.-D., Cogan, J.Z., Nuñez, J.K., Fields, A.P., Adamson, B., Itzhak, D.N., Li, J.Y., Mann,
M., Leonetti, M.D., et al. (2020). Pervasive functional translation of noncanonical human open
reading frames. Science *367*, 1140–1146. 10.1126/science.aay0262.

8. Prensner, J.R., Enache, O.M., Luria, V., Krug, K., Clauser, K.R., Dempster, J.M., Karger, A., Wang, L.,
Stumbraite, K., Wang, V.M., et al. (2021). Noncanonical open reading frames encode functional
proteins essential for cancer cell survival. Nat. Biotechnol., 1–8. 10.1038/s41587-020-00806-2.

9. van Heesch, S., Witte, F., Schneider-Lunitz, V., Schulz, J.F., Adami, E., Faber, A.B., Kirchner, M.,
Maatz, H., Blachut, S., Sandmann, C.-L., et al. (2019). The Translational Landscape of the Human
Heart. Cell *178*, 242-260.e29. 10.1016/j.cell.2019.05.010.

10. Laumont, C.M., Daouda, T., Laverdure, J.-P., Bonneil, É., Caron-Lizotte, O., Hardy, M.-P., Granados,
D.P., Durette, C., Lemieux, S., Thibault, P., et al. (2016). Global proteogenomic analysis of human
MHC class I-associated peptides derived from non-canonical reading frames. Nat. Commun. *7*,
10238. 10.1038/ncomms10238.

11. Jackson, R., Kroehling, L., Khitun, A., Bailis, W., Jarret, A., York, A.G., Khan, O.M., Brewer, J.R.,
Skadow, M.H., Duizer, C., et al. (2018). The translation of non-canonical open reading frames
controls mucosal immunity. Nature *564*, 434. 10.1038/s41586-018-0794-7.

12. Makarewich, C.A., and Olson, E.N. (2017). Mining for Micropeptides. Trends Cell Biol. *27*, 685–696.
10.1016/j.tcb.2017.04.006.

`

13. Anderson, D.M., Anderson, K.M., Chang, C.-L., Makarewich, C.A., Nelson, B.R., McAnally, J.R.,
Kasaragod, P., Shelton, J.M., Liou, J., Bassel-Duby, R., et al. (2015). A Micropeptide Encoded by a
Putative Long Noncoding RNA Regulates Muscle Performance. Cell *160*, 595–606.
10.1016/j.cell.2015.01.009.

14. Matsumoto, A., Pasut, A., Matsumoto, M., Yamashita, R., Fung, J., Monteleone, E., Saghatelian, A.,
Nakayama, K.I., Clohessy, J.G., and Pandolfi, P.P. (2017). mTORC1 and muscle regeneration are
regulated by the LINC00961-encoded SPAR polypeptide. Nature *541*, 228–232.
10.1038/nature21034.

15. Polycarpou-Schwarz, M., Groß, M., Mestdagh, P., Schott, J., Grund, S.E., Hildenbrand, C., Rom, J.,
Aulmann, S., Sinn, H.-P., Vandesompele, J., et al. (2018). The cancer-associated microprotein
CASIMO1 controls cell proliferation and interacts with squalene epoxidase modulating lipid droplet
formation. Oncogene *37*, 4750–4768. 10.1038/s41388-018-0281-5.

16. Housman, G., and Ulitsky, I. (2016). Methods for distinguishing between protein-coding and long
noncoding RNAs and the elusive biological purpose of translation of long noncoding RNAs. Biochim.
Biophys. Acta BBA - Gene Regul. Mech. *1859*, 31–40. 10.1016/j.bbagrm.2015.07.017.

17. Struhl, K. (2007). Transcriptional noise and the fidelity of initiation by RNA polymerase II. Nat. Struct.
Mol. Biol. *14*, 103–105. 10.1038/nsmb0207-103.

18. Pertea, M., Shumate, A., Pertea, G., Varabyou, A., Breitwieser, F.P., Chang, Y.-C., Madugundu, A.K.,
Pandey, A., and Salzberg, S.L. (2018). CHESS: a new human gene catalog curated from thousands of
large-scale RNA sequencing experiments reveals extensive transcriptional noise. Genome Biol. *19*,
208. 10.1186/s13059-018-1590-2.

19. Ponjavic, J., Ponting, C.P., and Lunter, G. (2007). Functionality or transcriptional noise? Evidence for
selection within long noncoding RNAs. Genome Res. *17*, 556–565. 10.1101/gr.6036807.

20. Kellis, M., Patterson, N., Endrizzi, M., Birren, B., and Lander, E.S. (2003). Sequencing and comparison
of yeast species to identify genes and regulatory elements. Nature *423*, 241. 10.1038/nature01644.

21. Ward, L.D., and Kellis, M. (2012). Evidence of Abundant Purifying Selection in Humans for Recently
Acquired Regulatory Functions. Science *337*, 1675–1678. 10.1126/science.1225057.

22. Kellis, M., Wold, B., Snyder, M.P., Bernstein, B.E., Kundaje, A., Marinov, G.K., Ward, L.D., Birney, E.,
Crawford, G.E., Dekker, J., et al. (2014). Defining functional DNA elements in the human genome.
Proc. Natl. Acad. Sci. *111*, 6131–6138. 10.1073/pnas.1318948111.

23. Oshiro, G., Wodicka, L.M., Washburn, M.P., Yates, J.R., Lockhart, D.J., and Winzeler, E.A. (2002).
Parallel Identification of New Genes in Saccharomyces cerevisiae. Genome Res. *12*, 1210–1220.
10.1101/gr.226802.

24. Blandin, G., Durrens, P., Tekaia, F., Aigle, M., Bolotin-Fukuhara, M., Bon, E., Casarégola, S.,
Montigny, J. de, Gaillardin, C., Lépingle, A., et al. (2000). Genomic Exploration of the
Hemiascomycetous Yeasts: 4. The genome of Saccharomyces cerevisiae revisited. FEBS Lett. *487*,
31–36. 10.1016/S0014-5793(00)02275-4.

`

25. Bazzini, A.A., Johnstone, T.G., Christiano, R., Mackowiak, S.D., Obermayer, B., Fleming, E.S., Vejnar,
C.E., Lee, M.T., Rajewsky, N., Walther, T.C., et al. (2014). Identification of small ORFs in vertebrates
using ribosome footprinting and evolutionary conservation. EMBO J. *33*, 981–993.
10.1002/embj.201488411.

26. Crappé, J., Van Criekinge, W., Trooskens, G., Hayakawa, E., Luyten, W., Baggerman, G., and
Menschaert, G. (2013). Combining in silico prediction and ribosome profiling in a genome-wide
search for novel putatively coding sORFs. BMC Genomics *14*, 648. 10.1186/1471-2164-14-648.

27. Durand, É., Gagnon-Arsenault, I., Hallin, J., Hatin, I., Dubé, A.K., Nielly-Thibault, L., Namy, O., and
Landry, C.R. (2019). Turnover of ribosome-associated transcripts from de novo ORFs produces gene-
like characteristics available for de novo gene emergence in wild yeast populations. Genome Res.
29, 932–943. 10.1101/gr.239822.118.

28. Ruiz-Orera, J., Verdaguer-Grau, P., Villanueva-Cañas, J.L., Messeguer, X., and Albà, M.M. (2018).
Translation of neutrally evolving peptides provides a basis for de novo gene evolution. Nat. Ecol.
Evol. *2*, 890. 10.1038/s41559-018-0506-6.

29. Olexiouk, V., Crappé, J., Verbruggen, S., Verhegen, K., Martens, L., and Menschaert, G. (2016).
sORFs.org: a repository of small ORFs identified by ribosome profiling. Nucleic Acids Res. *44*, D324–
D329. 10.1093/nar/gkv1175.

30. Lee, J., Wacholder, A., and Carvunis, A.-R. (2021). Evolutionary Characterization of the Short Protein
SPAAR. Genes *12*, 1864. 10.3390/genes12121864.

31. Mudge, J.M., Ruiz-Orera, J., Prensner, J.R., Brunet, M.A., Calvet, F., Jungreis, I., Gonzalez, J.M.,
Magrane, M., Martinez, T.F., Schulz, J.F., et al. (2022). Standardized annotation of translated open
reading frames. Nat. Biotechnol. *40*, 994–999. 10.1038/s41587-022-01369-0.

32. Van Oss, S.B., and Carvunis, A.-R. (2019). De novo gene birth. PLoS Genet. *15*.
10.1371/journal.pgen.1008160.

33. Blevins, W.R., Ruiz-Orera, J., Messeguer, X., Blasco-Moreno, B., Villanueva-Cañas, J.L., Espinar, L.,
Díez, J., Carey, L.B., and Albà, M.M. (2021). Uncovering de novo gene birth in yeast using deep
transcriptomics. Nat. Commun. *12*, 604. 10.1038/s41467-021-20911-3.

34. Yagoub, D., Tay, A.P., Chen, Z., Hamey, J.J., Cai, C., Chia, S.Z., Hart-Smith, G., and Wilkins, M.R.
(2015). Proteogenomic Discovery of a Small, Novel Protein in Yeast Reveals a Strategy for the
Detection of Unannotated Short Open Reading Frames. J. Proteome Res. *14*, 5038–5047.
10.1021/acs.jproteome.5b00734.

35. Lu, S., Zhang, J., Lian, X., Sun, L., Meng, K., Chen, Y., Sun, Z., Yin, X., Li, Y., Zhao, J., et al. (2019). A
hidden human proteome encoded by 'non-coding' genes. Nucleic Acids Res. *47*, 8111–8125.
10.1093/nar/gkz646.

36. Ouspenskaia, T., Law, T., Clauser, K.R., Klaeger, S., Sarkizova, S., Aguet, F., Li, B., Christian, E.,
Knisbacher, B.A., Le, P.M., et al. (2020). Thousands of novel unannotated proteins expand the MHC I
immunopeptidome in cancer. bioRxiv, 2020.02.12.945840. 10.1101/2020.02.12.945840.

`

37. Malone, B., Atanassov, I., Aeschimann, F., Li, X., Großhans, H., and Dieterich, C. (2017). Bayesian
prediction of RNA translation from ribosome profiling. Nucleic Acids Res. *45*, 2960–2972.
10.1093/nar/gkw1350.

38. Ji, Z. (2018). RibORF: Identifying Genome-Wide Translated Open Reading Frames Using Ribosome
Profiling. Curr. Protoc. Mol. Biol. *124*, e67. 10.1002/cpmb.67.

39. Calviello, L., and Ohler, U. (2017). Beyond Read-Counts: Ribo-seq Data Analysis to Understand the
Functions of the Transcriptome. Trends Genet. *33*, 728–744. 10.1016/j.tig.2017.08.003.

40. Peter, J., Chiara, M.D., Friedrich, A., Yue, J.-X., Pflieger, D., Bergström, A., Sigwalt, A., Barre, B., Freel,
K., Llored, A., et al. (2018). Genome evolution across 1,011 Saccharomyces cerevisiae isolates.
Nature *556*, 339. 10.1038/s41586-018-0030-5.

41. Shen, X.-X., Opulente, D.A., Kominek, J., Zhou, X., Steenwyk, J.L., Buh, K.V., Haase, M.A.B.,
Wisecaver, J.H., Wang, M., Doering, D.T., et al. (2018). Tempo and Mode of Genome Evolution in the
Budding Yeast Subphylum. Cell *175*, 1533-1545.e20. 10.1016/j.cell.2018.10.023.

42. Choudhary, S., Li, W., and D. Smith, A. (2020). Accurate detection of short and long active ORFs
using Ribo-seq data. Bioinformatics *36*, 2053–2059. 10.1093/bioinformatics/btz878.

43. Cherry, J.M., Adler, C., Ball, C., Chervitz, S.A., Dwight, S.S., Hester, E.T., Jia, Y., Juvik, G., Roe, T.,
Schroeder, M., et al. (1998). SGD: Saccharomyces Genome Database. Nucleic Acids Res. *26*, 73–79.
10.1093/nar/26.1.73.

44. Gerashchenko, M.V., and Gladyshev, V.N. (2014). Translation inhibitors cause abnormalities in
ribosome profiling experiments. Nucleic Acids Res. *42*, e134–e134. 10.1093/nar/gku671.

45. Santos, D.A., Shi, L., Tu, B.P., and Weissman, J.S. (2019). Cycloheximide can distort measurements of
mRNA levels and translation efficiency. Nucleic Acids Res. *47*, 4974–4985. 10.1093/nar/gkz205.

46. Duncan, C.D.S., and Mata, J. (2017). Effects of cycloheximide on the interpretation of ribosome
profiling experiments in Schizosaccharomyces pombe. Sci. Rep. *7*, 10331. 10.1038/s41598-017-
10650-1.

47. Pelechano, V., Wei, W., Jakob, P., and Steinmetz, L.M. (2014). Genome-wide identification of
transcript start and end sites by transcript isoform sequencing. Nat. Protoc. *9*, 1740–1759.
10.1038/nprot.2014.121.

48. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Methodol. *57*, 289–300.
10.1111/j.2517-6161.1995.tb02031.x.

49. Ji, Z., Song, R., Regev, A., and Struhl, K. (2015). Many lncRNAs, 5'UTRs, and pseudogenes are
translated and some are likely to express functional proteins. eLife *4*, e08890. 10.7554/eLife.08890.

50. Moro, S.G., Hermans, C., Ruiz-Orera, J., and Albà, M.M. (2021). Impact of uORFs in mediating
regulation of translation in stress conditions. BMC Mol. Cell Biol. *22*, 29. 10.1186/s12860-021-
00363-9.

`

51. Scannell, D.R., Zill, O.A., Rokas, A., Payen, C., Dunham, M.J., Eisen, M.B., Rine, J., Johnston, M., and
Hittinger, C.T. (2011). The Awesome Power of Yeast Evolutionary Genetics: New Genome Sequences
and Strain Resources for the Saccharomyces sensu stricto Genus. G3 Genes Genomes Genet. *1*, 11–
25. 10.1534/g3.111.000273.

52. Ruiz-Orera, J., Hernandez-Rodriguez, J., Chiva, C., Sabidó, E., Kondova, I., Bontrop, R., Marqués-
Bonet, T., and Albà, M.M. (2015). Origins of De Novo Genes in Human and Chimpanzee. PLOS Genet.
11, e1005721. 10.1371/journal.pgen.1005721.

53. Firth, A.E. (2014). Mapping overlapping functional elements embedded within the protein-coding
regions of RNA viruses. Nucleic Acids Res. *42*, 12425–12439. 10.1093/nar/gku981.

54. Sealfon, R.S., Lin, M.F., Jungreis, I., Wolf, M.Y., Kellis, M., and Sabeti, P.C. (2015). FRESCo: finding
regions of excess synonymous constraint in diverse viruses. Genome Biol. *16*, 38. 10.1186/s13059-
015-0603-7.

55. Hardison, R.C. (2003). Comparative Genomics. PLOS Biol. *1*, e58. 10.1371/journal.pbio.0000058.

56. Dujon, B. (1996). The yeast genome project: what did we learn? Trends Genet. *12*, 263–270.
10.1016/0168-9525(96)10027-5.

57. Dujon, B., Alexandraki, D., André, B., Ansorge, W., Baladron, V., Ballesta, J.P.G., Banrevi, A., Bolle,
P.A., Bolotin-Fukuhara, M., Bossier, P., et al. (1994). Complete DNA sequence of yeast chromosome
XI. Nature *369*, 371–378. 10.1038/369371a0.

58. Chong, Y.T., Koh, J.L.Y., Friesen, H., Duffy, S.K., Cox, M.J., Moses, A., Moffat, J., Boone, C., and
Andrews, B.J. (2015). Yeast Proteome Dynamics from Single Cell Imaging and Automated Analysis.
Cell *161*, 1413–1424. 10.1016/j.cell.2015.04.051.

59. Koh, J.L.Y., Chong, Y.T., Friesen, H., Moses, A., Boone, C., Andrews, B.J., and Moffat, J. (2015).
CYCLoPs: A Comprehensive Database Constructed from Automated Analysis of Protein Abundance
and Subcellular Localization Patterns in Saccharomyces cerevisiae. G3 GenesGenomesGenetics *5*,
1223–1232. 10.1534/g3.115.017830.

60. Meurer, M., Duan, Y., Sass, E., Kats, I., Herbst, K., Buchmuller, B.C., Dederer, V., Huber, F., Kirrmaier,
D., Štefl, M., et al. (2018). Genome-wide C-SWAT library for high-throughput yeast genome tagging.
Nat. Methods *15*, 598–600. 10.1038/s41592-018-0045-8.

61. Dubreuil, B., Sass, E., Nadav, Y., Heidenreich, M., Georgeson, J.M., Weill, U., Duan, Y., Meurer, M.,
Schuldiner, M., Knop, M., et al. (2019). YeastRGB: comparing the abundance and localization of
yeast proteins across cells and libraries. Nucleic Acids Res. *47*, D1245–D1249. 10.1093/nar/gky941.

62. Li, D., Dong, Y., Jiang, Y., Jiang, H., Cai, J., and Wang, W. (2010). A *de novo* originated gene depresses
budding yeast mating pathway and is repressed by the protein encoded by its antisense strand. Cell
Res. *20*, 408–420. 10.1038/cr.2010.31.

63. Vakirlis, N., Hebert, A.S., Opulente, D.A., Achaz, G., Hittinger, C.T., Fischer, G., Coon, J.J., and
Lafontaine, I. (2018). A Molecular Portrait of De Novo Genes in Yeasts. Mol. Biol. Evol. *35*, 631–645.
10.1093/molbev/msx315.

`

64. Li, D., Yan, Z., Lu, L., Jiang, H., and Wang, W. (2014). Pleiotropy of the de novo-originated gene
MDF1. Sci. Rep. *4*, 7280. 10.1038/srep07280.

65. Vakirlis, N., Acar, O., Hsu, B., Castilho Coelho, N., Van Oss, S.B., Wacholder, A., Medetgul-Ernar, K.,
Bowman, R.W., Hines, C.P., Iannotta, J., et al. (2020). De novo emergence of adaptive membrane
proteins from thymine-rich genomic sequences. Nat. Commun. *11*, 1–18. 10.1038/s41467-020-
14500-z.

66. Omidi, K., Jessulat, M., Hooshyar, M., Burnside, D., Schoenrock, A., Kazmirchuk, T., Hajikarimlou, M.,
Daniel, M., Moteshareie, H., Bhojoo, U., et al. (2018). Uncharacterized ORF HUR1 influences the
efficiency of non-homologous end-joining repair in Saccharomyces cerevisiae. Gene *639*, 128–136.
10.1016/j.gene.2017.10.003.

67. Hajikarimlou, M., Moteshareie, H., Omidi, K., Hooshyar, M., Shaikho, S., Kazmirchuk, T., Burnside, D.,
Takallou, S., Zare, N., Jagadeesan, S.K., et al. (2020). Sensitivity of yeast to lithium chloride connects
the activity of YTA6 and YPR096C to translation of structured mRNAs. PLOS ONE *15*, e0235033.
10.1371/journal.pone.0235033.

68. Alesso, C.A., Discola, K.F., and Monteiro, G. (2015). The gene ICS3 from the yeast Saccharomyces
cerevisiae is involved in copper homeostasis dependent on extracellular pH. Fungal Genet. Biol. *82*,
43–50. 10.1016/j.fgb.2015.06.007.

69. Costanzo, M., VanderSluis, B., Koch, E.N., Baryshnikova, A., Pons, C., Tan, G., Wang, W., Usaj, M.,
Hanchard, J., Lee, S.D., et al. (2016). A global genetic interaction network maps a wiring diagram of
cellular function. Science *353*, aaf1420. 10.1126/science.aaf1420.

70. Kearse, M.G., and Wilusz, J.E. (2017). Non-AUG translation: a new start for protein synthesis in
eukaryotes. Genes Dev. *31*, 1717–1731. 10.1101/gad.305250.117.

71. Loughran, G., Zhdanov, A.V., Mikhaylova, M.S., Rozov, F.N., Datskevich, P.N., Kovalchuk, S.I.,
Serebryakova, M.V., Kiniry, S.J., Michel, A.M., O'Connor, P.B.F., et al. (2020). Unusually efficient CUG
initiation of an overlapping reading frame in POLG mRNA yields novel protein POLGARF. Proc. Natl.
Acad. Sci. *117*, 24936–24946. 10.1073/pnas.2001433117.

72. McVeigh, A., Fasano, A., Scott, D.A., Jelacic, S., Moseley, S.L., Robertson, D.C., and Savarino, S.J.
(2000). IS1414, an Escherichia coliInsertion Sequence with a Heat-Stable Enterotoxin Gene
Embedded in a Transposase-Like Gene. Infect. Immun. *68*, 5710–5715. 10.1128/IAI.68.10.5710-
5715.2000.

73. Wright, B.W., Yi, Z., Weissman, J.S., and Chen, J. (2022). The dark proteome: translation from
noncanonical open reading frames. Trends Cell Biol. *32*, 243–258. 10.1016/j.tcb.2021.10.010.

74. Xie, C., Bekpen, C., Künzel, S., Keshavarz, M., Krebs-Wheaton, R., Skrabar, N., Ullrich, K.K., and Tautz,
D. (2019). A de novo evolved gene in the house mouse regulates female pregnancy cycles. eLife *8*.
10.7554/eLife.44392.

75. Borneman, A.R., Gianoulis, T.A., Zhang, Z.D., Yu, H., Rozowsky, J., Seringhaus, M.R., Wang, L.Y.,
Gerstein, M., and Snyder, M. (2007). Divergence of Transcription Factor Binding Sites Across Related
Yeast Species. Science *317*, 815–819. 10.1126/science.1140748.

`

76. Wethmar, K. (2014). The regulatory potential of upstream open reading frames in eukaryotic gene
expression. WIREs RNA *5*, 765–768. 10.1002/wrna.1245.

77. Wu, Q., Wright, M., Gogol, M.M., Bradford, W.D., Zhang, N., and Bazzini, A.A. (2020). Translation of
small downstream ORFs enhances translation of canonical main open reading frames. EMBO J. *39*,
e104763. 10.15252/embj.2020104763.

78. Andjus, S., Szachnowski, U., Vogt, N., Hatin, I., Papadopoulos, C., Lopes, A., Namy, O., Wery, M., and
Morillon, A. (2022). Translation is a key determinant controlling the fate of cytoplasmic long non-
coding RNAs. 2022.05.25.493276. 10.1101/2022.05.25.493276.

79. Wery, M., Descrimes, M., Vogt, N., Dallongeville, A.-S., Gautheret, D., and Morillon, A. (2016).
Nonsense-Mediated Decay Restricts LncRNA Levels in Yeast Unless Blocked by Double-Stranded
RNA Structure. Mol. Cell *61*, 379–392. 10.1016/j.molcel.2015.12.020.

80. Liti, G., Carter, D.M., Moses, A.M., Warringer, J., Parts, L., James, S.A., Davey, R.P., Roberts, I.N.,
Burt, A., Koufopanou, V., et al. (2009). Population genomics of domestic and wild yeasts. Nature
458, 337–341. 10.1038/nature07743.

81. Liti, G., Ba, A.N.N., Blythe, M., Müller, C.A., Bergström, A., Cubillos, F.A., Dafhnis-Calas, F.,
Khoshraftar, S., Malla, S., Mehta, N., et al. (2013). High quality de novo sequencing and assembly of
the Saccharomyces arboricolus genome. BMC Genomics *14*, 69. 10.1186/1471-2164-14-69.

82. Naseeb, S., Alsammar, H., Burgis, T., Donaldson, I., Knyazev, N., Knight, C., and Delneri, D. (2018).
Whole Genome Sequencing, de Novo Assembly and Phenotypic Profiling for the New Budding Yeast
Species Saccharomyces jurei. G3 Genes Genomes Genet. *8*, 2967–2977. 10.1534/g3.118.200476.

83. Bauer, S., Grossmann, S., Vingron, M., and Robinson, P.N. (2008). Ontologizer 2.0--a multifunctional
tool for GO term enrichment analysis and data exploration. Bioinforma. Oxf. Engl. *24*, 1650–1651.
10.1093/bioinformatics/btn250.

84. Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. *32*, 1792–1797. 10.1093/nar/gkh340.

85. Dunham, M.J., Dunham, M.J., Gartenberg, M.R., and Brown, G.W. (2015). Methods in yeast genetics
and genomics: a cold spring harbor laboratory course manual (Cold Spring Harbor Laboratory Press).

86. Leinonen, R., Sugawara, H., Shumway, M., and Collaboration, on behalf of the I.N.S.D. (2011). The
Sequence Read Archive. Nucleic Acids Res. *39*, D19–D21. 10.1093/nar/gkq1019.

87. Leinonen, R., Akhtar, R., Birney, E., Bower, L., Cerdeno-Tárraga, A., Cheng, Y., Cleland, I., Faruque,
N., Goodgame, N., Gibson, R., et al. (2011). The European Nucleotide Archive. Nucleic Acids Res. *39*,
D28–D31. 10.1093/nar/gkq967.

88. Smith, T.F., and Waterman, M.S. (1981). Identification of common molecular subsequences. J. Mol.
Biol. *147*, 195–197. 10.1016/0022-2836(81)90087-5.

`

89. Löytynoja, A. (2014). Phylogeny-aware alignment with PRANK. In Multiple Sequence Alignment
Methods Methods in Molecular Biology., D. J. Russell, ed. (Humana Press), pp. 155–170.
10.1007/978-1-62703-646-7_10.

90. Ruiz-Orera, J., Messeguer, X., Subirana, J.A., and Alba, M.M. (2014). Long non-coding RNAs as a
source of new peptides. eLife *3*. 10.7554/eLife.03523.

91. Usaj, M., Tan, Y., Wang, W., VanderSluis, B., Zou, A., Myers, C.L., Costanzo, M., Andrews, B., and
Boone, C. (2017). TheCellMap.org: A Web-Accessible Database for Visualizing and Mining the Global
Yeast Genetic Interaction Network. G3 GenesGenomesGenetics *7*, 1539–1549.
10.1534/g3.117.040220.

92. Parikh, S.B., Castilho Coelho, N., and Carvunis, A.-R. (2021). LI Detector: a framework for sensitive
colony-based screens regardless of the distribution of fitness effects. G3 GenesGenomesGenetics
11, jkaa068. 10.1093/g3journal/jkaa068.

1424