# Selenium in wheat from farming to food

**Min Wang [a], Baoqiang Li [b], Shuang Li [a], Ziwei Song [a], Fanmei Kong [a *], Xiaocun Zhang [a *]**

[a] Shandong Agricultural University, No. 61, Daizong Street, Taian, 271018, Shandong

Province, China P.R.

[b] Linyi Academy of Agricultural Sciences, No. 351, Wuhe North Street, Lanshan District,

Linyi City, 276003, Shandong Province, China P.R.

[*] Emall: fmkong@sdau.edu.cn; xczhang@sdau.edu.cn.

**Abstract:** Selenium (Se) plays an important role in human health. Approximately 80% of the

world's population does not consume enough Se which recommended by WHO (World Health

Organization). Wheat is an important staple food and Se source for most people in the world.

This article summarizes literatures about Se from 1936 to 2020 to investigate Se in wheat

farming soil, wheat, and its derived foods. Se fortification and the recommended Se level in

wheat were also discussed. Results showed that Se contents in wheat farming soil, grain, and

its derived foods around the world were 3.8–552 (mean, 220.99), 0–8,270 (mean, 347.30), and

15–2,372 (mean, 211.86) $\mu g \cdot kg^{-1}$, respectively. Adopting suitable agronomic measures could

effectively realize Se fortification in wheat. The contents in grain, flour, and its derived foods

could be improved from 93.94 to 1,181.92, 73.06 to 1,007.75, and 86.90 to 587.61 $\mu g \cdot kg^{-1}$ in

average after leaf Se fertilizer application in the field. There was a significant positive

correlation between Se content in farming soil and grain, and it was extremely the same

between foliar Se fertilizer concentration rate and grain Se increased rate. The recommended

Se fortification level in cultivation of wheat in China, India, and Spain was 18.53–23.96,

2.65–3.37, and 3.93–9.88 $g \cdot hm^{-2}$ respectively. Milling processing and food type could greatly

affect Se content of wheat derived food and should be considered seriously to meet people's Se

requirement by wheat.

**Keywords:** Se, soil, grain, flour, fortification, bioavailability

Selenium (Se) is a metalloid element of group VI A in the fourth period of the periodic

table of the elements. It was first discovered by Swedish scientist Berzelius JJ in 1817 and

named it after the Selene, the goddess moon[1]. However, researchers discovered in 1934 that

the "alkaline soil disease" and "stumble" of animals were caused by excessive intake of Se.

Since then, research on Se has mainly been based on its toxicity[2]. In 1860s, Se was determined

as an essential element in animals and human bodies and has gradually gained widespread

attention worldwide[3]. Se is one of the 14 essential trace elements in human body which

constitutes a series of important Se-containing proteins and enzymes in galacturids[4] although

the total amount of it in the human body is only 14–20 mg[4]. Se is closely related to the health

and Se deficiency can cause more than 40 kinds of diseases, such as Keshan disease, large

bone nodule disease, anemia, infertility, and muscular dystrophy[6]. Moreover, excessive Se

intake can cause hair and nail loss, skin ulceration, nerve damage, and other toxic symptoms[7].

Therefore, a safe and suitable daily Se intake should be maintained and 50–400 µg/day of Se

intake is safe and suitable according to the recommendations of World Health Organization

(WHO)[8]. Se deficiency is widely observed in more than 40 countries around the world, and

approximately 80% of the population is Se deficient[9]. The most direct source to get Se is

through dietary intake though Se bioavailability in human body varies widely (10%–85%)[10].

Therefore, improving the Se deficient situation of people through safe Se-rich diet is an

effective and long-term task.

Wheat is the second largest food crop in population consumption[11], and it provides nearly

50% daily calorie intake in most developing countries and more than 70% calorie intake in

rural areas[12]. Wheat has strong Se enrichment capacity. It can produce high-security organic Se

by absorbing Se in soil or foliage and store them in grains[12]. Se is mainly combined with

protein in wheat grain and more evenly distributed throughout the whole grain than other

minerals[14]. Therefore, unlike other micronutrients, most of the Se found in wheat grains can be

retained in the flour, rather than being lost to the bran and aleurone layer during milling. Hence,

it is an effective way to improve Se intake of people through Se enriched wheat.

Se fortification of wheat is a comprehensive and multi-objective process. Generally, the

final effects of wheat Se fortification (Se content in the food) can be significantly affected

mainly by three processes, including grain Se enriching in the field, grain grinding, and

wheat-derived food processing. This article summarized reports related to the Se fortification

of wheat, and discussed the main influencing factors, possible regulation measures and some

relatively safe and suitable reference during the above three processes. The data of Se

bioavailability was also discussed.

**2 MATERIALS AND METHODS**

**2.1 Data collection**

The papers relation to Se of wheat grain, flour and wheat derived foods published

between 1936 and 2020 were used to set up the database by a literature search using Wiley,

ACS, Science Direct, Springer Link and China National Knowledge Infrastructure (CNKI).

Search keywords included different combinations, "wheat" AND "Selenium" or "flour" AND

"Selenium" or "food" AND "Selenium". Selection criteria included: a) Field experiment:

report wheat harvest time, wheat type and fertilization type, including soil pH and Se content;

b) wheat milling: report Se fertilizer type and Se content in different components after milling;

and c) Food processing: report the source of wheat flour, food type, and Se content in food.

**2.2 Data Classification**

In selected reports, we summarized these data from soil to grain, milling, then to food

under different conditions. A total of 66 research papers and 1066 paired observations were

obtained (Figure 1); Grain Se content in different countries came from 41 references (Table

S1); Se content of different wheat milling components were also obtained under different

peeling and flour milling processing were got in 20 papers (Table S2); Se content of six kinds

of different foods under different flour processing were obtained from 26 reports (Table S3).

Total of 117 paired observations for the effects of soil Se on wheat grain from 21 articles, and

188 paired data for the effects of different Se fertilizer on wheat grain from 21 articles were

extracted.

**2.3 Data management**

The database was established using Excel 2019 software. OriginPro 8.5 and PowerPoint

2019 software were used to draw graphics. All statistical analyses were performed using SPSS

18.0 software.

**3 RESULT AND DISCUSSION**

**3.1 Se content in wheat grain**

Se content in wheat grain was 0–1500 $\mu g \cdot kg^{-1}$ (110.98 $\mu g \cdot kg^{-1}$ in average) in the world[8,

15-50]. The average Se content was lower than that suggested by the reference of Se-rich grains

(200–300 $\mu g \cdot kg^{-1}$)[39]. The large variance of Se content in wheat grain was mainly attributed the

difference of wheat varieties, area, basic properties of soil, and Se biological strengthening

measures in the field.

**3.1.1 Variance of Se content among different wheat varieties**

Generally, the old and new varieties might have different potentials to accumulate Se in

grain. According to our data, grain Se content in wheat reported in 1936–2000 could reach

198.76 µg·kg$^{-1}$ in average, and this value was significantly higher than that in 2001–2020

(Figure 1a). "Green Revolution", which began in the 1950s, dramatically increased crop

production[51] and the "dilution effect" caused by the increase of yield might be the most

important reason for the decrease of Se content in new varieties. In addition, different wheat

varieties have different ability to accumulate Se under the same soil environment. For example,

black grain wheat Se content was 177.34 µg·kg$^{-1}$, which was significantly higher than that in

white grain (70.29 µg·kg$^{-1}$)[33].

**3.1.2 Grain Se content of wheat in different continents and countries**

No significant difference was observed among the average grain Se contents of wheat in

Asia[8, 15-20, 22-25, 27-29, 31, 33-40, 43] (63.36 µg·kg$^{-1}$), Africa[8, 15-20, 22-25, 27-29, 31, 33-40, 43] (64.15 µg·kg$^{-1}$), and

Europe[20, 22, 26, 29, 30, 41, 42, 44, 46, 48, 50] (85.98 µg·kg$^{-1}$), and these values which were all lower than the

normal level (100 µg·kg$^{-1}$[39]). There was a significant difference among the grain Se content in

South America[22, 50], North America[20, 22, 47, 49, 50], and Oceania[22, 45, 50] with values of 600.00, 286.23,

and 194.80 µg·kg$^{-1}$, respectively (Figure 2). These values were significantly higher than that in

Asia, Africa, and Europe. Grain Se content in different countries on the same continent varied

greatly too. For Europe, the grain Se content in Spain, Slovakia, Portugal, and Norway were

26.75–93.90 µg·kg$^{-1}$, but that in Italy was 133.44 µg·kg$^{-1}$.

### 3.1.3 Basic properties of soil affecting grain Se content of wheat

Soil Se content and pH are the basic properties of soil which can significantly affect Se content in wheat grain. The total soil Se content varied greatly among different regions. Se-deficient, Se-sufficient, Se-rich, and Se-high soil were considered at soil Se contents of 0–175, 175–450, 450–2,000, and 2,000–3,000 $\mu g \cdot kg^{-1}$, respectively[28]. The soil in most parts of the world is Se-deficient. For instance, soil Se contents in Italy, China, India, and South Africa are 130, 21.21 12.60, and 6.14 $\mu g \cdot kg^{-1}$, respectively. Globally, soil Se content in farming system is 10–2,000 $\mu g \cdot kg^{-1}$ (300 $\mu g \cdot kg^{-1}$ in average)[28], while that in farming soil of wheat is 3.80–552.00 $\mu g \cdot kg^{-1}$. There was a significant positive correlation between soil Se content and grain Se content (Figure 3). Se content of wheat grain from the soil with Se content of 0–200 $\mu g \cdot kg^{-1}$ was significantly lower than that from the soil with Se content of 200–400 $\mu g \cdot kg^{-1}$ and 400–600 $\mu g \cdot kg^{-1}$, but no significant difference was observed between that from the soil with Se contents of 200–400 $\mu g \cdot kg^{-1}$ and 400–600 $\mu g \cdot kg^{-1}$ (Figure 1a).

Soil pH also significantly affects grain Se content in wheat, and a significant difference was observed in wheat grain Se content among neutral soil (pH 6.5–7.5), alkaline soil (pH>7.5), and acid soil (pH<6.5). The grain Se content of wheat planted in neutral soil (pH 6.5–7.5) was the highest (229.42 $\mu g \cdot kg^{-1}$), whereas that in acid soil (pH<6.5) was the lowest (55.50 $\mu g \cdot kg^{-1}$) (Figure 1a). Acid soil occupies approximately 30% of the world's ice-free land area and occurs mainly in tropical and subtropical regions. These acid soils are more likely to be Se-deficient[53].

The Se content in soil is significantly positively correlated with the Se content in grain, so planting in Se sufficient soil is an important way to obtain Se-rich wheat. However, most

cultivated soils in the world are Se insufficient, so applying Se fertilizer to enhance Se content

in wheat grains is the preferred route to consider.

**3.1.4 Effects of different Se fertilization on grain Se content in wheat**

The process of improving grain Se by fertilizer in the field is the most important part in

Se fortification. Se fertilization treatment during wheat growth can significantly increase grain

Se content according to the data about the grain Se contents in wheat around the world since

1936 (Table S1). The grain Se content of wheat planted in natural soil without Se fertilization

treatment was about 0–1,500.00 µg·kg$^{-1}$ (110.98 µg·kg$^{-1}$ in average)[8, 15-50], and that reached

58.00–8,270.00 µg·kg$^{-1}$ (677.21 µg·kg$^{-1}$ in average) with Se fertilization treatment.

However, significant differences were observed in the effect of different Se fertilizer

treatments and most of the reports indicated that foliar Se fertilization treatment might be the

most effective way to improve grain Se content[8, 15, 20, 26, 29, 35, 36, 52]. Grain Se content with foliar Se

fertilization treatment was significantly higher than that of soil Se fertilization treatment and

CK (Figure 1a). Se content in the wheat grain with soil Se fertilization treatment was

190.00–6,090.00 µg·kg$^{-1}$ (528.87 µg·kg$^{-1}$ in average)[19, 28, 29, 38, 42, 48, 54], that with foliar Se

fertilization treatment was 58.00–8,270.00 µg·kg$^{-1}$ (729.69 µg·kg$^{-1}$ in average)[8, 13, 15, 16, 18-20, 23-31,

35-38, 52, 54], and that with seed Se soaking treatment was only 64.80–300.00 µg·kg$^{-1}$ (128.67

µg·kg$^{-1}$ in average)[39, 55]. A positive correlation was observed between foliar Se fertilizer

application amount and grain Se content growth rate, and this correlation can be expressed

with the equation $Y=22.20587 X$ (Figure 4a). But there was no significant correlation between

soil Se fertilization and grain Se content, which correlation can be expressed with the equation

$Y=0.0157 X$. The use of foliar Se fertilizer was an environmentally safe and effective because

of its low concentration in solution, effective absorption, and more accurate control of total Se

application. Using 100 g Se·hm$^{-2}$ for foliar fertilization, grain Se content is 1770 μg·kg$^{-1}$; but

the same Se dosage for soil fertilization, grain Se content is only 1064 μg·kg$^{-129}$. The foliar Se

fertilizer is the most popular method in Se fortification. Moreover, $Na_2SeO_3$ is the most

common form of foliar Se fertilizer for its lower toxicity than $Na_2SeO_4$.

The grain Se content in different countries increased significantly after Se fortification in

the field (Figure 4 b), and this value increased from 93.94 μg·kg$^{-1}$ to 1181.92 μg·kg$^{-1}$ after Se

fortification. A large variance in Se fortification effects was observed in different countries. For

example, the grain Se content increased in Pakistan, India, Turkey, Spain, South Africa,

Norway, Portugal, Slovakia, China, and Italy by 2.16–25.98 times and that in Pakistan and

Italy was 2.16 and 25.98 times, respectively.

In addition to the Se application amount, Se concentration is also an important factor

affecting the effect of foliar fertilizer for one wheat variety. Wheat Se content can reach the

standard of Se fortification by foliar fertilizer application (Se concentration: 40 mg Se·L$^{-1}$,

application rate: 750 kg·hm$^{-2}$) at the early filling stage. Hence, the determination of the

appropriate application concentration and amount of foliar fertilizer are both the key to the

effect of Se nutrient enhancement of wheat.

**3.2 Se in flour and its influencing factors**

The proportion of Se in wheat grains that can be directly ingested by the human body is

also a key link for human Se supplementation. Most of the wheat will be ground into flour, and

then be made into different foods. Wheat grain is generally divided into bran, embryo, and

endosperm, accounting for 14%–16%, 2%–3%, and 81%–84% of the grain weight,

respectively[29]. During milling, wheat grains are peeled off. The bran, which consists of the

seed coat, aleurone layer and embryo, are then stripped off. Se content in bran was

15.00–520.00 µg·kg$^{-1}$ (149.48 µg·kg$^{-1}$ in average)[13, 19, 26, 28, 45], which was significantly higher

than that in flour (101.02 µg·kg$^{-1}$ in average)[13, 19, 26, 28, 30, 41, 45, 56-67]. There were also some reports

indicated that Se content in bran is approximately three times higher than that in flour after

grinding[19, 26, 45]. It was clear that, there should have a part of Se loss during the milling process.

Different wheat milling methods remarkably affect flour Se content (Figure 1 b). Flour

can mainly be divided into two types, namely, whole wheat flour and standard flour, according

to the milling method. Se content in standard flour was significantly lower than that in whole

wheat flour. According to our results (Table S2), Se content in standard flour was

13.00–500.00 µg·kg$^{-1}$ (101.02 µg·kg$^{-1}$ in average)[13, 19, 26, 28, 30, 41, 45, 56-67], and the whole wheat flour

Se content was 35.60–650.00 µg·kg$^{-1}$ (227.58 µg·kg$^{-1}$ in average)[26, 57, 62, 64, 66, 68]. Flour road

during flour milling can be divided into six flour parts (i.e., three "Broken [B]" and three

"Reduced [R]" milling parts) and two kinds of bran parts (i.e., bran and shorts). A significant

difference was observed among the ingredients in different flour roads. The flour collected by

B1, R1, B2, R2, B3, and R3 was standard flour, the flour collected by B1, R1, B2, and B3 was

bread flour, and the flour from B1 and R1 was refined flour[26, 57, 62, 64, 66, 68]. The Se content in

different components after wheat milling can be arranged as follows: B flour >R flour. The Se

contents of flour from R1, R2, R3, B1, B2, and B3 were 106.85, 128.32, 207.90, 89.00, 96.82,

and 128.29 µg·kg$^{-1}$ in average, respectively. The Se content of flour in common flour roads was

B3>B2>R3>B1>R2>R1[57, 69]. The rate of Se concentration loss and Se content loss in wheat

milling were 17.19% and 46.40%, respectively. Most of the published literature used Se

concentration loss, and a few studies have focused on Se content loss. The peeling rate of

4%–5% can effectively increase standard flour Se content. When the peeling rate was greater

than 9%, no significant difference was observed in the Se content between standard flour and

unpeeled flour[19].

Se content in standard flour and the products in each flour road can be greatly affected by

grain Se content. Se contents in whole wheat flour and standard flour in the Se-rich grain with

foliar Se fortification treatment were significantly higher than that in the control treatments,

and increased from 227.58 $\mu g \cdot kg^{-1}$ to 1,110.03 $\mu g \cdot kg^{-1}$ and from 101.02 $\mu g \cdot kg^{-1}$ to 714.90

$\mu g \cdot kg^{-1}$ respectively (Figure 1b). The Se contents in the products of different flour roads for the

Se-enhancing grain from foliar Se fortification could be arranged as B2>R3>B1>R2>R1,

which was quite different with that of common flour. The average rates of Se concentration

and content loss in wheat milling were 15.04% and 38.56%, respectively, after Se fortification

treatment.

Therefore, milling methods might affect flour Se content, and foliar Se fortification can

greatly improve Se content in standard flour and change the Se distribution in products of

different flour roads.

**3.3 Se in wheat derived foods**

Flour is generally used for the preparation of foods, such as bread, noodles, and biscuits.

These staple foods play an important role in people's diet. For example, wheat-related

traditional foods account for 75% of the total wheat consumption in China[34]. Se content in

wheat-derived foods was 3.60–1130.00 $\mu g \cdot kg^{-1}$ with an average of 191.87 $\mu g \cdot kg^{-1}$[13, 30, 41, 45, 46, 57,

59-62, 64-67, 70-81]according to the data over the past 50 years (Table S3). There was a significant

difference for the Se content of different wheat-derived foods (Figure 5). Cooking foods has

the highest Se content which can reach 382.83 $\mu g \cdot kg^{-1}$, and it was significantly higher than that

in baked foods (244.98 $\mu g \cdot kg^{-1}$). In addition, the average Se contents in pasta, noodles, biscuits,

and bread also can achieve 325.45, 251.53, 181.65, and 137.51 $\mu g \cdot kg^{-1}$, respectively.

High temperature can result in the volatilization of Se[83]. Hence, Se loss in wheat-derived

foods processing is mainly occurs during heating, especially under the dual effect of

dissolution and heat volatilization when cooking, steaming, and frying the same food[82, 83].

Moreover, soaking steps in food fermentation also can reduce wheat-derived foods Se content.

For example, Se content in mung bean and millet would decrease by 7.17% and 2.62%,

respectively during soaking[83]. But no report about the change of Se in wheat grain during

soaking and the flour fermentation was retrieved.

In addition, increasing grain Se content through foliar Se fortification can effectively

improve Se content in wheat-derived foods (Figure 1c). For example, Se content in pasta

increased from 22.4 $\mu g \cdot kg^{-1}$ to 424.43 $\mu g \cdot kg^{-1}$ (17.94 times) and that in bread increased from

119.15 $\mu g \cdot kg^{-1}$ to 644.69 $\mu g \cdot kg^{-1}$ (4.41 times) using the flour derived from foliar Se

fortification. Hart et al.[57] found that foliar Se application rate is positively correlated with bread

Se content, and their relationship can be expressed using the formula $Y=19.2262\,X+95.95332$.

Food processing can greatly affect Se content in food, and increasing Se content in grain

through foliar Se fortification is an effective way to improve Se content in wheat-derived

foods.

**3.4 Se bioavailability**

Not all the Se obtained from foods can be used by the human body. The ratio of body

absorbed Se to total intake Se is called bioavailability[10]. The Se from food can be effectively

used by the human body only through the absorption in blood circulation or tissues and then

transformed into active substances. A large difference was observed for the bioavailability of

different Se types. Se in wheat-derived foods mainly can be divided into inorganic Se (i.e.,

selenite) and organic Se (i.e., SeMet). Organic SeMet is the main form in wheat grain, flour,

and wheat-derived foods, which accounts for 70%–90%[14, 84], 59%–82%[43, 78, 85, 86], and

42%–83%[43, 57, 85] of total Se, respectively.

Different forms of Se are digested and absorbed in different ways, causing different

bioavailability. Organic Se is the mainly forms which can be actively absorbed by human

small intestine, and the bioavailability generally fluctuates between 70% and 90%[43, 57, 85].

Inorganic Se can be absorbed via simple absorption (selenite) and passive diffusion (selenite),

and the bioavailability is generally less than 50%[87]. Moreover, the long-term consumption of

inorganic Se may result in toxic and side effects[10]. The molecular of $H_2Se$ is the key in the Se

metabolism of human body. The role of $H_2Se$ mainly includes : 1) $H_2Se$ first transforms to

selenite and SeMet form, and then synthesizes selenoprotein in liver and 2) SeMet and

SeMeCys catalyze $H_2Se$ to form different Se ions forms under continuous methylation, which

are discharged through respiration, urine, and sweat[88, 89].

Se bioavailability in human body can be predicted via *in vitro* simulated gastrointestinal

digestion model[88, 89], Caco-2 based cell model[91], animal models[92], and human Se bioavailability

detection[93]. Se bioavailability in wheat was 83% in rat feeding trials, while those in mushroom,

tuna, and beef kidneys were 5%, 57%, and 97%, respectively[92]. Se bioavailability in wheat

flour was also very high, which values was 75.4%–91.8%[94]. The consecutive consumption of

Se-enriched wheat for six weeks could significantly increase human serum Se content, but

Se-enriched fish had no obvious effect[93]. Generally, SeMet can be effectively absorbed by

human body, and its bioavailability may exceed 90%. By using *in vitro* gastrointestinal

digestion method to simulate bread digestion, soluble Se content after gastric and

gastrointestinal digestion accounted for 75% and 89% of total Se content in bread, respectively,

and SeMet accounted for 95.5% and 96.8% of soluble Se[93].

Se bioavailability in foods is 10%–85%[10]. These differences in bioavailability might be

ascribed to the following factors: 1) Se type in food - different Se types showed different

bioavailability. For instance, SeMet and SeMeCys have a bioavailable of up to 90%[92]. The

bioavailability of SeMet is approximately 1.5-fold higher than that of selenite[96]. 100% of

selenate can be taken up, but a significant fraction will be excreted in the urine. Yet, the direct

absorption of selenite is approximately 50%–60%[97]. 2) Food type - a high-fat diet can hinder

Se absorption, resulting in low Se bioavailability[90]. 3) Food processing - heat treatment not

only affects total Se content but also reduced Se antagonist content. However, it's worth noting

that proper heat treatment might improve Se bioavailability too[92]. 4) Antagonism with other

minerals - Ca intake in the human body would reduce Se bioavailability. Meanwhile, Vc can

transform selenate into insoluble Se compounds in a short time, thus reducing Se

bioavailability[98]. 5) Digestion and absorption capacity of human body. For instance,

gastrointestinal diseases will reduce Se bioavailability[98].

The reports above indicated that wheat-related foods were effective Se source for people

because most of the Se in wheat-related foods are organic SeMet which has high

bioavailability. So, improving Se content in the wheat grain by foliar Se fortification is the key

step and effective way to supplement Se in human body in most of the Se deficient areas.

**3.5 Suitable level of Se in wheat grain and in fertilizer applied in the field**

Both insufficient and excessive Se intake will have adverse effects on human health.

Therefore , it should be seriously considered about the suitable Se level in the grain of wheat

and the proper Se fertilizer quantity in the field which can be determined according to some

important factors including the proper Se intake quantity per capita, dietary pattern, wheat Se

accumulation efficiency and Se loss in food processing etc.

The range of Se intake recommended by WHO is 50–400 µg/day and this can be the

reference to determine the proper Se intake quantity per capita of people. Dietary pattern,

which refers to the combination of various forms of food ingredients used in people's actual

life, should also be considered[100, 101]. The representative dietary patterns are divided into

Western, Oriental, and Mediterranean dietary patterns. Western dietary patterns are mainly

characterized by the intake of large amounts of red meat, processed meat products, refined

cereals, desserts, French fries, and high-fat dairy products. Oriental dietary patterns are mainly

characterized by the intake of a large number of fruits, vegetables, legumes, whole wheat, an

appropriate amount of fish, and poultry. Mediterranean dietary patterns are mainly

characterized by the intake of monounsaturated fatty acids rich in olive and olive oil, whole

grains, fruits, vegetables and dairy products, weekly intake of fish, poultry, nuts, and legumes.

The distribution law of food Se content is as follows[62]: fish > meat > egg > cereal > fruits and

vegetables. Hence, the actual Se intake per capita in western dietary patterns countries could

reach the recommended level. The actual Se intake per capita in some Mediterranean dietary

pattern countries also could reach the recommended level, such as Italy, but some countries

such as Spain cannot not reach the recommended level. While the actual Se intake per capita in

Oriental dietary patterns countries, such as China and India, did not reach the recommended

level.

With wheat as Se source, the level of Se needed in wheat grains and the amount of Se to

be applied in the field should be determined.

*Daily insufficient Se intake = Recommended Se intake per capita-Actual Se intake per capita*

*Total Se loss rate*

$$= \frac{\textit{Unit weight of grain} \times \textit{Grain Se content-Unit weight of grain} \times \textit{Flour extraction} \times \textit{Flour Se content}}{\textit{Unit weight of grain} \times \textit{Grain Se content}} \times 100\%$$

$$\textit{Recommended grain intake per capita} = \frac{\textit{Recommended flour intake per capita}}{\textit{Flour extraction}}$$

*Suitable Se fortification Level in grain*

$$= \frac{\textit{Daily insufficient Se intake + Actual Se intake per capita + proportion of Se intake by grain (daily Se intake)}}{\textit{Recommended grain intake per capita} \times \textit{(1-Total Se loss rate)}}$$

The recommended Se intake per capita is 60–400 $\mu g \cdot day^{-1}$, the recommended Se in grain

is 150 g (assuming all for wheat), the actual Se intake is 43.90 $\mu g \cdot kg^{-1}$, and the grain daily Se

intake accounts for approximately 22.6% in China according to the national nutrition and

health guidelines. The upper limit of Se content in wheat grain is 300 $\mu g \cdot kg^{-1}$, and the scope of

Se content in wheat grain is 242.74–300.00 $\mu g \cdot kg^{-1}$ in China. According to the formula and the

parameters, the amount of Se fortification by foliar Se fertilizer in China should be

18.53–23.96 $g \cdot hm^{-2}$ to meet people's daily need. The result varied widely between different

countries, for example, the amount of Se fortification should be 2.65–3.37 and 3.93–9.88

$g \cdot hm^{-2}$ in India and Spain, respectively (Figure 7, Table 1 [8, 98, 102-107]).

**Table 1 Recommended amount of wheat Se fertilizer in different countries**

| Dietary pattern | Country | Recommended daily Se intake per capita (µg·d⁻¹) | Wheat grain Se content (µg·kg⁻¹) | Actual Se intake per capita (µg·d⁻¹) | Recommended wheat grain Se content (µg·kg⁻¹) | Recommended Se fertilizer applying amount (g·hm⁻²) |
|---|---|---|---|---|---|---|
| Oriental dietary patterns | China | 60.00 [100] | 47.46 | 43.90 [97] | 242.74-300.00 | 18.53-23.96 |
| | India | 50.00 [8] | 171.70 | 48.00 [98] | 272.76-300.00 | 2.65-3.37 |
| Mediterranean dietary patterns | Spain | 50.00 [8] | 93.90 | 49.24 [95] | 175.84-300.00 | 3.93-9.88 |
| | Italy | 50.00 [8] | 133.44 | 66.53 [96] | No need to fortification | No need to fortification |
| Western dietary patterns | Australia | 55.00 [92] | 191.83 | 57.00 [99] | No need to fortification | No need to fortification |
| | America | 55.00 [92] | 304.95 | 80.00 [98] | No need to fortification | No need to fortification |

## 4 CONCLUSION AND FUTURE TRENDS

All the reports now showed that Se nutrition enhancement of human body through the consumption of wheat in the diet is an effective and feasible way. But the effects were mainly influenced by three important processes, including 1) the acquisition of Se-rich wheat grains, 2) reduction of Se loss in the process of flour milling and food processing, 3) the improvement of bioavailability of Se in food. There were many confirmed conclusions we can make and also many further researches needed to conduct in the future based on the above three aspects.

1) The acquisition of Se-rich wheat grains. It was the most important step in Se fortification. Application of Se foliar fertilizer is the first choice which is more effective and economic to get Se-rich grain than soil fertilization according to the available reports. The

application of foliar fertilizer 750 kg·hm$^{-2}$ with Se concentration 40 mg·L$^{-1}$ at the early filling

stage can most likely make grain Se content reach 200–300 μg·kg$^{-1}$. How to improve Se

efficiency of soil in the cultivation of wheat is another aspect needed further concentration. It

was necessary to further investigate the relationship between soil properties and Se

bioavailability for the plants in the soil, and these research combined with other special

methods might be helpful to improve Se efficiency of soil. In addition, the wheat varieties are

another important factor which should be considered seriously. Many reports confirmed the

large variance between different wheat varieties[33]. The grain Se content of the varieties with

high Se accumulation potential was 742.33% higher than that of wheat varieties with low Se

accumulation[8]. Therefore, it's necessary to selection, breeding and application of wheat

varieties with strong Se enrichment ability. Some genes/varieties related with high

accumulation of Se in grain should be considered in breeding[8]. Few investigations were

conducted on the effects of other nutrients on wheat Se accumulation such as foliar Se

fortification increased grain Se and Zn content, but reduced grain Cu, Fe, K, Mg and S

content[109].

2) Reduction of Se loss in the process of flour milling and food processing. Flour Se

content can be retained to the maximum extent when the peeling rate is 4%–5% and the

milling ratio is 70%–80% , and try to choose low temperature method to keep Se content in

wheat derived-foods according to the available reports. But few reports focused on the

mechanism of the effects on Se loss in different food processing methods.

3) The improvement of bioavailability of Se in food. The metabolism and regulation of

different forms of Se in body and the interaction mechanism with other substances needed

further investigation. The information will contribute to the other ways to supplement Se of

human body. For example, the addition of food additives can improve Se bioavailability in

food. This process should also be considered in the food industry.

Se deficiency is a worldwide problem. It is an effective and feasible way to supplement

Se for people through Se fortification in wheat. We can adopt the strategy including the above

three aspects to get enough Se from wheat. But this strategy needs more public attention, more

research and application results to achieve this goal.

**5 CONFILCT OF INTEREST**

*The authors declare that the research was conducted in the absence of any commercial or*

*financial relationships that could be construed as a potential conflict of interest.*

**6 AUTHOR CONTRIBUTIONS**

*MW, SL and ZWS collected and analysed data. MW, FMK and XCZ wrote this paper. XCZ,*

*FMK and BQL conceived and modified this paper.*

**7 ACKNOWLEDGMENTS**

*This paper was supported by Agricultural Variety Improvement Project of Shandong*

*Province (2019LZGC001) and Special Project of Regional Science and Technology*

*Collaborative Innovation between District Science and Technology Department and Rikaze*

*Municipal Government in 2020 (Research and Demonstration Application of Bioaugmentation*

*Technology for trace nutrients of highland barley in Rikaze area).*

**8 REFERENCE**

(1) Schwarz, K.; Foltz, C. M. Selenium as an integral part of factor 3 against dietary necrotic

liver degeneration. *J Am Chem Soc.* **1957**, 79, 3292-3293.

(2) Zhang, J. J. The Biological functions of selenium and research development of

se-enriched foodstuff. *Studies of Trace Elements and Health*. **2006**, 23, 58-60. **(In Chinese).**

(3) Kilness, A. W. Selenium and public health. *South Dakota Journal of Medicine*. **1973**, 26,

17-19.

(4) Arteel, G. E.; Sies, H. The biochemistry of selenium and the glutathione system. *Environ.*

*Toxicol. Phar.* **2001**, 10, 153-158.

(5) Huang, Q. J.; Wang, S. S.; Zheng, W.; Dai, G. Z.; Yang, J. Research progress of

selenium-rich in Wheat. *J. Green Sci. Technol.* **2015**, 12, 73-76. **(In Chinese).**

(6) Rayman, M. P. The importance of selenium to human health. *The Lancet.* **2000**, 356,

233-241.

(7) Johnson, L. J.; Meacham, S. L.; Kruskall, L. J. The Antioxidants-Vitamin C, Vitamin E,

Selenium, and Carotenoids. *J Agromedicine*. **2003**, 9, 65-82.

(8) Liu, H.; Yang, Y. E.; Wang, C. H.; Li, F. C.; Li, K. Y.; Yang, N.; Wang, S.; Wang, H.; He,

J.; Dai, S. Selenium content of wheat grain and its regulation in different wheat production

regions of china. *Sci. Agric. Sin*. **2016**, 49, 1715-1728. **(In Chinese).**

(9) Stoffaneller, R.; Morse, N. L. A review of dietary selenium intake and selenium status in

Europe and the Middle East. *Nutrients*. **2015**, 7, 1494-1537.

(10) Li, Y. N.; Ma, L. Y.; Huang, K. L.; Luo, Y. B.; Mei, X. H. Research advancement on

bioavailability of selenium in dietary sources. *Food and Nutrition in China.* **2019**, 25, (09), 5-9.

**(In Chinese).**

(11) Cardoso, R. C.; Fernandes, Â.; Gonzaléz-Paramás, A. M.; Barros, L.; Ferreira, I. F. R.

Flour fortification for nutritional and health improvement: A review. *Food Res. Int.* **2019**, 125,

108576.

(12) Cakmak, I. Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant Soil.* **2008**, 302, 1-17.

(13) Wang, P.; Zhang, X. C.; Kong, F. M.; Shi, Y. X.; Dong, H. Z. Change research of selenium in the processing from grain to steamed bread. *Current Biotechnology.* **2016**, 6, 422-427. **(In Chinese).**

(14) Cubadda, F.; Aureli, F.; Ciardullo, S.; D'Amato, M.; Raggi, A.; Acharya, R.; Reddy, R. A. V.; Prakash, N. T. Changes in Selenium Speciation Associated with Increasing Tissue Concentrations of Selenium in Wheat Grain. *J. Agr. Food Chem.* **2010**, 58, 2295-2301.

(15) Wang, M.; Ali, F.; Wang, M. K.; Dinh, Q. T.; Liang, D. Understanding boosting selenium accumulation in Wheat (Triticum aestivum L.) following foliar selenium application at different stages, forms, and doses. *Environ. Sci. Pollut. R.* **2020**, 27, 717-728.

(16) Zhang, P. P.; Ma, H. X.; Yao, J. B.; Zhang, P. Effects of selenium foliar spray on selenium distribution in milling fractions in common wheat. *Acta Agric.Nucl. Sin.* **2019**, 33, 2254-2260. **(In Chinese).**

(17) Qin, C.; Shi, C.; Wan, Q. Y.; Wang, L.; Liu, A. Q.; An, C. X. Speciation analysis of selenium in selenium-enriched wheat by PLC-ICP-MS. *Food R&D.* **2019**, 40, 140-144. **(In Chinese).**

(18) Dong, S. F.; Sun, M.; Zhao, J. M.; Bai, X.; Lu, P. F.; Gao, Z. Q.; Ren, A. X.; Hou, F. F.; Yin, X. B. Effects of sodium selenite application at anthesis stage on the growth characteristics and Se accumulation in wheat plants. *J. Shanxi Agric. Univ. (Nat. Sci.).* **2019**, 39, 13-18. **(In Chinese).**

(19) Chen, B. N. Research on the effect of selenium application technology in winter wheat.

In: Huazhong Agricultural University. **2019**. **(In Chinese).**

(20) Zou, C.; Du, Y.; Rashid, A.; Ram, H.; Savasli, E.; Pieterse, P. J.; Ortiz-Monasterio, I.;

Yazici, A.; Kaur, C.; Mahmood, K.; Singh, S.; Le Roux, M. R.; Kuang, W.; Onder, O.; Kalayci,

M.; Cakmak, I. Simultaneous Biofortification of Wheat with Zinc, Iodine, Selenium, and Iron

through Foliar Treatment of a Micronutrient Cocktail in Six Countries. *J. Agr. Food Chem.*

**2019**, 67, 8096-8106.

(21) López-Bellido, F. J.; Sanchez, V.; Rivas, I.; López-Bellido, R. J.; López-Bellido, L.

Wheat grain selenium content as affected by year and tillage system in a rainfed Mediterranean

Vertisol. *Field Crop. Res.* **2019**, 233, 41-48.

(22) Ji, H. T.; Wang, M.; Cao, Y.; Li, X. L.; Jiang, L. F.; Ma, X. F. Correlation analysis of Fe,

Zn, Se mass fraction in Durum and Emmer Wheat. *Acta Agric Bore-occ Sin*. **2018**, 27, 772-778.

**(In Chinese).**

(23) Wang, J. M.; Chen, H. Y.; Guo, J. J.; Wang, M.; Du, B.; Shi, L.; Xu, Y. Y.; Jiang, S. C.;

Jiang, H. F.; Zhou, L.; Chen, H.; Xing, D. Y.; Xie, Y. M. Preliminary study on the effect of

selenium foilar fertilizer on agronomic characters, yield and selenium content of wheat.

*Journal of Yangtze University (Natural Science Edition)*. **2018**, 15, 1-4. **(In Chinese).**

(24) Feng, K.; Jia, L. L.; Jia, Y. H.; Erken, D. L. X. T.; Liu, J.; Shi, S. B. Effect of selenium

application on yield and selenium content of spring wheat under different plantin. *J. Triticeae*

*Crops*. **2018**, 38, 834-840. **(In Chinese).**

(25) Jin, Y. F.; Gao, B. D. Effect of selenium fertilizer on Se-enrich reaction and Se

absorption and distribution of spring wheat. *China Soils Fert*. **2018**, 274, 113-117. **(In**

**Chinese).**

(26) De Vita, P.; Platani, C.; Fragasso, M.; Ficco, D. B. M.; Colecchia, S. A.; Del Nobile, M.

A.; Padalino, L.; Di Gennaro, S.; Petrozza, A. Selenium-enriched durum wheat improves the

nutritional profile of pasta without altering its organoleptic properties. *Food Chem.* **2017**, 214,

374-382.

(27) Zhang, R.; Liu, M. S.; Wang, R. C.; Li, S. L.; Wang, N.; Wang, J. M. Effect of foliar

spraying of plant Se-nutrients on wheat yield and quality. *J. Triticeae Crops.* **2015**, 35, 856-859.

**(In Chinese).**

(28) Yu, R. Effects of different valence selenium and its applications on wheat yield and

selenium content in the typical selenium deficient area of the loess plateau. In: Northwest A&F

University. **2015**. **(In Chinese).**

(29) Galinha, C.; María, S. M.; Adriano, M. P.; Maria, C. F.; Coutinho, S.; Benvindo, M.;

Almeida, A. S.; María, T. P.; Yolanda, M.; Wolterbeek, H. T. Characterization of

selenium-enriched wheat by agronomic biofortification. *J Food Sci Tech Mys*. **2015**, 52,

4236-4245.

(30) Poblaciones, M. J.; Rodrigo, S.; Santamaría, O.; Chen, Y.; McGrath, S. P. Agronomic

selenium biofortification in Triticum durum under Mediterranean conditions: From grain to

cooked pasta. *Food Chem.* **2014**, 146, 378-384.

(31) Meng, L. M.; Yang, Z. G.; Zhang, K.; Sun, J. W.; Ji, T. H. Effect of application of race

element fertilizer on wheat yield and content of trace elements. *J Anhui Agric Sci*. **2014**, 14,

4283-4285. **(In Chinese).**

(32) Beladel, B.; Nedjimi, B.; Mansouri, A.; Tahtat, D.; Belamri, M.; Tchanchane, A.;

Khelfaoui, F.; Benamar, M. E. A. Selenium content in wheat and estimation of the selenium

daily intake in different regions of Algeria. *Appl. Radiat. Isotopes.* **2013**, 71, 7-10.

(33) Pei, Z. Y.; Wen, H. Q.; Zhang, L. S.; Cheng, T. L.; Li, X. Study on rich selenium black

kernel wheat donghei No.1. *Seed*. **2013**, 32, 74-76. **(In Chinese).**

(34) Wang, X. X.; Jia, A. X.; Yu, Z. L. Distribution of mineral elements in wheat milling.

*Modern Flour Milling Industry*. **2012**, 26, 23-26. **(In Chinese).**

(35) Zhang, X.; Bu, D. N.; Li, R. N. Effects of foliar spraying microelement fertilizers on

yield and quality of winter wheat. *J. Triticeae Crops*. **2012**, 32, 747-749. **(In Chinese).**

(36) Zhang, J. Y.; Zhang, P. P.; Ma, H. X.; Yao, J. B.; Geng, Z. M. Effects of spraying

micronutrients on yield,quality and micronutrient content in grain of wheat. *Acta Agric.*

*Jiangxi*. **2012**, 24, 64-66. **(In Chinese).**

(37) Zhang, Y. Effects of selenium and zinc fertilizer on yield and selenium absorption and

accumulation in spring wheat. *Guangxi Agric Sci*. **2012**, 43, 626-629. **(In Chinese).**

(38) Tang, Y. X.; Wang, H. M.; Yang, J. F.; Lv, Y. H. Studies on the selenium content and

selenium enriched technique of winter wheat in hebei province. *J. Triticeae Crops*. **2011**, 31,

347-351. **(In Chinese).**

(39) Tang, Y. X.; Wang, H. M.; Lv, Y. H.; Liu, Q. L.; Yang, J. F. Effects of selenium seed

soaking on growth, yield and seeds selenium content of wheat. *J. Triticeae Crops.* **2010**, 30,

731-734. **(In Chinese).**

(40) Lu, L.; Ji, Y. M.; Li, L. R.; Li, Z. L.; Wu, Y. Analysis of Fe, Zn and Se contents in

different wheat cultivars (Lines) planted in different areas. *Chinese Journal of Applied &*

*Environmental Biology.* **2010**, 16, 646-649. **(In Chinese).**

(41) Cubadda, F.; Aureli, F.; Raggi, A.; Carcea, M. Effect of milling, pasta making and

cooking on minerals in durum wheat. *J. Cereal Sci.* **2009**, 49, 92-97.

(42) Ducsay, L.; Lozek, O.; Varga, L. The influence of selenium soil application on its

content in spring wheat. *Plant Soil & Environment.* **2009**, 55, 80-84.

(43) Yadav, S. K.; Singh, I.; Sharma, A.; Singh, D. Selenium status in food grains of

northern districts of India. *J. Environ. Manage.* **2008**, 88, 770-774.

(44) Spadoni, M.; Voltaggio, M.; Carcea, M.; Coni, E.; Raggi, A.; Cubadda, F. Bioaccessible

selenium in Italian agricultural soils: Comparison of the biogeochemical approach with a

regression model based on geochemical and pedoclimatic variables. *Sci. Total Environ.* **2007**,

376, 160-177.

(45) Lyons, G.; Ortiz-Monasterio, I.; Stangoulis, J.; Graham, R. Selenium concentration in

wheat grain: Is there sufficient genotypic variation to use in breeding? *Plant & Soil.* **2005**, 269,

369-380.

(46) Díaz-Alarcón, J. P.; Navarro-Alarcón, M.; Lbpez-Garcia de la Serrana, H.;

López-Martínez, M. C. Determination of selenium in cereals, legumes and dry fruits from

southeastern Spain for calculation of daily dietary intake. *Sci. Total Environ.* **1996**, 184,

183-189.

(47) Wyatt, C. J.; Meléndez, J. M.; Acu A, N.; Rascon, A. Selenium (Se) in foods in northern

Mexico, their contribution to the daily Se intake and the relationship of Se plasma levels and

glutathione peroxidase activity. *Nutr. Res.* **1996**, 16, 949-960.

(48) Tveitnes, S.; Singh, B. R.; Ruud, L. Selenium concentration in spring wheat as

influenced by basal application and top dressing of selenium-enriched fertilizers. *Fertilizer*

*Research.* **1995**, 45, 163-167.

(49) Harn, M. H.; Kuennen, R. W.; Caruso, J. A.; Fricke, F. L. Determination of trace amounts of selenium in corn, lettuce, potatoes, soybeans, and wheat by hydride generation/condensation and flame atomic absorption spectrometry. *J. Agr. Food Chem.* **1981**, 29, 792–796.

(50) Robinson, W. O. Selenium Content of Wheat from Various Parts of the World. *Ind. Eng. Chem.* **1936**, 6, 736-738.

(51) Evenson, R. E.; Gollin, D. Assessing the Impact of the Green Revolution, 1960 to 2000. *Sci*. **2003**, 300, 758-762.

(52) Galinha, C.; Freitas, M. C.; Pacheco, A. M. G.; Coutinho, J.; Ma S, B.; Almeida, A. S. Determination of selenium in bread-wheat samples grown under a Se-supplementation regime in actual field conditions. *J Radioanal Nucl Ch*. **2012**, 291, 231-235.

(53) Von Uexküll, H. R.; Mutert, E. Global extent, development and economic impact of acid soils. *Springer Netherlands.* **1995**, 171, 1-15.

(54) Wang, J. W. Effects of selenium and zinc on yield and mineral nutrition of main crops in dryland. In: Northwest A&F University. **2012**. **(In Chinese).**

(55) Souza, G. A.; Hart, J. J.; Carvalho, J. G.; Rutzke, M. A.; Albrecht, J. C.; Guilherme, L. R. G.; Kochian, L. V.; Li, L. Genotypic variation of zinc and selenium concentration in grains of Brazilian wheat lines. *Plant Sci.* **2014**, 224, 27-35.

(56) Li, X. Z.; Wang, X. X.; Ma, S.; Ye, M. X. Effect of wheat peeling technology on selenium content in flour. *Cereals & Oils.* **2017**, 30, 50-54. **(In Chinese).**

(57) Hart, D. J.; Fairweather-Tait, S. J.; Broadley, M. R.; Dickinson, S. J.; Foot, I.; Knott, P.;

McGrath, S. P.; Mowat, H.; Norman, K.; Scott, P. R.; Stroud, J. L.; Tucker, M.; White, P. J.;

Zhao, F. J.; Hurst, R. Selenium concentration and speciation in biofortified flour and bread:

Retention of selenium during grain biofortification, processing and production of Se-enriched

food. *Food Chem.* **2011**, 126, 1771-1778.

(58) Matos-Reyes, M. N.; Cervera, M. L.; Campos, R. C.; Guardia, M. Total content of As,

Sb, Se, Te and Bi in Spanish vegetables, cereals and pulses and estimation of the contribution

of these foods to the Mediterranean daily intake of trace elements. *Food Chem.* **2010**, 122,

188-194.

(59) Wang, H. B.; Li, Q. L.; Liu, J. L.; Gan, P. Y.; Wang, Y. F.; Zhu, G. M. Development of

kappa-selenocarrageen in premix and determination of se content in wheat flour and its

products. *Cereal & Feed Industry*. **2008**, 6, 5-8. **(In Chinese).**

(60) Alamin, M. B.; Bejey, A. M.; Kučera, J.; Mizera, J. Determination of mercury and

selenium in consumed food items in Libya using instrumental and radiochemical NAA. *J.*

*Radioanal. Nucl. Ch.* **2006**, 270, 143-146.

(61) Klapec, T.; Mandi, M. L.; Grgi, J.; Primorac, L.; Krstanovi, V. Selenium in selected

foods grown or purchased in eastern Croatia. *Food Chem.* **2004**, 85, 445-452.

(62) Murphy, J.; Cashman, K. D. Selenium content of a range of Irish foods. *Food Chem.*

**2001**, 74, 493-498.

(63) Díaz-Alarcón, J. P.; Navarro-Alarcón, M.; de la Serrana, H. L.; López-Martínez, M. C.

Determination of selenium in cereals, legumes and dry fruits from southeastern Spain for

calculation of daily dietary intake. *Sci. Total Environ.* **1996**, 184, 183-189.

(64) Barclay, M. N. I.; MacPherson, A.; Dixon, J. Selenium Content of a Range of UK

Foods. *J. Food Compos. Anal.* **1995**, 8, 307-318.

(65) Oster, O.; Prellwitz, W. The daily dietary selenium intake of West German adults. *Biol.*

*Trace Elem. Res.* **1989**, 20, 1-14.

(66) Arthur, D. Selenium Content of Canadian Foods. *Canadian Institute of Food Science*

*and Technology Journal.* **1972**, 5, 165-169.

(67) Madaric, A.; Kadrabova, J.; Ginter, E. The selenium content of selected food from the

Slovak Republic. *Food Chem.* **1997**, 58, 29-32.

(68) Gao, X. Y.; Gao, Q. Z.; Zhu, Y. R.; Wang, C. Q. Determination oftrace selenium content

in nanyang color wheat. Science and Technology of Cereals. *Oils and Foods.* **2015**, 23, 67-70.

**(In Chinese).**

(69) Sager, M.; Werteker, M. Zur Elementverteilung in Mahlfraktionen von drei

Weizensorten an zwei niederösterreichischen Standorten. *Nutrition.* **2010**, 34, 357-367.

(70) He, Y. The development and antioxidation of selenium rich biscuits. *The Food Industry.*

**2019**, 40, 153-157. **(In Chinese).**

(71) Zhai, L. G.; Li, L. J.; Guo, Y. X.; Wang, J. Y.; Wang, B. C. Development of

selenium-enriched rye crisp biscuit. *Cereal & Feed Industry.* **2018**, 11, 29-33. **(In Chinese).**

(72) Wang, X. Y.; Zhang, Q. F. The development of nutrition rich selenium yeast bread.

*Cereals Oils.* **2015**, 28, 25-27. **(In Chinese).**

(73) Bryszewska, M. A.; Ambroziak, W.; Langford, N. J.; Baxter, M. J.; Colyer, A.; Lewis,

D. J. The Effect of Consumption of Selenium Enriched Rye/Wheat Sourdough Bread on the

Body's Selenium Status. *Plant Food. Hum. Nutr.* **2007**, 62, 121-126.

(74) Pappa, E. C.; Pappas, A. C.; Surai, P. F. Selenium content in selected foods from the

Greek market and estimation of the daily intake. *Sci. Total Environ.* **2006**, 372, 100-108.

(75) Holben, D. H.; Smith, A. M. The diverse role of selenium within selenoproteins: a

review. *J Am Diet Assoc.* **1999**, 99, 836-843.

(76) Finley, J. W.; Matthys, L.; Shuler, T.; Korynta, E. Selenium content of foods purchased

in North Dakota. *Nutr. Res.* **1996**, 16, 723-728.

(77) Aro, A.; Kumpulainen, J.; Alfthan, G.; Voshchenko, A. V.; Ivanov, V. N. Factors

affecting the selenium intake of people in Transbaikalian Russia. *Biol. Trace Elem. Res.* **1994**,

40, 277-285.

(78) Eurola, M. H.; Ekholm, P. I. S. H.; Ylinen, M. E.; Varo, P. T.; Koivistoinen, P. E.

Selenium in Finnish foods after beginning the use of selenate-supplemented fertilisers. *J. Sci.*

*Food Agr.* **1991**, 56, 57-70.

(79) Olson, O. E.; PALMER, I. S. Selenium in Foods Purchased or Produced in South

Dakota. *J. Food Sci.* **1984**, 49, 446-452.

(80) Ferretti, R. J.; Levander, O. A. Effect of milling and processing on the selenium content

of grains and cereal products. *J Agric Food Chem.* **1974**, 22, 1049.

(81) Laila, H.; Bruggeman, J. Selenium analysis of selected Egyptian foods and estimated

daily intakes among a population group. *Food Chem.* **1999**, 65, 527-532.

(82) Guo, H.; Huang, X. S.; Hui, J. R.; Chen, Y. H.; Shen, Q. Effect of processing on

selenium content of wheat. *Food Sci Tech-brazil.* **2017**, 42, 185-188. **(In Chinese).**

(83) Khanam, A.; Platel, K. Influence of domestic processing on the bioaccessibility of

selenium from selected food grains and composite meals. *J Food Sci Tech Mys.* **2016**, 53,

1634-1639.

(84) Kirby, J. K.; Lyons, G. H.; Karkkainen, M. P. Selenium speciation and bioavailability in

biofortified products using species-unspecific isotope dilution and reverse phase ion

pairing-inductively coupled plasma-mass spectrometry. *J Agric Food Chem.* **2008**, 56, 1772.

(85) Hawkesford, M. J.; Zhao, F. Strategies for increasing the selenium content of wheat. *J.*

*Cereal Sci.* **2007**, 46, 282-292.

(86) D Az Huerta, V.; Hinojosa Reyes, L.; Marchante-Gay N, J. M.; Fern Ndez S Nchez, M.

A. L.; Sanz-Medel, A. Total determination and quantitative speciation analysis of selenium in

yeast and wheat flour by isotope dilution analysis ICP-MS. *J. Anal. Atom. Spectrom.* **2003**, 18,

1243-1247.

(87) Thiry, C.; Ruttens, A.; De Temmerman, L.; Schneider, Y.; Pussemier, L. Current

knowledge in species-related bioavailability of selenium in food. *Food Chem.* **2012**, 130,

767-784.

(88) Wrobel, J. K.; Power, R.; Toborek, M. Biological activity of selenium: Revisited.

*International Union of Biochemistry and Molecular Biology Life.* **2016**, 68, 97-105.

(89) Gerald, F. C. Biomarkers of Selenium Status. *Nutrients.* **2015**, 7, 2209-2236.

(90) Shen, L.; Van Dyck, K.; Luten, J.; Deelstra, H. Diffusibility of selenate, selenite,

seleno-methionine, and seleno-cystine during simulated gastrointestinal digestion. *Biol Trace*

*Elem Res.* **1997**, 58, 55-63.

(91) Artursson, P.; Karlsson, J. Correlation between oral drug absorption in humans and

apparent drug permeability coefficients in human intestinal epithelial (Caco-2) cells. *Biochem.*

*Bioph. Res. Co.* **1991**, 175, 880-885.

(92) Thomson, C. D. Assessment of requirements for selenium and adequacy of selenium

status: a review. *Eur. J. Clin. Nutr.* **2004**, 58, 391-402.

(93) Meltzer, H. M.; Bibow, K.; Paulsen, I. T.; Mundal, H. H.; Norheim, G.; Holm, H. Different bioavailability in humans of wheat and fish selenium as measured by blood platelet response to increased dietary Se. *Biol. Trace Elem. Res.* **1993**, 36, 229-241.

(94) Govasmark, E.; Brandt-Kjelsen, A.; Szpunar, J.; Bierla, K.; Vegarud, G.; Salbu, B. Bioaccessibility of Se from Se-enriched wheat and chicken meat. *Pure Appl. Chem.* **2010**, 82, 461-471.

(95) Sánchez-Martínez, M.; Pérez-Corona, T.; Cámara, C.; Madrid, Y. Preparation and Characterization of a Laboratory Scale Selenomethionine-Enriched Bread. Selenium Bioaccessibility. *J Agric Food Chem.* **2015**, 63, 120-127.

(96) Burk, R. F.; Hill, K. E. Regulation of Selenium Metabolism and Transport. Annu. Rev. Nutr. **2015**, 35, 109-34.

(97) Roman, M.; Jitaru, P.; Barbante, C. Selenium biochemistry and its role for human health. *Metallomics.* **2014**, 6, 25-54.

(98) Rahavi, E. B.; Altman, J. M.; Stoody, E. E. *Dietary Guidelines for Americans, 2015–2020: National Nutrition Guidelines*. **2019**.

(99) Zheng, S. J.; Yu, R. A.; Liu, X. L.; Zhou, J. C. Research progress on mechanisms of absorption and transportation of selenium. *Chin Occup Med.* **2013**, 40, 456-460. **(In Chinese).**

(100) Cheng, M. W. Dietary patterns and their association with metabolic ayndrome among adults in China (2006-2009). *National Institute for Nutririon and Health, Chinese Center for Diseases Control and Prevention.* **2017**. **(In Chinese).**

(101) Zhang, J. G. Changes in dietary patterns and their association with general and central

obesity among adults in China (1991-2009). *Chinese Center for Disease Control and Prevention.* **2013**. **(In Chinese).**

(102) Filippini, T.; Cilloni, S.; Malavolti, M.; Violi, F.; Malagoli, C.; Tesauro, M.; Bottecchi, I.; Ferrari, A.; Vescovi, L.; Vinceti, M. Dietary intake of cadmium, chromium, copper, manganese, selenium and zinc in a Northern Italy community. *J. Trace Elem. Med. Bio.* **2018**, 50, 508-517.

(103) Rivas, A.; Romero, A.; Mariscal-Arcas, M.; Monteagudo, C.; Lopez, G.; Lorenzo, M. L.; Ocana-Peinado, F. M.; Olea-Serrano, F. Association between dietary antioxidant quality score (DAQs) and bone mineral density in Spanish women. *Nutr Hosp.* **2012**, 27, 1886-93.

(104) Gao, J.; Liu, Y.; Huang, Y.; Lin, Z. Q.; Banuelos, G. S.; Lam, H. W.; Yin, X. Daily selenium intake in a moderate selenium deficiency area of Suzhou. China. *Food Chem.* **2011**, 126, 1088-1093.

(105) Organization, G. W. H. Vitamin and mineral requirements in human nutrition. *Geneva World Health Organization.* **2005**.

(106) Ren, F.; Guo, X. Characteristics, toxicity and present situation of selenium in some countries. *Foreign Medical Sciences (Section of Medgeography).* **2004**, 3, 117-119. **(In Chinese).**

(107) WS/T 578.3-2017. Chinese dietary reference intakes-Part 3: Trace element. **(In Chinese).**

(108) Graham, R.; Senadhira, D.; Beebe, S.; Iglesias, C.; Monasterio, I. Breeding for micronutrient density in edible portions of staple food crops: conventional approaches. *Field Crop. Res.* **1999**, 60, 57-80.

(109) Sun, F. Y.; Chen, X.; Ma, Q. C.; Li, L., Wu, H. Y.; Li, T. Evaluation on Selenium

Biofortification Potential of Commercial Wheat Cultivars in Middle and Lower Reaches of the

Yangtze River. *J Agr Sci Tech-iran.* **2018**, 20, 83-90. **(In Chinese).**

**Figure 1** Comparison of Se content in wheat grain (A), flour (B) and food (C)

**Note**: Data originated from 66 studies; FSF: foliar Se fertilizer treatment, SSF: Soil Se

fertilizer treatment; CK: unfortified Se treatment; The year is based on the harvest year of

wheat; Data are presented as mean ± 95% CIs, with n in parentheses. Asterisks indicate

significant differences (*p < 0.05, \*\*p < 0.01, and \*\*\*p < 0.001)*; Different letters indicate

significant differences between mean effect size for groups within each category.

**Figure 2** Selenium content of wheat among different continents

**Note**: Data originated from 42 studies; The letters a, b, c and d indicate a significant

difference at the level of $P < 0.05$; n is the total number of samples.

**Figure 3** Relationship between soil selenium content and grain selenium content

**Note**: Data originated from 21 studies; n is the total number of samples.

**Figure 4** Relationship between foliar fertilization application rate and grain selenium content

(A) and increase times of grain selenium content after fortification in different countries (B)

**Note**: (A) Data originated from 21 studies; Selenium was added in the form of $Na_2SO_3$

solution; n is the total number of samples; (B) Data originated from 41 studies.

695    **Figure 5** Selenium content in different wheat derived foods

696        **Note**: Data originated from 26 studies; The letters a and b indicate a significant difference

697    at the level of $P < 0.05$; n is the total number of samples.

699    **Figure 6** The change of selenium from grain to human body and the appropriate level of

700    selenium fortification

702

703                              TOC graphic

**:Foliar Se fertilizer**

**:Soil Se**

Flour

Bread

Biscuits

Noodles

Pasta

Selenium content ($\mu g \cdot kg^{-1}$)

**(A)**

Y=22.20587X
R²=0.99975
n=188

Grain Selenium growth rate (%)

Foliar selenium application rate (g·hm⁻²)

**(B)**

| Country | Increase times of grain selenium content (Times) |
|---|---|
| Italy | 25.98 |
| South Africa | 11.53 |
| Portugal | 11.41 |
| Slovakia | 11.39 |
| Norway | 9.47 |
| China | 8.41 |
| Spain | 8.28 |
| Turkey | 5.80 |
| Pakistan | 2.42 |
| India | 2.16 |