Single cell analysis reveals X chromosome upregulation is not global and primarily belongs to ancestral genes in pre-gastrulation embryos

Naik C H, Chandel D, and Gayen S*

Department of Molecular Reproduction, Development and Genetics, Indian Institute of Science, Bangalore-560012, India.

*Correspondence: srimonta@iisc.ac.in

Abstract

Recent studies have provided substantial evidence supporting Ohno's hypothesis that upregulation of active X chromosome genes balances the dosage of X-linked gene expression relative to autosomal genes. However, the dynamics of X-chromosome upregulation (XCU)

during early development remain poorly understood. Here, we have profiled the dynamics of XCU in different lineages of female pre-gastrulation mouse embryos at single cell level through allele-specific single cell RNA-seq analysis. We found dynamic XCU upon initiation of random X-chromosome inactivation (XCI) in epiblast cells and cells of extraembryonic lineages, which undergo imprinted XCI, also harbored upregulated active-X chromosome. On the other hand, the extent of XCU remains controversial till date. While it is thought that it is a global phenomenon, some studies suggested that it affects dosage sensitive genes only. Interestingly, through profiling gene-wise dynamics of XCU, we found that X-upregulation is not global and primarily belongs to ancestral X-linked genes. However, it is not fully restricted to ancestral X-linked genes as some of newly acquired X-linked genes also undergo XCU, suggesting evolution of XCU to the newly acquired genes as well. Importantly, we found that occupancy of RNApolII, H3k4me3, H4k16ac and H3K36me3 is enhanced at loci of the upregulated/ancestral genes compared to non-upregulated/newly acquired X-linked genes. Moreover, upregulated X-linked genes showed increased burst frequency compared to the non-upregulated genes. Altogether, our study provides significant insight into the gene-wise dynamics, mechanistic basis, and evolution of XCU during early development.

Keywords: X chromosome upregulation, X chromosome inactivation, Sex chromosome evolution, Dosage compensation, Y degeneration, Pre-gastrulation.

Introduction

In therian mammals, sex is determined by the sex chromosomes: XX in female and XY in male. During the evolution of sex chromosomes from a pair of autosomes, the Y chromosome's degradation led to the dosage imbalance between X and autosomal genes in males [1]. In 1967, Ohno hypothesized that the expression of X-linked genes in XY cells was upregulated to two-fold to correct this imbalance [2]. Subsequently, this X-chromosome upregulation (XCU) was inherited in females and thereby introduced an extra dosage of the X chromosome in female cells. Therefore, to counteract the extra dosage from X-chromosome in female cells, the evolution of X-chromosome inactivation (XCI) happened, a process that silent one of the X-chromosome in female mammals [3]. However, Ohno's hypothesis was not accepted well for a long time due to the lack of proper experimental evidence supporting the existence of upregulated active-X chromosome. The first evidence of X-upregulation came through the studies based on microarray analysis; however, subsequently, it was challenged through RNA-seq based analysis [4–9]. Down the line, several independent studies came out both in supporting and refuting Ohno's hypothesis [10–19]. Although, Ohno's hypothesis is still contested, recent studies by us and others have shown the presence of upregulated active-X chromosome (X_{2a}) in different cell types *in vivo* and *in vitro* [20]. However, the dynamics of X-chromosome upregulation (XCU) during early development remain poorly understood. Here, we have profiled the dynamics of XCU in different lineages of female pre-gastrulation mouse embryos at single cell level through allele-specific single cell RNA-seq analysis.

On the other hand, the extent of XCU, i.e., whether XCU is chromosome wide like XCI or restricted to specific genes, remains controversial. Many studies implicated that XCU might be global, though direct evidence showing gene-wise dynamics of upregulation is not well understood [11,14]. In contrary, several studies implicated that XCU affects dosage-sensitive genes such as components of macromolecular complexes, signal transduction pathways or encoding for transcription factors [21,22]. To get better insight about this, here we have explored gene-wise dynamics of XCU in pre-gastrulation mouse embryos. We found that XCU is neither global nor restricted to dosage sensitive genes only. Furthermore, we have investigated if XCU is related to the evolutionary timeline of X-linked genes. Based on the evolutionary time point X-linked genes can be categorized into two classes: ancestral X-linked genes, which were present originally from the beginning of sex chromosome evolution and newly acquired X-linked genes. Whether X-chromosome upregulation happens to the newly acquired X-linked genes or restricted to mostly in ancestral genes is another intriguing question. Here, we have analyzed the XCU kinetics in ancestral vs. newly acquire X-linked genes to understand the evolution of XCU. Finally, we have explored the mechanistic basis of XCU.

Results

Dynamic active-X upregulation upon random XCI in epiblast cells of pre-gastrulation embryos

In pre-gastrulation mouse embryos, while extraembryonic cells have imprinted inactive X, epiblast cells are at the onset of random XCI [23–27]. Therefore, to investigate the dynamics of XCU, we segregated the cells of E5.5, E6.25, and E6.5 mouse embryos into the three lineages: epiblast (EPI), extraembryonic ectoderm (ExE), and visceral endoderm (VE) based on lineage-specific marker gene expression by t-distributed stochastic neighbour embedding (t-SNE) analysis as described earlier (Fig. 1A). We investigated status of XCU in these

Figure 1: Dynamic X-chromosome upregulation in different lineages of pre-gastrulation embryos. (A)
Schematic outline of the workflow: profiling of active X-upregulation in different lineages (EPI: Epiblast, ExE:
Extraembryonic ectoderm, and VE: Visceral endoderm) of pre-gastrulation hybrid mouse embryos (E5.5, E6.25,
and E6.50) at the single-cell level using published scRNA-seq dataset. Hybrid mouse embryos were obtained from
crossing between two divergent mouse strains C57 and CAST. (B) Classification of cells based on XCI state
through profiling of fraction maternal expression of X-linked genes in the single cells of different lineages of pre-
gastrulation embryos (EPI, ExE, and VE). Ranges of fraction of maternal expression was considered for different
category cells: X_aX_a = 0.4-0.6, X_aX_p= 0.6-0.9/ 0.1-0.4 and X_aX_i = 0.9-1/0-0.1. (C) X:A ratios represented as violin

plots for the XaXa, XaXp, XaXi cells of different lineages of pre-gastrulation embryos. (D) Allelic expression
levels of X-linked and autosomal genes in XaXa, XaXp, XaXi cells of different lineages of pre-gastrulation
embryos.

different lineages of pre-gastrulation embryos through performing allelic/non-allelic gene
expression analysis using the available scRNA-seq dataset of E5.5, E6.25, and E6.5 hybrid
mouse embryos [28] (Fig. 1A). These embryos were derived from two divergent mouse strains
(C57BL/6J and CAST/EiJ) and therefore harbored polymorphic sites across the genome,
allowing us to profile gene expression with allelic resolution (Fig. 1A). First, we categorized
the cells of different lineages of the embryos based on their XCI status: cells with no XCI
(X_aX_a), partial or undergoing XCI (X_aX_p), and complete XCI (X_aX_i) through profiling the
fraction of maternal allele expression (Fig. 1B). We found that in the EPI lineage lot of cells
belongs to X_aX_p/X_aX_a indicating these cells are onset of random X-inactivation, whereas cells
of VE/ExE lineage mostly belong to X_aX_i category since these cells underwent of imprinted
X-inactivation (Fig. 1B). As expected, autosomal genes showed almost equivalent
paternal/maternal allele expression, thus validating our allele-specific analysis (Fig S1A).
Next, to check the upregulation dynamics, we profiled X:A ratio in the individual cells of
different stages/lineages of embryos. If a diploid female cell (X_aX_i) has upregulated active X,
the X:A ratio should be more than 0.5 and closer to 1. We found that X:A ratio of X_aX_p/X_aX_i
cells is always > 0.5 and close to 1 despite XCI, indicating the presence of dynamic X-
upregulation from the active X chromosome in X_aX_p/X_aX_i cells (Fig. 1C). For X:A analysis,
we made sure that there are no significant differences between the X-linked and autosomal
gene expression distribution (Method; Supplementary file 1). Next, we compared the allelic
expression pattern from autosome and X-chromosome in individual cells of each lineage. As
expected, we found that the active X expression is always significantly higher than the
autosomal allelic expression in X_aX_p/X_aX_i cells, corroborating upregulation of gene expression
from the active X -chromosome (Fig. 1D). On the other hand, there were no significant
differences in active Xs and autosomal allelic expression in X_aX_a cells of epiblast lineage
suggesting no upregulation of X-chromosome in absence of XCI (Fig. 1D). Altogether, these
analyses suggested dynamic active XCU upon initiation of random XCI in epiblast cells of pre-
gastrulation embryos.

133 X-chromosome upregulation is primarily confined to the ancestral X-linked genes

Our X:A analysis in pre-gastrulation embryos and differentiation mESCs revealed that the X:A
ratio of XaXi cells is always lower compared to the XaXa cells (Fig. 1C). This data hinted that
X-upregulation of X-linked genes in XaXi cells is partial or all genes do not undergo
upregulation. To explore this further, we investigated whether X-chromosome upregulation
occurs globally or restricted to certain class of genes. To test this, we profiled gene-wise
dynamics of XCU by comparing the expression of X-linked genes from the active X
chromosome of X_aX_i cells with the same active allele of X_aX_a cells in EPI E6.5 (Fig. 2A). If
active allele of a gene is upregulated in X_aX_i cells, it will show increased expression from the
active allele in X_aX_i cells compared to the same active allele of X_aX_a cells. We found that while
many X-linked genes showed increased expression from the active-X allele in X_aX_i cells
compared to the X_aX_a, there were significant number of genes which did not show such
increased expression suggesting that all genes do not undergo upregulation (Fig. 2A).
Moreover, while some genes showed robust upregulation from the active-X allele, the others
were moderately upregulated. Altogether, this result suggested that X-upregulation is not

Figure 2: X-chromosome upregulation is not global. (A) Comparison of expression (log normalized reads) of individual X-linked genes from the active allele between the XaXa and XaXi EPI cells of E6.50 embryos (B) Plot representing intersections of allele-wise upregulation of X-linked genes in E6.5 EPI cells.

global or do not occur chromosome-wide. On the other hand, surprisingly we found that an adequate number of upregulated X-linked genes showed allele-specificity as they showed upregulation from either C57 or CAST as an active-X suggesting regulation of upregulation of active allele may occur in parent of origin specific manner (Fig. 2B).

Next, we investigated the mechanistic basis of why some genes undergo upregulation and others are not. To find this, we categorized the X-linked genes in E6.5 EPI to three categories based on their active X-upregulation status. X-linked genes with robust upregulation from both C57 and CAST allele (Category 1), robust upregulation from one allele but moderate from the other allele (Category 2) and no upregulation from either of the allele (Category3) (Fig. 3A). First, we explored if there any relevance of the location of the genes on the X-chromosome with the upregulation. To test this, we profiled the position of the upregulated (Cat1 and Cat 2) and non-upregulated genes (Cat 3). We found that all these three categories' genes are located across the X-chromosome and there is no specific pattern of position of upregulated vs non-upregulated genes indicating chromosomal positioning of X-linked genes is not relevant to X-upregulation (Fig 3B). Next, we asked if there any differences of expression level of these three classes of genes and found no significant differences (Fig. 3C). Moreover, the X-linked genes in the three categories did not show any distinction with respect to their location in topologically associated domains (TADs) (Fig. 3D). As the retrotransposon elements play a profound role in gene regulation, we wanted to see if the genes undergoing upregulation from active-X chromosome has any link to the retrotransposon enrichment. However, the three categories of X-linked genes showed no correlation with the density of LINE elements (Fig. 3E). SINEs on the other hand, were significantly enriched in not upregulated (Cat3) genes compared to the robustly upregulated (Cat1) genes (Fig. 3E). Previous reports stated that XCU is restricted only to dosage-sensitive genes such as genes encoding for large protein complexes, transcription factors, proteins involve in signal transduction etc. Therefore, we analyzed if the upregulated genes in EPI E6.5 is mostly belonging such dosage sensitive genes or not. However, we found that while there are some upregulated genes fall into dosage sensitive category, many genes are not (Supplementary file 2). Moreover, there are many dosage sensitive genes among the non-upregulated genes. Altogether, there are no significant differences of distribution of dosage sensitive / insensitive genes among the upregulated or non-upregulated genes suggesting XCU is not restricted to dosage sensitive genes only (Supplementary file 2). Next, we collated the three categories of X-linked genes with their one-to-one orthologs in chicken (ancestral X-linked genes) and found that most of the robustly upregulated (Cat1) and moderately upregulated (Cat2) genes belong to this ancestral class of X-linked genes (79% in Cat1 and 60% in Cat2) and only a few belong to the class of acquired X-linked genes. While the genes that showed no upregulation had equivalent distribution to two classes of ancestral and acquired X-linked genes (Fig. 3F). Altogether, these data suggest that the process of active-X upregulation is mostly confined to the ancestral class of X-linked genes (Fig. 3F).

Upregulated / ancestral X-linked genes show higher enrichment of active chromatin marks

Previous studies have reported that the X linked genes from active X-chromosome have greater enrichment of the active chromatin marks compared to the autosomal genes [14]. Therefore, we investigated if the different active chromatin marks (H3K4me3, H3K36me3, H4K16ac and RNA PolII) are differentially enriched at the transcriptional start site and gene body of upregulated (Cat1 and Cat2) vs not upregulated (Cat3) genes on the active X-chromosome (Fig.

Figure 3: X-upregulation is mostly confined to ancestral X-linked genes. (A) Shematic representation of different categories of X-linked genes in E6.5 EPI based on their allele-wise upregulation status. X-linked genes with robust upregulation from both C57 and CAST allele (Category 1), robust upregulation from one allele but moderate from the other allele (Category 2) and no upregulation from either of the allele (Category3). (B) Genomic position of the three categories of X-linked genes on the X-chromosome. (C) Comparison of expression

levels (log2 TPM) between the three categories of X-linked genes. (D) Histograms representing the percent of the
three categories of X-linked genes located in TADs. (E) Density of LINE and SINE repeats in the three categories
of X-linked genes. p-values of Wilcoxon rank test comparing the retrotransposon enrichment in the three
categories (*p-value < 0.05; ***) (F) Heatmap representing the segregation of the three categories of X-linked genes
to ancestral and acquired classes of X-linked genes in E6.5 EPI cells.

4A). Interestingly, we found that the average density of H3K4me3, H4K16ac and RNA PolII
showed notable increase in the upregulated X-linked genes (Cat1 and Cat2) compared to the
not upregulated genes (Cat3) both around the TSS and gene body regions (Fig. 4A). However,
for H3K36me3 there were slight difference in enrichment. Additionally, we compared the
enrichment of these active marks between ancestral vs acquired X-linked genes on the active
X-chromosome and found that the ancestral class of X-linked genes showed higher enrichment
for these marks compared to the newly acquired X-linked genes (Fig. 4B). Finally, we profiled
the allelic transcriptional burst kinetics between the upregulated and not upregulated X-linked
genes in the X_aX_i cells of E6.5 EPI. Excitingly, the upregulated X-linked genes (Cat1 and Cat2)
showed significantly higher burst frequency compared to the not upregulated X-linked genes
(Cat3) (Fig. 4C). However, burst sizes were not significantly different between the two
categories of X-linked genes (Fig. 4C).

Discussion

In 1967, Ohno hypothesized that gene loss from the Y chromosome led to the upregulation of
X-chromosomes in both sexes, which subsequently led to the inactivation of one X-
chromosome in female mammals to ensure proper X-dosage. However, Ohno's hypothesis was
contested for a long time due to the lack of appropriate experimental evidence. Recent studies
have provided evidence supporting active X upregulation in different cell types *in vivo* and *in*
vitro. However, the dynamics of X-upregulation during early development remain poorly
understood. Here, we have profiled the XCU dynamics at single cell level in female pre-
gastrulation mouse embryos. We found that in EPI cells (E5.5, E6.25, E6.5), which are at the
onset of random XCI, X upregulation is dynamically linked with the X-inactivation. VE and
ExE of pre-gastrulation embryos, which undergo imprinted X-inactivation also showed
presence of upregulated active-X chromosome. Altogether, these results suggested that two X-
chromosomes expression state is highly plastic in nature towards balancing the optimal X
chromosome dosage during early development.

One intriguing question is whether X-upregulation occurs globally or restricted to certain class
of genes. It is thought that upon X-chromosome inactivation, numerous trans-acting factors
leave the inactivating-X and shift to the active-X thereby leading to global increase in
expression from the active-X chromosome. In contrast, it is thought that XCU is restricted to
dosage sensitive genes only. Interestingly, we found that though XCU is dynamically linked to
XCI, XCU is not global like XCI. Indeed, we found that the X:A ratio of XaXi cells is always
lower compared to the XaXa cells. Importantly, through assaying gene-wise dynamics of XCU
we found that many X-linked genes do not undergo upregulation. Moreover, we found that
upregulated genes are not restricted to dosage sensitive genes. Altogether, these results
suggested that XCU is neither global nor restricted to dosage sensitive genes only.
Interestingly, we found that XCU is mostly confined to the ancestral class of X-linked genes.
However, it is not restricted to ancestral X-linked gene only as some of newly acquired X-
linked genes also undergo X-upregulation, suggesting evolution of X-upregulation to the newly
acquired genes as well.

Figure 4: Upregulated X-linked genes show higher enrichment of different active chromatin marks and higher burst frequency compared to non-upregulated genes. Density plots showing the enrichment of RNA

PolII S5P, active chromatin marks H3K4me3, H3K36me3 and H4K16ac in the (A) upregulated vs not upregulated
X-linked genes and (B) ancestral vs acquired classes of X-linked genes on the active-X chromosome. (C)
Comparison of allelic burst frequency and burst size of the upregulated vs not upregulated X-linked genes in the
E6.5 EPI cells (Wilcoxon rank test: *p-value < 0.01; *** and *p-value < 0.001; ****). (D) Model representing that
X-chromosome upregulation is mostly confined to the ancestral X-linked genes and mechanism of XCU.

On the other hand, mechanism of XCU, remain poorly understood. We found that the active
histone marks such as H3K4me3, H3K36me3, H4K16ac as well as RNA PolII are highly
enriched on the genes subjected to X-upregulation compared to the not-upregulated genes (Fig.
4A). Furthermore, these marks are seen to have higher average density on the ancestral X-
linked genes compared to the newly acquired class of X-linked genes (Fig. 4B). Interestingly,
we found that the transcriptional burst frequencies are significantly enhanced in upregulated
genes compared to the non-upregulated genes. On the other hand, transcriptional burst sizes
remained almost similar between the two categories of X-linked genes (Fig. 4C). Taken
together, our results unveils that the distinctive enrichment of active chromatin marks and
higher transcriptional burst frequencies might be one of the major regulators for XCU (Fig.
4D).

In summary, our study confers significant support to Ohno's hypothesis. In addition, we show
that although the two X-chromosomes' expression state is highly plastic in nature towards
balancing the optimal X chromosome dosage during development, XCU is not global like XCI.
Importantly, we found that XCU is primarily confined to ancestral X-linked genes only.

Methods

Data acquisition: Pre-gastrulation embryo single-cell RNA-seq dataset for E5.5 and E6.25
generated from hybrid mouse embryos (C57BL/6J × CAST/EiJ) and E6.5 from hybrid mouse
with the reciprocal cross (CAST/EiJ × C57BL/6J) was retrieved under the accession code-
GSE109071 [28].

Read alignment and counting: RNA-seq reads were aligned to the mouse reference genome
mm10 using STAR [29]. The reads aligned were then counted using HTSeq-count. To avoid
the dropout events due to low amount of mRNA sequenced within single cells, we used a
statistical imputation method scImpute [30], which identifies the likely dropouts without
introducing any bias in the rest of the data. Expression levels of transcripts was computed using
Transcripts per million (TPM) method.

Sexing of the embryos: For sex-determination of the pre-gastrulation embryos, an embryo was
classified as male if the sum of the read count for the Y-linked genes (Usp9y, Uty, Ddx3y,
Eif2s3y, Kdm5d, Ube1y1, Zfy2, Zfy1) in each cell of an embryo was greater than 12, rest were
considered as female embryos.

X:A ratio: We calculated the X:A ratio for different lineages of pre-gastrulation embryo by
dividing the median expression (TPM) of X-linked genes with the median expression (TPM)
of the autosomal genes. For this analysis, we considered those X-linked and autosomal genes
having ≥ 0.5 TPM. We also applied an upper TPM threshold which corresponded to the lowest
90 th centile of TPM expression to avoid any differences between the X-linked and autosomal
gene expression distribution. Also, statistical Kolmogorov-Smirnov's test was performed
which again validated the non-significant differences in the levels of gene expression

distribution between X and autosomal genes. We excluded the escapees of X-inactivation and the genes in the pseudo autosomal region from our analysis.

Allele-specific expression analysis: We first constructed in silico CAST specific parental genome by incorporating CAST/EiJ specific SNPs (https://www.sanger.ac.uk/science/data/mouse-genomes-project) into the mm10 genome using VCF tools. Reads were mapped onto C57BL/6J (mm10) reference genome and in silico CAST genomes using STAR allowing no multi-mapped reads. To exclude any false positives, we only considered those genes with at least 2 informative SNPs and minimum 3 reads per SNP site. We took an average of SNP-wise reads to have the allelic read counts. After normalization of allelic read counts, we considered only those genes for downstream analysis which had at least 1 average reads across the cells of each lineage from a specific developmental stage for pre-gastrulation embryos. Further, only those single-cells were considered for downstream analysis which showed at least 10 X-linked gene expressions. Allelic ratio was obtained individually for each gene using formula = (Maternal/Paternal reads) ÷ (Maternal reads + Paternal reads).

Classification of ancestral and acquired X-linked genes: The ancestral X-linked genes are identified as those having one-to-one orthologs in chicken and were retrieved from the Ensembl biomart release 102. All the remaining X-linked genes are considered as the acquired class of X-linked genes.

Topologically associated domains (TADs), LINE and SINE analysis: The coordinates of TADs called in mouse ES cells by GMAP method were retrieved from the TADKB database for our analysis [31]. For LINE and SINE element analysis, we used the table containing the genomic coordinates of repeat elements from the UCSC table browser, the remasker table ("rmsk") for mm10 reference mouse genome assembly. The density of LINE and SINE elements was then calculated for a gene as the ratio of the total length of the retrotransposons present to the length of that gene.

Transcriptional burst kinetics: We used SCALE to profile allelic transcriptional burst kinetics [32]. For this analysis, we used genes with at least 5 average read counts across the cells of EPI E6.5.

ChIP-seq analysis: To estimate the enrichment of H3K4me3, H3K36me3, RNA PolII S5P, and H4K16ac in our gene categories, we retrieved the available ChIP-seq datasets for MEF cells from GSE33823 [33] , GSE36905 [34] for H3K4me3, H3K36me3, RNAPolII S5P and GSE97459 [35] for H4K16ac.

Gene function analysis: We identified association of the genes with different biological function through Gene Ontology (Biological process, Molecular Function and Cellular Component, Signal related function (http://geneontology.org/) [36], Transcription factor databases (http://bioinfo.life.hust.edu.cn/AnimalTFDB/#!/ , https://sunlab.cpy.cuhk.edu.hk/mTFkb/) [37,38], involved in protein complex (http://mips.helmholtz-muenchen.de/corum/#) [39], Dosage sensitive genes (human) (https://www.clinicalgenome.org/) [40], Housekeeping genes database (https://housekeeping.unicamp.br/, Li et al., 2017) [41,42].

Statistical tests and plots: All statistical tests and plots were performed in R version 3.6.3.

Author's Contribution

SG conceptualized and supervised the study. HCN and DC did bioinformatic analyses. SG, DC, and HCN wrote, edited, and proofread the manuscript. The final manuscript was edited and approved by all the authors.

Acknowledgments

Study is supported by DBT grant (BT/PR30399/BRB/10/1746/2018), DST-SERB (CRG/2019/003067), DBT-Ramalingaswamy fellowship (BT/RLF/Re-entry/05/2016) and Infosys Young Investigator award to SG. We also thank DST-FIST [SR/FST/LS11-036/2014(C)], UGC-SAP [F.4.13/2018/DRS-III (SAP-II)] and DBT-IISc Partnership Program Phase-II (BT/PR27952-INF/22/212/2018) for infrastructure and financial support.

References:

1. Graves JAM. Evolution of vertebrate sex chromosomes and dosage compensation. Nature Reviews Genetics. 2016. pp. 33–46. doi:10.1038/nrg.2015.2

2. Ohno S. Sex Chromosomes and Sex-linked Genes. Springer-Verlag, Berlin, Heidelberg, New York. 1967;68: 1375. doi:10.7326/0003-4819-68-6-1375_2

3. Lyon MF. Gene action in the X-chromosome of the mouse (mus musculus L.). Nature. 1961;190: 372–373. doi:10.1038/190372a0

4. Di KN, Disteche CM. Dosage compensation of the active X chromosome in mammals. Nat Genet. 2006;38: 47–53. doi:10.1038/ng1705

5. Gupta V, Parisi M, Sturgill D, Nuttall R, Doctolero M, Dudko OK, et al. Global analysis of X-chromosome dosage compensation. J Biol. 2006;5. doi:10.1186/jbiol30

6. Lin H, Gupta V, Vermilyea MD, Falciani F, Lee JT, O'Neill LP, et al. Dosage compensation in the mouse balances up-regulation and silencing of X-linked genes. PLoS Biol. 2007;5: 2809–2820. doi:10.1371/journal.pbio.0050326

7. Xiong Y, Chen X, Chen Z, Wang X, Shi S, Wang X, et al. RNA sequencing shows no dosage compensation of the active X-chromosome. Nat Genet. 2010;42: 1043–1047. doi:10.1038/ng.711

8. Johnston CM, Lovell FL, Leongamornlert DA, Stranger BE, Dermitzakis ET, Ross MT. Large-scale population study of human cell lines indicates that dosage compensation is virtually complete. PLoS Genet. 2008;4: 0088–0098. doi:10.1371/journal.pgen.0040009

9. Talebizadeh Z, Simon SD, Butler MG. X chromosome gene expression in human tissues: Male and female comparisons. Genomics. 2006;88: 675–681. doi:10.1016/j.ygeno.2006.07.016

10. Larsson AJM, Coucoravas C, Sandberg R, Reinius B. X-chromosome upregulation is driven by increased burst frequency. Nat Struct Mol Biol. 2019;26: 963–969. doi:10.1038/s41594-019-0306-y

11. Deng X, Hiatt JB, Nguyen DK, Ercan S, Sturgill D, Hillier LW, et al. Evidence for

compensatory upregulation of expressed X-linked genes in mammals, Caenorhabditis
elegans and Drosophila melanogaster. Nat Genet. 2011;43: 1179–1185.
doi:10.1038/ng.948

12. Sangrithi MN, Royo H, Mahadevaiah SK, Ojarikre O, Bhaw L, Sesay A, et al. Non-
Canonical and Sexually Dimorphic X Dosage Compensation States in the Mouse and
Human Germline. Dev Cell. 2017;40: 289-301.e3. doi:10.1016/j.devcel.2016.12.023

13. Li X, Hu Z, Yu X, Zhang C, Ma B, He L, et al. Dosage compensation in the process of
inactivation/reactivation during both germ cell development and early embryogenesis
in mouse. Sci Rep. 2017;7. doi:10.1038/s41598-017-03829-z

14. Deng X, Berletch JB, Ma W, Nguyen DK, Hiatt JB, Noble WS, et al. Mammalian X
upregulation is associated with enhanced transcription initiation, RNA half-life, and
MOF-mediated H4K16 acetylation. Dev Cell. 2013;25: 55–68.
doi:10.1016/j.devcel.2013.01.028

15. Lin H, Halsall JA, Antczak P, O'Neill LP, Falciani F, Turner BM. Relative
overexpression of X-linked genes in mouse embryonic stem cells is consistent with
Ohno's hypothesis. Nat Genet. 2011; doi:10.1038/ng.992

16. De Mello JCM, Fernandes GR, Vibranovski MD, Pereira L V. Early X chromosome
inactivation during human preimplantation development revealed by single-cell RNA-
sequencing. Sci Rep. Springer US; 2017;7: 1–12. doi:10.1038/s41598-017-11044-z

17. Kharchenko P V., Xi R, Park PJ. Evidence for dosage compensation between the X
chromosome and autosomes in mammals. Nature Genetics. 2011. pp. 1167–1169.
doi:10.1038/ng.991

18. Chen X, Zhang J. The X to autosome expression ratio in haploid and diploid human
embryonic stem cells. Mol Biol Evol. 2016;33: 3104–3107.
doi:10.1093/molbev/msw187

19. Julien P, Brawand D, Soumillon M, Necsulea A, Liechti A, Schütz F, et al.
Mechanisms and evolutionary patterns of mammalian and avian dosage compensation.
PLoS Biol. 2012; doi:10.1371/journal.pbio.1001328

20. Mandal S, Chandel D, Kaur H, Majumdar S, Arava M, Gayen S. Single-Cell Analysis
Reveals Partial Reactivation of X Chromosome instead of Chromosome-wide
Dampening in Naive Human Pluripotent Stem Cells. Stem Cell Reports. 2020;14:
745–754. doi:10.1016/j.stemcr.2020.03.027

21. Pessia E, Makino T, Bailly-Bechet M, McLysaght A, Marais GAB. Mammalian X
chromosome inactivation evolved as a dosage-compensation mechanism for dosage-
sensitive genes on the X chromosome. Proc Natl Acad Sci U S A. 2012;109: 5346–
5351. doi:10.1073/pnas.1116763109

22. Pessia E, Engelstädter J, Marais GAB. The evolution of X chromosome inactivation in
mammals: The demise of Ohno's hypothesis? Cellular and Molecular Life Sciences.
2014. pp. 1383–1394. doi:10.1007/s00018-013-1499-6

23. Gayen S, Maclary E, Buttigieg E, Hinten M, Kalantry S. A Primary Role for the Tsix
lncRNA in Maintaining Random X-Chromosome Inactivation. Cell Rep. 2015;11:
1251–1265. doi:10.1016/j.celrep.2015.04.039

24.   Harris C, Cloutier M, Trotter M, Hinten M, Gayen S, Du Z, et al. Conversion of
426   random x-inactivation to imprinted x-inactivation by maternal prc2. Elife. 2019;8.
427   doi:10.7554/eLife.44258

428   25.   Maclary E, Hinten M, Harris C, Sethuraman S, Gayen S, Kalantry S. PRC2 represses
429   transcribed genes on the imprinted inactive X chromosome in mice. Genome Biol.
430   2017;18. doi:10.1186/s13059-017-1211-5

431   26.   Gayen S, Maclary E, Hinten M, Kalantry S. Sex-specific silencing of X-linked genes
432   by Xist RNA. Proc Natl Acad Sci U S A. 2016;113. doi:10.1073/pnas.1515971113

433   27.   Sarkar MK, Gayen S, Kumar S, Maclary E, Buttigieg E, Hinten M, et al. An Xist-
434   activating antisense RNA required for X-chromosome inactivation. Nat Commun.
435   2015;6. doi:10.1038/ncomms9564

436   28.   Cheng S, Pei Y, He L, Peng G, Reinius B, Tam PPL, et al. Single-Cell RNA-Seq
437   Reveals Cellular Heterogeneity of Pluripotency Transition and X Chromosome
438   Dynamics during Early Mouse Development. Cell Rep. 2019;26: 2593--2607.e3.
439   doi:10.1016/j.celrep.2019.02.031

440   29.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
441   Ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.
442   doi:10.1093/bioinformatics/bts635

443   30.   Li WV, Li JJ. An accurate and robust imputation method scImpute for single-cell
444   RNA-seq data. Nat Commun. 2018;9. doi:10.1038/s41467-018-03405-7

445   31.   Liu T, Porter J, Zhao C, Zhu H, Wang N, Sun Z, et al. TADKB: Family classification
446   and a knowledge base of topologically associating domains. BMC Genomics 2019
447   201. BioMed Central; 2019;20: 1–17. doi:10.1186/S12864-019-5551-2

448   32.   Jiang Y, Zhang NR, Li M. SCALE: Modeling allele-specific gene expression by
449   single-cell RNA sequencing. Genome Biol. 2017;18. doi:10.1186/s13059-017-1200-8

450   33.   Yildirim E, Sadreyev RI, Pinter SF, Lee JT. X-chromosome hyperactivation in
451   mammals via nonlinear relationships between chromatin states and transcription. Nat
452   Struct Mol Biol. 2012;19: 56–62. doi:10.1038/nsmb.2195

453   34.   Pinter SF, Sadreyev RI, Yildirim E, Jeon Y, Ohsumi TK, Borowsky M, et al.
454   Spreading of X chromosome inactivation via a hierarchy of defined Polycomb stations.
455   Genome Res. Cold Spring Harbor Laboratory Press; 2012;22: 1864–1876.
456   doi:10.1101/GR.133751.111

457   35.   Ye A, Kim H, Kim J. PEG3 control on the mammalian MSL complex. 2017;
458   doi:10.1371/journal.pone.0178363

459   36.   H M, A M, D E, X H, PD T. PANTHER version 14: more genomes, a new PANTHER
460   GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res. Nucleic
461   Acids Res; 2019;47: D419–D426. doi:10.1093/NAR/GKY1038

462   37.   H H, YR M, LH J, QY Y, Q Z, AY G. AnimalTFDB 3.0: a comprehensive resource
463   for annotation and prediction of animal transcription factors. Nucleic Acids Res.
464   Nucleic Acids Res; 2019;47: D33–D38. doi:10.1093/NAR/GKY822

465   38.   Sun K, Wang H, Sun H. mTFkb: a knowledgebase for fundamental annotation of
466   mouse transcription factors. Sci Reports 2017 71. Nature Publishing Group; 2017;7: 1–

11. doi:10.1038/s41598-017-02404-w

39. M G, J R, B B, I D-K, G F, G F, et al. CORUM: the comprehensive resource of
mammalian protein complexes-2019. Nucleic Acids Res. Nucleic Acids Res; 2019;47:
D559–D563. doi:10.1093/NAR/GKY973

40. HL R, JS B, LD B, CD B, JP E, MJ L, et al. ClinGen--the Clinical Genome Resource.
N Engl J Med. N Engl J Med; 2015;372: 2235–2242. doi:10.1056/NEJMSR1406261

41. Li B, Qing T, Zhu J, Wen Z, Yu Y, Fukumura R, et al. A Comprehensive Mouse
Transcriptomic BodyMap across 17 Tissues by RNA-seq. Sci Reports 2017 71. Nature
Publishing Group; 2017;7: 1–10. doi:10.1038/s41598-017-04520-z

42. Hounkpe BW, Chenou F, de Lima F, De Paula EV. HRT Atlas v1.0 database:
redefining human and mouse housekeeping genes and candidate reference transcripts
by mining massive RNA-seq datasets. Nucleic Acids Res. Oxford Academic; 2021;49:
D947–D955. doi:10.1093/NAR/GKAA609