**Cardiomyocyte adenyl cyclase 8 overexpression increases locomotor activity in mice by modulating EEG-gamma oscillations**

*Jacopo Agrimi [a, b], Danilo Menicucci [c], Jia-Hua Qu [b], Marco Laurino [d], Chelsea D Mackey [b], Laila Hasnain [b], Yelena S Tarasova[b], Kirill V Tarasov[b], Ross A McDevitt [h], Donald B Hoover [e, g], Angelo Gemignani* [c], Nazareno Paolocci* [a, f], Edward G Lakatta* [b]*

a) *Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA*
b) *Laboratory of Cardiovascular Sciences, National Institute on Aging, National Institutes of Health Biomedical Research Center (BRC), Baltimore, MD, USA*
c) *Department of Surgical, Medical, Molecular Pathology, and Critical Care Medicine, University of Pisa, Pisa, Italy*
d) *Institute of Clinical Physiology, National Research Council, Pisa, Italy*
e) *Department of Biomedical Sciences, James H. Quillen College of Medicine, East Tennessee State University, Johnson City TN*
f) *Department of Biomedical Sciences, University of Padova, Padova, Italy*
g) *Center of Excellence in Inflammation, Infectious Disease and Immunity, East Tennessee State University, Johnson City, TN*
h) *The Comparative Medicine Section, National Institute on Aging, National Institutes of Health, Baltimore, MD, USA*

*\*Contributed equally*

**Keywords:** Adenylyl cyclase type 8, EEG Gamma Rhythm, Granger Causality, Locomotor Activity, Heart-Brain Communication, Sympathetic Stress

Correspondence to:

**Nazareno Paolocci, M.D., Ph.D.**

Division of Cardiology, Traylor 911
Johns Hopkins Hospital
720 Rutland Avenue
Baltimore, MD, 21212
npaoloc1@jhmi.edu

**Edward G. Lakatta, MD.**

Laboratory of Cardiovascular Sciences,
National Institute on Aging,
National Institutes of Health Biomedical Research Center (BRC),
Baltimore, MD, 21224
lakattae@grc.nia.nih.gov

**Abstract**

The central nervous system modulates heart function on a beat-to-beat basis via increasingly understood mechanisms. Conversely, whether and how humoral/functional cardiac variations shape brain activity and adaptive behavior remains unclear. This study shows that mice overexpressing adenylyl cyclase type 8 in myocytes (TGAC8), characterized by persistently elevated heart rate/contractility, also display increased locomotion. This effect is sustained by enhanced gamma rhythms, as evidenced by simultaneous behavioral and EEG/ECG monitoring. These changes are specific because they are not paralleled by other behavioral changes, such as heightened anxiety-like behavior. In unison, TGAC8 mice hippocampus exhibits upregulated GABA-A receptors, whose activation chiefly accounts for gamma activity generation. Moreover, the Granger causality analysis between ECG and EEG attests to the causal involvement of the autonomic component of the heartbeat in shaping EEG gamma oscillations in a bottom-up modality. Mechanistically, TGAC8 harbors elevated circulating dopamine/DOPA levels of cardiac origin and upregulated hippocampal D5 dopamine receptor levels. In synergy with the GABA-A receptor, D5 activation favors hippocampal inhibitory currents that drive EEG gamma oscillations. These studies, therefore, inform how heart-initiated functional and/or humoral modifications reverberate back to the brain to modulate specific primary behavioral responses, such as locomotion.

**Significance**

The brain is continuously aware of the functional status of many bodily organs, modulating, for instance, the heart's activity beat-by-beat. Conversely, the impact of cardiac activity modifications on brain function and behavior is less understood. We disclose that augmenting myocyte adenyl cyclase 8 (AC8) activity in mice increases heart function and locomotor activity. Elevated cardiac AC8 levels lead to higher circulating dopamine and DOPA, two hormones crucially involved in movement control, and increased expression of the hippocampus's GABA-A and D5 receptors; the latter are all critical for gamma activity generation. Thus, the brain interprets changes in cardiac AC8 activity as a "sustained exercise-like" situation and responds by activating areas that eventually command an increase of locomotion.

## Introduction

Damasio's somatic theory of emotions postulates that afferent somatic signals from the body's peripheral districts are integrated into subcortical and cortical regions, shaping emotional and behavioral responses needed to restore the body's homeostasis (1). The bidirectional relationship between the heart and the brain falls entirely within this loop. Indeed, besides the autonomic nervous system's descending control of cardiac function, information is also processed by the intrinsic cardiac nervous system (the so-called "little brain of the heart") that communicates back to the brain via ascending fibers located in the spinal cord and vagus nerve (2). These afferent impulses reach relay stations such as the medulla, hypothalamus, thalamus, and, ultimately, the cerebral cortex, carrying sensory information (2). Despite numerous observational clues supporting the possibility that changes in cardiac activity can alter behavior (3–5), definitive mechanistic evidence validating this view and putting forth pathways potentially accounting for these *bottom-up* phenomena remains to be gained.

Mice harboring a cardiac-specific overexpression of adenylyl cyclase (AC) type 8 (TGAC8) display a persistently elevated heart rate (HR), reduced HR variability (HRV), and increased contractility owing to enduringly elevated cardiac-intrinsic cAMP-PKA-Ca$^{2+}$ signaling (6, 7). Moreover, the heart of these transgenic mice escapes top-down autonomic surveillance by blocking beta-adrenergic signaling and catecholamine production to escape harmful additional sympathetic stress (6). Further, the endocrine portfolio of the TGAC8 heart showcases an accentuated dopamine anabolism mirrored by elevated circulating levels of dopamine and DOPA (6), and the latter can permeate the blood-brain barrier (8). With this tool and evidence in hand, in the present study, we sought to model Damasio's theory of the somatic marker in an experimental setting, proving that bioelectrical, mechanical, and endocrine signaling, borne in the periphery, can shape the activity of specific brain areas and consequently, adaptive behaviors.

## Results

 **Mice with cardiac overexpression of AC8 display increased locomotor activity**

***Figure 1.*** *TGAC8 mice display a general increase in locomotor behavior. (A) Plot of the animals movements during OF test, WT vs TGAC8 mouse; (B) Open Field test, total distance travelled; (C) Open Field test, average speed; (D) Plot of the animals movements during Elevated Plus Maze test, WT vs TGAC8 mouse; (E) Elevated Plus Maze test, number of entries in the open arms of the labyrinth; (F) Elevated Plus Maze test, time spent in the open arms of the labyrinth; (G) Elevated Plus Maze test, total entries in the open arms of the labyrinth; (H) Elevated Plus Maze test, total distance travelled; (I) Elevated Plus Maze test, average speed; (J) Elevated Plus Maze test, maximum speed; (K) Elevated Plus Maze test, total time freezing; (L) Elevated Plus Maze test, freezing episodes; (M) Fear Conditioning Test, time freezing in a contextual frame; (N) Fear Conditioning Test, time freezing in a cued frame . Results are shown as violins plot, N=7, unpaired t-test has been performed between WT and TGAC8 groups, *p<0.05, †p<0.01, ‡p<0.001.*

Whether chronically elevated heart rhythm and/or contractility impact fundamental brain
functions is currently unclear. Hence, we first set out to determine whether persistently elevated
cardiac chronotropy/inotropy emanating from AC8 overexpression would affect two main
behavioral variables, i.e., locomotion and exploratory behavior (i.e, anxiety/like behavior): two
major behaviors associated with heart rate modifications (9). When challenged with an open field
(OF) test, TGAC8 mice traveled more distance and at a higher speed than their age-matched
littermates (+43% and +38%, respectively, p<0.01) (Fig. 1 B, C). Next, we determined their
exploratory activity and anxiety-like behavior using the elevated plus maze (EPM) test. First, the
EPM test confirmed the ability of TGCA8 mice to cover a longer distance at a higher average speed
(Fig.1H, I), and with a maximum speed (Fig.1J) sizably higher than in the controls. Second, this test
revealed a considerable decrease in time and episodes of freezing behavior in the transgenic mice
(-45%, *p<0.05) (Fig. 1K, L). However, TGAC8 and WT did not differ in the two main parameters
evaluating anxiety-like behavior, i.e., the number and times of entry in the open arms (Fig. E, F),
results confirmed by the light-dark box test (Sup. Mat.). Further to unveil a possible anxious
behavior, we performed the fear conditioning test, which, in contrast to the EPM, cannot be biased
by concomitant alterations in locomotor activity (10). Thus, there was no sizable difference
between the WT and TGAC8 on the basis of these two main trial parameters, time freezing
contextual and cued (Fig. M, N). Hence, these findings indicate that TGAC8 mice - harboring
persistently elevated chronotropic/inotropic activity at the heart level – display a hyperkinetic
behavior without corollary anxiety.

**Cardiac AC8 transgenesis increases EEG-gamma frequencies, shifting the movement control**
**from theta to fast rhythms**

To gain initial mechanistic clues explaining TGAC8 hyperlocomotion, we implanted dual lead
telemetric devices into TGAC8 and aged-matched WT littermates. This approach enabled us to
monitor EEG and ECG activities simultaneously (time series), as well as actigraphy (Fig. 2A). The 24
hours recording of the activity confirmed that transgenic mice spent more time moving (active)
compared to control littermates (Fig. 2B). Then, we focused on the ECG traces recorded during
the mice's active state. Consistent with previous findings (5, 6), overexpressing AC8 only in cardiac
cells increased heart rate and contractility while rendering the heart unresponsive to autonomic

nerve system (ANS) surveillance (6). Here, however, we validated and expanded this evidence.

Indeed, we observed that TGAC8 mice persistently elevated chronotropic activity paired with a

marked drop in both time domains and high- and low-frequency domains of HR variability. The

**Figure 2.** *Double ECG and EEG recording, Heart Rate Variability is impaired, while EEG-gamma rhythms are upregulated in TGAC8 mice. (A) Schematic of F20-EET double implant for telemetry recording in mice; (B) Actigraphy, time mobile; (C-F) Heart Rate Variability parameters measured in vivo from telemetric ECG recording: (C) Heart Rate; (D) Low Frequency; (E) High Frequency; (F) Low Frequency/High Frequency ratio; (G) EEG frequencies bands (from δ to γ3) measured in vivo from telemetric EEG recording. Results are shown as violin plots, N=7, unpaired t-test has been performed between WT and TGAC8 groups, *p<0.05, †p<0.01, ‡p<0.001. (H) Linear regression model between activity (time mobile recorded by actigraphy) and θ power; (I) Linear regression model between activity (time mobile recorded by actigraphy) and γ power.*

latter was recorded by an ECG branch of the double telemetry implant (p<0.001*** TGAC8 vs. WT)

(Fig. 2D-F). Additionally, when examining the active portion (animals moving) of the 24-hrs EEG

traces, we discovered that the TGAC8 mice had an overall increase in gamma activity, i.e., γ1/2/3,

spanning from 30 *Hz* to 160 *Hz* (Fig. 2G).

Theta and gamma activities are crucial determinants of overall locomotor behavior in

species such as mice (11, 12). Therefore, we next correlated the active movement recorded by the

actigraphy with these bands using a linear regression model to determine the degree of

association between activity and EEG patterns. We found that the theta bands power (theta-2 and

theta-1) is nicely associated with the activity pattern of WT mice ($R^2$=0.8), as shown previously

(11). Surprisingly, however, the theta band did not influence locomotion in TGAC8 mice ($R^2 = 0$)

(Fig. 2H). Conversely, the three gamma bands power were reliably associated to active state

(moving animal) in TGAC8 mice ($R^2 = 0.7$) while maintaining a moderate fit for movements in WT

mice ($R^2 = 0.5$) (Fig. 2I).

In aggregate, these data suggest that cardiac-selective overexpression of AC8 can drive

an increase in myocardial performance and concurrently bring about changes in the EEG central

patterns, especially within the gamma frequency range. In turn, these modifications can alter the

mouse locomotor behavior. Indeed, the current data uncover a shift in the control of locomotion

of TGAC8 from the classical theta band to a more prominent role of gamma rhythms.

***The flow of information between the heart and brain is augmented in TGAC8***

*Figure 3.* TAC8 mice show a blunted response to sympathomimetic agents, while maintaining parasympathetic reactivity. (A) Representative picture and diagram of isolated atria preparation; (B) Change (in percentage) of heart rate after Isoproterenol administration, n=7-8, data are shown as violin plots, unpaired t-test between WT and TGAC8, ‡p<0.001; (C) Change (in percentage) of heart rate after carbachol administration, n=4-5, log dose-response curve.

Parallel to the descending control exerted by the ANS on cardiac function, the peripheral information generated within the heart is communicated to

the brain via the intrinsic cardiac nervous system through ascending autonomic fibers (2). We next
explored the capacity of the TGAC8 hearts to respond to autonomic input. More in detail, we
assessed the intrinsic response of the sinoatrial node (SAN) to external sympathetic and
parasympathetic stimuli in isolated atria driven by spontaneous SAN impulses (Fig. 3A). Despite a
substantial decreased isoproterenol response (Fig. 3B), the parasympathetic effects evoked by
carbachol administration were intact (Fig. 3C). Next, we evaluated the baroreflex *in vivo* via
phenylephrine administration. Again, both WTs and TGAC8s responded to the stimulation
similarly (Fig. 3D). Thus, we confirmed that, despite some alterations, the autonomic
communication between the heart and the brain is still in place in TGAC8 mice.

We next determined whether the bidirectional communication, in terms of interaction
between ECG and EEG signals, is altered in TGAC8. To this end, we used the Granger causality
approach to determine whether and to what extent heart-generated electrical signals causally
influence electrical cerebral activities (EEG patterns) and vice-versa (13) (Fig.4A, D).

**Figure 4.** *The causal interaction between heart and brain is increased in TGAC8 mice. (A) Granger Causality (GC) brain-*
*>heart estimate. The amplitude of the envelope of the band-filtered EEG signal (local amplitude) was extracted and used*

We filtered the ECG traces for Low (LF) and High Frequencies (HF), which substantially represent

the autonomic modulation of heartbeat intervals. Analogously, EEG were also filtered to obtain a

signal for each band (from δ up to γ3). The GC analysis takes advantage of the variability expressed

over time by both brain and heart rhythm. Indeed, it aims to derive the mutual causal influence

between heart rate sources (LF and HF) and EEG band activities, and vice versa. Hence, for both

ECG and EEG band filtered signals we considered the local amplitude, i.e. envelopes showed in

figures 4 A, and D. In terms of brain-to-heart influence, we observed a significant increase in the

causal interaction for each EEG band (except for delta) towards LF-filtered ECG RR intervals (Fig.

4B). This is not surprising when considering the necessity of the brain to modulate the elevated

HR and contractility of TGAC8 negatively. However, no significant changes were observed in the

HF-filtered ECG RR intervals range. Concerning the relation between LF-filtered ECG and EEG, we

observed a marked rise in the exchange of information for all the three gamma bands, i.e., the LF

component of heartbeat influences and guides changes in TGAC8s EEG patterns in the range of

gamma (Fig. 4E). However, we noticed no differences between TGAC8 and WT when examining

HF-filtered ECG (Fig. 4F).

We interpret these data to indicate that the flow of information between the heart and brain in

TGAC8 is increased. This increment is particularly evident between the LF component of the ECG

and the gamma frequencies of the EEG, likely accounting for the gamma rhythms rise observed

in TGAC8.

## Changes in GABA and glutamate receptors account for increased gamma activity in the hippocampus of TGAC8 mice

***Figure 5.*** *Transcriptomic/proteomic analysis of WT and TGAC8 hippocampus and SA node. (A) vulcano plots of transcriptomic, proteomic and phosphoproteomic analysis of TGAC8 mice hippocampus: protein coding RNAs, proteins, phosphopeptides, proteins/phosphopeptides ratio downregulated in blue, upregulated in red; (B) GABAergic synapse scheme, upregulated receptors in red, downregulated in blue; (C) glutamatergic synapse scheme, upregulated receptors in red, downregulated in blue; (D) Dopamine receptor 5 (D5) transcrption levels in the hippocampus; (E) Dopamine receptor 4 (D4) transcrption levels in the SA node. Results in D and E are shown as mean and SD, n=4, unpaired t-test between WT and TGAC8, †p<0.01, ‡p<0.001.*

Because EEG recording indicated that TGCA8 mice displayed a marked increase in gamma power -from 30 *Hz* to 160 *Hz* – when these mice were active (Fig.2 G), we next determined what long-term neurochemical adaptations may subtend these changes. The hippocampal formation is one of the primary hubs of gamma activity generation, especially high-speed gamma ripples (14). The earliest model of gamma oscillations generation is based on the reciprocal connections between pools of excitatory pyramidal neurons and inhibitory interneurons, i.e., resulting from a GABAergic/glutamatergic interplay (15). Against this background, we performed a transcriptomic/proteomic analysis to determine eventual changes (long-term adaptations) in the transcription and expression of GABA and glutamate receptors' transcription and expression levels. The receptor GABA-A is now recognized to play a significant role in gamma activity generation (15). Consistent with this acquired evidence, we found the increased EEG gamma

frequency bands in TGAC8 transgenic group coupled with a marked up-regulation of GABA-A.
Moreover, the GABA-B, known to inhibit gamma wave generation (16) were downregulated in
TGAC8 (Fig.5B). There is a close relationship between GABA-dependent signaling and glutamate
receptors synapses in gamma rhythms generation (15, 17). Accordingly, the excitatory receptor
mGLU1/5 was upregulated in TGAC8. In contrast, the inhibitory mGLU4/6 and mGLU4/8 were both
downregulated in these animals (Fig. 5C). Taken together, these data link the augmented EEG
gamma bands to long-term adaptation/remodeling of the GABA/glutamate receptor signaling
involved in gamma wave generation, at least at the hippocampal level.

**TGAC8 mice show an increment of dopamine signaling within the heart, circulating plasma,**
**and hippocampus**

The sinoatrial node of the TGAC8s mice displays elevated tyrosine hydroxylase (TH) expression
levels with reduced transcription of dopamine beta-hydroxylase: effects converging to augment
dopamine bioavailability, as previously shown (6). Moreover, the same study demonstrated a
sizable rise in circulating L-Dopa and dopamine in TGAC8 (6). Here, we assessed dopamine
receptor transcription levels in the sinoatrial node cells and observed a marked increase in the
transcription of the D4 receptor (Fig. 5E). Of note, at the cardiac level, this isoform modulates the
activity of the autonomic nervous system (18). This evidence fits very well with the altered
bidirectional heart-brain communication detected in the Granger causality analysis. On these
grounds, and bearing in mind that DOPA can pass the blood-brain barrier, we hypothesized that
increased levels of cardiac-derived DOPA coupled with altered *bottom-up* autonomic signaling
account, at least partly, for the hyperlocomotion manifested by the TGAC8 mice. We evaluated
dopamine receptor levels in the hippocampus to explore this hypothesis further. Indeed, recent
studies proved that the interaction between GABA-A and dopamine D5 receptors is critical to
evoking a rise in GABAergic inhibitory postsynaptic currents (IPSCs) (19), an event at the
foundation of gamma rhythms generation. In keeping with our initial hypothesis, we found that
the expression of D5 receptors was increased in the TGAC8 hippocampus (Fig. 5D).

## Discussion

**Figure 6. Mechanistic framework of the study.** *At the cardiac level, mice overexpressing adenylyl cyclase type 8 in cardiomyocytes display heightened dopamine anabolism, thus elevated circulating dopamine/DOPA levels, paralleled by the mRNA transcript upregulation of D4 dopamine receptors - primary modulators of cardiac autonomic activity - in the SA node. Centrally, the hippocampus of the TGAC8 mice retains increased expression of both GABA-A and D5 dopamine receptors. The activation of the former chiefly accounts for gamma activity generation, while the latter exerts a permissive (synergistic) action on GABA-A activation, hence favoring overall hippocampal gamma rhythms generation. Owing to this heightened (bottom-up) heart-brain interaction (confirmed by the Granger causality analysis of the ECG and EEG traces), AC8 transgenesis at the myocyte levels initiates and substains enhanced locomotor activity.*

The central nervous system continuously monitors peripheral internal (and external) body changes (1); for example, the degree of contraction of visceral muscles, heart rate, and metabolite levels in the internal milieu. Collected by the interoceptive system, these modifications signal back to sensory CNS regions encrypting body region representations to command physiological reactions. The latter ultimately modifies body functions to cope with stress conditions and/or maintain organismal homeostasis (1). Our observations fit nicely in this conceptual framework while providing unprecedented experimental evidence that buttresses and expands the somatic theory of emotions (20). Capitalizing on a mouse model that overexpresses AC8 in cardiomyocytes, here we report that TGAC8 mice manifest increased locomotor behavior sustained by enhanced EEG gamma rhythms. In these mice, the analysis of causality interplay (Granger) between ECG and EEG signals attests to a causal involvement of the autonomic component of the heartbeat in shaping EEG gamma oscillations. Mechanistically, the heart of TGAC8 mice showcases heightened dopamine anabolism translating into elevated circulating dopamine/DOPA, paralleled by an upregulation of D4 dopamine receptor. The latter is a primary modulator of cardiac autonomic activity. Centrally, the hippocampus of the TGAC8 mice retains upregulated expression of both GABA-A and D5 dopamine receptors; the activation of the former chiefly accounts for gamma activity generation, while that of the latter exerts a permissive

255 (synergistic) action on GABA-A activation, favoring hippocampal inhibitory currents generations
256 (Fig. 6).

**Overexpressing AC8 in cardiomyocytes triggers a locomotion status akin to perpetual**
**exercise**

Mice overexpressing AC8 in cardiomyocytes (TGAC8 mice) exhibit a global and persistent
increase in locomotor activity independent of changes in other behavioral variables, such as
anxiety. We ground our conclusion on the following experimental evidence: 1) actigraphy (24 hrs.
recording) showing increased activity in TGAC8s; 2) open field and elevated plus maze showing
increased moving time and distance traveled; 3) superior motor performance, i.e., higher average
speed and higher maximum speed and decreased freezing. Of relevance, both OF, EPM, and light-
dark box reveal the absence of anxiety-like behavior in these mice. This conclusion is reinforced
by the fear-conditioning test, an anxiety-centered test completely independent of motor
influences.

We advance the possibility that the rise in locomotor activity found in the TGAC8 is
"functionally justified" and congruent with chronic peripheral physiological modifications, such as
the persistently elevated heart rate. Put simply, the TGAC8 mouse heart experiences and must
handle a status akin to "perpetual exercise" where the brain, in keeping with the somatic theory
of emotions, adapt its outflow, switching the animal behavior from a state of rest to an active one,
such as running. This scenario also corresponds to the significant cardiac workload sustained by
the TGAC8 mice. Moreover, that these transgenes showcase hyperlocomotion but not anxiety is,
in our opinion, not so surprising. Indeed, their cardiac chronotropic rhythm is regular and
continuous, mimicking a "perpetual exercise" status. Conversely, sudden and irregular bouts of
heart rhythm perturbations typically subtend an anxious behavior, such as palpitations and/or
frank arrhythmias (21, 22). We posit that the modified pattern of brain-heart communication
observed in the TGAC8 mice occurs in anticipation of exercise performance but in the reverse
order. Normally, the brain commands the heart to increase its beating rate, ultimately preparing
the body to augment physical efforts (23). In stark contrast, in the present experimental setting,
the AC8-driven cardiac humoral (endocrine) and functional alterations signal back to the brain to

initiate locomotion. In essence, we postulate that, as with other behavioral domains such as
emotion control, a heart-brain loop also exists to start and/or fine-tune locomotion.

***Gamma rhythms, locomotion, and heart-brain causal interaction***

In rodents, hippocampal theta and gamma rhythms have been linked to locomotion, controlling
both its initiation and speed (11, 12). Heightened running speed is accompanied by significant,
systematic rises in the frequency of hippocampal network oscillations spanning the entire gamma
range (30–120 *Hz*) and beyond (12). Changes in frequency correlate strongly with firing rate
changes of individual interneurons mediated by GABA-A receptors, consistent with gamma
generation models (15). In our experimental setting, the 24 hr. EEG recordings documented that,
during the active phases, the TGAC8 mice experienced a marked rise in gamma frequencies from
30 *Hz* to 160 *Hz*. When correlating the animal motor activity with theta and gamma frequencies,
we found a prominent correlation between theta frequencies and movement in WTs. Conversely,
such correlation was no longer evident (completely abolished) in TGAC8 mice where gamma
activity correlated with locomotion.

Delving further into this finding, we performed some transcriptomic and proteomic
analyses of the hippocampus, i.e., the fundamental hub of gamma activity generation and a
primary player in locomotor activity control (11, 14). We observed a biomolecular profile that
nicely mirrors what was detected at the EEG level. Indeed, all receptors accounting for gamma
activity generation (15, 17) were upregulated (especially GABA-A and mGLUr 1/5), while those
implicated in gamma inhibition (16), i.e., GABA-B and mGLUr 4/8, were down-regulated.

Finally, to establish a causal link between the EEG pattern changes and cardiac activity
stemming from AC8 cardiac overexpression, we evaluated the extent and characteristics of the
heart/brain bidirectional communication, applying the Granger Causality approach to EEG and
ECG traces. The main finding is that the autonomic component of the heartbeat (especially LF)
causally influences the gamma frequencies in a *bottom-up modality*. Specifically, our analysis takes
advantage of the variability expressed over time by both the heart rhythm (decomposed in LF and
HF) and EEG frequency bands and checks whether amplitude variations of the former are
predictive of changes in the latter.

Prior functional magnetic resonance imaging studies using the Granger approach revealed
the hippocampus as a pivotal structure in the top-down control of the cardiac output. However,
the same study failed to identify causal interactions in the opposite direction, i.e., from the heart
to the brain, likely because performed in animals under resting conditions (24). However, recent
work by Candia-Rivera and collaborators unveiled the temporal dynamics of brain and cardiac
activities in human subjects who underwent emotional elicitation through videos. This research
demonstrates, via computational modeling, that emotional stimuli modulate heartbeat activity,
stimulating a specific cortical response (3). These observations corroborate both Ledoux's and
Damasio's theories on emotions. However, in the study by Candia-Rivera and colleagues, the
original stimulus, i.e., the trigger of the adaptive brain response, was not peripheral (somatic), in
principle. In our study, we specifically overexpressed AC8 peripherally in the heart, resulting in
heart rate elevation, heart contractility, and alterations in heart and plasma neurotransmitter
levels. While escaping top-down sympathetic control, these alterations would modify the heart-
brain communication in a mere bottom-up modality.

***Dopamine and the heart-brain loop***

Sinoatrial node cells of the TGAC8s mice harbor markedly elevated levels of tyrosine
hydroxylase (TH) with reduced transcription of dopamine beta-hydroxylase, as shown previously
(6). And these changes lead to elevation of cardiac and systemic DA/DOPA availability (6). Here,
we add to this mosaic two new pieces: first, TGAC8 sinoatrial node cells display a marked rise in
D4 transcription levels (Fig. 4E); second, these transgenic mice have an increased hippocampal
transcription of D5 receptor (Fig. 4D). In aggregate, the new evidence may have the following
functional consequences. First, a recent study shows how atrial D4 can modulate vagal activity
(18) (i.e., heart-to-brain and brain-to-heart interaction). This fits very well with what we have
observed at the level of bidirectional heart-brain communication by Granger analysis. Second, the
incremented levels of circulating DOPA output by the TGAC8 heart can permeate the brain-blood-
barrier (BBB), reaching central dopaminergic stations fundamental for the generation and
maintenance of locomotor activity such as the basal ganglia (25). Third, the D5 receptor,
upregulated in the TGAC8 hippocampus, is one of the primary activators of GABA-A, the main
catalyst of gamma activity (15). Accordingly, recent work shows that functional cross-talk exists

between GABA-A and D5 in hippocampal neurons, favoring inhibitory current generations (18).

Thus, the present results support the novel idea that cardiac dopaminergic signaling shapes heart-

to-brain communication through direct endocrine mechanisms (systemic DOPA production) and

autonomic ones (D4 modulation on afferent vagal transmission), as well as indirect ones, such as

hippocampal upregulation of the D5 receptors.

***Limitations and studies in perspective***

The chosen experimental model, i.e., persistent intracardiac AC8 overexpression, can be perceived

as a limitation of the present study because it is unlikely that AC8 could be so selectively

upregulated in a clinical setting. However, such an artifact is instrumental to the current research's

conceptual and methodological strut: to generate a peripheral physiological variation to assess

how cardiac autologous humoral/functional signaling reverberates back to the brain. Future

studies could use, for instance, a loss-of-function approach instead, determining, for example, the

repercussions of cardiac-specific deletion of AC8 on heart properties and locomotion to further

validate the current findings. Admittedly, we focused primarily on the hippocampus because it

regulates locomotor activity while simultaneously generating gamma EEG activity (9). However,

we believe that the results of our study will stimulate future studies focusing on other brain

regions/areas that are crucially involved in the directional heart-brain communication (for

example, the nucleus tractus solitarius or the hypothalamus) or movement control (e.g., basal

ganglia and septum).

**Conclusions**

These studies inform how heart-initiated specific humoral/functional changes (in the present case,

elevated dopamine/DOPA signaling and increased myocardial performance due to AC8

overexpression) trigger modifications of just as definite brain areas (hippocampus) to shape

specific adaptive responses, such as locomotion. These data reinforce and expand the relevance

of somatic theories of emotions.

**Materials and Methods**

**Animals**

All studies were executed in agreement with the Guide for the Care and Use of Laboratory Animals published by the National Institutes of Health (NIH Publication no. 85-23, revised 1996). The experimental procedures were approved by the Animal Care and Use Committee of the National Institutes of Health (protocol #441-LCS-2016). A breeder pair of TG$^{AC8}$ mice, generated by ligating the murine α-myosin heavy chain promoter to a cDNA coding for human AC8 (26), were a gift from Nicole Defer/Jacques Hanoune, Unite de Recherches, INSERM U-99, Hôpital Henri Mondor, F-94010 Créteil, France. Wild type (WT) littermates, bred from the C57BL/6 background, were used as controls.

**Behavioral tests**

All the behavioral tests were conducted on mice with transmitters already implanted. Total time mobile or immobile was evaluated during the 24 hours of EEG/ECG recording phase through actigraphy (included in the double implant). All the other behavioral assays were conducted adopting a one-week interval between tests to allow the mice to recover. Tests were performed at the same time of day, with no reversed dark/light cycle.

**Open field**

Open field (OF) is a gold standard behavioral test used to study locomotor activity, exploratory behavior, and anxiety-like behavior in rodents (27). The animal is free to explore a circumscribed environment (50cm x 50cm arena surrounded by walls) for 10 minutes, and its locomotor activity and tendency to explore open spaces or hide in the peripheral corners of the arena is evaluated. The analysis of the behavioral parameters has been performed using ANY-maze software (Stoelting Co., IL, USA).

**Elevated plus maze**

Elevated plus maze (EPM) is another consolidated test to study rodents' anxiety-like behavior and locomotor activity. Elevated Plus Maze test exploits the rodent's conflict between aversion to open

spaces (i.e., hides in closed arms of a labyrinth) and instinct to explore new environments (e.g.,

exploration of open arms of the same maze). The testing apparatus consisted of four white arms

at a 90∘ angle from each other. Tall walls surrounded two "closed" arms, and two "open" arms had

no walls (27). Animals were placed at the center of the apparatus facing an open arm and allowed

to explore for 5 minutes freely. The analysis of the behavioral parameters has been performed

employing ANY-maze.

**Light-Dark Box**

The light/dark box (LDB) test is based on the innate aversion of mice to illuminated spaces and

the spontaneous exploratory activity in response to minor stressors, such as a new environment

and light. This experimental task allows the evaluation of the level of anxiety-like behavior

experienced by the animals. The test apparatus comprises a small, dark, and larger illuminated

compartment (28). The analysis of the behavioral parameters has been performed through ANY-

maze software.

**Fear Conditioning Test**

The fear conditioning (FC) test measures the capacity of rodents to learn and recall an association

between environmental cues and aversive experiences. Testing was conducted in standard

conditioning chambers (*MED-Associates; St Albans, VT*) equipped with a stainless-steel rod floor

that delivered an aversive shock, a ventilation fan emitting background white noise, video

recording camera, and a white light that illuminated the chamber. During the 8-minute training

session mice were placed in the training context and allowed to explore the environment freely.

After two minutes, an auditory cue (80 dB) was presented for 30s, with an electric footshock (0.35

mA) delivered continuously during the final 2s of the auditory cue. The tone-shock pairing was

repeated on two separate occasions (at 240s and 360s). Mice were left undisturbed in the chamber

for 90s and then returned to the home cage to conclude training. Approximately 24 hours later,

the conditioning acquisition mice were individually retrieved and placed in the same training

context for five minutes for contextual conditioning testing. One hour later, mice were tested for

cued conditioning. In this paradigm, mice were first acclimated to a new room for one hour, with

red room lighting. Cued fear conditioning was measured in a different context that utilized a white

foam floor covering the electrical grid, a modified chamber shape from a black triangle insertion, and white noise was absent. During testing, mice were exposed to the same auditory tone for 3 minutes after an initial habituation period of 3 minutes. Freezing behavior during the test was recorded as an index of associative learning (10).

**Telemetry double implant to simultaneously monitor the EEG and ECG**

Telemetric radio transmitters (F20-EET; Data Sciences International (DSI), St. Paul, MN) were surgically implanted in young (3-4 months) WT and TG$^{AC8}$ mice as described (29). Briefly, two surface electrodes, a positive electrode (parietal cortex; AP, -2.0 mm; L, 2.0 mm) and a reference electrode (cerebellum; AP, -6.0 mm; L, 2.0 mm), were passed subcutaneously to the cranial base and placed directly on the dura mater. Two additional bipotential electrodes were routed subcutaneously via a vertical midline incision overlying the abdomen with leads situated in the right upper chest and lower left abdomen below the heart to monitor continuous heart rate (Fig.1). Following a two-weeks recovery period, 24-hr free behaving electrocardiogram (ECG) and electroencephalogram (EEG) were recorded at a sampling rate of 1000 Hz and 500 Hz, respectively, with simultaneous activity recording using the Dataquest ART acquisition system (DSI, version 4.36). Average activity counts were obtained every 10s.

**Signal analysis**

**ECG analysis**

Scoring of wake-sleep states (NeuroScore, DSI, version 3.2.0) allowed for selecting all 10 seconds wake epochs and extracting signals.

After appropriate pre-processing (see Supplemental Information), the series of heartbeat time intervals (RR series) was processed to derive HRV features. All features were signal-length independent (30) estimated on 10 seconds epochs to account for highly frequent wake/sleep switches, thus describing the short-time heartbeat dynamics. More in detail, we considered: the mean value of RR intervals and, from frequency spectrum analysis (Hamming-windowed FFT), the power in low frequency (LF: 0.15–1.5 Hz) and high frequency (HF: 1.5–4 Hz) bands. We also estimated some nonlinear features that provide additional information about cardiac dynamics.

According to previous works studying non-linearity in a heartbeat series (6), we characterize the

RR series auto similarity through the Detrended Fluctuation Analysis (DFA).

**EEG Spectral analysis**

For each 10 seconds wake epoch, a spectral analysis was performed on the EEG signal. The mean

power spectrum density characterizing the wake state of each animal was evaluated by applying

a Hamming-windowed Fast Fourier Transform (FFT) and averaging them. Power values were

measured in dB, thus, data from FFT were log-transformed (31).

**Brain/heart communication**

The brain/heart communication study was based on a Granger Causality (GC) analysis. $GC_{x->y}$ is a

measure of the contribution of the past of the x(t) time series to the prediction of the present

value of y(t), compared to the contribution of the past of the *y(t) time series* in the prediction of

its own present value. In our application (Fig. 4) x(t) and y(t) are signals related to brain and heart

functioning, and the GC has allowed estimating heart rhythm influence on brain EEG rhythms and

vice versa (by switching x and y). Specifically, our analysis takes advantage of the variability

expressed over time by both the brain rhythms and heart rhythms and checks whether amplitude

variations of the former are predictive of changes in the latter or vice versa. The GC analysis was

conducted using the Causal Connectivity Toolbox (32), which assumes a linear model and order

within the model, estimated using the Akaike Information Criterion. Herein, in unison with the GC

applicability criteria, EEG epochs satisfying the stationarity test (33) were retained for the analysis

(more than 95% retained). For each mouse, the median of their epochs model orders was chosen

as the representative order. The 95[th] percentile of 'animals' model orders was taken as the overall

order (13): from data collected in this experiment, this corresponded to 20 samples, equivalent to

2 seconds. The overall model order was applied to the GC estimates of all wake epochs for all

mice (both groups). Finally, the validity of the model order was verified by estimating the model

consistency (percentage of data correlation structure explained by the model)[19]. Consistency

values higher than 75% were considered satisfactory, and all periods had mean consistencies of

90% or above.

**RNA sequencing (RNA-seq) and transcriptomic data analysis**

We used six hippocampal formations from six transgenic mice and six hippocampal formations
from six wild-type littermates for the RNA sequencing experiment. The differential expression
gene (DEG) analysis was performed using the DESeq2 package in R language. In total, the
expression of 17542 protein-coding mRNAs was identified. The adjusted p-value and fold change
calculated by DESeq2 (34) were used to draw a volcano plot to figure out the expression changes
in transcriptome. Specifically, 927 genes were downregulated (-log10(adjusted p-value) > 1.3 and
log2(fold change) < 0) and 666 upregulated ((-log10(adjusted p-value) > 1.3 and log2(fold
change) > 0).

**Mass spectrometry and proteomic and phosphoproteomic data analysis**

4 hippocampla tissues from 4 transgenic mice and 4 hippocampal tissues from 4 wild-type
littermates were used for the mass spectrometry experiment. Samples were labeled with the 10-
plex tandem mass tag (TMT) according to Thermo Scientific's TMT Mass Tagging kits protocol.
About 5% of the labeled tryptic peptides from each of the 24-fractions were used for global
proteomics analysis, while the remaining 95% in the 24-fractions were then pooled into 12-
fractions and were subjected to subsequent TiO2-enrichment with a Titansphere Phos-kit (GL
biosciences Inc.). All MS and MS/MS raw spectra of TMT experiments from each set were
processed and searched using Sequest HT algorithm within the Proteome Discoverer 2.2 (PD2.2
software, Thermo Scientific). The subsequent analyses were adapted from our previous work (35).
Specifically, the raw counts of total proteins were normalized to the total counts in each sample,
and log2 was transformed. At last, p-values were calculated from the normalized and transformed
counts, and fold changes between TG and WT were calculated upon the average expression
between the two groups. For quantitative phosphopeptides analysis, additional phosphorylation
on Ser, Thr, Tyr residues were specified as variable modifications. We performed the subsequent
analyses in two strategies, one for the phosphopeptides and another for the ratio between
phosphopeptides/total proteins (also called normalized phosphorylation). For the for the ratio
between phosphopeptides/total proteins part, the raw count of phosphorylation was first divided
by the raw count of its corresponding protein to obtain the raw normalized phosphorylation
count. Then, the raw phosphopeptides or raw normalized phosphorylation counts were
normalized to the total counts in each sample, and log2 transformed. At last, p-values were

calculated from the normalized and transformed counts, and fold changes between TG and WT

were calculated upon the average expression between the two groups.

The p-value and fold change in total proteins, phosphopepetides, and phosphopepetides/total

proteins were used to draw volcano plots to determine the expression changes in proteome and

phosphoproteome. Specifically, 84 proteins were downregulated (-log10(p-value) > 1.3 and

log2(fold change) < 0) and 123 upregulated ((-log10(p-value) > 1.3 and log2(fold change) > 0);

158 phosphopeptides were downregulated (-log10(p-value) > 1.3 and log2(fold change) < 0) and

286 upregulated ((-log10(p-value) > 1.3 and log2(fold change) > 0); 196 ratios of

phosphopeptides to proteins were downregulated (-log10(p-value) > 1.3 and log2(fold change)

< 0) and 290 upregulated ((-log10(p-value) > 1.3 and log2(fold change) > 0). The normalized and

transformed counts and fold changes between TG and WT were calculated upon the average

expression between the two groups. For quantitative phosphopeptides analysis, additional

phosphorylation on Ser, Thr, Tyr residues were specified as variable modifications. We performed

the subsequent analyses in two strategies, one for the phosphopeptides and another for the ratio

between phosphopeptides/total proteins (also called normalized phosphorylation). For the for the

ratio between phosphopeptides/total proteins part, the raw count of phosphorylation was first

divided by the raw count of its corresponding protein to obtain the raw normalized

phosphorylation count. Then, the raw phosphopeptides or raw normalized phosphorylation

counts were normalized to the total counts in each sample, and log2 transformed. At last, p-values

were calculated from the normalized and transformed counts, and fold changes between TG and

WT were calculated upon the average expression between the two groups. The p-value and fold

change in total proteins, phosphopeptides, and phosphopeptides/total proteins were used to

draw volcano plots to determine the expression changes in proteome and phosphoproteome.

Specifically, 84 proteins were downregulated (-log10(p-value) > 1.3 and log2(fold change) < 0)

and 123 upregulated ((-log10(p-value) > 1.3 and log2(fold change) > 0); 158 phosphopeptides

were downregulated (-log10(p-value) > 1.3 and log2(fold change) < 0) and 286 upregulated ((-

log10(p-value) > 1.3 and log2(fold change) > 0); 196 ratios of phosphopeptides to proteins were

downregulated (-log10(p-value) > 1.3 and log2(fold change) < 0) and 290 upregulated ((-log10(p-

value) > 1.3 and log2(fold change) > 0).

## RT-qPCR

RT-qPCR of the hippocampus and SA tissue was performed to determine the transcript abundance of dopamine receptors (n=4 WT and 4 TG$^{AC8}$ mice). RNA was extracted from the hippocampus with TRIzol™ Reagent (Thermo Fisher Scientific, Waltham MA) and DNAse on column digestion. The cDNA was prepared using MMLV reverse transcriptase (Promega, Madison, WI) using 500 ng of total RNA per reaction. RT-qPCR was performed using a QuantStudio 6 Flex Real-Time PCR System (Thermo Fisher Scientific, Waltham MA) with a 384-well platform. The reaction was performed with a FastStart Universal SYBR Green Master Kit with Rox (Roche, Indianapolis, IN) using the manufacturer's recommended conditions; the sizes of amplicons were verified. Each well contained 0.5 µl of cDNA solution and 10 µl of the reaction mixture. Each sample was quadruplicated. Preliminary reactions were performed to determine the efficiency of amplification. RT-qPCR analysis was performed using the ddCt method. *Hprt* was used as a housekeeping gene and results represented as Mean of Relative Quantification (RQ) normalized to *Hprt* +/- SD (7). Primers were selected with Primer Express 3.0 software (Applied Biosystems).

## Statistical analysis procedures

Results are presented as violin plots. All parametric data were analyzed by unpaired t-tests between the WT and TGAC8 groups. Correlations between groups of values were evaluated through multiple linear regressions (36).

## Funding

This research was supported by the Intramural Research Program of the NIH, National Institute on Aging (USA).

1. A. Damasio, G. B. Carvalho, The nature of feelings: evolutionary and neurobiological origins. *Nat Rev Neurosci* **14**, 143–152 (2013).

2. J. A. Armour, The little brain on the heart. *Cleve Clin J Med* **74 Suppl 1**, S48-51 (2007).

3.      D. Candia-Rivera, V. Catrambone, J. F. Thayer, C. Gentili, G. Valenza, Cardiac sympathetic-
vagal activity initiates a functional brain-body response to emotional arousal. *Proc Natl Acad Sci*
*U S A* **119**, e2119599119 (2022).

4.      J. Agrimi, *et al.*, Obese mice exposed to psychosocial stress display cardiac and
hippocampal dysfunction associated with local brain-derived neurotrophic factor depletion.
*EBioMedicine* **47**, 384–401 (2019).

5.      B. S. McEwen, P. J. Gianaros, Central role of the brain in stress and adaptation: Links to
socioeconomic status, health, and disease: Central links between stress and SES. *Annals of the*
*New York Academy of Sciences* **1186**, 190–222 (2010).

6.      J. M. Moen, *et al.*, Overexpression of a Neuronal Type Adenylyl Cyclase (Type 8) in
Sinoatrial Node Markedly Impacts Heart Rate and Rhythm. *Front Neurosci* **13**, 615 (2019).

7.      K. V. Tarasov, *et al.*, "A Remarkable Adaptive Paradigm Of Heart Performance And
Protection Emerges In Response To The Constitutive Challenge Of Marked Cardiac-Specific
Overexpression Of Adenylyl Cyclase Type 8" (Physiology, 2022)
https:/doi.org/10.1101/2022.05.20.491883 (August 5, 2022).

8.      P. G. Ravenstijn, H.-J. Drenth, M. J. O'Neill, M. Danhof, E. C. de Lange, Evaluation of
blood-brain barrier transport and CNS drug metabolism in diseased and control brain after
intravenous L-DOPA in a unilateral rat model of Parkinson's disease. *Fluids Barriers CNS* **9**, 4
(2012).

9.      D. Ho, *et al.*, Heart Rate and Electrocardiography Monitoring in Mice. *CP Mouse Biology*
**1**, 123–139 (2011).

10.      H. Shoji, K. Takao, S. Hattori, T. Miyakawa, Contextual and Cued Fear Conditioning Test
Using a Video Analyzing System in Mice. *JoVE*, 50871 (2014).

11.      F. Bender, *et al.*, Theta oscillations regulate the speed of locomotion via a hippocampus
to lateral septum pathway. *Nat Commun* **6**, 8521 (2015).

12.      O. J. Ahmed, M. R. Mehta, Running Speed Alters the Frequency of Hippocampal Gamma
Oscillations. *Journal of Neuroscience* **32**, 7373–7383 (2012).

13.      A. B. Barrett, *et al.*, Granger causality analysis of steady-state electroencephalographic
signals during propofol-induced anaesthesia. *PLoS One* **7**, e29072 (2012).

14.      A. Oliva, A. Fernández-Ruiz, E. Fermino de Oliveira, G. Buzsáki, Origin of Gamma
Frequency Power during Hippocampal Sharp-Wave Ripples. *Cell Rep* **25**, 1693-1700.e4 (2018).

15.      G. Buzsáki, X.-J. Wang, Mechanisms of gamma oscillations. *Annu Rev Neurosci* **35**, 203–
225 (2012).

16.    L. S. Leung, B. Shen, GABAB receptor blockade enhances theta and gamma rhythms in
the hippocampus of behaving rats. *Hippocampus* **17**, 281–291 (2007).

17.    G. Govindaiah, *et al.*, Group I metabotropic glutamate receptors generate two types of
intrinsic membrane oscillations in hippocampal oriens/alveus interneurons. *Neuropharmacology*
**139**, 150–162 (2018).

18.    J. Á. García-Pedraza, *et al.*, Dopamine D4 receptor subtype activation reduces the rat
cardiac parasympathetic discharge. *Pflugers Arch - Eur J Physiol* **472**, 1693–1703 (2020).

19.    F. Maingret, L. Groc, Characterization of the Functional Cross-Talk between Surface
GABAA and Dopamine D5 Receptors. *Int J Mol Sci* **22**, 4867 (2021).

20.    J. E. LeDoux, R. Brown, A higher-order theory of emotional consciousness. *Proc. Natl.*
*Acad. Sci. U.S.A.* **114** (2017).

21.    A. V. Abbott, Diagnostic approach to palpitations. *Am Fam Physician* **71**, 743–750 (2005).

22.    L. Carnevali, *et al.*, Low vagally-mediated heart rate variability and increased susceptibility
to ventricular arrhythmias in rats bred for high anxiety. *Physiology & Behavior* **128**, 16–25 (2014).

23.    R. Brown, U. Kemp, V. Macefield, Increases in muscle sympathetic nerve activity, heart
rate, respiration, and skin blood flow during passive viewing of exercise. *Front. Neurosci.* **7**
(2013).

24.    A. Duggento, *et al.*, Globally conditioned Granger causality in brain–brain and brain–
heart interactions: a combined heart rate variability/ultra-high-field (7 T) functional magnetic
resonance imaging study. *Phil. Trans. R. Soc. A.* **374**, 20150185 (2016).

25.    J. L. Lanciego, N. Luquin, J. A. Obeso, Functional neuroanatomy of the basal ganglia. *Cold*
*Spring Harb Perspect Med* **2**, a009621 (2012).

26.    L. Lipskaia, *et al.*, Enhanced cardiac function in transgenic mice expressing a Ca(2+)-
stimulated adenylyl cyclase. *Circ Res* **86**, 795–801 (2000).

27.    N. Sestakova, A. Puzserova, M. Kluknavsky, I. Bernatova, Determination of motor activity
and anxiety-related behaviour in rodents: methodological aspects and role of nitric oxide.
*Interdisciplinary Toxicology* **6**, 126–135 (2013).

28.    J. Agrimi, *et al.*, Psychosocial Stress Hastens Disease Progression and Sudden Death in
Mice with Arrhythmogenic Cardiomyopathy. *J Clin Med* **9**, E3804 (2020).

29.    R. Wan, S. Camandola, M. P. Mattson, Intermittent Food Deprivation Improves
Cardiovascular and Neuroendocrine Responses to Stress in Rats. *The Journal of Nutrition* **133**,
1921–1929 (2003).

30.    O. Michel, Holger kantz and Thomas schreiber, nonlinear time series analysis. *J Biol Phys*
**24**, 79–80 (1998).

31.    A. Zaccaro, A. Piarulli, L. Melosini, D. Menicucci, A. Gemignani, Neural Correlates of Non-
ordinary States of Consciousness in Pranayama Practitioners: The Role of Slow Nasal Breathing.
*Front. Syst. Neurosci.* **16**, 803904 (2022).

32.    A. K. Seth, A MATLAB toolbox for Granger causal connectivity analysis. *Journal of
Neuroscience Methods* **186**, 262–273 (2010).

33.    D. Kwiatkowski, P. C. B. Phillips, P. Schmidt, Y. Shin, Testing the null hypothesis of
stationarity against the alternative of a unit root. *Journal of Econometrics* **54**, 159–178 (1992).

34.    M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

35.    J.-H. Qu, *et al.*, "The reprogrammed mouse heart phosphoproteome in response to the
chronic stress of markedly high cardiac-specific adenylyl cyclase type 8 overexpression"
(Bioinformatics, 2022) https:/doi.org/10.1101/2022.04.29.488779 (August 9, 2022).

36.    D. A. Freedman, *Statistical Models: Theory and Practice*, 2nd Ed. (Cambridge University
Press, 2009) https:/doi.org/10.1017/CBO9780511815867 (August 2, 2022).

**Supplemental Information**

642
643    *Figure 7: Light-dark box test, TGAC8 mice do not show signs of anxiety-like behavior. A) Light-dark box test, transitions to
644    light; B) Light-dark box test, time in the light zone. Results are shown as violins plot, N=7, unpaired t-test has been
645    performed between WT and TGAC8 groups.*