Paninvasion severity assessment of a U.S. grape pest to disrupt the global wine market

Nicholas A. Huron*

Jocelyn E. Behm

Matthew R. Helmus

Integrative Ecology Lab, Department of Biology, Temple University, Philadelphia, PA 19122

USA.

*   corresponding author, nahuron@temple.edu

Abstract

Economic impacts from plant pests are often felt at the regional scale, yet some impacts expand

to the global scale through the alignment of a pest's invasion potentials. Such globally invasive

species (i.e., paninvasives) are like the human pathogens that cause pandemics. Like pandemics,

assessing paninvasion risk for an emerging regional pest is key for stakeholders to take early

actions that avoid market disruption. Here, we develop the paninvasion severity assessment

framework and use it to assess a rapidly spreading regional U.S. grape pest, the spotted lanternfly

planthopper (*Lycorma delicatula;* SLF), to spread and disrupt the global wine market. We found

that SLF invasion potentials are aligned globally because important viticultural regions with

suitable environments for SLF establishment also heavily trade with invaded U.S. states. If the

U.S. acts as an invasive bridgehead, Italy, France, Spain, and other important wine exporters are

likely to experience the next SLF introductions. Risk to the global wine market is high unless

stakeholders work to reduce SLF invasion potentials in the U.S. and globally.

Introduction

Invasive plant pests cause substantial economic impacts[1], but most pests and their

impacts are confined to specific regions. For a regional pest to become a globally invasive

species that disrupts global markets (i.e., paninvasive), ecological and economic factors that

determine the pest's transport, establishment, and impact potentials must be aligned at the global

scale (Fig. 1, Supplementary Methods)[2]. First, paninvasive pests have high transport potential

because they can be easily transported among regions, often through global trade[3]. Second,

paninvasive pests have high establishment potential, because their environmental needs for

population growth are met in many regions[4]. Third, paninvasive pests have high impact potential,

because invaded regions have sizeable agricultural production and industries vulnerable to the

pest[5]. If these invasion potentials are correlated across multiple regions globally for an emerging

regional pest, there is a high risk of the pest spreading to cause supply crashes in regional

markets that cascade to disrupt global markets[6].

Despite the importance of identifying emerging paninvasives, existing approaches lack a

cohesive and universal framework for rapidly assessing and effectively communicating such risk

to stakeholders[7]. To address this gap, we developed the paninvasion severity assessment

framework by adapting the U.S. CDC pandemic severity assessment framework to invasion

process theory, which describes translocations of species in terms of transport, establishment,

and impact potentials (Fig. 1, 2)[2,8–10]. Although invasive species frameworks are increasingly

adapted to understand infectious diseases like COVID-19[11–16], adapting public-health

frameworks to invasion science is novel and leverages an increasingly universal risk vocabulary

(Fig. 2)[17]. Under this framework, we assessed the paninvasion risk of the spotted lanternfly

planthopper (Hemiptera: *Lycorma delicatula*; SLF, Fig. 1). SLF was introduced to South Korea

and Japan in the early 2000s and then to the U.S. (ca. 2014) on goods imported from its native

China[18]. SLF has rapidly spread from Pennsylvania to several other states, presenting increased

opportunities for stepping-stone, bridgehead invasions to additional regions[19–21]. SLF greatly

impacts grape production[22–26] and has been presented to the public as one of the worst invasive

species to establish in the U.S. in a century[27–29], but its paninvasion risk has not been assessed[19].

SLF likely has high global transport potential because it lays inconspicuous egg masses

on plants, stone, and trade infrastructure (e.g., containers, railcars, pallets), which facilitates

long-distance transport when eggs are laid on exported items (Fig. 1a–c). Landscaping stone

imported from China was the likely vector of the U.S. invasion[18]. Following successful transport,

SLF global establishment potential is likely enhanced by the cosmopolitan distribution of its

preferred host plant, the tree-of-heaven (*Ailanthus altissima*, TOH, Fig. 1d–f)[21]. The native

ranges of SLF and TOH overlap in China, but for >250 years TOH has escaped cultivation into

disturbed habitats and agricultural margins in temperate, subtropical, and Mediterranean regions

globally (Fig. 1 map). Once established, SLF likely has high global impact potential to wine

markets because grape is an equally suitable host. SLF develops at similar rates when fed grape

or TOH, and fecundity increases when fed a mixed diet of these two preferred hosts[25,30–33]. Asian

vineyard production is impacted by SLF infestations[34,35]; and SLF-invaded U.S. vineyards have

reported vine deaths, >90% yield losses, and closure (Fig. 1g–i)[21,23,36].

To assess paninvasion risk, we calculated invasion potentials from estimates of SLF

transport, establishment, and impact potentials from the U.S. invaded region to uninvaded U.S.

states and countries using trade, species distribution models, and grape and wine production data.

We then mapped invasion potentials, calculated alignment correlations, and estimated risk to the

$300B global wine market[37].

Results

Spotted lanternfly invaded range

To assess SLF paninvasion risk, we first estimated the current U.S. invaded range (ca.

2020). We aggregated distributional data from multiple sources, including announcements made

by state departments of agriculture on cargo interceptions that did not lead to established

populations (i.e., regulatory incidents). By 2020, SLF had established in nine U.S. states, with

clear incidents of long-distance transport and establishment in Virginia, New York, and

Pennsylvania. In eight additional states, individuals were intercepted in cargo and on transported

goods originating from states with established SLF. California had intercepted the most with >40

individual SLF on 35 flights found during cargo inspections, but all were dead or moribund and

not egg masses[38]. No international reports of regulatory incidents from the U.S. have been

published. These regulatory incidents suggest that cargo with SLF were frequently transported

from the invaded range in the U.S. northeast to at least as far as to the U.S. west coast (Fig. 3).

Alignment of invasion potentials

We estimated the three invasion potentials—transport, establishment, impact—for 50

U.S. states and 223 countries. For transport potential, we used the total metric tonnage of goods

imported from invaded states[39,40]. The current SLF spread in the U.S. (Fig. 3) was explained by

this transport potential metric (Supplementary Methods, Supplementary Table 1). States with the

highest transport potential were mostly in the eastern US, but Illinois, Texas, and California also

heavily traded with the invaded states, indicating that SLF had high potential to be transported

both regionally and transcontinentally (Fig. 4a, 5a). Globally, transport potentials were highest in

several European countries, Canada, and Brazil (Fig. 4b, 5b).

We based establishment potential on an ensemble estimate of species distribution models

(SDMs) built on SLF and TOH geolocations (see Supplementary Methods for SDM methods,

Fig. 4)[41]. Our ensemble estimate of SLF establishment potential was spatially similar to other

SDMs[42,43] and physiologically based demographic models of SLF[44,45]. However, our estimate

indicated urban landscapes as likely establishment locations and showed fine spatial-scale

variation in establishment potential (see our interactive map https://ieco-lab.github.io/slfrsk/).

For each state and country, we extracted the mean, median and max predicted suitability to

estimate establishment potential (Supplementary Methods). Most U.S. states had high

establishment potential (Fig. 4a, 5a), and all the countries with highest transport potential also

had the highest establishment potential (Fig. 4b, 5b).

We estimated SLF impact potential as the annual average tonnage of grapes and wine

produced for each U.S. state and country[46–48]. States and countries with many important

viticultural regions were geographically clustered (Fig. 4). Those with the highest impact

potentials included: California, Washington, and Oregon on the U.S. west coast; New York and

Pennsylvania on the U.S. east coast (Fig. 4a, 5a); and Italy, France, and Spain in western Europe

(Fig. 4b, 5b). After discovery of the Pennsylvania invasion in 2014, it only took a few years for

neighboring states to be invaded as well (Fig. 3). Thus, should one region on the U.S. west coast

or in western Europe become invaded by SLF, neighboring states and countries are likely to also

become invaded quickly.

SLF invasion potentials across states and countries were aligned (Fig. 5). Alignment

correlations calculated as Spearman's rank correlations (ρ statistic) among transport,

establishment, and impact potentials were positive for impact potential measured as state grape

production ($\rho = 0.41$, $P<0.005$, Fig. 5a), state wine production ($\rho = 0.52$, $P<0.001$), country

grape production ($\rho = 0.67$, *P*<0.001, Fig. 5b), and country wine production ($\rho = 0.63$, *P*<0.001).

This alignment of potentials is clear in the invasion-potential alignment plots (Fig. 5). Major

grape producing regions fall in the upper-right quadrant of the plots where regions have both

high transport and high establishment potentials.

Paninvasion risk

We estimated the risk of SLF to disrupt the global wine market to be an 8 out of 10 (Fig.

6). To derive this value, we regressed country grape production on country transport and

establishment potentials. Each predicted value from this multivariate regression can be

considered an estimate of the risk of SLF to invade and impact a country's grape production. We

then rescaled these predicted values from 1–10 and correlated them to wine export market size (ρ

$= 0.66$, *P*<0.001). To place SLF on a scale of paninvasion severity, we rescaled the correlation

coefficient, ρ, from 1–10, so that 1 is a complete negative correlation and 10 is a complete

positive correlation between predicted impact and market size. Low values on this scale indicate

that the global market is buffered against a paninvasion, while high values indicate that a

paninvasion is likely unless mitigation actions are taken to reduce invasion potentials.

Discussion

The risk of a spotted lanternfly (SLF) paninvasion is high and coordinated effort should

be made to reduce its transport, establishment, and impact potentials globally. In the U.S., efforts

to reduce SLF transport potential are primarily through quarantine and inspection of goods. SLF

is a regulated plant pest and the U.S. Department of Agriculture (USDA) is working towards

implementing consistent, science-based, and nation-wide transport protocols[21,49,50]. We

recommend that estimates of SLF transport potential be updated regularly as more states become

invaded. Finer spatial and temporal scale data are needed on high transport potential pathways

such as rail, landscaping stone, and live tree shipments to better forecast long-distance

transport[18,21,34]. Egg masses in diapause are the most likely life stage to result in new satellite

populations if transported. Thus, models should integrate annual trade dynamics with SLF annual

phenology[44,51]. Such models would identify goods to quarantine and inspect that are transported

out of the invaded range when eggs are in diapause before spring hatch[52].

The main host plant for SLF is the invasive tree of heaven (TOH) and reduction of

establishment potential in the U.S. focuses on removing and treating TOH with herbicide[18]. The

USDA and state departments of agriculture are the main agencies tasked with SLF control.

However, these agencies lack resources to manage TOH at both broad and local scales needed to

reduce establishment potential, and businesses and private citizens are increasingly burdened

with TOH removal costs[21]. Future research should focus on cost-effective, targeted TOH

biocontrol methods that can be implemented at broad spatial scales [53]. We suggest increased

public funding for local TOH management around vineyards and properties at risk of

transporting SLF[21]. Currently, TOH seeds are inexpensive and easily purchased online. We

suggest eliminating the horticultural sale of TOH globally. The U.S. federal and state

governments should put TOH on their noxious weed lists, making it unlawful to sell, grow, or

move this weedy invasive[54]. As SLF spreads, it will encounter new hosts besides TOH. To

anticipate novel hosts, association studies that track SLF survivorship, development speed, and

attraction are needed across a wide range of potential hosts, especially for those in landscapes

surrounding high transport and impact potential locations like railyards and vineyards[22,30,31,55].

Reduction to SLF impact potential currently relies on reducing populations with tree-

band trapping and broad-spectrum insecticides (e.g., carbamates, organophosphates, pyrethroids,

neonicotinoids) that have high nontarget mortality[18,23,56,57]. However, existing management

practices do not prevent vineyard reinfestations and pesticide application often overlaps with

grape harvest when adults move into vineyards[36]. Damaged vines can be pruned but grape yield

is reduced[58]. More research on long-term control methods for established populations is critical

for reducing SLF impact potential. First, trapping technologies that reduce bycatch must be

refined and widely deployed[57,59]. Second, biocontrol agents that specialize on SLF, such as

parasitoid wasps and fungus show promise, but more work is needed to understand non-target

attack rates [60–62]. Finally, like targeted mRNA vaccines developed to reduce the impact of SARS-

CoV-2 on human health[63], SLF-specific RNAi insecticides have the ability to control outbreaks

in vineyards and beyond[21,64]. Although SLF impacts to vineyards within its invaded range are

significant, to date, SLF has yet to invade a major viticultural area. Its actual impact on such

regions with larger, wealthier, and interconnected wine economies is thus unknown. It is also

unclear whether market elasticity might weaken or strengthen the disruption of a SLF

paninvasion to the global wine market. It behooves governments to heed the paninvasion risk of

this grapevine pest.

When a pest like SLF with high paninvasion risk emerges, coordinated governmental

efforts can mitigate global market disruptions. For example, the Great Wine Blight of the late

19th century caused by grapevine phylloxera (Hemiptera: *Daktulosphaira vitifoliae*) was the

largest shock to the global wine market ever recorded[65]. Phylloxera decimated European

vineyards, but the market recovered due to pest management solutions whose development were

coordinated by high-level officials in the French federal government[65].

For SLF in the U.S., early federal coordination was hampered. The U.S. National

Invasive Species Council (NISC) contains high-level federal officials (e.g., Secretaries of State,

Interior, Agriculture, Defense) who coordinate reduction of invasive species impact[66]. In 2019,

NISC funding was cut and the Invasive Species Advisory Committee (ISAC) was dissolved. The

ISAC comprised scientific experts who advised the NISC by producing memoranda and white

papers on emerging invasive species issues[67]. These actions decreased U.S. capacity to respond

to emerging paninvasive species[68]. In 2021, the ISAC was reinstated, and NISC's funding is

expected to be fully restored[69]. Based on our SLF risk assessment, we suggest the reinstated

ISAC produce memoranda and white papers on solutions for SLF as soon as possible. SLF risk

must be communicated to the NISC, who can mobilize the resources needed to reduce its impact.

The paninvasion severity assessment framework is a stakeholder communication tool to

assess if an invasive can cause market, environmental, and human-health disruptions at the

global scale (Fig. 1). It does so by equating pathogen transmission, infectivity, and virulence—

well-known to the public due to COVID-19 pandemic—with invasive species transport,

establishment, and impact potentials (Fig. 2). Paninvasion assessments produce accessible maps

(Fig. 3, 4), scatter plots (Fig. 5), and easy to understand risk values (Fig. 6). Going forward,

invasion potentials for other species are likely to increasingly align and coordinated

governmental efforts will be needed to reduce such potentials in the U.S. and internationally. It is

prudent that when an invasive species is found, the severity of its paninvasion risk be assessed.

Methods

Here, we provide a methodological discussion of the paninvasion severity assessment

framework (Fig. 2) and its application to spotted lanternfly (SLF, Fig. 1). To make the SLF

assessment easy to refine once new data and insights are available, we provide both an open-

source R package that includes all data to reproduce all results (https://ieco-lab.github.io/slfrsk/)

and a Google Earth Engine application to map SLF paninvasion severity from global to local

scales (https://ieco.users.earthengine.app/view/ieco-slf-riskmap). These open-science tools also

are adaptable to other emerging regional invasives at risk of paninvasion[e.g., 70]. Here, we focused

on agricultural and economic data most relevant to assess pest risk, but for non-pest invasives,

data on environmental and human health may be a higher priority.

Paninvasion Severity Assessment Framework

Although the invasion process can be divided into many stages, the paninvasion severity

assessment framework focuses on the three main stages most often estimated in invasion risk

assessments[2] and that are analogous to the disease potentials that public-health scientists quantify

for pathogens (Fig. 2)[8]. When a pathogen with pandemic risk emerges, public health scientists

place it within scaled measures of transmissibility and infectivity (often combined and termed

transmissibility), and virulence (clinical severity) to assess its risk[8,9]. For example, when SARS-

CoV-2 emerged during the COVID-19 pandemic, the initial understanding was that different age

groups had similar potentials to transmit and become infected (Fig. 2a, y-axis), but different age

groups varied in their clinical severity once infected (Fig. 2a, x-axis)[8,71,72]. To adapt this public-

health framework to invasion process theory[2,73–76], we equated transmission, infectivity, and

virulence potentials of a pathogen across different human populations to the transport,

establishment, and impact potentials of an invasive species across different regions (see colored

arrows between Fig. 2a and b). For example, in Fig. 2b we placed several hypothetical regions

that together indicate strong alignment (i.e., multivariate correlation) among invasion potentials

across the regions. In this example, predicted invasion risk (Fig. 2c, x-axis) for these three

hypothetical regions is strongly correlated to a measure of their contributions to a global market

(Fig. 2c, y-axis), indicating an overall high paninvasion risk.

Paninvasion assessments comprise four steps (Fig. 2d): 1) estimate invasion potentials, 2)

calculate alignment of invasion potentials, 3) quantify paninvasion risk, and 4) articulate caveats,

which we describe in detail for SLF below and in the Supplementary Methods.

Step 1: Estimate Invasion Potentials

Transport Potential

Transport potential is a measure of propagule pressure[77]. The prevailing hypothesis on

SLF transport potential is that regions that import more tonnage of commodities from the

invaded U.S. region also import more total tonnage of goods and trade infrastructure (e.g., cargo

containers, pallets, and railcars) that inadvertently transport SLF propagules, such as egg masses,

long-distances[18,21,26,34,78,79]. To estimate which states were invaded and identify SLF

transportation events, we obtained a database of SLF records from the USDA and aggregated

first-find and regulatory incident reports[e.g., 38]. As of December 2020, the invaded states were

Connecticut, Delaware, Maryland, New Jersey, New York, Ohio, Pennsylvania, Virginia, and

West Virginia (Fig. 3). We estimated transport potentials from the U.S. invaded region as the

$\log_{10}$ of the average annual metric total tonnage of all goods imported between 2012–2017 by

states and countries from the invaded U.S. states. This date range encapsulates both pre- and

post-introduction of SLF to the U.S. and maximized temporal overlap across different data

sources. Tonnage data from these invaded states were from the U.S. Freight Analysis Framework

for interstate imports[39] and from the U.S. Trade Online database for international imports[40], both

accessed on June 14, 2019. Current SLF spread in the U.S. was explained by our transport

potential metric suggesting that using total tonnage is a valid metric of transport potential for

SLF (Supplementary Methods, Supplementary Table 1).

Establishment Potential

Establishment potential is the set of species-specific and environmental characteristics of

a region that determine if a transported species can generate a self-sustaining population[2]. We

determined SLF establishment potential from an ensemble estimate from three global species

distribution models (SDMs): a multivariate SDM of TOH (*sdm_toh*), a multivariate SDM of SLF

(*sdm_slf1*), and a univariate SDM of SLF that modeled SLF presence on the predicted values

from *sdm_toh (sdm_slf2)*. Models were constructed using MaxEnt ver. 3.4.1 according to

unbiased niche modeling best practices (see Supplementary Methods)[80–82]. Specifically, our

SDMs were built from unique, error checked, and spatially rarefied presence records: *sdm_toh*

on 8,022 TOH presence records, and *sdm_slf1* and *sdm_slf2* on 325 SLF presence records

obtained from GBIF on October 20, 2020. To find the best-fit models that explained TOH and

SLF presences, we identified a subset of six covariates, from 22 candidate covariates[83–86], that

minimized model collinearity: annual mean temperature, mean diurnal temperature range, annual

precipitation, precipitation seasonality, elevation, and access to cities. We fit *sdm_toh* and

sdm_slf1 with these six covariates. *sdm_toh* represents our best estimate of the global distribution

of TOH, thus we fit *sdm_slf2* that modeled SLF suitability from the predicted values of *sdm_toh*.

As such, *sdm_slf2* represents suitability that considers a primary plant host (TOH)[25] that is also

invasive but likely not at equilibrium[87] and the same abiotic covariates as *sdm_slf1* (*sdm_toh*

uses the same covariates). We evaluated model performance with *k*-fold cross-validation,

specifically the receiver operating characteristic of the AUC (area under the curve) and omission

error[88–90].

281   Each model estimated suitability at a 30-arc-second (at the equator approximately 1 km^2)

global resolution with pixel values scaled 0–1, which we averaged across models per pixel to

produce one ensemble image and intersected this image with state and country polygons[82].

Establishment potential for the 50 U.S. states and 223 countries was estimated as the maximum

pixel value for each state and country. Results and conclusions with mean and median pixel

values instead of max were similar (see https://ieco-lab.github.io/slfrsk/).

Impact Potential

289   We used log$_{10}$-transformed average annual production tonnages of grapes and wine as

two separate estimates of impact potential. For consistency, we used grape and wine production

during the same span of time as transport potential estimates, 2012–2017. Grape production for

countries was from the Food and Agriculture Organization of the United Nations crop database[46]

and for states from the USDA National Agricultural and Statistics Service commodity

database[47], both accessed on January 24, 2020. Wine production in metric tons was from

FAOSTAT for countries[46], accessed on June 21, 2019, and from the Alcohol and Tobacco Tax

and Trade Bureau (TTB) for states[48], accessed on June 22, 2019, which was in gallons but we

converted it to metric tons assuming 3.776e-3 t/gallon[91]. Major viticultural regions (Fig. 4) were

aggregated and georeferenced from a TTB U.S. state data set[92] and the global viticultural regions

Wikipedia list[93] to better visualize impact within states and countries, both accessed on April 22,

2020.

Step 2: Calculate Alignment Correlations

To consider how all three invasion potentials may coincide for regions, we calculated

alignment correlations for states and countries separately. Alignment was calculated for each of

the two measures of impact potential as Spearman rank correlations between impact potential

and the predicted values from linear regressions models of each impact potential regressed on

transport and establishment potentials together[94]. We then visualized these multiple multivariate

correlations as quadrant plots following a stakeholder-friendly and approachable format adapted

from the pandemic severity assessment framework[8,9].

Step 3: Quantify Paninvasion Risk

To determine if invasion risk for countries corresponds with economic impact to the

global wine industry, we investigated the relationship between wine market size and predicted

risk of invasion for individual countries. We estimated wine market size for 223 countries

(including some that export but do not produce wine) as the value of wine exports corresponding

with the years for our trade data (2012–2017, $\log_{10}$ USD) downloaded from the FAOSTAT

detailed trade matrix[46], accessed August 31, 2020. Then, we regressed country grape production

on transport and establishment potentials with multiple linear regression. Each predicted value

from this regression can be considered an estimate of the risk of SLF to invade and impact a

country's grape production. We rescaled these estimates from 1–10 to create an easily interpreted

estimate of risk and then correlated these predicted values to wine export market size. To place

overall SLF paninvasion severity on a clear scale for both researchers and stakeholders, we

simply rescaled the Pearson correlation from 1–10, so that 1 is a complete negative correlation

and 10 is a complete positive correlation between country risk and wine export market size.

Step 4: Articulate Caveats

Paninvasion risk assessments should be performed iteratively as the invasion process

continues across regions and responses are mobilized. Early assessments of emergent pests have

great utility to support early responses but often come with caveats. The fourth step of the

framework is to articulate the caveats of a current assessment to guide future research. These

caveats should explicitly consider the assumptions made when estimating invasion potentials and

how those potentials and risk could change as the invasion process continues. Below we

articulate caveats of our SLF assessment to provide a basis for future refined assessments of SLF

to disrupt the global wine market.

We calculate paninvasion risk of SLF via stepping-stone transport from the eastern US.

However, major wine producing nations also heavily trade with China, Japan, and South Korea,

where SLF is also established. Total SLF transport potential is thus greater than our estimates,

meaning paninvasion risk is higher than what we report here. Future work should account for

global trade network dynamics to other nations with established SLF populations. However,

comprehensive surveys are first needed on the distribution of SLF in these other countries and

the trade emanating from regions with established populations. Further research should focus on

the identification of which industries and commodities are most likely to increase long-distance

spread. This requires linking trade dynamic models to phenological models to indicate if high

risk trade is occurring at the same time as egg laying, because eggs are the life-stage most likely

to be transported long-distances[21]. Additionally, refinement of SLF propagule pressure

dynamics[77], specifically propagule number and ratio of successful to failed transportation events,

can improve estimates of transport potential.

Establishment potential should be improved as additional data and models become

available. Because we use MaxEnt, a presence only, correlative SDM method to estimate

establishment, we measure suitability for SLF in a way that does not account for how SLF

population density and possible plastic and adaptive responses to novel environmental conditions

in invaded regions may affect establishment success. Omission of population density and other

demographic variables can hinder accurate prediction, especially for SDMs[95], and whenever

possible, priority should be placed on using them alongside models that rely on pest physiology

to predict establishment potential[96]. For SLF, two physiologically based models[44,45] largely

correspond with our SDM-based establishment potentials and thus support the global

establishment potential for SLF we report here. Early assessments of paninvasion severity are

unlikely to account for plasticity and adaptation that is common for invasive pests, especially

when combined with variation in local weather patterns and climate change. Indeed, a recent

analysis suggests that SLF will experience increased suitable habitat and a greater impact in

China in the future due to climate change[97]. The expected effect of climate change is likely more

complicated for SLF, which has a flexible life cycle that can include but does not require,

temperature-linked diapause for overwintering in cooler regions[24,98,99]. Survivorship appears

greater without such diapause[98], and thus establishment potential may be even higher than

expected in warmer climes.

Variation in weather, climate change, host preference, and pest density can influence pest

impacts. Such factors often act at different scales and in a spatially heterogenous manner. For

example, SLF prefers grapes, but the degree to which it does over alternative hosts near

vineyards remains poorly known, which is important, since SLF appear to have their highest

densities at vineyard edge habitats[100]. The vulnerability of viticultural regions may be affected by

the prevalence of particular grape cultivars or alternative hosts, but additional research is

necessary to elucidate SLF feeding presence. Similarly, the relationship of SLF density within a

vineyard and density in the surrounding landscape remains poorly known, which is in turn likely

to be influenced by weather patterns and plant phenology[36]. As SLF host preference and its

relationship to landscape variables become better understood, they should be incorporated into

considerations of impact potential.

Lastly, to refine the SLF paninvasion risk assessment, future work should calculate

invasion potentials for other grape pests like phylloxera to place SLF on an absolute scale of risk

severity[101]. Our assessment of SLF relativizes invasion potentials with the assumption that

regions with high potentials relative to other regions also have high absolute potentials. SLF has

broad environmental suitability, a flexible life cycle, ability to lay many discrete egg masses on

numerous substrates (Fig. 1b), observed rapid spread (Fig. 3) and realized impacts to grape and

wine production[21–26,36], so its absolute potentials are likely very high. However, absolute

potentials can only be assessed by comparing multiple paninvasive species, like what is done for

pathogens. When a pathogen with pandemic potential emerges, the pandemic severity

assessment framework compares the severity of the potentials of the current outbreak pathogen

to past pandemic producing pathogens[8,9,71]. The next step towards a mature paninvasion

framework is to estimate invasion potentials for current paninvasive species, so that the

likelihood of a paninvasion for any emerging regional pest can be placed on an absolute scale of

severity.

Data availability

All datasets generated during and/or analyzed in this study are available as described in

the methods and research compendium (https://ieco-lab.github.io/slfrsk) or can be obtained as a

part of the companion R package (https://github.com/ieco-lab/slfrsk). Cumbersome datasets (e.g.,

SDM files) are available in a Data Dryad repository (https://doi.org/10.5061/dryad.msbcc2g1b).

Code availability

All code written and used for this study are available as an R package that is publicly

available at: https://github.com/ieco-lab/slfrsk.

References

1. Ristaino, J. B. *et al.* The persistent threat of emerging plant disease pandemics to global food

security. *Proc. Natl. Acad. Sci.* **118**, (2021).

2. Blackburn, T. M. *et al.* A proposed unified framework for biological invasions. *Trends Ecol.*

Evol. **26**, 333–339 (2011).

3. Chapman, D., Purse, B. V., Roy, H. E. & Bullock, J. M. Global trade networks determine the

distribution of invasive non-native species. *Glob. Ecol. Biogeogr.* **26**, 907–917 (2017).

4. Liebhold, A. M. *et al.* Plant diversity drives global patterns of insect invasions. *Sci. Rep.* **8**,

1–5 (2018).

5. Bradshaw, C. J. A. *et al.* Massive yet grossly underestimated global costs of invasive insects.

Nat. Commun. **7**, 1–8 (2016).

6. Wyckhuys, K. A. G. *et al.* Biological control of an invasive pest eases pressures on global

commodity markets. *Environ. Res. Lett.* **13**, 094005 (2018).

7. Leung, B., Finnoff, D., Shogren, J. F. & Lodge, D. Managing invasive species: Rules of

thumb for rapid assessment. *Ecol. Econ.* **55**, 24–36 (2005).

8. Reed, C. *et al.* Novel framework for assessing epidemiologic effects of influenza epidemics

and pandemics. *Emerg. Infect. Dis.* **19**, 85 (2013).

9. Qualls, N. *et al.* Community mitigation guidelines to prevent pandemic influenza—United

States, 2017. *MMWR Recomm. Rep.* **66**, 1 (2017).

10. Grarock, K., Lindenmayer, D. B., Wood, J. T. & Tidemann, C. R. Using invasion process

theory to enhance the understanding and management of introduced species: A case study

reconstructing the invasion sequence of the common myna (*Acridotheres tristis*). *J. Environ.*

Manage. **129**, 398–409 (2013).

11. Nuñez, M. A., Pauchard, A. & Ricciardi, A. Invasion science and the global spread of SARS-

CoV-2. *Trends Ecol. Evol.* **35**, 642–645 (2020).

12. Ogden, N. H. *et al.* Emerging infectious diseases and biological invasions: A call for a one

health collaboration in science and management. *R. Soc. Open Sci.* **6**, 181577 (2019).

13. Hatcher, M. J., Dick, J. T. A. & Dunn, A. M. Disease emergence and invasions. *J. Ecol.* **26**,

1275–1287 (2016).

14. Bright, C. Invasive species: Pathogens of globalization. *Foreign Policy* **1**, 50–64 (1999).

15. Simberloff, D., Meyerson, L. & Fefferman, N. Invasive species policy and COVID-19.

(2020).

16. Comizzoli, P., Pagenkopp Lohan, K. M., Muletz-Wolz, C., Hassell, J. & Coyle, B. The

interconnected health initiative: A Smithsonian framework to extend one health research and

education. *Front. Vet. Sci.* **8**, (2021).

17. Katella, K. Our new COVID-19 vocabulary—what does it all mean? > Stories at Yale

Medicine. *Yale Medicine* https://www.yalemedicine.org/stories/covid-19-glossary/ (2020).

18. Parra, G., Moylett, H. & Bulluck, R. USDA-APHIS-PPQ-CPHST Technical working group

summary report spotted lanternfly, *Lycorma delicatula* (White, 1845). (2018).

19. Floerl, O., Inglis, G. J., Dey, K. & Smith, A. The importance of transport hubs in stepping-

stone invasions. *J. Appl. Ecol.* **46**, 37–45 (2009).

20. Barringer, L. E., Donovall, L. R., Spichiger, S.-E., Lynch, D. & Henry, D. The first New

World record of *Lycorma delicatula* (Insecta: Hemiptera: Fulgoridae). *Entomol. News* **125**,

20–23 (2015).

21. Urban, J. M. Perspective: Shedding light on spotted lanternfly impacts in the USA. *Pest*

Manag. Sci. **76**, 10–17 (2020).

22. Nixon, L. J. *et al.* Survivorship and development of the invasive *Lycorma delicatula*

(Hemiptera: Fulgoridae) on wild and cultivated temperate host plants. *Environ. Entomol.*

nvab137 (2021) doi:10.1093/ee/nvab137.

23. Urban, J. M., Calvin, D. & Hills-Stevenson, J. Early response (2018–2020) to the threat of

spotted lanternfly, *Lycorma delicatula* (Hemiptera: Fulgoridae) in Pennsylvania. *Ann.*

Entomol. Soc. Am. **114**, 709–718 (2021).

24. Du, Z. *et al.* Global phylogeography and invasion history of the spotted lanternfly revealed

by mitochondrial phylogenomics. *Evol. Appl.* (2021) doi:https://doi.org/10.1111/eva.13170.

25. Lee, J.-E. *et al.* Feeding behavior of *Lycorma delicatula* (Hemiptera: Fulgoridae) and

response on feeding stimulants of some plants. *Korean J. Appl. Entomol.* **48**, 467–477

(2009).

26. Lee, D.-H., Park, Y.-L. & Leskey, T. C. A review of biology and management of *Lycorma*

delicatula (Hemiptera: Fulgoridae), an emerging global invasive species. *J. Asia-Pac.*

Entomol. **22**, 589–596 (2019).

27. Roush, R. How we can contain the spotted lanternfly — maybe the worst invasive pest in

generations | Opinion. *https://www.inquirer.com* (2018).

28. Imbler, S. The dreaded lanternfly, scourge of agriculture, spreads in New Jersey. *The New*

York Times (2020).

29. Morrison, R. Invasive insects: The top 4 'most wanted' list. *Entomology Today*

https://entomologytoday.org/2018/06/21/invasive-insects-the-top-4-most-wanted-list/ (2018).

30. Murman, K. *et al.* Distribution, survival, and development of spotted lanternfly on host

plants found in North America. *Environ. Entomol.* **49**, 1270–1281 (2020).

31. Derstine, N. T. *et al.* Plant volatiles help mediate host plant selection and attraction of the

spotted lanternfly (Hemiptera: Fulgoridae): A generalist with a preferred host. *Environ.*

Entomol. **49**, 1049–1062 (2020).

32. Dechaine, A. C. *et al.* Phenology of *Lycorma delicatula* (Hemiptera: Fulgoridae) in Virginia,

USA. *Environ. Entomol.* nvab107 (2021) doi:10.1093/ee/nvab107.

33. Uyi, O. *et al.* Spotted lanternfly (Hemiptera: Fulgoridae) can complete development and

reproduce without access to the Ppreferred host, *Ailanthus altissima*. *Environ. Entomol.*

(2020) doi:10.1093/ee/nvaa083.

34. Park, M., Kim, K.-S. & Lee, J.-H. Genetic structure of *Lycorma delicatula* (Hemiptera:

Fulgoridae) populations in Korea: Implication for invasion processes in heterogeneous

landscapes. *Bull. Entomol. Res.* **103**, 414–424 (2013).

35. Dara, S. K., Barringer, L. & Arthurs, S. P. *Lycorma delicatula* (Hemiptera: Fulgoridae): A

new invasive pest in the United States. *J. Integr. Pest Manag.* **6**, 1–6 (2015).

36. Leach, H. & Leach, A. Seasonal phenology and activity of spotted lanternfly (*Lycorma*

delicatula) in Eastern U.S. vineyards. *J. Pest Sci.* **93**, 1215–1224 (2020).

37. International Organisation of Vine and Wine. *2019 Statistical Report on World*

Vitiviniculture. 23 (2019).

38. California Department of Food and Agriculture. *Pest detection advisory no. PD17-2020*

spotted lanternfly PD/EP activity summary 2020. 1–7 (2020).

39. Oak Ridge National Lab. Freight analysis framework version 4. http://faf.ornl.gov/fafweb/

(2017).

40. U.S. Census Bureau. U.S.A. Trade Online. https://usatrade.census.gov/index.php?do=login

(2019).

41. Derived dataset GBIF.org. Filtered export of GBIF occurrence data. (2021)

doi:10.15468/DD.KS6ACS.

42. Jung, J.-M., Jung, S., Byeon, D. & Lee, W.-H. Model-based prediction of potential

distribution of the invasive insect pest, spotted lanternfly *Lycorma delicatula* (Hemiptera:

Fulgoridae), by using CLIMEX. *J. Asia-Pac. Biodivers.* **10**, 532–538 (2017).

43. Wakie, T. T., Neven, L. G., Yee, W. L. & Lu, Z. The establishment risk of *Lycorma*

delicatula (Hemiptera: Fulgoridae) in the United States and globally. *J. Econ. Entomol.* **113**,

306–314 (2020).

44. Lewkiewicz, S. M., De Bona, S., Helmus, M. R. & Seibold, B. Temperature sensitivity of

pest reproductive numbers in age-structured PDE models, with a focus on the invasive

spotted lanternfly. *ArXiv211211448 Q-Bio* (2021).

45. Maino, J. L., Schouten, R., Lye, J. C., Umina, P. A. & Reynolds, O. L. Mapping the life-

history, development, and survival of spotted lantern fly in occupied and uninvaded ranges.

InReview 1–18 (2021) doi:10.21203/rs.3.rs-400798/v1.

46. FAOSTAT. FAOSTAT statistical database. http://www.fao.org/faostat/en/#data/QC (2019).

47. USDA National Agricultural Statistics Service. National agricultural statistics service - quick

stats. https://quickstats.nass.usda.gov/ (2019).

48. U.S. Alcohol and Tobacco Tax and Trade Bureau. Wine statistics.

https://www.ttb.gov/wine/wine-stats.shtml (2019).

49. Crowe, J. Spotted lanternfly control program in the Mid-Atlantic region environmental

assessment. *USDA APHIS Rep.* 46 (2018).

50. US Animal and Plant Health Inspection Service. USDA provides $7.1 million to

Pennsylvania to support projects that protect agriculture and natural resources.

https://www.aphis.usda.gov/wcm/connect/APHIS_Content_Library/SA_Newsroom/SA_Ne

ws/SA_By_Date/SA-2019/pennsylvania-

funding?presentationtemplate=APHIS_Design_Library%2FPT_Print_Friendly_News_releas

e (2019).

51. Jones, C. M. *et al.* Iteratively forecasting biological invasions with PoPS and a little help

from our friends. *Front. Ecol. Environ.* (2021) doi:10.1002/fee.2357.

52. Smyers, E. C. *et al.* Spatio-Temporal Model for Predicting Spring Hatch of the Spotted

Lanternfly (Hemiptera: Fulgoridae). *Environ. Entomol.* **50**, 126–137 (2021).

53. Brooks, R. K., Wickert, K. L., Baudoin, A., Kasson, M. T. & Salom, S. Field-inoculated

Ailanthus altissima stands reveal the biological control potential of *Verticillium nonalfalfae*

in the Mid-Atlantic region of the United States. *Biol. Control* **148**, 104298 (2020).

54. Commonwealth of Pennsylvania. Pennsylvania Bulletin. **49**, 2705–2902 (2019).

55. Barringer, L. & Ciafré, C. M. Worldwide feeding host plants of spotted lanternfly, with

significant additions from North America. *Environ. Entomol.* **49**, 999–1011 (2020).

56. Leach, H., Biddinger, D. J., Krawczyk, G., Smyers, E. & Urban, J. M. Evaluation of

insecticides for control of the spotted lanternfly, *Lycorma delicatula*, (Hemiptera:

Fulgoridae), a new pest of fruit in the Northeastern U.S. *Crop Prot.* **124**, 104833 (2019).

57. Francese, J. A. *et al.* Developing traps for the spotted lanternfly, *Lycorma delicatula*

(Hemiptera: Fulgoridae). *Environ. Entomol.* **49**, 269–276 (2020).

58. Penn State Extension. Spotted lanternfly management in vineyards.

https://extension.psu.edu/spotted-lanternfly-management-in-vineyards (2021).

59. Nixon, L. J. *et al.* Development of Behaviorally Based Monitoring and Biosurveillance Tools

for the Invasive Spotted Lanternfly (Hemiptera: Fulgoridae). *Environ. Entomol.* **49**, 1117–

1126 (2020).

60. Liu, H. & Mottern, J. An old remedy for a new problem? Identification of *Ooencyrtus*

kuvanae (Hymenoptera: Encyrtidae), an egg parasitoid of *Lycorma delicatula* (Hemiptera:

Fulgoridae) in North America. *J. Insect Sci.* **17**, 1–6 (2017).

61. Yang, Z.-Q., Choi, W.-Y., Cao, L.-M., Wang, X.-Y. & Hou, Z.-R. A new species of

Anastatus (Hymenoptera: Eulpelmidae) from China, parasitizing eggs of *Lycorma delicatula*

(Homoptera: Fulgoridae). *Zool. Syst.* **40**, 290–302 (2015).

62. Clifton, E. H. *et al.* Applications of *Beauveria bassiana* (Hypocreales: Cordycipitaceae) to

control populations of spotted lanternfly (Hemiptera: Fulgoridae), in semi-natural landscapes

and on grapevines. *Environ. Entomol.* **49**, 854–864 (2020).

63. Hogan, M. J. & Pardi, N. mRNA Vaccines in the COVID-19 Pandemic and Beyond. *Annu.*

Rev. Med. **73**, 17–39 (2022).

64. Whyard, S., Singh, A. D. & Wong, S. Ingested double-stranded RNAs can act as species-

specific insecticides. *Insect Biochem. Mol. Biol.* **39**, 824–832 (2009).

65. Ordish, G. *The great wine blight*. (Charles Scribner's Sons, 1972).

66. About the Council. https://www.doi.gov/invasivespecies/about-nisc (2016).

67. Invasive Species Advisory Committee Products. https://www.doi.gov/invasivespecies/isac-

resources (2015).

68. Simberloff, D. *et al.* U.S. action lowers barriers to invasive species. *Science* **367**, 636–636

(2020).

69. Exec. Order No. 14048, A. of J. R. B., Jr. *Executive Order on Continuance or*

Reestablishment of Certain Federal Advisory Committees and Amendments to Other

Executive Orders. (2021).

70. Zhu, G., Illan, J. G., Looney, C. & Crowder, D. W. Assessing the ecological niche and

invasion potential of the Asian giant hornet. *Proc. Natl. Acad. Sci.* **117**, 24646–24648

(2020).

71. Freitas, A. R. R. *et al.* Assessing the severity of COVID-19. *Epidemiol. E Serviços Saúde* **29**,

1–5 (2020).

72. Prevent Epidemics. *COVID-19 Key COVID-19 metrics based on the latest available science*.

https://preventepidemics.org/covid19/science/weekly-science-review/ (2020).

73. Lockwood, J. L., Hoopes, M. F. & Marchetti, M. P. *Invasion ecology*. (Wiley-Blackwell,

2013).

74. Ehler, L. E. Invasion biology and biological control. *Biol. Control* **13**, 127–133 (1998).

75. Ludsin, S. A. & Wolfe, A. D. Biological invasion theory: Darwin's contributions from *The*

Origin of Species. *BioScience* **51**, 780 (2001).

76. Schulz, A. N., Lucardi, R. D. & Marsico, T. D. Strengthening the ties that bind: An

evaluation of cross-disciplinary communication between invasion ecologists and biological

control researchers in entomology. *Ann. Entomol. Soc. Am.* **114**, 163–174 (2021).

77. Lockwood, J. L., Cassey, P. & Blackburn, T. The role of propagule pressure in explaining

species invasions. *Trends Ecol. Evol.* **20**, 223–228 (2005).

78. Liu, H. Oviposition substrate selection, egg mass characteristics, host preference, and life

history of the spotted lanternfly (Hemiptera: Fulgoridae) in North America. *Environ.*

Entomol. **48**, 1452–1468 (2019).

79. Liu, H. Seasonal development, cumulative growing degree-days, and population density of

spotted lanternfly (Hemiptera: Fulgoridae) on selected hosts and substrates. *Environ.*

Entomol. **49**, 1171–1184 (2020).

80. Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E. & Blair, M. E. Opening the black

box: An open-source release of Maxent. *Ecography* **40**, 887–893 (2017).

81. Araújo, M. B. & New, M. Ensemble forecasting of species distributions. *Trends Ecol. Evol.*

22, 42–47 (2007).

82. Araújo, M. B. *et al.* Standards for distribution models in biodiversity assessments. *Sci. Adv.*

5, eaat4858 (2019).

83. Fick, S. E. & Hijmans, R. J. WorldClim 2: New 1-km spatial resolution climate surfaces for

global land areas. *Int. J. Climatol.* **37**, 4302–4315 (2017).

84. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution

interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**, 1965–1978 (2005).

85. Weiss, D. J. *et al.* A global map of travel time to cities to assess inequalities in accessibility

in 2015. *Nature* **553**, 333–336 (2018).

86. Simard, M., Pinto, N., Fisher, J. B. & Baccini, A. Mapping forest canopy height globally

with spaceborne lidar. *J. Geophys. Res. Biogeosciences* **116**, (2011).

87. Sladonja, B., Sušek, M. & Guillermic, J. Review on Invasive Tree of Heaven (Ailanthus

altissima (Mill.) Swingle) Conflicting Values: Assessment of Its Ecosystem Services and

Potential Biological Threat. *Environ. Manage.* **56**, 1009–1034 (2015).

88. Fielding, A. H. & Bell, J. F. A review of methods for the assessment of prediction errors in

conservation presence/absence models. *Environ. Conserv.* **24**, 38–49 (1997).

89. Pearson, R. G., Raxworthy, C. J., Nakamura, M. & Peterson, A. T. Predicting species

distributions from small numbers of occurrence records: A test case using cryptic geckos in

Madagascar. *J. Biogeogr.* **34**, 102–117 (2007).

90. Anderson, R. P. & Gonzalez, I. Species-specific tuning increases robustness to sampling bias

in models of species distributions: An implementation with Maxent. *Ecol. Model.* **222**, 2796–

2811 (2011).

91. AVCALC. Density of alcoholic beverage, wine, table, all (food). https://www.aqua-

calc.com/page/density-table/substance/alcoholic-blank-beverage-coma-and-blank-wine-

coma-and-blank-table-coma-and-blank-all (2019).

92. U.S. Alcohol and Tobacco Tax and Trade Bureau. Established AVAs.

https://www.ttb.gov/wine/established-avas (2019).

93. Wikipedia. https://en.wikipedia.org/wiki/List_of_wine-producing_regions. (2020).

94. Allison, P. D. *Multiple regression: A primer*. (Pine Forge Press, 1999).

95. Ponti, L. *et al.* Biological invasion risk assessment of *Tuta absoluta*: Mechanistic versus

correlative methods. *Biol. Invasions* **23**, 3809–3829 (2021).

96. Briscoe, N. J. *et al.* Forecasting species range dynamics with process-explicit models: Matching methods to applications. *Ecol. Lett.* **22**, 1940–1956 (2019).

97. Wang, C.-J. *et al.* Risk assessment of insect pest expansion in alpine ecosystems under climate change. *Pest Manag. Sci.* **77**, 3165–3178 (2021).

98. Keena, M. A. & Nielsen, A. L. Comparison of the hatch of newly laid *Lycorma delicatula* (Hemiptera: Fulgoridae) eggs from the United States after exposure to different temperatures and durations of low temperature. *Environ. Entomol.* 1–8 (2021) doi:10.1093/ee/nvaa177.

99. Xin, B. *et al.* Exploratory survey of spotted lanternfly (Hemiptera: Fulgoridae) and its natural enemies in China. *Environ. Entomol.* **50**, 36–45 (2020).

100. Leach, A. & Leach, H. Characterizing the spatial distributions of spotted lanternfly (Hemiptera: Fulgoridae) in Pennsylvania vineyards. *Sci. Rep.* **10**, 1–9 (2020).

101. Granett, J., Walker, M. A., Kocsis, L. & Omer, A. D. Biology and management of grape phylloxera. *Annu. Rev. Entomol.* **46**, 387–412 (2001).

Acknowledgments

We thank Julie Urban, Heather Leach, Nadège Bélouard, Stefani Cannon, Seba De Bona, Jason Gleditsch, Stephanie Lewkiewicz, Victoria Ramirez, Payton Phillips, Timothy Swartz and the Integrative Ecology Lab at Temple University for comments and feedback on earlier drafts of this manuscript. This work was funded by the United States Department of Agriculture Animal and Plant Health Inspection Service Plant Protection and Quarantine under Cooperative Agreements AP19PPQS&T00C251 and AP20PPQS&T00C136; the United States Department of Agriculture National Institute of Food and Agriculture Specialty Crop Research Initiative Coordinated Agricultural Project Award 2019-51181-30014; and the Pennsylvania Department of Agriculture under agreements 44176768, 44187342, and C9400000036.

Competing interests

645   The authors declare no competing interests.

Author Contributions

648   All authors conceived and designed the study. NAH acquired and analyzed data. All authors
649   contributed materials/analysis tools and wrote the paper.

650
Figures and Tables

Fig. 1: Our concept of paninvasive species is based on invasion process theory. Any species
has potentials to complete three sequential stages (depicted by the arrow) to become invasive in a
region (i.e., transport to the region, establishment in the region, impact to the region's
economy)[2]. For a pest paninvasion to occur, a pest must be transported and establish in suitable
regions globally where it impacts susceptible agriculture disrupting markets at a global scale
(depicted by the bracket). Here, we focus on estimating invasion stage potentials for the spotted
lanternfly planthopper (*Lycorma delicatula*, SLF), an emerging U.S. pest at risk of disrupting the
global wine market. Photos introduce SLF biology. Gravid females (**a**) lay eggs on many
surfaces like stone (**b**) and transport infrastructure (**c**). Once eggs hatch, SLF develop by molting
through three black and one red nymphal instars (**d**) before molting into winged adults (top, **a**).
SLF feed on phloem of many plants but develop quickly on tree-of-heaven (*Ailanthus altissima*,
TOH, **e, f**), a paninvasive tree commonly found in habitat fragments around railroads (**f**) and
warehouses (**c**). SLF also heavily feed on grape (**g**), reducing yield (**h**) and contributing to vine
death (**i**). Photo credits: S. Cannon (**e**), M. Helmus (top, **a, d**), H. Leach (b, **c, g–i**), G. Parra (**f**).
The map is of SLF (orange) and TOH (blue) presences (ca. 2020) we used to estimate
paninvasion risk (see Methods).

Fig. 2: The paninvasion severity assessment framework is adapted from the U.S. CDC pandemic severity assessment framework used to estimate the risk of emerging human pathogens. For pandemics (**a**), quadrant plots of pathogen transmissibility and infectivity (combined on one axis) vs. pathogen virulence (clinical severity) are used to compare the risk of a pathogen across different populations or age groups[8,9,11]. For paninvasions (**b**), invasion potentials for an emerging regional invasive species are estimated (**d** Step 1) by equating pathogen transmission with transport potential, infectivity with establishment potential, and virulence with impact potential (follow the arrows) across regions (black circles) to construct quadrant plots that depict their alignment based on multivariate correlations (**d** Step 2; see Methods). Next, paninvasion risk (**c**) is estimated from the correlation between regional invasion risk estimated from the multivariate regression of invasion potentials (**d** Step 3; see Methods) and the size of regional markets that could be disrupted. The steps of the paninvasive severity assessment framework (**d**), culminating by articulating caveats in the current assessment that direct future research (**d** Step 4) that provide data to inform the next assessment iteration.

Fig. 3: Spotted lanternfly (*Lycorma delicatula*, SLF) had invaded nine states in the U.S. by 2020. Colored polygons are counties with established SLF populations. Grey-filled polygons are counties where SLF have been transported but have not established. These regulatory incidents include any egg cases, living, moribund, and dead individuals found in cargo. Red outlined states have established SLF populations.

Fig. 4: There is strong spatial alignment of spotted lanternfly (*Lycorma delicatula*) transport, establishment, and impact potentials. Arrows point from the U.S. invaded region to the top ten states (**a**) and countries (**b**) with the highest transport potentials. Purple points are locations of important viticultural areas.

Fig. 5: Spotted lanternfly (*Lycorma delicatula*) invasion-potentials are aligned for wine and

grape producing regions. States (**a**) and countries (**b**) that produce most of the global supply of

grapes (point size) and wine (point fill color) also have high transport and establishment

potentials. Invaded regions and the top-10 grape producing regions are labeled; box plots split

the distributions of establishment potential and transport potential into high (purple) and low

(white) wine producing regions (center line is the median, box limits are the upper and lower

quartiles, and whiskers are 1.5x the interquartile range); dashed lines divide the data into high-

high, high-low, and low-low potential quadrants; and axes are scaled and formatted as suggested

by the pandemic severity assessment framework[8,9]. Point color indicates high and low wine

producing regions. Point and label size depict wine production in metric tons.

709
710

Fig. 6: Spotted lanternfly (*Lycorma delicatula*) paninvasion risk is an 8 out of 10 due to the correlation between country wine export market size and invasion risk. Dashed lines are one-to-one guidelines at -0.5, 0, 0.5 intercepts. Point color indicates high and low wine producing regions. Point and label size depict wine production in metric tons.