Comparison of Neutralizing Antibody Titers Elicited by mRNA and

Adenoviral Vector Vaccine against SARS-CoV-2 Variants

Takuya Tada[1,4], Hao Zhou[1,4], Marie I. Samanovic[2,3,4], Belinda M. Dcosta[1], Amber

Cornelius[2,3], Mark J. Mulligan[1,2,3,5] and Nathaniel R. Landau[1,5][**]

Affiliation:

[1]Department of Microbiology, NYU Grossman School of Medicine, New York, NY, USA.

[2]Department of Medicine, NYU Grossman School of Medicine, New York, NY, USA.

[3]NYU Langone Vaccine Center, NYU Grossman School of Medicine, New York, NY, USA.

[4]Contributed equally to this study

[5]Contributed equally to this study

[**]Corresponding author:

Nathaniel R. Landau, Ph.D.

NYU Grossman School of Medicine

430 East 29th Street, Alexandria West Building, Rm 509, New York, NY 10016

Email: nathaniel.landau@med.nyu.edu

Phone: (212) 263-9197

Abstract

The increasing prevalence of SARS-CoV-2 variants has raised concerns regarding possible decreases in vaccine efficacy. Here, neutralizing antibody titers elicited by mRNA-based and an adenoviral vector-based vaccine against variant pseudotyped viruses were compared. BNT162b2 and mRNA-1273-elicited antibodies showed modest neutralization resistance against Beta, Delta, Delta plus and Lambda variants whereas Ad26.COV2.S-elicited antibodies from a significant fraction of vaccinated individuals were of low neutralizing titer (IC_{50} <50). The data underscore the importance of surveillance for breakthrough infections that result in severe COVID-19 and suggest the benefit of a second immunization following Ad26.COV2.S to increase protection against the variants.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) vaccines from two

vaccine platforms have been granted U.S. Food and Drug Administration (FDA)

Emergency Use Authorization: mRNA-based (Pfizer and Moderna) and adenoviral

vector-based (Johnson & Johnson (J&J)), all of which have been shown to be highly

effective. The mRNA-based vaccines were 94-95% effective in preventing COVID-19[1]

whereas the adenoviral vector-based J&J vaccine had 66.9% efficacy in preventing

moderate to severe disease[2]. However, the ongoing emergence of highly transmissible

variants with mutations in the spike protein raises concerns regarding possible decreases

in vaccine effectiveness due to spike protein antigenic variability.

SARS-CoV-2 variants have been classified by the World Health Organization (WHO)

based on increased transmissibility and/or pathogenicity as variants of concern (VOC;

Alpha (B.1.1.7), Beta (B.1.351), Gamma (B.1.1.248) and Delta (B.1.617.2) and variants

of interest (VOI; Epsilon (B.1.427/B.1.429), Iota (B.1.526), and Delta plus (AY.1) and

Lambda (C.37)[3]. The increased transmissibility and/or pathogenicity of the variants is due,

at least in part, to mutations in the spike protein RBD that increase its affinity for ACE2

on target cells. Mutations in the Beta, Gamma and Delta variant spike RBDs have been

shown to cause partial resistance to neutralization by the serum antibodies of vaccinated

and convalescent individuals and therapeutic monoclonal antibodies[4-11].

This study compared the neutralization titers of serum antibodies from individuals

immunized with three U.S. FDA Emergency use authorization vaccines (BNT162b2,

mRNA-1273 and Ad26.COV2.S) against viruses with the VOC and Lambda spike proteins.

The study groups were controlled for age, clinical co-morbidity, history of pre-vaccination

infection and sera were collected on similar days post-vaccination. The results

demonstrate a high level of cross-neutralization by antibodies elicited by BNT162b2 and

mRNA-1273 on the variants but significantly decreased neutralization by those elicited by

the single dose Ad26.COV2.S.

Variant pseudotyped lentiviruses. The Delta plus spike contains K417N, L452R and

T478K in the RBD **(Figure S1A)**. The Lambda spike protein contains novel L452Q and

F490S mutations in the RBD **(Figure S1A)**. We previously described the production of

lentiviruses pseudotyped by the Alpha, Beta, Gamma and Delta spike proteins and here

report the generation of pseudotypes with the Delta plus and Lambda variant spike

proteins and the individual constituent mutations. The variant spike proteins were well

expressed, proteolytically processed and incorporated into lentiviral virions at a level

similar to that of the parental D614G spike protein in the producer cells and virions **(Figure**

S1B). The measurement of neutralizing antibody titers with such pseudotypes has been

shown to yield results consistent with those obtained with the live virus plaque reduction

neutralization test[12].

Reduced sensitivity of virus with variant spikes to neutralization by convalescent

sera and mRNA vaccine-elicited antibodies. Sera from individuals who had been

infected prior to the emergence of the variants (collected 32-57 days post symptom onset)

neutralized virus with the D614G spike protein with an average IC_{50} titer of 346 and

neutralized the Alpha variant with a similar titer (IC_{50} of 305). Neutralizing titers for Beta,

Delta, Delta plus and Lambda variants were decreased 3.2-4.9-fold relative to D614G,

indicative of a modest resistance to neutralization **(Figure 1A, Table S1)**. The sera of

individuals vaccinated with BNT162b2 and mRNA-1273 that were collected 7-days post-

second injection – a peak antibody response timepoint - neutralized virus with the D614G

spike with significantly higher titer (1835 and 1594, respectively) relative to the

convalescent sera, and the antibodies cross-reacted on the variants with a modest 2.5-

4.0-fold decrease in titer **(Figure 1A)**. The resistance of the Beta variant was attributed

to the E484K mutation whereas resistance of the Delta variant was attributed to the L452R

mutation **(Figure S2)**. The resistance of the lambda variant was attributed to both the

L452Q and F490S mutations **(Figure S2).**

Resistance of viruses with variant spike proteins to neutralization by Ad26.COV2.S-

elicited antibodies. We next compared the neutralizing titers of antibodies elicited by the

BNT162b2 and mRNA-1273 mRNA vaccines with that of the Ad26.COV2.S adenoviral

vector-based vaccine. The sera analyzed were collected from individuals at similar time-

points post-final injection, on average (90 days for BNT162b2, 80 days for mRNA-1273

and 82 days for Ad26.COV2.S; **Table S2**) and from individuals of similar age and with

similar clinical co-morbidities **(Table S2)**. None of the participants had a history of COVID-

19 pre- or post-vaccination and all were negative for antibodies against the SARS-CoV-

2 N protein **(Table S2)**. The results showed that BNT162b2 sera neutralized virus with

the D614G and Alpha spikes with an average titer of 695 and 626. Compared to the

D614G, the neutralizing titer against Beta was decreased 6.1-fold and Delta plus was

decreased 2.7-fold. Results for the mRNA-1273 vaccine were similar with a 3.3-fold

decrease in neutralizing titer for Delta plus and 4.6-fold for Beta. Ad26.COV2.S sera

neutralized D614G and Alpha variants with average IC_{50} titers of 221 and 232,

respectively, and neutralized the variants with titers that were decreased by 5.4-fold for

Delta plus to 6.7-fold for the Beta variant as compared to D614G **(Figure 1B).**

Presentation of the data grouped by variant shows the decreased neutralizing titers

against the variants by sera of the Ad26.COV2.S-vaccinated individuals **(Figure 1C)**.

The L452R/Q mutation of the Delta plus and Lambda spike proteins increases

infectivity and affinity for ACE2. Measurement of the infectivity of the pseudotyped

viruses, normalized for particle number, showed that the Lambda variant spike protein

increased viral infectivity by 2-fold **(Figure 2A)**, an increase equivalent to that of the Delta

and Delta plus variants. The increase was due to the L452Q mutation and was similar to

that of the L452R found in the Delta and Delta plus variants. The other mutations (Δ246-

252, G75V-T76I, F490S and T859N) had no significant effect on infectivity **(Figure 2A)**.

Measurement of the relative affinity of the variant spike proteins for ACE2 using sACE2

neutralization assay showed that variant spikes had a 3-fold increase in sACE2 binding

(Figure 2B). This increase was confirmed in a virion:ACE2 binding assay **(Figure 2C)**.

The increase was caused by the L452R and L452Q mutation and were similar to the

increase caused by the N501Y mutation[13,14].

Neutralization by REGN10933 and REGN10987. Analysis of REGN10933 and

REGN10987 monoclonal antibodies that constitute the REGN-COV2 therapy showed that

REGN10933 had decreased activity against the Beta variant spike which resulted in a

127-fold decrease in neutralizing titer. REGN10933 also had decreased activity against

the Delta plus variant which resulted in a 92.7-fold decrease in neutralizing titer. The

resistance to REGN10933 was attributed to K417N and E484K **(Figure S3)**. REGN10933

neutralized virus with the Delta variant spike with a 12-fold decrease in titer which had

only a minor effect on the activity of the cocktail. REGN10987 showed a minor reduction

in neutralizing titer of virus with the Beta, Delta, Delta plus and Lambda variant spikes but

this had little effect on neutralization of the virus by the cocktail **(Figure 2D).** The

resistance of variants to REGN10987 was attributed to the L452R/Q **(Figure S3)**.

Discussion

Several reports have shown partial resistance of SARS-CoV-2 VOCs to vaccine-elicited

antibodies[4-11]. The data shown here extend those findings to the Delta plus and Lambda

variants. Delta plus and Lambda, VOIs, both displayed a degree of resistance to mRNA

vaccine-elicited antibodies similar to that of the Beta and Delta variants. In sera collected

~3 months post-second immunization, BNT162b2 and mRNA-1273 mRNA vaccine-

elicited antibodies neutralized the variants with a modest 3-fold average decrease in titer

resulting in an average IC_{50} of about 1:600, a titer that is greater than that of convalescent

sera and likely, in combination with post-vaccination T- and B-cell memory responses, to

provide durable protection. Ad26.COV2.S vaccination-elicited neutralizing antibodies

showed a more pronounced decrease in neutralizing titer against the variants, raising the

potential for decreased protection against the VOCs and the Lambda variant. Vaccination

with Ad26.COV2.S resulted in IC_{50} titers against Beta, Delta, Delta plus and Lambda

variants that decreased 5-7-fold, resulting in mean neutralizing antibody titers of 33, 30,

41, and 36 against viruses with the Beta, Delta, Delta plus and Lambda variant spikes,

respectively, which according to mathematical modeling, could result in decreased

protection against infection[15]. Modeling predicts that 50% protection from infection is

provided by a titer that is 20% that of the mean convalescent titer. In this study, given a

mean convalescent titer of 346 **(Table S1)**, 50% protection would correspond to an IC_{50}

of 69. The titer required to protect against severe disease was shown to be 3% that of the

mean titer of convalescent sera which in this study corresponds to a titer of 10. In a

published report of phase 3 trial data, a single dose of Ad26.COV2.S, 28 days post

administration, provided 64.0% protection against moderate to severe disease and 81.7%

against severe-critical COVID-19 in a country where 95% of circulating SARS-CoV-2 was

the Beta variant[2]. The authors considered possible roles for non-neutralizing antibody Fc-

mediated effector functions and the role of the T cell response in maintaining protection

against the partially neutralizing antibody-resistant Beta variant.

The data reported here differ somewhat from those reported by Barouch *et al*. and

Jongeneelen *et al*. who found that Ad26.COV2.S-elicited antibody titers were mostly

maintained against the variants[16,17]. In addition, Alter *et al*. reported a 5-fold decrease in

neutralizing antibody titer against Beta and 3.3-fold decrease against the Gamma variant

by the sera from Ad26.COV2.S vaccination[18] which were less pronounced than those

reported here. While the studies used similar assays to measure antibody neutralization

and analyzed sera collected at a similar time-point post-immunization, it is possible that

differences in the study populations accounted for the experimental differences.

Several recent studies have shown that boosting a single immunization of the

ChAdOx1nCoV-19 adenoviral vector vaccine with BNT162b2 resulted in high neutralizing

titer against the VOCs[19-21]. It is likely that neutralizing antibody titers against the VOCs

elicited by the single shot Ad26.COV2.S could similarly be improved by boosting with a

second immunization or by a heterologous boost with one of the mRNA vaccines. While

a single dose vaccination has advantages, the benefit provided by a second immunization

may be well worth the inconvenience.

The data presented here emphasize the importance of surveillance for breakthrough

infections with the increased prevalence of highly transmissible variants. If an increase in

breakthrough infections accompanied by severe COVID-19 is found following adenovirus

vector or mRNA vaccination, this would provide a rationale for public health policy-makers

and manufacturers to consider booster immunizations that would increase protection

against the VOCs and Lambda variant. As such a need is not currently evident, the public

health apparatus should focus on primary immunization in the U.S. and globally.

Methods

Clinical Samples

Convalescent sera were collected 32-57 days post-symptom onset. For the early time-point, BNT162b2 and Moderna-vaccinated sera were collected on day 28 and 35, respectively, 7 days post-second immunization. For the later time-point, BNT162b2-vaccinated sera were on average collected 90 days post-second immunization and mRNA-1273-vaccinated sera were collected on average 80 days post-second immunization. Ad26.COV2.S-vaccinated sera were collected, on average, 82 days post-immunization **(Table S2)**. Blood was drawn at the NYU Vaccine Center with written consent under IRB approved protocols (IRB 18-02035 and IRB 18-02037)**.** REGN10933 and REGN10987 were generated as previously described[22].

SARS-CoV-2 spike lentiviral pseudotypes

Lentiviruses pseudotyped by variant SARS-CoV-2 spikes were produced as previously reported[23] and normalized for reverse transcriptase (RT) activity. Neutralization titers of sera, monoclonal antibody and soluble ACE2 (sACE2)[24] were determined as previously described[23].

sACE2 pull-down assay

sACE2-bound-beads were mixed with pseudotyped virions as previously described[24]. The amount of virus bound was quantified by immunoblot analysis of bound p24.

Statistical Analysis

All experiments were in technical duplicates or triplicates. Statistical significance was

determined by two-tailed, unpaired t-test with confidence intervals shown as the mean ±

SD or SEM. (*P≤0.05, **P≤0.01, ***P≤0.001, ****P≤0.0001). Spike protein structure

(7BNM)[25] was downloaded from the Protein Data Bank.

Acknowledgements

The work was funded in part by grants from the NIH to N.R.L. (DA046100, AI122390 and

AI120898) and to M.J.M. (UM1AI148574). T.T. was supported by the Vilcek/Goldfarb

Fellowship Endowment Fund. M.J.M. and M.I.S. were partially supported by NYU

Grossman SOM institutional support.

Author contributions

T.T. and N.R.L. designed the experiments. H.Z., T.T. and B.M.D. carried out the

experiments and analyzed data. T.T., H.Z. and N.R.L. wrote the manuscript. M.I.S. and

M.J.M. designed and supervised the specimen selection, clinical information collection

and the N ELISAs, and provided key reagents and useful insights. All authors provided

critical comments on manuscript.

Declaration of Interests.

The authors declare no competing interests except M.J.M. who received research

grants from Lilly, Pfizer, and Sanofi, and serves on advisory boards for Pfizer and

Meissa Vaccines

Figure legends

Figure 1. Comparison of neutralization titers of variant spike protein pseudotyped

viruses by convalescent sera, antibodies elicited by BNT162b2, mRNA-1273,

Ad26.COV2.S.

(A) Neutralization of variant spike protein pseudotyped viruses by convalescent serum

(n=8) (left). Neutralizing titers of serum samples from BNT162b2 vaccinated individuals

(n=15) (middle). Neutralizing titers of serum samples from mRNA-1273 vaccinated

donors (n=6) (right). The serum was collected at early time point (7 days after second

immunization). The neutralization IC_{50} from individual donors is shown. Significance is

based on two-sided t-test.

(B) Comparison of neutralization of variants by convalescent serum (n=8, the same

donors in A), BNT162b2 vaccinated individuals (n=9), mRNA-1273 vaccinated donors

(n=8), Ad26.COV2.S vaccinated donors (n=10), sera from vaccinated individuals were

collected at later time points (90, 80, 82 days on average after last immunization of each

vaccine, see the table S2) . Each line shows individual donors.

(C) Comparison of neutralization potency of each vaccine by different SARS-CoV-2

variants. The neutralization IC_{50} from individual donors vaccinated by BNT162b2 (yellow),

mRNA-1273 (pink), Ad26.COV2.S (black) is shown. Significance is based on two-sided

t-test.

Figure 2. Neutralization of variant spike protein pseudotyped viruses by

monoclonal antibodies and sACE2.

(A) Infectivity of virus pseudotyped by variant and D614G spike proteins. Viruses were

normalized for RT activity and applied to target cells. Infectivity of viruses pseudotyped

with the variant proteins or the individual Lambda mutations were tested on ACE2.293T.

Luciferase activity was measured two days post-infection. Significance was based on two-

sided t-test.

(B) Neutralization of variant spike protein variants by sACE2. Viruses pseudotyped with

variant spike proteins were incubated with a serially diluted recombinant sACE2 and then

applied to ACE2.293T cells. Each plot represents the percent infectivity of D614G and

other mutated spike pseudotyped virus. The diagram shows the IC_{50} for each curve.

(C) Nickel beads were coated for 1 hour with 1, 0.5 and 0.1 μg of sACE2 proteins.

Unbound protein was removed and SARS-CoV-2 variant pseudotyped virions (D614G,

Delta, Lambda) were incubated with the beads. After 1 hour, the bound virions were

analyzed on an immunoblot with antibody p24 antibody. Beads-bound p24 (ng) was

calculated and indicated in the bottom (left). Input virions were analyzed on an

immunoblot with anti-p24 antibody (middle). Input sACE2 proteins were analyzed on an

immunoblot with anti-His-tag antibody (right).

(D) Neutralization of Beta, Delta, Delta plus and Lambda variant spike protein variants by

REGN10933 and REGN10987 monoclonal antibodies. Neutralization of D614G and

variant pseudotyped viruses by REGN10933 (left), REGN10987 (middle), and 1:1 ratio of

REGN10933 and REGN10987 (right). The IC_{50} values of REGN10933, REGN10987 and

the cocktail is shown in the table.

Supplemental Figure S1.

The structure of variant spikes and immunoblot analysis of spike proteins.

(A) The domain structure of the SARS-CoV-2 spike is diagrammed with Delta (B.1.617.2),

Delta plus (AY.1), Lambda (C.37) variant amino acid residues indicated. NTD, N-terminal

domain; RBD, receptor-binding domain; RBM, receptor-binding motif; SD1 subdomain 1;

SD2, subdomain 2; CS, cleavage site; FP, fusion peptide; HR1, heptad repeat 1; HR2,

heptad repeat 2; TM, transmembrane region; IC, intracellular domain. Key mutations are

shown in 3D structure (top view).

(B) Immunoblot analysis of the Delta (B.1.617.2), Delta plus (AY.1), single point mutated

of Lambda (C.37) variant, Lambda (C.37) variant spike proteins in transfected 293T cells.

Pseudotyped viruses were produced by transfection of 293T cells. Two days post-

transfection, virions were analyzed on an immunoblot probed with anti-spike antibody and

anti-HIV-1 p24. The cell lysates were probed with anti-spike antibody and anti-GAPDH

antibodies as a loading control.

Supplemental Figure S2.

Neutralization titers of spike protein pseudotyped viruses (single point mutations)

by convalescent sera, antibodies elicited by BNT162b2, mRNA-1273.

(A) Neutralization of variant spike protein (single point mutations) pseudotyped viruses by

convalescent serum (n=8). Dots represent the IC_{50} of single donors.

(B) Neutralizing titers of serum samples from BNT162b2 vaccinated individuals (n=15).

The serum was collected at early time point (7 days after second immunization). Each dot

represents the IC_{50} for a single donor.

(C) Neutralizing titers of serum samples from mRNA-1273 vaccinated donors (n=6). The

serum was collected at early time point (7 days after second immunization). The

neutralization IC_{50} from individual donors is shown. Significance is based on the two-sided

t-test.

Supplemental Figure S3.

Neutralization titers of spike protein pseudotyped viruses (single point mutations)

by monoclonal antibodies.

Neutralization of variant spike protein variants (single point mutations) by REGN10933

and REGN10987 monoclonal antibodies. The IC_{50} of REGN10933, REGN10987 and the

cocktail is shown in the table.

References

1.   Polack, F.P.*, et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine.

New England Journal of Medicine **383**, 2603-2615 (2020).

2.   Sadoff, J.*, et al.* Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against

Covid-19. *New England Journal of Medicine* **384**, 2187-2201 (2021).

3.   World Health Organization. (2021).

4.   Garcia-Beltran, W.F.*, et al.* Multiple SARS-CoV-2 variants escape neutralization

by vaccine-induced humoral immunity. *Cell* (2021).

5.   Tada, T.*, et al.* Convalescent-Phase Sera and Vaccine-Elicited Antibodies Largely

Maintain Neutralizing Titer against Global SARS-CoV-2 Variant Spikes. *mBio* **12**,

e0069621 (2021).

6.   Wang, P.*, et al.* Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7.

Nature (2021).

7.   Wu, K.*, et al.* Serum Neutralizing Activity Elicited by mRNA-1273 Vaccine. *New*

England Journal of Medicine **384**, 1468-1470 (2021).

8.   Xie, X.*, et al.* Neutralization of SARS-CoV-2 spike 69/70 deletion, E484K and

N501Y variants by BNT162b2 vaccine-elicited sera. *Nature Medicine* **27**, 620-621

(2021).

9.   Liu, J.*, et al.* BNT162b2-elicited neutralization of B.1.617 and other SARS-CoV-2

variants. *Nature* (2021).

10.   Choi, A.*, et al.* Serum Neutralizing Activity of mRNA-1273 against SARS-CoV-2

Variants. *bioRxiv*, 2021.2006.2028.449914 (2021).

11.   Planas, D*., et al.* Reduced sensitivity of infectious SARS-CoV-2 variant B.1.617.2

to monoclonal antibodies and sera from convalescent and vaccinated individuals.

bioRxiv, 2021.2005.2026.445838 (2021).

12.   Noval, M.G*., et al.* Antibody isotype diversity against SARS-CoV-2 is associated

with differential serum neutralization capacities. *Scientific Reports* **11**, 5538 (2021).

13.   Gu, H*., et al.* Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine

efficacy. *Science* **369**, 1603-1607 (2020).

14.   Starr, T.N*., et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding

Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310

e1220 (2020).

15.   Khoury, D.S*., et al.* Neutralizing antibody levels are highly predictive of immune

protection from symptomatic SARS-CoV-2 infection. *Nature Medicine* **27**, 1205-

1211 (2021).

16.   Jongeneelen, M*., et al.* Ad26.COV2.S elicited neutralizing activity against Delta

and other SARS-CoV-2 variants of concern. *bioRxiv*, 2021.2007.2001.450707

(2021).

17.   Barouch, D.H*., et al.* Durable Humoral and Cellular Immune Responses Following

Ad26.COV2.S Vaccination for COVID-19. *medRxiv*, 2021.2007.2005.21259918

(2021).

18.   Alter, G*., et al.* Immunogenicity of Ad26.COV2.S vaccine against SARS-CoV-2

variants in humans. *Nature* (2021).

19.   Tenbusch, M*., et al.* Heterologous prime-boost vaccination with ChAdOx1 nCoV-

19 and BNT162b2 mRNA. *medRxiv*, 2021.2007.2003.21258887 (2021).

20.   Gross, R.*, et al.* Heterologous ChAdOx1 nCoV-19 and BNT162b2 prime-boost

361   vaccination elicits potent neutralizing antibody responses and T cell reactivity.

362   *medRxiv*, 2021.2005.2030.21257971 (2021).

363   21.   Barros-Martins, J.*, et al.* Immune responses against SARS-CoV-2 variants after

364   heterologous and homologous ChAdOx1 nCoV-19/BNT162b2 vaccination. *Nature*

365   *Medicine* (2021).

366   22.   Tada, T.*, et al.* Decreased neutralization of SARS-CoV-2 global variants by

367   therapeutic   anti-spike   protein   monoclonal   antibodies.   *bioRxiv*,

368   2021.2002.2018.431897 (2021).

369   23.   Tada, T.*, et al.* Neutralization of viruses with European, South African, and United

370   States SARS-CoV-2 variant spike proteins by convalescent sera and BNT162b2

371   mRNA vaccine-elicited antibodies. *bioRxiv*, 2021.2002.2005.430003 (2021).

372   24.   Tada, T.*, et al.* An ACE2 Microbody Containing a Single Immunoglobulin Fc

373   Domain Is a Potent Inhibitor of SARS-CoV-2. *Cell Rep* **33**, 108528 (2020).

374   25.   Benton, D.J.*, et al.* The effect of the D614G substitution on the structure of the

375   spike glycoprotein of SARS-CoV-2. *Proc Natl Acad Sci U S A* **118**(2021).

Figure 1

	IC50 (ng/ml)				
	D614G (B.1)	Beta (B.1.351)	Delta (B.1.617.2)	Delta plus (AY.1)	Lambda (C.37)
REGN10933	3.3	420.4	39.1	306.3	2.9
REGN10987	9.3	16.4	28.8	29.7	33.7
REGN-COV2	4.4	33.1	9.8	19.5	7.0

Figure 2

Supplemental Figure S1

Supplemental Figure S2

A

	IC50 (ng/ml)									
	D614G	G75V-T76I	Δ246-252	L452Q	F490S	T859N	E484K	L452R	K417N	T478K
REGN10933	3.3	5.8	3.5	19.3	1.9	4.3	157.6	32.7	373.3	9.5
REGN10987	9.3	10.7	9.7	40.5	11.2	11.5	11.7	63.3	3.3	13.0
REGN-COV2	4.4	5.6	4.5	9.9	0.2	5.5	13.9	69.0	29.3	11.4

Supplemental Figure 3

Table S1. Neutralization of variants by convalescent sera, BNT162b2 and mRNA-1273 elicited antibodies

7 days post-second vaccination.

	Convalescent					
	IC$_{50}$(serum dilution)					
donor	D614G (B.1)	Alpha (B.1.1.7)	Beta (B.1.351)	Delta (B.1.617.2)	Delta plus (AY.1)	Lambda (C.37)
1	251	312	69	56	112	94
2	176	223	91	84	121	124
3	77	140	68	185	132	29
4	406	375	38	51	68	121
5	602	146	57	3	77	27
6	383	416	119	34	142	199
7	520	556	100	119	89	187
8	359	273	25	94	100	61
Mean (SD)	346 (174)	305 (142)	71 (32)	78 (56)	105 (26)	105 (66)

		BNT162b2					
		IC$_{50}$(serum dilution)					
donor	Days post last vaccine	D614G (B.1)	Alpha (B.1.1.7)	Beta (B.1.351)	Delta (B.1.617.2)	Delta plus (AY.1)	Lambda (C.37)
1	7	1915	1994	877	914	575	834
2	7	697	615	228	231	169	191
3	7	2572	2026	1366	950	1088	1244
4	7	939	925	145	507	171	123
5	7	1445	1717	161	416	361	167
6	7	2205	2069	413	370	614	935
7	7	1689	1259	918	560	1769	735
8	7	3189	2676	1045	1095	762	1032
9	7	1352	1720	456	594	363	451
10	7	1170	1355	604	669	796	635
11	7	672	729	219	398	592	238
12	7	571	841	364	441	480	259
13	7	3338	3099	1245	1463	1241	926
14	7	3486	3181	591	1042	685	1200
15	7	2294	2257	654	1092	1138	1888
Mean (SD)		1835 (986)	1764 (822)	619 (394)	716 (354)	720 (436)	724 (502)

		mRNA-1273					
		IC$_{50}$(serum dilution)					
donor	Days post last vaccine	D614G (B.1)	Alpha (B.1.1.7)	Beta (B.1.351)	Delta (B.1.617.2)	Delta plus (AY.1)	Lambda (C.37)
1	7	1380	1186	532	500	382	472
2	7	1963	1852	362	614	731	1185
3	7	1010	833	351	273	1055	209
4	7	1305	779	298	419	234	427
5	7	1879	2395	535	638	411	880
6	7	2028	1990	322	568	615	946
Mean (SD)		1594 (419)	1506 (668)	400 (106)	502 (138)	571 (296)	687 (373)

Table S2. Neutralization of viruses by sera from BNT162b2, mRNA-1273 and Ad26.COV.S vaccinated individuals.

						BNT162b2					
						IC$_{50}$(serum dilution)					
donor	Days post last vaccine	Anti-N ELISA	Age	Sex	Comorbidities	D614G	Alpha	Beta	Delta	Delta plus	Lambda
1	84	-	39	F	None	575	427	51	141	215	167
2	52	-	23	F	None	1338	1055	82	314	296	101
3	101	-	26	F	Asthma	1101	829	258	362	598	209
4	109	-	33	F	None	562	750	138	243	186	111
5	60	-	35	F	Hypothyroidism, Psoriasis	1024	930	53	239	391	284
6	81	-	42	F	Asthma	258	279	32	103	248	39
7	108	-	26	F	None	580	485	247	95	133	396
8	107	-	24	M	None	372	520	104	77	147	78
9	110	-	35	M	None	445	362	60	148	67	95
Mean (SD)	90 (22)		31			695 (369)	626 (272)	114 (85)	191 (102)	253 (161)	164 (114)

						mRNA-1273					
						IC$_{50}$(serum dilution)					
donor	Days post last vaccine	Anti-N ELISA	Age	Sex	Comorbidities	D614G	Alpha	Beta	Delta	Delta plus	Lambda
1	89	-	26	M	None	984	1043	108	173	364	257
2	92	-	53	M	None	972	703	237	207	239	273
3	61	-	67	M	Prediabetes	774	544	87	68	264	139
4	93	-	33	F	None	509	443	58	209	82	91
5	44	-	32	M	None	856	579	273	203	365	258
6	100	-	29	F	None	1038	1014	305	295	312	274
7	52	-	33	F	None	990	968	145	322	213	152
8	105	-	55	F	Asthma	537	485	246	184	160	391
Mean (SD)	80 (24)		41			833 (209)	722 (249)	182 (94)	208 (77)	250 (99)	229 (96)

						Ad26.COV2.S					
						IC$_{50}$(serum dilution)					
donor	Days post last vaccine	Anti-N ELISA	Age	Sex	Comorbidities	D614G	Alpha	Beta	Delta	Delta plus	Lambda
1	57	-	42	F	None	46	55	22	31	41	21
2	58	-	28	F	None	133	101	5	28	46	47
3	66	-	36	F	None	500	130	ND	ND	ND	ND
4	92	-	33	F	None	333	257	23	31	8	24
5	87	-	39	F	Prediabetes	244	205	19	42	31	36
6	72	-	32	M	None	268	308	79	34	63	59
7	92	-	39	F	None	251	377	44	46	38	70
8	71	-	75	F	None	298	648	ND	7	ND	ND
9	105	-	30	M	None	38	45	18	37	31	13
10	115	-	33	F	None	98	194	50	15	68	20
Mean (SD)	82 (20)		39			221 (144)	232 (182)	33 (24)	30 (12)	41 (19)	36 (21)