Fighting cancer with oncolytic viral therapy: identifying threshold parameters for success

Sana Jahedi[1], Lin Wang[2], and James Watmough[3]

Department of Mathematics and Statistics, University of New Brunswick, NB, Canada

Abstract

We model interactions between cancer cells and free virus during oncolytic viral therapy. One of our main goals is to identify parameter regions which yield treatment failure or success. We show that the tumor size under therapy at a certain time is less than the tumor size without therapy. We determine the minimum tumor size by the therapy and parameter regions under which this minimum is attained. Our analysis shows there are two thresholds for the horizontal transmission rate: a "Control threshold", the threshold above which treatment is efficient, and an "optimum threshold", the threshold beyond which infection prevalence reaches 100% and the tumor shrinks to its smallest size. Moreover, we explain how changes in the virulence level of the free virus alters the optimum threshold and the minimum tumor size. We identify a threshold for the virulence level of the virus and show how this threshold depends on timescale of virus dynamics. Our results suggests that when timescale of virus dynamics is fast, administration of a more virulent virus leads into more tumor reduction. When viral timescale is slow, a higher virulence will have drawbacks on the results, such as high amplitude oscillations. Furthermore, our numerical observation depicts fast and slow dynamics. Our numerical simulations indicate there exists a two-dimensional globally attracting surface that includes unstable manifold of the interior equilibrium. All solutions with positive initial conditions rapidly approach this two-dimensional attracting surface. In contrast, the trajectories on the attracting surface slowly tend to the periodic solution.

Keywords: oncolytic virus, tumor size, virulent virus, Hopf-bifurcation, tumor control, mathematical model, oscillatory behavior

Highlights

- The assumption that the viral load is in quasi-steady state is relaxed and infected cells are assumed to be mitotic.

- Our model supports that the tumor size under therapy over time is always lower in comparison with the case with no therapy.

- Minimum tumor size, control threshold and optimum threshold of the therapy are identified.

- Our analysis shows optimal virulence level of oncolytic virus depends on time scale of virus dynamics.

- When virus dynamic is slow, highly virulent virus causes long remission before relapse.

1. Introduction

The idea of using viruses as a cancer treatment formed when doctors noticed that some cancer patients experienced a short period of remission after a viral infection [1]. Oncolytic viruses are natural or modified

---

[1]Corresponding author: s.jahedi@unb.ca
[2]lwang2@unb.ca
[3]watmough@unb.ca

viruses that specifically target cancerous cells, replicate in them and eventually kill the infected cancer cells via lysis. Oncolytic viruses can also be armed with some cytokines or chemokines to activate antitumor immune cells [2, 3, 4, 5, 6], but here we only focus on their use as oncolytic agents.

To date, clinical trials conducted by using various oncolytic viruses have shown that these agents are less toxic than conventional therapies [7, 8], which makes this therapy popular. However, the therapeutic success of an oncolytic virus is limited [9, 10], and the complete eradication of a tumor has rarely been achieved when an oncolytic virus has been used as a single agent [11, 12]. This motivates us to study interactions between virus and cancer cells during oncolytic viral therapy in order to identify barriers and find ways to overcome them. Here, we use mathematical modeling to address the following questions: what are the outcomes of oncolytic viral therapy? under which condition is a specific outcome possible? can oncolytic virus be used as a cure or only as a containment method? and how does the virulence of the virus affect the outcome of the therapy? Moreover, we examine how to optimize the efficiency of viral therapy by identifying which parameter regions will lead to best outcome under the therapy.

Several mathematical models of oncolytic virus dynamics have been developed [13, 14, 15, 16, 17]. Komarova and Wodarz [18] studied the interactions between infected and uninfected cancer cells under the assumption that the turnover of the free virus is fast enough to maintain a quasi-steady state. They showed there is an optimum value for the death rate of infected cells at which the tumor reaches a minimum size. Long term oscillatory behavior has not been observed in their model, but in vivo studies of oncolytic viruses have suggested oscillatory behavior [9]. This encourages us to consider a three dimensional model, by relaxing the quasi-steady state assumption, to capture all possible dynamics during an oncolytic viral therapy.

Similar to many viruses, oncolytic viruses come in two types: enveloped and naked. Some example of naked oncolytic viruses are adenovirus [19] and reovirus [20]. For examples of enveloped oncolytic viruses we can refer to vesicular stomatitis virus [21], type 1 herpes simplex virus [22], and zika virus [23]. Enveloped oncolytic viruses can be easier for the host immune cells to remove than naked oncolytic viruses [24]. New virions of enveloped viruses can leave the membrane of their host without lysis, but naked viruses only leave the host cells via lysis. Hence, enveloped oncolytic viruses seem to be less efficient than naked oncolytic viruses. Our Model describes interaction of a naked oncolytic virus with, infected and uninfected, cancer cells during an oncolytic viral therapy.

Similar to Komarova and Wodarz [18], we consider using a mitotic virus. In other words, there are two pathways for transmission of infection: horizontal transmission, where an uninfected cell encounters free virus and becomes infected and vertical transmission, where infected cells produce infected daughter cells through mitosis.

In some cancers, larger tumors have an increased risk of metastasis. Koscielny et al [25] showed that the incidence of metastasis increases as the size of a breast tumor increases. In a study on survival experience of 1894 patients with invasive breast tumor, Narod [26] showed that increase in tumor size decreases 15-year survival rate of patients. Hence, the size to which tumor shrinks is one of the important factors to identify how successful a cancer treatment is. A treatment that causes considerable tumor shrinkage may be used as a tumor control method and in combination with other therapies may lead to tumor eradication. This inspires us to investigate how a virus should be engineered in order to shrink steady state of the tumor size in comparison to the case that no treatment occurs. It is important to know for which area in parameter space tumor size is minimum. As we will show, there exists a threshold for the horizontal rate of infectivity above of which the tumor size reaches its minimum.

In order to optimize the therapy, one needs to know how virulent the virus should be. Virulence is the negative impact caused by a viral infection on the functionality of host cells [27], such as a lower cell division rate or a higher mortality rate after infection. Here, we consider the use of an aggressive virus and explicitly include a rate associated with the effect of infection on cancer cell viability into the model. Our goal is to assess how the rate associated with virulence affects final tumor size. Recent studies suggest that very strong levels of viral aggression are not recommended [18]. Here, we show that when the virus becomes more aggressive, the system starts to show oscillatory behavior. We derive a threshold for virulence above which a periodic solution exists. Our numerical observations suggest this periodic solution is globally stable: that is to say, it attracts every solution that starts from a positive point which is not on the stable manifold of the positive equilibrium.

Capturing fast and slow dynamics are very beneficial to understand outcome of the treatment in longer term. Specially, when the fast behavior is followed immediately with a slow behavior. For example, when dynamics are fast and the outcome of treatment is followed only for a short time, it may sound that the treatment is successful in the sense that a huge tumor shrinkage is observed on the fast time scale, but tumors may appear again on the slower time scale. When the dynamics are too slow, tumor growth may not be noticeable when solutions are only observed for a short time, but in practice within some years tumor may get big enough to be a concern again. We show for which parameter region and for which state spaces fast and slow dynamics occur.

2. Model and Assumptions

In order to achieve a tractable model that helps us to focus on our research questions mentioned earlier we make the following assumptions.

A1: To model nutrient limitations we assume the mitosis rates are density dependent, dropping linearly to zero at a maximal tumor size K. We denote the per capita mitosis rates of uninfected and infected cancer cells at zero density by $\bar{r}$ and $\bar{s}$ respectively.

A2: As we mentioned before we assume there are two pathways for transmission of infection: β is the horizontal rate of infectivity per virus and $\bar{s}$, the mitosis rate of infected cancer cells, is the vertical transmission rate, see Figure 1.

A3: Furthermore, we assume vertical transmission is perfect. Meaning an infected cell never produces an uninfected daughter cell.

A4: Uninfected and infected cancer cells die at rates $\bar{d}$ and $\bar{d} + \bar{a}$ respectively and these rates are assumed to be lower than the per capita mitosis rates $\bar{r}$ and $\bar{s}$ respectively.

A5: The virus is virulent, meaning

- infection reduces cell growth; the self-renewal rate of infected cancer cells, $\bar{s}$, is lower than self-renewal rate of uninfected cancer cells, $\bar{r}$, i.e. $\bar{s} < \bar{r}$.

- infection elevates apoptosis; parameter $\bar{a} > 0$, denotes the elevation in death rate of cancer cells after infection.

A6: When an infected cell dies through apoptosis, no new virions are released, or the virions that are released in this way are not infectious (meaning the assembly of the virus has not been completed). We further assume that no virion can leave the membrane of an infected cell without lysis. As Figure 1 illustrates the free virus makes $\bar{q}$ copies of itself in each infected cell and the infected cell is lysed with rate $\bar{a}$. Therefore, $\bar{a}\bar{q}$ new virions are produced per each lysed infected cell.

Under the above assumptions we have the following model:

$$\dot{X} = \bar{r}X(1 - \frac{X+Y}{K}) - \bar{d}X - \beta XV \tag{1a}$$

$$\dot{Y} = \bar{s}Y(1 - \frac{X+Y}{K}) - \bar{d}Y - \bar{a}Y + \beta XV \tag{1b}$$

$$\dot{V} = \bar{a}\bar{q}Y - \bar{m}V \tag{1c}$$

where X, Y, and V denote the population density of uninfected cancer cells, infected cancer cells, and free virus, respectively. $\bar{m}$ in Eq. (1c) is the clearance rate of the free virus and finally, $X + Y$ denotes the tumor size. Figure 1 illustrates the viral cycle. First, viral incidence occurs between an uninfected cancer cell X and the free virus V at rate β. Once the free virus penetrates into the target cell, target cells becomes infected. Then the free virus replicates itself till the membrane of the infected cell is disrupted and the new virions are released in the extracellular space.

As we mentioned in assumption (A6), Model (1) demonstrators dynamics of a naked oncolytic virus. Therefore, it is assumed new virions only are released to extracellular area by lysis. Therefore, the rate at which new virions are reproduced is $\bar{a}\bar{q}$. However, when using an enveloped virus, new virions can leave the membrane of an infected cell without lysis. We also assumed that no new virions are released when an infected

cell goes through natural apoptosis, or the number of new virions released in this way are rare that can be ignored. In order for Model (1) to describe dynamics of any type of oncolytic virus one should change equation $\dot{V} = \bar{a}\bar{q}V - \bar{m}V$ to $\dot{V} = \bar{q}V - \bar{m}V$ (in this case $\bar{q}$ denotes the rate at which new virions are produced by each infected cell). Under such a change, mathematically there will be no difference in the way we analyze the results. Since we report the results for the rescaled model. For both scenarios, with different rescalings, same dimensionless model can be obtained. However, if we extend Model (1) to consider interactions with virus specific immune responses, depending in which types of virus specific immune response we consider, the outcome of therapy may be different depending on which kind of oncolytic virus (naked vs enveloped) is considered.

Figure 1: **A schematic description of the Model** (1). X, Y and V denote population densities of uninfected cancer cells, infected cancer cells and the free virus, respectively.

To simplify our analysis, we rescale the parameters and variables of Model (1) as follows:

$$x = \frac{X}{K}, \quad y = \frac{Y}{K}, \quad v = \frac{V\bar{m}}{K\bar{q}\bar{a}}, \quad t = \bar{r}\mathcal{T}$$

$$d = \frac{\bar{d}}{\bar{r}}, \quad a = \frac{\bar{a}}{\bar{r}}, \quad m = \frac{\bar{m}}{\bar{r}},$$

$$b = \frac{\beta K\bar{q}\bar{a}}{\bar{r}\bar{m}}, \quad s = \frac{\bar{s}}{\bar{r}},$$

Also for simplification we introduce two parameter combinations $X_U = 1 - d$ and $Y_I = 1 - (a + d)/s$. After applying these changes Model (1) turns to the following dimensionless model, the algebraic calculations are provided in Appendix A.1.

$$\dot{x} = x\big(X_U - (x + y)\big) - bvx, \tag{2a}$$

$$\dot{y} = sy\big(Y_I - (x + y)\big) + bvx, \tag{2b}$$

$$\dot{v} = m(y - v). \tag{2c}$$

Notice that when m is a large (small) number according to Eq. 2c, $|\dot{V}|$ is large (small, respectively). Therefore, throughout this work we use parameter m as a measure of time scale of virus dynamics.

Assumptions (A4) and (A5) in the terms of new parameters can be restated as follows;

(H1): $s < 1$; this holds since from assumption (A5), $\bar{s} < \bar{r}$, hence $s = \frac{\bar{s}}{\bar{r}} < 1$.

(H2): $X_U > Y_I > 0$; these hold because according to assumption (A4), $\bar{d} < \bar{r}$ and $\bar{d} + \bar{a} < \bar{s}$. Hence,

$d = \dfrac{\bar{d}}{\bar{r}} < 1$ and $\dfrac{d+a}{s} = \dfrac{\frac{\bar{d}}{\bar{r}} + \frac{\bar{a}}{\bar{r}}}{\frac{\bar{s}}{\bar{r}}} < 1$. Thus, $X_U > 0$ and $Y_I > 0$. According to assumption (H1), $s < 1$.

Therefore, $sd < 1(d+a)$. Hence, $1 - d > 1 - \dfrac{d+a}{s}$, meaning $X_U > Y_I$.

When modeling a biological phenomenon since the variables are representing population densities, we should show the solutions always stay non-negative and bounded. Next Lemma says each solution of Model (2) starting in $\mathcal{W}$ will remain in $\mathcal{W}$. The shaded wedge shape area in Figure 2(a) represents $\mathcal{W}$. Since $\mathcal{W}$ is a bounded subset of the first octant, therefore solutions of Model (2) are bounded and non-negative.

Lemma 1. For Model (2) the set

$$\mathcal{W} = \{(x, y, v) : 0 \leq x + y \leq X_U, 0 \leq x, y, v \leq X_U\} \tag{3}$$

is positively invariant and globally attractive.

Proof. To show $\mathcal{W}$ is positively invariant, it is enough to show a trajectory that starts from a point on the boundaries of $\mathcal{W}$ either remains on the boundaries of $\mathcal{W}$ for all $t \geq 0$ or it tends inward $\mathcal{W}$. Each solution of Model (2) that starts with an initial point on the boundaries of $\mathcal{W}$ which is not located on the v axis, x axis or on the $y - v$ plane, points inward of $\mathcal{W}$. Each solution that starts from an initial point on the boundaries of $\mathcal{W}$, which is located on the v axis, the x axis or $y - v$ plane tends to $(0, 0, 0)$, $(X_U, 0, 0)$ or $(0, Y_I, Y_I)$, respectively (for details see Appendix A.2). $(0, 0, 0)$, $(X_U, 0, 0)$ and $(0, Y_I, Y_I)$ are all in boundaries of $\mathcal{W}$. Therefore, $\mathcal{W}$ is positively invariant.

When $x + y \geq X_U$ and $x, y > 0$, Eqs. (2a) and (2b) yield the following equation.

$$\frac{d(x+y)}{dt} = x\big(X_U - (x+y)\big) + sy\big(Y_I - (x+y)\big) \leq -sy(X_U - Y_I). \tag{4}$$

The above equation suggests if $x + y \geq X_U$ and $x, y > 0$, then $\dfrac{d(x+y)}{dt} < 0$. In other words, trajectory of each solution starting from a point $(x(0), y(0), v(0))$ that $x(0) + y(0) \geq X_U$ and $x(0), y(0) > 0$, moves in the direction of decreasing $x + y$, therefore, it tends inward $\mathcal{W}$.

When $v \geq X_U$ and $v > y$, then $\dot{v} < 0$. Hence v decreases and also any solution starting on the $x + y = X_U$ plane points towards wedge in the direction of decreasing $x + y$. Therefore each solution starting from an initial point $(x(0), y(0), v(0))$, for which $x(0) + y(0) < X_U$ and $v(0) > X_U$, tends to a point in $\mathcal{W}$.

Therefore, each solution with an initial point (x_0, y_0, v_0) such that $x_0, y_0, v_0 \geq 0$ and $(x_0, y_0, v_0) \notin \mathcal{W}$, eventually tends to a point in $\mathcal{W}$. $\square$

Here after we only consider solutions with initial points in $\mathcal{W}$. The shaded plane in the Figure 2(b) is the plane $y = v$ intersection with $\mathcal{W}$. All equilibria of Model (2) are on this plane.

Figure 2: $\mathcal{W}$ **is positively invariant and globally attractive.** (a) This plot demonstrates that trajectory of each solution that starts from a point in the first octant eventually gets attracted to $\mathcal{W}$. (b) This plot is same as plot (a), but it also includes the $y = v$ plane's intersection with $\mathcal{W}$. When a solutions starts above of this plane, $y < v$ so v decreases, and when a solution starts below this plane v increases.

The tumor size, meaning the total density of infected and uninfected cancer cells determines whether the therapy is successful. The next proposition shows the tumor size under oncolytic viral therapy over time always remains lower than the tumor size with no therapy over time.

Let $\bar{x}(t)$, $t \geq 0$, be the tumor size without therapy. That is, $\bar{x}(t)$ is the solution of Model (2) with initial condition $(x(0), y(0), v(0))$, where $x(0) > 0$, $y(0) = v(0) = 0$. Here $x(t) + y(t)$, $t \geq 0$ shows the tumor size with therapy. That is, the solution to Model (2) with initial condition $(x(0), y(0), v(0))$, where $x(0) > 0$, $v(0) > 0$, $y(0) \geq 0$. Then we have the following.

Proposition 1. Under Model (2) assume the virus is introduced at time $t = 0$, either directly into cells or into the extracellular environment. Then the tumor size under therapy $x(t) + y(t)$ at time $t > 0$ will be less than the tumor size without therapy $\bar{x}(t)$. I.e. if $x(0) + y(0) = \bar{x}(0)$ and $v(0) > 0$, $y(0) \geq 0$, then $x(t) + y(t) < \bar{x}(t)$ for all $t > 0$.

Proof. According to Eq. (2b), for each solution of Model (2) that starts with $x(0) > 0$ and $v(0) > 0$ and $y(0) = 0$, y variable is initially increasing $\dot{y}(0) > 0$. Hence, we conclude that $y(t) > 0$ for all $t > 0$. Now suppose the contrary assumption, that there exists $t^* > 0$ such that $x(t^*) + y(t^*) \geq \bar{x}(t^*)$, i.e. $\frac{x(t^*) + y(t^*)}{\bar{x}(t^*)} \geq 1$. Define $\rho(t) = \frac{x(t) + y(t)}{\bar{x}(t)}$; $t \geq 0$. Notice that $\rho(t)$ is a continuous function. Now we will show $\rho(t)$ is strictly decreasing on the set of all t for which $\rho(t) \geq 1$. Eqs. (2a) and (2b) yield to

$$\frac{d(x + y)}{dt} = x\big(X_U - (x + y)\big) + sy\big(Y_I - (x + y)\big).$$

Since $y > 0$ for all $t > 0$, $Y_I < X_U$, and $s < 1$,

$$\frac{d(x + y)}{dt} < x\big(X_U - (x + y)\big) + sy\big(X_U - (x + y)\big) = (x + sy)\big(X_U - (x + y)\big) < (x + y)\big(X_U - (x + y)\big).$$

Since $x + y \neq 0$,

$$\frac{1}{x + y}\frac{d(x + y)}{dt} < X_U - (x + y).$$

For all $t > 0$ such that $\rho(t) \geq 1$ (i.e. $x(t) + y(t) \geq \bar{x}(t)$),

$$\frac{1}{x + y}\frac{d(x + y)(t)}{dt} < X_U - (x + y)(t) \leq X_U - \bar{x}(t) = \frac{1}{\bar{x}}\frac{d\bar{x}(t)}{dt}.$$

Hence, when $\rho(t) \geq 1$, $\frac{d}{dt}(Ln(\rho(t))) = \frac{\dot{\rho}(t)}{\rho(t)} < 0$. Therefore, $\rho(t)$ is strictly decreasing on the set of all $t > 0$ for which $\rho(t) \geq 1$. But $\rho(t)$ is a continuous function and $\rho(0) = 1$. Therefore, ρ is strictly decreasing on the set of all $t \geq 0$ for which $\rho(t) \geq 1$. Therefore $\rho(0) > \rho(t^*) \geq 1$, contradicting our assumption. $\qquad\square$

In the proof of the next proposition, we use comparison theorem [28], for convenience of the readers the version of comparison theorem that we use is provided in Appendix A.4.

Proposition 2.

(I) The tumor size under the therapy $x(t) + y(t)$ will remain between Y_I and X_U if the initial tumor size is between Y_I and X_U. I.e,

$$Y_I \leq x(t) + y(t) \leq X_U; \text{ for all } t > 0, \tag{5}$$

if $Y_I \leq x(0) + y(0) \leq X_U$.

(II) For each solution $(x(t), y(t), v(t))$ with an initial condition $(x(0), y(0), v(0))$, where $x(0) > 0, y(0) \geq 0$ there exists $t^* > 0$ such that $Y_I \leq x(t^*) + y(t^*) \leq X_U$.

Proof of (I). Eqs. (2a) and (2b) yield the following equation.

$$\frac{d(x+y)}{dt} = x\big(X_U - (x+y)\big) + sy\big(Y_I - (x+y)\big). \tag{6}$$

Since $x, y \geq 0$, $Y_I < X_U$ and $s < 1$ therefore, $sy\big(Y_I - (x+y)\big) \leq y\big(X_U - (x+y)\big)$. Hence,

$$\frac{d(x+y)}{dt} \leq (x+y)(X_U - (x+y)).$$

Therefore, if $x(0) + y(0) \leq X_U$, then according to the comparison theorem, $x(t) + y(t) \leq X_U$ for all $t \geq 0$.

Similarly, using $s(x+y) \leq x + sy$ from Eq. (6)

$$\frac{d(x+y)}{dt} \geq sx\big(Y_I - (x+y)\big) + sy\big(Y_I - (x+y)\big) = s(x+y)\big(Y_I - (x+y)\big).$$

Therefore, if $x(0) + y(0) \geq Y_I$, then according to comparison theorem, $x(t) + y(t) \geq Y_I$ for all $t \geq 0$.

Proof of (II). Now assume $(x(t), y(t), v(t))$ is a solution with $(x(0), y(0), v(0))$, where $x(0), y(0) > 0$. We will show there exist $t^* > 0$ such that $Y_I \leq x(t^*) + y(t^*) \leq X_U$.

As we showed in the proof of Lemma 1, $[0, X_U]$ is globally attracting, so we only need to show that for each solution $(x(t), y(t), v(t))$ with initial condition $(x(0), y(0), v(0))$ where $0 < x(0) + y(0) < Y_I$ and $x(0), y(0) > 0$ there exists t^* such that $x(t^*) + y(t^*) \geq Y_I$.

By contrary assume that the set $U := \{(x(t), y(t), v(t)) \in \mathcal{W} : 0 < x(0) + y(0) < Y_I, x(0), y(0) > 0\}$ is positively invariant. If $x + y < Y_I$ and $x, y > 0$ according to Eq. (6), $\frac{d(x+y)}{dt} > 0$, therefore, $x(t) + y(t)$ is increasing along each solution $(x(t), y(t), v(t)) \in U$ and it is bounded above by Y_I. Hence, $\lim_{t \to \infty} x(t) + y(t) = Y_I$, for any $(x(t), y(t), v(t)) \in U$. But Y_I is not in U, this contradicts by U being positively invariant. $\square$

3. Equilibria and their stability

Model (2) can have four equilibria in $\mathcal{W}$, which we will refer to as follows: the tumor eradication equilibrium, $E_0 = (0, 0, 0)$, the treatment failure equilibrium, $E_U = (X_U, 0, 0)$, 100% infection prevalence equilibrium, $E_I = (0, Y_I, Y_I)$ and the partially infected equilibrium, $E_p = (\bar{x}, \bar{y}, \bar{v})$, where

$$\bar{x} = \frac{s(-Y_I b + X_U - Y_I)}{b(b + 1 - s)} \tag{7a}$$

$$\bar{y} = \frac{X_U b - s(X_U - Y_I)}{b(b + 1 - s)} \tag{7b}$$

$$\bar{v} = \bar{y}. \tag{7c}$$

Next lemma tells under which conditions the above equilibria exist.

Lemma 2. E_0, E_U, E_I always exist. E_p is positive if and only if

$$\frac{s(X_U - Y_I)}{X_U} < b < \frac{X_U - Y_I}{Y_I} \tag{8}$$

Proof. According to assumption (H2), $X_U > 0$ and $Y_I > 0$, thus E_U and E_I always exist.

If $\bar{x} = 0$, then simple calculations shows $E_p = E_I$. Similarly, if $\bar{y} = 0$, then $E_p = E_U$. But, we are interested in the case that both $\bar{x}$ and $\bar{y}$ are positive. According to assumption (H1), $s < 1$. Thus, $b + 1 - s > 0$. Hence, E_p is positive if and only if both followings are satisfied:

1. $-Y_I b + X_U - Y_I > 0$

2. $X_U b - s(X_U - Y_I) > 0$

Therefore, E_p is positive if and only if $\frac{s(X_U - Y_I)}{X_U} < b < \frac{X_U - Y_I}{Y_I}$. $\qquad\square$

We will refer to E_p as positive equilibrium. Next lemma summarizes the conditions under which each boundary equilibrium is locally asymptotically stable.

Lemma 3.

1. E_0 is always unstable

2. $E_U = (X_U, 0, 0)$ is locally stable if and only if $b < \frac{s(X_U - Y_I)}{X_U}$.

3. $E_I = (0, Y_I, Y_I)$ is locally stable if and only if $\frac{X_U - Y_I}{Y_I} < b$.

Proof of (1). Eigenvalues of Jacobian of Model (2) at E_0 are $-m, X_U, sY_I$. According to assumption (H2), X_U, Y_I are both positive. Since E_0 has two positive eigenvalues, so it is always unstable.

Proof of (2). Jacobian of Model (2) at E_U always has a negative eigenvalue, $-X_U$. The other two remaining eigenvalues are eigenvalues of the following matrix.

$$A = \begin{bmatrix} s(Y_I - X_U) & bX_U \\ m & -m \end{bmatrix}$$

$\mathrm{tr}A = -m + s(Y_I - X_U)$ and $\det A = -mX_U(b - \frac{s(X_U - Y_I)}{X_U})$. According to assumption (H2), $Y_I < X_U$. Hence, trA is always negative. Therefore, the local stability of E_U is only determined by the sign of $\det A$. Hence, E_U is locally stable if and only if $\det A > 0$. Thus, E_U is locally stable if and only if $b < \frac{s(X_U - Y_I)}{X_U}$.

Proof of (3). Jacobian of Model (2) at equilibrium E_I has eigenvalues $\lambda_{I1} = -m$, $\lambda_{I2} = -sY_I$, and $\lambda_{I3} = X_U - Y_I - bY_I$. The first two eigenvalues are negative, so E_I is locally stable if and only if λ_{I3} is negative. In other words, E_I is locally stable if and only if $b > \frac{X_U - Y_I}{Y_I}$ $\qquad\square$

Proposition 2 and Lemma 3 together show when $b > \frac{s(X_U - Y_I)}{X_U}$ tumor shrinkage occurs and when $b < \frac{s(X_U - Y_I)}{X_U}$ treatment fails. According to Lemma 3 when $b < \frac{s(X_U - Y_I)}{X_U}$, the treatment failure equilibrium E_U is stable and our numerical observations illustrate when an equilibrium is locally stable, it is also globally stable. I.e., each solution starting from a positive initial point will tend to that stable equilibrium. Therefore, when $b < \frac{s(X_U - Y_I)}{X_U}$, regardless of initial tumor size, administrating oncolytic virus has no effect and tumor grows to maximum possible tumor size X_U. Furthermore, Proposition 2 shows eventual tumor size is always smaller than or equal to X_U, but Lemma 3 says when $b > \frac{s(X_U - Y_I)}{X_U}$, eventual tumor size is never X_U. Hence, when $b > \frac{s(X_U - Y_I)}{X_U}$ eventually tumor size is smaller than X_U. Therefore, effective result by therapy will be observable when $b > \frac{s(X_U - Y_I)}{X_U}$. Hence, we call $\boldsymbol{b = \frac{s(X_U - Y_I)}{X_U}}$ **tumor control threshold**.

Lemma 3 says when $b > \frac{X_U - Y_I}{Y_I}$, 100% infection prevalence equilibrium E_I is stable. In this case eventually tumor size is Y_I. Additionally, Proposition 2 says Y_I is the optimum tumor size for a specific virus (fixed set of parameters), therefore we call $\boldsymbol{b = \frac{X_U - Y_I}{Y_I}}$ **optimum threshold** of b for therapy.

Figure 3 is a graphical representation of Proposition 2. This figure shows regardless of time scale of virus dynamics, tumor size under therapy remains between Y_I and X_U. This figure shows tumor size under therapy decreases when the rescaled infectivity rate b increases. Notice that according to (H1), $s < 1$ and according to (H2), $Y_I < X_U$, therefore $\frac{s}{X_U} < \frac{1}{Y_I}$, multiplying this inequality by $(X_U - Y_I)$ implies $\frac{s(X_U - Y_I)}{X_U} < \frac{X_U - Y_I}{Y_I}$. I.e, control threshold is always strictly lower than optimum threshold. Hence, Figure 3 is a generic bifurcation diagram of Model (2).

(a) Fast viral time scale   (b) Slow viral time scale

Figure 3: **The outcome of therapy is enhanced when horizontal rate of infection transmission increases.** In both plots maximum and minimum tumor size is sketched. When an equilibrium is globally stable eventually all the solutions will tend to it, therefore in that case maximum and minimum tumor size will be same. As both plots represent, when b is smaller than control threshold, treatment fails and when b is greater than optimum threshold, minimum tumor size by therapy Y_I is obtained. In both cases the tumor size under therapy remains between Y_I and X_U. When viral time scale is fast, same as plot(a), no long term oscillatory behavior will be observable and when viral time scale is slow, same as plot(b), sustained oscillation is observable. As plot (b) indicates even when sustained oscillations are observable tumor size remains between Y_I and X_U. This simulation is conducted under the set of parameters $X_U = 0.9$, $Y_I = 0.08$, and $s = 0.5$. $m = 1$ in plot(a) and $m = 0.1$ in plot(b). In this plot, a dashed (solid) curve means for that value of the parameter, the corresponding equilibrium is unstable (stable).

4. Hopf-bifurcation and sustained oscillations

A Hopf-bifurcation occurs in a system when by changes in one of the parameters of the system periodic solutions appear, i.e., in one side of the bifurcation point there is no limit cycle, but as the bifurcation parameter passes through bifurcation point, periodic solution appears [29]. For convenience of the reader Hopf bifurcation Theorem is provided in Appendix A.3. In this section we will identify the parameter space at which Hopf bifurcation occurs.

Verifying of occurrence of a Hopf-bifurcation is often involved with algebraic calculation of eigenvalues. In 1998, Liu [30] proposed criterion for occurrence of Hopf bifurcation without calculation of eigenvalues, for a general system of ordinary differential equation of order n. Here, we use this criterion in the proof of Theorem 2. Hence, below we restate the criterion "Routh-Hopf" for a three dimensional system of ordinary differential equations and provide a simpler proof for this special case.

Consider a three dimensional system of ordinary differential equations is given, meaning

$$\dot{U} = f(U, m), \tag{9}$$

where $U \in \mathbb{R}^3$ and $m \in \mathbb{R}$.

Assume $(U_{eq}(m), m) \in \mathbb{R}^{3+1}$ is a curve of stationary points for the system (9). Denote the characteristic polynomial of $Df(U_{eq}(m), m)$ by P_m. Meaning

$$P_m(\lambda) := \det\left(\lambda I - Df\big(U_{eq}(m), m\big)\right)$$

P_m is a degree three polynomial therefore, it can be written as $P_m(\lambda) = \lambda^3 + P_2(m)\lambda^2 + P_1(m)\lambda + P_0(m)$.

Define Routh function as $H(m) := P_2(m)P_1(m) - P_0(m)$ [31].

Theorem 1 (Routh-Hopf Criterion). Assume that $P_0 \neq 0$. A Hopf bifurcation occurs at $m = m^*$ for
the system (9) if the two following conditions are satisfied.

(i) $H(m^*) = 0$,

(ii) $\dfrac{dH}{dm}\big|_{m^*} \neq 0$.

Proof. Let us assume that P_m has two complex roots $\lambda_1 = a + bi$, $\lambda_2 = a - bi$ and a real root c, where $a = a(m)$, $b = b(m)$ and $c = c(m)$. Hence, P_m can be written as

$$P_m = \big(\lambda - a + bi\big)\big(\lambda - a - bi\big)\big(\lambda - c\big)$$

After simplification we have

$$P_m = \lambda^3 - \big(2a + c\big)\lambda^2 + \big(a^2 + 2ca + b^2\big)\lambda - \big(a^2 + b^2\big)c. \tag{10}$$

238   From Equation (10), Routh function $H = P_2 P_1 - P_0$ can be written as $H = -2a \cdot (a^2 + 2ac + b^2 + c^2)$.
239   Therefore, $H = -2a\big((a+c)^2 + b^2\big)$. Since $P_0 \neq 0$, therefore H is zero if and only if $a = 0$. Also, Since
240    $P_0 \neq 0$, a and b cannot both be zero simultaneously.

241   Assume that $a(m^*) = 0$. We will show $\dfrac{da}{dm}\big|_{m^*} \neq 0$ if and only if $\dfrac{dH}{dm}\big|_{m^*} \neq 0$

As we showed above $H = -2a\big((a+c)^2 + b^2\big)$, therefore

$$\frac{dH}{dm}\big|_{m^*} = -2\frac{da}{dm}\big|_{m^*}\big((a(m^*) + c(m^*))^2 + b(m^*)^2\big) - 2a(m^*)\frac{d}{dm}\big((a+c)^2 + b^2)\big)\big|_{m^*}$$

Since $a(m^*) = 0$, therefore

$$\frac{dH}{dm}\big|_{m^*} = -2\frac{da}{dm}\big|_{m^*}\big((a+c)^2 + b^2\big)$$

242   Since $P_0 \neq 0$, then $(a+c)^2 + b^2 \neq 0$, therefore

$$\frac{dH}{dm}\big|_{m^*} = 0 \text{ if and only if } \frac{da}{dm}\big|_{m^*} = 0. \tag{11}$$

243   We have now shown (i) and (ii) imply the conditions of occurrence of a Hopf bifurcation (see Appendix
244   A.3), so by Hopf bifurcation theorem, a Hopf-bifurcation occurs when (i) and (ii) hold. $\square$

Theorem 2. Let $\dfrac{s(X_U - Y_I)}{X_U} < b < \dfrac{X_U - Y_I}{Y_I}$. Define,

$$\Delta := -\Big((\bar{x} + s\bar{y} + b\bar{x})^2 - \bar{x}(\bar{x} + \bar{y})b^2\Big) - 2\sqrt{\big(\bar{x} + s\bar{y} + b\bar{x})(\bar{x} + s\bar{y})(\bar{x} + \bar{y})\bar{x}b},$$

245   where $\bar{x}$ and $\bar{y}$ are same as Eq. (7). Then we have the following.

246   (i) If $\Delta < 0$, then E_p is a stable equilibrium.

247   (ii) If $\Delta > 0$, then two Hopf bifurcations occur as the parameter m changes. We denote these Hopf-
248   bifurcation points by m_1 and m_2, where $m_1 < m_2$.

(iii) If $\Delta > 0$, then E_p is a stable equilibrium if and only if $m \in (-\infty, m_1) \cup (m_2, \infty)$

Proof. The Jacobian of Model (2) at $E_p = (\bar{x}, \bar{y}, \bar{v})$ is

$$
J_C = \begin{bmatrix} -\bar{x} & -\bar{x} & -b\bar{x} \\ (b-s)\bar{y} & -b\bar{x} - s\bar{y} & b\bar{x} \\ 0 & m & -m \end{bmatrix}
$$

The characteristic equation for J_C is

$$
\begin{aligned}
P &= \det(\lambda I - J_C) \\
&= \lambda^3 + \underbrace{(b\bar{x} + s\bar{y} + m + \bar{x})}_{a_2} \lambda^2 + \underbrace{(b\bar{x}^2 + b\bar{x}\bar{y} + ms\bar{y} + m\bar{x})}_{a_1} \lambda + \underbrace{m\bar{x}\bar{y}b(b + 1 - s)}_{a_0}
\end{aligned}
\tag{12}
$$

According to assumption (H1), $s < 1$. Therefore, a_1, a_2 and a_3 are always positive. Routh function H for the above characteristic polynomial is as follows.

$$
H = a_2 a_1 - a_0 = \underbrace{(\bar{x} + s\bar{y})}_{A} m^2 + \underbrace{\left((\bar{x} + s\bar{y} + b\bar{x})^2 - \bar{x}(\bar{x} + \bar{y})b^2 \right)}_{B} m
\tag{13}
$$
$$
+ \underbrace{b(\bar{x} + s\bar{y} + b\bar{x})(\bar{x} + \bar{y})\bar{x}}_{C}
$$

Since $A, C > 0$, $H = 0$ has positive solutions for m if and only if $B < 0$ and $B^2 - 4AC \geq 0$. In other words, H has positive solutions for m if and only if $-B \geq 2\sqrt{AC}$. Notice that $\Delta = -B - 2\sqrt{AC}$. Hence, $H = 0$ only has solutions if and only if $\Delta \geq 0$.

When $\Delta < 0$, H is always positive. As we mentioned above a_1, a_2 and a_3 are also positive. Thus, according to Routh-Hurwitz stability criterion [31] equilibrium E_p is locally asymptotically stable. This proves part (i) of the theorem.

When $\Delta > 0$, the quadratic H has two distinct positive roots m_1 and m_2, where $m_1 < m_2$. Of course, $\frac{dH}{dm}(m) \neq 0$ for $m = m_1, m_2$. Therefore according to Routh-Hopf Criterion, Hopf-bifurcations occur as m changes through m_1 and m_2. This proves part (ii) of the theorem.

When $\Delta > 0$, and $m \in (-\infty, m_1) \cup (m_2, \infty)$, then Routh function is positive, therefore according to Routh-Hurwitz criterion E_p is locally asymptotically stable. This proves part (iii). $\qquad \square$

Note that when $\Delta = 0$, then the Routh function has a repeated root. Hence, there exists an m^* such that $H(m^*) = 0$, but $\frac{\partial H}{\partial m}(m^*) = 0$. Thus, the conditions for the occurrence of Hopf bifurcation with respect to parameter m, are not satisfied at $m = m^*$. Therefore, when $\Delta = 0$, for any $m \neq m^*$ E_p is an stable equilibrium and it is a non-hyperbolic equilibrium at $m = m^*$. In our numerical simulations we observed that for $m = m^*$, the equilibrium E_p is globally stable, and every solution starting from a positive initial point tends to E_p very slowly, in the way that oscillating solutions are decaying very slowly, tending eventually to E_p. Notice that when function H changes sign, the real part of eigenvalues of E_p are changing the sign. Therefore, any bifurcation curve bifurcating from E_p, should also cross $H = 0$. Since H is a function of all the parameters of Model (2), one may choose a different bifurcation parameter, here we have chosen m, since we are interested in effect of time scale of virus dynamics on the outcome of therapy.

5. Numerical observations

In this section we report some observations that are illustrated numerically.

As it is mentioned above in Theorem 2 when $\Delta > 0$ and $m_1 < m < m_2$ the equilibrium E_p is unstable and a locally stable limit cycle exists. In Figure 4 we have conducted a simulation such that a limit cycle exists: the black closed curve is the limit cycle, the black circle is equilibrium E_p, and the brown curve passing through equilibrium E_p is its stable manifold. Our numerical observations suggest that when a limit cycle exists, it is almost everywhere globally stable. I.e., each solution starting from a positive initial point that is not located on the stable manifold of the equilibrium E_p tends to the limit cycle. In Figure 4, each solution starting from a point on the brown curve tends to E_p. Each solution starting from a positive initial point which is not on the brown curve tends to the limit cycle. To have a more clear Figure here only some solutions (green and red curves) are sketched. The red curves start from a solution outside of the limit cycle near to the stable manifold of E_p, they jump into the area enclosed by the limit cycle and spiral out from the nearby of the equilibrium E_p and tend to the limit cycle. The green curve oscillations' amplitude gets smaller and spirals in towards the limit cycle. As Figure illustrates the disk which is traced by the red curve, starting very near by E_p and bounds to the limit cycle is the unstable manifold of the equilibrium E_p. The red disk together with the equilibrium E_p and its boundary, the limit cycle forms a global attracting surface which we denote it by G. Each solution with positive initial condition tends to G.

Figure 4: **The global attracting surface (a)** The black curve represents a stable limit cycle and the black circle is E_p. Each solution starting from a positive point (except for the steady state E_p and its one dimensional stable manifold) approaches to this limit cycle. The red disk (traced out by the red curves) is the unstable manifold of E_p and the limit cycle is its boundary. Plot **(b)** is the same figure as plot (a) which has been rotated in order to show the stable manifold of equilibrium E_p. The brown curve shows one dimensional stable manifold of E_p. Every solution starting from a point on the brown curve gets attracted to the E_p. The disk centered at equilibrium E_p with its boundary, the limit cycle, is a global attracting surface. This simulation is conducted for $s = 0.5$, $X_U = 0.9$, $Y_I = 0.08$, $m = 0.1$, and $b \sim 5.35$. This b is the mid point of the interval $\left(\frac{s(X_U - Y_I)}{X_U}, \frac{X_U - Y_I}{Y_I}\right)$, on which E_P exists.

Figure 5 shows bifurcation of tumor size with respect to changes in parameter m, when $\frac{X_U - Y_I}{X_U} < b < \frac{X_U - Y_I}{Y_I}$. Our simulations show that when $m < m_1$ or $m > m_2$ the fixed point $E_p = (\bar{x}, \bar{y}, \bar{y})$ is globally stable, strictly speaking it attracts all the solutions starting from a positive initial point. Therefore, when $m < m_1$ or $m > m_2$, the steady state tumor size is $T_p = \bar{x} + \bar{y}$. When $m_1 < m < m_2$, all the solutions starting from a positive point which is not on the stable manifold of E_p tends to the limit cycle, so in this case the tumor size is oscillating, so it has a maximum, denoted by the upper curve $max(T_{per})$ and a minimum, denoted by the lower curve $min(T_{per})$. As this figure illustrates the Hopf- bifurcation is global in the sense that as m changes a continuous path of periodic solutions sources from m_1 and sinks in m_2 [32].

Figure 5: **Bifurcation with respect to m:** This plot shows how the oscillation in tumor size changes when variable m changes. As figure describes when m passes through m_1, partially infected equilibrium, $E_p = (\bar{x}, \bar{y}, \bar{y})$ becomes unstable and a stable limit cycle appears. $T_p = \bar{x} + \bar{y}$, shows the tumor size when E_p is stable, and $max(T_{per})$ and $min(T_{per})$ shows the maximum and minimum of tumor size when the limit cycle exists. When m passes through m_2 the stable limit cycle disappears and again E_p becomes stable. This simulation is done for the set of parameters s=0.5, X_U=0.9, Y_I=0.08, and $b \sim 5.35$. This b is the mid point of the interval $(\frac{s(X_U - Y_I)}{X_U}, \frac{X_U - Y_I}{Y_I})$, on which E_P is positive.

5.1. Fast-slow dynamics of tumor size

Another interesting result which is observed numerically is that Model (1) shows fast and slow behavior. As we mentioned above there is a range of parameters for which there is a unique stable periodic solution to which every trajectory starting from a positive point, except for a single curve (the stable manifold of the steady state), is attracted. However, not every solution gets attracted with the same speed and since the speed of remission is a crucial factor in the therapy, it motivates us to investigate for which regions of state spaces fast and slow behavior occur.

An example of slow dynamics is shown in Figure 6, where two solutions are shown approaching a periodic orbit very slowly. Trajectories oscillate with a period about $56.6606 \sim 57$, but takes so long to get close to the periodic orbit. One (green) approaches with decreasing amplitude and the other (red) approaches with increasing amplitude. The black curve is equilibrium E_p. The red curve starts with an initial condition near to E_p slowly spirals out and tends to the stable periodic solution. The figure suggests that it takes up to 83 periods for the red curve and almost 38 periods for the green curve to tend to the stable periodic solution. The two points marked by yellow circles indicate that for the amplitude of red curve to grow from 10% of amplitude of periodic orbit to 90% of amplitude of periodic orbit takes about 51 periods, which means in practice, if the solution is only observed for a short period of time, only very slowly growing oscillations will be seen.

Figure 6: **Slow dynamics**. The black curve is the equilibrium E_p. As figure suggests the red and green curves tend very slowly to the stable periodic orbit. Period of periodic orbit is 57. The green curve after 38 periods and the red curve after 83 periods tend to stable limit cycle. The points marked by yellow circles on the red curve show it takes about 51 periods for the amplitude of red curve to grow to 80% of amplitude of stable periodic orbit. This figure suggests that there exists an area of state spaces in which slow dynamics occurs. This simulation is conducted for $X_U = 0.9$, $s = 0.5$, $Y_I = 0.08$, $m = 0.1$ and $b \sim 5.35$. This b is the mid point of the interval $(\frac{s(X_U - Y_I)}{X_U}, \frac{X_U - Y_I}{Y_I})$, on which E_P exists.

Figure 7 which is a modified version of Figure 4 shows fast dynamics. Under the set of parameters that this simulation is conducted equilibrium E_p, which is marked with black circle, has three eigenvalues $e_1 = -0.212$ and $e2, e3 = 0.001 \pm 0.11I$. Therefore, in a neighborhood of E_P solutions oscillate with the period $\frac{2\pi}{\text{Im}(e_2)} \sim 57$. To show fast attraction to the global attracting surface a thousand random points, marked by yellow circles, are chosen. Each solution starting from a point marked yellow, terminates in a purple point after 57 days. In Figure 7 same plot has been provided from two different views to provide a better observation that all the purple points are located on the slow manifold. Slow manifold includes the global attracting surface and extends outside of the limit cycle including the solid green area. Solutions on the slow manifold tend very slowly to the limit cycle, as we discussed in Figure 6.

Figure 7: **Fast attraction and collapse to the global attracting surface over one period**. The two plots are the same simulation shown in different views. The black closed curve bounding the red disk is an attracting limit cycle and the black circle near the center of the red disk is the equilibrium E_p. In this figure thousand random points, yellow dots, are chosen, and the positions of their trajectories after 57 days are shown with purple dots. All the purple dots appear to lie on a 2-dimensional surface that contains the limit cycle, E_p and the unstable manifold of E_p(The global attracting surface is the disk centered at Equilibrium E_p and includes the disk boundary which is the limit cycle). This shows each trajectory tends quickly to the global attracting surface.

5.2. Dependency of virulence threshold on viral time scale and horizontal transmission rate

Recall parameter $Y_I = 1 - \frac{a+d}{s}$. Changes in parameters a and s lead to changes in Y_I. A more aggressive virus causes more increase in death rate and more decrease in growth rate of cancer cells after infection, i.e. higher a and lower s. Notice that when a increases or s decreases, Y_I decreases. Hence, lower Y_I values means a stronger virus is administrated. Thus, we can use Y_I as a measure of level of cytotoxicity of virus. It is important to know how virulent a virus should be engineered so that more tumor shrinkage under therapy can be obtained. Our simulations below suggest that identifying the optimum range for virulence level of virus depends on both the viral time scale and infection horizontal transmission rate.

Figure 8 is a contour plot of maximum tumor size with respect to parameters b and Y_I in two scenarios; (a) fast viral time scale, $m = 1$ (b) slow viral time scale, $m = 0.1$. This figure recommends when parameter b and Y_I are chosen in the area below solid curve (this curve is labeled by $b = \frac{s(X_U - Y_I)}{X_U}$), then therapy is not effective regardless of viral time scale. Additionally, when parameters b and Y_I are chosen in the area above of the dotted curve (this curve is labeled by $b = \frac{X_U - Y_I}{Y_I}$), for a fixed b value, by decreasing Y_I, tumor size decreases, regardless of viral time scale.

Figure 8 suggests the viral time scale affects outcome of therapy when parameters Y_I and b are chosen in the area between the solid and dotted curves.

As Figure 8a illustrates when viral time scale is fast, the best range for parameter b and Y_I is the area between solid curve and the dotted curve.

When viral time scale is slow, as Figure 8b represents the most tumor shrinkage is obtained when the parameters Y_I and b are chosen along the dashed curve. The dashed curve labeled by $H = 0$ shows an implicit plot of roots of Routh function (Eq. (13)). Assume $b = 5.35$ is fixed, in this case $H(Y_I = 0.08) = 0$. When $b = 5.35$ is fixed and Y_I decreases and passes through the value 0.08, the maximum tumor size increases. Therefore, Figure 8b suggests when viral timescale is slow, high level of virulence has drawbacks on the outcome of viral therapy. Therefore, when viral time scale is slow, the best range for parameters Y_I and b is the area between the three curves.

(a) fast viral time scale   (b) slow viral time scale

Figure 8: **Changes in the maximum tumor size with respect to b and Y_I.** This figure represents how tumor size changes when parameters b and Y_I changes. Parameter values are $s = 0.5$, $X_U = 0.9$, and $m = 1$ in plot (a), and $m = 0.1$ in plot (b). Plot (a) shows when virus dynamics is fast, the maximum tumor shrinkage under therapy is possible for lower Y_I and higher b values . Plot (b) shows when virus dynamics is slow, lower Y_I values will not guarantee the best outcome. In this case, the smallest tumor size is achieved when parameter b and Y_I are chosen such that $H = 0$.

In Figure 9, five different values of Y_I are chosen, to find out the effective level of virus cytotoxicity for a fixed b value. The value of Y_I for which the red curve is simulated is $Y_I = \frac{X_U}{1+b}$. The value of Y_I for which black curve is plotted, when $m = 0.1$, is the value of Y_I for which $H = 0$.

In the case fast viral time scale, in other words when m is high enough that it is not in the interval (m_1, m_2), no periodic orbit exists. In this case as Figure 8a and Figure 9a suggest for $Y_I \leq \frac{X_U}{b+1}$, changes in Y_I does not affect the tumor size much. The blue, black and green curve in Figure 9a are all corresponding to the cases where $Y_I < \frac{X_U}{b+1}$. There is not much difference in their eventual tumor size with the case when $Y_I = \frac{X_U}{b+1}$. However, the tumor size is eventually smaller when $Y_I = \frac{X_U}{b+1}$. Therefore, for higher values of m, and fixed b value, the best range for Y_I is $(0, \frac{X_U}{1+b}]$ and the most tumor shrinkage is achievable at $Y_I = \frac{X_U}{1+b}$.

Figure 9b shows the tumor size over time when the viral time scale is slow and parameter b is fixed. As figure illustrates when $Y_I < \frac{X_U}{1+b}$, system shows oscillatory behavior: both green and black curves are decaying oscillations. The value of Y_I for which the black curve is simulated is when $H = 0$, therefore a Hopf-bifurcation occurs at this point, $Y_I = 0.088$. Therefore, the amplitude of the oscillations takes very long time to tend to zero. Hence, in practice if solutions are only observed for a short time, black curve seems like a stable periodic solution. When $Y_I = 0.01 < 0.088$ (the blue curve), the tumor size is periodic with a very high amplitude. As the small box in the right corner of Figure 9b suggests there is not much difference in the eventual tumor size for the black, green and red curves. If the solution be observed for a very long period of time, eventually the red curve would be lower, but in practice it is not possible. Therefore, when the viral time scale is slow, for a fixed b value, the best range for Y_I is between the value of Y_I at which H equals zero and $Y_I = \frac{X_U}{1+b}$. In this case the most tumor shrinkage is observed at the value of Y_I for which $H = 0$.

(a)

(b)

Figure 9: **Effect of cytotoxicity Y_I on the tumor size.** Plot (a) $m = 1$ and Plot (b) $m = 0.1$. As Plot (a) shows for higher values of m, when $Y_I \leq \frac{X_U}{b+1}$, eventually the tumor size is smaller. Therefore, the best range for parameter Y_I, $0 < Y_I < \frac{X_U}{1+b}$. For $m = 0.1$, as plot (b) suggest when values of Y_I decrease, dynamics becomes oscillatory and the amplitude of oscillation increase by decreasing Y_I. For $Y_I = 0.01$ stable oscillation occurs with a very high amplitude. As the small panel at the right top corner of plot (b) shows, both green and black curve or decaying oscillation but it takes very long time for the amplitude of the black curve to tends to zero. When $m = 0.1$, the best range for $0.088 < Y_I \leq \frac{X_U}{b+1}$.

6. Discussion

This paper offers a very simple mathematical model, Model (1), to analyze the interactions between cancer cells and free viruses during an oncolytic viral therapy. Even though Model (1) is simple, it captures some important features about oncolytic viral therapy: features that help to set expectations about the outcome of the treatment and how to engineer the vaccine in order to get better results. For example, Lemma 3 shows the tumor eradication equilibrium, E_0, is always unstable. This means complete tumor eradication is not possible. This suggest an oncolytic virus that only has an oncolysis effect cannot be used as a cure for cancer. However, Proposition 1 implies that the tumor size under therapy over time is always lower compare to the case with no therapy. Therefore, oncolytic viral therapy can be used to control the speed of growth of a tumor. Proposition 2 and Lemma 3 together show that oncolytic viral therapy can be used to considerably reduce the tumor size. The proposition suggests oncolytic viral therapy may be used as an intervention method to control the size of tumor, before a conventional therapy can be followed up.

One of our goals in this study was to optimize the therapy by recommending the parameter regions in which the therapy has the best outcome. To do this we focused our analysis on some parameter combinations: a measure of virulence level, Y_I, a rescaled horizontal transmission rate, b, and the viral clearance rate relative to the mutation rate of uninfected cancer cells, m.

To ease the communication with the reader throughout the paper we denoted the stable steady state tumor size when therapy fails, or if no treatment occurs, by X_U and the minimal steady state tumor size under the therapy by Y_I. In Proposition 2, we showed that the interval $[Y_I, X_U]$ is a global attracting set for the tumor size. Specifically, for a specific virus, i.e. for a fixed set of parameters, the tumor size by therapy is eventually bigger or equal to Y_I. Therefore, Y_I is the minimum tumor size under the therapy.

Lemma 3 and Theorem 2 specify two possible outcomes for the treatment: 1-failure, 2-partial success or tumor control. Partial success means considerable reduction in tumor size. Partial success includes two scenarios: 2(a) tumor partially gets infected 2(b) 100% infection prevalence occurs.

1-Failure: Treatment fails when the horizontal rate of infection transmission is smaller than control threshold. In this case therapy has no impact and tumor grows to the maximum size X_U, same as when no therapy would take place. long term tumor reduction will be possible when horizontal rate of infection transmission is above of control threshold.

2-Partial success:
2(a). **partial infection:** This scenario occurs when the infection transmission rate is between control threshold and optimum threshold. In this case, there are two possible scenarios; if viral time scale is slow, the positive equilibrium is unstable and every solution starting from a positive point which is not on the stable manifold of the positive equilibrium tends to the stable limit cycle. When viral time scale is fast the positive equilibrium is globally stable and no limit cycle exists. See Figures 4 and 5. In this case, as Propositions 1 and 2, and Lemma 2 suggest since horizontal infection transmission is above of the control threshold therapy reduces the tumor size, in comparison with the case that no therapy occurs, but it is not the minimum possible tumor size under the therapy.

2(b). **100% infection prevalence:** The maximum tumor shrinkage by therapy is obtained in this scenario which happens if infection horizontal transmission is equal to optimum threshold.

In Model (1), we assumed (assumptions (A2) and (A4)) cancer cells are mitotic and the growth rate of infected cancer cells is smaller than their apoptosis rate. If we relax this assumption, meaning either the infected cells do not self-renew or the self-renewal rate of infected cancer cells be lower than their apoptosis rate, then 100% infection prevalence will not be possible. Therefore, only two scenarioes would be possible: failure or partial infection. Furthermore, the tumor shrinkage by therapy will be less in comparison with our model which considers two pathways to infection transmission.

In this study interactions with immune cells and spatial heterogeneity (neither morphological nor phenotyp-
ical), are not discussed. Model (1) can be extended to consider the effect of virus specific immune response
that seems to be one of the main barriers in the success of oncolytic viral therapy. In our work in progress,
by extending the model to also consider interactions of virus specific immune responses with cancer cells and
the free virus, in most cases no longer oscillatory behavior is observable. In the presence of virus specific
immune responses, the reduction in tumor size will be lower in comparison with the case no interaction with
immune cells is taken into account, but oncolytic therapy seems yet promising as considerable reduction in
tumor size is still observable.

One of the assumptions under which we started the study was assumption (A5). One might question how
virulent the virus should be? As we discussed before parameter Y_I, the minimum tumor size, can also be
used as a measure of level of cytotoxicity of the virus. Lower values of Y_I means virus is more virulent.
As Figures 8 and 9 illustrate electing the best range for virulence level also depends on the parameter m.
Parameter m is the relative clearance rate of the free virus, and gives the timescale of the virus dynamics.
When parameter m is large, meaning $\bar{m} >> \bar{r}$, as earlier studies by Komarova and Wodarz [18] suggested
the system is in quasi-steady state. Meaning, $y - v$ rapidly approaches zero, therefore Model 1 can be
reduced to the "slow" x-y subsystem. The maximum level for cytotoxicity recommended by Komarova and
Wodarz [18] in terms of parameters of Model (2) is equivalent to $Y_I = \frac{X_U}{b+1}$ (notice that maximum level
of cytotoxicity means lowest value recommended for Y_I). However, our results show that if the parameter
m is high enough that does not belong to the interval (m_1, m_2) (see Theorem 2 for m_1, and m_2), then
$0 < Y_I \leq \frac{X_U}{b+1}$. Our simulations in Figures 8 and 9 recommend the highest value for Y_I is $X_U/(b+1)$.
Although, there is not a huge difference in the amount of tumor shrinkage corresponding to Y_I values within
the interval $(0, X_U/(b+1))$. If $m \in (m_1, m_2)$, then Y_I should be chosen between the values of Y_I for which
the Routh function $H = 0$ (see Eq. (13)), and $\frac{X_U}{b+1}$. This suggest when m is lower, in other terms, virus
dynamics is slow, a very high level of virulence is not recommended.

As we mentioned above oncolytic viral therapy can be used as a method for controlling tumor size. What
we suggested above about parameter regions is mostly considered when the viral therapy is going to be used
alone. Therefore, we suggest the parameters are chosen in the way that sustained oscillations do not occur,
specifically parameters should be chosen in the way $H \geq 0$. In contrast, if oncolytic viral therapy is going
to be followed by another therapy, one may reconsider the lower values of Y_I for which sustained oscillations
occur. As we observed in Figure 9b, when the virus is highly virulent and proper initial dosage of virus is
applied, early tumor shrinkage occurs to less than 11% of the initial tumor size and the tumor remains in
this size for almost a year before starting to relapse. For example, when $Y_I = 0.088$ (the black curve), tumor
size reaches to 9.8% of initial tumor size at $t = 266$ or the most aggressive case when $Y_I = 0.01$ tumor size
reaches to 2% of initial tumor size at $t = 150$. This raises multiple questions, such as, what happens if the
treatment switches to another therapy before tumor relapse? which treatment would be best as the followed
up treatment? We suggest that a virulent virus can be used, but before tumor relapses the second therapy
must be followed up.

Appendix A. Appendix

Appendix A.1. Derivation of dimensionless Model (2)

Here we derive the dimensionless Model (2). First we recall Model (1).

$$
\begin{aligned}
\frac{dX}{dT} &= \bar{r}X(1 - \frac{X+Y}{K}) - \bar{d}X - \beta XV \\
\frac{dY}{dT} &= \bar{s}Y(1 - \frac{X+Y}{K}) - \bar{d}Y - \bar{a}Y + \beta XV \\
\frac{dV}{dT} &= \bar{a}\bar{q}Y - \bar{m}V
\end{aligned}
$$

454 Now let $x = \frac{X}{K}$, $y = \frac{Y}{K}$, $v = \frac{\bar{m}}{K\bar{q}\bar{a}}V$ and $t = \bar{r}T$, then

$$\frac{d(Kx)}{dt} = Kx(1 - x - y) - \frac{\bar{d}}{\bar{r}}Kx - \frac{\beta}{\bar{r}}(Kx)(\frac{K\bar{q}\bar{a}}{\bar{m}}v),$$

$$\frac{d(Ky)}{dt} = \frac{\bar{s}}{\bar{r}}Ky(1 - x - y) - \frac{\bar{d}}{\bar{r}}Ky - \frac{\bar{a}}{\bar{r}}Ky + \frac{\beta}{\bar{r}}(Kx)(\frac{K\bar{q}\bar{a}}{\bar{m}}v),$$

$$\frac{d(\frac{K\bar{q}\bar{a}}{\bar{m}})}{v})dt = \frac{\bar{a}\bar{q}}{\bar{r}}Ky - \frac{\bar{m}}{\bar{r}}(\frac{K\bar{q}\bar{a}}{\bar{m}}v).$$

After some algebraic simplifications the above system of equations turns to

$$\frac{dx}{dt} = x(1 - x - y) - \frac{\bar{d}}{\bar{r}}x - \frac{\beta}{\bar{r}}(\frac{K\bar{q}\bar{a}}{\bar{m}})xv,$$

$$\frac{dy}{dt} = \frac{\bar{s}}{\bar{r}}y(1 - x - y) - \frac{\bar{d}}{\bar{r}}y - \frac{\bar{a}}{\bar{r}}y + \frac{\beta}{\bar{r}}(\frac{K\bar{q}\bar{a}}{\bar{m}})xv,$$

$$\frac{dv}{dt} = \frac{\bar{m}}{\bar{r}}y - \frac{\bar{m}}{\bar{r}}v.$$

Now we will call dimensionless parameters as follows:

$$a = \frac{\bar{a}}{\bar{r}}, \quad d = \frac{\bar{d}}{\bar{r}}, \quad s = \frac{\bar{s}}{\bar{r}}, \quad m = \frac{\bar{m}}{\bar{r}}, \quad b = \frac{\beta}{\bar{r}}(\frac{K\bar{q}\bar{a}}{\bar{m}})$$

Using the new dimensionless parameters the above system of equation gets the following form.

$$\frac{dx}{dt} = x(1 - x - y) - dx - bxv,$$

$$\frac{dy}{dt} = sy(1 - x - y) - dy - ay + bxv,$$

$$\frac{dv}{dt} = m(y - v).$$

Now let $X_U = 1 - d$ and $Y_I = 1 - \frac{a+d}{s}$, then Model (1) gets the following dimensionless form.

$$\frac{dx}{dt} = x(X_U - (x + y)) - bxv$$

$$\frac{dy}{dt} = sy(Y_I - (x + y)) + bxv$$

$$\frac{dv}{dt} = m(y - v)$$

Appendix A.2. Detailed proof of lemma 1

In this section we will show what happens in each boundary of the set $\mathcal{W}$.

On the x axis Model (2) simplifies to: $\frac{dx}{dt} = x(X_U - x), \frac{dy}{dt} = 0, \frac{dv}{dt} = 0$. Hence, each solution starting from a point on the x axis with $x \geq 0$ remains on the x axis. Except $E_0 = (0,0,0)$ which is an equilibrium of the system, each solution $(x(t),0,0)$ with $x(0) > 0$ tends to $E_U = (X_U,0,0)$ as t increases.

On the y axis Model (2) simplifies to: $\frac{dx}{dt} = 0, \frac{dy}{dt} = sy(Y_I - y), \frac{dv}{dt} = my$. Since $\frac{dx}{dt} = 0$, $x(t) = 0$ for all $t \geq 0$.

The solution to ode $\frac{dy}{dt} = sy(Y_I - y)$, with initial point $y(0)$, is $\frac{Y_I y(0)}{y(0) + (Y_I - y(0))e^{-sY_I t}}$. Therefore, if $y(0) > 0$, then $y(t) > 0$ for all $t \geq 0$.

Since $\dfrac{dv}{dt} = my > 0$ when $y > 0$, v is increasing. Therefore, $v(t) \geq 0$ for all ≥ 0.

Hence, each solution starting from a point $(0, y_0, 0)$ with $y_0 > 0$ remains on the $y - v$ plane.

On the v axis Model (2) simplifies to: $\dfrac{dx}{dt} = 0$, $\dfrac{dy}{dt} = 0$, $\dfrac{dv}{dt} = -mv$. Therefore, a solution starting on the v

axis, remains on the v axis. Additionally, since $\dfrac{dv}{dt} < 0$ when $v > 0$, each solution starting from a point on

the v axis with $v_0 > 0$, tends to E_0.

On the x-v plane $y = 0$. Hence, Model (2) simplifies to: $\dfrac{dx}{dt} = x(X_U - x) - bxv$, $\dfrac{dy}{dt} = bxv$, $\dfrac{dv}{dt} = -mv$.

Therefore, $v(t) = v(0)e^{-mt}$. v is non-negative and decreasing. Since $\dfrac{dy}{dt} > 0$ when $x, v > 0$, starting from an

initial point with $x(0), v(0) > 0$, y increases. Therefore, y remains positive. Thus, any solution starting from

a point in the x-v, except the points on the x and v axes, tends inward W, i.e. in the direction of increasing

y.

On the x-y plane $v = 0$. Thus, Model (2) simplifies to; $\dot{x} = x(X_U - (x + y))$, $\dot{y} = sy(Y_I - (x + y))$, $\dot{v} = my$.

When $y(0) > 0$, $\dot{v} > 0$. Hence, v is increasing. Each solution that starts from a point $(x(0), y(0), 0) \in W$

where $y(0) > 0$, points inward the W as t increases.

On the y-v plane, $x = 0$. Thus, Model (2) simplifies to: $\dot{x} = 0$, $\dot{y} = sy(Y_I - y)$, $\dot{v} = m(y - v)$.

Since $\dot{x} = 0$, therefore $x(t) = 0$ for all $t \geq 0$. Hence, each solution starting on this plane, is tangent to the plane. Solution to the foregoing system of differential equations with an initial point $(0, y(0), v(0))$ is

$$x(t) = 0,$$
$$y(t) = \frac{Y_I y(0)}{y(0) + (Y_I - y(0))e^{-sY_I t}},$$
$$v(t) = me^{-mt} \int_0^t y(z)e^{mz} dz + v(0)e^{-mt}.$$

478 Therefore, each solution that starts from a point in the y-v plane with $y_0 \neq 0$ goes eventually to E_I.

479 Figure A.10 shows the vector field on the y-v plane.

Figure A.10: Any solution that starts in the y-v plane will remain the y-v plane. All the solutions starting in y-v plane with $y(0) > 0$, eventually tends to E_I.

480 The intersection of W with the plane $v = X_U$ is above of $y = v$ plane, this boundary only meets the plane

481 $y = v$ at point $(0, X_U, X_U)$). Hence, $\dot{v} < 0$ every where on this boundary except point $(0, X_U, X_U)$. Therefore,

v is decreasing. Therefore, any solution starting on this plane points inward of $\mathcal{W}$. Since $(0, X_U, X_U)$ is located in y-v plane, the solution starting from $(0, X_U, X_U)$ tends to E_I.

When $\mathcal{W}$ meets the plane $x(0) + y(0) = X_U$, $\dot{x} + \dot{y} = sy(Y_I - X_U)$. According to assumption (H2), $Y_I < X_U$. Thus, $x + y$ is decreasing when $y \neq 0$, so each solution that starts from a point on the intersection of $\mathcal{W}$ with plane $x + y = X_U$, points inward $\mathcal{W}$. $\square$

Appendix A.3. Hopf Bifurcation.

Consider a three dimensional system of ordinary differential equations is given, meaning

$$\dot{U} = f(U, \mu), \tag{A.1}$$

where $U \in \mathbb{R}^3$ and $\mu \in \mathbb{R}$.

Hopf Bifurcation Theorem. A Hopf-bifurcation occurs at (U^*, μ^*) for the system (A.1) if there exists a curve of stationary points $(U_{eq}(\mu), \mu) \in \mathbb{R}^{3+1}$ such that $(U_{eq}(\mu^*), \mu^*) = (U^*, \mu^*)$ and $Df\big(U_{eq}(\mu), \mu\big)$ has a pair of complex eigenvalues $\lambda(\mu) = R(\mu) \pm iI(\mu)$, such that

HB(1) $R(\mu^*) = 0$ and $I(\mu^*) \neq 0$,

HB(2) $\dfrac{dR}{d\mu}(\mu^*) \neq 0$,

and the real eigenvalue of $Df\big(U_{eq}(\mu), \mu\big)$ is nonzero for all values of μ.

Appendix A.4. Comparison Theorem

Theorem 3 (First Comparison Theorem [28])**.** Let $x(t)$ and $y(t)$ be continuously differentiable functions from $[0, +\infty)$ to $\mathbb{R}$, satisfying

$$\frac{dx}{dt} < \frac{dy}{dt} \qquad \text{where } x(0) < y(0), \tag{A.2}$$

then, $x(t) < y(t)$ for all $t > 0$.

Acknowledgments

We thank James Yorke and Kamran kaveh for their helpful comments. James Watmough was partially supported by NSERC (RGPIN-2017-05760) and CIHR, the Mathematical Modelling of COVID-19 Task Force. Lin Wang was partially supported by NSERC (RGPIN-2020-04143). James Watmough and Lin Wang were also partially supported by New Brunswick Health Research Fund.

References

[1] Kelly E, Russell SJ. History of Oncolytic Viruses: Genesis to Genetic Engineering. Molecular Therapy 2007;15(4):651–9. doi:`10.1038/sj.mt.6300108`.

[2] Bridle BW, Stephenson KB, Boudreau JE, Koshy S, Kazdhan N, Pullenayegum E, et al. Potentiating Cancer Immunotherapy Using an Oncolytic Virus. Molecular Therapy 2010;18(8):1430–9. doi:`10.1038/mt.2010.98`.

[3] Jenner AL, Yun CO, Yoon A, Coster ACF, Kim PS. Modelling combined virotherapy and immunotherapy: strengthening the antitumour immune response mediated by IL-12 and GM-CSF expression. Letters in Biomathematics 2018;5(sup1):S99–. doi:`10.1080/23737867.2018.1438216`.

[4] de Graaf J, de Vor L, Fouchier R, van den Hoogen B. Armed oncolytic viruses: A kick-start for anti-tumor immunity. Cytokine & Growth Factor Reviews 2018;41:28–39. doi:`10.1016/j.cytogfr.2018.03.006`.

[5] Gujar S, Bell J, Diallo JS. SnapShot: Cancer Immunotherapy with Oncolytic Viruses. Cell 2019;176(5):1240–1240.e1. doi:`10.1016/j.cell.2019.01.051`.

[6] Rivadeneira DB, DePeaux K, Wang Y, Kulkarni A, Tabib T, Menk AV, et al. Oncolytic viruses engineered to enforce leptin expression reprogram tumor-infiltrating t cell metabolism and promote tumor clearance. Immunity 2019;51(3):548–560.e4. doi:`10.1016/j.immuni.2019.07.003`.

[7] Bell JC, Lichty B, Stojdl D. Getting oncolytic virus therapies off the ground. Cancer Cell 2003;4(1):7–11. doi:`10.1016/S1535-6108(03)00170-3`.

[8] Macedo N, Miller DM, Haq R, Kaufman HL. Clinical landscape of oncolytic virus research in 2020. Journal for Immunotherapy of Cancer 2020;8(2). doi:`10.1136/jitc-2020-001486`.

[9] Dingli D, Offord C, Myers R, Peng KW, Carr TW, Josic K, et al. Dynamics of multiple myeloma tumor therapy with a recombinant measles virus. Cancer Gene Therapy 2009;16(12):873–82. doi:`10.1038/cgt.2009.40`.

[10] Zhang Q, Liu F. Advances and potential pitfalls of oncolytic viruses expressing immunomodulatory transgene therapy for malignant gliomas. Cell Death & Disease 2020;11(6):1–11. doi:`10.1038/s41419-020-2696-5`.

[11] Harrison D, Sauthoff H, Heitner S, Jagirdar J, Rom WN, Hay JG. Wild-type adenovirus decreases tumor xenograft growth, but despite viral persistence complete tumor responses are rarely achieved–deletion of the viral E1b-19-kD gene increases the viral oncolytic effect. Human Gene Therapy 2001;12(10):1323–32. doi:`10.1089/104303401750270977`.

[12] Fountzilas C, Patel S, Mahalingam D. Review: Oncolytic virotherapy, updates and future directions. Oncotarget 2017;8(60). doi:`10.18632/oncotarget.18309`.

[13] Bajzer Z, Carr T, Josić K, Russell SJ, Dingli D. Modeling of cancer virotherapy with recombinant measles viruses. Journal of Theoretical Biology 2008;252(1):109–22. doi:`10.1016/j.jtbi.2008.01.016`.

[14] Jenner A, Coster A, Kim P, Frascoli F. Treating cancerous cells with viruses: insights from a minimal model for oncolytic virotherapy. Letters in Biomathematics 2018;5:1–20. doi:`10.1080/23737867.2018.1440977`.

[15] Komarova NL, Wodarz D. ODE models for oncolytic virus dynamics. Journal of Theoretical Biology 2010;263(4):530–43. doi:`10.1016/j.jtbi.2010.01.009`.

[16] Wodarz D. Viruses as Antitumor Weapons: Defining Conditions for Tumor Remission. Cancer Research 2001;61(8):3501–7.

[17] Wodarz D. Computational approaches to study oncolytic virus therapy: insights and challenges. Gene Therapy and Molecular Biology 2004;:10.

[18] Komarova NL, Wodarz D. Mitotic Virus Transmission and Immune Responses. In: Targeted Cancer Treatment in Silico. Modeling and Simulation in Science, Engineering and Technology; New York, NY: Birkhäuser, New York, NY; 2014. ISBN 978-1-4614-8301-4. doi:`10.1007/978-1-4614-8301-4_12`.

[19] Yamamoto Y, Nagasato M, Yoshida T, Aoki K. Recent advances in genetic modification of adenovirus vectors for cancer treatment. Cancer Science 2017;108(5):831–7. doi:`10.1111/cas.13228`.

[20] Berkeley RA, Steele LP, Mulder AA, Wollenberg DJMvd, Kottke TJ, Thompson J, et al. Antibody-Neutralized Reovirus Is Effective in Oncolytic Virotherapy. Cancer Immunology Research 2018;6(10):1161–73. doi:`10.1158/2326-6066.CIR-18-0309`.

[21] Felt SA, Grdzelishvili VZ. Recent advances in vesicular stomatitis virus-based oncolytic virotherapy: a 5-year update. The Journal of General Virology 2017;98(12):2895–911. doi:`10.1099/jgv.0.000980`.

[22] Fu X, Rivera A, Tao L, De Geest B, Zhang X. Construction of an Oncolytic Herpes Simplex Virus That Precisely Targets Hepatocellular Carcinoma Cells. Molecular Therapy 2012;20(2):339–46. doi:`10.1038/mt.2011.265`.

[23] Trus I, Berube N, Jiang P, Rak J, Gerdts V, Karniychuk U. Zika Virus with Increased CpG Dinucleotide Frequencies Shows Oncolytic Activity in Glioblastoma Stem Cells. Viruses 2020;12(5):579. doi:`10.3390/v12050579`.

[24] Zheng M, Huang J, Tong A, Yang H. Oncolytic Viruses for Cancer Therapy: Barriers and Recent Advances. Molecular Therapy - Oncolytics 2019;15:234–47. doi:`10.1016/j.omto.2019.10.007`.

[25] Koscielny S, Tubiana M, Lê MG, Valleron AJ, Mouriesse H, Contesso G, et al. Breast cancer: Relationship between the size of the primary tumour and the probability of metastatic dissemination. British Journal of Cancer 1984;49(6):709–15. doi:`10.1038/bjc.1984.112`.

[26] Narod SA. Tumour size predicts long-term survival among women with lymph node-positive breast cancer. Current oncology 2012;19(5):249–53. doi:`10.3747/co.19.1043`.

[27] Nowak MA. The Mathematical Biology of Human Infections. Conservation Ecology 1999;3(2).

[28] Liao X, Wang L, Yu P. Stability of dynamical systems. 5; 1st ed.; Amsterdam ; Boston: Elsevier; 2007.

[29] Kuznetsov YA. Elements of Applied Bifurcation Theory. 2nd ed.; New York: Springer; 1998.

[30] Liu WM. Criterion of Hopf Bifurcations without Using Eigenvalues. Journal of Mathematical Analysis and Applications 1994;182(1):250–6. doi:`10.1006/jmaa.1994.1079`.

[31] Routh EJ. On Laplace's Three Particles, with a Supplement on the Stability of Steady Motion. Proceedings of the London Mathematical Society 1874;s1-6(1):86–97. doi:`10.1112/plms/s1-6.1.86`.

[32] Mallet-Paret J, Yorke JA. Snakes: Oriented families of periodic orbits, their sources, sinks, and continuation. Journal of Differential Equations 1982;43(3):419–50. doi:`10.1016/0022-0396(82)90085-7`.